1

**Single-cell transcriptomic profiling of the whole colony of *Botrylloides diegensis*:**

**Insights into tissue specialization and blastogenesis**

**Authors:** Berivan Temiz[1], Michael Meier[1-2], and Megan J. Wilson[1*]

[1]Developmental Biology and Genomics Laboratory, Department of Anatomy, Otago School

of Biomedical Sciences, University of Otago, P.O. Box 56, Dunedin 9054, New Zealand[1]

[2]Department of Pathology, Otago School of Biomedical Sciences, University of Otago,

Dunedin 9054, New Zealand[2]

**\* Corresponding Author:** Megan J. Wilson

**Mail:** meganj.wilson@otago.ac.nz

**Orcid IDs of the Authors**

Berivan Temiz - https://orcid.org/0000-0002-0499-8301

Megan J. Wilson - https://orcid.org/0000-0003-3425-5071

Running title: Single-Cell Profiling of *B. diegensis* Colony

Key words: ascidian, progenitor, single-cell sequencing, *Botrylloides*

Summary statement: Single-cell RNA sequencing of *Botrylloides diegensis* revealed cellular heterogeneity, identified 29 major cell clusters, and provided insights into tissue specialization and blastogenesis.

**Abstract**

*Botrylloides diegensis* is a colonial ascidian that has been the focus of developmental, evolutionary, and regeneration research. In this study, we performed single-cell RNA sequencing (scRNA-seq) of an entire *B. diegensis* colony, including zooids, buds, and vascular tunics, to resolve cellular heterogeneity and identify cell and tissue markers. We identified 29 major cell clusters within the colony and used *in situ* hybridization to examine the spatial expression of cluster marker genes. Numerous tissue types were identified at the molecular level, including blood cells and zooid tissues such as the branchial epithelium, stomach, and endostyle. Distinct cluster markers were identified for specific regions of the stomach epithelium, highlighting the specialization of these regions and the strength of using scRNA-seq to explore their functionality. Cell trajectory projections highlighted the early appearance of progenitor clusters, whereas more differentiated zooid-related tissues appeared later in the developmental path. This study provides a valuable resource for understanding the development, tissue function, and regeneration of *B. diegensis.* This demonstrates the power of scRNA-seq to define cell types and tissues in complex colonial organisms.

**Introduction**

Single-cell RNA sequencing (scRNA-seq) can quantify the total transcriptome of individual cells in the tissue/organism of interest. This information can be used to classify, characterize, and distinguish each cell at the mRNA level, allowing for the identification of different cell types, developmental states, and trajectories. Many studies have mapped the transcriptomic composition of diverse animals, including invertebrates such as planaria, sponges, *Hydra*, and ascidian species. These studies are essential to understand how cell types differ from each other

through the activation or repression of specific pathways; therefore, they can explain how cells

possess pluripotency, commit to becoming specific biological units, access categorical

morphology, and become a part of tissue during development or regeneration.

Colonial ascidians are sessile marine chordates (Fig. 1A), which can be categorized into three

main parts:1) zooids, buds, and budlets as the asexually developing body; 2) the vascular

network, including veins, blood cells, and vascular termini/ampullae; and 3) tunic, the

gelatinous matrix covering all colonies (Berrill, 1947a; Blanchoud et al., 2017) (Fig. 1B and

C). At least 11 blood cell types have been identified in the hemolymph and are grouped into

five cell types:1) undifferentiated cells, hemoblasts, and differentiating cells; 2) immunocytes,

hyaline amebocytes, macrophage-like cells, granular amebocytes, and morula cells; 3)

transport cells, compartment amebocytes, and compartment cells; 4) mast cell-like cells or

granular cells; and 5) storage cells, pigment cells, or nephrocytes(Blanchoud et al., 2017; Cima

et al., 2001; Hirose et al., 2003; Wright and Ermak, 1982). The zooid, as a filter-feeding

individual, consists of various anatomical structures, including the atrial epithelium, atrial

siphon, branchial epithelium, bud, endostyle, gonads, intestine, neural complex, peribranchial

sacs, pericardium, pyloric gland, stigmata, and the stomach (Fig. 1D and E) (Anselmi et al.,

2022; Berrill, 1947a; Blanchoud et al., 2017; Holland, 2016; Kawamura and Sunanaga, 2010).

During asexual reproduction and blastogenesis, a new zooid is produced by budding from the

atrial epithelium of the parental zooid (Berrill, 1947a; Manni and Burighel, 2006; Mukai et al.,

1978). This budding process, also known as blastogenesis or palleal budding, is

morphologically similar to whole-body regeneration (WBR) (Berrill, 1951; Rinkevich et al.,

2007). Thus, studying intact whole colonies will help capture the critical aspects of their

development, tissue function, and regeneration.

Several sc-RNA-seq studies have been performed with the solidary ascidian *Ciona intestinalis*,

mainly during the embryonic stage, showing the conservation of developmental and functional

programs in chordate evolution (Cao et al., 2019; Horie et al., 2018; Sharma et al., 2019; Zhang

et al., 2020). *B. diegensis* is an emerging model for development, regeneration, and stem cell

studies owing to its small genome size, short replenishment period, and excellent example of

WBR (Blanchoud et al., 2018). Therefore, investigating the composition of mature colonies is

crucial to understand the cell types and tissues of *B. diegensis*.

This study aimed to develop a protocol and resource for *B. diegensis* to identify cell and tissue

markers. Single-cell transcriptomic profiling of a mature (blastogenic stage A) *B. diegensis*

colony was performed. Consequently, several cell and tissue types were separated based on

their unique gene expression profiles, and their spatial expression was validated using *in situ*

hybridization.

**Figure 1. The study system: *Botryllodies diegensis.*** A. Simplified phylogeny represents the position
of tunicates as the closest chordate group to vertebrates. B. Example *B. diegensis* colony attached to a
glass slide. C. Orange morphs of *the B. diegensis* colony are illustrated. The red lines indicate shared
circulatory networks. The vascular termini (ampulla) are elliptical vascular ends. The individuals
(zooids) are lined side by side in a ladder-like arrangement in the gelatinous matrix (tunic). D and E.

Simplified zooid anatomy, dorsal (D) and lateral (E) views. Abbreviations: intestine (in), stomach (st),
gonad (go), neural complex (nc), endostyle (en), branchial epithelium (be).
99

## Results and Discussion

*B. diegensis* colonies were obtained from a local marina and kept in the laboratory. We

performed scRNA-seq using acetic acid-methanol (ACME) tissue dissociation and fixation

methods (Fig. 2A). Cells from a *B. diegensis* colony at stage A of the blastogenic cycle were

fixed and dissociated using ACME (Fig S1 A-B). This was followed by FACS sorting to

remove cell aggregates and debris (Fig.S1E-L). Chromium$^{TM}$ Single-cell 3' (10X Genomics)

was used for single-cell barcoding. The barcoded reads were sequenced in both directions for

paired-end sequencing. Library quality control was performed using FastQC software (Fig.

S2). High-quality reads were mapped back to the reference genome using STARSolo software.

The summary statistics for mapping are provided in Table S1. A single-cell transcriptome

library containing approximately 58 million reads was generated. The percentage of the aligned

reads with valid barcodes was 96%. The rate of reads that mapped to unique genes was 73%.

In total, 6353 cells were detected, with a mean of 1586 UMIs and 481 genes per cell. A total of

13621 genes were identified (Table S1).

Cluster analysis was performed to group single-cell transcriptomes based on similarity using

the Seurat software. Genes showing high variation between cells were calculated to obtain the

best signal in cluster differences, and 2,000 genes were included by default. Twenty-nine

distinct clusters were identified using nonlinear dimensional reduction with Uniform Manifold

Approximation and Projection (UMAP) (Fig 2B). The cell numbers for each cluster are shown

in Fig. S3. The highest cell number was observed for cluster 0, with 779 cells, and the lowest

cell number was 25 for cluster 28. The mean cell number/cluster ratio was 208. The average

number of genes in each cluster was 291.

**Figure 2. Overview of the experimental pipeline and clustering results.** A. Single cells were prepared using
the ACME maceration method (Garcia-Castro et al., 2021). Cells were stained with DRAQ5 and DRAQ5+ cells
were sorted by FACS. Cells were captured in droplets using a 10X chromium system. Single-cell libraries were
prepared and sequenced. After mapping the transcripts to the genome, clustering analysis was performed to
identify the cell types. B. UMAP clustering analysis revealed 29 single-cell clusters. Each cluster was color-
coded. C. Heatmap of the top five marker genes in each cluster. Abbreviations: acetic-methanol (ACME).

## Identification of Cluster Marker Genes

The FindMarkers function was used for differential expression analysis within the Seurat

package to identify the top genes that were potential markers for each cluster (Fig. 2C).

Differentially expressed genes (DEGs) were recorded as fold changes (LFC). The full DEG list

for each cluster is provided in File S1. DEGs were annotated to their closest human orthologs.

Genes that did not have vertebrate or human matches were also calculated for each cluster

(Table S2). Gene ontology (GO) analysis was conducted for each cluster to aid the functional

characterization of the clusters. The closest vertebrate orthologs that matched the DEGs were

used to generate gene lists for GO analysis (File S2).

**Using single-cell data to identify regional tissue markers**

The expression patterns of the top marker genes of Clusters 4, 5, and 14 were determined by

*in situ* hybridization (Fig. 3). *Ctrb1* (*g03753*) was one of the most highly expressed transcripts

(~8 LFC) in Cluster 4, which was exclusively present in this cluster compared to other clusters

(Fig 3A). *Ctrb1* encodes a serine protease enzyme linked to the acinar-like exocrine glandular

cells involved in digestion (Perillo et al., 2016). A probe was designed to determine the cell-

type expression of *Ctrb1,* and a strong staining signal was detected in the stomach of the zooid

(Fig 3A). These *Ctrb1+* epithelial cells were a subset of cells located in the outer curling of the

stomach folds (Fig. 3A). No staining was observed in other tissues or vascular cells in mature

colonies (Fig. 3A).

*Kng1 (g08491)* was identified as a marker for Cluster 14 (File S1) and was present to a lesser

extent in Clusters 2, 4, and 5. A 610 bp fragment of *Kng1* was cloned and sequenced for use

during *in situ* hybridization. Intense *Kng1* staining was observed within the lateral edges of the

stomach via *in situ* hybridization in adult tissue sections (Fig 3B). Thus, Cluster 14 was

identified as a part of the stomach epithelium.

Cluster 10 marker g14532, a cornifelin-like gene (*Cnfn)*), showed specific expression in

stigmata cells (Fig. 3C). This gene is localized to microtubules in humans, particularly in the

epidermis and oral mucosa (Wagner et al., 2019), and is associated with cell-cell adhesion. It

contains a cysteine-rich domain known as a PLAC8 domain.

*Tubb* was broadly upregulated in branchial epithelial tissues of the digestive tract (Fig. 3D).

Tubb is a tubulin beta protein that functions in the microtubules of the cytoskeleton, controlling

cell shape, movement, and transport within the cell (Sewell et al., 2024). Cilia are microtubule-

based organelles, and the zone 1 cilium has a different axonemal structure than the other zones

(*Ciona* endostyle) (Konno and Inaba, 2020), which may indicate that these cells express

distinct combinations of microtubule genes.

**Figure 3. Identification of *B. diegensis* digestive tract clusters.** A. Violin plot showing *Ctrb1* as a marker of
cluster 4. *In situ* expression of *Ctrb1* mRNA was observed in the cells on the outer side of each stomach fold. B.
*Kng1+* cells longitudinal stomach folds. C. Single-cell cluster expression profile for *Cnfn*, g04846, the top marker
gene for Cluster 11. The *Cnfn* probe-stained cells are found in rows (asterixis) that line the branchial chamber,
known as stigmata cells. D. Cluster expression for *Tubb* was identified as a top marker gene for Clusters 10 and
11, although it was expressed by cells found in several other Clusters. *Tubb* mRNA was found in the stomach
epithelium, branchial epithelial cells, and epithelia covering the oocytes/eggs of developing buds. E. Based on the
*in situ* results, an Illustration of a stomach fold showing the location of *Tubb+, Ctrb1+* and *Kng1+* cells in the
stomach epithelium. Abbreviations: endostyle (end), stomach (stm), inner stomach fold (isf), outer stomach fold
(osf), stigmata (sti), tunic (t), intestine (int).

**Figure 4. GO and pathway analyses for cell Clusters associated with digestive functions.** A. Metascape
network of enriched terms, colored by Cluster number and predicted tissue type. Each node is colored according

to the number of genes (from each list) associated with that term. B. A clustered heatmap of the top enriched GO
terms colored by p-values. Each row represents an enriched GO term, and each column represents the gene list
for Clusters 14, 4, 5, 10 and 11.

GO and pathway analyses were conducted for genes highly expressed in digestive- and

branchial-associated tissues (Clusters 4, 5, 10, 11, and 14) (Fig. 4). Clusters 10 and 11 were

overrepresented in the pathways linked to cilia assembly and movement (Fig. 4). This aligns

with the mRNA expression of *Cnfn* and *Tubb,* markers of cell clusters 11 and 10, respectively,

which showed intense staining in cilia-rich cell types (Fig. 3C and D). Annotation of clusters

4, 14, and 5 confirmed the enrichment of biological processes associated with food breakdown,

such as metabolic and catabolic oxidoreductase activity and cell secretion (Fig. 4).

**Identification of Endostyle clusters**

Endostyle, a tissue similar to the pharynx, filters food, synthesizes hormones, and provides

immune defences (Takagi et al., 2022). They are also believed to possess stem cell niches or

hematopoietic properties (Voskoboynik et al., 2008b). Genes known to be expressed in the

endostyle of other ascidian species (*Ciona, Styela*, and *Botryllus*) were selected and identified

using the single-cell dataset (Table S4). Clusters 6, 3, and 8 were deemed potential endostyle

clusters based on the expression of known endostyle markers. In the ascidian *Ciona*, galectin

is expressed in various regions of the endostyle (Parrinello et al., 2015; Parrinello et al., 2017).

The *B. diegensis* genome contains multiple *Lgal* genes (File S3) whose transcripts were

detected in clusters 3, 4, and 10 (File S3) (Fig. 5A). In *Styela*, *Itnl1/Fcn1* mRNAs were enriched

in clusters 8 and 3 and zones 6 and 7 of the endostyle (Jiang et al., 2023). *Muc5a* and *VWF*

transcripts were found in Clusters 3 and 10(Sasaki et al., 2003; Yamagishi et al., 2022).

Glutathione peroxidase, a common endostyle and branchial sac enzyme (Kobayashi et al.,

1983), was observed in our dataset in Clusters 3 and 6 (Fig. 5A).

GO and pathway analyses for Cluster 3 associated genes revealed enrichment for processes

such as immunity, ECM, carbohydrate binding, and cell adhesion (Fig 5B). Jiang et al. (2023)

also found a list of similar GO terms including ribosomal, thyroid hormone, immune function,

digestive function (such as mucus), and neurosecretory genes (such as semaphorin 1a) for

*Styela clava* (Jiang et al., 2023).

**Figure 5. Identification of Endostyle Clusters**. A. UMAP feature density plots showing the highest
concentration of cells expressing each of these tunicate endostyle genes. B. GO and pathway terms colored by p-
value as determined by Metascape. C. UMAP plot for cluster marker 3, *Fbn1*. D. *In situ* expression pattern for
Cluster 3 marker *Fbn1*, in zone 8 of the endostyle. E. *Tubb* mRNA was detected in zone 1 cells of the endostyle.
F. Schematic showing the different zones of the endostyle, highlighting the spatial expression of *Fbn1* and *Tubb*.
Abbreviations: zone 8 (z8), zone 1 (z1), stigmata (st), ventral (v), dorsal (d). Scale bar is 50 μm.
223

The top marker gene for Cluster 3, g09775, (Fig. 5C) encodes fibrillin (Fbn1), a protein with

fibrillin repeats and an EGF domain that shares 44% identity with the Human Fbn1/2/3

proteins. Fbn proteins are found in connective tissues and maintain the elasticity of the tissues.

It is critical for microfibril synthesis, as it binds to calcium and regulates TGF-β release,

growth, and development (Chaudhry et al., 2007; Handford, 2000; Reinhardt et al., 1996). *Fbn1*

expression was observed in zone 8 of the endostyle (Fig. 5D). This region of the endostyle

functions with zone 7 in immune activities, with high iodine and peroxidase activities (Jiang et

al., 2023; Sasaki et al., 2003) (Alesci et al., 2022). Iodine metabolism has been linked to thyroid

gland evolution (Fujita and Nanba, 1971). We also found that the *Tubb* probe (identified as a

ciliated cell marker in Fig. 3D) marked a subset of endostyle cells. *In situ* hybridization showed

that its mRNA was present in zone 1 cells of the endostyle (Fig. 5E). These cells have long

cilia, which together with mucus, aid in trapping food particles from the water current (Holley,

1986). Based on this information, Cluster 3 was assigned as an endostyle cell cluster.

**Identification of Blood cell clusters**

The top Cluster 25 gene g07537 is predicted to encode a FAD-dependent oxidoreductase

domain-containing protein (Foxred2) (Didion et al., 2002). This transcript was also detected in

some of the cluster 12 cells (Fig. 6A). Foxred2 functions to balance redox states and contributes

to the generation of reactive oxygen species (ROS), which are associated with endoplasmic

reticulum (ER) stress. Strong Foxred2 staining was observed in the thin endothelium lining

244 blood vessels and in a small number of immunocytes with a few large granules (Fig. 6A).

245 Additionally, staining was observed in the developing heart tube cells as they became thinner.

246 In ascidians, the expression of Foxred2-like proteins in the vascular lining and granular

247 immunocytes suggests that it may be crucial for maintaining redox balance and metabolic

248 processes critical for vascular function and immune response. This may include protecting cells

249 from oxidative stress, aiding in antimicrobial defence, detoxifying harmful substances, and

250 regulating inflammation. GO analysis (Fig. S10) revealed associations with heme binding,

251 biosynthesis, oxidoreductase activity, and SLC-mediated transmembrane transport within

252 Cluster 25. SLC-mediated transport is important for thin-monolayer barriers, such as the blood-

253 brain barrier, to mediate the transport of substances across the endothelium (Morris et al.,

254 2017).

256 The expression of the Cluster 13 marker, Glutathione S-Transferase Alpha 3 (*Gst3A*) *g11109,*

257 was also detected in some cells and distributed throughout the other clusters (Fig 6B).

258 Staining was observed in cytotoxic cells, including granular amoebocytes (Fig.6B, asterisks),

259 compartment amoebocytes, and morula-like cells (Fig. 6B). Morula cells are variable in size,

260 and the extent of staining may have been influenced by vacuole size. Only a few significantly

261 enriched GO terms were identified because of the low number of marker genes in this cluster.

262 However, it included immune system processes (Fig. S10).

**A** Cluster 25

*g07537 Foxred2*

**B** Cluster 13

*g11109 Gst3a*

**C** Cluster 17

*g13310 Csmd3-like*

**D** Cluster 8

*g05125 Notch-like*

**Figure 6. Candidate blood cell clusters.** A. UMAP feature plot for Cluster 25 marker, *Foxred2*. Detection of Foxred2 mRNA by in situ hybridisation identified vascular endothelium (blue arrow), cells of the developing heart and some blood cells (asterisk). B. Cluster 13, g11109 *Gst3a*. C. Cluster 17 marker, g13310 (*Csmd3*). *In situ* hybridization with *Csmd3,* with mRNA-positive vesicles indicated (asterisks). D. Cluster 8 marker *g05125* (*Notch-like*). Only a few scattered cells in the vascular stained with the *g05125* probe (arrows). Abbreviations: stomach (stm), vessel endothelium (et), blood vessel (bv), endostyle (end), compartment amoebocyte (ca), compartment cell (cc), zone 2 (z2), zone 4 (z4) The scale bars are 10 μm.

The gene g13310 was identified as a cell marker for Cluster 7 (Fig. 6C). This gene is predicted to encode a protein ortholog of CSMD3 with multiple Sushi and von Willebrand factor type A (CUB) domains, typically found in transmembrane receptors or adhesion proteins. Sushi or complement control domains are involved in the immune system (Ermis Akyuz and Bell, 2022). Excessive activation of the complement pathway is prevented by Sushi domain-containing proteins, which bind to activated C3/4 components to target them for degradation (Ojha et al., 2019). *Csmd3+* cells were detected in the circulation (Fig. 6C), which appeared to be phagocytic cells, including macrophage-like cells and hyaline amoebocytes. Gene ontology analysis revealed an overrepresentation of terms related to the cytoskeleton, apoptosis, and immunity (Fig. S9).

Cluster 8 top marker gene, *g05125* (Fig. 6D), codes for a large protein that shares 30% identity with NOTCH1, NOTCH2, and SNED1 due to multiple Sushi and EGF domains. Several Notch-like genes are present in the *B. diegensis* genome. *In situ* hybridization detected expression in a small number of storage and/or granular cells; larger cells were characterized by multiple small vesicles (Fig. 6D). These cells resemble mast cells and are predicted to release inflammatory factors such as histamine and chemokines (Blanchoud et al., 2017; Cima et al., 2001). Gene Ontology and pathway analysis indicated terms related to peroxidase and leukocyte-mediated immunity (Fig. S11).

**Developmental Trajectory analysis**

To identify the cells forming the bud disc of the peribranchial epithelium, we examined cells

with thickened epithelium (Berrill, 1947b) and budlet/vesicle precursors. We focused on

identifying the cells with the highest expression of transcription factors found in disc cells,

*Pitx1, Otx, Nk4*, and *Runx* (Langenbacher et al., 2015; Ricci et al., 2016; Tiozzo et al., 2005).

The subset of cells within Cluster 6, likely the peribranchial cluster, showed the highest joint

density, indicating that cells co-expressed these four genes (Fig. 7C). Using this as the root, the

predicted cell trajectory was plotted using the Monocle3. Pseudotime trajectory paths indicate

the progression and pathways of these cells, going from this putative stem cell cluster to more

differentiated states later in the pseudotime. This analysis connected all Seurat clusters, with

the origin located in Cluster 6 (Fig. 7C). To extend this further, we determined whether several

previously studied candidate stem cell markers were present in the scRNA-seq dataset,

including *Itga6*, *Notch2*, *Vasa*, and *Piwi* (Kassmer et al., 2020; Kawamura and Sunanaga, 2010;

Rinkevich et al., 2010; Rosental et al., 2018). *Vasa, Piwi1*, and *Piwi2* were missing from the

dataset. *Notch2* and *Itga6* were detected in the cells scattered across several clusters (Fig. S12).

The proportion of dense cells was the same as that of bud disc genes (Fig. S12 and Fig. 7).

Additionally, orthologs of the reprogramming factors (Yamanaka factors) (Takahashi and

Yamanaka, 2006, 2016), *Oct3/4 (*POU3/4 proteins), *Sox2* (SOXB subgroup of Sox factors),

*Klf4*, and *cMyc* were found in the same group of cells within cluster 6 (Fig. S13). This further

supports the notion that a subset of Cluster 6 cells likely has stem cell properties and represents

the bud disc.

**Figure 7. Cell populations and their relationships.** A. Summary of the main tissue and organs formed. B.
Example schema showing the relationships between tissues. The bud arises from a small group of cells (3-4 cells)
within the epithelium of the peribranchial chambers, which appear in the immature zooid before the stigmata
perforate. These cells form the bud disc. (Berrill, 1947). The disc separates from the epithelium to form a closed
vesicle surrounded by the outer epidermis obtained from the parent zooid. Germ cells segregate early from the
inner vesicle and form oocytes and sperm. On the dorsal side of the vesicle, the invagination forms a neural
placode. The heart and digestive tracks form on the posterior side, and primordia form first as outpockets of the
branchial epithelium. C. UMAP density plot showing co-expression of *Pitx, Otx, Nk4*, and *Runx* in cells located
in Cluster 6. Cells with higher joint densities were more orange in color. D. Monocle 3 trajectory analysis. The
root is the progenitor (within cluster 6, likely the peribranchial epithelium). Abbreviations: endostyle (en),
stigmata (st), peribranchial chamber (pbc), branchial chamber (bc), intestine (in), stomach (sm), testes (tt), ovary
(o), bud disc (bd).
329

The trajectory of the colony dataset was analyzed to understand the relationship between cell

clusters and differentiation time using Monocle3 (Fig. 7D). The trajectory of cell development

was charted on UMAP, with the candidate progenitor bud disc cells of the peribranchial cluster

(Cluster 6) serving as the root (Fig. 7C). Pseudotime trajectories reveal that the clusters,

predicted to be bud, peribranchial and brachial epithelial cells, emerge early. In contrast,

clusters associated with specialized tissues, such as the stomach epithelium and neuronal

clusters, were later in the pseudotime (Fig. 7D). Epithelial tissues within the zooid appeared

early in the trajectory, preceding the development of stomach and endostyle tissues (left

branch). The third pathway leads to immune cell formation. Overall, the development of zooid-

related tissues later in the trajectory was consistent with blastogenesis (Fig. 7A and B)

**Figure 8. Identification of potential differentiation pathways using CellRank.** A. The PAGA plot displays each cell cluster as a node, with branching lines indicating potential differentiation pathways among the clusters. B. A tSNE projection where cells are colored by pseudotime, showing the progression of cell states over time. Darker colors represent earlier pseudotime states, whereas lighter colors represent later pseudotime states. The Cluster IDs are given for the later states. A UMAP projeciont is shown in Fig. S15. C and D. Velocity plots using the tSNE projection, where arrows represent the direction of RNA velocity, indicating the predicted future states of the cells based on their transcriptional activity and pseudotime. The cells are colored using the Seurat Cluster ID. D. An annotated version of plot C shows detailed identification of specific cell types and their predicted differentiation pathways.

To explore the relationships between clusters, we generated PAGA plots to investigate the underlying structure and infer the potential differentiation pathways (Fig. 8A). This suggests that multiple common progenitor populations may give rise to differentiated cell types. The relationships between different clusters of cells during colony asexual reproduction were visualized, with each node representing a distinct cluster identity. Lines connecting the nodes illustrate lineage relationships between clusters. By examining central nodes with numerous connections, such as nodes 20, 21, 0, and 6, we can identify potential progenitor states that can

differentiate into several other states. These clusters had numerous outgoing edges, indicating

potential branching points in the differentiation trajectories (Fig. 8A). Finally, the cell clusters

at the ends of the branches represent the terminally differentiated states, and this largely

matches the later cell Clusters identified in pseudotime analyses (Fig. 8C and 7D).

The velocity (scvelo) plot utilizes the spliced-to-unspliced mRNA ratio to indicate the direction

of the cell transition. The length of the arrows in the velocity plot represents the speed of the

transitions, with longer arrows indicating faster transitions and shorter arrows indicating slower

or more stable transitions. This provides dynamic information about cellular processes.

Additionally, the arrows converged at specific points, offering further insights into the

directionality and progression of cell states. Clusters with numerous outgoing arrows are

highlighted, emphasizing their roles in cellular differentiation (Fig. 8C). This plot was

consistent with the PAGA plot, particularly with multiple transition events from clusters 0, 6,

21, and 20 (Fig. 8A), suggesting that cells within these clusters serve as central nodes with

multiple connections (Fig. 8D).

**Fig. 9. Cell fate dynamics.** A. UMAP plot showing diffusion pseudotime (dpt_pseudotime). The color scale

represents pseudotime, cells colored yellow appear later in the predicted cell trajectory pathway. B. Fate

probability UMAP, shown as terminal clusters. C. Aggregate fate probabilities fate bar graphs. These display the probabilities of different clusters (22, 11, 15, 19, and 6) progressing towards the terminal states, as predicted by CellRank. D. A selection of driver genes is displayed alongside a gene whose expression is highest at the earliest pseudotime on the lineage towards the terminal state fate (File S4).

Finally, CellRank was used to examine the cell fate dynamics (Fig. 9). It has been designed to deal with complex data, such as data produced from developing and regeneration tissues, with many initial, intermediate, and terminal cell states (Lange et al., 2022). Figures 9C and S14 display a bar graph of the aggregate fate probabilities of cells from the Seurat cluster. These cells will progress to a terminal state, and the clusters with multiple bars of varying height have different probabilities of transitioning to different end states, such as Clusters 0, 5, 6, 8, 20, 18, 21, 22, 26, and 28. In contrast, clusters with cells that transition to one fate predominantly have a single high bar. For example, all cells in Cluster 11 transitioned to the ciliated stigmata cell type, according to GO and *in situ* expression (Fig. 3C).

We used CellRank to identify potential driver genes strongly associated with the terminal state that may serve as key regulators or markers of cell fate (Fig. 9E and File 4). These genes exhibited high expression levels at the earliest pseudotime but were low in the final terminal state. Such genes may play a role in initiating or driving the early stages of differentiation and then decreasing their expression as cells reach their final differentiated state. One example of a potential driver gene is the *Lin2*8 ortholog, a highly conserved RNA-binding protein that regulates stem cell differentiation and proliferation in organisms ranging from nematodes to mammals (Wu et al., 2022). This gene is expressed in undifferentiated pluripotent cells and is downregulated as cells progress toward a specialized cell fate or undergo reprogramming (Pieknell et al., 2022). Another potential driver gene was *the Atf4/*5 ortholog. In vertebrates, this transcription factor is associated with stress-induced responses and the differentiation of stem cells into secretory cells. It plays a role in epithelial differentiation and stresses immune

response (Barrera-Lopez et al. 2024). *B. diegensis g11626* encodes a DExD/H-box polypeptide,

39 B (DDx39b)-like RNA helicase. DDx39b regulates mRNA splicing events, including the

processing of transcripts of genes required for immune and myocyte cell fate (Hirano et al.,

2023; Zhang et al., 2018). Another potential driver gene is high mobility group box 1 (*Hmgb1*),

a chromatin-associated protein secreted by immune cells and involved in differentiating

mesenchymal stem cells into vascular cells (Meng et al., 2018).

Together, these analyses provided strong supporting evidence for the inferred differentiation

pathways. They identified key transitional cell clusters and several candidate progenitor cell

populations.

**Discussion**

Using sc-RNA-seq, we gained insights into the cellular content of a *B. diegensis* colony. This

technology identifies cluster markers by deconstructing the cellular heterogeneity of the filter-

feeding colony tissues. The colonial tissue was resolved into 29 clusters, including zooid and

vascular tissues. As a more ancestral chordate, *B. diegensis* occupies an essential phylogenetic

space with vertebrates, highlighting its significance in evolutionary and developmental

research. This sc-RNAseq resource paves the way for examining the cellular composition of

the colonial chordate, *B. diegensis.*

Based on the marker expression profiles that were highly and distinctively expressed in each

cluster, the colonial cellular profiles showed conserved gene regulatory programs and

functions. In contrast, clusters not enriched for a particular group of markers or functional

annotations may indicate an intermediate cell state. Although we found some blood cell

markers, some predicted hemocyte clusters showed intermediate and non-specific gene

expression patterns. These functions were not specifically enriched for the blood type because

of functional annotations based on the closest vertebrate relatives. Similarly, some blood

populations of *B. schlosseri* did not exhibit specific functions and contained unannotated genes

with invertebrate homologues (Rosental et al., 2018).

Trajectory analysis of cell fate during blastogenesis suggested that multiple initial and

intermediate states are present in the colony at any given time. In ascidians, bud development

is initiated by the emergence of bud primordium through the thickening of the peribranchial

epithelium via dedifferentiation or transdifferentiation (Manni et al., 2007). The peribranchial

epithelium corresponded to Cluster 6 in our study. Our gene expression analysis in Cluster 6

supports the hypothesis that the peribranchial epithelium is a critical source of progenitor cells

for bud development. The endostyle has been proposed as a stem cell niche in zooids,

suggesting a more transitional stem cell source for the colony (Voskoboynik et al., 2008a).

Specialized progenitors associated with the vascular lining and epithelial tissues have also been

identified. This demonstrates the ability of epithelial tissues to function as a source of

progenitors, preserving progenitor-like characteristics. This is significant because it indicates

that *Botrylloides* stem cells may not be fixed in one location, but can transition between

different niches within the organism. Additionally, the presence of transitional stem cell niches,

such as endostyle, highlights the dynamic nature of stem cell populations in these organisms.

Adult stem cells (ASCs) are involved in various epithelial tissues in invertebrates. For example,

in sponges, epithelial cells function similarly to stem cells and play a role in tissue repair

(Rinkevich et al., 2022). This observation suggests a conserved mechanism across different

invertebrate species where epithelial cells have multipotent capabilities essential for

regeneration and repair. This comparison underscores the importance of epithelial and vascular

progenitors in colonial ascidians and other invertebrates, indicating a potentially conserved

evolutionary strategy for tissue regeneration and maintenance. We hypothesized that epithelial-

and vascular-associated progenitors play key roles in the development and maintenance of

colonial ascidians.

Assigning cellular identities in scRNA-seq is inherently challenging because of the complexity

and variability of the gene expression profiles. The study was based on a limited set of

approximately 6,000 cells, which may not capture the full diversity of cell types in the

organism. Low-abundance transcripts may be underrepresented, leading to a potential bias in

gene expression data and overlooking genes expressed at low levels. Obtaining cells from the

tunic, which is rich in cellulose, presents a significant challenge, leading to the absence of

certain tunic cell types. The timing of colony collection might have missed some stages of the

sexual cycle, potentially omitting important cell types or developmental stages. For example,

the stages of oogenesis will be missed as *B. diegensis* only sexually reproduces briefly at the

end of the summer in New Zealand.

*In situ* hybridization, which is useful for the spatial localization of gene expression, is difficult

to quantify accurately, limiting its utility in confirming single-cell RNA sequencing (scRNA-

seq) findings. No one cluster was identified as marked by gene expression commonly attributed

to stem-like cells in colonial ascidians (Ballarin and Rosner, 2022). Specifically, previously

identified germline markers for colonial and solitary tunicates, such as *Piwi* or *Vasa*

(Kawamura et al., 2011; Rinkevich et al., 2010), were missing from the dataset. This absence

could be due to technical dropout, a common issue in scRNA-seq where lowly expressed genes

might not be detected in every cell, leading to their apparent absence in the data (Kharchenko

et al., 2014). The genes in *B. diegensis* were named and assigned functions for gene ontology

analysis based on their closest vertebrate orthologs, assuming similar functions in ascidians.

However, many genes lack orthologs, making functional prediction more challenging.

**SUMMARY**

In this study, we performed single-cell RNA sequencing (scRNA-seq) on an entire *B. diegensis*

colony, including zooids, buds, and the vascular tunic, to identify cell and tissue markers and

resolve cellular heterogeneity. The analysis identified 29 major cell clusters within the colony,

and *in situ* hybridization was used to examine the spatial expression of the cluster marker genes.

Various tissue types have been identified at the molecular level, including blood cells and zooid

tissues such as the branchial epithelium, stomach, and endostyle. Distinct cluster markers were

found in specific regions of the stomach epithelium, highlighting their specialization.

Trajectory estimations were aligned with predictions based on the early appearance of

progenitor clusters, whereas zooid-related tissues appeared later in the developmental path.

This study provides a valuable resource for understanding the development, function, and

regeneration of *B. diegensis*, and demonstrates the power of scRNA-seq in defining cell types

and tissues in complex colonial organisms.

**Methods**

**Animal husbandry**

Colonies of *B. diegensis* were sampled from Otago Harbor in New Zealand (45°52'18.1″''S,

170°31'37.6''″ E). The animals were attached to 5x7 cm glass slides. The tanks were aerated,

and the colonies were fed a shellfish diet (a blend of marine microalgae). The colonies were

fed every two days, and the seawater was replaced. Colonies were confirmed to be *B. diegensis*

by COI barcoding (Temiz et al., 2023).

**Single-cell preparation & FACS**

Single cells of *B. diegensis* were prepared using the ACME dissociation method described by García-Castro et al. (2021). Fresh ACME solution was prepared for fixation and separation of tissue cells. It contained DNase/RNase-free distilled water, methanol, glacial acetic acid, and glycerol at a ratio of 13:3:2:2. The mature colony was in stage A of blastogenesis with active filtering, newly emerged primary buds, and no signs of secondary buds. The colony was placed in a Petri dish on a glass slide, where it was attached using a microtome razor blade with minimal disturbance. The animal was washed with 1 ml of 7.5% N-acetyl l-cysteine in 1XPBS. This solution removes excess seawater and protects RNA. One mL ACME was added, and the colony was minced well using a single-edged razor blade.

The suspension and all the larger tissue pieces were placed in a 1.5 mL tube. The tube was placed in a rotator to apply seesaw motion at ~30-40 rpm for 1 h at room temperature. The sample was pipetted twice and strained with a 40 μm cell strainer into an ice-covered 50 mL Falcon tube. The cells were cooled to prevent RNA degradation. The suspension was centrifuged at 1000 $g$ for 5 min at 4°C. The supernatant was removed, and the pellet was apparent. One mL of 1% BSA-1XPBS with RNase inhibitor was added to the pellet, and the tube was flicked to mix. As the second washing step to remove all ACME, 1 mL of 1% BSA-1XPBS with RNase inhibitor was added to the tube, and the tubes were flicked. A 70 μm flowmi cell strainer was used to strain the cells to decrease the number of aggregates. The sample was centrifuged at 1500 $\times$ $g$ for 5 min at 4°C, and the cell pellet was visible.

The cells were checked on a hemocytometer by staining with trypan blue (1:1). The approximate number of cells was estimated, and their integrity was investigated. ACME fixed cells were stained with the DNA dye, DRAQ5™, to sort intact single cells from cellular debris

and aggregates (eBioscience 0.66 μL/mL of 5 mM stock). After staining, cells were incubated

in the dark and on ice for an hour. The stained cells were sorted using a BD FACSAria Fusion

flow cytometer (BD Biosciences) with a red laser (640 nm). In total, 50,000 cells were sorted

in collection buffer containing 1X PBS-1% BSA-RNAse inhibitor (40 U/mL) (Fig. S1). The

cells were visualized under the far-red channel of a Nikon Ti2 Inverted fluorescence

microscope (Fig. S1). The cells were preserved at -20°C by adding DMSO (10% final

concentration).

**sc-RNA-seq via 10X Genomics & sequencing**

Frozen cells were thawed on ice and centrifuged at 1500 × g and 4°C. The supernatant was

used to wash away the DMSO, followed by adding 1XPBS-1%BSA-RNase inhibitor (40

U/mL). Sorted cells were counted on a hemocytometer using trypan blue in a 1:1 ratio and

diluted to yield approximately 6000 cells for further processing. The cells underwent

centrifugation at 1500 × g at 4°C, then 10X Master Mix was added to the single cells.

Subsequently, the cells were loaded onto a 10x Genomics Chromium chip. Gel bead emulsion

(GEM) generation, barcoding, reverse transcription, cDNA amplification, and library

preparation were carried out according to the 10X Genomics protocol for 3' Gene Expression

(v3) user guide (10X Genomics CG000183 Rev C). The target cell recovery rate was set to

3000. After the GEM generation, a post-GEM-RT clean-up procedure was performed. cDNA

amplification was performed for 16 cycles. The resulting cDNA was analyzed using a Qubit

fluorometer and agarose gel electrophoresis. Samples were indexed for library preparation and

library concentrations were quantified using KAPA PCR. The combined cDNA libraries were

sequenced with a length of 150 bp on an Illumina NovaSeq 6000 sequencing platform at the

Australian Genome Research Facility Ltd.

**Read alignment & cluster analysis**

Quality checks of the library were performed using FastQC (Andrews, 2010) (Supplementary Fig. S22). Raw sequencing reads were mapped to the reference *B. diegensis* genome using STARSolo (Kaminow et al., 2021). The genome file of *B. diegensis* (formerly *B. leachii*) was downloaded from the Aniseed Database (http://www.aniseed.cnrs.fr) (Blanchoud et al., 2018). Gene models were identified using the StringTie software (Pertea et al., 2016). First, genome indices were created using STAR 2.7.9 (--genome SAindexNbases 12)(Dobin et al., 2013). The indices were then used to map the raw sequences to the *B. diegensis* genome using STAR 2.7.9, which masks the polyA tail during alignment; therefore, no prior trimming was performed. These run options were selected for the default barcode lengths using a droplet-type algorithm (soloUMIlen 12, soloType Droplet). No barcode read length defined (-- soloBarcodeReadLength 0). The empty droplets were then filtered (soloCellFilter EmptyDrops_CR). Finally, sequences with barcodes present within the barcode whitelist were selected while mapping (soloCBwhitelist). The mapping statistics are listed in Table S1.

After mapping, clustering was performed using Seurat 4.0.1(Stuart et al., 2019) and R 4.1.3. Before clustering, filtering was performed by selecting transcriptomes with 200–2000 genes expressed in at least three cells. The counts were normalized to the total counts using log normalization, and the scale factor was set to 10000. Variability was identified within the 2000 genes using the *FindVariableFeatures* function with vst as the selection method. The linear dimensional reduction method was applied to the single-cell transcriptome using principal component analysis. The first 50 PCs for the mature colony were selected based on elbow and jackstraw analyses. Clustering was executed with a 0.8 resolution determined using the clustree R package (Zappia and Oshlack, 2018). The data were plotted using the nonlinear dimensional

reduction method and presented using the Uniform Manifold Approximation and Projection

(UMAP).

Differential expression analysis was performed using Seurat package to identify the top genes

that were potential markers for each cluster. Top genes were considered significant if Padj <

0.05 and had a $\log_2$FC (fold change) value of > 0.5 (Supplementary File 1). Single-cell

transcriptomes were saved in the Seurat object, which can be used to check for feature

expression. Finally, gene annotation was performed using Metascape (Zhou et al., 2019). The

pathway and GO term enrichment were calculated by comparing them to the background genes

(all the expressed genes in the dataset with an orthologue, 6039 genes).

Pseudo-time estimations of single cells were calculated using Monocle 3 (Trapnell et al., 2014),

Scanpy 1.10.1 (Wolf et al., 2018), scVelo 0.3.2 (Bergen et al., 2020) and CellRank v. 2.0 (Lange

et al., 2022). Code availability: https://github.com/MJWilsonOtago/scRNAseqBotrylloides/ .

**Probe synthesis, *in situ* hybridization & image acquisition**

Probe preparation was initiated by cloning the target genes. Primers were designed to amplify

the genes of interest (Table S2). The amplified fragments were inserted into pCRII-TOPO (Life

Technologies). Plasmids were sequenced to verify the correct fragment. Digoxygenin (DIG)-

labelled sense and antisense probes were synthesized using a 10× DIG RNA labelling mix and

SP6/T7 RNA polymerases (Sigma-Aldrich) via *in vitro* transcription. In situ hybridisation was

performed as described previously (Zondag et al., 2019). Images were acquired using a Nikon

TiE with 60x magnification.

**Data availability**

The authors declare that all data supporting this study's findings are available in the article and its supplementary information files or through https://megan-wilson-otago.shinyapps.io/Botryllodies_scRNAseq/ .

**Acknowledgements**

The authors would like to thank the Gemmell Lab of the University of Otago, particularly Joanne Gillum, for their contribution to the single-cell optimization. We also thank Bridget Fellows, Devon Gamble, Justine Gapuz, and Ed Moody for their comments on the final draft. This study was supported by funding from the Department of Anatomy and a University of Otago Research Grant. B.T. was supported by an Anatomy Department Ph.D. scholarship from the University of Otago.

**Contributions**

B.T.: Data acquisition and interpretation, and manuscript writing. M.M.: funding acquisition, data acquisition, interpretation, and manuscript writing. M.J.W.: Study design, data acquisition and interpretation, funding acquisition, and manuscript writing.

**Competing interests**

The authors declare no competing interests.

**References**

Alesci, A., Pergolizzi, S., Lo Cascio, P., Capillo, G., Lauriano, E.R., 2022. Localization of vasoactive intestinal peptide and toll-like receptor 2 immunoreactive cells in endostyle of urochordate Styela plicata (Lesueur, 1823). Microsc Res Tech 85, 2651-2658.

Andrews, S., 2010. FastQC: a quality control tool for high throughput sequence data. Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom.

Anselmi, C., Kowarsky, M., Gasparini, F., Caicci, F., Ishizuka, K.J., Palmeri, K.J., Raveh, T., Sinha, R., Neff, N., Quake, S.R., Weissman, I.L., Voskoboynik, A., Manni, L., 2022. Two distinct

evolutionary conserved neural degeneration pathways characterized in a colonial chordate. Proceedings of the National Academy of Sciences 119, e2203032119.

Bergen, V., Lange, M., Peidli, S., Wolf, F.A., Theis, F.J., 2020. Generalizing RNA velocity to transient cell states through dynamical modeling. Nature biotechnology 38, 1408-1414.

Berrill, N.J., 1947a. The developmental cycle of *Botrylloides*. Q J Microsc Sci. 3, 393-407.

Berrill, N.J., 1947b. The Developmental Cycle of Botrylloides. Q J Microsc Sci 88, 393-407.

Berrill, N.J., 1951. Regeneration and budding in tunicates. Biol Rev 26, 456-475.

Blanchoud, S., Rutherford, K., Zondag, L., Gemmell, N.J., Wilson, M.J., 2018. De novo draft assembly of the Botrylloides leachii genome provides further insight into tunicate evolution. Sci Rep 8, 5518.

Blanchoud, S., Zondag, L., Lamare, M.D., Wilson, M.J., 2017. Hematological Analysis of the Ascidian Botrylloides leachii (Savigny, 1816) During Whole-Body Regeneration. The Biological bulletin 232, 143-157.

Cao, C., Lemaire, L.A., Wang, W., Yoon, P.H., Choi, Y.A., Parsons, L.R., Matese, J.C., Levine, M., Chen, K., 2019. Comprehensive single-cell transcriptome lineages of a proto-vertebrate. Nature 571, 349-354.

Chaudhry, S.S., Cain, S.A., Morgan, A., Dallas, S.L., Shuttleworth, C.A., Kielty, C.M., 2007. Fibrillin-1 regulates the bioavailability of TGFβ1. The Journal of cell biology 176, 355-367.

Cima, F., Perin, A., Burighel, P., Ballarin, L., 2001. Morpho-functional characterization of haemocytes of the compound ascidian Botrylloides leachi (Tunicata, Ascidiacea). Acta Zoologica 82, 261-274.

Didion, S.P., Ryan, M.J., Didion, L.A., Fegan, P.E., Sigmund, C.D., Faraci, F.M., 2002. Increased superoxide and vascular dysfunction in CuZnSOD-deficient mice. Circ Res 91, 938-944.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., Gingeras, T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15-21.

Ermis Akyuz, E., Bell, S.M., 2022. The Diverse Role of CUB and Sushi Multiple Domains 1 (CSMD1) in Human Diseases. Genes (Basel) 13.

Fujita, H., Nanba, H., 1971. Fine structure and its functional properties of the endostyle of ascidians, Ciona intestinalis. A part of phylogenetic studies of the thyroid gland. Z Zellforsch Mikrosk Anat 121, 455-469.

García-Castro, H., Kenny, N.J., Iglesias, M., Álvarez-Campos, P., Mason, V., Elek, A., Schönauer, A., Sleight, V.A., Neiro, J., Aboobaker, A., 2021. ACME dissociation: a versatile cell fixation-dissociation method for single-cell transcriptomics. Genome biology 22, 1-34.

Garcia-Castro, H., Kenny, N.J., Iglesias, M., Alvarez-Campos, P., Mason, V., Elek, A., Schonauer, A., Sleight, V.A., Neiro, J., Aboobaker, A., Permanyer, J., Irimia, M., Sebe-Pedros, A., Solana, J., 2021. ACME dissociation: a versatile cell fixation-dissociation method for single-cell transcriptomics. Genome biology 22, 89.

Handford, P., 2000. Fibrillin-1, a calcium binding protein of extracellular matrix. Biochimica et Biophysica Acta (BBA)-Molecular Cell Research 1498, 84-90.

Hirano, M., Galarza-Munoz, G., Nagasawa, C., Schott, G., Wang, L., Antonia, A.L., Jain, V., Yu, X., Widen, S.G., Briggs, F.B.S., Gregory, S.G., Ko, D.C., Fagg, W.S., Bradrick, S., Garcia-Blanco, M.A., 2023. The RNA helicase DDX39B activates FOXP3 RNA splicing to control T regulatory cell fate. eLife 12.

Hirose, E., Shirae, M., Saito, Y., 2003. Ultrastructures and classification of circulating hemocytes in 9 botryllid ascidians (Chordata: Ascidiacea). Zoological science 20, 647-656.

Holland, L.Z., 2016. Tunicates. Current Biology 26, R146-R152.

Holley, M.C., 1986. Cell shape, spatial patterns of cilia, and mucus-net construction in the
ascidian endostyle. Tissue Cell 18, 667-684.

Horie, R., Hazbun, A., Chen, K., Cao, C., Levine, M., Horie, T., 2018. Shared evolutionary origin
of vertebrate neural crest and cranial placodes. Nature 560, 228-232.

Jiang, A., Zhang, W., Wei, J., Liu, P., Dong, B., 2023. Transcriptional Analysis of the Endostyle
Reveals Pharyngeal Organ Functions in Ascidian. Biology (Basel) 12.

Kaminow, B., Yunusov, D., Dobin, A., 2021. STARsolo: accurate, fast and versatile
mapping/quantification of single-cell and single-nucleus RNA-seq data. bioRxiv.

Kassmer, S.H., Langenbacher, A.D., De Tomaso, A.W., 2020. Integrin-alpha-6+ Candidate stem
cells are responsible for whole body regeneration in the invertebrate chordate Botrylloides
diegensis. Nature communications 11, 4435.

Kawamura, K., Sunanaga, T., 2010. Hemoblasts in colonial tunicates: Are they stem cells or
tissue-restricted progenitor cells? Development, growth & differentiation 52, 69-76.

Kawamura, K., Tiozzo, S., Manni, L., Sunanaga, T., Burighel, P., De Tomaso, A.W., 2011. Germline
cell formation and gonad regeneration in solitary and colonial ascidians. Developmental
Dynamics 240, 299-308.

Kharchenko, P.V., Silberstein, L., Scadden, D.T., 2014. Bayesian approach to single-cell
differential expression analysis. Nat Methods 11, 740-742.

Kobayashi, H., Tsuneki, K., Akiyoshi, H., Kobayashi, Y., Nozaki, M., Ouji, M., 1983. Histochemical
distribution of peroxidase in ascidians with special reference to the endostyle and the
branchial sac. Gen Comp Endocrinol 50, 172-187.

Lange, M., Bergen, V., Klein, M., Setty, M., Reuter, B., Bakhti, M., Lickert, H., Ansari, M.,
Schniering, J., Schiller, H.B., Pe'er, D., Theis, F.J., 2022. CellRank for directed single-cell fate
mapping. Nat Methods 19, 159-170.

Langenbacher, A.D., Rodriguez, D., Di Maio, A., De Tomaso, A.W., 2015. Whole-mount
fluorescent in situ hybridization staining of the colonial tunicate Botryllus schlosseri. Genesis
53, 194-201.

Manni, L., Burighel, P., 2006. Common and divergent pathways in alternative developmental
processes of ascidians. BioEssays : news and reviews in molecular, cellular and developmental
biology 28, 902-912.

Manni, L., Zaniolo, G., Cima, F., Burighel, P., Ballarin, L., 2007. Botryllus schlosseri: a model
ascidian for the study of asexual reproduction. Developmental dynamics: an official
publication of the American Association of Anatomists 236, 335-352.

Meng, X., Chen, M., Su, W., Tao, X., Sun, M., Zou, X., Ying, R., Wei, W., Wang, B., 2018. The
differentiation of mesenchymal stem cells to vascular cells regulated by the HMGB1/RAGE
axis: its application in cell therapy for transplant arteriosclerosis. Stem Cell Res Ther 9, 85.

Morris, M.E., Rodriguez-Cruz, V., Felmlee, M.A., 2017. SLC and ABC Transporters: Expression,
Localization, and Species Differences at the Blood-Brain and the Blood-Cerebrospinal Fluid
Barriers. AAPS J 19, 1317-1331.

Mukai, H., Sugimoto, K., Taneda, Y., 1978. Comparative studies on the circulatory system of
the compound ascidians, *Botryllus, Botrylloides* and *Symplegma*. Journal of morphology 157,
49-77.

Ojha, H., Ghosh, P., Singh Panwar, H., Shende, R., Gondane, A., Mande, S.C., Sahu, A., 2019.
Spatially conserved motifs in complement control protein domains determine functionality in
regulators of complement activation-family proteins. Commun Biol 2, 290.

Parrinello, D., Sanfratello, M.A., Vizzini, A., Parrinello, N., Cammarata, M., 2015. Ciona intestinalis galectin (CiLgals-a and CiLgals-b) genes are differentially expressed in endostyle zones and challenged by LPS. Fish Shellfish Immunol 42, 171-176.

Parrinello, D., Sanfratello, M.A., Vizzini, A., Testasecca, L., Parrinello, N., Cammarata, M., 2017. The Ciona intestinalis immune-related galectin genes (CiLgals-a and CiLgals-b) are expressed by the gastric epithelium. Fish Shellfish Immunol 62, 24-30.

Perillo, M., Wang, Y.J., Leach, S.D., Arnone, M.I., 2016. A pancreatic exocrine-like cell regulatory circuit operating in the upper stomach of the sea urchin *Strongylocentrotus purpuratus* larva. BMC evolutionary biology 16, 1-15.

Pertea, M., Kim, D., Pertea, G.M., Leek, J.T., Salzberg, S.L., 2016. Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nature protocols 11, 1650-1667.

Pieknell, K., Sulistio, Y.A., Wulansari, N., Darsono, W.H.W., Chang, M.Y., Ko, J.Y., Chang, J.W., Kim, M.J., Lee, M.R., Lee, S.A., Lee, H., Lee, G., Jung, B.H., Park, H., Kim, G.H., Kim, D., Cho, G., Kim, C.H., Ly, D.D., Park, K.S., Lee, S.H., 2022. LIN28A enhances regenerative capacity of human somatic tissue stem cells via metabolic and mitochondrial reprogramming. Cell Death Differ 29, 540-555.

Reinhardt, D.P., Keene, D.R., Corson, G.M., Pöschl, E., Bächinger, H.P., Gambee, J.E., Sakai, L.Y., 1996. Fibrillin-1: organization in microfibrils and structural properties. Journal of molecular biology 258, 104-116.

Ricci, L., Cabrera, F., Lotito, S., Tiozzo, S., 2016. Redeployment of germ layers related TFs shows regionalized expression during two non-embryonic developments. Developmental biology 416, 235-248.

Rinkevich, B., Ballarin, L., Martinez, P., Somorjai, I., Ben-Hamo, O., Borisenko, I., Berezikov, E., Ereskovsky, A., Gazave, E., Khnykin, D., 2022. A pan-metazoan concept for adult stem cells: the wobbling Penrose landscape. Biol Rev 97, 299-325.

Rinkevich, Y., Douek, J., Haber, O., Rinkevich, B., Reshef, R., 2007. Urochordate whole body regeneration inaugurates a diverse innate immune signaling profile. Developmental biology 312, 131-146.

Rinkevich, Y., Rosner, A., Rabinowitz, C., Lapidot, Z., Moiseeva, E., Rinkevich, B., 2010. Piwi positive cells that line the vasculature epithelium, underlie whole body regeneration in a basal chordate. Developmental biology 345, 94-104.

Rosental, B., Kowarsky, M., Seita, J., Corey, D.M., Ishizuka, K.J., Palmeri, K.J., Chen, S.-Y., Sinha, R., Okamoto, J., Mantalas, G., 2018. Complex mammalian-like haematopoietic system found in a colonial chordate. Nature 564, 425-429.

Sasaki, A., Miyamoto, Y., Satou, Y., Satoh, N., Ogasawara, M., 2003. Novel endostyle-specific genes in the ascidian Ciona intestinalis. Zoolog Sci 20, 1025-1030.

Sewell, M.T., Legue, E., Liem, K.F., Jr., 2024. Tubb4b is required for multi-ciliogenesis in the mouse. Development 151.

Sharma, S., Wang, W., Stolfi, A., 2019. Single-cell transcriptome profiling of the *Ciona* larval brain. Developmental biology 448, 226-236.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck III, W.M., Hao, Y., Stoeckius, M., Smibert, P., Satija, R., 2019. Comprehensive integration of single-cell data. Cell 177, 1888-1902. e1821.

Takagi, W., Sugahara, F., Higuchi, S., Kusakabe, R., Pascual-Anaya, J., Sato, I., Oisi, Y., Ogawa, N., Miyanishi, H., Adachi, N., 2022. Thyroid and endostyle development in cyclostomes provides new insights into the evolutionary history of vertebrates. BMC biology 20, 76.

Takahashi, K., Yamanaka, S., 2006. Induction of pluripotent stem cells from mouse embryonic
and adult fibroblast cultures by defined factors. Cell 126, 663-676.
Takahashi, K., Yamanaka, S., 2016. A decade of transcription factor-mediated reprogramming
to pluripotency. Nature reviews. Molecular cell biology 17, 183-193.
Temiz, B., Clarke, R.M., Page, M., Lamare, M., Wilson, M.J., 2023. Identification and
characterisation of Botrylloides (Styelidae) species from Aotearoa New Zealand coasts. New
Zealand Journal of Marine and Freshwater Research, 1-19.
Tiozzo, S., Christiaen, L., Deyts, C., Manni, L., Joly, J.S., Burighel, P., 2005. Embryonic versus
blastogenetic development in the compound ascidian Botryllus schlosseri: insights from Pitx
expression patterns. Developmental dynamics : an official publication of the American
Association of Anatomists 232, 468-478.
Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J., Livak, K.J.,
Mikkelsen, T.S., Rinn, J.L., 2014. The dynamics and regulators of cell fate decisions are revealed
by pseudotemporal ordering of single cells. Nature biotechnology 32, 381-386.
Voskoboynik, A., Soen, Y., Rinkevich, Y., Rosner, A., Ueno, H., Reshef, R., Ishizuka, K.J., Palmeri,
K.J., Moiseeva, E., Rinkevich, B., 2008a. Identification of the endostyle as a stem cell niche in
a colonial chordate. Cell stem cell 3, 456-464.
Voskoboynik, A., Soen, Y., Rinkevich, Y., Rosner, A., Ueno, H., Reshef, R., Ishizuka, K.J., Palmeri,
K.J., Moiseeva, E., Rinkevich, B., Weissman, I.L., 2008b. Identification of the endostyle as a
stem cell niche in a colonial chordate. Cell stem cell 3, 456-464.
Wagner, T., Beer, L., Gschwandtner, M., Eckhart, L., Kalinina, P., Laggner, M., Ellinger, A.,
Gruber, R., Kuchler, U., Golabi, B., Tschachler, E., Mildner, M., 2019. The Differentiation-
Associated Keratinocyte Protein Cornifelin Contributes to Cell-Cell Adhesion of Epidermal and
Mucosal Keratinocytes. The Journal of investigative dermatology 139, 2292-2301 e2299.
Wolf, F.A., Angerer, P., Theis, F.J., 2018. SCANPY: large-scale single-cell gene expression data
analysis. Genome biology 19, 15.
Wright, R.K., Ermak, T.H., 1982. Cellular defense systems of the Protochordata, Phylogeny and
Ontogeny. Springer, pp. 283-320.
Wu, K., Ahmad, T., Eri, R., 2022. LIN28A: A multifunctional versatile molecule with future
therapeutic potential. World J Biol Chem 13, 35-46.
Yamagishi, M., Huang, T., Hozumi, A., Onuma, T.A., Sasakura, Y., Ogasawara, M., 2022.
Differentiation of endostyle cells by Nkx2-1 and FoxE in the ascidian Ciona intestinalis type A:
insights into shared gene regulation in glandular- and thyroid-equivalent elements of the
chordate endostyle. Cell Tissue Res 390, 189-205.
Zappia, L., Oshlack, A., 2018. Clustering trees: a visualization for evaluating clusterings at
multiple resolutions. Gigascience 7, giy083.
Zhang, L., Yang, Y., Li, B., Scott, I.C., Lou, X., 2018. The DEAD-box RNA helicase Ddx39ab is
essential for myocyte and lens development in zebrafish. Development 145.
Zhang, T., Xu, Y., Imai, K., Fei, T., Wang, G., Dong, B., Yu, T., Satou, Y., Shi, W., Bao, Z., 2020. A
single-cell analysis of the molecular lineage of chordate embryogenesis. Science advances 6,
eabc4773.
Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A.H., Tanaseichuk, O., Benner, C.,
Chanda, S.K., 2019. Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. Nature communications 10, 1-10.
Zondag, L., Clarke, R.M., Wilson, M.J., 2019. Histone deacetylase activity is required for
Botrylloides leachii whole-body regeneration. J Exp Biol 222.
814