# Protein restriction associated with high fat induces metabolic dysregulation without obesity in juvenile mice

Amélie Joly[1], Jean-Louis Thoumas[1], Anne Lambert[1], François Leulier[1], Filipe De Vadder[1†]

[1] Institut de Génomique Fonctionnelle de Lyon, École Normale Supérieure de Lyon, CNRS UMR 5242, UCBL Lyon-1, F-69007 Lyon, France
[†] To whom correspondence should be addressed: filipe.de_vadder@ens-lyon.fr

Running head (max 60 characters including spaces):

Diet-induced metabolic dysfunction in lean juveniles

**Abstract:** max 250 words

Dysregulation of energy metabolism, including hyperglycemia, insulin resistance and fatty liver have been reported in a substantial proportion of lean children. However, non-obese murine models recapitulating these features are lacking to study the mechanisms underlying the development of metabolic dysregulations in lean children. Here, we develop a model of diet-induced metabolic dysfunction without obesity in juvenile mice by feeding male and female mice a diet reflecting Western nutritional intake combined with protein restriction (mWD) during 5 weeks after weaning. mWD-fed mice (32% fat, 8% protein) do not exhibit significant weight gain and have moderate increase in adiposity compared to control mice (16% fat, 20% protein). After 3 weeks of mWD, juvenile mice have impaired glucose metabolism including hyperglycemia, insulin resistance and glucose intolerance. mWD also triggers hepatic metabolism alterations, as shown by the development of simple liver steatosis. Both male and female mice fed with mWD displayed metabolic dysregulation, which a probiotic treatment with *Lactiplantibacillus plantarum* WJL failed to improve. Overall, mWD-fed mice appear to be a good preclinical model to study the development of diet-induced metabolic dysfunction without obesity in juveniles.

**Keywords:** Metabolic dysfunction, Lean, Juvenile, MASLD

**Introduction**:

Childhood represents a critical developmental window during which environmental cues, notably nutrition, significantly influence body mass index and energy metabolism (1). Notably, pediatric overweight and obesity result from an energy intake exceeding a child's requirement, and affects almost 40 million children under 5 worldwide (2). It is estimated that 6-39% of obese children will develop metabolic dysregulation, including elevated blood glucose, insulin resistance and metabolic dysfunction-associated steatotic liver disease (MASLD), which elevate the risk of metabolic disorders such as type 2 diabetes and cardiovascular diseases (3, 4). Even though its prevalence is much higher in obese and overweight children, metabolic dysregulation can also occur independently of BMI increase in children (5–9). Indeed, a recent retrospective study has shown that MASLD may affect 8% of

lean American adolescents (10). Likewise, the prevalence of insulin resistance, low HDL cholesterol and elevated plasma glucose among non-overweight and non-obese children can reach 2.2%, 8.1% and 6.6% respectively (11). However, the mechanisms underlying the development of these metabolic alterations in lean children are poorly understood.

Rodent models recapitulating the features of metabolic dysfunction without obesity are key for comprehending these mechanisms and their short and long-term effects on organismal physiology. However, current rodent models do not capture the diverse spectrum of metabolic dysfunctions observed in humans, nor accurately reflect the nutritional context under which these metabolic alterations develop in lean children. Specifically, diets used to model pediatric metabolic dysfunction without obesity are often extremely deficient in specific amino acids (12–14) or contain levels of fats or sugars that are too high to reflect the average intake in a Western diet (15, 16). Moreover, these murine models have been extensively described in adults, but to date very few rodent studies have investigated their effects in juvenile animals (17).

In this study, we aimed to bridge this gap by developing a model of metabolic dysfunction without obesity in juvenile mice, with a nutritional challenge consistent with the dietary intakes of children. To this end, we adapted the new total western diet (ntWD), which is a rodent diet designed to reflect dietary intake of overweight Americans and contains physiological amounts of fat and sugars from different sources (18). Previous work has shown that the ntWD induces substantial weight gain, but not obesity in adult mice, without clear modification of glucose metabolism (18, 19). Building on the fact that dietary amino acid restriction in juvenile mice is associated with glucose metabolism alterations (20–22), as well as hepatic steatosis onset (23, 24), we designed a modified version of the ntWD with a reduced amount of protein that we called the modified western diet (mWD). In juvenile mice, this diet induced features of a metabolic dysregulation including hyperglycemia, insulin resistance and hepatic steatosis, without substantial body mass index increase. The mWD therefore provides a model of diet-induced metabolic dysfunction without obesity in juvenile mice.

## Material and methods:

### Animals and diet:

Animal experiments have been carried out in accordance with the European Community Council Directive of September 22[nd], 2010 (2010/63/EU) regarding the protection of animals used for scientific purposes. All experiments have been approved by a local ethic committee (CECCAPP) and by the French Ministry of Research (APAFIS #40481-2022103017295302 v3).

Adult conventional C57Bl/6N male and female mice were purchased from Charles River. After one week of acclimatation, animals were mated and their offspring was used for experiments. The litters were reduced to 6 pups per dam at post-natal day 2. At post-natal day 21, male or female offspring were randomly weaned on control AIN93G diet containing 63.6% of carbohydrates, 15.9% of lipids and 20.5% of proteins (CD) or on the experimental diet corresponding to a modified version of the new total Western Diet (18) with moderate protein restriction and containing 58.3% of carbohydrates, 33.4% of lipids and 8.3% of proteins (mWD). Details of diet composition are provided in Table S1. Animals were weighed and measured once a week after flash anesthesia with 3% isoflurane, from weaning until post-natal day 56. Food intake was calculated on a weekly basis by subtracting the amount of remaining food from the amount provided the week before. At post-natal day 56, animals were fasted during 6h starting at 8 AM, flash anesthetized with 3% isoflurane and killed by cervical dislocation. Liver, perigonadal adipose tissue and subcutaneous adipose tissue were weighed and liver was kept in PBS-30% sucrose to be processed for histology or snap-frozen in liquid nitrogen and kept at −80°C until further analysis.

Probiotic treatment: Mice were treated 5 times a week with $10^9$ CFU/100 µL of *Lactiplantibacillus plantarum* WJL, or a placebo solution containing maltodextrin, administered using a pipette in the mouth of the mouse. Animals were held until they swallowed the solution.

### Glucose metabolism assessment:

An oral glucose tolerance test was performed at post-natal day 42, after 3 weeks of feeding with experimental diets. Animals were fasted during 6h starting at 8 AM and then gavaged with 2 mg/g of D-(+)-glucose (Sigma Aldrich, G70021). Glycemia was

measured from the tail vein using a Verio Reflect glucometer (OneTouch) before (0

min) and 15, 30, 45, 60, 90 and 120 min after glucose gavage. In addition, blood

samples were collected from the tip of the tail vein at 0, 15 and 30 min using

Microvette© CB 300 Z (Starstedt) and centrifuged 5 min at 10,000 g. Serum was

stored at −20°C until insulin measurement using Ultra Sensitive Mouse Insulin ELISA

kit (Crystal Chem, 90080), following the manufacturer's instructions.

<u>Lipid metabolism assessment:</u>

Liver triglycerides were measured using the GPO method (Biolabo, 87319) according

to the manufacturer's instructions.

Liver sections were stained with Oil Red O as previously described (24). Briefly, liver

samples were dehydrated overnight in 30% sucrose (Roth, 4621.1), embedded in

OCT (Leica, 14020108926) and kept at −80°C until further processing. 10 µm-thick

sections were cut using cryostat (Leica, CM 3050S), fixed in 4% paraformaldehyde

(Sigma, P6148) and stained with Oil Red O (Sigma, O-0625). Sections were imaged

using a DM6000 microscope (Leica) at 20x magnification. 10 images were taken per

mouse, on 4 different slides, choosing random fields on each slide. Images were

analyzed with Fiji software (25), using a modified version of the macro developed by

Nishad et al. (26), with the following parameters: threshold (0 – 95) and particle size

$(10 – 250 \ \mu m^2)$.

<u>RNA extraction and RT-qPCR:</u>

Total RNA from liver samples was extracted and purified using the Nucleospin RNA

kit (Macherey-Nagel, 740955.250). RNA quality and quantity was assessed using

Nano Drop 2000 (Thermo Fisher Scientific) and 1 µg of RNA was retrotranscribed

with the Sensifast cDNA synthesis kit (Meridian Biosciences, BIO-65053).

Quantitative PCR was performed on a Bio-rad CFX96 apparatus using Takyon[TM] No

ROX SYBR Mastermix blue dTTP (Eurogentec, UF-NSMT-B0701), 1/5 cDNA

dilutions of the reverse transcription product and specific primer sets (Table S2).

Analysis was performed according the MIQE guidelines (27). Amplification efficiency

was calculated for each primer set using cDNA serial dilutions and melting curves for

amplicons were analyzed to ensure unique and specific amplification. *Tbp*, *Rpl32* and

*Actb* were used as control genes after confirming their stability using the geNorm tool

available in the CFX Manager$^{TM}$ software (Biorad). Data are expressed as geometric
mean relative to the expression of control genes.
138
<u>Statistical analysis:</u>
Statistical analysis was performed with GraphPad Prism software version 10.2.1
(GraphPad software, San Diego, California, USA, www.graphpad.com) and using
parametric Student's t-test or the non-parametric Mann-Whitney test. Male and
Female datasets were analyzed separately to assess specifically the effect of the diet
in each sex without the interaction between the two parameters. Data were
considered to show evidence for an effect when the p-value was inferior to 0.05.
146
147
148　**Results:**
149
150　　1.　mWD does not induce obesity in juvenile mice
151
To explore the effects of modified western diet (mWD) feeding on physiology of
juvenile mice, we fed mice with mWD and as control CD diet for 5 weeks. Both male
and female mice fed with mWD during 5 weeks after weaning did not develop obesity
or overweight (Fig 1 A-B), contrary to what has been observed in response to classic
high-fat feeding (28–30). Indeed, males displayed similar weight-for-length ratio
compared to control diet-fed mice (Fig 1C-D). mWD-fed females gained slightly more
weight than CD animals, since they weighed on average 1.2 g more than controls at
post-natal day 56 (Fig 1B). In line with that, analysis of covariance (ANCOVA)
revealed that mWD females had slightly higher weight-for-length ratio than CD
females (Fig 1D). Despite a modest weight increase, we can consider that mWD fed
females did not develop overweight or obesity, similar to males. Consistently, mWD
did not induce changes in subcutaneous fat ratio (Fig 1E, G). However, juvenile male
but not female mice fed a mWD seemed to accumulate more visceral adipose tissue,
evaluated by the mass of perigonadal fat (Fig 1F, G), a type of visceral fat, which is a
typical feature of metabolic dysfunction (31).
The lack of weight gain in mWD fed animals was not due to a modification of caloric
intake, as cumulative food intake was similar between mWD and CD animals (Fig 1I-
J). However, this seemingly similar caloric intake hid discrepancies in major nutrient

intake, since mWD-fed mice consumed twice as much fat and half as much protein
as control mice (Figure S1). Therefore, mWD during the juvenile period is not
obesogenic despite increased accumulation of visceral fat in a sex-dependent
manner.
174
175       2.  mWD triggers glucose metabolism alterations that are independent of sex
176
We then characterized glucose metabolism in juvenile male and female mice after
3 weeks of mWD or CD feeding. Both males and females developed hyperglycemia
with respectively 38% and 47% increase in fasting glycemia compared to controls
(Figure 2 A, F). However, mWD feeding did not trigger changes in fasting insulinemia
(Figure 2B, G). In accordance, homeostatic model assessment of insulin resistance
(HOMA-IR) calculation showed evidence for insulin resistance in these animals
(Figure 2C, H). Thus, mWD feeding induces alterations of glucose metabolism during
fasting. An oral glucose tolerance test demonstrated that both mWD-fed males and
females had impaired glucose tolerance (Figure 2D, E, I, J). Altogether, these data
indicate that mWD feeding during the juvenile period triggers a complete
dysregulation of glucose homeostasis, independently of any weight gain and obesity.
188
189       3.  mWD feeding induces simple hepatic steatosis development
190
In addition to glucose metabolism dysfunction, lean children and adolescent can
develop MASLD (10). We thus investigated the extent to which mWD altered liver
metabolism. mWD-fed animals did not develop hepatomegaly (Figure 3A, C).
However, significant liver triglyceride accumulation was detected in response to
juvenile mWD feeding (Figure 3B, D), indicating that these animals suffer from liver
steatosis. Consistently, Oil Red O staining revealed increased lipid droplet size upon
mWD (Figure 3E-F).
Dyslipidemia, defined as elevated circulating levels of triglycerides or decreased
levels of high-density lipoprotein (32) is included in the scope of symptoms
associated to metabolic dysfunction. Circulating triglyceride levels were similar in the
portal vein and cava vein of mWD and CD fed animals (Figure S2A, B, D, E),
providing no evidence for dyslipidemia development in our model. These data
suggest no increased triglyceride absorption from the gut nor modified liver

triglyceride export. This is consistent with previous findings supporting the idea that liver steatosis is associated with *de novo* lipid synthesis in the liver (33). However, we did not detect modified expression of *Fasn* and *Acaa2*, two enzymes involved in triglyceride synthesis and oxidation respectively (Figure S2C, F).

MASLD is a disease that evolves throughout time from simple steatosis to liver fibrosis, cirrhosis and hepatocellular carcinoma (34). Hepatic fibrosis is associated with the appearance of several histological features including extracellular matrix development (34). To test whether mWD feeding induced liver fibrosis in addition to steatosis, we looked for markers of extracellular matrix accumulation. We detected no increased expression of several genes encoding collagen subunits (*Col1a2*, *Col3a1*, *Col4a1*) nor of genes coding for the extracellular matrix components desmotubulin (*Des*) and vimentin (*Vim*,) in the liver of male and female mice fed mWD (Figures 3G, H). Therefore, 5 weeks of mWD feeding during the juvenile period appears to lead to the development of simple liver steatosis, but not to fibrosis.

4. A chronic treatment with the probiotic bacterium LpWJL fails to rescue metabolic dysfunction induced by mWD

Modulating the gut microbiome through probiotic supplementation represents a promising therapeutic approach that has been shown to favorably impact metabolic parameters (35). In particular, the probiotic bacterium *Lactiplantibacillus plantarum$^{WJL}$* (LpWJL) has demonstrated beneficial effects on energy metabolism and growth of juvenile malnourished mice (20, 36). We thus hypothesized that LpWJL could ameliorate the features of metabolic dysfunction triggered by mWD feeding during mouse juvenile period. We chronically treated juvenile mWD-fed mice with $10^8$ CFU of LpWJL or with a placebo during 5 weeks after weaning. LpWJL did not induce modification of body weight and subcutaneous adipose tissue proportion in males and females (Fig 4A-B, G-H). Interestingly, male, but not female, mice treated with LpWJL displayed decreased visceral adipose tissue ratio (Fig 4C, I), a feature that was specifically altered by mWD in males (Fig 1F). However, LpWJL treatment failed to rescue glucose metabolism and hepatic alterations induced by mWD (Figure 4D-F, J-L). These data demonstrate that the beneficial effect of LpWJL treatment on the juvenile host is highly dependent on the nutritional context, with a positive impact upon chronic protein and lipid malnutrition but not upon mWD (20). Future

238    investigations will have to evaluate the ability of other specific probiotic strains or
239    consortia to ameliorate metabolic abnormalities in non-obese juvenile mice.
240
241    **Discussion:**
242
The prevalence of children suffering from metabolic dysfunction without obesity is
non-negligible and these children have an increased risk of mortality and
development of metabolic diseases (5, 9). Thus, the establishment of rodent models
recapitulating the features of this disorder is crucial to study the mechanistic basis of
its onset and evolution. In this study, we describe a modified version of the new total
western diet (18) with moderate protein restriction, which triggers the development of
metabolic dysfunction in juvenile mice without inducing obesity.
250
Juvenile male and female mice fed a mWD during 5 weeks did not develop obesity,
contrary to what has been described in this strain of mouse in response to feeding
with a high fat diet during the juvenile period (28–30). However, mWD-fed animals
had metabolic dysfunction, including hyperglycemia, impaired glucose tolerance,
insulin resistance and simple liver steatosis, but did not present evidence of liver
fibrosis. In murine models, fibrosis usually takes several months to develop. For
example, it was observed after 8 weeks of high-fat high-sucrose diet in a mouse
model of juvenile MASLD (17). In this study, we wanted to address metabolic defects
induced by mWD during the juvenile and peri-weaning period. We could hypothesize
that hepatic fibrosis would develop upon longer feeding with mWD. Of note, it is
evaluated that approximately 25% of fatty livers will progress toward steatosis and
cirrhosis, the other 75% remaining simple steatosis (37). Therefore, the simple
steatosis that we capture in our model is representative of the majority of cases.
264
Metabolic syndrome (MetS) constitutes a cluster of symptoms, including elevated
blood glucose, insulin resistance, MASLD, hypertension and obesity, which elevate
the risk of metabolic disorders such as type 2 diabetes and cardiovascular diseases
(4). However, the criteria for MetS proposed by different organizations have not been
fully harmonized (38). The type of metabolic dysfunction that we describe in this
study, occurring independently of some typical MetS symptoms, highlights that
metabolic disturbances can manifest in different ways, and that the traditional criteria

for MetS may not fully capture the heterogeneity of metabolic dysfunction, especially
in lean populations. Overall, mWD induces metabolic defects that recapitulate many
features of MetS in juveniles, but further work will be required to fully characterize
this model, and in particular its relevance for the cardiovascular disorders associated
with MetS.

Consumption of a diet presenting a western-type nutrient profile by adult male mice
does not induce metabolic disorder nor weight gain (18). Here, we show that
modifying this nutrient profile by decreasing the amount of protein is sufficient to
induce metabolic dysfunction in male and female juvenile mice. Interestingly, we
recently showed that male mice fed mWD between 35 and 119 post-natal days did
not develop obesity, hyperglycemia nor glucose intolerance (19), highlighting that the
post-weaning period is key for the development of metabolic alterations.

Several other rodent models of diet-induced metabolic dysfunction without obesity
have been previously described. However, they are associated with nutrient
composition that are not physiologically relevant to human health and usually induce
weight loss. Mice fed methionine- and choline-deficient diets have impaired
phosphatidylcholine synthesis and VLDL assembly, which causes hepatic triglyceride
accumulation and MASLD (13, 14). However, this diet induces significant body
weight loss (12, 13) and does not trigger insulin resistance (14). Besides, choline-
deficient amino acid defined diets mimic histological features of MASLD and avoid
extreme weight loss but do not induce insulin resistance nor impaired glucose
tolerance (13). To our knowledge, our model is the first to induce metabolic
dysfunction without obesity in juvenile mice, using a diet that does not involve
extreme deficiencies or excesses in essential nutrients.

Metabolic dysfunctions often are sexually dimorphic. For example, men are at higher
risk of developing more severe liver fibrosis compared to pre-menopausal but not
post-menopausal women (39). In addition, females are less susceptible to develop
metabolic dysfunction than males in many rodent models of diet-induced MetS (40,
41). Surprisingly, we did not detect strong sex-specific response to mWD feeding,
since both males and females had altered glucose metabolism and developed
hepatic steatosis to a similar extent. It is well established that estrogens play a critical

role in protecting females against metabolic syndrome (42, 43). In our study, juvenile female mice experienced nutritional challenge before the onset of sexual maturation, possibly overriding the protective effects of sex steroids. Therefore, our model provides an interesting opportunity to study metabolic dysfunction in both sexes.

Gut microbiota has emerged as a contributor of metabolic dysfunction development, including insulin resistance, dyslipidemia and hepatic steatosis (4, 44). Interestingly, among children with obesity, those suffering from metabolic dysfunction have specific alterations of the gut microbiota composition (3, 45, 46). While previous studies have demonstrated that probiotic treatment can improve the metabolic alterations of obese children (47, 48), we did not found beneficial effects of the strain *Lactiplantibacillus plantarum* WJL (LpWJL) in mWD-fed juvenile mice. Positive outcome of treatment with LpWJL have previously been demonstrated upon low-protein low-fat feeding in juvenile mice (20), emphasizing the strong connection between probiotic effectiveness and nutritional context.

In conclusion, our study shows that feeding juvenile mice a diet that recapitulates western dietary intake combined with moderate protein restriction induces metabolic dysregulation without obesity, independently of sex. The results of this study might be used to model metabolic alterations in young children and gain new insights in the molecular mechanisms underpinning the development of this understudied pathology.

Bibliography:

1.      **Silventoinen K, Jelenkovic A, Sund R, Hur Y-M, Yokoyama Y, Honda C, Hjelmborg J vB, Möller S, Ooki S, Aaltonen S, Ji F, Ning F, Pang Z, Rebato E, Busjahn A, Kandler C, Saudino KJ, Jang KL, Cozen W, Hwang AE, Mack TM, Gao W, Yu C, Li L, Corley RP, Huibregtse BM, Christensen K, Skytthe A, Kyvik KO, Derom CA, Vlietinck RF, Loos RJ, Heikkilä K, Wardle J, Llewellyn CH, Fisher A, McAdams TA, Eley TC, Gregory AM, He M, Ding X, Bjerregaard-Andersen M, Beck-Nielsen H, Sodemann M, Tarnoki AD, Tarnoki DL, Stazi MA, Fagnani C, D'Ippolito C, Knafo-Noam A, Mankuta D, Abramson L, Burt SA, Klump KL, Silberg JL, Eaves LJ, Maes HH, Krueger RF, McGue M, Pahlen S, Gatz M, Butler DA, Bartels M, van Beijsterveldt TC, Craig JM, Saffery R, Freitas DL, Maia JA, Dubois L, Boivin M, Brendgen M, Dionne G, Vitaro F, Martin NG, Medland SE, Montgomery GW, Chong Y, Swan GE, Krasnow R, Magnusson PK, Pedersen NL, Tynelius P, Lichtenstein P, Haworth CM, Plomin R, Bayasgalan G, Narandalai D, Harden KP, Tucker-Drob EM, Öncel SY, Aliev F, Spector T, Mangino M, Lachance G, Baker LA, Tuvblad C, Duncan GE, Buchwald D, Willemsen G, Rasmussen F, Goldberg JH, Sørensen TI, Boomsma DI, Kaprio J.** Genetic and environmental effects on body mass index from infancy to the onset of

adulthood: an individual-based pooled analysis of 45 twin cohorts participating in the
COllaborative project of Development of Anthropometrical measures in Twins
(CODATwins) study. *Am J Clin Nutr* 104: 371–379, 2016. doi: 10.3945/ajcn.116.130252.
2.        **UNICEF**, **WHO**, **World Bank Group Joint Child Malnutrition**. Levels and trends
in child malnutrition. 2023.
3.        **Carrizales-Sánchez AK**, **García-Cayuela T**, **Hernández-Brenes C**, **Senés-**
**Guerrero C**. Gut microbiota associations with metabolic syndrome and relevance of its study
in pediatric subjects. *Gut Microbes* 13: 1960135, 2021. doi:
10.1080/19490976.2021.1960135.
4.        **Dabke K**, **Hendrick G**, **Devkota S**. The gut microbiome and metabolic syndrome. *J*
*Clin Invest* 129: 4050–4057, 2019. doi: 10.1172/JCI129194.
5.        **Friend A**, **Craig L**, **Turner S**. The prevalence of metabolic syndrome in children: a
systematic review of the literature. *Metab Syndr Relat Disord* 11: 71–80, 2013. doi:
10.1089/met.2012.0122.
6.        **Bitew ZW**, **Alemu A**, **Tenaw Z**, **Alebel A**, **Worku T**, **Ayele EG**. Prevalence of
Metabolic Syndrome among Children and Adolescents in High-Income Countries: A
Systematic Review and Meta-Analysis of Observational Studies. *BioMed Research*
*International* 2021: e6661457, 2021. doi: 10.1155/2021/6661457.
7.        **Kelishadi R**, **Hovsepian S**, **Djalalinia S**, **Jamshidi F**, **Qorbani M**. A systematic
review on the prevalence of metabolic syndrome in Iranian children and adolescents. *J Res*
*Med Sci* 21: 90, 2016. doi: 10.4103/1735-1995.192506.
8.        **Ahrens W**, **Moreno LA**, **Mårild S**, **Molnár D**, **Siani A**, **De Henauw S**, **Böhmann J**,
**Günther K**, **Hadjigeorgiou C**, **Iacoviello L**, **Lissner L**, **Veidebaum T**, **Pohlabeln H**, **Pigeot**
**I**. Metabolic syndrome in young children: definitions and results of the IDEFICS study. *Int J*
*Obes* 38: S4–S14, 2014. doi: 10.1038/ijo.2014.130.
9.        **Guzmán-Guzmán IP**, **Salgado-Bernabé AB**, **Muñoz Valle JF**, **Vences-Velázquez**
**A**, **Parra-Rojas I**. Prevalence of metabolic syndrome in children with and without obesity.
*Med Clin (Barc)* 144: 198–203, 2015. doi: 10.1016/j.medcle.2015.05.031.
10.      **Sweeny KF**, **Lee CK**. Nonalcoholic Fatty Liver Disease in Children. *Gastroenterol*
*Hepatol (N Y)* 17: 579–587, 2021.
11.      **Sharma V**, **Coleman S**, **Nixon J**, **Sharples L**, **Hamilton□Shield J**, **Rutter H**,
**Bryant M**. A systematic review and meta□analysis estimating the population prevalence of
comorbidities in children and adolescents aged 5 to 18 years. *Obes Rev* 20: 1341–1349, 2019.
doi: 10.1111/obr.12904.
12.      **Leclercq IA**, **Farrell GC**, **Field J**, **Bell DR**, **Gonzalez FJ**, **Robertson GR**. CYP2E1
and CYP4A as microsomal catalysts of lipid peroxides in murine nonalcoholic steatohepatitis.
*J Clin Invest* 105: 1067–1075, 2000. doi: 10.1172/JCI8814.
13.      **Haczeyni F**, **Yeh MM**, **Ioannou GN**, **Leclercq IA**, **Goldin R**, **Dan YY**, **Yu J**, **Teoh**
**NC**, **Farrell GC**. Mouse models of non-alcoholic steatohepatitis: A reflection on recent
literature. *Journal of Gastroenterology and Hepatology* 33: 1312–1320, 2018. doi:
10.1111/jgh.14122.
14.      **Rinella ME**, **Green RM**. The methionine-choline deficient dietary model of
steatohepatitis does not exhibit insulin resistance. *Journal of Hepatology* 40: 47–51, 2004.
doi: 10.1016/j.jhep.2003.09.020.
15.      **Moreno-Fernández S**, **Garcés-Rimón M**, **Vera G**, **Astier J**, **Landrier JF**, **Miguel**
**M**. High Fat/High Glucose Diet Induces Metabolic Syndrome in an Experimental Rat Model.
*Nutrients* 10: 1502, 2018. doi: 10.3390/nu10101502.
16.      **Xu R**, **Pan J**, **Zhou W**, **Ji G**, **Dang Y**. Recent advances in lean NAFLD. *Biomedicine*
*& Pharmacotherapy* 153: 113331, 2022. doi: 10.1016/j.biopha.2022.113331.
17.      **Marin V**, **Rosso N**, **Ben MD**, **Raseni A**, **Boschelle M**, **Degrassi C**, **Nemeckova I**,

Nachtigal P, Avellini C, Tiribelli C, Gazzin S. An Animal Model for the Juvenile Non-Alcoholic Fatty Liver Disease and Non-Alcoholic Steatohepatitis. *PLOS ONE* 11: e0158817, 2016. doi: 10.1371/journal.pone.0158817.

18.    Monsanto SP, Hintze KJ, Ward RE, Larson DP, Lefevre M, Benninghoff AD. The new total Western diet for rodents does not induce an overweight phenotype or alter parameters of metabolic syndrome in mice. *Nutrition Research* 36: 1031–1044, 2016. doi: 10.1016/j.nutres.2016.06.002.

19.    Thoumas J-L, Cavaroc A, Sery D, Leulier F, Vadder FD. Suboptimal refeeding compensates stunting in a mouse model of juvenile malnutrition. bioRxiv: 2024.03.25.586077, 2024.

20.    Schwarzer M, Gautam UK, Makki K, Lambert A, Brabec T, Joly A, Šrůtková D, Poinsot P, Novotná T, Geoffroy S, Courtin P, Hermanová PP, Matos RC, Landry JJM, Gérard C, Bulteau A-L, Hudcovic T, Kozáková H, Filipp D, Chapot-Chartier M-P, Šinkora M, Peretti N, Boneca IG, Chamaillard M, Vidal H, De Vadder F, Leulier F. Microbe-mediated intestinal NOD2 stimulation improves linear growth of undernourished infant mice. *Science* 379: 826–833, 2023. doi: 10.1126/science.ade9767.

21.    Borck PC, Batista TM, Vettorazzi JF, Camargo RL, Boschero AC, Vieira E, Carneiro EM. Protein malnutrition after weaning disrupts peripheral clock and daily insulin secretion in mice. *The Journal of Nutritional Biochemistry* 50: 54–65, 2017. doi: 10.1016/j.jnutbio.2017.08.013.

22.    Soriano S, Gonzalez A, Marroquí L, Tudurí E, Vieira E, Amaral AG, Batista TM, Rafacho A, Boschero AC, Nadal A, Carneiro EM, Quesada I. Reduced Insulin Secretion in Protein Malnourished Mice Is Associated with Multiple Changes in the β-Cell Stimulus-Secretion Coupling. *Endocrinology* 151: 3543–3554, 2010. doi: 10.1210/en.2010-0008.

23.    Van Zutphen T, Ciapaite J, Bloks VW, Ackereley C, Gerding A, Jurdzinski A, De Moraes RA, Zhang L, Wolters JC, Bischoff R, Wanders RJ, Houten SM, Bronte-Tinkew D, Shatseva T, Lewis GF, Groen AK, Reijngoud D-J, Bakker BM, Jonker JW, Kim PK, Bandsma RHJ. Malnutrition-associated liver steatosis and ATP depletion is caused by peroxisomal and mitochondrial dysfunction. *Journal of Hepatology* 65: 1198–1208, 2016. doi: 10.1016/j.jhep.2016.05.046.

24.    Hu G, Ling C, Chi L, Thind MK, Furse S, Koulman A, Swann JR, Lee D, Calon MM, Bourdon C, Versloot CJ, Bakker BM, Gonzales GB, Kim PK, Bandsma RHJ. The role of the tryptophan-NAD□+□pathway in a mouse model of severe malnutrition induced liver dysfunction. *Nat Commun* 13: 7576, 2022. doi: 10.1038/s41467-022-35317-y.

25.    Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. Fiji: an open-source platform for biological-image analysis. *Nat Methods* 9: 676–682, 2012. doi: 10.1038/nmeth.2019.

26.    Nishad A, Naseem A, Rani S, Malik S. Automated qualitative batch measurement of lipid droplets in the liver of bird using ImageJ. *STAR Protocols* 4: 102466, 2023. doi: 10.1016/j.xpro.2023.102466.

27.    Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 55: 611–622, 2009. doi: 10.1373/clinchem.2008.112797.

28.    Podrini C, Cambridge EL, Lelliott CJ, Carragher DM, Estabel J, Gerdin A-K, Karp NA, Scudamore CL, Ramirez-Solis R, White JK, Sanger Mouse Genetics Project. High-fat feeding rapidly induces obesity and lipid derangements in C57BL/6N mice. *Mamm Genome* 24: 240–251, 2013. doi: 10.1007/s00335-013-9456-0.

29.     **Lizier M**, **Bomba L**, **Minuti A**, **Chegdani F**, **Capraro J**, **Tondelli B**, **Mazza R**,
**Callegari ML**, **Trevisi E**, **Rossi F**, **Marsan PA**, **Lucchini F**. The nutrigenomic investigation
of C57BL/6N mice fed a short-term high-fat diet highlights early changes in clock genes
expression. *Genes Nutr* 8: 465–474, 2013. doi: 10.1007/s12263-013-0344-8.
30.     **Deol P**, **Fahrmann J**, **Yang J**, **Evans JR**, **Rizo A**, **Grapov D**, **Salemi M**,
**Wanichthanarak K**, **Fiehn O**, **Phinney B**, **Hammock BD**, **Sladek FM**. Omega-6 and
omega-3 oxylipins are implicated in soybean oil-induced obesity in mice. *Sci Rep* 7: 12488,
2017. doi: 10.1038/s41598-017-12624-9.
31.     **Akagiri S**, **Naito Y**, **Ichikawa H**, **Mizushima K**, **Takagi T**, **Handa O**, **Kokura S**,
**Yoshikawa T**. A Mouse Model of Metabolic Syndrome; Increase in Visceral Adipose Tissue
Precedes the Development of Fatty Liver and Insulin Resistance in High-Fat Diet-Fed Male
KK/Ta Mice. *J Clin Biochem Nutr* 42: 150–157, 2008. doi: 10.3164/jcbn.2008022.
32.     **Jellinger PS**, **Handelsman Y**, **Rosenblit PD**, **Bloomgarden ZT**, **Fonseca VA**,
**Garber AJ**, **Grunberger G**, **Guerin CK**, **Bell DSH**, **Mechanick JI**, **Pessah-Pollack R**,
**Wyne K**, **Smith D**, **Brinton EA**, **Fazio S**, **Davidson M**, **Zangeneh F**, **Bush MA**.
AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS AND AMERICAN
COLLEGE OF ENDOCRINOLOGY GUIDELINES FOR MANAGEMENT OF
DYSLIPIDEMIA AND PREVENTION OF CARDIOVASCULAR DISEASE -
EXECUTIVE SUMMARYComplete Appendix to Guidelines available at
http://journals.aace.com. *Endocr Pract* 23: 479–497, 2017. doi: 10.4158/EP171764.GL.
33.     **Taguchi Y**, **Toyoshima Y**, **Tokita R**, **Kato H**, **Takahashi S-I**, **Minami S**.
Triglyceride synthesis in hepatocytes isolated from rats fed a low-protein diet is enhanced
independently of upregulation of insulin signaling. *Biochemical and Biophysical Research*
*Communications* 490: 800–805, 2017. doi: 10.1016/j.bbrc.2017.06.120.
34.     **Chan W-K**, **Chuah K-H**, **Rajaram RB**, **Lim L-L**, **Ratnasingam J**, **Vethakkan SR**.
Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD): A State-of-the-Art
Review. *J Obes Metab Syndr* 32: 197–213, 2023. doi: 10.7570/jomes23052.
35.     **Xavier-Santos D**, **Bedani R**, **Lima ED**, **Saad SMI**. Impact of probiotics and
prebiotics targeting metabolic syndrome. *Journal of Functional Foods* 64: 103666, 2020. doi:
10.1016/j.jff.2019.103666.
36.     **Schwarzer M**, **Makki K**, **Storelli G**, **Machuca-Gayet I**, **Srutkova D**, **Hermanova P**,
**Martino ME**, **Balmand S**, **Hudcovic T**, **Heddi A**, **Rieusset J**, **Kozakova H**, **Vidal H**,
**Leulier F**. Lactobacillus plantarum strain maintains growth of infant mice during chronic
undernutrition. *Science* 351: 854–857, 2016. doi: 10.1126/science.aad8588.
37.     **Williams CD**, **Stengel J**, **Asike MI**, **Torres DM**, **Shaw J**, **Contreras M**, **Landt CL**,
**Harrison SA**. Prevalence of nonalcoholic fatty liver disease and nonalcoholic steatohepatitis
among a largely middle-aged population utilizing ultrasound and liver biopsy: a prospective
study. *Gastroenterology* 140: 124–131, 2011. doi: 10.1053/j.gastro.2010.09.038.
38.     **Ramírez-Mejía MM**, **Qi X**, **Abenavoli L**, **Romero-Gómez M**, **Eslam M**, **Méndez-**
**Sánchez N**. Metabolic dysfunction: The silenced connection with fatty liver disease. *Ann*
*Hepatol* 28, 2023. doi: 10.1016/j.aohep.2023.101138.
39.     **Yang JD**, **Abdelmalek MF**, **Pang H**, **Guy CD**, **Smith AD**, **Diehl AM**, **Suzuki A**.
Gender and Menopause Impact Severity of Fibrosis Among Patients with Nonalcoholic
Steatohepatitis. *Hepatology* 59: 1406–1414, 2014. doi: 10.1002/hep.26761.
40.     **Kucsera D**, **Tóth VE**, **Gergő D**, **Vörös I**, **Onódi Z**, **Görbe A**, **Ferdinandy P**, **Varga**
**ZV**. Characterization of the CDAA Diet-Induced Non-alcoholic Steatohepatitis Model: Sex-
Specific Differences in Inflammation, Fibrosis, and Cholesterol Metabolism in Middle-Aged
Mice. *Front Physiol* 12: 609465, 2021. doi: 10.3389/fphys.2021.609465.
41.     **Chaix A**, **Deota S**, **Bhardwaj R**, **Lin T**, **Panda S**. Sex- and age-dependent outcomes
of 9-hour time-restricted feeding of a Western high-fat high-sucrose diet in C57BL/6J mice.

*Cell Rep* 36: 109543, 2021. doi: 10.1016/j.celrep.2021.109543.

42.    **Mauvais-Jarvis F**. Estrogen and androgen receptors: regulators of fuel homeostasis
and emerging targets for diabetes and obesity. *Trends in Endocrinology & Metabolism* 22:
24–33, 2011. doi: 10.1016/j.tem.2010.10.002.

43.    **Finan B**, **Yang B**, **Ottaway N**, **Stemmer K**, **Müller TD**, **Yi C-X**, **Habegger K**,
**Schriever SC**, **García-Cáceres C**, **Kabra DG**, **Hembree J**, **Holland J**, **Raver C**, **Seeley RJ**,
**Hans W**, **Irmler M**, **Beckers J**, **de Angelis MH**, **Tiano JP**, **Mauvais-Jarvis F**, **Perez-Tilve**
**D**, **Pfluger P**, **Zhang L**, **Gelfanov V**, **DiMarchi RD**, **Tschöp MH**. Targeted estrogen
delivery reverses the metabolic syndrome. *Nat Med* 18: 1847–1856, 2012. doi:
10.1038/nm.3009.

44.    **Thomas MS**, **Blesso CN**, **Calle MC**, **Chun OK**, **Puglisi M**, **Fernandez ML**. Dietary
Influences on Gut Microbiota with a Focus on Metabolic Syndrome. *Metabolic Syndrome and*
*Related Disorders* 20: 429–439, 2022. doi: 10.1089/met.2021.0131.

45.    **Gallardo-Becerra L**, **Cornejo-Granados F**, **García-López R**, **Valdez-Lara A**, **Bikel**
**S**, **Canizales-Quinteros S**, **López-Contreras BE**, **Mendoza-Vargas A**, **Nielsen H**, **Ochoa-**
**Leyva A**. Metatranscriptomic analysis to define the Secrebiome, and 16S rRNA profiling of
the gut microbiome in obesity and metabolic syndrome of Mexican children. *Microb Cell*
*Fact* 19: 1–18, 2020. doi: 10.1186/s12934-020-01319-y.

46.    **Alcazar M**, **Escribano J**, **Ferré N**, **Closa-Monasterolo R**, **Selma-Royo M**, **Feliu A**,
**Castillejo G**, **Luque V**, **Obemat2.0 Study Group**. Gut microbiota is associated with
metabolic health in children with obesity. *Clin Nutr* 41: 1680–1688, 2022. doi:
10.1016/j.clnu.2022.06.007.

47.    **Chen A-C**, **Fang T-J**, **Ho H-H**, **Chen J-F**, **Kuo Y-W**, **Huang Y-Y**, **Tsai S-Y**, **Wu S-**
**F**, **Lin H-C**, **Yeh Y-T**. A multi-strain probiotic blend reshaped obesity-related gut dysbiosis
and improved lipid metabolism in obese children. *Front Nutr* 9: 922993, 2022. doi:
10.3389/fnut.2022.922993.

48.    **Li Y**, **Liu T**, **Qin L**, **Wu L**. Effects of probiotic administration on overweight or obese
children: a meta-analysis and systematic review. *J Transl Med* 21: 525, 2023. doi:
10.1186/s12967-023-04319-9.

Acknowledgements: The authors thank the animal facility of IGFL (PEHR) where

animal experiments were conducted, and the associated group for animal welfare

(SBEA). They also acknowledge Héloïse Dusson-Andreu for her help in setting up

liver histology.

Funding: This work was supported by funding from Fondation pour la Recherche

Médicale (Equipe FRM EQU202203014629). AJ is supported by a grant from

Fondation pour la Recherche Médicale (FDT202304016501).

Author contributions: AJ, FDV and FL conceived and designed research. AJ, JLT and

AL performed experiments. AJ analyzed data. AJ and FDV interpreted results of

experiments. AJ drafted manuscript. AJ, FDV and FL edited and revised manuscript.

All authors approved final version of manuscript.

**Figure legends**

Figure 1: Modified western diet feeding to juvenile male and female mice does not induce obesity.

(A-B) Body weight of male and female mice fed control diet (CD) or modified western diet (mWD) between weaning (postnatal day 21, P21) and P56. (C-D) Length-versus-weight of CD- and mWD-fed males and females. (E-H) Weight of subcutaneous adipose tissue (SAT) and visceral adipose tissue (VAT) measured at P56 and expressed per gram of body weight. (I-J) Cumulative food intake.

Data are expressed as mean ± SEM.

* $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$. (A-H): unpaired t-test. (I-J): Mann-Whitney test

Figure 2: Male and female mice fed mWD after weaning have impaired glucose metabolism.

Fasting glycemia (A, F) and insulinemia (B, G) of control (CD) or mWD-fed males and females at post-natal day 42 (P42). (C, H) Homeostatic model assessment of insulin resistance (HOMA-IR) calculated at P42. (D, I) Oral glucose tolerance test performed on CD and mWD males and females at P42 and area under the curve for this test (E, J).

Data are expressed as mean ± SEM.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, unpaired t-test.

Figure 3: mWD feeding during the juvenile period triggers simple hepatic steatosis in both males and females.

Weight (A, C) and triglyceride concentration (B,D) in the liver of CD- and mWD-fed males (blue) and females (green) at post-natal day 56 (P56). (E, F) Representative images of Oil Red O stained liver sections and quantification of mean lipid droplet sizes. Scale bars: 50 µm. (G, H) Liver *Col1a2*, *Col3a1*, *Col4a1*, *Des* and *Vim* expression in CD- and mWD-fed males and females.

(A-F): Data are expressed as mean ± SEM. (G, H): Data are expressed as geometric mean ± geometric SD.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. (B, D): unpaired t-test. (E, F): Mann-Whitney test.

<u>Figure 4</u>: LpWJL treatment fails to improve metabolic dysfunction induced by mWD
feeding during the juvenile period.
Weight of body (A, G), subcutaneous adipose tissue (B, H) and visceral adipose
tissue (C, I) of male (blue) and female (green) mice fed mWD and treated with
placebo or with LpWJL at post-natal day 56. (D, J) Plasma glucose after 6 hours of
food deprivation and oral glucose tolerance test (E, K) performed at post-natal day
42. (F, L) Liver triglyceride content.
Data are expressed as mean ± SEM.
* p<0.05, unpaired t-test.
581
582 <u>Figure S1</u>: mWD-fed animals consume more lipids and less protein than CD-fed
583 animals
584 Cumulative food (A-B), carbohydrate (C-D), lipid (E-F) and protein (G-H) intake of
585 male (blue) and female (green) animals fed CD or mWD.
586 Data are expressed as mean ± SEM.
587
<u>Figure S2</u>: Circulating triglycerides and hepatic expression of lipid metabolism
enzymes are not affected by mWD feeding.
Triglyceride concentration in the serum from cava vein (A, D) and portal vein (B, E) of
male (blue) and female (green) animals fed CD or mWD. (C, F): Liver Fasn and
Acaa2 expression in CD and mWD-fed males and females.
(A, B, D, E): Data are expressed as mean ± SEM. (C, F): Data are expressed as
geometric mean ± geometric SD.
595
596