Range geographies, not functional traits, explain convergent range and phenology shifts under climate change

Catherine Sirois-Delisle[1]*, Susan C.C. Gordon[1], Jeremy T. Kerr[1]

[1]*University of Ottawa, 75 Laurier Ave E, Ottawa, ON K1N 6N5, Canada*

Correspondence: csiroisd@uottawa.ca

Abstract

Climate change may introduce conditions beyond species' tolerances; to survive, species must

avoid these extremes. Phenological shifts are one strategy, as species move their activity or life

history events in time to avoid extreme conditions. Species may also shift in space, moving their

ranges poleward to escape extremes. However, whether species are more likely to exhibit one or

both strategies, and whether this can be predicted based on a species' functional traits, is

unknown. Using a powerful macroecological dataset of European and North American odonate

observations, we assessed range and phenology shifts between two time periods (1980-2002 and

2008-2018) to measure the strength and direction of the association between responses. Species

with the greatest poleward range shifts also showed the largest phenological shifts toward earlier

annual activity periods, with half of all species shifting in both space and time. This response

was consistent across continents, despite highly divergent land use and biogeographical histories

in these regions. Surprisingly, species' range and phenology shifts were not related to functional

traits; rather, southern species shifted their range limits more strongly, while increasing

temperature variability hindered range shifts. By reducing risk through phenological shifts, the

resulting larger populations may be more likely to disperse and expand species' ranges. While

species shifting in both space and time may be more resilient to extreme conditions, we

identified a small number of species (approximately 10%) that failed to shift at all; these species

are likely to be particularly vulnerable to climate change, and should be prioritized for

conservation intervention.

Introduction

Climate change alters climatic means and increases the frequency of extreme weather events, exposing species to conditions outside of their tolerances, and often leading to population declines (Goulson, 2019; IPCC, 2021). Species may avoid extreme conditions by dispersing to new areas where conditions pose fewer weather-related challenges, often leading to poleward range expansion (Davis et al., 2005; Lawlor et al., 2024). Species' biological timing could also shift, through adaptation or phenotypic plasticity, with earlier warming advancing the timing of early-season activities and life-history events (Davis et al., 2005; Hällfors et al., 2024; Parmesan and Yohe, 2003). Both species' geographical ranges and seasonal timing depend strongly on climate and habitat conditions, with shifts in space and time permitting species to remain within the limits of their ecological niches (Figure 1; Chen et al., 2011; Menzel et al., 2006; Parmesan and Yohe, 2003). This allows populations to grow, despite changing environments, and reduces the risk of climate debt and extinction (Devictor et al., 2012; Franks et al., 2018; Lustenhouwer et al., 2018; Saino et al., 2010; Souza et al., 2023; Urban, 2015).

Positive population trends can be stronger in species that shift both their range and phenology (Hällfors et al., 2024). Greater phenological plasticity under warmer spring temperatures may increase reproductive success, leading to greater population growth and range expansions (Macgregor et al., 2019), as positive or stable trends in species abundance and habitat availability are essential for range shifts (Mair et al., 2014; Platts et al., 2019). However, there could also be a trade-off between phenological and geographical shifts (Amano et al., 2014; Hassall, 2015; Socolar et al., 2017). Species with greater dispersal abilities may have less need for phenological shifts as they track their climatic niche through space, while weaker dispersers may be confronted with greater selective pressure to shift phenology within their range (Amano

et al., 2014; Hassall, 2015; Socolar et al., 2017). Since range shifts can also be driven by

extirpations at species' trailing range edge (Parmesan et al., 1999), greater phenological shifts

may mitigate the need for range shifts, as species better tolerate new climatic conditions. Cross

continental studies report converging effects of climate change on species' range shifts and

abundances (Kerr et al., 2015; Stephens et al., 2016), but potential relationships between

phenological and geographic responses have not yet been investigated at continental scales.

Traits, such as dispersal, may determine species' spatial and temporal responses to

climate change (Chen et al., 2011; Kharouba et al., 2009; Schuetz et al., 2019; Zografou et al.,

2021). However, these relationships generalize poorly taxonomically and geographically, and

cross-continental tests have not been attempted (Angert et al., 2011; Buckley and Kingsolver,

2012; Estrada et al., 2016; MacLean and Beissinger, 2017).

Geographic locations and environmental characteristics of species' ranges may also

predict range shifts, as species with high latitude ranges have been shown to exhibit smaller

range shifts (MacLean and Beissinger, 2017), while increasing local temperature and loss of

natural landcover may drive range retractions (Pacifici et al., 2020). However, exposure to

extreme climate events (e.g. drought, heat waves, or storms) within species ranges may disrupt

species' dispersal abilities and capacities to tolerate new conditions (Kerr, 2020; Román-Palacios

and Wiens, 2020). Exposure to thermal anomalies can rapidly change entire communities and

create shifts towards new ecosystems, sometimes leading to local declines (Day et al., 2018;

Grant et al., 2017; Harris et al., 2018; Román-Palacios and Wiens, 2020).

While global change research on insects often emphasizes butterfly and bee taxa, recently

assembled databases of odonate observations provide opportunities to answer new questions

(Kalkman et al., 2018). Dragonflies and damselflies respond rapidly to climate change, both

spatially (Ball-Damerow et al., 2015; Grewe et al., 2013; Hickling et al., 2005; Powney et al.,

2015; Rapacciuolo et al., 2017; Sirois-Delisle and Kerr, 2022) and temporally (Hassall, 2015;

Hassall et al., 2007; McCauley et al., 2018), making them an excellent indicator species (Hassall,

2015). There is some evidence that functional traits relate to odonates' interspecific variation in

range shifts (Angert et al., 2011; Grewe et al., 2013), phenology shifts (Diamond et al., 2011;

Gutiérrez and Wilson, 2021; Zografou et al., 2021), extinction risks (Cardillo et al., 2008;

Cooper et al., 2008; Fritz et al., 2009; Suhonen et al., 2022), and rates of decline and expansion

within limited geographic scopes (Powney et al., 2015; Rapacciuolo et al., 2017). While

relationships between morphological traits and range boundaries have been shown for some

groups (i.e. Rundle et al., 2007), these may depend on species' geographic context. For example,

lentic species (i.e. species breeding in slow moving water like lakes or ponds) tend to have larger

ranges and higher dispersal capacities than lotic species (i.e. fast moving water-dwelling

species), and may track their climate niches more closely (Grewe et al., 2013; Hof et al., 2006).

Lotic habitat may be more difficult to reach, and lotic species may face stronger anthropogenic

pressures in places where their diversity is high (Kalkman et al., 2018), affecting species abilities

to track changes in climatic seasonal timing. Lentic species have been shown to expand their

range boundaries more than lotic species, but only among species with more southern

distributions (Grewe et al., 2013). Abundances of European odonates with ranges extending

south from sample landscapes have been found to respond more positively to climate change

than species with ranges extending north from sample locations, due to their geographic position

and thermal tolerance (Powney et al., 2015). Species previously limited by tolerance to low

temperatures could expand their ranges following warming (Devictor et al., 2008).

In this study, we tested whether species with stronger geographical range shifts also

advanced their emergence phenologies, or if one response offsets the need for the other. We also

asked whether biological traits, range geography (i.e. southerly vs. northerly), or temperature

variability predict range shifts at species' northern range limits, and whether these factors can

also predict shifts in species emergence phenology. We predicted that species would exhibit

shifts in both geography and phenology, as we expect species shifting in phenology to have

larger populations, increasing likelihood of dispersal and successful range shifts. We predicted

that generalist species would shift their phenologies and geographies, as generalists outperform

specialists as climate and land uses change (Ball-Damerow et al., 2015, 2014; Hassall and

Thompson, 2008; Powney et al., 2015; Rapacciuolo et al., 2017); in contrast, we expect

specialists to fail to shift in both space and time, making them more vulnerable to extirpation and

extinction. Alternatively, species might respond to rapid climate changes in ways that reflect

their geographical position, indicating that where they are found is a better predictor of their

conservation risk from global change than their intrinsic biological characteristics.

Methods

Biological records

We assembled ~2 million observations records for North American and European

odonate species collected between 1980 and 2018. Data sources included online dataset

aggregators GBIF (http://gbif.org/) and Canadensys (http://www.canadensys.net/), Odonata

Central (Abbott, 2020), and other institutions (see Acknowledgments). While odonates were

sampled opportunistically, biases associated with data that are not systematically collected are

less likely to affect trends at large spatial and temporal scales, particularly if data are obtained

from multiple independent sources (Pyke and Ehrlich, 2010a; Zattara and Aizen, 2021a). We

removed records with incomplete or missing species identification, year, or locality information.

We selected unique observations for species, location, year, and Julian day of collection, and

restricted the data to continental North America and Europe. We mapped species-specific

observations using ArcGIS software (ESRI, 2019), and qualitatively verified species ranges. In

the event that the same species was found in both Europe and North America, we selected

records from the continent containing the largest number of observations (Supplementary

Information).

We assigned species presences to 100 X 100 km quadrats to identify the best sampled

species. We excluded species found in fewer than 50 quadrats to increase the likelihood of

accurately predicting the position of species' northern range boundaries. We retained ~1,100,000

records comprising 76 species (Figure 2). Observation records were separated into two time

periods, to compare species' recent phenologies and northern range limits (2008 to 2018) to

conditions in a historical time period (1980 to 2002).

Climate data

Temperature variability during a species' flight season may impact its ability to establish

in new locations, or shift its emergence timing the following year. We downloaded a high-

resolution gridded dataset for monthly average daily maximum temperature from the Climatic

Research Unit (New et al., 2002). We extracted average values per 100 X 100 km quadrat across

the months of April to October, covering the flight period for odonates in this study, for each

year included in the historical and recent periods. We calculated the standard deviation per

quadrat for each time period and averaged these values per species and time period to measure

interannual temperature variability during species' flight season across their range. We used the

difference between recent and historical measures as an estimate of change in temperature

variability.

Spatial and temporal change metrics

To limit potential effects of temporal and spatial biases, we generated range and

phenology shift metrics with specific criteria for quadrats and species selection (Bartomeus et al.,

2018; Gaiji et al., 2013; García-Roselló et al., 2015); we retained 76 species for range shift

estimates (Supplementary Information). Species' northern range boundaries were calculated

using the mean of the 10 most northern points of each species range in both the historic (1980 –

2002) and recent (2008 – 2018) time periods, measured in kilometers from the equator (as in

Kerr et al., 2015). We used the difference between range limit positions in the historic recent

time periods to estimate species' northern range limit shifts.

Since spatiotemporal biases sometimes inflate range shift measurements (Kujala et al.,

2013), we used null models to test whether observed range shift estimates were robust, and the

extent to which those responses differed from expectations arising because of rising sampling

intensity over time. 1000 randomized datasets were created with the same number of species-

specific geographical points per time period as the number of actual observations. Maximum and

minimum latitude and longitude values were held constant relative to observed values. Northern

range limit shifts were calculated as the difference in the mean of the 10 most northern points

between the historic and recent time periods. We used generalized linear models (GLMs) to test

whether species-specific range limit shifts in each iteration predicted range shifts measured from

the observation data (Figure S1, S2). We found that observed northern range limit shifts are not

consistent with expectations derived from changes in sampling intensity.

We estimated species-specific emergence phenology for each time period in 200 X 200

km quadrats; using larger quadrats increases probabilities of detecting signals of emergence

phenology, which may otherwise be lost due to gaps in data density. We retained quadrats that

contained at least 25 observations for a given species in both time periods. To estimate

phenology per area, we used the *weib.limit* function of the *WeibullR* R package (Pearse et al.,

2017). This function uses the Weibull distribution to estimate the Julian day of a species' first

appearance, and is especially useful to measure the timing of phenological events in sparsely

sampled datasets. Techniques such as using the average of the n^{th} first observations of

phenological events, or the n^{th} percentile flight dates (Brooks et al., 2014; Robbirt et al., 2011),

tend to overestimate the timing of biological events due to temporal bias towards later days in the

species' active period (Pearse et al., 2017). We retained emergence estimates between March 1^{st}

and September 1^{st}, as well as species and quadrats that showed a difference in emergence

phenology of -25 to 25 days, -30 to 30 days, or -35 to 35 days between both time periods, to

include only phenology shifts that could be biologically meaningful to environmental climate

change. 68 species found across 63 quadrats met these criteria. Large changes in phenology are

likely explained by other anthropogenic or natural factors, or could occur due to noise in the

data, since these phenology calculations per region are extremely data intensive. We calculated

the difference in the day of emergence per quadrat between both time periods, as well as mean

phenology change across all quadrats for each species. The number of quadrats per species used

to calculate their mean phenological shift varied between 2 and 46.

We used null models to assess whether our approach to estimating phenology shifts was

robust, as potential issues may arise due to spatiotemporal biases in the underlying data (Kujala

et al., 2013). We constructed 1000 randomized datasets of species' hypothetical days of

occurrence, using the same number of quadrats, and observations within quadrats, as in each

time period of the observation records. We assigned the maximum and minimum Julian day of

occurrence from the observation records to limit values in the randomized datasets. We applied

the same method and criteria of inclusion to the randomized datasets as we did to measure

phenology shifts from the observation data. GLMs were built to test whether phenology shifts

calculated using 1000 random datasets predicted the phenology shifts that we measured. No

discernable pattern emerged, indicating that observed shifts in phenology are not consistent with

expectations derived from differences in sampling intensity over time (Figure S3, S4).

Ecological traits

To assemble trait data for the 76 species in the database, we used field guides (R.A.

Cannings, 2002; C.D. Jones et al., 2008; D. Paulson, 2012) and existing trait databases (G. D.

Powney et al., 2014; Waller et al., 2019a). We considered any evolved morphological,

physiological, behavioural, or life history characteristic as a trait (Beissinger and Riddell, 2021).

Geographic or climatic characteristics (e.g. range size, geographic position of species' ranges,

climatic conditions of the range) were not considered traits, as they are not products of evolution,

but rather consequences of environmental, ecological, biogeographical, or evolutionary patterns

linked to range shifting mechanisms (Beissinger and Riddell, 2021). Such variables may predict

species' responses to climate change nonetheless, and can add significant value to predictive

models (MacLean and Beissinger, 2017).

Along with two geographic and climatic attributes, including temperature variability and

distribution region (northern, southern, or widespread), we selected four traits likely to be

biologically relevant to spatial and temporal responses to climate change: flight duration,

breeding habitat type (lotic, lentic, or both), egg laying habitat (exophytic vs. endophytic), and

body size. Species' flight period was measured as the total number of days of the flight period,

estimated from the approximate time of the month of average first and last appearances.

Breeding habitat was assigned according to a species' uses of lotic, lentic, or both habitat types.

Body size corresponded to the mean length of the abdomen of each species. We excluded

overwintering stage and range size from our analysis as data were incomplete for many species,

and excluded migratory behavior as nearly all species included in the study were non-migratory.

We tested for correlations amongst all predictors by calculating the Predictive Power

Statistic (PPS) and Pearson correlations among traits (van der Laken, 2021): we found no

evidence of correlation.

Statistical analyses

We conducted statistical analyses using R Statistical Software (R Core Team, 2019). All

continuous variables were transformed into Z-scores using the *scale* function in R. First, we

investigated whether there was a relationship between species' range and phenological shifts. We

used a by-species GLM and a Markov chain Monte Carlo generalized linear mixed model

(MCMCglmm; Hadfield, 2010); we compared frequentist and Bayesian approaches to improve

the robustness of the results. We included a term to account for phylogeny in the MCMCglmm

model, as species that are closely related are likely to have similar traits. We used the

phylogenetic tree published by the Odonate Phenotypic Database (Waller et al., 2019a), which

was generated using a combination of DNA-sequences and species' taxonomy based on

morphology (Waller and Svensson, 2017). Trace and density plots for the MCMCglmm model

revealed no issues with autocorrelation or model convergence (Figure S7).

Next, we investigated whether biological traits, range geography, or temperature

variability predicted range shifts at species' northern range limits, and whether the same

predictors explaining range expansions could also predict shifts in species emergence phenology.

We constructed two sets of GLMs, in addition to two sets of MCMCglmms accounting for

phylogeny; one of each with changes in species' northern range limits as the response variable,

and the other with changes in emergence phenology as the response variable. We included

species traits, species range geography, and temperature variability experienced by species as

predictor variables. Trace and density plots for the MCMCglmm models did not indicate

limitations related to autocorrelation or model convergence (Figure S8).

In addition to the inclusion of phylogeny in statistical models to account for potential data

non-independence, we measured the phylogenetic signal in range and phenological shifts. We

used the *phylosig* function of the *Phytools* package version 0.7-70 (Revell, 2012), which

calculates phylogenetic signal using Pagel's lambda and Blomberg's K.

Results

Relationship between range and phenology shifts

Most species (69% of those included) expanded their northern range limits toward higher

latitudes (mean range expansions of 180 km). The average range expansion across all species

was 63 km northward (Figure S5). Some species showed range retractions (i.e. *Libeullula*

luctuosa); small range retractions may result from sampling variability or stochastic population

fluctuations along the northern range edge. Most species (60% of assessed species) maintained or

advanced their emergence phenology (mean of -2.71 days in emergence phenology shifts among

all species; Figure S6). Fewer species were included in phenology analyses than analyses of

range shifts due to the data intensity required to capture phenology shifts at a study site (N=66

vs. N = 76). Many species (50%) showed both advancing emergence phenology and range

expansions, while 10% of species showed neither range nor phenological shifts relative to

historical baselines. Species that did not respond to climate change temporally or spatially

included *Coenagrion mercuriale*, a European species that is listed as near threatened by the

IUCN Red List (IUCN, 2021), and is projected to lose 68% of its range by 2035 (Jaeschke et al.,

2013).

Species that shifted their northern range limits to higher latitudes also showed stronger

advances in their emergence phenology (Figure 3), a result that was consistent in GLM and

Bayesian analyses ($p < 0.05$; Table 1; $R^2 = 17\%$ and 15%, for GLM and MCMCglmm models,

respectively). This trend converged across continents, and is consistent among both dragonflies

and damselflies, although there is considerable interspecific variation in the magnitude of spatial

and temporal shifts. Accounting for phylogeny did not improve model predictions and did not

explain greater model variance (Table 1).

Drivers of range and phenology shifts

Range geography and climate variability, rather than ecological traits, predicted range

shifts in both North America and Europe, with range geography consistently the strongest

predictor. Species' ecological traits did not relate to the extent of observed phenological or

geographical range shifts in tests using GLM and MCMCglmm models. Species with more

southern distributions shifted their northern range limits towards higher latitudes more than

northern and widespread species ($p < 0.05$; Table 2; $R^2 = 23\%$ and 20%, for GLM and

MCMCglmm models, respectively). Species experiencing smaller changes in interannual

temperature variability also had a higher likelihood of northern range limit shifts ($p <$

0.05). Results from the GLM and MCMCglmm models were qualitatively similar, however a

smaller amount of model variance was explained when phylogeny was accounted for. Neither

range geography nor climate variability predicted emergence phenology shifts. Even though both

response types are related, this suggests that the mechanisms underlying phenological shifts are

different than those underlying range shifts.

A phylogenetic signal may indicate that there are traits that determine species' spatial and

temporal responses to changing climate that were not measured in this study. Yet we detected no

phylogenetic signal using Pagel's lambda or Blomberg's K in either geographical range or

phenological responses (Table 2), despite pronounced recent warming. Adding a phylogeny term

to the MCMCglmm models also failed to produce a pattern different to the GLMs, and model

performance was not improved when accounting for phylogeny in our assessment of northern

range shifts.

Discussion

In one of the first studies to investigate both shifts of phenology and range at a

continental scale, we find that dragonfly and damselfly species show pronounced geographical

and phenological shifts that converged across Europe and North America. Species expanding

their ranges poleward also emerged earlier in the spring on both continents (Figure 3), with shifts

predicted by range geography and climate variability, not ecological traits. These results suggest

that some species are 'universal winners' with respect to climate change: they demonstrate the

flexibility to respond both temporally and spatially to the onset of rapid climate change.

Conversely, species that show neither geographic nor phenological shifts may be particularly

vulnerable to climate change.

We found no evidence for a tradeoff between range and phenology shifts; instead, half of

species shifted both range and phenology. Earlier seasonal timing allows species to stay within

their climatic limits. This may lead to higher local population growth (Macgregor et al., 2019),

although earlier emergence could expose individuals to early season weather extremes

(McCauley et al., 2018). As only a small proportion of odonate adults undertake long range

dispersal (Conrad et al., 1999), greater local population sizes should contribute to higher

dispersal rates (Mair et al., 2014), facilitating range shifts (Kerr, 2020; Leroux et al., 2013). This

is consistent with results from other taxa: among British butterflies, early emergence increased

population growth and facilitated range shifts for species with multiple generations per year

(Macgregor et al., 2019); Finnish butterfly species with the greatest population growth rates

shifted both their phenology and ranges (Hällfors et al., 2021). Such population growth or

maintenance, and therefore the potential for range shifts, is only possible if habitat is available

(Mair et al., 2014). Future work should consider habitat availability alongside range and

phenology shifts, as it may help explain why some species are able to shift their phenology but

not their range.

Southern species were more likely to expand their ranges northward than northern or

widespread species. Species' ability to maintain large populations may be impaired in northern

latitudes, where rates of climate change are high (IPCC, 2021), hindering dispersal and

colonization that are necessary for range expansions (Mair et al., 2014). Southern species may

have narrower niche breadths than widespread or northern species, and may respond more

rapidly to climate change to track this narrower niche (Hällfors et al., 2024). As more southern

species expand northwards they may be better able to tolerate local temperature extremes due to

their warmer thermal niches, and should be better able to maintain populations than species with

northern distributions (Day et al., 2018), facilitating further range shifts.

Increasing frequency and severity of extreme weather limited species' geographical range

responses (Table 2). This trend was independent of traits that are mechanistically linked to

species' climate change responses, such as dispersal ability or habitat preference. Extreme

temperatures can reduce population sizes, leading to local extinctions (Román-Palacios and

Wiens, 2020), and reducing the likelihood of range expansions (Mair et al., 2014). In odonates,

experimental evidence has demonstrated that larval mortality rises with short-term extreme

weather (Mccauley et al., 2015). Individuals that shift phenologies earlier in the season to avoid

climate extremes could still be exposed to harmful conditions (Iler et al., 2021); for example,

odonate populations that respond to unusually warm spring temperatures may experience high

mortality if temperatures return to seasonal conditions. Species that experience extreme

conditions may then be unable to successfully shift in time, reducing population sizes, and

reducing the likelihood of range shifts.

In contrast to previous work demonstrating that range and phenology shifts are at least

partially determined by species traits (i.e. Sunday et al., 2015; Zografou et al., 2021), no

functional trait, or combination of traits, explained species' shifts. While we could not capture all

functional traits in this analysis, our results are consistent with other work that identifies climate

velocity and sensitivity as the best predictors of range shifts and thermal preferences tracking in

marine systems (Pinsky et al., 2013; Schuetz et al., 2019). Species' tolerances to increasingly

variable temperatures, rather than traits, also help predict extinction risk during climate change

(Kerr, 2020). More work is needed to determine if traits consistently fail to predict range shifts,

particularly at large spatial scales, although reliable observations at these scales are often

lacking. Detailed mechanistic models that incorporate multiple independent processes may help

untangle these patterns, although challenges with parameterization remain (Urban et al., 2022).

The geographic positions of species' ranges determine the local pressures and

environmental factors to which they are exposed (MacLean and Beissinger, 2017; Pacifici et al.,

2020), potentially masking the effects of traits (Schuetz et al., 2019). This process could cause

trait-related trends to differ across levels of biological organization (Srivastava et al., 2021), from

local populations (where traits might be critical) to biogeographical extents (where traits might

be unrelated to range or phenological shifts; Grewe et al., 2013; Gutiérrez and Wilson, 2021;

Sunday et al., 2015; Zografou et al., 2021). Tests of local adaptation in critical traits, like thermal

tolerances among larval life stages, would be valuable.

Given that species' traits did not predict temporal or geographic responses, it is

unsurprising that species' responses were also independent of phylogenetic history. The

phylogenetic approach did not improve model predictions in any model that we tested, and there

was no phylogenetic signal in either response according to Pagel's lambda and Blomberg's K

(Table 2). These results are consistent with previous work that found no phylogenetic trend in

local odonate population extinctions (Suhonen et al., 2022). There may be strong variation in

thermal niches among closely related species: species that are geographically isolated adapt to

different local climates, while populations that coexist may experience divergent selection within

their climate tolerances (Schuetz et al., 2019).

While more work is needed to determine if the range and phenology shifts reported here

are associated with increases in species' abundances, our results suggest that there are clear

'winners' and 'losers' under climate change. Climate 'winners', species that are shifting in space

and time, may require more limited conservation intervention. Species expanding their ranges

could be better supported if habitat area and connectivity are conserved, facilitating climate-

driven range shifts (Littlefield et al., 2019). Species only shifting their phenologies may require

further study, as phenology shifts may have positive or negative impacts on abundance (Iler et

al., 2021). Climate 'losers', species that are failing to shift in both space and time, may require

more direct conservation intervention, such as managed relocation (Richardson et al., 2009). Our

analysis of phenology and range shifts should be repeated in other taxa, as it may offer a method

of identifying conservation actions among species groups.

Here, we showed that odonate species exhibit convergent responses of range and

phenology shifts across continents. While species with southern distributions were more likely to

shift their ranges, increasing temperature variation limited geographical range responses among

species in both Europe and North America. Simultaneous consideration of shifts in range and

phenology is a powerful and necessary approach to measure species' apparent vulnerability to

rapid global changes. Understanding how range and phenology shifts vary across species, and

what drives this variation, is increasingly urgent as climate change alters local and regional

environmental conditions. This study gives us new insight into patterns of range and phenology

shifts, as well as suggesting processes behind the patterns, offering a valuable method of

understanding species' vulnerability to change that should be explored in other taxa.

References

Amano T, Freckleton RP, Queenborough SA, Doxford SW, Smithers RJ, Sparks TH, Sutherland

WJ. 2014. Links between plant species' spatial and temporal responses to a warming

climate. *Proceedings of the Royal Society B: Biological Sciences* **281**:20133017.

doi:10.1098/rspb.2013.3017

Angert AL, Crozier LG, Rissler LJ, Gilman SE, Tewksbury JJ, Chunco AJ. 2011. Do species'

traits predict recent shifts at expanding range edges? *Ecology Letters* **14**:677–689.

doi:10.1111/j.1461-0248.2011.01620.x

Ball-Damerow JE, M'Gonigle LK, Resh VH. 2014. Changes in occurrence, richness, and

biological traits of dragonflies and damselflies (Odonata) in California and Nevada over

the past century. *Biodivers Conserv* **23**:2107–2126. doi:10.1007/s10531-014-0707-5

Ball-Damerow JE, Oboyski PT, Resh VH. 2015. California dragonfly and damselfly (Odonata)

database: temporal and spatial distribution of species records collected over the past

century. *Zookeys* 67–89. doi:10.3897/zookeys.482.8453

Bartomeus I, Stavert JR, Ward D, Aguado O. 2018. Historical collections as a tool for assessing

the global pollination crisis. *Philosophical Transactions of the Royal Society B:*

Biological Sciences **374**:20170389. doi:10.1098/rstb.2017.0389

Beissinger SR, Riddell EA. 2021. Why Are Species' Traits Weak Predictors of Range Shifts?

Annual Review of Ecology, Evolution, and Systematics **52**:47–66. doi:10.1146/annurev-

ecolsys-012021-092849

Brooks SJ, Self A, Toloni F, Sparks T. 2014. Natural history museum collections provide

information on phenological change in British butterflies since the late-nineteenth

century. *Int J Biometeorol* **58**:1749–1758. doi:10.1007/s00484-013-0780-6

Buckley LB, Kingsolver JG. 2012. Functional and Phylogenetic Approaches to Forecasting

Species' Responses to Climate Change. *Annual Review of Ecology, Evolution, and*

Systematics **43**:205–226. doi:10.1146/annurev-ecolsys-110411-160516

Cannings R.A. 2002. Introducing the Dragonflies of British Columbia and Yukon. Royal British

Columbia Museum.

Cardillo M, Mace GM, Gittleman JL, Jones KE, Bielby J, Purvis A. 2008. The predictability of

extinction: biological and external correlates of decline in mammals. *Proceedings of the*

Royal Society B: Biological Sciences **275**:1441–1448. doi:10.1098/rspb.2008.0179

Chen I-C, Hill JK, Ohlemüller R, Roy DB, Thomas CD. 2011. Rapid Range Shifts of Species

Associated with High Levels of Climate Warming. *Science* **333**:1024–1026.

Conrad KF, Willson KH, Harvey IF, Thomas CJ, Sherratt TN. 1999. Dispersal characteristics of

seven odonate species in an agricultural landscape. *Ecography* **22**:524–531.

doi:10.1111/j.1600-0587.1999.tb01282.x

Cooper N, Bielby J, Thomas GH, Purvis A. 2008. Macroecology and extinction risk correlates of

frogs. *Global Ecology and Biogeography* **17**:211–221. doi:10.1111/j.1466-

8238.2007.00355.x

Davis MB, Shaw RG, Etterson JR. 2005. Evolutionary responses to changing climate. *Ecology*

86:1704–1714. doi:10.1890/03-0788

Day PB, Stuart-Smith RD, Edgar GJ, Bates AE. 2018. Species' thermal ranges predict changes

in reef fish community structure during 8 years of extreme temperature variation.

Diversity and Distributions **24**:1036–1046. doi:10.1111/ddi.12753

Devictor V, Julliard R, Couvet D, Jiguet F. 2008. Birds are tracking climate warming, but not

fast enough. *Proceedings of the Royal Society B: Biological Sciences* **275**:2743–2748.

doi:10.1098/rspb.2008.0878

Devictor V, van Swaay C, Brereton T, Brotons L, Chamberlain D, Heliölä J, Herrando S, Julliard

R, Kuussaari M, Lindström Å, Reif J, Roy DB, Schweiger O, Settele J, Stefanescu C,

Van Strien A, Van Turnhout C, Vermouzek Z, WallisDeVries M, Wynhoff I, Jiguet F.

2012. Differences in the climatic debts of birds and butterflies at a continental scale.

Nature Clim Change **2**:121–124. doi:10.1038/nclimate1347

Diamond SE, Frame AM, Martin RA, Buckley LB. 2011. Species' traits predict phenological

responses to climate change in butterflies. *Ecology* **92**:1005–1012. doi:10.1890/10-1594.1

ESRI. 2019. ArcGIS (Version 10.7.1).

Estrada A, Morales-Castilla I, Caplat P, Early R. 2016. Usefulness of Species Traits in Predicting

Range Shifts. *Trends in Ecology & Evolution* **31**:190–203.

doi:10.1016/j.tree.2015.12.014

Franks SE, Pearce-Higgins JW, Atkinson S, Bell JR, Botham MS, Brereton TM, Harrington R,

Leech DI. 2018. The sensitivity of breeding songbirds to changes in seasonal timing is

linked to population change but cannot be directly attributed to the effects of trophic

asynchrony on productivity. *Global Change Biology* **24**:957–971. doi:10.1111/gcb.13960

Fritz SA, Bininda-Emonds ORP, Purvis A. 2009. Geographical variation in predictors of

mammalian extinction risk: big is bad, but only in the tropics. *Ecology Letters* **12**:538–

549. doi:10.1111/j.1461-0248.2009.01307.x

Gaiji S, Chavan V, Ariño AH, Otegui J, Hobern D, Sood R, Robles E. 2013. Content assessment

of the primary biodiversity data published through GBIF network: Status, challenges and

potentials. *Biodiversity Informatics* **8**. doi:10.17161/bi.v8i2.4124

García-Roselló E, Guisande C, Manjarrés-Hernández A, González-Dacosta J, Heine J, Pelayo-

Villamil P, González-Vilas L, Vari RP, Vaamonde A, Granado-Lorencio C, Lobo JM.

2015. Can we derive macroecological patterns from primary Global Biodiversity

Information Facility data? *Global Ecology and Biogeography* **24**:335–347.

doi:10.1111/geb.12260

Goulson D. 2019. The insect apocalypse, and why it matters. *Current Biology* **29**:R967–R971.

doi:10.1016/j.cub.2019.06.069

Grant PR, Grant BR, Huey RB, Johnson MTJ, Knoll AH, Schmitt J. 2017. Evolution caused by

extreme events. *Philosophical Transactions of the Royal Society B: Biological Sciences*

372:20160146. doi:10.1098/rstb.2016.0146

Grewe Y, Hof C, Dehling DM, Brandl R, Brändle M. 2013. Recent range shifts of European

dragonflies provide support for an inverse relationship between habitat predictability and

dispersal. *Global Ecology and Biogeography* **22**:403–409. doi:10.1111/geb.12004

Gutiérrez D, Wilson RJ. 2021. Intra- and interspecific variation in the responses of insect

phenology to climate. *Journal of Animal Ecology* **90**:248–259. doi:10.1111/1365-

2656.13348

Hadfield JD. 2010. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The

MCMCglmm R Package. *Journal of Statistical Software* **33**:1–22.

doi:10.18637/jss.v033.i02

Hällfors MH, Heikkinen RK, Kuussaari M, Lehikoinen A, Luoto M, Pöyry J, Virkkala R,

Saastamoinen M, Kujala H. 2024. Recent range shifts of moths, butterflies, and birds are

driven by the breadth of their climatic niche. *Evolution Letters* **8**:89–100.

doi:10.1093/evlett/qrad004

Hällfors MH, Pöyry J, Heliölä J, Kohonen I, Kuussaari M, Leinonen R, Schmucki R, Sihvonen

P, Saastamoinen M. 2021. Combining range and phenology shifts offers a winning

strategy for boreal Lepidoptera. *Ecology Letters* **24**:1619–1632. doi:10.1111/ele.13774

Harris RMB, Beaumont LJ, Vance TR, Tozer CR, Remenyi TA, Perkins-Kirkpatrick SE,

Mitchell PJ, Nicotra AB, McGregor S, Andrew NR, Letnic M, Kearney MR, Wernberg

T, Hutley LB, Chambers LE, Fletcher M-S, Keatley MR, Woodward CA, Williamson G,

Duke NC, Bowman DMJS. 2018. Biological responses to the press and pulse of climate

trends and extreme events. *Nature Clim Change* **8**:579–587.

doi:10.1038/s41558-018-0187-9

Hassall C. 2015. Odonata as candidate macroecological barometers for global climate change.

Freshwater Science **34**:1040–1049. doi:10.1086/682210

Hassall C, Thompson DJ. 2008. The effects of environmental warming on Odonata: a review.

International Journal of Odonatology **11**:131–153. doi:10.1080/13887890.2008.9748319

Hassall C, Thompson DJ, French GC, Harvey IF. 2007. Historical changes in the phenology of

British Odonata are related to climate. *Global Change Biology* **13**:933–941.

doi:10.1111/j.1365-2486.2007.01318.x

Hickling R, Roy DB, Hill JK, Thomas CD. 2005. A northward shift of range margins in British

Odonata. *Global Change Biology* **11**:502–506. doi:10.1111/j.1365-2486.2005.00904.x

Hof C, Brändle M, Brandl R. 2006. Lentic odonates have larger and more northern ranges than

lotic species. *Journal of Biogeography* **33**:63–70. doi:10.1111/j.1365-2699.2005.01358.x

Iler AM, CaraDonna PJ, Forrest JRK, Post E. 2021. Demographic Consequences of Phenological

Shifts in Response to Climate Change. *Annual Review of Ecology, Evolution, and*

Systematics **52**:221–245. doi:10.1146/annurev-ecolsys-011921-032939

IPCC. 2021. Climate Change 2021: The Physical Science Basis. Contribution of Working Group

I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.

Cambridge University Press.

IUCN. 2021. The IUCN Red List of Threatened Species.

Jaeschke A, Bittner T, Reineking B, Beierkuhnlein C. 2013. Can they keep up with climate

change? – Integrating specific dispersal abilities of protected Odonata in species

distribution modelling. *Insect Conservation and Diversity* **6**:93–103. doi:10.1111/j.1752-

4598.2012.00194.x

Jones C.D., Kingsley A, Burke P, Holder M. 2008. Field guide to the dragonflies and damselflies

of Algonquin Provincial Park and the surrounding area, 1st ed. The Friends of the

Algonquin Park.

Jones Colin D., Kingsley A, Burke P, Holder M. 2008. Field guide to the dragonflies and

damselflies of Algonquin Provincial Park and the surrounding area, 1st ed. Whitney,

Ontario: The Friends of the Algonquin Park.

Kalkman VJ, Boudot J-P, Bernard R, De Knijf G, Suhling F, Termaat T. 2018. Diversity and

conservation of European dragonflies and damselflies (Odonata). *Hydrobiologia*

811:269–282. doi:10.1007/s10750-017-3495-6

Kerr JT. 2020. Racing against change: understanding dispersal and persistence to improve

species' conservation prospects. *Proceedings of the Royal Society B: Biological Sciences*

287:20202061. doi:10.1098/rspb.2020.2061

Kerr JT, Pindar A, Galpern P, Packer L, Potts SG, Roberts SM, Rasmont P, Schweiger O, Colla

SR, Richardson LL, Wagner DL, Gall LF, Sikes DS, Pantoja A. 2015. Climate change

impacts on bumblebees converge across continents. *Science* **349**:177–180.

doi:10.1126/science.aaa7031

Kharouba HM, Algar AC, Kerr JT. 2009. Historically calibrated predictions of butterfly species'

range shift using global change as a pseudo-experiment. *Ecology* **90**:2213–2222.

doi:10.1890/08-1304.1

Kujala H, Vepsäläinen V, Zuckerberg B, Brommer JE. 2013. Range margin shifts of birds

revisited – the role of spatiotemporally varying survey effort. *Global Change Biology*

19:420–430. doi:10.1111/gcb.12042

Lawlor JA, Comte L, Grenouillet G, Lenoir J, Baecher JA, Bandara RMWJ, Bertrand R, Chen I-

C, Diamond SE, Lancaster LT, Moore N, Murienne J, Oliveira BF, Pecl GT, Pinsky ML,

Rolland J, Rubenstein M, Scheffers BR, Thompson LM, van Amerom B, Villalobos F,

Weiskopf SR, Sunday J. 2024. Mechanisms, detection and impacts of species

redistributions under climate change. *Nat Rev Earth Environ* **5**:351–368.

doi:10.1038/s43017-024-00527-z

Lenoir J, Bertrand R, Comte L, Bourgeaud L, Hattab T, Murienne J, Grenouillet G. 2020.

Species better track climate warming in the oceans than on land. *Nat Ecol Evol* **4**:1044–

1059. doi:10.1038/s41559-020-1198-2

Leroux SJ, Larrivée M, Boucher-Lalonde V, Hurford A, Zuloaga J, Kerr JT, Lutscher F. 2013.

Mechanistic models for the spatial spread of species under climate change. *Ecological*

Applications **23**:815–828. doi:10.1890/12-1407.1

Littlefield CE, Krosby M, Michalak JL, Lawler JJ. 2019. Connectivity for species on the move:

supporting climate-driven range shifts. *Frontiers in Ecology and the Environment*

17:270–278. doi:10.1002/fee.2043

Lustenhouwer N, Wilschut RA, Williams JL, van der Putten WH, Levine JM. 2018. Rapid

evolution of phenology during range expansion with recent climate change. *Global*

Change Biology **24**:e534–e544. doi:10.1111/gcb.13947

Macgregor CJ, Thomas CD, Roy DB, Beaumont MA, Bell JR, Brereton T, Bridle JR, Dytham C,

Fox R, Gotthard K, Hoffmann AA, Martin G, Middlebrook I, Nylin S, Platts PJ, Rasteiro

R, Saccheri IJ, Villoutreix R, Wheat CW, Hill JK. 2019. Climate-induced phenology

shifts linked to range expansions in species with multiple reproductive cycles per year.

Nat Commun **10**:4455. doi:10.1038/s41467-019-12479-w

MacLean SA, Beissinger SR. 2017. Species' traits as predictors of range shifts under

contemporary climate change: A review and meta-analysis. *Global Change Biology*

23:4094–4105. doi:10.1111/gcb.13736

Mair L, Hill JK, Fox R, Botham M, Brereton T, Thomas CD. 2014. Abundance changes and

habitat availability drive species' responses to climate change. *Nature Clim Change*

4:127–131. doi:10.1038/nclimate2086

Mccauley SJ, Hammond JI, Frances DN, Mabry KE. 2015. Effects of experimental warming on

survival, phenology, and morphology of an aquatic insect (Odonata). *Ecological*

Entomology **40**:211–220. doi:10.1111/een.12175

McCauley SJ, Hammond JI, Mabry KE. 2018. Simulated climate change increases larval

mortality, alters phenology, and affects flight morphology of a dragonfly. *Ecosphere*

9:e02151. doi:10.1002/ecs2.2151

Menzel A, Sparks TH, Estrella N, Koch E, Aasa A, Ahas R, Alm-Kübler K, Bissolli P,

Braslavská O, Briede A, Chmielewski FM, Crepinsek Z, Curnel Y, Dahl Å, Defila C,

Donnelly A, Filella Y, Jatczak K, Måge F, Mestre A, Nordli Ø, Peñuelas J, Pirinen P,

Remišová V, Scheifinger H, Striz M, Susnik A, Van Vliet AJH, Wielgolaski F-E, Zach S,

Zust A. 2006. European phenological response to climate change matches the warming

pattern. *Global Change Biology* **12**:1969–1976. doi:10.1111/j.1365-2486.2006.01193.x

Nagelkerke NJD. 1991. A Note on a General Definition of the Coefficient of Determination.

Biometrika **78**:691–692. doi:10.2307/2337038

New M, Lister D, Hulme M, Makin I. 2002. A high-resolution data set of surface climate over

global land areas. *Climate Research* **21**:1–25. doi:10.3354/cr021001

Pacifici M, Rondinini C, Rhodes JR, Burbidge AA, Cristiano A, Watson JEM, Woinarski JCZ,

Di Marco M. 2020. Global correlates of range contractions and expansions in terrestrial

mammals. *Nat Commun* **11**:2840. doi:10.1038/s41467-020-16684-w

Parmesan C, Ryrholm N, Stefanescu C, Hill JK, Thomas CD, Descimon H, Huntley B, Kaila L,

Kullberg J, Tammaru T, Tennent WJ, Thomas JA, Warren M. 1999. Poleward shifts in

geographical ranges of butterfly species associated with regional warming. *Nature*

399:579–583. doi:10.1038/21181

Parmesan C, Yohe G. 2003. A globally coherent fingerprint of climate change impacts across

natural systems. *Nature* **421**:37–42. doi:10.1038/nature01286

Paulson D. 2012. Dragonflies and Damselflies of the East. Princeton University Press.

Pearse WD, Davis CC, Inouye DW, Primack RB, Davies TJ. 2017. A statistical estimator for

determining the limits of contemporary and historic phenology. *Nat Ecol Evol* **1**:1876–

1882. doi:10.1038/s41559-017-0350-0

Pinsky ML, Worm B, Fogarty MJ, Sarmiento JL, Levin SA. 2013. Marine Taxa Track Local

Climate Velocities. *Science* **341**:1239–1242. doi:10.1126/science.1239352

Platts PJ, Mason SC, Palmer G, Hill JK, Oliver TH, Powney GD, Fox R, Thomas CD. 2019.

Habitat availability explains variation in climate-driven range shifts across multiple

taxonomic groups. *Sci Rep* **9**:15039. doi:10.1038/s41598-019-51582-2

Powney G, Brooks S, Barwell L, Bowles P, Fitt R, Pavitt A, Spriggs R, Isaac N. 2014.

Morphological and geographical traits of the British Odonata. *Biodiversity Data Journal*

2:e1041. doi:10.3897/bdj.2.e1041

Powney GD, Brooks SJ, Barwell LJ, Bowles P, Fitt RNL, Pavitt A, Spriggs RA, Isaac NJB.

2014. Morphological and Geographical Traits of the British Odonata. *Biodivers Data J*

e1041. doi:10.3897/BDJ.2.e1041

Powney GD, Cham SSA, Smallshire D, Isaac NJB. 2015. Trait correlates of distribution trends in

the Odonata of Britain and Ireland. *PeerJ* **3**:e1410. doi:10.7717/peerj.1410

Pyke GH, Ehrlich PR. 2010. Biological collections and ecological/environmental research: a

review, some observations and a look to the future. *Biological Reviews* **85**:247–266.

doi:10.1111/j.1469-185X.2009.00098.x

R Core Team. 2019. R: A language and environment for statistical computing.

Rapacciuolo G, Ball-Damerow JE, Zeilinger AR, Resh VH. 2017. Detecting long-term

occupancy changes in Californian odonates from natural history and citizen science

records. *Biodivers Conserv* **26**:2933–2949. doi:10.1007/s10531-017-1399-4

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other

things). *Methods in Ecology and Evolution* **3**:217–223.

doi:10.1111/j.2041-210X.2011.00169.x

Richardson DM, Hellmann JJ, McLachlan JS, Sax DF, Schwartz MW, Gonzalez P, Brennan EJ,

Camacho A, Root TL, Sala OE, Schneider SH, Ashe DM, Clark JR, Early R, Etterson JR,

Fielder ED, Gill JL, Minteer BA, Polasky S, Safford HD, Thompson AR, Vellend M,

Ehrlich PR. 2009. Multidimensional Evaluation of Managed Relocation. *Proceedings of*

the National Academy of Sciences of the United States of America **106**:9721–9724.

Robbirt KM, Davy AJ, Hutchings MJ, Roberts DL. 2011. Validation of biological collections as

a source of phenological data for use in climate change studies: a case study with the

orchid Ophrys sphegodes. *Journal of Ecology* **99**:235–241.

doi:10.1111/j.1365-2745.2010.01727.x

Román-Palacios C, Wiens JJ. 2020. Recent responses to climate change reveal the drivers of

species extinction and survival. *Proceedings of the National Academy of Sciences*

117:4211–4217. doi:10.1073/pnas.1913007117

Rundle SD, Bilton DT, Abbott JC, Foggo A. 2007. Range size in North American Enallagma

damselflies correlates with wing size. *Freshwater Biology* **52**:471–477.

doi:10.1111/j.1365-2427.2006.01712.x

Saino N, Ambrosini R, Rubolini D, von Hardenberg J, Provenzale A, Hüppop K, Hüppop O,

Lehikoinen A, Lehikoinen E, Rainio K, Romano M, Sokolov L. 2010. Climate warming,

ecological mismatch at arrival and population decline in migratory birds. *Proceedings of*

the Royal Society B: Biological Sciences **278**:835–842. doi:10.1098/rspb.2010.1778

Schuetz JG, Mills KE, Allyn AJ, Stamieszkin K, Bris AL, Pershing AJ. 2019. Complex patterns

of temperature sensitivity, not ecological traits, dictate diverse species responses to

climate change. *Ecography* **42**:111–124. doi:10.1111/ecog.03823

Sirois-Delisle C, Kerr JT. 2022. Climate change aggravates non-target effects of pesticides on

dragonflies at macroecological scales. *Ecological Applications* **32**:e2494.

doi:10.1002/eap.2494

Socolar JB, Epanchin PN, Beissinger SR, Tingley MW. 2017. Phenological shifts conserve

thermal niches in North American birds and reshape expectations for climate-driven

range shifts. *Proceedings of the National Academy of Sciences* **114**:12976–12981.

doi:10.1073/pnas.1705897114

Souza KS, Fortunato DS, Jardim L, Terribile LC, Lima-Ribeiro MS, Mariano CÁ, Pinto-

Ledezma JN, Loyola R, Dobrovolski R, Rangel TF, Machado IF, Rocha T, Batista MG,

Lorini ML, Vale MM, Navas CA, Maciel NM, Villalobos F, Olalla-Tarraga MÂ,

Rodrigues JFM, Gouveia SF, Diniz-Filho JAF. 2023. Evolutionary rescue and geographic

range shifts under climate change for global amphibians. *Front Ecol Evol* **11**.

doi:10.3389/fevo.2023.1038018

Srivastava DS, Coristine L, Angert AL, Bontrager M, Amundrud SL, Williams JL, Yeung ACY,

de Zwaan DR, Thompson PL, Aitken SN, Sunday JM, O'Connor MI, Whitton J, Brown

NEM, MacLeod CD, Parfrey LW, Bernhardt JR, Carrillo J, Harley CDG, Martone PT,

Freeman BG, Tseng M, Donner SD. 2021. Wildcards in climate change biology.

Ecological Monographs **91**:e01471. doi:10.1002/ecm.1471

Stephens PA, Mason LR, Green RE, Gregory RD, Sauer JR, Alison J, Aunins A, Brotons L,

Butchart SHM, Campedelli T, Chodkiewicz T, Chylarecki P, Crowe O, Elts J, Escandell

V, Foppen RPB, Heldbjerg H, Herrando S, Husby M, Jiguet F, Lehikoinen A, Lindström

Å, Noble DG, Paquet J-Y, Reif J, Sattler T, Szép T, Teufelbauer N, Trautmann S, van

Strien AJ, van Turnhout CAM, Vorisek P, Willis SG. 2016. Consistent response of bird

populations to climate change on two continents. *Science* **352**:84–87.

doi:10.1126/science.aac4858

Suhonen J, Ilvonen JJ, Korkeamäki E, Nokkala C, Salmela J. 2022. Using functional traits and

phylogeny to understand local extinction risk in dragonflies and damselflies (Odonata).

Ecology and Evolution **12**:e8648. doi:10.1002/ece3.8648

Sunday JM, Pecl GT, Frusher S, Hobday AJ, Hill N, Holbrook NJ, Edgar GJ, Stuart-Smith R,

Barrett N, Wernberg T, Watson RA, Smale DA, Fulton EA, Slawinski D, Feng M,

Radford BT, Thompson PA, Bates AE. 2015. Species traits and climate velocity explain

geographic range shifts in an ocean-warming hotspot. *Ecology Letters* **18**:944–953.

doi:10.1111/ele.12474

Urban MC. 2015. Accelerating extinction risk from climate change. *Science* **348**:571–573.

doi:10.1126/science.aaa4984

Urban MC, Travis JMJ, Zurell D, Thompson PL, Synes NW, Scarpa A, Peres-Neto PR,

Malchow A-K, James PMA, Gravel D, De Meester L, Brown C, Bocedi G, Albert CH,

Gonzalez A, Hendry AP. 2022. Coding for Life: Designing a Platform for Projecting and

Protecting Global Biodiversity. *BioScience* **72**:91–104. doi:10.1093/biosci/biab099

van der Laken P. 2021. ppsr: Predictive Power Score.

Waller JT, Svensson EI. 2017. Body size evolution in an old insect order: No evidence for

Cope's Rule in spite of fitness benefits of large size. *Evolution* **71**:2178–2193.

doi:10.1111/evo.13302

Waller JT, Willink B, Tschol M, Svensson EI. 2019b. The odonate phenotypic database, a new

open data resource for comparative studies of an old insect order. *Scientific Data* **6**:1–6.

doi:10.1038/s41597-019-0318-9

Zattara EE, Aizen MA. 2021. Worldwide occurrence records suggest a global decline in bee

species richness. *One Earth* **4**:114–123. doi:10.1016/j.oneear.2020.12.005

Zografou K, Swartz MT, Adamidis GC, Tilden VP, McKinney EN, Sewall BJ. 2021. Species

traits affect phenological responses to climate change in a butterfly community. *Sci Rep*

11:3283. doi:10.1038/s41598-021-82723-1

Table 1: Fixed effects estimates and associated statistics from the generalized linear model and

generalized mixed effects model (accounting for phylogeny; for credible intervals, see Table S3)

of the relationship between range shifts and emergence phenology change. The continent term

shows effects of the North American continent compared to the European continent as the

reference level. N gives the number of species involved in the model, and an asterisk indicates

statistical significance of the variable in question (p-value < 0.05). The pseudo R^2 type is

Nagelkerke (Nagelkerke, 1991).

695
696

	Phenology shift (N = 66)			
	GLM		**MCMCglmm**	
Predictors	**Est.**	**p**	**Post.m**	**p**
(intercept)	0.12	0.53	0.12	-0.49
Range shift	-0.45	<0.01*	-0.45	<0.01*
Continent	-0.22	0.44	-0.22	0.42
Model evaluation				
AIC/DIC	185.39		185.43	
Null model	193.13		193.12	
Pseudo R^2	17.08%		14.90%	

Table 2: Fixed effects estimates and associated statistics from the generalized linear model and

generalized mixed effects model (accounting for phylogeny; for credible intervals, see Table S3)

of drivers of odonate range shifts. N indicates the number of modelled species, an asterisk

indicates statistical significance of the variable in question, and a dash symbol shows that the

variable was excluded from the final model. The pseudo R^2 type is Nagelkerke (Nagelkerke,

1991). For the categorical variables breeding habitat type and range geography, we used lotic

habitat type and Northern range as reference levels, respectively. Phenology shifts were unrelated

to any of the predictor variables that we tested.

	Range shift (N = 76)			
	GLM		MCMCglmm	
Predictors	**Est.**	**p**	**Post.m**	**p**
(intercept)	-0.61	0.03*	-0.62	0.04*
Southern range	0.90	<0.01*	0.91	<0.01*
Widespread	0.32	0.34	0.32	0.36
T ° variability	-0.34	<0.01*	-0.34	<0.01*
Both hab. Types	-	-	-	-
Lentic habitat	-	-	-	-
Flight duration	-	-	-	-
Oviposition	-	-	-	-
Body size	-	-	-	-
Model evaluation				
AIC/DIC		208.99		209.08
Null model		221.51		221.49
Pseudo R^2		22.72%		20.18%
Phylogenetic signal				
Pagel's lambda (p)		0.02 (0.63)		
Blomberg's K (p)		0.08 (0.77)		

Figure 1: Representation of temporal and geographical limits characterising the ecological niche

of a hypothetical odonate species. Points show 250 individuals according to their Julian day of

emergence, latitudinal position, and temperatures to which each individual is exposed. **(A)** Points

represent historical observations, triangle symbols show observations following a shift towards

higher latitudes, and plus signs show observations following a shift towards earlier emergence

dates, necessary to maintain temperatures to which species were exposed historically, after

temperatures warmed across the species' range. Warm and cool colors show hot and cold

temperatures, respectively. **(B)** Grey points represent historical observations, and the colored

points show observations after warming without allowing shifts through time or space. In this

scenario, we assumed that the hottest temperature tolerated by the species corresponds to the

hottest temperature experienced in its historical niche. In this case, individuals are exposed to

hotter temperatures overall while declining in areas outside species' thermal tolerances after

warming.

750 **Figure 2**: Richness of 76 odonate species sampled in North America and Europe between 1980

751 and 2018 per 100 X 100 km quadrat. Dark red indicates high species richness, while light pink

752 indicates low species richness.

Figure 3: Relationship between range shifts and emergence phenology shifts among North

American and European odonate species (N = 66). The shaded area shows mean latitudinal range

shifts of terrestrial taxa (Lenoir et al., 2020), calculated as the yearly mean dispersal rate of 1.11

+/- 0.96 km per year over 38 years.

Supplementary Information

<u>Odonate data</u>

The full raw dataset includes ~2 million Odonata occurrence records. We first removed

inadequate records from our primary dataset. Inadequate records were identified as those with

incomplete information for species identification, year, or locality, inaccurate georeferenced

points, and duplicate records. We retained ~1,100,000 unique species-location-date observations.

There are 452,929 records in the first time period (1980 – 2002) and 641,590 observations in the

second time period (2008 – 2018).

If a species was found on both continents, we only kept observations from the continent

that was the most densely sampled. This simplified analyses in terms of calculating range and

phenology shifts. It would not be logical to perform a cross-continental comparison when data

points for a single species exist in both places and are merged together to create a single

measurement of range shifts or location-specific measures of phenological change. This problem

cannot be solved by treating species as different units because that leads to uninterpretable

outcomes (and creation of pseudo-replication) in the context of phylogenetic analyses. Most

importantly, it was not possible to calculate phenology shifts for both continents in these species

because of the data intensity needed to make the calculations and the strict criteria we set for

inclusion of species and quadrats in the analysis. Future work may focus on species found on

both continents if the data and statistical analyses permit.

<u>Phenology and range position estimates</u>

We have not tried to interpret noise in phenology shift estimates, which could occur due to

sampling issues, or to another biologically meaningful eco-evolutionary response. Our methods

do not enable tests of those ideas, however. For example, *Aeshna umbrosa* has a surprisingly

high phenology shift (>25 days) but a very strong range retraction (<-300 km). For this species,

we retained 8 quadrats to calculate mean phenology shift estimate. It is possible that a section of

the range was lost in which emergence was especially early, compared to other regions, due to

local adaptations. This pattern may affect results where species appear to shift emergence dates

earlier/later, but phenology shifts are affected by parts of the range that remain occupied.

We put in place criteria to make sure to include species with range shifts likely to result

from climate change effects, rather than from sampling issues or to other anthropogenic

pressures such as land use change, or sudden land use intensification. *Nehalennia irene* was thus

removed, as its range positions were highly unusual, and likely due to other factors than global

change (>800 km). *Libellula quadrimaculata* was removed due to sampling intensity

discrepancies between time periods, having over 10,000 extra points in the second period

compared to the first.

Temporal and spatial bias are likely to be present in opportunistic data, but they are less

likely to impact long term factors of species' distributions if including data from as many sources

as possible, and that span across large geographic and temporal scopes (Pyke and Ehrlich, 2010;

Zattara and Aizen, 2021). Here, we put in place several criteria in careful interpretation of

results, as described in the Methods section of the main text. Further, among preliminary

examinations of the data, we test for the effect of sampling onto our phenology and range shift

estimates. We found no effect of sampling on range of phenology responses that we calculated

here. We tested a linear mixed model with species as a random effect that assesses whether

sampling intensity change per quadrat (where sampling intensity corresponds to sampling per

species/quadrat) predicts species' phenology change per quadrat. The p-value of the predictor

variable, sampling intensity change, was not significant. Secondly, we built a Generalized Linear

Model to assess whether greater sampling overall (measured as the number of quadrats in which

each species is sampled) predicts species' range shifts that we computed. The p-value of the

predictor variable, sampling change per species, was not significant. If spatial bias was present

here, we may expect the number of quadrats in which species are found to increase the position

of species' northern range boundaries.

<u>Model information and statements</u>

We report the full model statements below as executed with the *MCMCglmm* R package,

including the priors, and the model of trait evolution to account for phylogeny. Prior structure

followed standard practice according to continuous or binary response variable. The family

"gaussian" or "threshold" was selected depending on the continuous or binary nature of the data,

respectively. The number of iterations, burnin, and thinning parameters were tested and

confirmed with a visual assessment of model convergence using the trace and density plots. It is

common practice for the burnin parameter to correspond to 10% of the number of iterations.

Model 1

Aphylo ← vcv(phylogeny, model = "Brownian", corr = T)

Ainv ← inverseA(phylogeny, nodes = "TIPS", scale = F)$Ainv

prior ← list(R=list(V=1,nu=0.002))

MCMCglmm(Phenological shifts ~ Range shifts + Continent, ginverse = list(species = Ainv),

family = "gaussian", prior = prior, DIC = T, nitt = 150,000, thin = 100, burnin = 15,000, data)

Model 2

Aphylo ← vcv(phylogeny, model = "Brownian", corr = T)

Ainv ← inverseA(phylogeny, nodes = "TIPS", scale = F)$Ainv

prior ← list(R=list(V=1,nu=0.002, fix = 1))

MCMCglmm(Range shifts ~ Breeding habitat type + Distribution region + Flight duration + Egg

laying habitat + Body size + Temperature variability, ginverse = list(species = Ainv), family =

"threshold", prior = prior, DIC = T, nitt = 150,000, thin = 100, burnin = 15,000, data)

Model 3

Aphylo ← vcv(phylogeny, model = "Brownian", corr = T)

Ainv ← inverseA(phylogeny, nodes = "TIPS", scale = F)$Ainv

prior ← list(R=list(V=1,nu=0.002))

MCMCglmm(Phenology shift ~ Breeding habitat type + Distribution region + Flight duration +

Egg laying habitat + Body size + Temperature variability, ginverse = list(species = Ainv), family

= "gaussian", prior = prior, DIC = T, nitt = 150,000, thin = 100, burnin = 15,000, data)

Table S1: 76 species sampled across North America and Europe between 1980 and 2018

followed our criteria for quality observation records for inclusion in our analysis of geographical

shifts. Species Northern Range Limits (NRL) are shown in this table, as well as range limit

shifts. All range limit values are shown in kilometers from the equator. We used the 10 most

northern points of sampling in each time period to identify species' NRL, as detailed in the

Methods section of the main text.

Species	Continent	NRL (1980 - 2002)	NRL (2008 - 2018)	NRL shift
Aeshna constricta	North-America	5500.81	5692.6	191.79
Aeshna cyanea	Europe	6880.39	7018.26	137.87
Aeshna eremita	North-America	7527.81	7394.66	-133.16
Aeshna grandis	Europe	7504.84	7607.68	102.83
Aeshna interrupta	North-America	7235.65	7286.83	51.18
Aeshna juncea	Europe	7747.56	7812.16	64.6
Aeshna mixta	Europe	6566.18	6776.3	210.12
Aeshna umbrosa	North-America	6691.19	6209.63	-481.55
Anax imperator	Europe	6263.92	6548.57	284.65
Anax junius	North-America	5539.31	5666.11	126.8
Argia apicalis	North-America	4922.17	5219.38	297.22
Argia fumipennis	North-America	5235.2	5168.12	-67.08
Argia moesta	North-America	5533.74	5301.98	-231.77
Basiaeschna janata	North-America	5649.43	5655.77	6.34
Brachytron pratense	Europe	6737.42	6905.46	168.04
Calopteryx maculata	North-America	5329.56	5359.13	29.57
Calopteryx virgo	Europe	7275.9	7549.35	273.45
Ceriagrion tenellum	Europe	5900.55	5950.63	50.08
Coenagrion hastulatum	Europe	7528.94	7651.51	122.57
Coenagrion mercuriale	Europe	5913.57	5867.06	-46.52
Coenagrion puella	Europe	6904.7	6869.58	-35.12
Coenagrion pulchellum	Europe	7076.45	7152.81	76.36
Coenagrion resolutum	North-America	7516.44	7408.7	-107.73
Cordulegaster boltonii	Europe	7237.43	7346.93	109.49
Cordulia aenea	Europe	7400.12	7434.02	33.9
Cordulia shurtleffii	North-America	7499.82	7421.98	-77.84
Enallagma antennatum	North-America	5037.73	5347.01	309.28

Enallagma basidens	North-America	4768.61	4850.05	81.44
Enallagma carunculatum	North-America	5950.86	5592.47	-358.38
Enallagma civile	North-America	5322.59	5473.78	151.19
Enallagma cyathigerum	Europe	7594.97	7733.06	138.09
Enallagma ebrium	North-America	6341.31	5907.69	-433.62
Enallagma exsulans	North-America	5655.94	5253.94	-402
Enallagma hageni	North-America	6113.97	5865.52	-248.45
Enallagma signatum	North-America	5208.92	5281.09	72.17
Epitheca cynosura	North-America	5541.19	5582.56	41.37
Erythemis simplicicollis	North-America	5271	5292.65	21.65
Erythromma najas	Europe	7171.87	7372.13	200.25
Gomphus vulgatissimus	Europe	6890.76	7010.12	119.37
Hetaerina americana	North-America	5044.41	5244.49	200.08
Ischnura cervula	North-America	5973.57	5526.83	-446.74
Ischnura hastata	North-America	4725.07	4897.71	172.65
Ischnura perparva	North-America	5608.92	5438.53	-170.4
Ischnura posita	North-America	5084.08	5143.79	59.71
Ischnura pumilio	Europe	6213.77	6769.62	555.85
Ischnura verticalis	North-America	5579.68	5750.9	171.22
Ladona julia	North-America	5908.68	5871.84	-36.83
Lestes congener	North-America	6240.48	5803.75	-436.74
Lestes disjunctus	North-America	7294.75	7314.26	19.51
Lestes dryas	Europe	6824.89	7108.33	283.44
Lestes rectangularis	North-America	5263.85	5552.64	288.79
Lestes unguiculatus	North-America	5803.91	6030.62	226.71
Leucorrhinia dubia	Europe	7675.72	7749.01	73.29
Leucorrhinia hudsonica	North-America	7500.27	7446.85	-53.42
Libellula depressa	Europe	6810.85	7015.37	204.52
Libellula fulva	Europe	6602.18	6882.5	280.32
Libellula luctuosa	North-America	5337.86	5320.1	-17.76
Libellula pulchella	North-America	5556.31	5692.14	135.83
Orthetrum coerulescens	Europe	6744.55	6903.01	158.46
Pachydiplax longipennis	North-America	5495.02	5467.11	-27.92
Pantala flavescens	North-America	5190.25	5510.15	319.9
Perithemis tenera	North-America	4929.38	5192.47	263.09
Plathemis lydia	North-America	5539.41	5594.8	55.38
Platycnemis pennipes	Europe	6934.22	7125.72	191.5
Pyrrhosoma nymphula	Europe	7131.07	7313.37	182.3
Rhionaeschna californica	North-America	5641.53	5458.21	-183.32
Rhionaeschna multicolor	North-America	5581	5477.34	-103.66

Somatochlora metallica	Europe	7718.67	7743.21	24.54
Somatochlora semicircularis	North-America	6533.02	5994.13	-538.89
Sympetrum corruptum	North-America	5585.68	5733.2	147.52
Sympetrum danae	Europe	7268.54	7565.29	296.75
Sympetrum internum	North-America	7179.45	7121.38	-58.07
Sympetrum pallipes	North-America	5655.65	5527.84	-127.8
Sympetrum sanguineum	Europe	6663.43	6970.56	307.13
Sympetrum striolatum	Europe	6972.22	7098.74	126.52
Sympetrum vulgatum	Europe	6897.33	7274.78	377.45

855 **Table S2:** 66 species sampled across North America and Europe between 1980 and 2018

856 followed our criteria for quality observation records for inclusion in our analysis of emergence

857 phenology shifts. Mean phenological shifts (PS) is measured in the number of Julian days

858 comparing both time periods, as estimated using the Weibull distribution (See Methods). We also

859 report the number of 200 X 200 quadrats used to calculate phenology estimates per species.

Species	Number of quadrats	Mean PS
Aeshna cyanea	37	-8.05
Aeshna grandis	43	6.73
Aeshna juncea	34	-1.11
Aeshna mixta	22	-12.95
Aeshna umbrosa	8	27.42
Anax imperator	27	4.05
Anax junius	19	-12.21
Argia fumipennis	8	-3.39
Argia moesta	11	-7.62
Basiaeschna janata	2	-4.69
Brachytron pratense	24	-8.18
Calopteryx maculata	15	-4.14
Calopteryx virgo	41	4.35
Ceriagrion tenellum	14	-3.88
Coenagrion hastulatum	20	-4.97
Coenagrion mercuriale	11	5.80
Coenagrion puella	33	-1.00
Coenagrion pulchellum	29	1.94
Coenagrion resolutum	2	0.33
Cordulegaster boltonii	34	4.08
Cordulia aenea	31	-7.72
Cordulia shurtleffii	4	1.78
Enallagma basidens	2	0.52
Enallagma carunculatum	8	5.10
Enallagma civile	11	1.82
Enallagma cyathigerum	46	-9.81
Enallagma ebrium	7	-1.34
Enallagma exsulans	8	-2.83
Enallagma hageni	5	-4.79
Enallagma signatum	6	1.51

Epitheca cynosura	4	-8.82
Erythemis simplicicollis	15	0.01
Erythromma najas	35	-4.82
Gomphus vulgatissimus	12	-9.14
Hetaerina americana	6	-10.15
Ischnura cervula	3	10.65
Ischnura perparva	2	8.85
Ischnura posita	12	-5.55
Ischnura pumilio	14	-0.97
Ischnura verticalis	16	2.39
Ladona julia	9	4.29
Lestes congener	9	6.03
Lestes disjunctus	4	10.48
Lestes dryas	6	5.13
Lestes rectangularis	12	0.81
Leucorrhinia dubia	18	-6.05
Leucorrhinia hudsonica	4	-13.17
Libellula depressa	30	-6.13
Libellula fulva	16	-2.50
Libellula luctuosa	15	-5.11
Libellula pulchella	14	-6.43
Orthetrum coerulescens	22	4.11
Pachydiplax longipennis	13	-3.84
Perithemis tenera	5	-13.60
Plathemis lydia	14	0.30
Platycnemis pennipes	30	-0.40
Pyrrhosoma nymphula	39	-19.40
Rhionaeschna californica	2	-8.41
Rhionaeschna multicolor	4	-11.68
Somatochlora metallica	31	7.91
Sympetrum danae	32	-3.34
Sympetrum internum	5	3.05
Sympetrum pallipes	3	9.67
Sympetrum sanguineum	28	-13.30
Sympetrum vulgatum	16	-13.56

860
861

Table S3: Ecological and geographical traits of 76 North American and European odonate

species used in this work. Field guides (Cannings, 2002; Jones et al., 2008; Paulson, 2012) and

existing trait databases (Powney et al., 2014; Waller et al., 2019) were used to build this dataset.

Habitat type represents species' breeding habitat, and can have a value of lentic, lotic, or both

types. Distribution shows the general geographic position of each species' range, which can be

widespread (W), southern (S), northern (N), southern and widespread (SW), or northern and

widespread (NW). Oviposition type corresponds to egg laying inside plants (endophytic) as

opposed to directly in water or on plants (exophytic). Body size is measured as body length in

mm. In the case that body length was given as a maximum and minimum value, we used the

average of both values.

Species	Habitat	Distribution	Flight	Oviposition	Body size
Aeshna constricta	Both	W	2.5	Endophytic	69
Aeshna cyanea	Lentic	S	3	Endophytic	71.5
Aeshna eremita	Both	N-W	3	Endophytic	72.5
Aeshna grandis	Both	N	4	Endophytic	73.5
Aeshna interrupta	Both	W	3	Endophytic	66.5
Aeshna juncea	Lentic	N	4.5	Endophytic	75.5
Aeshna mixta	Both	S	2.5	Endophytic	60
Aeshna umbrosa	Both	W	3	Endophytic	68.5
Anax imperator	Lentic	S	2.5	Endophytic	76.5
Anax junius	Both	S-W	7	Endophytic	74
Argia apicalis	Both	S-W	4	Endophytic	36.5
Argia fumipennis	Both	S-W	3.5	Endophytic	31.5
Argia moesta	Lotic	S-W	3	Endophytic	39.5
Basiaeschna janata	Both	W	1	Endophytic	59
Brachytron pratense	Lentic	S	2	Endophytic	58.5
Calopteryx maculata	Lotic	S-W	3	Endophytic	48
Calopteryx virgo	Lotic	W	3.5	Endophytic	47
Ceriagrion tenellum	Both	S	3	Endophytic	30
Coenagrion hastulatum	Lentic	N	2.5	Endophytic	32
Coenagrion mercuriale	Lotic	S	2.5	Endophytic	29.5
Coenagrion puella	Lentic	S	3.5	Endophytic	34

Coenagrion pulchellum	Both	S	2	Endophytic	36
Coenagrion resolutum	Lentic	N-W	2.5	Endophytic	30
Cordulegaster boltonii	Lotic	W	3.5	Endophytic	77
Cordulia aenea	Lentic	S	2	Exophytic	51
Cordulia shurtleffii	Both	N-W	2	Exophytic	46
Enallagma antennatum	Both	S-W	2	Endophytic	30
Enallagma basidens	Both	S-W	4.5	Endophytic	24.5
Enallagma carunculatum	Both	W	3.5	Endophytic	31.5
Enallagma civile	Both	S-W	3.5	Endophytic	33.5
Enallagma cyathigerum	Both	W	3.5	Endophytic	32
Enallagma ebrium	Both	N-W	2.5	Endophytic	31
Enallagma exsulans	Both	S-W	3.5	Endophytic	34
Enallagma hageni	Both	N-W	2.5	Endophytic	30
Enallagma signatum	Both	S-W	2.5	Endophytic	32.5
Epitheca cynosura	Both	S-W	2.5	Endophytic	40.5
Erythemis simplicicollis	Both	S-W	2.5	Endophytic	41
Erythromma najas	Both	S	3	Endophytic	33
Gomphus vulgatissimus	Lotic	S	2	Exophytic	47.5
Hetaerina americana	Lotic	S-W	5	Endophytic	42
Ischnura cervula	Both	W	6	Endophytic	27.5
Ischnura hastata	Lentic	S-W	3.5	Endophytic	24
Ischnura perparva	Both	S-W	5	Endophytic	26.5
Ischnura posita	Both	S-W	3.5	Endophytic	25
Ischnura pumilio	Lentic	S	3	Endophytic	29
Ischnura verticalis	Both	W	3.5	Endophytic	26.5
Ladona julia	Lentic	N-W	2	Endophytic	41.5
Lestes congener	Lentic	W	2.5	Endophytic	36.5
Lestes disjunctus	Both	W	2.5	Endophytic	37.5
Lestes dryas	Lentic	S	2	Endophytic	37.5
Lestes rectangularis	Both	S-W	2.5	Endophytic	45
Lestes unguiculatus	Both	W	3	Endophytic	37.5
Leucorrhinia dubia	Lentic	N	2.5	Exophytic	33.5
Leucorrhinia hudsonica	Lentic	N-W	2	Endophytic	29.5
Libellula depressa	Lentic	S	2.5	Exophytic	43.5
Libellula fulva	Lentic	S	2	Exophytic	43.5
Libellula luctuosa	Lentic	S-W	2.5	Exophytic	46
Libellula pulchella	Both	W	2.5	Endophytic	54.5
Orthetrum coerulescens	Lotic	S	3	Exophytic	40.5
Pachydiplax longipennis	Both	S-W	3	Endophytic	35.5
Pantala flavescens	Both	S	3	Exophytic	48.5

Perithemis tenera	Both	S-W	4.5	Exophytic	22.5
Plathemis lydia	Both	S-W	3	Exophytic	45
Platycnemis pennipes	Lotic	S	2.5	Endophytic	36
Pyrrhosoma nymphula	Lentic	W	1.5	Endophytic	34.5
Rhionaeschna californica	Lentic	S-W	4	Endophytic	60.5
Rhionaeschna multicolor	Both	S-W	2	Endophytic	67
Somatochlora metallica	Both	W	1.5	Exophytic	53
Somatochlora semicircularis	Lentic	W	2	Exophytic	49.5
Sympetrum corruptum	Both	S-W	5	Exophytic	40.5
Sympetrum danae	Lentic	W	2	Exophytic	31.5
Sympetrum internum	Lentic	W	4	Exophytic	33.5
Sympetrum pallipes	Both	S-W	2	Endophytic	36
Sympetrum sanguineum	Lentic	S	3.5	Exophytic	36.5
Sympetrum striolatum	Both	W	5	Exophytic	39.5
Sympetrum vulgatum	Lentic	S	2	Exophytic	37.5

878　**Table S4**: Credible intervals of all MCMCglmm models testing predictions regarding the range

879　and phenology shifts across 66 odonate species in North America and Europe. These models are

880　detailed in *Model information and statements* of the Supplementary Information.

	Lower credible interval	Upper credible interval
Phenological shifts ~ Range shifts		
(Intercept)	-0.26	0.49
Range shifts	-0.71	-0.16
Continent	-0.75	0.30
Range shifts ~ Range geography + T ° variability		
(intercept)	-1.13	-0.07
Southern range	0.33	1.50
Widespread	-0.32	0.96
T ° variability	-0.53	-0.12
Both hab. Types	-	-
Lentic habitat	-	-
Flight duration	-	-
Oviposition	-	-
Body size	-	-

890 **Figure S1**: P-values and coefficients of 1000 GLM iterations testing whether range shifts as

891 calculated from random datasets predict the range shifts measured in the study. Each point shows

892 the results of a single GLM model, with measured range shifts as the dependant variable and

893 randomized range shifts as the independent variable.

Figure S2: Observed range shifts in km from the equator, against randomized predicted values

according to 4 random datasets. Points represent species and each pane contains a different set of

random data in calculations of randomized range shifts. There is no consistent relationship

among 1000 iterations.

Figure S3: P-values and coefficients of 1000 GLM iterations testing whether phenology shifts as

calculated from random datasets predict the phenology shifts measured in the study. Each point

shows the results of a single GLM model, with measured phenology shifts as the dependant

variable and randomized phenology shifts as the independent variable.

922   **Figure S4**: Observed phenology shifts in Julian day, against randomized predicted values

923   according to 4 random datasets. Points represent species and each pane contains a different set of

924   random data in calculations of randomized phenology shifts. There is no consistent relationship

925   among 1000 iterations.

Figure S5: Distribution of values computed for range shifts of 76 European and North American species between a recent time period (2008 - 2018) and a historical time period (1980 - 2002). See Methods section of the main text for full details on data assembly, and steps undertaken to produce these preliminary results.

Figure S6: Distribution of values computed for phenology shifts of 66 European and North

American species between a recent time period (2008 - 2018) and a historical time period (1980 -

2002). See Methods section of the main text for full details on data assembly, and steps

undertaken to produce these preliminary results.

939   <u>Figure S7, A</u>

Trace of (Intercept)

Iterations

Density of (Intercept)

N = 1350   Bandwidth = 0.04731

Trace of RangeShift

Iterations

Density of RangeShift

N = 1350   Bandwidth = 0.03503

Trace of ContinentNAM

Iterations

Density of ContinentNAM

N = 1350   Bandwidth = 0.06703

<u>Figure S7, B</u>

Figure S7: Panels A and B show the trace and density estimates of a phylogenetic mixed effects

model exploring the relationship between range and phenology shifts in North American and

European odonates (N = 66). 150,000 iterations were run to produce these results. These plots

verify model convergence and absence of autocorrelation within the explanatory variables.

<u>Figure S8, A</u>

Trace of (Intercept)

Iterations

Density of (Intercept)

N = 1350   Bandwidth = 0.06702

Trace of DistributionS

Iterations

Density of DistributionS

N = 1350   Bandwidth = 0.07843

Trace of DistributionW

Iterations

Density of DistributionW

N = 1350   Bandwidth = 0.08535

Trace of TemperatureVariability

Iterations

Density of TemperatureVariability

N = 1350   Bandwidth = 0.02616

952   Figure S8, B

Trace of units

Density of units

954   **Figure S8**: Panels A and B show the trace and density estimates a phylogenetic mixed effects

955   model testing whether ecological traits, and geographic and climatic attributes predict range

956   shifts in North American and European odonates (N = 76). 150,000 iterations were run to

957   produce these results. Results of the best model, according to DIC, are shown here. These plots

958   verify model convergence and absence of autocorrelation within the explanatory variables.