1 **The timing of preceding tactile inputs modulates cortical processing**

2 Wenyu Wan[1,2*], K. Richard Ridderinkhof[2], Arko Ghosh[1*]

4 [1]Cognitive Psychology Unit, Institute of Psychology, Leiden University, The Netherlands

5 [2]Department of Psychology, University of Amsterdam, The Netherlands

9 *Corresponding authors:

10 Wenyu Wan & Arko Ghosh

11 Institute of Psychology

12 Leiden University Wassenaarseweg 52

13 Leiden, 2333 AK, the Netherlands.

14 Email: w.wan@fsw.leidenuniv.nl [Wenyu Wan]

15 Email: a.ghosh@fsw.leidenuniv.nl [Arko Ghosh]

Summary

Tactile experiences in the real world are rich in texture and temporal patterns. While the role

of texture in driving somatosensory cortical activity is well established, there is emerging

evidence that somatosensory activity is sensitive to tactile temporal statistics (i.e., the time

intervals that separate stimuli in trains of successive tactile pulses). Cortical processing of

tactile pulses may be shaped by preceding pulses, but this influence will vary with inter-

stimulus intervals. It is possible that tactile information lingers only briefly in primary

somatosensory cortex and will influence processing of the next pulse only at short intervals;

in secondary somatosensory cortex it lingers longer, allowing it to shape the processing of

even multiple successive pulses over more prolonged intervals. Here we recorded scalp

EEG signals from somatosensory cortex in response to a train of tactile pulses presented to

the fingertips with varying inter-stimulus intervals spanning 100 to 10,000 ms. We traced

cortical tactile processing through its early (<75 ms), intermediate (75 to 150 ms) and late

stages (150 to 300 ms) . The early and late stages of somatosensory activity were similarly

shaped by the preceding pulse; this influence declined with increasing inter-stimulus interval.

The intermediate stage of somatosensory activity was sensitive to both the previous and the

penultimate pulses, a sensitivity that was again modulated by their temporal dynamics. Our

findings suggest that somatosensory cortex integrates complex temporal patterns during its

intermediary processing stages, allowing previous and even penultimate stimuli to modulate

ongoing processing of current stimuli.

Introduction

The peripheral nervous system encodes texture by using the timing of tactile inputs in the millisecond scale (Weber et al., 2013). In the central nervous system, the cortex is capable of separating sequences of tactile information according to their temporal pattern in the scale of milliseconds to seconds (Bale et al., 2017). This capacity may serve diverse sensory functions – from localising touch to learning behaviourally relevant patterns (Godde et al., 2000; Nguyen et al., 2016). In the real world, the temporal patterns cut across various time scales. For instance, the inter-touch interval distribution captured on the smartphone touchscreen is heavy tailed ranging from milliseconds to hours (Pfister and Ghosh, 2020). The somatosensory cortex may separate rich tactile sequences according to the recent history of tactile events – i.e., the previous and the penultimate intervals.

There is scattered evidence that the previous interval shapes somatosensory cortical processing, but the role of the penultimate interval is less clear. When the interval to the previous tactile stimulation is shortened, the somatosensory cortical response is attenuated Espenhahn et al. (2020). A similar pattern of results is also found for studies using electrical stimulation of peripheral nerves (Kisley and Cornwell, 2006; Spooner et al., 2020). The measurements are usually limited to few select intervals (such as continuous tactile stimulations separated by 150 and 1050 ms in Espenhahn et al. (2020) and paired tactile stimulations separated by 30, 60 and 150 ms in Wühle et al. (2011)). For very short (<150 ms) intervals, the suppression may be under the influence of cortical excitability, with higher attenuation corresponding to lower excitability (Lenz et al., 2012; Shagass and Schwartz, 1964). By using MEG and electrical stimulation, the signal attenuation has been localised to the primary and secondary somatosensory cortex (Hamada et al., 2002). The primary somatosensory cortex shows attenuation for intervals up to 100 ms, whereas the secondary somatosensory cortex shows attenuated signals for intervals up to 500 ms. A similar pattern of results is found for tactile stimulation, albeit the exploration only spanned intervals of 30 to 150 ms (Wühle et al., 2011). These results suggest that the tactile information lingers longer in the secondary than in the primary somatosensory cortex. This idea of lingering activity in the secondary somatosensory cortex is further supported by invasive neural recordings in non-human primates while they discriminate the frequency difference between two consecutive tactile stimuli (Hernández et al., 2010). The sustained information processing of preceding stimuli in the secondary somatosensory cortex may allow both the previous and penultimate pulses to shape the somatosensory processing of the current pulse, as well as its modulation by the length of the respective inter-stimulus intervals, as long as the total sequence occurs within 500 ms. If and how the previous and penultimate tactile pulses at intervals spanning >500 ms shape the cortical processing is not clear.

The conventional analytical framework – involving only a few select intervals – is limited in its
ability to address the role of rich temporal statistics in cortical processing. Furthermore, this
framework can only capture a simple relationship between the temporal statistics and brain
responses (e.g., the shorter the interval, the higher the cortical signal attenuation). By
contrast, the framework of joint-interval distribution (JID) allows us to separately measure
tactile experiences according to their next-interval dynamics (i.e., the duration of the previous
interval relative to the duration of the penultimate interval). This framework has been
previously used to capture the rich temporal statistics of spiking neurons (Eagan and
Partridge, 1989; Fitzurka and Tam, 1999) and smartphone touchscreen interactions (Ceolini
et al., 2022a; Ceolini and Ghosh, 2023; Ceolini et al., 2022b; Duckrow et al., 2021; Eagan
and Partridge, 1989; Fitzurka and Tam, 1999). According to this framework the stimulation
triad can be separated according to the joint properties of the previous interval (say k) and
the penultimate interval (say k-1), and then gathered in two-dimensional bins (b_n × b_n
number of bins) spanning the JID. Here, an event-related potential (ERP) signal that reflects
somatosensory (Espenhahn et al., 2020) can be estimated separately for each of the two-
dimensional bins (henceforth called JID-ERP). This approach can capture more complex
relationships than when using only few select intervals. For example, patterns associated
with short consecutive intervals may be differently attenuated compared to the patterns
associated with short followed by long intervals.

The different stages of sensory cortical processing may be distinctly influenced by these
temporal statistics. This influence can be captured using the signal latencies of the ERPs
derived from the separate two-dimensional bins of the JID. Therefore, this JID-ERP
framework can in theory capture a rich variety of influences of the temporal statistics on brain
activity. To reveal these typical dynamics, the high-dimensional feature space (b_n × b_n bins of
the JID-ERP × T signal processing latencies of the ERP) can be reduced by using non-
negative matrix factorization (NNMF) (Lee and Seung, 1999). NNMF is a dimensionality
reduction technique that can help reveal hidden patterns in the data by decomposing a
matrix into lower-rank matrices (not unlike principal components, but resulting in additive
rather than orthogonal components). The reduction can yield low-dimensional and
interpretable prototypical patterns (so-called meta patterns). This method of reduction has
been used to reduce a range of complex datasets from identifying facial features in images to
human behavioural patterns captured on the smartphone (Ceolini and Ghosh, 2023; Lee and
Seung, 1999). Here, the method can reduce the JID-ERP into a few interpretable prototypical
temporal structures (*meta-JIDs*) and their corresponding timing (*meta-ERPs*). The *meta-JIDs*
represent how the -cortical signals are modulated by preceding tactile pulses as a function of

the penultimate and previous intervals, whereas the *meta-ERPs* represent how the patterns

of modulations are distributed over the different signal processing stages.

In this study, we captured the EEG signals evoked by tactile pulse stimulations separated by

a range of previous and penultimate intervals spanning 100 to 10,000 ms. The stimulations

were presented at the fingertips. We used the JID framework yielding 5 × 5 two-dimensional

bins. The ERPs spanned an epoch from –200 ms prior to the tactile pulse to 299 ms

following the pulse. This yielded a combined JID-ERP feature space of 5 × 5 × 500 two-

dimensional bins. We reduced the JID-ERP after stimuli onset (5 × 5 × 299 two-dimensional

bins) into interpretable prototypical patterns by using NNMF. To preview, these *meta-JIDs*

and *meta-ERPs* revealed that the intermediate stages of cortical processing play a role that

stands in contrast to the early or late stages.

Result

Tactile evoked potentials and the distinct stages of cortical somatosensory
processing

This report focused on the high-dimensional JID-ERP approach, but we first described the
simple event-related potential signal disregarding the next-interval dynamics to provide an
overview of the signal components observed here. Note, throughout this section we focused
on the findings from the stimulations presented at the index fingertip (with other stimulation
sites described in the supplementary information). In response to tactile pulse stimulation at
the index fingertip, the evoked responses showed short-latency signals at the electrodes
placed over the somatosensory cortex, while longer-latency signals were observed over the
central electrodes (Supplementary Movie 1 & 4). As expected, the short-latency signals from
the electrodes located in somatosensory cortex showed components with peak latencies of
~50 ms (positive peak), ~75 ms (negative peak), ~100 ms (positive peak), ~150 ms (negative
peak), and ~200 ms (positive peak). For subsequent analysis using the JID-ERP approach,
we focused on the electrode with the largest amplitude (latency of 25-75 ms) over the
somatosensory cortex (Fig. 1B). In the rest of the report we use the terms *early*, *intermediate*
and *late* stages of the sensory processing to refer to the latencies of 1-75, 75-150, and 150-
300 ms, respectively. For the ERP stemming from the central electrode see Supplementary
Fig. 1. Across both the somatosensory and central electrodes, similar patterns as for the
index finger stimulation were also captured in response to stimulation of the thumb and index
finger (Supplementary Fig. 2, 3, 4, 5).

Mass-univariate analysis reveals the influence of recent temporal statistics on tactile
evoked potentials

To address how the tactile temporal statistics shapes the EEG signals, we separated the
tactile ERPs according to the next-interval dynamics preceding the stimulation (JID-ERP).
Here, a matrix of ERPs was separated in two-dimensional bins, binned based on the
previous and the penultimate intervals (Fig. 1C). We analysed the differences in the ERP
signal amplitudes across the JID for all the latencies of the ERP (JID-ERP). At each latency,
the amplitudes were *z*-scored across the JID, and one-sample *t*-tests (*z*-scores vs. 0)
revealed the two-dimensional bins with distinctly higher or lower activations (Fig. 1D). Note,
in this approach the bins are considered independently of each other (mass-univariate
statistics). In the pre-stimulation (but post ERP baseline) period, there was no consistent
difference between the two-dimensional bins. The differences became evident at the early
stages (~50 ms post stimulation), such that the bins with short previous intervals <500 ms

showed attenuated signals compared to the rest of the bins. The attenuation appeared irrespective of the penultimate interval. At this stage, there was evidence for signal enhancement for bins composed of short (<500 ms) penultimate intervals followed by long (>2,000 ms) intervals. The subsequent stage of processing, the intermediate stage, showed a more dynamic pattern of signal modulations which altered from one ERP peak to another. The late stage showed a pattern of modulations similar to the early stages – where the two-dimensional bins with <500 ms previous intervals showed attenuated signals. In this stage, there was enhancement of the signals with previous intervals >2,000 ms. A similar pattern of results was observed when the thumb or the little finger was stimulated (Supplementary Figure 2 & 4, Supplementary Movie 2 & 3).

We also analysed the ERPs over the central electrode, consisting of a broad signal peaking at ~ 200 ms. This entire signal was attenuated at the two-dimensional bins with previous interval <500 ms. This pattern was observed for all three stimulation locations (Supplementary Movie 4, 5 & 6, Supplementary Figure 1, 3 & 5).

Similarities in the pattern of JID-ERP modulation at the early and late stages of somatosensory processing

The JID-ERP revealed modulation of the EEG signals according to the two-dimensional bins based on the previous and penultimate intervals. As described above, the modulation varied according to the stage of processing for the somatosensory signals. We quantified the similarities of the JID-ERP at any given latency to the rest of the latencies by using correlations (Fig. 2A). For the index finger stimulations, this analysis revealed strong similarities between the JID-ERP patterns at the early and late stages, but the intermediate stage appeared largely unrelated to the other processing stages. There was a notable persistence of patterns within the different stages of processing – but not for the intermediate stages. The distinction of the intermediate stage was most prominent for the index finger, followed by the thumb and the little finger (Supplementary Figure 2C & 4C). For the ERP over the central electrode, the JID-ERP patterns showed similar modulations across the intermediate and late stages (Supplementary Figure 3C & 5C).

Dimensionality reduction: Interval patterns relevant for somatosensory processing across the different stages of processing

By using simple analytical frameworks, i.e., mass-univariate statistics and correlation analysis, we established the dominant patterns underlying the JID-ERP, but these methods

are not designed to extract time-variant complex patterns. Therefore, we next used data-
driven dimensionality reduction to extract the key patterns of the rich JID-ERP. Essentially,
the JID-ERP magnitudes (absolute value of the EEG signals, see Methods) spanning all the
latencies were reduced in terms of a few *meta-JIDs*, i.e., prototypical interval-based
modulations of the two-dimensional bins, and the corresponding *meta-ERPs*, i.e., the extent
to which prototypical patterns were present at any given latency (Fig. 3A). We applied this
analysis both on the grand averaged JID-ERP signal and on the individual JID-ERP traces,
and also to the signals from the somatosensory and the central electrodes. Here we focus on
the results from the index finger (for the thumb and little finger please see Supplementary
Figure 6 & 7).

For the grand averaged signal from the somatosensory electrode, the dimensionality
reduction revealed three prototypical patterns. First, we observed a prototypical pattern
where short penultimate followed by short previous intervals were attenuated and the
attenuation diminished for longer previous intervals. According to the corresponding *meta-
ERPs* this pattern was present in the early and late stages of somatosensory processing.
Second, we observed a more complex prototypical pattern where penultimate intervals
<1,000 ms and previous intervals between 500 and 1,000 ms showed the larger amplitudes.
This pattern was present in both the early and intermediate stages of processing. Third, we
observed a pattern where previous intervals <500 ms showed larger amplitudes than longer
previous intervals, and within this range, the longer penultimate intervals showed larger
amplitudes. This pattern was largely constrained to the period of transition to and from the
intermediate stage.

We performed a similar analysis at the level of each individual and then clustered the results
based on the *meta-ERPs*. The corresponding *meta-JIDs* were averaged across the clusters.
This approach revealed three prototypical patterns. First, we observed a pattern similar to the
first prototypical pattern identified using the grand average signal. Here, the signals were
attenuated for short consecutive intervals and enhanced for the temporally distant
consecutive intervals. This pattern was present in the early and the late stages of
somatosensory processing. Second, we observed a pattern similar to the first pattern but
with a sharper attenuation for previous intervals <500ms and with strong presence in the late
stages of the processing. Third, there were *meta-ERPs* which indicated peculiar modulation
at the intermediate stages but the averaged *meta-JIDs* revealed only a marginal
enhancement for long penultimate intervals followed by short previous intervals – the
marginally central tendency is indicative of inter-individual variability at this stage.

The grand averaged signal at the central electrode revealed two prototypical patterns. The
first was present only in the early stages of somatosensory processing. This pattern
appeared complex with a tendency for enhanced processing of long penultimate intervals
followed by short intervals (Supplementary Figure 7A). The second was present in the
intermediate and late stages. This pattern was characterised by an attenuation of
stimulations with short previous intervals. The individual-participant signals revealed an
additional complex prototypical pattern which was specific to the intermediate stage
(Supplementary Figure 7A).

Discussion

We analysed cortical signals in response to a train of tactile inputs on the hand, with the interval between stimulations spanning ~100 to 10,000 ms. Signal processing in the somatosensory cortex was shaped by both the penultimate and previous tactile pulses, and this influence was modulated by the respective inter-stimulus intervals spanning up to 8,000 ms. The intervals modulated the cortical processing by using rules which varied according to the stages of processing. By using a series of analytical approaches, we showed that the early and the late stages of cortical processing were similarly influenced by the interval patterns. In both of these stages, the signals were attenuated when the previous intervals were shorter than 500 ms. The influence of the temporal statistics at the intermediate stages was modulated by a more complex combination of the penultimate and previous intervals. Our findings indicate that the ability to separate tactile information according to temporal structures in the scales of milliseconds to seconds is distributed across different cortical networks.

One of the prominent patterns we observed included a sharp attenuation of the early cortical signals for previous intervals <500 ms. This is consistent with prior studies (Hamada et al., 2002; Zhu et al., 2007). For instance, inter-touch intervals separated by 2,000 ms resulted in larger amplitude in early stages of the ERP (<75 ms) compared to intervals separated by <500 ms; the late-stage ERPs (75-150 ms) were virtually abolished by the shorter intervals (Zhu et al., 2007). One possibility is that this attenuation reflects a form of inhibition induced when processing the first input (not unlike prepulse inhibition of startle responses; see, e.g., Braff et al., 2001). In our observations, the penultimate intervals played only a marginal role in further attenuating signals; the putative inhibitory processes appeared short-lived (i.e., lasts <500 ms). The data driven dimensionality reduction revealed an enhancement of the signal amplitudes at the offset of the early processing stage (latency of ~75 ms). This may be explained by a rebound of activation, as often seen post-inhibition (Qi et al., 2016). Finally, there may be a concern that the attenuation observed does not reflect altered neural activity but rather how the EEG signal evoked by the second stimuli summates with the previous stimuli. However, this is unlikely as we saw no signal changes in the segment that appeared after the baseline period but prior to the second stimuli, and there was temporal jitter in the intervals within each bin.

Another pattern of modulation consisted of amplitude gradient across the 'diagonal' of the joint-interval distribution: with stronger signal attenuation for consecutive short-short intervals than for consecutive long-long intervals. The gradient spanned ~100 to 8,000 ms. We speculate that this pattern reflects mechanisms that help sustain neural representations

towards temporal integration of event series, such as short-term memory or perceptual
priming (Contreras et al., 2013; Han et al., 2008). Animal studies suggest that this kind of
reverberation of brain activity for past events can last up to a few minutes (Contreras et al.,
2013). Based on the ERP signal latencies, we suggest that the activity in the early stages is
echoed or recurred in the late stages resulting in the similarity of prototypical pattern between
those two stages (Auksztulewicz et al., 2012; de Lafuente and Romo, 2006). Alternatively,
some top-down modulation at the sensory cortex (Ruff, 2013) may exist in the late-stage
processing to get ready for the arriving stimuli.

Our results also indicated a function of the intermediary stages which is distinct from the
early and late stages of somatosensory processing. According to the correlational analysis,
the JID-ERP was distinct at these stages as opposed to the early and late stages.
Furthermore, the dimensionality reduction also revealed complex prototypical patterns largely
unique to the intermediary stages. Our results are in line with previous studies indicating that
complex temporal integration follows early somatosensory cortical processing (Rossi-Pool et
al., 2021). However, exactly which mechanisms underscore these remains unclear. It is
possible that stimulation induced inhibition, sustained neural representation, and links to
extra-somatosensory processes (such as for motor control) all contribute to the distinct
processing at the intermediary stages (Forss and Jousmaki, 1998; Haggard and Whitford,
2004).

In this study we separated the ERP signals according the dynamics of the tactile stimulations.
The patterns captured in the so-called JID-ERP suggest that tactile processing is shaped
according to the temporal context. This parallels the emerging view from both human and
animal studies that cortical sensory processing is state-dependent (Parajuli et al., 2023;
Russell et al., 2024; Taga et al., 2018; Zagha et al., 2013).

Our study provides a new perspective on studying how brain states interact with continuous
sensory stimuli from outside. Our findings lead to the possibility that early stages of cortical
processing may leverage the timing of previous and penultimate inputs to recognise the
current state. Such a process may work in tandem with other feed-forward processes in early
sensory processing to prioritise select input features (Westerberg et al., 2023). The sensory
processing revealed here was largely consistent across the different fingers – the thumb, the
index and little finger – indicating that the discovered state-dependent processing may
generalise to the rest of the hand. The JID-ERP approach introduced here may be extended
to other body parts or even other sensory modalities to reveal the shared mechanisms
underlying state dependent cortical sensory processing.

Methods

Participants

This study is part of a large data collection effort involving long-term smartphone behavioral
logs and EEG in the laboratory. Participants were recruited from a pool accumulated through
the agestudy.nl data collection platform. This platform leverages the Dutch brain registry
(Zwan et al., 2021). Sixty-four healthy participants (age range: 20-81; 40 female) were
enrolled. However, the self-reported health status of 4 participants altered leading to
eliminations (one had tinnitus, one had stroke, one had arthrosis, and one had a psychiatric
condition). All participants provided the informed consent and the study was approved by the
Ethics Committee of Psychology, Leiden University (ERB-reference number: 2020-02-14-
Ghosh, dr. A.-V2-2044).

Tactile Stimulation

We used tactile stimulations while recording the EEG signals (see below). Continuous train of
tactile stimuli were applied to participants' three fingertips (right thumb, index finger, and little
finger, in different blocks) by using miniature electromagnetic solenoid-type stimulator
(Dancer Design, North Yorkshire, England). The stimulator was controlled by using the
parallel port of a computer and MATLAB (MathWorks, Natick, USA). We delivered 1800
stimulations, with each stimulation lasting 10 ms-long and the stimulations were
suprathreshold according to verbal reporting conducted while testing the attached stimulators.
The stimulations were time-stamped by the EEG amplifier based on the cop (see below). The
inter-stimulus interval varied from 100 to 10,000 ms. In order to cover any noise generated
by tactile stimulators and make participants feel less bored during measurement, we played
a documentary series (David Attenborough's Africa) on a screen and delivered the audio
stream via a neck speaker. Participants were encouraged to have a short break every ~15
minutes and the stimulation session lasted about 1 h.

EEG recordings and preprocessing

EEG data was collected by using 64 channels EEG caps (62 scalp electrodes, 2 ocular
electrodes) with customized equidistant layout (EasyCap Gmbh, Wörthsee,
Germany)(Haenzi et al., 2014). The EEG signals were gathered referenced to the vertex and
amplified with BrainAmp amplifier (Brain Products GmbH, Gilching, Germany). The signals
were recorded and digitalized at the sampling rate of 1KHz. The participants arrived for the

measurement with washed hair and scalp, and we further degreased the skin at the contact sites by using alcohol swabs. We applied Supervisc gel (Easycap) to obtain an electrical contact between the skin and electrode, and targeted an impedance under 10 kOhm for each electrode.

The data was processed using the EEGLAB toolbox (Delorme and Makeig, 2004) on MATLAB 2022b (MATLAB, Mathworks, Natick). Any recorded electrode with impedance >10 kOhm was excluded. The data was band filtered between 0.1 and 75 Hz. Eye-blink related EEG signal artefacts were removed by using independent component analysis (Pontifex et al., 2017). Next, the data was further filtered between 1 and 45 Hz. An excluded channels were subsequently interpolated. The data was epoched using a temporal window of 500 ms: - 200 to 300 ms from the onset of the stimuli. All epochs with signals beyond the set ±80 µV threshold were excluded. The data was re-referenced to the average from scalp electrodes. Note, the baseline was corrected for each epoch by subtracting the signal mean from -200 to -50 ms. The electrode with the largest short-latency signals (latency of 25-75 ms) placed over the left somatosensory cortex and with longer latency signals (latency of 150-300 ms) located in the central areas were chosen for each participant.

We excluded 2 participants due to corrupted EEG data files and six participants were further removed based on a signal amplitude threshold (10th percentile of the population distribution). This yielded 52 participants for the subsequent ERP analysis.

JID-ERP

We adapted the joint-interval distribution (JID) approach to separate the tactile stimulations according to their next interval dynamics. We introduced this approach for the study of real-world touchscreen behavioral dynamics (Ceolini and Ghosh, 2023; Duckrow et al., 2021) .

Here, a previous interval (say *k*) is described in conjunction with the penultimate interval (say *k*-1). The tactile joint intervals (stimulation triads) were binned into 5 × 5 two-dimensional bins with edges (upper bound) of 0.5, 1, 2, 4, 8 s. Then, for each bin separately we estimated the tactile-related potential (ERP) by averaging the signals evoked by the last event of the triad. The threshold number of events necessary to estimate the ERP was set to 10. After these steps were conducted for a given individual, for some of the analysis we additionally pooled across the individuals by creating a grand-average of the binned ERPs. Note, to reveal whether a particular bin's amplitude was higher or lower than other bins at a given latency of the ERP, the JID-ERP was normalised by using *z*-score at each of the latencies.

Mass-univariate statistical analysis

To reveal which two-dimensional bins were consistently suppressed or facilitated, we used
mass-univariate one-sample *t*-tests of the *z*-scored amplitudes at all the time points. Multiple
comparisons were corrected by using false discovery rate (FDR, using fdr() from eeglab
toolbox ALG NAME) applied across the entire JID-ERP (α = 0.05). To quantify the similarities
of the JID-ERP at any given latency to the rest of the latencies, a cross-correlation analysis
was employed based on *z*-transformed grand-averaged JID-ERP using the Spearman
method. Multiple comparisons were corrected by using the FDR method (α = 0.05). The
values along the diagonal of the cross-correlation matrix (ρ = 1, window of 25 ms from the
diagonal) was eliminated from the FDR.

Interpretable data-driven dimensionality reduction

In the mass-univariate analysis above, each two-dimensional bin of the JID-ERP is
considered independently from the other bins. However, the patterns in the JID-ERP may
simultaneously encompass various bins in a complex manner. To extract such patterns we
leveraged non-negative matrix factorization (NNMF). This method was originally
demonstrated in facial image analysis, revealing key facial features (Lee and Seung, 1999),
and extended to the study of behavioral JID to discover prototypical behaviours (Ceolini and
Ghosh, 2023). Here we applied a variant of the method, stable and reproducible NNMF (star
NNMF) described here (Ceolini and Ghosh, 2023) . In brief, at the level of each individual, we
applied the NNMF on a matrix of JID-ERP, where the EEG signal magnitudes (absolute value)
was *z*-scored at all ERP latencies. Because this method requires non-negative inputs, we
subtracted minimum of the *z*-transformed matrix from the entire matrix. The dimensional
reduction method yielded a lower-rank approximation in terms of prototypical patterns: how
the signals were modulated across the two-dimensional bins, *meta-JIDs*, and the
corresponding latencies where the pattern was present, *meta-ERPs*. In the search for the
optimal cluster between 2 to 10 ranks, and evaluated the optimal cluster number by
calculating the error between real matrix and reconstructed data. The stable reduction was
established based on 100 repetitions. We further clustered similar *meta-ERPs* across the
population by using *k*-means; optimal cluster size search between 3 to 7 clusters, 1000
repetitions, with *silhouette* method using evalclusters.mat as implemented in MATLAB. The
NNMF was applied also applied to the grand-averaged EEG JID-ERP matrix using the same
steps as described above but the *k*-means clustering was not applicable in this case.

Data, Materials, and Software Availability

Pseudo-anonymized MATLAB.mat file containing processed JID-ERP data is made available on dataverse.nl upon publication; The custom-written scripts used towards this report are shared on: https://github.com/CODELABCODELIB/JID_ERP_Tactile_2024.

Acknowledgements

This research was funded by Velux Stiftung (No.1283, awarded to A.G. with K.R.R as co-applicant). We thank Ruchella Kock for her help in programming. We thank Beste Yavuz, Lysanne Groenewegen, Barbora Michalidesová, Lorenzo Van Hoorde, and etc. for assistance in EEG data collection. We thank all participants for contributing with their time and effort.

Author contributions

A.G. conceived the study. A.G. and W.W. designed the study. A.G. programmed the experiments and performed the EEG pre-processing. W.W acquired and analysed the data with the aid of A.G.. W.W drafted with the aid of A.G.. K.R.R, A.G., W.W. edited the manuscript.

Declaration of interests

A.G. is a co-founder and chairman of QuantActions AG, and is an advisor for Axite B.V. W.W. and K.R.R. had no conflicts to disclose.

Reference

Auksztulewicz, R., Spitzer, B., Blankenburg, F., 2012. Recurrent neural processing and
somatosensory awareness. Journal of Neuroscience 32, 799-805.
Bale, M.R., Bitzidou, M., Pitas, A., Brebner, L.S., Khazim, L., Anagnou, S.T., Stevenson,
C.D., Maravall, M., 2017. Learning and recognition of tactile temporal sequences by mice
and humans. Elife 6, e27333.
Braff, D.L., Geyer, M.A., Swerdlow, N.R., 2001. Human studies of prepulse inhibition of
startle: normal subjects, patient groups, and pharmacological studies. Psychopharmacology
(Berl) 156, 234-258.
Ceolini, E., Brunner, I., Bunschoten, J., Majoie, M.H., Thijs, R.D., Ghosh, A., 2022a. A model
of healthy aging based on smartphone interactions reveals advanced behavioral age in
neurological disease. Iscience 25.
Ceolini, E., Ghosh, A., 2023. Common multi-day rhythms in smartphone behavior. NPJ
Digital Medicine 6, 49.
Ceolini, E., Kock, R., Band, G.P., Stoet, G., Ghosh, A., 2022b. Temporal clusters of age-
related behavioral alterations captured in smartphone touchscreen interactions. Iscience 25.
Contreras, E.J.B., Schjetnan, A.G.P., Muhammad, A., Bartho, P., McNaughton, B.L., Kolb, B.,
Gruber, A.J., Luczak, A., 2013. Formation and reverberation of sequential neural activity
patterns evoked by sensory stimulation are enhanced during cortical desynchronization.
Neuron 79, 555-566.
de Lafuente, V., Romo, R., 2006. Neural correlate of subjective sensory experience gradually
builds up across cortical areas. Proceedings of the National Academy of Sciences 103,
14266-14271.
Delorme, A., Makeig, S., 2004. EEGLAB: an open source toolbox for analysis of single-trial
EEG dynamics including independent component analysis. Journal of neuroscience methods
134, 9-21.
Duckrow, R., Ceolini, E., Zaveri, H., Brooks, C., Ghosh, A., 2021. Artificial neural network
trained on smartphone behavior can trace epileptiform activity in epilepsy. IScience 102538.
Eagan, K.P., Partridge, L.D., 1989. The sequential-interval state space: a means of
displaying temporal information in neuron firing. Journal of neuroscience methods 27, 245-
252.
Espenhahn, S., Yan, T., Beltrano, W., Kaur, S., Godfrey, K., Cortese, F., Bray, S., Harris, A.D.,
2020. The effect of movie-watching on electroencephalographic responses to tactile
stimulation. Neuroimage 220, 117130.
Fitzurka, M.A., Tam, D.C., 1999. A joint interspike interval difference stochastic spike train
analysis: detecting local trends in the temporal firing patterns of single neurons. Biological
cybernetics 80, 309-326.
Forss, N., Jousmaki, V., 1998. Sensorimotor integration in human primary and secondary
somatosensory cortices. Brain Res 781, 259-267.
Godde, B., Stauffenberg, B., Spengler, F., Dinse, H.R., 2000. Tactile coactivation-induced
changes in spatial discrimination performance. Journal of Neuroscience 20, 1597-1604.
Haenzi, S., Stefanics, G., Lanaras, T., Calcagni, M., Ghosh, A., 2014. Altered cortical
activation from the hand after facial botulinum toxin treatment. Ann Clin Transl Neurol 1, 64-
68.
Haggard, P., Whitford, B., 2004. Supplementary motor area provides an efferent signal for
sensory suppression. Brain Res Cogn Brain Res 19, 52-58.
Hamada, Y., Otsuka, S., Okamoto, T., Suzuki, R., 2002. The profile of the recovery cycle in
human primary and secondary somatosensory cortex: a magnetoencephalography study.
Clinical neurophysiology 113, 1787-1793.
Han, F., Caporale, N., Dan, Y., 2008. Reverberation of recent visual experience in
spontaneous cortical waves. Neuron 60, 321-327.
Hernández, A., Nácher, V., Luna, R., Zainos, A., Lemus, L., Alvarez, M., Vázquez, Y.,
Camarillo, L., Romo, R., 2010. Decoding a perceptual decision process across cortex.
Neuron 66, 300-314.

Kisley, M.A., Cornwell, Z.M., 2006. Gamma and beta neural activity evoked during a sensory
gating paradigm: effects of auditory, somatosensory and cross-modal stimulation. Clinical
neurophysiology 117, 2549-2563.
Lee, D.D., Seung, H.S., 1999. Learning the parts of objects by non-negative matrix
factorization. nature 401, 788-791.
Lenz, M., Tegenthoff, M., Kohlhaas, K., Stude, P., Höffken, O., Tossi, M.A.G., Kalisch, T.,
Dinse, H.R., 2012. Increased excitability of somatosensory cortex in aged humans is
associated with impaired tactile acuity. Journal of Neuroscience 32, 1811-1816.
Nguyen, E.H., Taylor, J.L., Brooks, J., Seizova-Cajic, T., 2016. Velocity of motion across the
skin influences perception of tactile location. J Neurophysiol 115, 674-684.
Parajuli, A., Gutnisky, D., Tandon, N., Dragoi, V., 2023. Endogenous fluctuations in cortical
state selectively enhance different modes of sensory processing in human temporal lobe.
Nature Communications 14, 5591.
Pfister, J.-P., Ghosh, A., 2020. Generalized priority-based model for smartphone screen
touches. Physical Review E 102, 012307.
Pontifex, M.B., Miskovic, V., Laszlo, S., 2017. Evaluating the efficacy of fully automated
approaches for the selection of eyeblink ICA components. Psychophysiology 54, 780-791.
Qi, H.-X., Reed, J.L., Franca, J.G., Jain, N., Kajikawa, Y., Kaas, J.H., 2016. Chronic
recordings reveal tactile stimuli can suppress spontaneous activity of neurons in
somatosensory cortex of awake and anesthetized primates. Journal of neurophysiology 115,
2105-2123.
Rossi-Pool, R., Zainos, A., Alvarez, M., Parra, S., Zizumbo, J., Romo, R., 2021. Invariant
timescale hierarchy across the cortical somatosensory network. Proceedings of the National
Academy of Sciences 118, e2021843118.
Ruff, C.C., 2013. Sensory processing: who's in (top-down) control? Annals of the New York
Academy of Sciences 1296, 88-107.
Russell, L.E., Fişek, M., Yang, Z., Tan, L.P., Packer, A.M., Dalgleish, H.W., Chettih, S.N.,
Harvey, C.D., Häusser, M., 2024. The influence of cortical activity on perception depends on
behavioral state and sensory context. Nature Communications 15, 2456.
Shagass, C., Schwartz, M., 1964. Recovery functions of somatosensory peripheral nerve
and cerebral evoked responses in man. Electroencephalography and clinical
neurophysiology 17, 126-135.
Spooner, R.K., Eastman, J.A., Wiesman, A.I., Wilson, T.W., 2020. Methodological
considerations for a better somatosensory gating paradigm: The impact of the inter-stimulus
interval. Neuroimage 220, 117048.
Taga, G., Watanabe, H., Homae, F., 2018. Developmental changes in cortical sensory
processing during wakefulness and sleep. Neuroimage 178, 519-530.
Weber, A.I., Saal, H.P., Lieber, J.D., Cheng, J.-W., Manfredi, L.R., Dammann III, J.F.,
Bensmaia, S.J., 2013. Spatial and temporal codes mediate the tactile perception of natural
textures. Proceedings of the National Academy of Sciences 110, 17107-17112.
Westerberg, J.A., Schall, J.D., Woodman, G.F., Maier, A., 2023. Feedforward attentional
selection in sensory cortex. Nat Commun 14, 5993.
Wühle, A., Preissl, H., Braun, C., 2011. Cortical processing of near-threshold tactile stimuli in
a paired-stimulus paradigm–an MEG study. European Journal of Neuroscience 34, 641-651.
Zagha, E., Casale, A.E., Sachdev, R.N., McGinley, M.J., McCormick, D.A., 2013. Motor
cortex feedback influences sensory processing by modulating network state. Neuron 79,
567-578.
Zhu, Z., Disbrow, E.A., Zumer, J.M., McGonigle, D.J., Nagarajan, S.S., 2007. Spatiotemporal
integration of tactile information in human somatosensory cortex. BMC neuroscience 8, 1-14.
Zwan, M.D., van der Flier, W.M., Cleutjens, S., Schouten, T.C., Vermunt, L., Jutten, R.J., van
Maurik, I.S., Sikkes, S.A.M., Flenniken, D., Howell, T., Weiner, M.W., Scheltens, P., Prins,
N.D., 2021. Dutch Brain Research Registry for study participant recruitment: Design and first
results. Alzheimers Dement (N Y) 7, e12132.

 Figure Legends

Figure 1. The temporal dynamics of tactile inputs shape cortical somatosensory processing.
(A) Schematic overview of the approach used to link the next-interval dynamics of tactile
stimulations to cortical processing. A train of tactile stimuli with varying intervals (from 100 to
10,000 ms) was applied to the index fingertip. The stimulations were separated according to
the joint properties of the previous interval (k) and the penultimate interval (k-1) into 5 × 5
two-dimensional bins. (B) At the level of each individual, the electrode with the largest
amplitude at short latencies was chosen. The electrode locations were shown as insert. The
plot showed the grand averaged tactile evoked potentials across the sample. The blue, red,
and yellow bars marked the early (from 1-75 ms), intermediate (from 75-150 ms), and late
stage (from 150-300 ms), respectively. The grand average was shown using a red line and
the evoked potentials from each individual are shown in grey. The grey contour marked the
baseline to which individual ERPs were aligned. (C) An example of separating the ERPs
according to the next-interval dynamics (the JID-ERP). (D) Population level JID-ERP based
on the grand average EEG signals. The first column showed the ERP signal disregarding the
next-interval dynamics to help approximate the stage of sensory processing. The second
column showed the JID-ERP of signal amplitudes at different latencies. The third column re-
scaled the amplitudes (z-score) across the current ERP temporal latency. The fourth column
revealed which two-dimensional bins were consistently suppressed or enhanced based on
mass-univariate one-sample t-tests of the z-scored amplitudes (multiple comparisons was
corrected by using false discovery rate (FDR) applied across the entire JID-ERP).

Figure 2. The JID-ERP pattern is shaped by temporal dynamics at distinct stages. (A)
Schematic overview of the cross-correlation analysis for calculating the similarity of each
time point of JID-ERP. Bar colors are identical to Fig 1. (B) Cross-correlation analysis results
of tactile evoked JID-ERP of the index finger. The left panel is the coefficient plot, and the
right panel is the p statistic plot (p<.05, FDR).

Figure 3. The typical JID-ERP patterns across different stages. (A) A sketch of the NNMF
dimensionality reduction method. This method can yield low-dimensional and interpretable
prototypical patterns, which can indicate what the pattern is (*meta-JIDs*) and when the
pattern appeared (*meta-ERPs*). (B) Three typical JID-ERP patterns at the population level,
derived from the grand averaged JID-ERP. (C) Three typical JID-ERP patterns at the
individual-participant level. We obtained the typical JID-ERP pattern for each participant and
then clustered meta-JIDs using k-means (see Methods). Finally, we averaged *meta-JIDs* and
meta-ERPs for each cluster. Note that a different scale was used here compared to (B). The

averaged *meta-JIDs* was depicted by a black line, while the *meta-JIDs* for each participant
was shown in grey.

Supplementary Figure 1. Supplementary Figure 1. The temporal dynamics of tactile inputs
from index fingertip can shape the cortical processing. (A) At the level of each individual, the
electrode with the largest amplitude at long latencies (150-300ms) was chosen. The central
electrode locations were shown as insert. The grand averaged tactile evoked potentials
across the sample were shown using a red line and the evoked potentials from each
individual were shown in grey. Blue bar marked the early stage (from 1-75ms), red bar
marked the intermediate stage (from 75-150ms), and yellow bar marked the late stage (from
150-300ms). (B) Population level JID-ERP based on the grand average EEG signals. The
first column showed the ERP signal disregarding the next interval dynamics to help
approximate the stage of sensory processing. The second column showed JID-ERP signal
amplitudes at different latencies. The third column re-scaled the amplitudes (z-score) across
the current ERP temporal latency. The fourth column revealed which two-dimensional bins
were consistently suppressed or enhanced based on mass-univariate one-sample t-tests of
the z-scored amplitudes (multiple comparisons was corrected by using false discovery rate,
FDR applied across the entire JID-ERP). (C) Cross-correlation analysis results of tactile
evoked JID-ERP of index finger. The left panel was the coefficient plot, and the right panel
was the p statistic plot (survived by p<.05 and FDR).

Supplementary Figure 2. The temporal dynamics of tactile inputs from thumb fingertip can
shape the cortical processing. (A) At the level of each individual, the electrode with the
largest amplitude at short latencies (25-75ms) was chosen. The electrode location placed
over left somatosensory cortex is shown as insert. The grand averaged tactile evoked
potentials across the sample. Blue bar marked the early stage (<75ms), red bar marked the
intermediate stage (from 75-150ms), and yellow bar marked the late stage (from 150-300ms).
The grand averaged potentials were shown using a red line and the potentials from each
individual were shown by grey lines. (B) (C) Please see the supplementary Figure1 for
legend.

Supplementary Figure 3. The temporal dynamics of tactile inputs from thumb fingertip can
shape the cortical processing at the central electrode location. Please see the supplementary
Figure1 for legend.

Supplementary Figure 4 . The temporal dynamics of tactile inputs from index fingertip can
shape the cortical processing at the electrode placed over the left somatosensory cortex.
Please see the supplementary Figure2 for legend.

Supplementary Figure 5. The temporal dynamics of tactile inputs from index fingertip can
shape the cortical processing at the central electrode location. Please see the supplementary
Figure1 for legend.

Supplementary Figure 6. The typical JID-ERP patterns across different stages for other two
fingers with the electrode placed over the left somatosensory cortex. (A) Typical *JID-ERP*
patterns for thumb. The frequency for each electrode was shown as insert topography plot.
The first row of right side showed the population level results based on the grand averaged
EEG signals across the sample. The second row showed the individual level results after
clustering all participants *meta-ERPs* and averaging *meta-JIDs* and *meta-ERPs* for each
clusters. *Meta-ERPs* for each participant was shown as grey line and the averaged *meta-*
JIDs was shown as dark line. Different scales were used for averaged *meta-JIDs* compared
to (A). (B) Typical *JID-ERP* patterns for little finger. Same legend with (A).

Supplementary Figure 7. The typical JID-ERP patterns across different stages for all fingers
with the electrode placed over the central electrode. (A) Typical JID-ERP patterns for index
finger. (B) Typical JID-ERP patterns for thumb. (C) Typical JID-ERP patterns for little finger.
Please see the supplementary Figure6 for the detailed legend.

Supplementary Movie 1. Grand averaged tactile evoked potentials of electrode placed over
somatosensory cortex for index finger. Population level JID-ERP based on the grand
average EEG signals. Re-scaled the amplitudes (*z*-score) across the current ERP temporal
latency. Mass-univariate one-sample t-tests of the z-scored amplitudes (multiple comparisons
was corrected by using false discovery rate (FDR) applied across the entire JID-ERP).

Supplementary Movie 2. Grand averaged tactile evoked potentials of electrode placed over
somatosensory cortex for thumb. Please see the supplementary Movie 1. for the detailed
legend.

Supplementary Movie 3. Grand averaged tactile evoked potentials of electrode placed over
somatosensory cortex for little finger. Please see the supplementary Movie 1. for the detailed
legend.

Supplementary Movie 4. Grand averaged tactile evoked potentials of central electrode for
index finger. Please see the supplementary Movie 1. for the detailed legend.

Supplementary Movie 5. Grand averaged tactile evoked potentials of central electrode for
thumb. Please see the supplementary Movie 1. for the detailed legend.

Supplementary Movie 6. Grand averaged tactile evoked potentials of central electrode for
little finger. Please see the supplementary Movie 1. for the detailed legend.

A

ρ=0.165(p>.05)

105ms

55ms

ρ=0.688(p<.05)

200ms

55ms

Latency[ms]

ρ

Latency[ms]

B

Latency[ms]

ρ

Latency[ms]

p<.05, FDR

Latency[ms]

Latency[ms]