Evidence for homing behaviour consistent with path integration in *Dascyllus trimaculatus*

Freddie Murley[1], Adelaide Sibeaux[1], Theresa Burt de Perera[1]

[1] Department of Biology, University of Oxford, 11a Mansfield Road, Oxford, OX1 3SZ, UK

Corresponding author: freddie.murley@gmail.com

Abstract

Path integration is a strategy that allows animals to monitor their movement relative to a starting point using directional and distance cues. It has been observed in a wide range of terrestrial species, but evidence of this behaviour in fish is still lacking. In contrast to most animals shown to navigate via path integration, fish are not surface bound but exist within a three-dimensional medium. This may present additional challenges when monitoring their position relative to a specific point. I developed a novel experimental paradigm to test whether the coral reef-dwelling domino damselfish *Dascyllus trimaculatus* could use path integration to navigate back to a shelter location. This consisted of a circular pool with a shelter and landmarks near the edge and a trap containing food in the centre. My results show that these fish follow homing trajectories back to the previous location of the removed shelter without using landmarks after a foraging trip into the central trap. Error in homing trajectory increased with outward path length, consistent with the use of path integration. These results add *D. trimaculatus* to the growing list of path-integrating species, opening new avenues for investigating the evolution and neural underpinnings of this navigational strategy.

Introduction

Path integration is a powerful navigational strategy that allows animals to return to a home location efficiently at the end of an outward journey. A path integrating animal must keep track of its directional change and distance travelled on an outward trip using movement relative to direction cues and odometer cues. This is essential to continually update a homing vector, which is the most direct path back to a home location (Müller & Wehner, 1988a; Patel & Cronin, 2020b, Figure 1).

The greater the distance a path-integrating animal travels on an outward journey, the more error is predicted to accumulate around its estimate of home location. This is due to the cumulative effect of small errors in directional change and distance measurements over the course of the trip due to inherent sensorimotor system and environmental noise (Cheung, 2014; Cheung et al., 2007). To compensate for the accumulation of errors, many animals perform search behaviour if they don't reach the exact location of their home using their self-generated homing vector (Patel & Cronin, 2020c). Searching animals often follow ever increasing loops around the end of their homing vector, returning to the start of their search at the end of each loop (Wehner & Srinivasan, 1981). The total journey of a path integrating animal leaving its shelter for a foraging trip can therefore be broken down into (1) an outward trip, (2) a homeward path following a homing vector, and (3) search behaviour if the homing vector is inaccurate.

Figure 1 - *How an animal can keep track of its location relative to its home and calculate the most direct route back via path integration.*

Animals can use allothetic or idiothetic directional cues for path integration. Allothetic directional cues are derived from the environment and include stable external directional cues like compasses based on the Sun's position or Earth's magnetic field. Idiothetic cues are generated from self-motion and

can incorporate information from the proprioceptive or vestibular systems. Error accumulates more quickly if using idiothetic cues than if using allothetic cues for path integration (Cheung et al., 2007). The use of idiothetic directional cues requires the cumulative use of rotation estimates, leading to a rapid accumulation of error. In contrast, the use of allothetic cues allows the constant calibration of an animal's direction estimate relative to a stable external cue, so directional error should not accumulate. Irrespective of the type of directional cue used, errors around the estimation of distance travelled should always accumulate linearly with increasing outward path length. Therefore, animals using allothetic directional cues to path integrate should exhibit a linear accumulation of error around their estimated home location with increasing outward path length. In contrast, animals using idiothetic cues should exhibit a more rapid accumulation of error with increasing outward path directional change as well (Cheung, 2014; Heinze et al., 2018).

Path integration is a taxonomically widespread strategy likely to be used by animals living in habitats where visual landmark use is unreliable. Studies in desert ants demonstrated path integration using a sky compass and a stride integrator as an odometer (Wehner & Müller, 2006; Wittlinger et al., 2007), research in golden hamsters revealed the use of a single light source as a directional cue and self-motion information as an odometer for path integration (Etienne et al., 1990), investigations in stomatopods observed path integration underwater for the first time, albeit in a surface-bound animal (Patel & Cronin, 2020b), and studies in cichlid fish found initial evidence consistent with the use of path integration in aquatic vertebrates (Sibeaux et al., 2024). However, clear evidence for the use of path integration in fish is still lacking, and research in this group is neglected. Teleost fish represent half of all vertebrate species and are a highly diverse clade, but we have limited information on the diversity of their local navigational strategies (Volff, 2005). Additionally, fish exist in a three-dimensional medium and can move with six degrees of freedom, which may present additional challenges when navigating compared to surface-bound animals (Jeffery et al., 2013). Fish are therefore excellent models for navigation research, and studying path integration in fish could help determine how navigational strategies differ between animals which move in two- versus three-dimensional space.

An alternative to path integration is navigating by landmarks. This can range in complexity from simpler behaviours like beaconing, where an animal uses a landmark to identify the position of a goal, to more complex ones like piloting, where animals follow a sequence of landmarks to find a goal (Reese, 1989; Warburton, 1990). Landmarks can generally be classified as local, which directly indicate a specific location, or global, which can be used along with other cues to calculate a location (Braithwaite & Burt de Perera, 2006).

Many fish species live in coral reefs, which are landmark-rich environments. However, navigation by path integration could be beneficial for coral reef-dwelling fish in several situations. First, conditions with reduced visibility, such as increased water turbidity or plankton blooms, could make visual landmark use unreliable (Wenger & McCormick, 2013). Second, if landmarks become altered or

unstable due to erosion, coral growth, or other disturbances, having additional navigational strategies available, like path integration, could be advantageous. Furthermore, while coral reefs are structurally complex overall, they may contain relatively featureless sections where landmark cues are scarce. In these cases, path integration could aid fish in finding their way back to their home range or shelter when other navigational cues are limited or disrupted. Evidence for path integration in coral reef fish would outline its use as a navigational strategy and provide insights into the evolution of this behavior and its underlying neural mechanisms.

I developed a novel experimental paradigm to test whether the coral reef-dwelling domino damsel fish (*Dascyllus trimaculatus*) could navigate back to a shelter location using path integration. Domino damsels are a very territorial species, with their mortality rate related to intraspecific competition for shelter (Holbrook & Schmitt, 2002). Shelter use is an important part of domino damsel life history, as shelters or rocky surfaces are used as a substrate to lay eggs. These fish tend to leave their shelter to forage and return to it for safety. I expected that it would be beneficial for domino damsels to path integrate, as it would allow them to return to safety rapidly and efficiently when encountering a predator or competitor. I therefore hypothesise that domino damsels can navigate back to a shelter location using path integration.

Methods

Overview

I tested whether domino damsel fish can navigate back to a shelter using path integration. I first allowed individuals to acclimatise to a circular experimental pool with two landmarks and a shelter overnight. The next morning, I placed a trap with food rewards in the centre of the pool. I recorded the fish as they swam from their shelter into the trap. Once trapped, I removed the landmarks and shelter from the pool. Fish were then released from the trap and free navigation movements were recorded and analysed.

Animal Husbandry

I used 10 domino damsel fish (*Dascyllus trimaculatus*) in experiments sourced from a supplier (Tropical Marine Centre, Solesbridge Lane, Chorleywood, Hertfordshire, WD3 5SX). Individuals were wild caught and kept by the supplier for one month before being housed in the laboratory.

I housed individuals in separate 0.35 m x 0. 32 m x 0.60 m tanks within a single flow-through marine system. I enriched each tank with 1 cm depth of coral gravel substrate, a large rock, a cave shelter (13 cm x 12 cm x 8 cm Aqua One Marble Cave), and a 30 cm tall plastic plant. I kept the fish on a 12 h light/12 h dark cycle with fluorescent light, and fed individuals with crisp (TetraPRO Energy Multi-Crisps) in the morning and mysis shrimp in the afternoon to provide supplementary nutrients. I maintained water parameters at healthy levels for this species (Temperature: 26.0 °C, Salinity

112 (specific gravity): 1.024, pH: 8.2, KH: 7-8 dKH, Nitrite: 0 ppm, Ammonia: 0 ppm, Nitrate: <20 ppm)

113 by carrying out water tests, water changes, and tank cleans once per week. Room temperature,

114 water temperature, and individual health were checked at each feed. I quarantined all individuals in

115 this system for 4 weeks before starting experiments.

Experimental Apparatus

I ran experiments in a 2 m-diameter blue plastic circular pool (Figure 2), which I filled with 10 cm

depth coral gravel substrate and 20 cm depth water (from water surface to top of substrate). I kept

pool water at identical parameters to the home tank system. To help maintain these parameters, I

placed a water heater and a filter pump in the pool (Figure 2).

During the acclimation period and in experiments, I placed two 14.0 cm x 14.6 cm x 18.1 cm

polyurethane resin replica green brain coral landmarks and a shelter (14 cm x 11 cm x 8 cm Aqua

One Marble Cave) in the pool. To reduce the shelter's saliency, I covered it with coral gravel substrate

using silicone. In this experiment it was important for the shelter to be inconspicuous to prevent fish

from remembering it as a salient landmark or beacon. In addition to covering the shelter with

substrate, the shelter was slightly buried (i.e. positioned below the substrate level) forcing the fish to

Figure 2 - *Picture taken on the GoPro showing the experimental pool prepared for fish acclimation.*

search for it when exploring the experimental pool. To reduce the likelihood of individuals using global visual cues from the laboratory room, I surrounded the pool with white curtains (Figure 2).

To record footage during acclimation and experiments, I positioned a GoPro Hero 8 Black 2 m above the centre of the pool (recording at 1440p, 30 fps, Linear FOV, see full view in Figure 2). To provide a low-latency live video feed of individual fish location, I positioned a USB webcam next to the GoPro and connected it to a laptop outside of the curtains. This second camera allowed me to monitor fish movement in real time for trapping.

Acclimation

To allow fish to acclimatise to the pool overnight, I moved individuals from their home tank to the experimental pool in the afternoon the day before an experiment. I ensured that water conditions were identical before moving an individual into the pool to acclimatise by measuring temperature, salinity, and pH in the tank and pool. Once moved, I fed the fish and recorded the following hour of the acclimation period, allowing me to calculate an exploratory score for each individual.

Experimental Setup

Before beginning an experimental trial, I measured pool water temperature. To eliminate confounding navigational cues within the pool, I removed the water heater and filter pump for the duration of the experimental trial. Stable external directional cues are essential for precise path integration in many species and the location of the sun, or even a single light source, can be used as a directional cue in animal path integration (Etienne et al., 1990; Heinze et al., 2018; Patel & Cronin, 2020b). To provide a directional cue analogous to the sun, I positioned a spotlight outside the experimental enclosure at a height of 128 cm and a horizontal distance 48 cm away from the pool, facing the curtains with a 20° negative inclination. This created a localised bright spot that the fish could use as a directional cue or heading indicator for path integration (Goodyear & Ferguson, 1969; Guilford & Taylor, 2014).

I then placed the trap into the centre of the pool (Figure 3a). The trap consisted of a 12.5 cm diameter by 25.3 cm height white opaque plastic cylinder with a 6.4 cm diameter entrance hole cut into one side 1.6 cm from the bottom. I used the trap to contain the fish during the trial and prevent it from directly observing any experimental manipulations. To get the trap to appear more like a natural shelter, I covered the exterior and interior with the coral gravel substrate. There was a 17.5 cm by 7.5 cm rectangular white opaque plastic door with attached steel weights behind the trap's entrance hole, which opened or closed by sliding along plastic rails (Figure 3a). I was able to open, close, and remove the trap remotely by using fishing lines attached via a pulley system. To provide additional enrichment and encourage the fish to explore the central area of the pool, I placed a plastic plant by the door (Figure 3a). I then started each trial by setting the GoPro to record, adding mysis shrimp inside the trap, and remotely raising the door.

Experimental Procedure

I recorded fish movement in each trial for up to two hours, and ended experimental trials if individuals
did not enter the trap in this period. As soon as I observed fish entering the trap on the live low-
latency USB camera feed, I closed the trap remotely. I then removed the landmarks, the shelter, and
the plastic plant (Figure 3b). To prevent fish from using chemical cues from the water or visual
patterns in the substrate to identify the shelter location, I also mixed and smoothed out any disturbed
substrate and water near the landmark and shelter locations. Finally, I released the fish by lifting the
trap vertically out of the water using the pulley system. I recorded five minutes of fish free movement
before ending the trial and used a large jug to return the fish to its home tank.

Fish Tracking and Trajectory Analyses

172 Over 6 weeks of testing and 21 experimental trials attempted, only 6 of the 10 fish tested entered
173 the trap. The following analyses have therefore been performed on the trajectories collected from
174 these 6 individuals.

Figure 3 - Diagram showing (a) the layout of the pool at the start of an experimental trial,
and (b) the layout of the pool after removing the shelter and landmarks and releasing the fish
by vertically lifting out the trap.

175 Fish movement in the acclimation and pre-trapping part of the experimental trial was tracked using
176 AnimalTA (Chiara & Kim, 2023). Fish movement after release from the trap was recorded manually
177 in MATLAB (MathWorks, 2023), and coordinates were extracted every second for one minute. I
178 tracked post-release movement manually as the background subtraction tracking method used by
179 AnimalTA was not suitable once the trap was removed from the water during a trial.

I focused on and analysed three navigational periods in detail:

1) Acclimation: To characterise fish exploratory behaviour during the acclimation period, I calculated average speed (cm/s), total distance travelled (cm), and area of the pool explored (cm^2) in AnimalTA. I also recorded exploration latency (the first time at which each fish exited the shelter and became entirely visible on camera, s) and measured the greatest distance it swam from its shelter (cm). Each fish was given an exploratory index calculated by multiplying normalised versions of the above metrics.

2) Last outward trajectory: To characterise fish behaviour during the experiment before the fish was trapped, I analysed fish movement during the last outward trajectory. This is defined as the period between the fish leaving its shelter for the last time during the trial and it being trapped in the centre. I calculated the length (cm) and time taken (s) for the last outward trajectories in AnimalTA. To obtain measures of the tortuosity of the last outward trajectories, I calculated their straightness indices and sinuosities in R using the 'trajr' package (version 1.5.1; McLean & Skowron Volponi, 2018). The straightness index is defined as $\frac{D}{L}$, where D is the beeline distance between the first and last points in the trajectory and L is the path length travelled (Batschelet, 1981). Sinuosity, defined by Benhamou (2004), uses the mean step length, the mean cosine of turning angles, and the coefficient of variation of step length (standard deviation/mean) to calculate a measure of tortuosity.

3) Homeward path: To characterise fish behaviour after being released from the trap, I analysed manually recorded fish coordinates (extracted with MATLAB) using the 'trajr' package in R (McLean & Skowron Volponi, 2018; R Core Team, 2022). I used the above metrics of straightness index and sinuosity to determine the transition point between the homeward path ending and the initiation of search behaviour. I identified this transition through visual inspection of a graph of straightness index and sinuosity over time, and defined the end of the homeward path based on when two sequential coordinates showed a breakdown in straightness index or increase in sinuosity after a plateau (Figure 4).

Figure 4 - *Plots of how the straightness index (blue) and sinuosity (red) of the post-release path of each fish develop over time. The first shared large deviation in straightness index or sinuosity (dotted line) was used to define the end point of an individual's homeward path.*

Once the homeward paths were identified, I calculated the homeward path direction (angle between first and last homeward path coordinates, degrees), beeline distance (straight line distance between first and last homeward path coordinates, cm), length (cm), time (s), mean speed (cm/s), straightness index, and sinuosity. I then used the homeward path direction to calculate homeward direction error (absolute angular difference between homeward path direction and shelter direction, degrees). I also calculated the homeward beeline distance error (absolute difference between shelter distance and homeward beeline distance, cm) and the homeward path length error (absolute difference between shelter distance and homeward path length, cm).

Statistical analysis

All statistics were carried out in R (version 4.2.2; R Core Team, 2022). I used circular statistics to analyse direction data, and the significance threshold α was set at 0.05.

1 – Homeward path direction analysis

To investigate whether fish homeward path directions followed a uniform, unimodal, or bimodal distribution, I used a model-based approach with maximum likelihood with the R package 'CircMLE' (version 0.3.0; Fitak & Johnsen, 2017). The fish direction data were converted to circular class data using the package 'circular' (version 0.5.0; Agostinelli & Lund, 2023). These directions were then

222 compared to ten direction models and the best fitted model was estimated using AIC values (for
223 more details see Fitak & Johnsen, 2017). I also conducted a Rayleigh test of uniformity to determine
224 if the distribution of homeward path directions differed significantly from a uniform distribution.

2 – Homeward path direction relative to shelter

I next analysed whether homeward path directions differed significantly from direction to the shelter.
First, I conducted a Watson's two-sample test of homogeneity between the distribution of homeward
path directions and a simulated distribution centred on the shelter location with the same
concentration parameter as the homeward path directions. The concentration parameter used
(3.095) was estimated during the maximum likelihood analysis with 'CircMLE' (Table S1).

Second, I calculated a circular 95% confidence interval around the mean homeward path direction
to see if this contained the shelter direction. Mean direction was calculated using the R package
'circular' (version 0.5.0; Agostinelli & Lund, 2023), and confidence intervals were calculated by
bootstrapping with replacement over 1000 iterations using the package 'boot' (version 1.3.28; Angelo
Canty & B. D. Ripley, 2024).

3 – Effect of last outward trajectory behaviour on homeward path behaviour

I next tested how homeward path behaviour was influenced by last outward trajectory behaviour by
fitting a series of linear models using the R package 'lme4' (Bates et al., 2015). I tested whether a
selection of biologically relevant last outward trajectory metrics set as explanatory variables (last
outward trajectory length, time, straightness index, and sinuosity) had any influence on homeward
path behaviour (response variables included: homeward direction error, homeward beeline distance
error, homeward path length error, time, mean speed, straightness index, and sinuosity). For all linear
models fitted see Table S2.

4 - Homeward path distance relative to shelter

245 I then analysed whether homeward path beeline distance or homeward path length differed
246 significantly from distance to the shelter. I conducted a Welch Two Sample t-test between homeward
247 path beeline distances and shelter distance. I also conducted a Wilcoxon rank sum exact test
248 between homeward path lengths and shelter distance, as homeward path lengths were not normally
249 distributed (Shapiro-Wilk normality test: W = 0.77291, p = 0.0331).

5 - Effect of acclimation on homeward path behaviour

I then tested how homeward path behaviour was influenced by acclimation behaviour by fitting a
series of linear models. I tested whether a selection of biologically relevant acclimation metrics set
as explanatory variables (acclimation distance travelled, area explored, and exploratory index) had
any influence on homeward path behaviour (response variables included: homeward direction error,
homeward beeline distance error, homeward path length error, total time, mean speed, straightness
index, and sinuosity). For all linear models fitted see Table S2.

6 – Power analyses

I finally conducted post hoc power analyses to determine whether my experiment had an appropriate sample size to observe significant effects of exploratory and outward path behaviour on homing behaviour. I used the package 'lme4' to fit my models and determined the power of these models with the '*powerSim*' function in the R package 'simr' (Green & MacLeod, 2016). For the results of the power analyses, see the supplementary material.

Results

1 – Homeward path directions followed a unimodal distribution

The directions fish swam on their homeward paths were best described by a unimodal distribution following a circular maximum likelihood approach (ΔAIC = 0, φ_1 = 6.049, κ_1 = 3.095, Table S1, Figure S1). The remaining circular distribution models fitted poorly described the homeward path direction data (ΔAIC $\geq$ 2.145, Table S1). I confirmed the non-uniformity of homeward path directions with a Rayleigh test (r = 0.8168, p = 0.0112). The homeward and search paths of each fish are shown in Figure 5a.

2 - Homeward path directions were oriented towards the shelter location

I did not find any significant difference between homeward path directions and shelter direction after conducting a Watson's two-sample test of homogeneity between the homeward path directions and a simulated distribution centred on the shelter location with the same concentration parameter as the homeward path directions (Test statistic = 0.0486, Level 0.05 critical value = 0.187). Additionally, the direction of the shelter location was contained within a circular 95% bootstrap confidence interval for the mean homeward path direction (Figure 5b).

Figure 5 - (a) *Homeward (red)* *and* *search (grey) paths of each fish after release from the trap. Food trap location (circle) and previous shelter location (triangle) are also shown.* ***(b) Shelter angle is contained within the 95% confidence interval of mean homeward path direction.*** *Homeward path directions (black dots)*, *mean homeward path direction (black arrow)*, *95% confidence interval of mean homeward path direction (red arrows)*, *and* *shelter direction (blue arrow)* *are shown.*

3 - Homeward direction error increased with last outward trajectory length

As hypothesised, homeward direction error increased with the length of the last outward trajectory.
The linear model fitted between last outward trajectory length and homeward direction error explains
a statistically significant and substantial proportion of variance (R^2 = 0.74, $F_{(1, 4)}$ = 11.42, p = 0.028,
adjusted R^2 = 0.60). Within this model the effect of last outward trajectory length is statistically
significant and positive (β = 0.01, 95% CI [1.90 x 10^{-3}, 0.02], $t_{(4)}$ = 3.38, p = 0.028; Std. β = 0.86, 95%
CI [0.15, 1.57], Figure 6). None of the other models fitted showed any significant effect of last outward
trajectory behaviour on homeward path behaviour (Table S2).

4 - Homeward path lengths and beeline distances were significantly greater than
shelter distance

Both homeward beeline distance and homeward path length were significantly greater than the
distance to the shelter. The Welch Two Sample t-test suggested that beeline length is significantly
greater than shelter distance (mean of beeline length = 88.17, mean of shelter distance = 78.00,
difference = 10.17, 95% CI [1.51, 18.83], t(6.05) = 2.87, p = 0.028; Cohen's d = 1.66, 95% CI [0.16,
3.07]). Additionally, the Wilcoxon rank sum exact test result suggested that homeward path length is
significantly greater than beeline length (W = 32.00, p = 0.026; r (rank biserial) = 0.78, 95% CI [0.34,
0.94]).

Figure 6 - Homeward path direction error increases with last outward trajectory length.

5 - Acclimation behaviour did not significantly influence homeward path behaviour

No significant effects of any of the biologically relevant acclimation explanatory variables (acclimation distance travelled, area explored, and exploratory index) were observed on any of the homeward path response variables (homeward path length error, homeward beeline distance error, total time, mean speed, straightness index, and sinuosity). Results for all of the linear models fitted can be found in Table S2.

Discussion

The results of this study provide evidence that the coral reef-dwelling domino damselfish (*Dascyllus trimaculatus*) can exhibit homing behaviour consistent with the use of path integration. After being released from a central location with no visual landmarks or their previous shelter present, the fish swam in a unimodally clustered distribution of directions that was not significantly different from the actual direction towards their previous shelter. This suggests the fish were attempting to navigate directly back to their shelter location rather than orienting randomly.

The error in the directional accuracy of the fish's homeward paths increased with the length of the last outward trajectory taken before being released from the central trap location. This relationship between outward path length and homing error fits the expectations of path integration and has been observed in many species, where errors in an internally calculated estimation of directional change

and distance covered accumulate over larger travel distances due to noise (Cheung, 2014; Heinze
et al., 2018; Merkle et al., 2006). Noise is present at all physical scales and impacts every level of
the nervous system. Its manifestations include fundamental quantum uncertainty, stochasticity of
neural spikes, and environmental perturbation, distortion, or obstruction of sensory cues (Faisal et
al., 2008). In path integration, noise can take the form of random environmental disturbances or
alterations of path integration inputs, computations, or outputs within the sensorimotor system.
These cannot be predicted or detected by the animal so result in error around a home vector. While
exploratory behaviour during acclimation did not significantly influence homing accuracy, effects may
emerge with larger sample sizes (see power analyses in the supplementary material).

Many other navigational strategies can be ruled out or seem unlikely based on the experimental
conditions used and results obtained. The use of local landmark-based navigational strategies like
piloting and beaconing is highly unlikely due to the removal of the shelter and landmarks as visual
cues. Additionally, the curtain surrounding the pool would limit the use of global landmarks outside
the arena, except for the provided spotlight. The lack of influence acclimation behaviour appears to
have on homeward movement, combined with the relatively homogeneous nature of the pool
environment, casts doubt on the possibility that fish navigate to their shelter by learning a map based
on cues in their environment (for example, based on the geometric relationships of objects present).
The use of olfactory navigation is also unlikely due to the deliberate mixing of water and substrate to
disrupt olfactory cues. While the fish could theoretically use magnetoreception, sound, or
electroreceptive cues for homing, there is no direct evidence supporting reliance on these strategies.
Though it is possible that geomagnetic cues could be used as a compass cue for path integration,
the use of a magnetic coordinate map to navigate is unlikely due to the small spatial scale of the
experimental pool and the nature of the homing behaviour observed. The use of hydrodynamic cues
to navigate is also improbable due to the still water conditions in the pool. As homing error increases
with longer outward path lengths, path integration emerges as the most viable and parsimonious
explanation for the observed homing behaviours in this experimental setting.

Path integration is a robust navigational strategy, allowing an animal to return to the starting point of
a journey without using familiar positional cues. It works through continuously estimating position
relative to a reference point using only directional and distance cues derived from an animal's own
locomotion. This enables an animal to efficiently return home along the most direct route at any place
or moment on its journey, even in featureless or unfamiliar terrain (Etienne & Jeffery, 2004; Müller &
Wehner, 1988b). Path integration can become more reliable when combined with other strategies,
like using landmarks or environmental cues to calibrate a homing vector (Etienne et al., 1996; Patel
& Cronin, 2020a). Due to its ability to be used with other navigational strategies and its presence in
disparate animal taxa, it is unsurprising that path integration might be one of many navigational
strategies employed by fish.

Despite the fact that animals can path integrate using purely self-generated (or idiothetic) directional cues, it is unlikely that this was the case for the individuals in this experiment. The fact that idiothetic rotational estimates must be used cumulatively, together with the impact of noise on these estimates, means idiothetic path integration leads to rapid accumulation of error around the animal's direction estimate. In contrast, the use of a stable external (or allothetic) directional cue allows the constant calibration of an animal's direction estimate relative to this cue, so directional error does not accumulate. Irrespective of the type of directional cue used, errors around an animal's estimation of distance travelled always accumulate linearly with increasing outward path length (Cheung et al., 2007; Heinze et al., 2018). Due to the lack of significant impact of last outward trajectory straightness or sinuosity on homeward direction error, together with the significant linear increase of homeward direction error with increasing last outward trajectory length, the use of stable external directional cues appears more likely than the use of self-generated idiothetic directional cues in this experiment.

Both the spotlight provided during experimental trials and the Earth's magnetic field could have served as directional cues for fish path integration in this experiment. Mosquitofish and rodents have been shown to use point light sources or sun-analogous cues to estimate their direction (Etienne et al., 1990; Goodyear & Ferguson, 1969). Therefore, the spotlight provided in this experiment may have functioned similarly as a global landmark or heading indicator that individuals could use as a directional cue for path integration (Guilford & Taylor, 2014). Additionally, the use of magnetoreception is widespread among fish, so individuals could have used geomagnetic cues as a compass for path integration (Naisbett-Jones & Lohmann, 2022). The use of other compass or directional cues, like the position of stars, skylight polarisation, a self-generated cognitive map, or odour cues, was highly unlikely due to the nature of the experimental conditions and behaviour observed.

As well as estimating direction, path integration also requires an animal to estimate distance travelled. Fish could use a wide variety of cues to estimate this, including optic flow, proprioception, energy expenditure, vestibular cues, or water flow. Recent research has demonstrated that fish can estimate their distance travelled using optic flow (Karlsson et al., 2022; Sibeaux et al., 2022). While other mechanisms like proprioception, energy expenditure, and vestibular cues can be used to estimate distance travelled in terrestrial animals, they are yet to be explored in fish (Jacob et al., 2017; Proffitt et al., 2003; Wittlinger et al., 2007). Lastly, although the use of water flow cues to estimate distance travelled may be plausible in fish, evidence supporting this behaviour is lacking.

While providing novel evidence supporting the use of path integration in coral reef fish, there are limitations to the current study that could be addressed through future research. First, the specific compass and odometer cues used for path integration in fish remain unidentified. Systematically manipulating potential external directional cues, like shifting the position of the artificial light source or altering the local magnetic field, could determine which directional cues are prioritised.

Additionally, altering potential odometer cues, like changing the visual characteristics of the pool
environment or altering the flow of water, could discern how these fish estimate distance travelled.

Second, future research could better characterise how error accumulates around the estimated
length of the home vector. While I found that homeward path length and homeward beeline distance
both differed significantly from shelter distance, it's possible that fish homing behaviour was
influenced by the relatively close proximity of the pool wall. Individuals may have associated the pool
wall with their shelter location, so may have continued toward the wall having followed a homing
vector. Similar behaviour has been observed in hamsters: when startled in an experiment, individuals
followed a homing direction towards the arena edge, then searched for their nest entrance by
following the arena border (Siegrist et al., 2003). Furthermore, any stress associated with being
trapped in the centre or the absence of their shelter may have induced a wall-following behavioural
response (thigmotaxis), leading to fish continuing on toward the wall after following their homing
vector (Champagne et al., 2010; Sharma et al., 2009). Future experiments in a larger arena could
position the shelter further from any walls to both limit any influence of walls on homing, and to more
definitively characterise path integration behaviour in fish. For example, this would allow better
investigation into how homeward path length and beeline distance compare to shelter distance, and
if error in these distance measures relative to shelter distance also accumulates linearly with outward
path length. This could also enable the characterisation of systematic search behaviour in fish
without the influence of a nearby wall.

Third, experiments introducing controlled outward paths with predetermined lengths and tortuosities,
instead of variably-terminating last outward trajectories, could more precisely characterise how
homing error accumulates with increasing outward path length and directional change. Encouraging
fish to swim through a series of controlled outward paths of predetermined lengths could more
precisely characterise how path integration error accumulates with increasing outward path length.
Additionally, allowing fish to swim through controlled outward paths of differing tortuosities could
identify whether fish use stable external directional cues or self-generated cues to estimate direction
when path integrating. However, restricting the movement of fish using controlled paths could lead
to biased results. Individuals might follow the boundaries of any controlled paths, and the necessary
recording of other stimuli (like the position of path boundaries) might distract from direction or
distance monitoring.

Furthermore, fish exist in a three-dimensional medium and can move with six degrees of freedom,
but this experiment only analyses movement in two dimensions. Future research could build on my
initial findings that fish follow homing trajectories consistent with path integration and use an
experimental arena with greater three-dimensional complexity to explore how this navigational
behaviour changes with greater use of an additional spatial dimension. Lastly, increased sample
sizes would likely provide greater statistical power to detect any weaker effects of acclimation or
outward path behaviour on homing behaviour (see power analysis in the supplementary materials).

By building upon this foundation with targeted experiments, we can gain deeper knowledge into the

specific sensorimotor mechanisms underpinning path integration in coral reef fish.

Despite these limitations, my results expand our understanding of path integration abilities to include

coral reef fish. Path integration has previously been demonstrated in terrestrial invertebrates like

desert ants, terrestrial vertebrates like rodents, and marine arthropods like stomatopod crustaceans

(Etienne et al., 1990; Müller & Wehner, 1988b; Patel & Cronin, 2020b). The finding that a coral reef

fish possesses similar path integration capabilities implies this navigational strategy either evolved

early in animals or potentially multiple times through convergent evolution across disparate taxa.

From an applied perspective, understanding the sensory cues and mechanisms behind fish path

integration could aid in predicting patterns of fish movement and distribution, which is relevant for

ecology, conservation, and fisheries management (Boström et al., 2011). Theoretically, details on

how path integration functions in fish could help reveal the evolutionary origins of this ability and

shed light on the neural processes governing spatial navigation systems.

In summary, this study provides novel evidence that the coral-reef dwelling domino damsel fish

Dascyllus trimaculatus possess the ability to follow a homing trajectory consistent with path

integration over spatial scales relevant to their ecology and life history. Future research could

investigate the specific odometer and compass cues these fish use for path integration. Overall, the

results add coral reef fish to the list of taxa employing this widespread navigation strategy.

References

Agostinelli, C., & Lund, U. (2023). *R package circular: Circular Statistics (version 0.5-0)*. https://CRAN.R-project.org/package=circular

Angelo Canty, & B. D. Ripley. (2024). *boot: Bootstrap R (S-Plus) Functions*.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Batschelet, Edward. (1981). *Circular statistics in biology* [Book]. Academic Press.

Benhamou, S. (2004). How to reliably estimate the tortuosity of an animal's path: *Journal of Theoretical Biology*, *229*(2), 209–220. https://doi.org/10.1016/j.jtbi.2004.03.016

Boström, C., Pittman, S., Simenstad, C., & Kneib, R. (2011). Seascape ecology of coastal biogenic habitats: advances, gaps, and challenges. *Marine Ecology Progress Series*, *427*, 191–217. https://doi.org/10.3354/meps09051

Braithwaite, V. A., & Burt de Perera, T. (2006). Short-range orientation in fish: How fish map space. *Marine and Freshwater Behaviour and Physiology*, *39*(1), 37–47. https://doi.org/10.1080/10236240600562844

Champagne, D. L., Hoefnagels, C. C. M., de Kloet, R. E., & Richardson, M. K. (2010). Translating rodent behavioral repertoire to zebrafish (Danio rerio): Relevance for stress research. *Behavioural Brain Research*, *214*(2), 332–342. https://doi.org/10.1016/j.bbr.2010.06.001

Cheung, A. (2014). Animal path integration: A model of positional uncertainty along tortuous paths. *Journal of Theoretical Biology*, *341*, 17–33. https://doi.org/10.1016/j.jtbi.2013.09.031

Cheung, A., Zhang, S., Stricker, C., & Srinivasan, M. V. (2007). Animal navigation: the difficulty of moving in a straight line. *Biological Cybernetics*, *97*(1), 47–61. https://doi.org/10.1007/s00422-007-0158-0

Chiara, V., & Kim, S. (2023). <scp>AnimalTA</scp> : A highly flexible and easy-to-use program for tracking and analysing animal movement in different environments. *Methods in Ecology and Evolution*, *14*(7), 1699–1707. https://doi.org/10.1111/2041-210X.14115

Etienne, A. S., & Jeffery, K. J. (2004). Path integration in mammals. *Hippocampus*, *14*(2), 180–192. https://doi.org/10.1002/hipo.10173

Etienne, A. S., Maurer, R., & Séguinot, V. (1996). Path Integration in Mammals and its Interaction With Visual Landmarks. *Journal of Experimental Biology*, *199*(1), 201–209. https://doi.org/10.1242/jeb.199.1.201

Etienne, A. S., Teroni, E., Hurni, C., & Portenier, V. (1990). The effect of a single light cue on homing behaviour of the golden hamster. *Animal Behaviour*, *39*(1), 17–41. https://doi.org/10.1016/S0003-3472(05)80723-7

Faisal, A. A., Selen, L. P. J., & Wolpert, D. M. (2008). Noise in the nervous system. *Nature Reviews Neuroscience*, *9*(4), 292–303. https://doi.org/10.1038/nrn2258

Fitak, R. R., & Johnsen, S. (2017). Bringing the analysis of animal orientation data full circle: model-based approaches with maximum likelihood. *Journal of Experimental Biology*. https://doi.org/10.1242/jeb.167056

Goodyear, C. P., & Ferguson, D. E. (1969). Sun-compass orientation in the mosquitofish, Gambusia affinis. *Animal Behaviour*, *17*(4), 636–640. https://doi.org/10.1016/S0003-3472(69)80005-9

Green, P., & MacLeod, C. J. (2016). <scp>SIMR</scp> : an R package for power analysis of
generalized linear mixed models by simulation. *Methods in Ecology and Evolution*, *7*(4), 493–
498. https://doi.org/10.1111/2041-210X.12504

Guilford, T., & Taylor, G. K. (2014). The sun compass revisited. *Animal Behaviour*, *97*, 135–143.
https://doi.org/10.1016/j.anbehav.2014.09.005

Heinze, S., Narendra, A., & Cheung, A. (2018). Principles of Insect Path Integration. *Current*
Biology, *28*(17), R1043–R1058. https://doi.org/10.1016/j.cub.2018.04.058

Holbrook, S. J., & Schmitt, R. J. (2002). Competition for Shelter Space Causes Density-Dependent
Predation Mortality in Damselfishes. *Ecology*, *83*(10), 2855. https://doi.org/10.2307/3072021

Jacob, P.-Y., Gordillo-Salas, M., Facchini, J., Poucet, B., Save, E., & Sargolini, F. (2017). Medial
entorhinal cortex and medial septum contribute to self-motion-based linear distance
estimation. *Brain Structure and Function*, *222*(6), 2727–2742. https://doi.org/10.1007/s00429-
017-1368-4

Jeffery, K. J., Jovalekic, A., Verriotis, M., & Hayman, R. (2013). Navigating in a three-dimensional
world. *Behavioral and Brain Sciences*, *36*(5), 523–543.
https://doi.org/10.1017/S0140525X12002476

Karlsson, C., Willis, J., Patel, M., & de Perera, T. B. (2022). Visual odometry of Rhinecanthus
aculeatus depends on the visual density of the environment. *Communications Biology*, *5*(1),
1045. https://doi.org/10.1038/s42003-022-03925-5

McLean, D. J., & Skowron Volponi, M. A. (2018). trajr: An R package for characterisation of animal
trajectories. *Ethology*, *124*(6), 440–448. https://doi.org/10.1111/eth.12739

Merkle, T., Knaden, M., & Wehner, R. (2006). Uncertainty about nest position influences systematic
search strategies in desert ants. *Journal of Experimental Biology*, *209*(18), 3545–3549.
https://doi.org/10.1242/jeb.02395

Müller, M., & Wehner, R. (1988a). Path integration in desert ants, *Cataglyphis fortis*. *Proceedings*
of the National Academy of Sciences, *85*(14), 5287–5290.
https://doi.org/10.1073/pnas.85.14.5287

Müller, M., & Wehner, R. (1988b). Path integration in desert ants, *Cataglyphis fortis*. *Proceedings*
of the National Academy of Sciences, *85*(14), 5287–5290.
https://doi.org/10.1073/pnas.85.14.5287

Naisbett-Jones, L. C., & Lohmann, K. J. (2022). Magnetoreception and magnetic navigation in
fishes: a half century of discovery. *Journal of Comparative Physiology A*, *208*(1), 19–40.
https://doi.org/10.1007/s00359-021-01527-w

Patel, R. N., & Cronin, T. W. (2020a). Landmark navigation in a mantis shrimp. *Proceedings of the*
Royal Society B: Biological Sciences, *287*(1936), 20201898.
https://doi.org/10.1098/rspb.2020.1898

Patel, R. N., & Cronin, T. W. (2020b). Mantis Shrimp Navigate Home Using Celestial and Idiothetic
Path Integration. *Current Biology*, *30*(11), 1981-1987.e3.
https://doi.org/10.1016/j.cub.2020.03.023

Patel, R. N., & Cronin, T. W. (2020c). Path integration error and adaptable search behaviors in a
mantis shrimp. *Journal of Experimental Biology*. https://doi.org/10.1242/jeb.224618

Proffitt, D. R., Stefanucci, J., Banton, T., & Epstein, W. (2003). The Role of Effort in Perceiving
Distance. *Psychological Science*, *14*(2), 106–112. https://doi.org/10.1111/1467-9280.t01-1-
01427

R Core Team. (2022). *R: A Language and Environment for Statistical Computing*. https://www.R-project.org/

Reese, E. S. (1989). Orientation behavior of butterflyfishes (family Chaetodontidae) on coral reefs: spatial learning of route specific landmarks and cognitive maps. *Environmental Biology of Fishes*, *25*(1–3), 79–86. https://doi.org/10.1007/BF00002202

Sharma, S., Coombs, S., Patton, P., & de Perera, T. B. (2009). The function of wall-following behaviors in the Mexican blind cavefish and a sighted relative, the Mexican tetra (Astyanax). *Journal of Comparative Physiology A*, *195*(3), 225–240. https://doi.org/10.1007/s00359-008-0400-9

Sibeaux, A., Karlsson, C., Newport, C., & Burt de Perera, T. (2022). Distance estimation in the goldfish (*Carassius auratus*). *Proceedings of the Royal Society B: Biological Sciences*, *289*(1984). https://doi.org/10.1098/rspb.2022.1220

Sibeaux, A., Newport, C., Green, J., Engelmann, J., & Perera, T. B. de. (2023). *Taking a shortcut: evidence of path integration in fish*. https://doi.org/10.21203/rs.3.rs-3234472/v1

Siegrist, C., Etienne, A. S., Boulens, V., Maurer, R., & Rowe, T. (2003). Homing by path integration in a new environment. *Animal Behaviour*, *65*(1), 185–194. https://doi.org/10.1006/anbe.2002.2036

The MathWorks Inc. (2023). *MATLAB Version: 23.2.0.2459199 (R2023b) Update 5*. The MathWorks Inc. https://www.mathworks.com

Volff, J.-N. (2005). Genome evolution and biodiversity in teleost fish. *Heredity*, *94*(3), 280–294. https://doi.org/10.1038/sj.hdy.6800635

Warburton, K. (1990). The use of local landmarks by foraging goldfish. *Animal Behaviour*, *40*(3), 500–505. https://doi.org/10.1016/S0003-3472(05)80530-5

Wehner, R., & Müller, M. (2006). The significance of direct sunlight and polarized skylight in the ant's celestial system of navigation. *Proceedings of the National Academy of Sciences*, *103*(33), 12575–12579. https://doi.org/10.1073/pnas.0604430103

Wehner, R., & Srinivasan, M. V. (1981). Searching behaviour of desert ants, genus Cataglyphis (Formicidae, Hymenoptera). *Journal of Comparative Physiology ? A*, *142*(3), 315–338. https://doi.org/10.1007/BF00605445

Wenger, A. S., & McCormick, M. I. (2013). Determining trigger values of suspended sediment for behavioral changes in a coral reef fish. *Marine Pollution Bulletin*, *70*(1–2), 73–80. https://doi.org/10.1016/j.marpolbul.2013.02.014

Wittlinger, M., Wehner, R., & Wolf, H. (2007). The desert ant odometer: a stride integrator that accounts for stride length and walking speed. *Journal of Experimental Biology*, *210*(2), 198–207. https://doi.org/10.1242/jeb.02657

Acknowledgements

We thank Christine Soper for her help with animal husbandry and all the technicians from the John Kreb Field station for helping with the fish. We thank Dr Cait Newport for lending us the experimental pool and spot light and all Fish Lab people for their feedback on the project. We thank the Human Frontier Science Program and the department of Biology of the University of Oxford for funding.

Supplementary Materials

Supplementary Tables and Figures

Table S1 - Output from CircMLE. The output from the 'circ_mle' function for all 10 models of orientation using default parameters. Homeward path directions were used as the input data.

Model	M2A	M2B	M2C	M3B	M4A	M1	M4B	M3A	M5B	M5A
Params	2	2	3	3	5	3	4	0	4	2
φ_1	6.049	5.858	6.052	5.939	5.853	2.73	5.984	NA	1.816	2.832
κ_1	3.095	8.738	2.999	5.381	5.495	2.526	4.472	0	2.426	2.72
λ	1	0.5	0.749	0.5	0.749	0.251	0.749	1	0.251	0.5
φ_2	NA	NA	NA	9.081	0.723	5.872	9.125	NA	6.092	5.974
κ_2	0	0	0	0	5.152	2.526	0.475	0	2.426	2.72
Likelihood	5.835	6.908	6.573	7.102	5.122	7.679	6.715	11.027	7.094	9.805
Convergence	0	0	0	0	0	0	0	0	0	0
AIC	15.67	17.815	19.146	20.203	20.243	21.358	21.43	22.055	22.189	23.61
AICc	19.67	21.815	31.146	32.203	Inf	33.358	61.43	22.055	62.189	27.61
BIC	15.253	17.399	18.521	19.579	19.202	20.734	20.597	22.055	21.356	23.193
ΔAIC	0	2.145	3.476	4.534	4.573	5.689	5.761	6.385	6.519	7.94
ΔAICc	0	2.145	11.476	12.534	Inf	13.689	41.761	2.385	42.519	7.94
ΔBIC	0	2.145	3.268	4.325	3.949	5.48	5.344	6.801	6.103	7.94
Relative Likelihoods	1	0.342	0.176	0.104	0.102	0.058	0.056	0.041	0.038	0.019
AIC Weights	0.5165	0.17672	0.09085	0.05354	0.05248	0.03005	0.0289	0.02122	0.01984	0.0097
Evidence Ratio	NA	2.923	5.686	9.649	9.842	17.19	17.82	24.348	26.038	52.987

572   *Table S2 - Summary of results for all linear models tested.*

Response variable	Explanatory variable(s)	R²	Adj. R²	p-value
Homeward direction error (degrees)	Acclimation distance travelled (cm)	0.09093	-0.13630	0.56140
	Acclimation area explored (cm²)	0.19630	-0.00462	0.37890
	Exploratory index	0.19860	-0.00172	0.37580
	Last outward trajectory length (cm)	0.74060	0.67580	**0.02780***
	Last outward trajectory time (s)	0.64820	0.56020	0.05328
	Last outward trajectory straightness index	0.12050	-0.09940	0.50020
	Last outward trajectory sinuosity	0.00180	-0.24770	0.93630
Homeward beeline distance error (cm)	Acclimation distance travelled (cm)	0.34940	0.18670	0.21660
	Acclimation area explored (cm²)	0.29240	0.11550	0.26790
	Exploratory index	0.38180	0.22720	0.19110
	Last outward trajectory length (cm)	0.32070	0.15090	0.24100
	Last outward trajectory time (s)	0.34600	0.18260	0.21900
	Last outward trajectory straightness index	0.06634	-0.16710	0.62220
	Last outward trajectory sinuosity	0.01156	-0.23440	0.83900
Homeward path length error (cm)	Acclimation distance travelled (cm)	0.21570	0.01959	0.35300
	Acclimation area explored (cm²)	0.47950	0.34930	0.12730
	Exploratory index	0.46530	0.33160	0.13550
	Last outward trajectory length (cm)	0.02824	-0.21470	0.75000
	Last outward trajectory time (s)	0.00411	-0.24490	0.90400
	Last outward trajectory straightness index	0.00391	-0.24510	0.90600
	Last outward trajectory sinuosity	0.16590	-0.04267	0.42300
Homeward time (s)	Acclimation distance travelled (cm)	0.12680	-0.09146	0.48840
	Acclimation area explored (cm²)	0.16130	-0.04844	0.43000
	Exploratory index	0.26760	0.08444	0.29330
	Last outward trajectory length (cm)	0.07201	-0.16000	0.60710
	Last outward trajectory time (s)	0.00534	-0.24330	0.89060
	Last outward trajectory straightness index	0.00018	-0.24980	0.97990
	Last outward trajectory sinuosity	0.04756	-0.19060	0.67800
Homeward mean speed (cm/s)	Acclimation distance travelled (cm)	0.00080	-0.24900	0.95750
	Acclimation area explored (cm²)	0.00754	-0.24060	0.87010
	Exploratory index	0.00622	-0.24220	0.88190
	Last outward trajectory length (cm)	0.03717	-0.20350	0.71440
	Last outward trajectory time (s)	0.00073	-0.24910	0.95940
	Last outward trajectory straightness index	0.00643	-0.24200	0.88000
	Last outward trajectory sinuosity	0.00303	-0.24620	0.91800
Homeward path straightness index	Acclimation distance travelled (cm)	0.08362	-0.14550	0.57830
	Acclimation area explored (cm²)	0.27620	0.09522	0.24830
	Exploratory index	0.23990	0.04990	0.32400
	Last outward trajectory length (cm)	0.10420	-0.11980	0.53260
	Last outward trajectory time (s)	0.05641	-0.17950	0.65040
	Last outward trajectory straightness index	0.00122	-0.24850	0.94765
	Last outward trajectory sinuosity	0.15110	-0.06110	0.44627

Response variable	Explanatory variable	R²	Adj. R²	p-value
Homeward path sinuosity	Acclimation distance travelled (cm)	0.05111	-0.18610	0.66670
	Acclimation area explored (cm²)	0.04035	-0.19960	0.70270
	Exploratory index	0.00089	-0.24890	0.95530
	Last outward trajectory length (cm)	0.04330	-0.19590	0.69240
	Last outward trajectory time (s)	0.05590	-0.18010	0.65190
	Last outward trajectory straightness index	0.00004	-0.24990	0.99040
	Last outward trajectory sinuosity	0.02820	-0.21470	0.75000

577 ***Figure S1 - Best fit model for homeward path directions.*** *Circular histogram (grey bars) and mean vector (red arrow)*
578 *of homeward path directions. The density (dotted line) and mean angles (dotted arrows) for the best fit model, model M2A,*
579 *are shown. The plot was produced using the 'plot_circMLE' function in the CircMLE package in R.*

Linear model power analyses

Linear Models

I conducted post hoc power analyses to determine whether my experiment was adequately powered to observe significant effects of exploratory and outward path behaviour on homing behaviour. I chose to analyse the power of four models in particular:

1. *homeward direction error ~ last outward trajectory length*. I analysed the power of this model to determine whether my experiment had an appropriate sample size to detect whether homeward direction error accumulates with outward path length. Error accumulation is expected to occur if path integration is used to navigate.

2. *homeward direction error ~ last outward trajectory straightness*. A significant effect of last outward trajectory tortuosity (here measured as straightness index) on homeward direction error could imply that individuals are using self-generated idiothetic directional cues rather than stable external directional cues when path integrating. I therefore analysed the power of this model and estimated an appropriate sample size to detect any effect.

3. *homeward direction error ~ last outward trajectory sinuosity*. Similarly to (2), a significant effect in this model could imply individuals are using idiothetic cues, as sinuosity is another measure of path tortuosity.

4. *homeward direction error ~ exploratory index*. A significant effect of exploratory index on homeward directional error would imply that fish could learn some aspects of the experimental pool before trials and use these learned cues to navigate. I therefore estimated an appropriate sample size to detect this effect.

Model 1: *homeward direction error ~ last outward trajectory length*

I fit a linear model between last outward trajectory length and homeward direction error using the package *lme4*, and determined the power of this model with the *powerSim()* function in the *simr* package:

```
lengthmodel1 <- lm(HomewardDirectionError ~ LOT_Length_cm, data=FishData)
# Fit linear model between homeward direction error and last outward traj
ectory length
summary(lengthmodel1)

##
## Call:
## lm(formula = HomewardDirectionError ~ LOT_Length_cm, data = FishData)
##
## Residuals:
##        1        2        3        4        5        6
## -22.107    5.729   13.446  -10.899    9.298    4.533
##
## Coefficients:
##               Estimate Std. Error t value Pr(>|t|)
## (Intercept)   2.252174   9.751774   0.231   0.8287
```

```
## LOT_Length_cm 0.010672   0.003158   3.380   0.0278 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 15.23 on 4 degrees of freedom
## Multiple R-squared:  0.7406, Adjusted R-squared:  0.6758
## F-statistic: 11.42 on 1 and 4 DF,  p-value: 0.0278

lengthmodel1power <- powerSim(lengthmodel1, nsim = 1000) # Calculate the
power of this linear model
lengthmodel1power

## Power for predictor 'LOT_Length_cm', (95% confidence interval):
##       71.40% (68.49, 74.18)
##
## Test: t-test
##
## Based on 1000 simulations, (0 warnings, 0 errors)
## alpha = 0.05, nrow = 6
##
## Time elapsed: 0 h 0 m 2 s
##
## nb: result might be an observed power calculation
```

Due to the power observed (71.40%) being less than the generally accepted level of 80%, I extended the model to 10 fish to estimate the power at this sample size:

```
lengthmodel2 <- extend(lengthmodel1, along="FishID", n=10) # Extend the l
inear model to include a sample size of 10 instead of 6
lengthmodel2power <- powerSim(lengthmodel2, nsim = 1000) # Calculate the
power of this extended model
lengthmodel2power

## Power for predictor 'LOT_Length_cm', (95% confidence interval):
##       98.70% (97.79, 99.31)
##
## Test: t-test
##
## Based on 1000 simulations, (0 warnings, 0 errors)
## alpha = 0.05, nrow = 10
##
## Time elapsed: 0 h 0 m 2 s
```

I then plotted a power curve to determine the minimum appropriate sample size for this model to have a power greater than 80%:

```
lengthcurve <- powerCurve(lengthmodel2, along="FishID", nsim = 1000) # Pl
ot a power curve to show the power for this range of sample sizes
plot(lengthcurve)
```

667 The resulting power curve suggests future experiments should use at least 8 fish to detect a
668 significant effect of last outward trajectory length on homeward direction error.

669 Model 2: *homeward direction error ~ last outward trajectory straightne*ss

670 I fit a linear model between last outward trajectory straightness and homeward direction error and
671 determined the power of this model:

```
672 straightnessmodel1 <- lm(HomewardDirectionError ~ LOT_Straightness, data=
673 FishData) # Fit linear model between homeward direction error and last ou
674 tward trajectory straightness
675 summary(straightnessmodel1)

676 ##
677 ## Call:
678 ## lm(formula = HomewardDirectionError ~ LOT_Straightness, data = FishDat
679 a)
680 ##
681 ## Residuals:
682 ##       1       2       3       4       5       6
683 ## -28.070  40.469   9.509 -23.873  -4.454   6.418
684 ##
685 ## Coefficients:
686 ##              Estimate Std. Error t value Pr(>|t|)
687 ## (Intercept)     36.30      16.37   2.217   0.0909 .
```

```
## LOT_Straightness   -112.44      151.90  -0.740   0.5002
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 28.05 on 4 degrees of freedom
## Multiple R-squared:  0.1205, Adjusted R-squared:  -0.0994
## F-statistic: 0.548 on 1 and 4 DF,  p-value: 0.5002

straightnessmodel1power <- powerSim(straightnessmodel1, nsim = 1000) # Ca
lculate the power of this linear model
straightnessmodel1power

## Power for predictor 'LOT_Straightness', (95% confidence interval):
##        9.50% ( 7.75, 11.49)
##
## Test: t-test
##
## Based on 1000 simulations, (0 warnings, 0 errors)
## alpha = 0.05, nrow = 6
##
## Time elapsed: 0 h 0 m 3 s
##
## nb: result might be an observed power calculation
```

Due to power observed (9.50%) being much less than the generally accepted level of 80%, I
extended the model to 100 fish to estimate the power at this sample size:

```
straightnessmodel2 <- extend(straightnessmodel1, along="FishID", n=100) #
Extend the linear model to include a sample size of 100 instead of 6
straightnessmodel2power <- powerSim(straightnessmodel2, nsim = 1000) # Ca
lculate the power of this extended model
straightnessmodel2power

## Power for predictor 'LOT_Straightness', (95% confidence interval):
##       89.10% (87.00, 90.96)
##
## Test: t-test
##
## Based on 1000 simulations, (0 warnings, 0 errors)
## alpha = 0.05, nrow = 100
##
## Time elapsed: 0 h 0 m 5 s
```

I then plotted a power curve to determine the minimum appropriate sample size for this model to
have a power greater than 80%:

```
straightnesscurve <- powerCurve(straightnessmodel2, along="FishID", nsim
= 1000) # Plot a power curve to show the power for this range of sample s
izes
plot(straightnesscurve)
```

731 The resulting power curve suggests future experiments should at least 90 fish to detect a significant

732 effect of last outward trajectory straightness on homeward direction error.

733 Model 3: *homeward direction error ~ last outward trajectory sinuosity*

734 I fit a linear model between last outward trajectory sinuosity and homeward direction error and

735 determined the power of this model:

```
736  sinuositymodel1 <- lm(HomewardDirectionError ~ LOT_Sinuosity, data=FishDa
737  ta) # Fit linear model between homeward direction error and last outward
738  trajectory sinuosity
739  summary(sinuositymodel1)

740  ##
741  ## Call:
742  ## lm(formula = HomewardDirectionError ~ LOT_Sinuosity, data = FishData)
743  ##
744  ## Residuals:
745  ##        1       2       3       4       5       6
746  ## -24.361  48.009  -6.911 -22.177  -4.996  10.436
747  ##
748  ## Coefficients:
749  ##              Estimate Std. Error t value Pr(>|t|)
750  ## (Intercept)    30.770     38.789   0.793    0.472
751  ## LOT_Sinuosity  -2.207     25.949  -0.085    0.936
752  ##
753  ## Residual standard error: 29.88 on 4 degrees of freedom
```

```
## Multiple R-squared:  0.001804,   Adjusted R-squared:  -0.2477
## F-statistic: 0.007231 on 1 and 4 DF,  p-value: 0.9363

sinuositymodel1power <- powerSim(sinuositymodel1, nsim = 1000) # Calculat
e the power of this linear model
sinuositymodel1power

## Power for predictor 'LOT_Sinuosity', (95% confidence interval):
##        4.20% ( 3.04,  5.64)
##
## Test: t-test
##
## Based on 1000 simulations, (0 warnings, 0 errors)
## alpha = 0.05, nrow = 6
##
## Time elapsed: 0 h 0 m 3 s
##
## nb: result might be an observed power calculation
```

Due to power observed (4.20%) being much less than the generally accepted level of 80%, I extended the model to 6,000 fish to estimate the power at this sample size:

```
sinuositymodel2 <- extend(sinuositymodel1, along="FishID", n=6000) # Exte
nd the linear model to include a sample size of 6000 instead of 6
sinuositymodel2power <- powerSim(sinuositymodel2, nsim = 1000) # Calculat
e the power of this extended model
sinuositymodel2power

## Power for predictor 'LOT_Sinuosity', (95% confidence interval):
##        78.30% (75.61, 80.82)
##
## Test: t-test
##
## Based on 1000 simulations, (0 warnings, 0 errors)
## alpha = 0.05, nrow = 6000
##
## Time elapsed: 0 h 0 m 5 s
```

As the power of the model only appears to approach 80% when using a massive sample size (at least 6,000 fish), future research aiming to determine whether fish use idiothetic directional cues or stable external directional cues to path integrate should use a different experimental design that tests use of idiothetic vs allothetic directional cues more directly.

Model 4: *homeward direction error ~ exploratory index*

I fit a linear model between exploratory index and homeward direction error and determined the power of this model:

```
exploratorymodel1 <- lm(HomewardDirectionError ~ ExploratoryIndex, data=F
ishData) # Fit linear model between homeward direction error and explorat
ory index
summary(exploratorymodel1)
```

```
## 
## Call:
## lm(formula = HomewardDirectionError ~ ExploratoryIndex, data = FishDat
## a)
## 
## Residuals:
##       1       2       3       4       5       6
## -2.933  39.429  -5.786 -35.034  -5.196   9.520
## 
## Coefficients:
##                  Estimate Std. Error t value Pr(>|t|)
## (Intercept)         80.92      58.33   1.387    0.238
## ExploratoryIndex   -73.51      79.02  -0.930    0.405
## 
## Residual standard error: 27.12 on 4 degrees of freedom
## Multiple R-squared:  0.1779, Adjusted R-squared:  -0.02767
## F-statistic: 0.8654 on 1 and 4 DF,  p-value: 0.4049

exploratorymodel1power <- powerSim(exploratorymodel1, nsim = 1000) # Calc
ulate the power of this linear model
exploratorymodel1power

## Power for predictor 'ExploratoryIndex', (95% confidence interval):
##       11.10% ( 9.22, 13.21)
## 
## Test: t-test
## 
## Based on 1000 simulations, (0 warnings, 0 errors)
## alpha = 0.05, nrow = 6
## 
## Time elapsed: 0 h 0 m 3 s
## 
## nb: result might be an observed power calculation
```

Due to power observed (11.10%) being much less than the generally accepted level of 80%, I extended the model to 60 fish to estimate the power at this sample size:

```
exploratorymodel2 <- extend(exploratorymodel1, along="FishID", n=60) # Ex
tend the linear model to include a sample size of 60 instead of 6
exploratorymodel2power <- powerSim(exploratorymodel2, nsim = 1000) # Calc
ulate the power of this extended model
exploratorymodel2power

## Power for predictor 'ExploratoryIndex', (95% confidence interval):
##       81.70% (79.16, 84.05)
## 
## Test: t-test
## 
## Based on 1000 simulations, (0 warnings, 0 errors)
## alpha = 0.05, nrow = 60
## 
## Time elapsed: 0 h 0 m 2 s
```

844   I then plotted a power curve to determine the minimum appropriate sample size for this model to

845   have a power greater than 80%:

```
846  exploratorycurve <- powerCurve(exploratorymodel2, along="FishID", nsim =
847  1000) # Plot a power curve to show the power for this range of sample siz
848  es
849  plot(exploratorycurve)
```

The resulting power curve suggests future experiments should include at least 60 fish to detect a

significant effect of exploratory behaviour on homeward direction error.

Conclusions

Although slightly underpowered, this experiment used a somewhat appropriate sample size to detect

whether homeward directional error accumulated with increasing outward path length. The

experiment was only slightly underpowered, with a power of 71% instead of the generally accepted

80%. Larger sample sizes or different experimental designs are needed to be able to more

conclusively determine whether homeward path behaviour is influenced by other outward path

behaviour or by exploratory behaviour in the acclimation period.