1   Host selection in European ectomycorrhizal Basidiomycetes affects their estimated

2   spore volumes.

4   Hironari Izumi

5   Microbial Ecology Unit, The Jurinji Buddhist temple, 481 Oshiocho, Ohranao,

6   Nishikyo-ku, Kyoto, 610-1133, Japan.

Abstract

Many ectomycorrhizal fungi show selective associations with certain host tree taxa.

Their significance has been demonstrated in the vegetative growth of the fungi, such as

structuring the fungal community of the mycorrhizal roots. On the other hand, our

understanding of the effects of host selection on reproductive traits such as pore

production is very limited. Using two independently published descriptions of host

associations and spore dimensions, the differences in spore volumes depending on the

associated host taxa were examined in species belonging to 10 ectomycorrhizal genera.

The species in *Suillus* were the most host-selective while those in *Inocybe* were

generalist. Some species in *Suillus*, *Lactarius*, *Russula,* and *Hebeloma* produced

significantly ($P<0.05$) larger spores when they associated with certain host taxa or

groups, including Pinaceae or broad-leaved hosts. However, such differences were not

found in *Leccinum* and *Tricholoma*. The ecological/evolutionary processes to facilitate

host selection through spore volumes were discussed.

Keywords: *Amanita*, *Cortinarius*, host-mycorrhizal interaction, host specificity,

Hygrophorus

Introduction

Ectomycorrhizas are a symbiotic lifestyle of the soil fungi with woody plant hosts,

including Pinaceae, Fagaceae, Betulaceae, and Salicaceae (Alexander, 2006). The recent

study reporting that more than half of the ectomycorrhizal fungi are associated with two

or fewer host plant genera (Voller et al., 2023) suggests many ectomycorrhizal fungi are

selective in their host plant taxa. The host selection of the ectomycorrhizal fungi plays a

pivotal role in structuring the fungi's belowground community (Ishida et al., 2007;

Matsuoka et al., 2020). Additionally, the host specificity has been demonstrated to

contribute to the dominance of the specialised fungi in the soil fungal community

(Tedersoo et al., 2024) by providing putative competitive advantages to the fungi

(Kennedy et al., 2020). Therefore, the importance of host selection of ectomycorrhizal

fungi on their belowground vegetative growth is well recognised whereas its influence

on aboveground reproductive traits, such as sporocarp and spore productions, is still

poorly understood, despite the fact that some of the reproductive traits of the

ectomycorrhizal fungi were influenced by the host tree. For instance, the host tree sizes

affect sporocarp production of associated ectomycorrhizal fungi (Nara et al., 2003) and

symbiotic ectomycorrhizal fungi yield larger sporocarps compared to solitary

saprotrophs through the possible supply of resources from the hosts (Bässler et al.,

2015). Therefore, it is likely that some other reproductive traits of the ectomycorrhizal

fungi, such as spore production, are affected by the associated host taxa.

In this study, I tested the possibility that the spore volumes show any differences

depending on the host taxa with which they are associated. Ten ectomycorrhizal genera,

including *Amanita*, *Cortinarius*, *Hebeloma*, *Hygrophorus*, *Inocybe*, *Lactarius*, *Leccinum*,

Russula, *Suillus*, *Tricholoma*, were selected for the present study because they are

proven to be ectomycorrhizal (Comandini et al., 2012), species-rich, show some host

specificity and produce large sporocarps with measurable spores. Additionally,

estimated sporocarp biomass of the species was compared between different host taxa to

assess the host effects on whole reproductive organ.

Materials and Methods

Descriptions of the associating host taxa and two-dimensional (length and width) spore

size measurements by species were extracted from the books by Læssøe and Petersen

(2019) and by Kibby (2020–2023). More than 500 (Kibby, 2020–2023) and 300

(Læssøe and Petersen, 2019) species descriptions were used for the estimation of the

spore volumes (Supplemental table 1).

To obtain a sufficient number of species for statistical analysis to evaluate the host

effects, the host tree taxa were classified by the family level for *Cortinarius*,

Hygrophorus, *Lactarius*, *Leccinum*, *Russula* and *Tricholoma*. For less selective fungi,

such as *Amanita*, *Hebeloma*, and *Inocybe*, the hosts were grouped into broad-leaved

only or both broad-leaved and coniferous species. For highly specific genus of *Suillus*,

the hosts were divided into genus level, such as *Larix* and *Pinus*. The comparison was

made between two major host tree taxa or groups within each ectomycorrhizal genus.

To estimate a degree of host specificity in the genus of interest, the proportions of the

species associated with a single genus of the host tree were calculated.

The midpoints of length and width measurements were employed to calculate the spore

volume for each species (Bässler et al., 2015), using the formula by Calhim et al. (2018).

For the volume estimation, the shape of the spores was assumed to be an ellipsoid

(Bässler et al., 2015), except for some species of *Amanita* whose shapes were spherical.

In that case, the volume was calculated by the formula of $V=4/3*3.14*r^3$.

For the comparisons of sporocarp biomass, the measurements were extracted from the

descriptions by Kibby only because Læssøe and Petersen did not provide such

information. The proxy values of sporocarp biomass were obtained by the squared mean

diameter of the cap (Bässler et al., 2015),

Brunner-Menzel test was performed to make statistical comparisons using Jamovi

software package (The Jamovi project, 2022).

Results

Among the fungal genera examined, *Suillus* was the most specific genus and all species

were associated with a single host tree genus. On the other hand, the least specific i.e.,

generalist, was *Inocybe* with less than 20% of a single genus associated species (Fig. 1).

The spore volumes of the species belong to *Suillus* were significantly (P<0.05) different,

depending on the host genera in both descriptions by Kibby and by Læssøe &Petersen.

However, those of *Inocybe* were statistically indistinguishable between the hosts by

either description. *Suillus*, *Russula*, *Lactarius* and *Hebeloma* were the genera that

statistically significant differences of spore volumes in specific host taxa were

supported by both descriptions. A general trend that the high proportions of the species

selecting specific host taxa accompanied with the robustness (supported by two

descriptions) of host dependent spore volume difference was observed.

In *Russula* and *Lactarius*, the spore volumes of the species associated with Pinaceae

hosts were significantly (P<0.05) larger (1.3 folds) than those with Fagaceae or

Betulaceae hosts in both descriptions (Table 1). Additionally, the species of *Hebeloma*

associated with broad-leaved hosts and those of *Suillus* with *Larix* produced

significantly larger (1.3 folds) spores than those with other hosts, such as both of

broad-leaved and coniferous or *Pinus* respectively (Table 1). Moreover, some species in

Amanita, Cortinarius and *Hygrophorus* produced larger spores (1.5 folds, 1.4 folds and

1.3 folds respectively) when they associated with Pinaceae, Fagaceae or broadleaved

hosts respectively although statistical significance was observed in only one of two

descriptions (Table 1).

To examine the possibility that observed larger spores in species of *Hebeloma*,

Lactarius, *Russula* and *Suillus* in association with the certain hosts is due to the

production of larger sporocarp, the sporocarp biomass was estimated for the same

species. There was no statistically significant difference of sporocarp biomass between

different host three taxa (Table 2), suggesting the host effect on spore volume is

specific.

Discussion

Because the fungi forming ectomycorrhizas with plants are diverse and consist of, at

least, 5400 species (Sharma, 2017), the number of species examined in this study is

rather small (500 species at most). Although such a small number would provide a

partial picture of the host effects on ectomycorrhizas interesting patterns emerge. Firstly,

degrees of host selection vary depending on the fungal taxa. Some species belonging to

the genera *Hebeloma*, *Lactarius*, *Russula*, and *Suillus* tend to be more host-selective

than those in Inocybe (Fig, 1). Secondary, they produce larger spores in association with

certain host taxa (Table 1). The host selectivity observed in this study can arise from

two distinct pathways. One is phynotypic flexibility of the fungi to the current

environmental conditions. The other is fungal genotypic evolution to increase fitness by

associating with particular host taxa. It is well known that overall sporocarp production

is influenced by rainfall and ambient temperature (Boddy et al., 2014). Thus, there is a

possibility that observed larger spores in association with certain hosts may reflect

phynotipic adaptation to facilitate survival under the conditions which the host grows.

However, phynotypic adaptation may not play critical role on the spore volume

difference because the result in this study showing no statistically significant difference

of sporocarp biomass between different suggests that the sporocarp sizes of the species

are consistent within the genus and it is likely that spore volumes are also genetically

fixed in the same genus. These lines of reasoning lead us to conclude that evolutionally

adaptation is more plausible for developing host specificity with larger spore volumes.

Then, what is the advantages to evolve producing larger spores? As the spores are

reproductive organ the advantages must be placed on how much the larger spore

volumes contribute fungal dispersal and survival. Indeed, some tropical *Amanita*

produce smaller spores for effective areal dispersal between sparsely distributed hosts

(Tulloss, 2005). The larger spores, on the other hand, would have merits for survival

rather than dispersal. In this study, *Larix* spp., the host of highly specific *Suillus*, grows

under dry, cold winter conditions in native high Alpine rages and usually performs

better in poor to medium nutrient sites as an early coloniser in the lower elevation (Da

Ronch et al., 2016). Additionally, it has been shown that the larger spores were thought

to retain more water and provide some protection against desiccation (Kauserud et al.,

2011) and spores become larger in resource-limited habitats (Halbwachs et al., 2017).

Therefore, larger spores in *Larix*-specific *Suillus* species may be the evolutionary

consequences selected by better survival of the larger spores in harsh environments.

Russula, *Lactarius* and *Hebeloma* have been known to be "late-stage" fungi (Smith &

Read, 2008; Ishida et al., 2008). The sporocarps of *Lactarius*, for instance, appear at

older stand ages in *Pinus sylvestris* forests (Bonet et al., 2004) and those of *Russula* and

Hebeloma are observed in a late stage of primary vegetative succession on volcanic Mt

Fuji (Nara et al., 2003). Whereas "late-stage" fruiting may be due to slow developments

of belowground fungal biomass, the late fruiting may be caused by delayed spore

germination, waiting for the time when the host tree reaches a certain age. The previous

report showing that the spores of these genera are difficult to produce mycorrhiza by

inoculations of the spores to seedlings (Nara, 2009) suggests spore germination may

require more mature host trees. Thus, larger spores can contribute to remaining viable

for a long time by waiting for the host with suitable ages to be around. This also

explains why *Inocybe* does not show host-specific differences in spore volumes.

Inocybe has been suggested one of early colonisers in primary vegetative succession

(Nara et al., 2003), which they can immediately colonise the host and do not wait for

colonisation for long.

How the spores become larger when the fungi are associated with certain host taxa is

completely unexplored but one possibility is that specific association may increase

physiological affinity between the fungi and the host, which facilitates to produce the

larger spores. The previous study has demonstrated that host tree allocated more carbon

to the specific *Suillus grevillei* colonised roots compared to non-specific *S. bovinus*

(Finlay,1989). Therefore, it is likely that host-specific fungi obtain more

photo-assimilated carbons from specific hosts and can produce larger spores.

The interesting genera are *Leccinum* and *Tricholoma*, which have high (>50%)

host-selective species but no statistically significant host effects on the spore volume.

Leccinum has been considered to shift their hosts recently (den Bakker et al., 2004),

which may not provide sufficient time to develop the host effects on spore volumes.

Tricholoma spp. have been shown to form mycorrhizas between oak and pine in vitro

(Yamanaka et al., 2014). Thus, the observed selectivity in the field descriptions may

reflect the environmental conditions, such as the availability of the host trees in the

locations the fungi grow. These two instances, of *Leccinum* and *Tricholoma,* suggests

that stable and exclusive association between the host and the fungi are required to

manifest the host effects on the spore volume.

The spore volumes of *Amanita*, *Cortinarius,* and *Hygrophorus* increased in association

with the hosts of broad-leaved or Pinaceae respectively (Table 1) alth ough only one of

the two descriptions, namely by Kiddy, was statistically supported. As Kiddy describes

mainly British species, this may reflect more specialisation to the host taxa of the

species in these genera in Britain, by geographical isolation from mainland Europe. For

example, the geographical isolation by growing on an island, such as Corsica, can

change the relationship between ectomycorrhizal fungi and their host tree by shifting the

host ranges of the fungi (Rochet et al., 2011). Thus, it is possible that the separation of

British species from those in mainland Europe may also influence the host effects on

spore volumes although this requires more detailed examinations.

Disclosure

The author declares no conflicts of interest.

Acknowledgment

This work was financially supported by the Jurinji Buddhist temple. Dr Melanie Roy of

PCI Ecology was acknowledged for her feedback and encouragement for early version

of the manuscript.

References

Alexander, I. J. (2006). Ectomycorrhizas-out of Africa? *New Phytologist*, 172,

589-591.   https://doi.org/10.1111/j.1469-8137.2006.01930.x

Bässler, C., Heilmann-Clausen,J., Karasch, K., Brandl, R., & Halbwachs, H. (2015)

Ectomycorrhizal fungi have larger fruit bodies than saprotrophic fungi. *Fungal

Ecology*, 17, 205-212. https://doi.org/10.1016/j.funeco.2014.06.005.

Boddy, L., Büntgen, U., Egli, S., Gange, A. C., Heegaard, E., Kirk, P. M., ... &

Kauserud, H. (2014). Climate variation effects on fungal fruiting. *Fungal*

Ecology, *10*, 20-33. https://doi.org/10.1016/j.funeco.2013.10.006

Bonet, J.A., Fischer, C.R. & Colinas, C. (2004). The relationship between forest age and

aspect on the production of sporocarps of ectomycorrhizal fungi in *Pinus*

sylvestris forests of the Central Pyrenees. *Forest Ecology and Management,* 203,

157-175. Doi:10.1016/j.foreco.2004.07.063

Calhim, S., Halme, P., Petersen, J.H., Læssøe, T., Bässler, C., & Heilmann-Clausen, J

(2018). Fungal spore diversity reflects substrate-specific deposition

challenges. *Scientifc Reports* 8, 5356 (2018).

https://doi.org/10.1038/s41598-018-23292-8

Comandini, O., Rinaldi, A.C., & Kuyper, D.W. (2012). Measuring and Estimating

Ectomycorrhizal Diversity: A Continuous Challenge. In Mycorrhiza: Occurrence in

Natural and Restored Environments. Pp165-200. Pagano, M ed. Nova Science

Publishers Inc. New York, USA.

Da Ronch, F., Caudullo, G., & de Rigo, T. D. (2016). *Larix decidua* and other larches in

Europe. In san-Mugel-Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T.,

Mauri, A. (eds.). *European Atlas of Forest Tree Species*. Pub. Off. Eu,

Luxembourg.

den Bakker, H.C., Zuccarello, G.C., Kuyper, T.W., & Noordeloos, M.E. (2004).

Evolution and host specificity in the ectomycorrhizal genus *Leccinum*. *New*

Phytologist, 163, 201-215. https://doi.org/10.1111/j.1469-8137.2004.01090.x

Finlay, R. (1989). Functional aspects of phosphorus uptake and carbon translocation in

incompatible ectomycorrhizal associations between *Pinus sylvestris* and *Suillus*

grevillei and *Boletinus cavipes*. *New Phytologist*, 112, 185– 192.

Halbwachs, H., Heilmann-Clausen, J., & Bässler, C. (2017). Mean spore size and shape

in ectomycorrhizal and saprotrophic assemblages show strong responses under

resource constraints. *Fungal Ecology*, 26, 59-64.

https://doi.org/10.1016/j.funeco.2016.12.001.

Ishida, T., Nara, K., & Hogetsu, T. (2007). Host effects on ectomycorrhizal fungal

communities: insight from eight host species in mixed conifer–broadleaf forests.

New Phytologist, 174, 430-440. https://doi.org/10.1111/j.1469-8137.2007.02016.x

Ishida TA, Nara K, Tanaka M, Kinoshita A, Hogetsu T. (2008). Germination and

infectivity of ectomycorrhizal fungal spores in relation to their ecological traits

during primary succession. *New Phytol*.180, 491-500. doi:

10.1111/j.1469-8137.2008.02572.x.

The Jamovi project (2022). *Jamovi*. (Version 2.3) [Computer Software]. Retrieved from
https://www.jamovi.org.

Kauserud, H., Heegaard, E., Halvorsen, R., Boddy, L., Høiland, K., & Stenseth, N.C.

(2011). Mushroom's spore size and time of fruiting are strongly related: is

moisture important? *Biology Letters*, 7, 273-6. doi: 10.1098/rsbl.2010.0820.

Kennedy, P.G., Gagne, J., Perez-Pazos, E., Lofgren, L.A., & Nguyen, N.H. (2020). Does

fungal competitive ability explain host specificity or rarity in ectomycorrhizal

symbioses? *PLoS ONE*,15, e0234099.

https://doi.org/10.1371/journal.pone.0234099

Kibby, G. (2020–2023). *Mushrooms and toaddtools of Britain & Europe*. 3rd Ed. Self

publication.

Læssøe, T., & Petersen, J. H. (2019). *Fungi of Temperate Europe*. Princeton University

Press.

Matsuoka S, Sugiyama Y, Tateno R, Imamura S, Kawaguchi E, Osono T. (2020).

Evaluation of host effects on ectomycorrhizal fungal community compositions in

a forested landscape in northern Japan. *Royal Society Open Science*, 19, 191952.

doi: 10.1098/rsos.191952.

Nara, K., Nakaya, H., Hogetsu, T. (2003), Ectomycorrhizal sporocarp succession and

production during early primary succession on Mount Fuji. *New Phytologist*, 158,

193-206. https://doi.org/10.1046/j.1469-8137.2003.00724.x

Nara K. (2009) Spores of ectomycorrhizal fungi: ecological strategies for germination

and dormancy. New Phytol, 181, 245-248. doi:

10.1111/j.1469-8137.2008.02691.x.

Rochet, J., Moreau, PA., Manzi, S., & Gardes, M. (2011). Comparative phylogenies and

host specialization in the alder ectomycorrhizal

fungi *Alnicola*, *Alpova* and *Lactarius* (Basidiomycota) in Europe. *BMC*

Evolutionary Bioogy,l 11, 40. https://doi.org/10.1186/1471-2148-11-40

Sharma, R. (2017). Ectomycorrhizal Mushrooms: Their Diversity, Ecology and

Practical Applications. In: Varma, A., Prasad, R., Tuteja, N. (eds) Mycorrhiza -

Function, Diversity, State of the Art. Springer, Cham.

https://doi.org/10.1007/978-3-319-53064-2_7

Smith, S.E. & Read, D (2008). *Mycorrhizal Symbiosis*. 3[rd] Ed. Academic Press.

https://doi.org/10.1016/B978-0-12-370526-6.X5001-6

Tedersoo, L., Drenkhan, R., Abarenkov, K., Anslan, S., Bahram, M., Bitenieks, K.,

Buegger, F., Gohar, D., Hagh-Doust, N., Klavina, D., Makovskis, K., Zusevica, A.,

Pritsch, K., Padari, A., Pölme, S., Rahimlou, S., Rungid, D., & Mikryukov, V.

(2024). The influence of tree genus, phylogeny, and richness on the specificity,

rarity, and diversity of ectomycorrhizal fungi. *Environmental Microbiology*

Reports, 16(2), e13253. Available from: https://doi.org/10.1111/1758-2229.13253

Tulloss, R. E. (2005). Amanita-distribution in the Americas, with comparison to eastern

and southern Asia and notes on spore character variation with latitude and

ecology. *Mycotaxon*, *93*(189), e231.

Voller, F., Ardanuy, A., Taylor, A. F. S., & Johnson, D. (2023). Maintenance of host

specialisation gradients in ectomycorrhizal symbionts. *New Phytologist*, 242,

1426-1435. https://doi.org/10.1111/nph.19395

Yamanaka, T., Ota ,Y., Konno, M., Kawai, M., Ohta, A., Neda, H., Terashima, Y., &

Yamada, A. (2014). The host ranges of conifer-associated *Tricholoma matsutake*,

Fagaceae-associated *T. bakamatsutake* and *T. fulvocastaneum* are wider in vitro

than in nature. *Mycologia*, 106, 97-406. doi: 10.3852/13-197.

Table 1 Comparisons of the estimated mean spore volumes (μm^3) of 10 ectomycorrhizal fungal genera occurred between two different host taxa or groups

Fungal genus	Host tree		P-value*	Reference
Amanita	Broad-leaved	Broad-leaved & coniferous		
	568 (21)	358 (14)	0.004	K
	632 (18)	417 (9)	0.101	L&P
Cortinarius	Pinaceae	Fagaceae		
	146 (87)	212 (66)	0.001	K
	168 (30)	188 (33)	0.169	L&P
Hebeloma	Broad-leaved	Broad-leaved & coniferous		
	261 (24)	196 (12)	0.036	K
	262 (8)	200 (4)	0.048	L&P
Hygrophorus	Pinaceae	Fagaceae		
	152 (13)	110 (11)	0.042	K
	138 (11)	112 (10)	0.338	L&P
Inocybe	Broad-leaved	Broad-leaved & coniferous		
	204 (67)	179 (25)	0.389	K
	203 (28)	173 (20)	0.112	L&P
Lactarius	Pinaceae	Betulaceae		
	204 (13)	147 (15)	0.002	K
	231 (14)	129 (11)	0.001	L&P
Leccinum	Betulaceae	Other genera		
	235 (5)	155 (4)	0.208	K
	232 (6)	174 (5)	0.24	L&P
Russula	Pinaceae	Fagaceae		

	249 (18)	187 (55)	0.001	K
	266 (14)	199 (39)	0.001	L&P
Suillus	*Larix*	*Pinus*		
	93.8 (6)	59.4 (8)	0.041	K
	103 (6)	65 (9)	0.021	L&P
Tricholoma	Pinaceae	Fagaceae		
	84 (20)	86.8 (13)	0.306	K
	76.2 (15)	85.1 (7)	0.381	L&P

* P-values were calculated by Brunner-Munzer test.

The numbers of species examined were indicated in round brackets.

In reference, K and L&P indicate Kibby (2020–2023) and Læssøe & Petersen (2019)

Table 2 Differences of estimated sporocarp sizes occurring between different host tree

Fungal genus	Compared host tree combination	P-value
Hebeloma	Broad-leaved/Broad-leaved & Coniferous	0.209
Lactarius	Pinaceae/Betulaceae	0.944
Russula	Pinaceae/Fagaceae	0.651
Suillus	Larix/Pine	0.548

Fig. 1 Proportions of the fungal species occurring with a single genus of the host tree. Two independent descriptions of the species by Kibby (2020–2023) and Læssøe and Petersen (2019) are shown in closed and open bars respectively. The star on the bar indicates the fungal descriptions that statistically supported (P<0.05) spore volume differences are observed between species occurring in different host taxa.