1   Host selection in European ectomycorrhizal Basidiomycetes affects their estimated

2   spore volumes.

4   Hironari Izumi

5   Microbial Ecology Unit, The Jurinji Buddhist temple, 481 Oshiocho, Ohranao,

6   Nishikyo-ku, Kyoto, 610-1133, Japan.

Abstract

Many ectomycorrhizal fungi show selective associations with certain host tree taxa.

Their significance has been demonstrated in the vegetative growth of the fungi, such as

structuring the fungal community of the mycorrhizal roots. On the other hand, our

understanding of the effects of host selection on reproductive traits such as pore

production is very limited. Using two independently published descriptions of host

associations and spore dimensions, the differences in spore volumes depending on the

associated host taxa were examined in 10 ectomycorrhizal genera. *Suillus* was the most

specific genus while *Inocybe* was a generalist. *Suillus*, *Lacterias*, *Russula,* and

Hebeloma produced significantly (P<0.05) larger spores when they associated with

certain host taxa or groups, including Pinaceae or broad-leaved hosts. However, such

differences were not found in *Leccinum* and *Tricoloma*. The possibility that the host

selection of ectomycorrhizal fungi provides an ecological advantage by producing larger

volumes was discussed.

Keywords: *Amanita*, *Cortinarius*, host-mycorrhizal interaction, host specificity,

Hygroporus

Ectomycorrhizas are a symbiotic lifestyle of the soil fungi with woody plant hosts,

including Pinaceae, Fagaceae, Betulaceae, and Salicaceae (Alexander, 2006). The recent

analysis reporting that more than half of the ectomycorrhizal fungi are associated with

two or fewer host plant genera (Voller et al., 2023) suggests many ectomycorrhizal

fungi are selective in their host plant taxa. The host selection of the ectomycorrhizal

fungi plays a pivotal role in structuring the fungi's belowground community (Ishida et

al., 2007; Matsuoka et al., 2020). Additionally, the host specificity has been

demonstrated to contribute to the dominance of the specialised fungi in the soil fungal

community (Tedersoo et al., 2024) by providing putative competitive advantages to the

fungi (Kennedy et al., 2020). Moreover, the host specificity facilitates nutrient exchange

between the host and the fungi (Finlay, 1989). Therefore, the importance of host

selection of ectomycorrhizal fungi on their vegetative growth in the soil is well

recognised whereas its influence on aboveground reproductive traits, such as spore

productions, is still poorly understood, despite the previous research demonstrating that

some of the reproductive traits of the ectomycorrhizal fungi were influenced by the host

tree. For instance, the host tree sizes affect sporocarp production of associated

ectomycorrhizal fungi (Nara et al., 2003) and symbiotic ectomycorrhizal fungi yield

larger sporocarps compared to solitary saprotrophs through the possible supply of

resources from the hosts (Bässler et al., 2015). Therefore, it is likely that the other

reproductive traits of the ectomycorrhizal fungi, such as spore production, are affected

by the specific host taxa.

In this study, I tested the possibility that the spore volumes show any differences

depending on the host taxa with which they are associated. Ten ectomycorrhizal genera,

including *Amanita*, *Cortinarius*, *Hebeloma*, *Hygroporus*, *Inocybe*, *Lactarius*, *Leccinum*,

Russula, *Suillus*, *Tricoloma*, were selected for the present study because they are proven

to be ectomycorrhizal (Comandini et al., 2012), species-rich, show some host specificity

and produce large sporocarps with measurable spores.

Descriptions of the associating host taxa and two-dimensional (length and width) spore

size measurements were extracted from the books by Læssøe and Petersen (2019) and

by Kibby (2020–2023).

To obtain a sufficient number of species for statistical analysis, the host tree taxa were

classified by the family level for *Cortinarius*, *Hygrophorus*, *Lactarius*, *Leccinum*,

Russula and *Tricoloma*. For less selective fungi, such as *Amanita*, *Hebeloma*, and

Inocybe, the hosts were grouped into broad-leaved only or both broad-leaved and

coniferous species. For highly specific *Suillus*, the hosts were divided into genus level,

such as *Larix* and *Pinus*. The comparison was made between two major host tree taxa or

groups for each ectomycorrhizal genus. To estimate a degree of host specificity in the

genus of interest, the proportions of the species associated with a single genus of the

host tree were calculated.

The midpoints of length and width measurements were employed to calculate the spore

volume for each species (Bässler et al., 2015), using the formula by Calhim et al. (2018).

For the volume estimations, the shape of the spores was assumed to be an ellipsoid

(Bässler et al., 2015), except for some species of *Amanita* whose shapes were spherical.

In that case, the volume was calculated by the formula of $V=4/3*3.14*r^3$.

Brunner-Menzel test was performed to make statistical comparisons using Jamovi

software package (The Jamovi project, 2022).

Among the fungal genera examined, *Suillus* was the most specific genus and all species

were associated with a single host tree genus. On the other hand, the least specific i.e.,

generalist, was *Inocybe* with less than 20% of a single genus associated species (Fig. 1).

The spore volumes in *Suillus* were significantly ($P<0.05$) influenced by the different

host genera in both descriptions by Kibby and by Læssøe &Petersen. However, those in

Inocybe were statistically indistinguishable between the hosts by either description.

Suillus, *Russula*, *Lacterius* and *Hebeloma* were the genera that statistically supported

host effects on spore volumes were detected in both descriptions. A general trend that

increased host specificity positively associated with the robustness (supported by two

descriptions) of host influence on the spore volume was observed. The interesting

genera were *Leccinum* and *Tricoloma*, which showed high (>50%) host selection but no

statistically significant host effects. *Leccinum* has been considered to shift their hosts

recently (den Bakker et al., 2004), which may not provide sufficient time to develop the

host effects on spore volumes. *Tricoloma* spp. have been shown to form mycorrhizas

between oak and pine in vitro (Yamanaka et al., 2014). Thus, the observed selectivity in

the field descriptions may reflect the environmental conditions such as the availability

of the host trees in the locations they grow. These two instances of *Leccinum* and

Tricoloma are revealing; suggesting that stable host-fungal associations and exclusive

affinity between the host and the fungi are required to manifest the host effects on the

spore volume.

In *Russula* and *Lactarius*, the spore volumes were significantly ($P<0.05$) larger with

Pinaceae hosts than those with Fagaceae or Betulaceae hosts in both descriptions (Table

1). Additionally, *Hebeloma* associated with broad-leaved hosts and *Suillus* with *Larix*

produced significantly larger spores than those with other hosts such as those found

with both broad-leaved and coniferous hosts or *Pinus* respectively (Table 1). Moreover,

Cortinarius, *Hygroporus*, and *Amanita* produced larger spores when they associated

with Pinaceae or broadleaved hosts although statistical significance was observed in

only one of two descriptions (Table 1). The previous studies have shown that the larger

spores were thought to retain more water and provide some protection against

desiccation (Kauserud et al., 2011) and spores become larger in resource-limited

conditions in the field (Halbwachs et al., 2017). These lines of evidence suggest the

possibility that having larger spores increases the chance of spore survival under harsh

and resource-constrained environments by preventing desiccation and storing larger

amounts of nutrients to be viable for a long period until encountering the host roots. In

line with this, producing larger spores by associating with particular host trees may

provide competitive advantages to the fungi for efficient ectomycorrhizal formation

when the conditions are unfavourable. Indeed, *Larix* spp., the host for highly specific

Suillus, grows under dry, cold winter conditions in native high Alpine rages and usually

performs better in poor to medium nutrient sites as an early coloniser in the lower

elevation (Da Ronch et al., 2016). Thus, producing larger spores in *Suillus* associated

with *Larix* spp. may reflect fungal adaptation for increasing the probability of survival

under such limited conditions.

Russula and *Lacterius* were known to be "late-stage" fungi (Smith & Read, 2008), and

their sporocarps appear at older stand ages in *Pinus sylvestris* forests (Bonet et al.,

2004). Whereas "late-stage" fruiting may be due to slow developments of belowground

fungal biomass, the late fruiting may be caused by delayed spore germination, waiting

for the time at which the host tree reaches a certain age. If this is the case, larger spores

can contribute to remaining viable for a longer period by waiting for the host with

suitable ages to be around.

The ecological significance of larger spores in *Hebeloma* is not yet clear while the

genera showed robust statistical support for host effects on spore volumes (Table 1).

The previous study showing that a specific association between *Suillus* and *Larix*

facilitated nutrient exchange (Finlay, 1989) suggests the possibility that *Hebeloma* has a

higher affinity to broad-laved hosts and produces larger spores by obtaining more

nutrients from the host tree.

The spore volumes of *Amanita*, *Cortinarius,* and *Hygrophorus* increased in association

with the hosts of broad-leaved or Pinaceae respectively (Table 1) although only one of

the two descriptions, namely by Kiddy, was statistically supported. As Kiddy describes

mainly British species, this may reflect more specialisations of these genera in Britain to

the particular host taxa by geographical isolation from mainland Europe. The

geographical isolation by growing on an island, such as Corsica, can change the

relationship between ectomycorrhizal fungi and their host tree by shifting the host

ranges of the fungi (Rochet et al., 2011). Such an instance suggests the possibility that

the separation of Britain from mainland Europe may also influence the host effects on

spore volumes although this requires more detailed examinations.

The results in this study showed that spore volumes of some ectomycorrhizal fungi were

affected by taxonomical differences of the host tree. It is interesting for future research

to ask the questions about whether other reproductive traits, such as time of fruiting or

fruitbody sizes may be affected by the different host taxa.

Disclosure

The author declares no conflicts of interest.

Acknowledgment

This work was financially supported by the Jurinji Buddhist temple.

References

Alexander, I. J. (2006). Ectomycorrhizas-out of Africa? *New Phytologist*, 172,

589-591. https://doi.org/10.1111/j.1469-8137.2006.01930.x

Bässler, C., Heilmann-Clausen,J., Karasch, K., Brandl, R., & Halbwachs, H. (2015)

Ectomycorrhizal fungi have larger fruit bodies than saprotrophic fungi. *Fungal*

Ecology, 17, 205-212. https://doi.org/10.1016/j.funeco.2014.06.005.

Bonet, J.A., Fischer, C.R. & Colinas, C. (2004). The relationship between forest age and

aspect on the production of sporocarps of ectomycorrhizal fungi in *Pinus*

sylvestris forests of the Central Pyrenees. *Forest Ecology and Management,* 203,

157-175. Doi:10.1016/j.foreco.2004.07.063

Calhim, S., Halme, P., Petersen, J.H., Læssøe, T., Bässler, C., & Heilmann-Clausen, J

(2018). Fungal spore diversity reflects substrate-specific deposition

challenges. *Scientifc Reports* 8, 5356 (2018).

https://doi.org/10.1038/s41598-018-23292-8

Comandini, O., Rinaldi, A.C., & Kuyper, D.W. (2012). Measuring and Estimating

Ectomycorrhizal Diversity: A Continuous Challenge. In Mycorrhiza: Occurrence in

Natural and Restored Environments. Pp165-200. Pagano, M ed. Nova Science

Publishers Inc. New York, USA.

Da Ronch, F., Caudullo, G., & de Rigo, T. D. (2016). *Larix decidua* and other larches in

Europe. In san-Mugel-Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T.,

Mauri, A. (eds.). *European Atlas of Forest Tree Species*. Pub. Off. Eu,

Luxembourg.

den Bakker, H.C., Zuccarello, G.C., Kuyper, T.W., & Noordeloos, M.E. (2004).

Evolution and host specificity in the ectomycorrhizal genus *Leccinum*. *New*

Phytologist, 163, 201-215. https://doi.org/10.1111/j.1469-8137.2004.01090.x

Finlay, R. (1989). Functional aspects of phosphorus uptake and carbon translocation in

incompatible ectomycorrhizal associations between *Pinus sylvestris* and *Suillus*

grevillei and *Boletinus cavipes*. *New Phytologist*, 112, 185– 192.

Halbwachs, H., Heilmann-Clausen, J., & Bässler, C. (2017). Mean spore size and shape

in ectomycorrhizal and saprotrophic assemblages show strong responses under

resource constraints. *Fungal Ecology*, 26, 59-64.

https://doi.org/10.1016/j.funeco.2016.12.001.

Ishida, T., Nara, K., & Hogetsu, T. (2007). Host effects on ectomycorrhizal fungal

communities: insight from eight host species in mixed conifer–broadleaf forests.

New Phytologist, 174, 430-440. https://doi.org/10.1111/j.1469-8137.2007.02016.x

The Jamovi project (2022). *Jamovi*. (Version 2.3) [Computer Software]. Retrieved from
https://www.jamovi.org.

Kauserud, H., Heegaard, E., Halvorsen, R., Boddy, L., Høiland, K., & Stenseth, N.C.

(2011). Mushroom's spore size and time of fruiting are strongly related: is

moisture important? *Biology Letters*, 7, 273-6. doi: 10.1098/rsbl.2010.0820.

Kennedy, P.G., Gagne, J., Perez-Pazos, E., Lofgren, L.A., & Nguyen, N.H. (2020). Does

fungal competitive ability explain host specificity or rarity in ectomycorrhizal

symbioses? *PLoS ONE*,15, e0234099.

https://doi.org/10.1371/journal.pone.0234099

Kibby, G. (2020–2023). *Mushrooms and toaddtools of Britain & Europe*. 3[rd] Ed. Self

publication.

Læssøe, T., & Petersen, J. H. (2019). *Fungi of Temperate Europe*. Princeton University

Press.

Matsuoka S, Sugiyama Y, Tateno R, Imamura S, Kawaguchi E, Osono T. (2020).

Evaluation of host effects on ectomycorrhizal fungal community compositions in

a forested landscape in northern Japan. *Royal Society Open Science*, 19, 191952.

doi: 10.1098/rsos.191952.

Nara, K., Nakaya, H., & Hogetsu, T. (2003), Ectomycorrhizal sporocarp succession and

production during early primary succession on Mount Fuji. *New Phytologist*, 158,

193-206. https://doi.org/10.1046/j.1469-8137.2003.00724.x

Rochet, J., Moreau, PA., Manzi, S., & Gardes, M. (2011). Comparative phylogenies and

host specialization in the alder ectomycorrhizal

fungi *Alnicola*, *Alpova* and *Lactarius* (Basidiomycota) in Europe. *BMC*

Evolutionary Bioogy,l 11, 40. https://doi.org/10.1186/1471-2148-11-40

Smith, S.E. & Read, D (2008). *Mycorrhizal Symbiosis*. 3[rd] Ed. Academic Press.

https://doi.org/10.1016/B978-0-12-370526-6.X5001-6

Tedersoo, L., Drenkhan, R., Abarenkov, K., Anslan, S., Bahram, M., Bitenieks, K.,

Buegger, F., Gohar, D., Hagh-Doust, N., Klavina, D., Makovskis, K., Zusevica, A.,

Pritsch, K., Padari, A., Pölme, S., Rahimlou, S., Rungid, D., & Mikryukov, V.

(2024). The influence of tree genus, phylogeny, and richness on the specificity,

rarity, and diversity of ectomycorrhizal fungi. *Environmental Microbiology*

Reports, 16(2), e13253. Available from: https://doi.org/10.1111/1758-2229.13253

Voller, F., Ardanuy, A., Taylor, A. F. S., & Johnson, D. (2023). Maintenance of host

specialisation gradients in ectomycorrhizal symbionts. *New Phytologist*, 242,

1426-1435. https://doi.org/10.1111/nph.19395

Yamanaka, T., Ota ,Y., Konno, M., Kawai, M., Ohta, A., Neda, H., Terashima, Y., &

Yamada, A. (2014). The host ranges of conifer-associated *Tricholoma matsutake*,

Fagaceae-associated *T. bakamatsutake* and *T. fulvocastaneum* are wider in vitro

than in nature. *Mycologia*, 106, 97-406. doi: 10.3852/13-197.

Table 1 Comparisons of the estimated mean spore volumes (μm^3) of 10 ectomycorrzal fungal genera occurred between two different host taxa or groups

Fungal genus	Host tree		P-value*	Reference
Amanita	Broad-leaved	Broad-leaved & coniferous		
	568 (21)	358 (14)	0.004	K
	632 (18)	417 (9)	0.101	L&P
Cortinarius	Pinaceae	Fageceae		
	146 (87)	212 (66)	0.001	K
	168 (30)	188 (33)	0.169	L&P
Hebeloma	Broad-leaved	Broad-leaved & coniferous		
	261 (24)	196 (12)	0.036	K
	262 (8)	200 (4)	0.048	L&P
Hygrophorus	Pinaceae	Fageceae		
	152 (13)	110 (11)	0.042	K
	138 (11)	112 (10)	0.338	L&P
Inocybe	Broad-leaved	Broad-leaved & coniferous		
	204 (67)	179 (25)	0.389	K
	203 (28)	173 (20)	0.112	L&P
Lactarius	Pinaceae	Betulaceae		
	204 (13)	147 (15)	0.002	K
	231 (14)	129 (11)	0.001	L&P
Leccinum	Betulaceae	Other genera		
	235 (5)	155 (4)	0.208	K
	232 (6)	174 (5)	0.24	L&P
Russula	Pinaceae	Fageceae		

	249 (18)	187 (55)	0.001	K
	266 (14)	199 (39)	0.001	L&P
Suillus	*Larix*	*Pinus*		
	93.8 (6)	59.4 (8)	0.041	K
	103 (6)	65 (9)	0.021	L&P
Tricoloma	Pinaceae	Fageceae		
	84 (20)	86.8 (13)	0.306	K
	76.2 (15)	85.1 (7)	0.381	L&P

* P-values were calculated by Brunner-Munzer test.

The numbers of species examined were indicated in round brackets.

In reference, K and L&P indicate Kibby (2020–2023) and Læssøe & Petersen (2019)

Fig. 1 Proportions of the fungal species occurring with a single genus of the host tree. Two independent descriptions of the species by Kibby (2020–2023) and Læssøe and Petersen (2019) are shown in closed and open bars respectively. The star on the bar indicates the fungal descriptions that statistically supported (P<0.05) spore volume differences are observed between species occurring in different host taxa.