Impaired fatty acid import or catabolism in macrophages restricts intracellular growth of

Mycobacterium tuberculosis

Nelson V. Simwela[1], Eleni Jaecklein [2], Christopher M. Sassetti[2], David G. Russell[1]*

1. Department of Microbiology and Immunology, College of Veterinary Medicine, Cornell

University, Ithaca, New York, USA

2. Department of Microbiology, UMass Chan Medical School, Worcester, Massachusetts,

USA

*Correspondence: dgr8@cornell.edu

Abstract

Mycobacterium tuberculosis (*Mtb*) infection of macrophages reprograms cellular metabolism to

promote lipid retention. While it is clearly known that intracellular *Mtb* utilize host derived lipids

to maintain infection, the role of macrophage lipid processing on the bacteria's ability to access

the intracellular lipid pool remains undefined. We utilized a CRISPR-Cas9 genetic approach to

assess the impact of sequential steps in fatty acid metabolism on the growth of intracellular *Mtb*.

Our analyzes demonstrate that macrophages which cannot either import, store or catabolize fatty

acids restrict *Mtb* growth by both common and divergent anti-microbial mechanisms, including

increased glycolysis, increased oxidative stress, production of pro-inflammatory cytokines,

enhanced autophagy and nutrient limitation. We also show that impaired macrophage lipid

droplet biogenesis is restrictive to *Mtb* replication, but increased induction of the same fails to

rescue *Mtb* growth. Our work expands our understanding of how host fatty acid homeostasis

impacts *Mtb* growth in the macrophage.

Introduction

Mycobacterium tuberculosis (*Mtb*), the causative agent of tuberculosis (TB), has caused disease and death in humans for centuries (WHO, 2023). *Mtb* primarily infects macrophages in the lung (Cohen et al., 2018; Wolf et al., 2007) wherein the bacterium relies on host derived fatty acids and cholesterol for synthesis of its lipid rich cell wall, and to produce energy and virulence factors (Peyron et al., 2008; Russell et al., 2009; Singh et al., 2012; Daniel et al., 2011; Muñoz-Elías and McKinney, 2005; Pandey and Sassetti, 2008; Brzostek et al., 2009). Within the lung microenvironment, resident alveolar macrophages preferentially oxidize fatty acids and are more permissive to *Mtb* growth while recruited interstitial macrophages are more glycolytic and restrictive of *Mtb* replication (Huang et al., 2018; Pisu et al., 2020; Pisu et al., 2021). Globally, *Mtb* infection modifies macrophage metabolism in a manner that enhances its survival. *Mtb* infected macrophages shift their mitochondrial substrate preference to exogenous fatty acids which drives the formation of foamy macrophages that are laden with cytosolic lipid droplets (Peyron et al., 2008; Russell et al., 2009; Cumming et al., 2018; Singh et al., 2012; Podinovskaia et al., 2013). Foamy macrophages are found in abundance in the central and inner layers of granulomas, a common histopathological feature of human TB (Russell et al., 2009; Kim et al., 2010). Interference with key regulators of lipid homeostasis, such as the miR-33 and the transcription factors peroxisome proliferator-activated receptor α (PPARα) and PPAR-γ enhances macrophage control of *Mtb* (Kim et al., 2017; Almeida et al., 2009; Ouimet et al., 2016). Moreover, compounds which modulate lipid metabolism such as the anti-diabetic drug metformin and some cholesterol lowering drugs are under investigation for host directed therapeutics (HDTs) against *Mtb* (Parihar et al., 2014; Singhal et al., 2014). Although the dependence of intracellular *Mtb* on host fatty acids and cholesterol is well documented (Wilburn et al., 2018), the impact of specific aspects of macrophage lipid metabolism on the bacteria remains opaque. In *Mtb* infected foamy macrophages, bacteria containing phagosomes are found in close apposition to intracellular lipid droplets (Peyron et al., 2008). It is believed that the bacterial induction of a foamy macrophage phenotype in host cells results in a steady supply of lipids that addresses the bacteria's nutritional requirements (Peyron et al., 2008; Russell et al., 2009; Singh et al., 2012; Daniel et al., 2011). In fact, intracellular *Mtb* has been shown to import fatty acids from host lipid droplet derived

triacylglycerols (Daniel et al., 2011). However, other studies indicate that macrophage lipid

droplet formation in response to *Mtb* infection can lead to the induction of a protective, anti-

microbial response (Knight et al., 2018). *Mtb* appears unable to acquire host lipids when lipid

droplets are induced by stimulation with interferon gamma (IFN-γ) (Knight et al., 2018).

Moreover, there is some evidence that lipid droplets can be sites for the production of host

protective pro-inflammatory eicosanoids (Knight et al., 2018; Daniel et al., 2011). Lipid droplets

can also act as innate immune hubs against intracellular bacterial pathogens by clustering anti-

bacterial proteins (Bosch et al., 2020). Inhibition of macrophage fatty acid oxidation by knocking

out mitochondrial carnitine palmitoyl transferase 2 (CPT2) or using chemical inhibitors of CPT2

also restrict intracellular growth of *Mtb* (Chandra et al., 2020; Huang et al., 2018). These data

demonstrate that modulation of the different stages in lipid metabolism inside *Mtb* infected

macrophages can result in conflicting outcomes.

We carried out a candidate-based, CRISPR mediated knockout of lipid import and metabolism

genes in macrophages to determine their roles in intracellular growth of *Mtb*. By targeting genes

involved in fatty acid import, sequestration, and metabolism in Hoxb8 derived conditionally

immortalized murine macrophages (Kiritsy et al., 2021), we show that impairing lipid homeostasis

in macrophages at different steps in the process negatively impacts the growth of intracellular

Mtb, albeit to differing degrees. The impact on *Mtb* growth in the mutant macrophages was

mediated through different mechanisms despite some common anti-microbial effectors. *Mtb*

infected macrophages deficient in the import of long chain fatty acids increased production of

pro-inflammatory markers such as interleukin 1β (IL-1β). In contrast, ablation of lipid droplet

biogenesis and fatty acid oxidation increased production of reactive oxygen species and limited

the bacteria's access to nutrients. We also found that suppression of *Mtb* growth in macrophages

that are unable to produce lipid droplets could not be rescued by exogenous addition of fatty

acids indicating that this is not purely nutritional restriction. Our data indicate that interference

of lipid metabolism in macrophages leads to suppression of *Mtb* growth via multiple routes.

Results

Knockout of fatty acid import and metabolism genes restricts *Mtb* growth in macrophages

To apply a holistic approach to assessing the role(s) of fatty acid metabolism on the intracellular growth of *Mtb*, we used a CRISPR genetic approach to knockout genes involved in lipid import (CD36, FATP1), lipid droplet formation (PLIN2) and fatty acid oxidation (CPT1A, CPT2) in mouse primary macrophages (Figure 1A). Deletion of CD36 or CPT2 from mouse macrophages has been shown to impair intracellular growth of *Mtb* (Hawkes et al., 2010; Chandra et al., 2020). But the role of specialized long chain fatty acid transporters (FATP1-6) on *Mtb* growth in macrophages is uncharacterized. FATP1 and FATP4 are the most abundant fatty acid transporter isoforms in macrophages (Nishiyama et al., 2018). PLIN2, or adipophilin, is known to be required for lipid droplet formation (Paul et al., 2008; Larigauderie et al., 2004). Five isoforms of mammalian perilipins (PLIN) are involved in lipid droplet biogenesis amongst which PLIN2 is the dominant isoform expressed in macrophages (Knight et al., 2018). However, macrophages derived from PLIN2$^{-/-}$ mice show no defects in production of lipid droplets nor do they impair intracellular growth of *Mtb* (Knight et al., 2018). We targeted each of these genes with at least 2 sgRNAs in Hoxb8 Cas9$^+$ conditionally immortalized myeloid progenitors (Kiritsy et al., 2021) to generate a panel of mutants that were deficient in the following candidates of interest; FATP1$^{-/-}$, PLIN2$^{-/-}$, CD36$^{-/-}$, CPT1A$^{-/-}$ and CPT2$^{-/-}$. Each individual sgRNA achieved >85% CRISPR-mediated deletion efficiency for all the 5 genes, as analyzed by the Inference for CRISPR Edits (ICE) tool (Conant et al., 2022) (Supplementary file 1). We verified the protein knockout phenotypes by flow cytometry and western blot analysis of differentiated macrophages derived from the CRISPR deleted Hoxb8 myeloid precursors (Figure 1 – figure supplement 1).

To confirm certain knockout phenotypes functionally, we checked lipid droplet biogenesis in PLIN2$^{-/-}$ macrophages in comparison to macrophages transduced with a non-targeting scramble sgRNA by confocal microscopy of BODIPY stained cells. Cells were cultured for 24 hours in the presence of exogenous oleate to enhance the formation of lipid droplets (Listenberger and Brown, 2007). We observed a complete absence of lipid droplet formation in PLIN2$^{-/-}$ macrophages as compared to controls (Figure 1 - figure supplement 2A). This is contrary to previous observations in macrophages derived from PLIN2 knockout mice which were reported

to have no defect in lipid droplet formation (Knight et al., 2018). We also assessed the ability of

CPT2$^{-/-}$ macrophages to oxidize fatty acids using the Agilent Seahorse XF Palmitate Oxidation

Stress Test. Scrambled sgRNA and CPT2$^{-/-}$ macrophages were cultured in substrate limiting

conditions and supplied with either bovine serum albumin (BSA) or BSA conjugated palmitate. As

shown in Figure 1 - figure supplement 2B, control macrophages were able to utilize and oxidize

palmitate in substrate limiting conditions indicated by a significant increase in oxygen

consumption rates (OCRs) in contrast to cells supplied with BSA alone. Addition of the CPT1A

inhibitor, etomoxir, inhibited the cell's ability to use palmitate in these conditions (Figure 1 - figure

supplement 2B). CPT2 knockout in CPT2$^{-/-}$ macrophages impaired the cell's ability to oxidize

palmitate to a degree comparable to etomoxir treatment as evidenced by baseline OCRs when

compared to scrambled sgRNA control (Figure 1 - figure supplement 2C).

We then assessed the different knockout mutant macrophages in their ability to support

the intracellular growth of *Mtb*. We infected macrophages with *Mtb* Erdman at a multiplicity of

infection (MOI) of 0.4 and assessed intracellular bacterial growth rates by counting colony forming

units (CFUs) 5 days post infection. All the 5 mutant macrophages significantly impaired *Mtb*

growth rates when compared to scrambled sgRNA as assessed by CFUs counts on day 5 (Figure

1C). PLIN2$^{-/-}$, FATP1$^{-/-}$ and CPT2$^{-/-}$ macrophages displayed the strongest growth restriction

phenotypes while CD36$^{-/-}$macrophages had a moderate, but significant, impact on *Mtb* growth.

We quantified intracellular bacteria on day 0, 3 hours post infection (Figure 1B), to ascertain that

subsequent differences on day 5 were not due to defects in initial phagocytosis. The moderate

growth restriction phenotypes of CD36$^{-/-}$ macrophages were consistent with previous findings

which reported similar impact on *Mtb* and *M. marinum* growth in macrophages derived from

CD36$^{-/-}$ mice (Hawkes et al., 2010). Impaired growth of *Mtb* in CPT1A$^{-/-}$ and CPT2$^{-/-}$ macrophages

is also consistent with previous reports that genetic and chemical inhibition of fatty acid oxidation

is detrimental to the growth of *Mtb* within macrophages (Huang et al., 2018; Chandra et al.,

2020).

174 ***Mtb* infected macrophages with impaired fatty acid import and metabolism display altered**
175 **mitochondrial metabolism and elevated glycolysis.**

176 Impairment of fatty acids metabolism by FATP1 knockout in macrophages rewires their substrate
177 bias from fatty acids to glucose (Johnson et al., 2016). We reasoned that deletion of genes
178 required for downstream processing of lipids (Figure 1A) could also reprogram macrophages and
179 increase glycolysis which could, in part, explain bacterial growth restriction. We analyzed the
180 metabolic states of 3 knockout macrophages (FATP1$^{-/-}$, PLIN2$^{-/-}$ and CPT2$^{-/-}$) in uninfected and *Mtb*
181 infected conditions by monitoring oxygen consumption rates (OCR) and extracellular acidification
182 rates (ECARs) using the Agilent Mito and Glucose Stress Test kits. All 3 mutant uninfected
183 macrophages displayed reduced mitochondrial respiration as evidenced by lower basal and spare
184 respiratory capacity (SRC) when compared to scrambled sgRNA controls (Figure 2 - figure
185 supplement 1A, 1B). *Mtb* infection proportionally reduced basal and SRC rates across all the
186 mutant macrophages and scrambled controls (Figure 2A, 2B) when compared to uninfected
187 macrophages, which is consistent with previous findings (Cumming et al., 2018). PLIN2$^{-/-}$
188 macrophages displayed the most marked reduction in mitochondrial activity in both uninfected
189 and infected conditions, while FATP1$^{-/-}$ macrophages were the least affected (Figure 2A, Figure 2
190 - figure supplement 1A). As reported previously (Johnson et al., 2016), uninfected FATP1$^{-/-}$
191 macrophages were more glycolytically active with higher basal and spare glycolytic capacity (SGC)
192 compared to scrambled controls (Figure 2 - figure supplement 1C, 1D). Uninfected PLIN2$^{-/-}$ and
193 CPT2$^{-/-}$ macrophages were also more glycolytically active, but to a greater degree than FATP1$^{-/-}$
194 (Figure 2 - figure supplement 1C, 1D). *Mtb* infection increased the glycolytic capacity of all the 3
195 mutant macrophages (Figure 2C, 2D). Overall, PLIN2$^{-/-}$ macrophages exhibited the highest
196 glycolytic capacity (Figure 2C, Figure 2 - figure supplement 1D). Our data indicates that
197 impairment of fatty acid metabolism at different steps significantly impacts mitochondrial
198 respiration and reprograms cells towards glycolysis. Increased glycolytic flux in macrophages has
199 been linked to the control of intracellular *Mtb* growth (Gleeson et al., 2016; Shi et al., 2015).
200 Metabolic realignment as a consequence of interference with lipid homeostasis, which results in
201 enhanced glycolysis may contribute to *Mtb* growth restriction in these mutant macrophages.

202

Knockout of lipid import and metabolism genes in macrophages activates AMPK and stabilizes HIF-1α

Mtb infection is known to induce increased glycolysis or the "Warburg effect" in macrophages, mouse lungs and human TB granulomas (Shi et al., 2015; Gleeson et al., 2016; Belton et al., 2016). Several studies have demonstrated that the Warburg effect is mediated by the master transcription factor hypoxia-inducible factor-1 (HIF-1) (Courtnay et al., 2015). During *Mtb* infection, HIF-1 is activated by production of reactive oxygen species (ROS), TCA cycle intermediates and hypoxia in the cellular microenvironments as a consequence of altered metabolic activities and increased immune cell functions (Li et al., 2024; Shi et al., 2015; Gleeson et al., 2016; Belton et al., 2016). We assessed HIF-1 stability in the 3 mutant macrophage lineages (FATP1$^{-/-}$, PLIN2$^{-/-}$ and CPT2$^{-/-}$) by monitoring total HIF-1α protein levels by western blot, having confirmed that they were all more glycolytically active than the scrambled controls (Figure 2, Figure 2 - figure supplement 1). Indeed, all the 3 mutant macrophages displayed significantly higher amounts of total HIF-1α when compared to scrambled controls in uninfected conditions and after *Mtb* infection for 48 hours (Figure 3A, 3B). We also checked the phosphorylation status of the adenosine monophosphate kinase (AMPK), a master regulator of cell energy homeostasis (Garcia and Shaw, 2017), in the mutant macrophages since Seahorse flux analyses indicated that they had impaired mitochondrial activities (Figure 2, Figure 2 - figure supplement 1). Western blot analysis revealed that in both *Mtb* infected and uninfected conditions, impaired fatty acid metabolism correlated with moderate activation of AMPK as indicated by increased AMPK phosphorylation (Figure 3C, 3D). These data point to a metabolic reprogramming of cells through activation of HIF-1α and AMPK to promote glycolysis. In energetically stressed cellular environments, activated AMPK promotes catabolic processes such as autophagy to maintain nutrient supply and energy homeostasis (Garcia and Shaw, 2017). Autophagy is also an innate immune defense mechanism against intracellular *Mtb* in macrophages (Gutierrez et al., 2004). We examined the levels of autophagic flux in the mutant macrophages by monitoring LC3I to LC3II conversion by western blot and by qPCR analysis of selected autophagy genes (*AMBRA1*, *ATG7*, *MAP1LC3B* and *ULK1)*. We observed a moderate increase in autophagic flux by western blot in uninfected FATP1$^{-/-}$ and CPT2$^{-/-}$ macrophages, which was amplified upon infection with *Mtb*

(Figure 2 - figure supplement 2A, 2B). Our qPCR analysis revealed that the 4 autophagy genes were upregulated in both *Mtb* infected and uninfected conditions in all the 3 mutants (Figure 2 - figure supplement 2C, 2D). These data suggest that impaired fatty acid import and metabolism in macrophages could be restricting *Mtb* growth by promoting autophagy. These data agree with previous observations that inhibition of fatty acid oxidation enhances macrophage xenophagic activity whichleads to improved control of *Mtb* (Chandra et al., 2020).

Exogenous oleate fails to rescue the *Mtb icl1*-deficient mutant in FATP1$^{-/-}$, PLIN2$^{-/-}$ and CPT2$^{-/-}$ macrophages

The mycobacterial isocitrate lyase (*icl1*) acts as an isocitrate lyase in the glyoxylate shunt and as a methyl-isocitrate lyase in the methyl-citrate cycle (MCC) (Gould et al., 2006; McKinney et al., 2000). *Mtb* uses the MCC to convert propionyl CoA originating from the breakdown of cholesterol rings and β-oxidation of odd chain fatty acids into succinate and pyruvate, which are eventually assimilated into the TCA cycle (Muñoz-Elías et al., 2006; Griffin et al., 2012). The buildup of propionyl CoA is toxic to *Mtb* and the bacteria relies on the MCC together with the incorporation of propionyl CoA to methyl-branched lipids in the cell wall as an internal detoxification system (Muñoz-Elías et al., 2006; Savvi et al., 2008). *Mtb* propionyl CoA toxicity is, in part, due to a cellular imbalance between propionyl CoA and acetyl CoA as an accumulation of the former or paucity of the latter results in the propionyl CoA mediated inhibition of pyruvate dehydrogenase (Lee et al., 2013). Consequently, *Mtb icl1* deficient mutants (*Mtb Δicl1*) are unable to grow in media supplemented with cholesterol or propionate, or intracellularly in macrophages (Lee et al., 2013). However, this growth inhibition could be rescued both in culture and in macrophages by exogenous supply of acetate or even chain fatty acids which can be oxidized to acetyl-CoA (Lee et al., 2013). We took advantage of this metabolic knowledge to assess whether exogenous addition of the even chain fatty acid oleate can rescue the intracellular growth of *Mtb Δicl1* mutants in our CRISPR knockout macrophages. Scrambled controls, FATP1$^{-/-}$, PLIN2$^{-/-}$ and CPT2$^{-/-}$ macrophages in normal macrophage media or media supplemented with oleate were infected with an *Mtb Δicl1* strain expressing mCherry at MOI 5. Bacterial growth measured by mCherry expression was recorded 5 days post infection. Consistent with previous observations (Lee et al., 2013), the *Mtb*

$\Delta icl1$ mutant failed to replicate in both mutant and scramble macrophages that were grown in

normal macrophage media as evidenced by baseline mCherry fluorescence (Figure 4A). Oleate

supplementation successfully rescued the *Mtb* $\Delta icl1$ mutant in scrambled control macrophages.

However, the growth restriction of the *Mtb* $\Delta icl1$ strain could not be alleviated by exogenous

addition of oleate to the mutant macrophages (FATP1$^{-/-}$, PLIN2$^{-/-}$ and CPT2$^{-/-}$) (Figure 4A). These

data suggest that impaired import (FATP1$^{-/-}$), sequestration (PLIN2$^{-/-}$) or β-oxidation of fatty acids

(CPT2$^{-/-}$) blocks *Mtb*'s ability to access and use cellular lipids.

Oleate supplementation in macrophages induces the formation of lipid droplets and we

were able to confirm the inability to produce lipid droplets in PLIN2$^{-/-}$ macrophages using this

approach (Figure 1 - figure supplement 2A). As an indirect measure to track the fate of

supplemented oleate in the mutant macrophages, we monitored lipid droplet biogenesis in

FATP1$^{-/-}$ and CPT2$^{-/-}$ macrophages to check if the inability to rescue the *Mtb* $\Delta icl1$ impaired growth

phenotypes in these mutant macrophages could be possibly related to disruptions in lipid droplet

biogenesis. Confocal analysis of BODIPY stained cells upon oleate supplementation revealed that

FATP1$^{-/-}$ macrophages also fail to generate lipid droplets (Figure 4B). In contrast, CPT2$^{-/-}$

macrophages produced more and larger lipid droplets in comparison to scrambled controls

(Figure 4B). These data suggest that the inhibition of *Mtb* growth in these mutant macrophages

is not merely through limitation of access to fatty acid nutrients.

Dual RNA sequencing to identify host and bacterial determinants of *Mtb* restriction in mutant

macrophage lineages.

We performed RNA sequencing of both host and bacteria in *Mtb* infected mutant macrophages

as a preliminary step in identification of pathways restricting bacterial growth. We infected

scrambled controls, FATP1$^{-/-}$, PLIN2$^{-/-}$ and CPT2$^{-/-}$ macrophages with the *Mtb* smyc'::mCherry

strain for 4 days and processed the samples for dual RNA sequencing as previously described

(Simwela et al., 2024). Principal component analysis (PCA) of host transcriptomes revealed a

clustering of all the 3 mutant macrophages away from scrambled controls (Figure 5A).

Interestingly, there was a separation in transcriptional responses between the 3 mutant

macrophages as CPT2$^{-/-}$ and PLIN2$^{-/-}$ macrophages clustered closer together and more distant

from FATP1$^{-/-}$ (Figure 5A). Overall, using an adjusted p-value < 0.05 and absolute log2 fold change

> 1.2, we identified 900 genes which were differentially expressed (DE) in PLIN2$^{-/-}$ macrophages

(589 up, 311 down), 817 genes which were DE in FATP1$^{-/-}$ macrophages (501 up, 315 down) and

189 genes which were DE in CPT2$^{-/-}$ (124 up, 65 down) (Supplementary file 2, Figure 5B).

Consistent with the PCA analysis, Venn diagram of the DE genes (Figure 5B) indicated divergent

responses in the 3 mutant macrophage populations. We performed pathway enrichment analysis

(Wu et al., 2021) of the DE genes to identify anti-microbial pathway candidates in the 3 mutant

macrophages. We found that defects in fatty acid uptake in FATP1$^{-/-}$ infected macrophages

upregulated pro-inflammatory pathways involved in MAPK and ERK signaling and production of

inflammatory cytokines (IFN-γ, interleukin-6, Interleukin-1α, β) (Figure 5D, Supplementary file 3).

The pro-inflammatory signatures of the FATP1$^{-/-}$ macrophages are consistent with previous

observations that demonstrated that a deficiency in FATP1 exacerbated macrophage activation *in*

vitro and *in vivo* (Johnson et al., 2016). FATP1 is a solute carrier family member (SLC27A1) and

Mtb infected FATP1$^{-/-}$ macrophages showed reduced expression of other solute carrier

transporters (SLC) such as FATP4 (SLC27A4), GLUT1 (SLC2A1) and 8 SLC amino acid transporters

(Figure 5 - figure supplement 1A, 1B, 2A, Supplementary file 2). Interestingly, *Mtb* infected FATP1^{-}

$^{/-}$ macrophage transcriptomes exhibited upregulation of macrophage scavenger receptors (MSR1)

and the ATP binding cassette transporter ABCC1 (Figure 5 - figure supplement 1A), both of which

can independently transport fatty acids into cells (Vogel et al., 2022; Raggers et al., 1999).

Meanwhile, inability generate lipid droplets in *Mtb* infected PLIN2$^{-/-}$ macrophages led to

upregulation in pathways involved in ribosomal biology, MHC class 1 antigen presentation,

canonical glycolysis, ATP metabolic processes and type 1 interferon responses (Figure 5C,

Supplementary file 3). In the downregulated PLIN2$^{-/-}$ DE gene set, enriched pathways included

those involved in production of pro-inflammatory cytokines; interleukin-6 and 8, IFN-γ and

interleukin-1 (Figure 5 - figure supplement 2B, Supplementary file 3). Oxidative phosphorylation

and processes involved in the respiratory chain electron transport were also significantly enriched

in PLIN2$^{-/-}$ downregulated genes. This suggests that *Mtb* infected PLIN2$^{-/-}$ macrophages increase

glycolytic flux and decrease mitochondrial activities, which is consistent with our metabolic flux

analysis data (Figure 2). Unlike FATP1$^{-/-}$ macrophages, PLIN2$^{-/-}$ macrophages are, however, broadly

anti-inflammatory as most pro-inflammatory genes were downregulated upon *Mtb* infection.

Further downstream in the lipid processing steps, inhibition of fatty acid oxidation in CPT2^{-}

$^{/-}$ macrophages upregulated pathways involved in MHC class 1 antigen presentation, response to

IFN-γ and interleukin-1 and T-cell mediated immunity (Figure 5 - figure supplement 3,

Supplementary file 3). There was a limited overlap in enriched pathways in the upregulated genes

between *Mtb* infected CPT2$^{-/-}$ and FATP1$^{-/-}$ macrophages such as those involved in the cellular

responses to interleukin-1 and IFN-γ. However, many pathways over-represented in CPT2$^{-/-}$

macrophages were common to PLIN2$^{-/-}$ macrophages (Figure 5C, Figure 5 - figure supplement 3,

Supplementary file 3). Similarly, both *Mtb*-infected PLIN2$^{-/-}$ *and* CPT2$^{-/-}$ macrophages were

downregulated in expression of genes involved in oxidative phosphorylation (Supplementary file

3). We confirmed expression levels of key genes by qPCR analysis of IL-1β and the type 1

interferon (IFN-β) response during the early time points of infection. Indeed, 4 hours post

infection, IL-1β and IFN-β were both upregulated in FATP1$^{-/-}$ macrophages as compared to

scrambled controls consistent with their pro-inflammatory phenotype (Figure 5 - figure

supplement 4A, 4B). On the contrary, PLIN2$^{-/-}$ macrophages downregulated IL-1β (Figure 5 - figure

supplement 4B). These data indicate that macrophages respond quite divergently to the deletion

of the different steps in fatty acid uptake, which implies that the intracellular pressures to which

Mtb is exposed may also differ.

Oxidative stress and nutrient limitation are major stresses experienced by *Mtb* in PLIN2$^{-/-}$ and

CPT2$^{-/-}$ macrophages

We also analyzed transcriptomes from intracellular *Mtb* from scrambled controls, FATP1$^{-/-}$, CPT2^{-}

$^{/-}$ and PLIN2$^{-/-}$ macrophages in parallel with host transcriptomes in Figure 5A. Using an adjusted

p-value of < 0.1 and an absolute log2 fold change > 1.4, 0 genes were DE in FATP1$^{-/-}$ macrophages,

105 *Mtb* genes were DE in PLIN2$^{-/-}$ macrophages (69 up, 36 down) and 10 genes were DE in CPT2^{-}

$^{/-}$ macrophages (3 up, 7 down) (Figure 6A). Despite being restrictive to *Mtb* growth (Figure 1C)

and appearing more pro-inflammatory (Figure 5D), FATP1$^{-/-}$ macrophages did not elicit a

detectable shift in the transcriptional response in *Mtb,* compared to control host cells. We

speculate that pro-inflammatory responses in FATP1$^{-/-}$ macrophages could be enough to restrict the growth of bacteria, but the resulting compensatory responses as evidenced by the upregulation of macrophage scavenger receptors (Figure 5 - figure supplement 1A) alleviate some of the stresses that a lack of fatty acid import could be duly exerting on the bacteria. PLIN2$^{-/-}$ macrophages appeared to elicit the strongest transcriptional response from *Mtb* which is consistent with our CFU data (Figure 1C) as PLIN2$^{-/-}$ macrophages exhibited the strongest growth restriction. Among the DE genes in *Mtb* from PLIN2$^{-/-}$ macrophages, a significant number of upregulated genes are involved in nutrient assimilation (Figure 6B). *Mtb* in PLIN2$^{-/-}$ macrophages upregulated CobU (Rv0254c) which is predicted to be involved in the bacteria's cobalamin (Vitamin B$_{12}$) biosynthesis. Vitamin B$_{12}$ is an important cofactor for the activity of *Mtb* genes required for cholesterol and fatty acid utilization (Campos-Pardos et al., 2024; Savvi et al., 2008). Genes involved in de novo long chain fatty acid synthesis (AccE5, Rv281) (Bazet Lyonnet et al., 2014), cholesterol breakdown (HsaD, Rv3569c) (Lack et al., 2010), β-oxidation of fatty acids (EchA18, Rv3373; FadE22, Rv3061c) (Schnappinger et al., 2003), purine salvage (Apt, Rv2584c) (Warner et al., 2014) and tryptophan metabolism (Lott, 2020) were also upregulated in *Mtb* from PLIN2$^{-/-}$ macrophages (Figure 6B). This metabolic re-alignment response is seen most frequently under nutrient limiting conditions (Huang et al., 2018; Pisu et al., 2020; Theriault et al., 2022). *Mtb* in PLIN2$^{-/-}$ macrophages also appears to experience a significant level of other cellular stresses as genes involved in DNA synthesis and repair, general response to oxidative stress and PH survival in the phagosome (DnaN, Rv0002; RecF, Rv0003; DinF, Rv2836c; Rv3242c, Rv1264) were upregulated (Figure 6B). Amongst the down-regulated genes in *Mtb* in PLIN2$^{-/-}$ macrophages, FurA (Rv1909c), a KatG repressor was the most significant (Figure 6B, Supplementary file 4). FurA downregulation derepresses the catalase peroxidase, KatG, which promotes *Mtb* survival in oxidative stress conditions (Zahrt et al., 2001). These data suggest that PLIN2$^{-/-}$ macrophages could be, in part, restricting *Mtb* growth by increasing production of ROS. The data also suggest that, contrary to a previous report (Knight et al., 2018), blocking lipid droplet formation in host macrophages does place increased nutritional and oxidative stress on intracellular *Mtb*.

CPT2$^{-/-}$ macrophages elicited a modest shift in transcriptional response from *Mtb*, and the

majority of the DE genes (8 out of 10) were also DE in PLIN2$^{-/-}$ macrophages (Supplementary file

4). This in in agreement with the host transcription response as CPT2$^{-/-}$ and PLIN2$^{-/-}$ macrophages

share similar candidate anti-bacterial responses (Figure 5A-C, Figure 5 - figure supplement 3). In

common with the *Mtb* transcriptome response in PLIN2$^{-/-}$ macrophages, upregulated genes in

Mtb isolated from CPT2$^{-/-}$ macrophages included those involved in response to oxidative stress

(DinF, Rv2836c) (Supplementary file 4). To substantiate some of these pathways, we assessed the

levels of total cellular ROS in FATP1$^{-/-}$, PLIN2$^{-/-}$ and CPT2$^{-/-}$ macrophages in both *Mtb* infected and

uninfected conditions by staining the cells with the Invitrogen CellROX dye and confocal

microscopy analysis of live stained cells. In both infected and uninfected conditions, all the 3

mutants displayed significantly elevated total cellular ROS when compared to scramble controls

(Fig. 6 - figure supplement 1A, 1B).

Inhibitors of lipid metabolism block intracellular growth of *Mtb* in macrophages but not in broth

culture

We next examined if pharmacological inhibitors would phenocopy the growth inhibition

phenotypes we observed with specific gene deletions. Compounds which modulate lipid

homeostasis are currently being investigated for HDT against TB, which is an area of considerable

interest (Kim et al., 2020). Such compounds include metformin, a widely used anti-diabetic drug

which activates AMPK, inhibits fatty acid synthesis and promotes β-oxidation of fatty acids

(Fullerton et al., 2013; Singhal et al., 2014). Chemical inhibition of fatty acid β-oxidation is already

known to promote macrophage control of *Mtb* (Chandra et al., 2020; Huang et al., 2018). We

targeted macrophage lipid homeostasis with trimetazidine (TMZ), an inhibitor of β-oxidation of

fatty acids, metformin, and an FATP1 inhibitor, FATP1 In (Matsufuji et al., 2013). We assessed the

impact of these compounds on extracellular *Mtb* cultured in broth over 9 days in the presence of

the inhibitors (DMSO, TMZ; 500 nM, Metformin; 2 mM, FATP1 In; 10 μM, Rifampicin; 0.5 μg/ml).

None of the 3 lipid metabolism inhibitors had a measurable effect on *Mtb* growth in liquid culture

media when compared to DMSO controls (Figure 7A). Treatment with rifampicin completely

blocked bacterial growth under the same conditions (Figure 7A). We next infected scrambled

sgRNA control macrophages with *Mtb* at MOI 0.5. Inhibitors were added to infected macrophages

3 hours post infection and bacterial CFUs were enumerated 4 days post treatment. Consistent

with previous observations (Chandra et al., 2020; Singhal et al., 2014), TMZ and metformin

significantly reduced bacterial loads in macrophages when compared to DMSO controls (Figure

7B). Similarly, FATP1 In also impacted the intracellular growth of *Mtb* (Figure 7B). The results

provide independent data that both genetic and chemical modulation of fatty acid metabolism at

different steps in the process negatively impacts intracellular growth of *Mtb*.

Discussion

It is clearly established that host derived fatty acids and cholesterol are important carbon sources

for *Mtb* (Wilburn et al., 2018). However, the relationship between *Mtb* and the infected host cell

lipid metabolism remains a subject of conjecture. In this report, we characterized the role of

macrophage lipid metabolism on the intracellular growth of *Mtb* by a targeted CRISPR mediated

knockout of host genes involved in fatty acid import, sequestration and catabolism. Macrophage

fatty acid uptake is mediated by the scavenger receptor CD36 and specialized long chain fatty acid

transporters, FATP1 and FATP4 (Deng et al., 2023). Earlier studies reported that a deficiency of

CD36 enhances macrophage control of *Mtb,* albeit to a modest degree (Hawkes et al., 2010).

Work from Hawkes et al.(Hawkes et al., 2010) indicated that the anti-microbial effectors in CD36

deficient macrophages were not due to bacterial uptake deficiencies, differences in the rate of

Mtb induced host cell death, production of ROS or pro-inflammatory cytokines (Hawkes et al.,

2010). We similarly observed a moderate *Mtb* growth restriction phenotype in our CRISPR

generated CD36$^{-/-}$ macrophages. However, a strong growth restriction of *Mtb* was observed when

we knocked out the long chain fatty acid transporter, FATP1. FATP1$^{-/-}$ macrophages displayed

altered metabolism characterized by stabilization of HIF-1α, activated AMPK, increased glycolysis

and reduced mitochondrial functions. Given that both CD36 and FATP1 perform similar functions,

it is expected that they should be some degree of compensation between the transporters when

either of the genes are deleted. Indeed, we found out that FATP1 knockout resulted in the

upregulation of other lipid import transporters (MSR1, ABCC1). This would be consistent with the

moderate anti-*Mtb* phenotypes in CD36$^{-/-}$ macrophages which could easily be compensated by

the presence of FATPs to alleviate the reduction in fatty acid supply experienced by intracellular

Mtb. However, the FATP1$^{-/-}$ macrophage phenotype appears to be more severe on *Mtb* and could

be exacerbated by an elevated pro-inflammatory response as has been reported previously both

in vitro and *in vivo* (Johnson et al., 2016).

After uptake into the cells, most fatty acids either undergo β-oxidation in the mitochondria to

provide energy or are esterified with glycerol phosphate to form triacylglycerols that may be

incorporated into lipid droplets in the endoplasmic reticulum (Deng et al., 2023). Over the last 2

decades, it has been believed that lipid droplets are a nutrient source for *Mtb* in macrophages

(Russell et al., 2009; Daniel et al., 2011; Peyron et al., 2008; Singh et al., 2012). However, recent

work indicates lipid droplets may serve as centers for the production of pro-inflammatory markers

and anti-microbial peptides (Knight et al., 2018; Bosch et al., 2020). It has also been reported that

bone marrow macrophages derived from PLIN2$^{-/-}$ mice did not have defects in the formation of

lipid droplets and supported robust intracellular *Mtb* replication (Knight et al., 2018). However,

our mutant macrophages with myeloid specific knockout of PLIN2 are unable to form lipid

droplets and are defective in supporting the growth of *Mtb*. The discrepancies with previous

observations (Knight et al., 2018) could be a consequence of compensatory responses to PLIN2

knockout in whole mice which when performed at embryonic level would allow for sufficient time

for the cells to recover by upregulating related PLIN isoforms. In fact, PLIN2$^{-/-}$ macrophages

displayed the strongest anti-*Mtb* phenotypes amongst our mutants exhibiting activated AMPK,

increased glycolysis and autophagy and impaired mitochondrial functions. *Mtb* isolated from

PLIN2$^{-/-}$ macrophages displayed signatures of severe nutrient limitation and oxidative stress

damage.

It has also been previously reported that chemical, genetic or miR33 mediated blockade

of fatty acid β-oxidation in macrophages induces lipid droplet formation (Chandra et al., 2020;

Ouimet et al., 2016) but that this enhanced lipid droplet formation does not correlate with

improved intracellular *Mtb* growth (Chandra et al., 2020). We observed a similar phenotype as

CPT2$^{-/-}$ macrophages that generated larger and more abundant lipid droplets than scrambled

control macrophages were still restrictive to *Mtb* growth. This implies that the presence or

absence of lipid droplets does not in itself indicate whether a macrophage will support or restrict

Mtb growth, and that the anti-microbial environment extends beyond simple nutrient availability.

In summary, our study shows that blocking macrophage's ability to import, sequester or

catabolize fatty acids chemically or by genetic knockout impairs *Mtb* intracellular growth. There

are shared features between potential anti-microbial effectors in macrophages which lack the

ability to import (FATP[-/-]) or metabolize fatty acids (PLIN2[-/-], CPT2[-/-]) such as increased glycolysis,

stabilized HIF-1α, activated AMPK, enhanced autophagy and production of ROS. However, there

are also intriguing points of divergence as FATP[-/-] macrophages are more pro-inflammatory while

PLIN2[-/-] macrophages appear to be broadly anti-inflammatory. The routes to *Mtb* growth

restriction in these mutant macrophages are clearly more complex than the bacteria's inability to

acquire nutrients. The data further emphasizes that targeting fatty acid homeostasis in

macrophages at different steps in the process (uptake, storage and catabolism) is worthy of

exploring in the development of new therapeutics against tuberculosis.

Materials and methods

All materials and methods are as described (Simwela et al., 2024) unless otherwise specified.

Flow cytometry and western blot analysis

Generation of CRISPR mutant Hoxb8 macrophages was carried out described previously (Simwela

et al., 2024). Antibodies used for both western blot and flow cytometry were as follows: rat anti-

Mouse CD36:Alexa Fluor®647 (Biorad, 10 μl/million cells), rabbit anti-PLIN2 (Proteintech, 1:1000),

rabbit anti-FATP1 (Affinity biosciences, 1:1000), rabbit anti-CPT1A antibody (Proteintech, 1:1000),

rabbit anti-CPT2 antibody (Proteintech, 1:1000), rabbit anti-HIF-1α antibody (Proteintech,

1:1000), rabbit anti-AMPKα(1:1000, Cell signalling Technology), rabbit anti-Phospho-

AMPKα(1:500, Cell signalling Technology), rabbit anti-LC3B (1:1000, Cell signalling Technology)

and rabbit anti-β Actin (1:1000, Cell Signalling Technology). For western blots, secondary

antibodies used were anti-rabbit/mouse StarBright Blue 700 (1:2500, Biorad). Blots were

developed and imaged as described previously (Simwela et al., 2024).

Staining for cellular lipid droplets

Staining for cellular lipid droplets

Macrophages monolayers in Ibidi eight-well chambers were supplemented with exogenous 400

µM oleate for 24 hours to induce the formation of lipid droplets (Listenberger and Brown, 2007).

Cells were then fixed in 4% paraformaldehyde and stained with BODIPY™ 493/503 (Invitrogen, 1

µg/ml) in 150 mM sodium chloride. Stained cells were mounted with media containing DAPI and

imaged using a Leica SP5 confocal microscope.

Seahorse XF palmitate oxidation stress test

A modified Seahorse mitochondrial stress test was used to measure macrophage's ability to

oxidize palmitate in substrate limiting conditions. 2 days before the assay, 1×10^5 cells were plated

in Seahorse cell culture mini plates. 1 day before the assay, macrophage media was replaced with

the Seahorse substrate limited growth media (DMEM without pyruvate supplemented with 0.5

mM glucose, 1 mM glutamine, 1% FBS and 0.5 mM L-Carnitine). On the day of the assay, substrate

limited media was replaced with assay media (DMEM without pyruvate supplemented with 2 mM

glucose and 0.5 mM L-Carnitine). In selected treatment conditions, cells were either supplied with

bovine serum albumin (BSA), BSA palmitate or BSA palmitate plus etomoxir (4 µM). Oxygen

consumption rates (OCRs) were measured using the Mito Stress Test assay conditions as

described previously (Simwela et al., 2024).

Rescue of the *Mtb* Δicl1 mutant in oleate supplemented media

The *Mtb* H37Rv Δ*icl1* mutant expressing mCherry (Lee et al., 2013) was used for the rescue

experiments. The strain was maintained in 7H9 OADC broth as previously described (Simwela et

al., 2024) in the presence of Kanamycin (25 µg/ml) and hygromycin (50 µg/ml). 24 hours before

infection, macrophages were cultured in normal macrophage media or media supplemented with

400 µM oleate. Cells were then infected with the *Mtb* Δ*icl1* mutant at MOI 5. The bacterial

mCherry signal was measured on day 0 and day 5 post infection on an Envision plate reader

(PerkinElmer). Oleate was maintained throughout the experiment in the rescue assay conditions.

Measurement of total cellular ROS

Uninfected or *Mtb* infected macrophages monolayers in Ibidi eight-well chambers were stained with the CellROX Deep Red dye (Invitrogen) as per manufacturer's staining protocol. Live stained cells were imaged on a Leica SP5 confocal microscope. Z-stacks were re-constructed in ImageJ from which mean fluorescence intensities (MFI) for individual cells were obtained.

Datasets

The RNA-seq data from the Dual RNA-seq analysis of infected mouse macrophages,542 which includes both macrophage and *Mtb* reads are available in GEO: GSE270571

Acknowledgements

We would like to thank Dr. Jen K. Grenier and Ann E. Tate from the Cornell BRC Transcriptional Regulation and Expression Facility for their help with the development of dual RNA-Seq protocols. This work was supported by grants from the National Institutes of Health (AI155319, AI162598, and OD032135), Bill and Melinda Gates Foundation and the Mueller Health Foundation to D.G.R. EJ was supported by T32AI007349.

References

Almeida, P. E., Silva, A. R., Maya-Monteiro, C. M., Töröcsik, D., D'avila, H., Dezsö, B., et al. 2009. *Mycobacterium bovis* bacillus Calmette-Guérin infection induces TLR2-dependent peroxisome proliferator-activated receptor gamma expression and activation: functions in inflammation, lipid metabolism, and pathogenesis. J Immunol*, ***183,*** 1337-1345.

Bazet Lyonnet, B., Diacovich, L., Cabruja, M., Bardou, F., Quémard, A., Gago, G. & Gramajo, H. 2014. Pleiotropic effect of AccD5 and AccE5 depletion in acyl-coenzyme A carboxylase activity and in lipid biosynthesis in Mycobacteria. PLoS One*, ***9,*** e99853.

Belton, M., Brilha, S., Manavaki, R., Mauri, F., Nijran, K., Hong, Y. T., et al. 2016. Hypoxia and tissue destruction in pulmonary TB. Thorax*, ***71,*** 1145-1153.

Bosch, M., Sánchez-Álvarez, M., Fajardo, A., Kapetanovic, R., Steiner, B., Dutra, F., et al. 2020.

Mammalian lipid droplets are innate immune hubs integrating cell metabolism and host defense.

Science, **370**, eaay8085

Brzostek, A., Pawelczyk, J., Rumijowska-Galewicz, A., Dziadek, B. & Dziadek, J. 2009.

Mycobacterium tuberculosis is able to accumulate and utilize cholesterol. J Bacteriol, **191,** 6584-

6591.

Campos-Pardos, E., Uranga, S., Picó, A., Gómez, A. B. & Gonzalo-Asensio, J. 2024. Dependency on

host Vitamin B12 has shaped *Mycobacterium tuberculosis* Complex evolution. Nat Commun, **15,**

2161.

Chandra, P., He, L., Zimmerman, M., Yang, G., Köster, S., Ouimet, M., et al. 2020. Inhibition of fatty

acid oxidation promotes macrophage control of *Mycobacterium tuberculosis*. mBio, **11**, e01139-

20

Cohen, S. B., Gern, B. H., Delahaye, J. L., Adams, K. N., Plumlee, C. R., Winkler, J. K., et al. 2018.

Alveolar macrophages provide an early *Mycobacterium tuberculosis* niche and initiate

dissemination. Cell Host Microbe, **24,** 439-446.e4.

Conant, D., Hsiau, T., Rossi, N., Oki, J., Maures, T., Waite, K., et al. 2022. Inference of CRISPR edits

from Sanger trace data. CRISPR J, **5,** 123-130.

Courtnay, R., Ngo, D. C., Malik, N., Ververis, K., Tortorella, S. M. & Karagiannis, T. C. 2015. Cancer

metabolism and the Warburg effect: the role of HIF-1 and PI3K. Mol Biol Rep, **42,** 841-851.

Cumming, B. M., Addicott, K. W., Adamson, J. H. & Steyn, A. J. 2018. *Mycobacterium tuberculosis*

induces decelerated bioenergetic metabolism in human macrophages. Elife, **7**, e39169

Daniel, J., Maamar, H., Deb, C., Sirakova, T. D. & Kolattukudy, P. E. 2011. *Mycobacterium*

tuberculosis uses host triacylglycerol to accumulate lipid droplets and acquires a dormancy-like

phenotype in lipid-loaded macrophages. PLoS Pathog, **7,** e1002093.

Deng, L., Kersten, S. & Stienstra, R. 2023. Triacylglycerol uptake and handling by macrophages:

from fatty acids to lipoproteins. Progress in Lipid Research*, **92,** 101250.

Fullerton, M. D., Galic, S., Marcinko, K., Sikkema, S., Pulinilkunnil, T., Chen, Z. P., et al. 2013. Single

phosphorylation sites in ACC1 and ACC2 regulate lipid homeostasis and the insulin-sensitizing

effects of metformin. Nat Med*, **19,** 1649-1654.

Garcia, D. & Shaw, R. J. 2017. AMPK: mechanisms of cellular energy sensing and restoration of

metabolic balance. Mol Cell*, **66,** 789-800.

Gleeson, L. E., Sheedy, F. J., Palsson-Mcdermott, E. M., Triglia, D., O'leary, S. M., O'sullivan, M. P.,

et al. 2016. Cutting edge: *Mycobacterium tuberculosis* induces aerobic glycolysis in human

alveolar macrophages that is required for control of intracellular bacillary replication. J Immunol*,

196, 2444-2449.

Gould, T. A., Van De Langemheen, H., Muñoz-Elías, E. J., Mckinney, J. D. & Sacchettini, J. C. 2006.

Dual role of isocitrate lyase 1 in the glyoxylate and methylcitrate cycles in *Mycobacterium*

tuberculosis. Mol Microbiol*, **61,** 940-947.

Griffin, J. E., Pandey, A. K., Gilmore, S. A., Mizrahi, V., Mckinney, J. D., Bertozzi, C. R. & Sassetti, C.

M. 2012. Cholesterol catabolism by *Mycobacterium tuberculosis* requires transcriptional and

metabolic adaptations. Chem Biol*, **19,** 218-227.

Gutierrez, M. G., Master, S. S., Singh, S. B., Taylor, G. A., Colombo, M. I. & Deretic, V. 2004.

Autophagy is a defense mechanism inhibiting BCG and *Mycobacterium tuberculosis* survival in

infected macrophages. Cell*, **119,** 753-766.

Hawkes, M., Li, X., Crockett, M., Diassiti, A., Finney, C., Min-Oo, G., et al. 2010. CD36 deficiency

attenuates experimental Mycobacterial infection. BMC Infect Dis*, **10,** 299.

Huang, L., Nazarova, E. V., Tan, S., Liu, Y. & Russell, D. G. 2018. Growth of *Mycobacterium*

tuberculosis in vivo segregates with host macrophage metabolism and ontogeny. J Exp Med*, **215,**

1135-1152.

Johnson, A. R., Qin, Y., Cozzo, A. J., Freemerman, A. J., Huang, M. J., Zhao, L., et al. 2016. Metabolic
reprogramming through fatty acid transport protein 1 (FATP1) regulates macrophage
inflammatory potential and adipose inflammation. Mol Metab*, ***5,** 506-526.

Kim, J. S., Kim, Y. R. & Yang, C. S. 2020. Host-directed therapy in tuberculosis: targeting host
metabolism. Front Immunol*, ***11,** 1790.

Kim, M. J., Wainwright, H. C., Locketz, M., Bekker, L. G., Walther, G. B., Dittrich, C., et al. 2010.
Caseation of human tuberculosis granulomas correlates with elevated host lipid metabolism.
EMBO Mol Med*, ***2,** 258-274.

Kim, Y. S., Lee, H. M., Kim, J. K., Yang, C. S., Kim, T. S., Jung, M., et al. 2017. PPAR-α Activation
mediates innate host defense through induction of TFEB and lipid catabolism. J Immunol*, ***198,**
3283-3295.

Kiritsy, M. C., Mccann, K., Mott, D., Holland, S. M., Behar, S. M., Sassetti, C. M. & Olive, A. J. 2021.
Mitochondrial respiration contributes to the interferon gamma response in antigen-presenting
cells. Elife*, ***10**, e65109

Knight, M., Braverman, J., Asfaha, K., Gronert, K. & Stanley, S. 2018. Lipid droplet formation in
Mycobacterium tuberculosis infected macrophages requires IFN-γ/HIF-1α signaling and supports
host defense. PLoS Pathog*, ***14,** e1006874.

Lack, N. A., Yam, K. C., Lowe, E. D., Horsman, G. P., Owen, R. L., Sim, E. & Eltis, L. D. 2010.
Characterization of a carbon-carbon hydrolase from *Mycobacterium tuberculosis* involved in
cholesterol metabolism. J Biol Chem*, ***285,** 434-443.

Larigauderie, G., Furman, C., Jaye, M., Lasselin, C., Copin, C., Fruchart, J. C., et al. 2004. Adipophilin
enhances lipid accumulation and prevents lipid efflux from THP-1 macrophages: potential role in
atherogenesis. Arterioscler Thromb Vasc Biol*, ***24,** 504-510.

Lee, W., Vanderven, B. C., Fahey, R. J. & Russell, D. G. 2013. Intracellular *Mycobacterium*
tuberculosis exploits host-derived fatty acids to limit metabolic stress. J Biol Chem*,* **288,** 6788-
6800.

Li, C., Wang, J., Xu, J. F., Pi, J. & Zheng, B. 2024. Roles of HIF-1α signaling in *Mycobacterium*
tuberculosis infection: new targets for anti-TB therapeutics? Biochem Biophys Res Commun*,* **711,**
149920.

Listenberger, L. L. & Brown, D. A. 2007. Fluorescent detection of lipid droplets and associated
proteins. Curr Protoc Cell Biol*,* **Chapter 24,** Unit 24.2.

Lott, J. S. 2020. The tryptophan biosynthetic pathway is essential for *Mycobacterium tuberculosis*
to cause disease. Biochem Soc Trans*,* **48,** 2029-2037.

Matsufuji, T., Ikeda, M., Naito, A., Hirouchi, M., Kanda, S., Izumi, M., et al. 2013. Arylpiperazines
as fatty acid transport protein 1 (FATP1) inhibitors with improved potency and pharmacokinetic
properties. Bioorg Med Chem Lett*,* **23,** 2560-2565.

Mckinney, J. D., Höner Zu Bentrup, K., Muñoz-Elías, E. J., Miczak, A., Chen, B., Chan, W. T., et al.
2000. Persistence of *Mycobacterium tuberculosis* in macrophages and mice requires the
glyoxylate shunt enzyme isocitrate lyase. Nature*,* **406,** 735-738.

Muñoz-Elías, E. J. & Mckinney, J. D. 2005. *Mycobacterium tuberculosis* isocitrate lyases 1 and 2
are jointly required for in vivo growth and virulence. Nat Med*,* **11,** 638-644.

Muñoz-Elías, E. J., Upton, A. M., Cherian, J. & Mckinney, J. D. 2006. Role of the methylcitrate cycle
in *Mycobacterium tuberculosis* metabolism, intracellular growth, and virulence. Mol Microbiol*,*
60, 1109-22.

Nishiyama, K., Fujita, T., Fujimoto, Y., Nakajima, H., Takeuchi, T. & Azuma, Y.-T. 2018. Fatty acid
transport protein 1 enhances the macrophage inflammatory response by coupling with ceramide
and c-Jun N-terminal kinase signaling. Int Immunopharmacol*,* **55,** 205-215.

Ouimet, M., Koster, S., Sakowski, E., Ramkhelawon, B., Van Solingen, C., Oldebeken, S., et al. 2016. *Mycobacterium tuberculosis* induces the miR-33 locus to reprogram autophagy and host lipid metabolism. Nat Immunol*, **17,** 677-686.

Pandey, A. K. & Sassetti, C. M. 2008. *Mycobacterial* persistence requires the utilization of host cholesterol. Proc Natl Acad Sci U S A*, **105,** 4376-4380.

Parihar, S. P., Guler, R., Khutlang, R., Lang, D. M., Hurdayal, R., Mhlanga, M. M., et al. 2014. Statin therapy reduces the *Mycobacterium tuberculosis* burden in human macrophages and in mice by enhancing autophagy and phagosome maturation. J Infect Dis*, **209,** 754-763.

Paul, A., Chang, B. H., Li, L., Yechoor, V. K. & Chan, L. 2008. Deficiency of adipose differentiation-related protein impairs foam cell formation and protects against atherosclerosis. Circ Res*, **102,** 1492-1501.

Peyron, P., Vaubourgeix, J., Poquet, Y., Levillain, F., Botanch, C., Bardou, F., et al. 2008. Foamy macrophages from tuberculous patients' granulomas constitute a nutrient-rich reservoir for *M. tuberculosis* persistence. PLoS Pathog*, **4,** e1000204.

Pisu, D., Huang, L., Grenier, J. K. & Russell, D. G. 2020. Dual RNA-seq of Mtb-infected macrophages in vivo reveals ontologically distinct host-pathogen interactions. Cell Rep*, **30,** 335-350 e4.

Pisu, D., Huang, L., Narang, V., Theriault, M., Le-Bury, G., Lee, B., et al. 2021. Single cell analysis of *M. tuberculosis* phenotype and macrophage lineages in the infected lung. J Exp Med*, **218,** e20210615

Podinovskaia, M., Lee, W., Caldwell, S. & Russell, D. G. 2013. Infection of macrophages with *Mycobacterium tuberculosis* induces global modifications to phagosomal function. Cell Microbiol*, **15,** 843-859.

Raggers, R. J., Van Helvoort, A., Evers, R. & Van Meer, G. 1999. The human multidrug resistance protein MRP1 translocates sphingolipid analogs across the plasma membrane. J Cell Sci*, **112 (Pt 3),** 415-422.

Russell, D. G., Cardona, P. J., Kim, M. J., Allain, S. & Altare, F. 2009. Foamy macrophages and the
progression of the human tuberculosis granuloma. Nat Immunol, **10,** 943-948.

Savvi, S., Warner, D. F., Kana, B. D., Mckinney, J. D., Mizrahi, V. & Dawes, S. S. 2008. Functional
characterization of a Vitamin B12-dependent methylmalonyl pathway in *Mycobacterium*
tuberculosis: implications for propionate metabolism during growth on fatty acids. J Bacteriol,
190, 3886-3895.

Schnappinger, D., Ehrt, S., Voskuil, M. I., Liu, Y., Mangan, J. A., Monahan, I. M., et al. 2003.
Transcriptional adaptation of *Mycobacterium tuberculosis* within Macrophages: insights into the
phagosomal environment. J Exp Med, **198,** 693-704.

Shi, L., Salamon, H., Eugenin, E. A., Pine, R., Cooper, A. & Gennaro, M. L. 2015. Infection with
Mycobacterium tuberculosis induces the Warburg effect in mouse lungs. Sci Rep, **5,** 18176.

Simwela, N. V., Johnston, L., Bitar, P. P., Eleni, J., Altier, C., Sassetti, C. M. & Russell, D. G. 2024.
Genome-wide screen of *Mycobacterium tuberculosis* infected macrophages identified the
GID/CTLH complex as a determinant of intracellular bacterial growth. bioRxiv,
2024.05.06.592714.

Singh, V., Jamwal, S., Jain, R., Verma, P., Gokhale, R. & Rao, K. V. 2012. *Mycobacterium*
tuberculosis-driven targeted recalibration of macrophage lipid homeostasis promotes the foamy
phenotype. Cell Host Microbe, **12,** 669-81.

Singhal, A., Jie, L., Kumar, P., Hong, G. S., Leow, M. K., Paleja, B., et al. 2014. Metformin as adjunct
antituberculosis therapy. Sci Transl Med, **6,** 263ra159.

Theriault, M. E., Pisu, D., Wilburn, K. M., Lê-Bury, G., Macnamara, C. W., Michael Petrassi, H., et
al. 2022. Iron limitation in *M. tuberculosis* has broad impact on central carbon metabolism.
Commun Biol, **5,** 685.

Vogel, A., Brunner, J. S., Hajto, A., Sharif, O. & Schabbauer, G. 2022. Lipid scavenging
macrophages and inflammation. Biochim Biophys Acta Mol Cell Biol Lipids, **1867,** 159066.

Warner, D. F., Evans, J. C. & Mizrahi, V. 2014. Nucleotide metabolism and DNA replication.
Microbiol Spectr*, **2***, MGM2-0001-2013

WHO, 2023. Global tuberculosis report 2023. Geneva, Switzerland World Health Organisation.

Wilburn, K. M., Fieweger, R. A. & Vanderven, B. C. 2018. Cholesterol and fatty acids grease the
wheels of *Mycobacterium tuberculosis* pathogenesis. Pathog Dis*, **76***, fty021.

Wolf, A. J., Linas, B., Trevejo-Nuñez, G. J., Kincaid, E., Tamura, T., Takatsu, K. & Ernst, J. D. 2007.
Mycobacterium tuberculosis infects dendritic cells with high frequency and impairs their function
in vivo. J Immunol*, **179,*** 2509-2519.

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., et al. 2021. clusterProfiler 4.0: A universal
enrichment tool for interpreting omics data. Innovation (Camb)*, **2,*** 100141.

Zahrt, T. C., Song, J., Siple, J. & Deretic, V. 2001. Mycobacterial FurA is a negative regulator of
catalase-peroxidase gene katG. Mol Microbiol*, **39,*** 1174-1185.

Figure legends

Figure 1. Knockout of fatty acid import and metabolism genes restricts *Mtb* growth in macrophages. (A) Schematic of lipid import and metabolism genes in macrophages. Genes targeted for CRISPR-Cas9 mediated knockout are highlighted in red. (B-C) Scramble or indicated mutant macrophages were infected with the *Mtb* Erdman strain at MOI 0.4. CFUs were plated 3 hours post infection (B) and on Day 5 (C) to determine intracellular *Mtb* replication rates. n=6; ****, $P < 0.0001$.

Figure 2. *Mtb* infected macrophages with impaired fatty acid import and metabolism display reduced mitochondrial activities and are more glycolytic. (A) Seahorse flux analyses of scramble or PLIN2$^{-/-}$, FATP1$^{-/-}$ and CPT2$^{-/-}$ macrophages infected with *Mtb* Erdman strain at MOI 1 24 hours post infection. Oxygen consumption rates (OCR) were measured using the Cell Mito Stress Test Kit (Agilent). Oligo, oligomycin; FCCP, fluoro-carbonyl cyanide phenylhydrazone; Rot/A, rotenone and antimycin A. (B) Comparison of basal respiration and spare respiratory capacity (SRC) from A. SRC was calculated by subtracting the normalized maximal OCR from basal OCR. n=3 (2 technical repeats per replicate); ****, $P < 0.0001$. (C) Extracellular acidification rates (ECARs) of scramble or PLIN2$^{-/-}$, FATP1$^{-/-}$ and CPT2$^{-/-}$ macrophages infected with *Mtb* as in A. ECARs were measured using the Agilent Seahorse Glycolysis Stress Test kit. 2DG, 2-Deoxy-D-glucose. (D) Comparison of basal glycolysis and spare glycolytic capacity (SGC) in the indicated mutant macrophages. SGC was calculated as SRC above. n=3 (2 technical repeats per replicate); *, $P < 0.05$, ****, $P < 0.0001$.

Figure 3. Knockout of fatty acid import and metabolism genes in macrophages activate AMPK and stabilizes HIF-1α. (A) Western blot analysis of HIF-1α in uninfected and *Mtb* infected scramble or PLIN2$^{-/-}$, FATP1$^{-/-}$ and CPT2$^{-/-}$ macrophages. In *Mtb* infected conditions, cells were infected with the bacteria at MOI 1 for 48 hours before preparation of cell lysates. (B) Western blot analysis of total and phosphorylated AMPK in uninfected and *Mtb* infected scramble or PLIN2$^{-/-}$, FATP1$^{-/-}$ and CPT2$^{-/-}$ *macrophages*. Cell lysates were prepared as in A.

Figure 4. Supplementation with exogenous oleate fails to rescue the *Mtb* Δicl1 mutant in PLIN2$^{-/-}$, FATP1$^{-/-}$ and CPT2$^{-/-}$ macrophages. (A) Scramble or indicated mutant macrophages were infected with the *Mtb* H37Rv Δicl1 mutant expressing mCherry at MOI 5. Oleate supplementation (400 μM) was commenced 24 hours before infection in the treatment group, removed during *Mtb* infection and re-added 3-hours post infection for the entire duration of the experiment. Growth kinetics of *Mtb* were measured by monitoring mCherry fluorescence using a plate reader. n=4; ****, *P* < 0.0001. (B) Uninfected scramble or FATP1$^{-/-}$ and CPT2$^{-/-}$ macrophages were supplemented with 400 μM oleate for 24 hours. Cells were then fixed for 20 minutes and stained for lipid droplet inclusions using the Bodipy 493/503 dye. DAPI was used as a counterstain to detect nuclei.

Figure 5. Dual RNA sequencing to identify host and bacterial determinants of *Mtb* restriction in macrophages with fatty acid import and metabolism knockout genes. (A) Principal component analysis (PCA) of scramble or PLIN2$^{-/-}$, FATP1$^{-/-}$ and CPT2$^{-/-}$ macrophages transcriptomes infected with the *Mtb* smyc'::mCherry strain at MOI 0.5 4 days post infection. (B) Venn diagram of DE gene sets (Supplementary file 2) in PLIN2$^{-/-}$, FATP1$^{-/-}$ and CPT2$^{-/-}$ mutant macrophages as compared to scramble showing overlapping genes. DE genes cutoff; abs (log$_2$ fold change) > 0.3, adjusted p-value < 0.05. (C-D) Tree plot of top 80 enriched gene ontology terms (biological process) in *Mtb* infected PLIN2$^{-/-}$ and FATP1$^{-/-}$ upregulated genes.

Figure 6. Nutritional and oxidative stress define the core transcriptome response of *Mtb* inside PLIN2$^{-/-}$ macrophages. (A) Schematic showing DE genes in *Mtb* transcriptomes isolated from PLIN2$^{-/-}$, FATP1$^{-/-}$ and CPT2$^{-/-}$ macrophages in Figure 5A. (B) Heatmaps of nutritional and oxidative stress DE genes in PLIN2$^{-/-}$ macrophages. Arrows show genes which are also DE in CPT2$^{-/-}$ macrophages in a similar trend (Supplementary file 4).

Figure 7. Inhibitors of fatty acid transport and metabolism block intracellular growth of *Mtb* in macrophages. (A) Growth of *Mtb* in liquid broth in the absence of drug (DMSO) or presence of metformin, FATP1 inhibitor (FATP1 In, 10 μM) and the β-oxidation of fatty acid inhibitor,

Trimetazidine (TMZ, 500 nM). *Mtb* Erdman was grown to log phase and diluted to OD_{600} 0.01 in

7H9 media in the presence of the above inhibitors. Growth kinetics were monitored by OD_{600}

measurements using a plate reader. Rifampicin (RIF) at 0.5 μg/ml was used as a total killing

control. (B) Scramble macrophages were infected with *Mtb* Erdman at MOI 0.5. Inhibitors were

added 3 hours post infection following which CFUs were plated 4 days post infection. n=5; **, P

< 0.01; ****, P < 0.0001.

A

B

Scramble

FATP1⁻ᐟ⁻ g4

CPT2⁻ᐟ⁻ g2

BODIPY 493/503:DAPI

PLIN2⁻ᐟ⁻ up

FATP1⁻ᐟ⁻ up

PLIN2⁻ᐟ⁻ macrophages intracellular *Mtb* DE genes: up 69, down 36

FATP1⁻ᐟ⁻ macrophages intracellular *Mtb* DE genes: up 0, down 0

CPT2⁻ᐟ⁻ macrophages intracellular *Mtb* DE genes: up 3, down 7