Cryptosporidium oocysts and *Giardia* cysts at two South African wastewater

treatment plants

Thandubuhle Gonose[1, 2]*; Hein Du Preez[1]; Fanus Venter[2]; and Monique Grundlingh[1]

[1]Rand Water Analytical Services, PO Box 3526, Vereeniging, South Africa

*E-mail: spuraza@yahoo.com

[2]University of Pretoria, Department of Microbiology and Plant Pathology, Pretoria

ABSTRACT

Cryptosporidium and *Giardia* are one of the most prominent parasitic protozoans found in wastewater treated by wastewater treatment plants (WWTPs). They cause Cryptosporidiosis and Giardiasis respectively, which can be fatal in immuno-compromised individuals. Thus, their removal by WWTPs is essential. This study aimed at investigating the prevalence of *Crytposporium* and *Giardia* through treatment processes at two South African wastewater treatment plants and assessing the efficiency of the WWTPs in removing them. A total of 162 samples (54 raw influents, 36 primary treated effluents, 36 secondary treated effluents and 36 final treated effluents) were collected from two South African WWTPs over a period of one year. Results obtained from the study show that *Cryptosporidium* was detected in 98% (159/162) of the samples collected, whilst Giardia was detected in 99% (160/162) of the samples collected from both wastewater treatment plants. Both *Crytposporium* and *Giardia* were detected in 100% (n=54) of raw influent samples. The same result was obtained in primary treated effluent, where both *Cryptosporidium* and *Giardia* was detected in 100% (n=36) of the samples collected. In secondary treated effluent, 36 samples were collected and *Cryptosporidium* was detected in 94% of the samples, whereas *Giardia* was detected in 100% of the samples. In the final treated effluent samples, both *Cryptosporidium* and *Giardia* where detected 97% (n=36) of the samples collected. Detection of *Cryptosporidium* and *Giardia* in most of the wastewater treatment stages indicated these parasites are prevalent in the two WWTPs. Although the two WWTPs showed high removal of the two parasites in wastewater, their presence in the final treated effluent indicated the inefficiency of treatment processes in their removal. This highlights the potential of these WWTPs in contaminating receiving water sources.

Keywords: Prevalence, wastewater treatment, *Giardia*, *Cryptosporidium*, wastewater treatment plants

IMPORTANCE OF THE STUDY

This study will specifically evaluate the removal of Cryptosporidium oocysts and Giardia cysts by two South African WWTPs and will estimate their ability of removing them. The study will also attempt to give insight into the potential and extend of treated effluent in polluting surface water resources with respect to Cryptosporidium and Giardia. Identification of the potential sources of the pollution is one of the significant parts in resolving this problem. Hence the finding of the study are very important in an endeavour to eliminate pollution of water source.

1. INTRODUCTION

Cryptosporidium and *Giardia* are protozoan parasites that infect the gastrointestinal tract of a broad range of vertebrate animals such as birds, reptiles, mammals, amphibians and fishes (Putignani and Menichella, 2010). In humans, *Cryptosporidium* and *Giardia* infect the distal and proximal regions of the small intestines and multiply within the hosts (Bukhari and Clancy, 2000). The infection result into a disease called Cryptosporidiosis (*Cryptosporidium*) and Giardiasis (*Giardia*) which is characterised by watery diarrhoea that could be accompanied by stomach cramps, nausea and fever (Health Canada, 2012). Infected humans excrete large amounts of *Cryptosporidium* and *Giardia*, in transmissive stages called oocysts and cysts respectively, into the environment via their feacal excretion (Putignani and Menichella, 2010).

Feacal excretion of large numbers of *Cryptosporidium* oocysts and *Giardi*a cysts by infected hosts makes these parasites some of the most frequently isolated parasites from wastewater (also referred to as sewage) (Fayer et al. 2000). Furthermore, *Cryptosporidium* oocysts and the *Giardia* cysts excreted by the infected hosts are very robust and can remain viable under different environmental conditions for long periods. They are also able to resist oxidizing disinfectants typically used by water suppliers to remove pathogens (Carmena, 2010). Because of these features, *Cryptosporidium* and *Giardia* are amongst the most important pathogens in the water industry.

Several studies have isolated both *Cryptosporidium* and *Giardia* from raw and treated wastewater at wastewater treatment plants in different parts of the world (Sroka et al. 2013; Castro-Hermida et al. 2010; Cheng et al. 2009; Dungeni and Momba, 2010). Isolation of the two parasites in raw wastewater could be an indication of the infection of the population by *Cryptosporidium* and *Giardia* (Robertson et al. 2000). Although high counts of *Cryptosporidium* and *Giardia* in raw wastewater is important, their detection through the treatment process and in the final treated wastewater that is discharged into water sources is of serious concern as it highlights the risk to public health. This is because the treated wastewater is discharged into surface water sources such as dams and rivers which are sometimes used by consumers for different purposes such as drinking, recreation and agriculture (Teklehaimanot et al. 2015).

Frequent detection of *Cryptosporidium* oocysts and *Giardia* cysts in the final treated effluent has shown that a number of WWTPs are not equipped to effectively remove *Cryptosporidium* oocysts and *Giardia* cysts from wastewater (Cheng et al. 2009). Various studies have

confirmed this as both *Cryptosporidium* oocysts and *Giardia* cysts are regularly detected in the

treated wastewater effluent that is discharge by WWTPs into water sources (Cheng et al. 2009;

McCuin and Clancy, 2006; Dungeni and Momba, 2010; Robertson et al. 2006).

Factors that contribute to the inefficiency of WWTPs in removing *Cryptosporidium* oocysts

and *Giardia* cysts include poor management of WWTPs, the use of ineffective treatment

processes by the WWTPs and uncontrolled wastewater inputs (Nikiema et al. 2013). These

factors together with financial challenges play a major role in the failure of WWTPs to produce

effluent that is free of pathogens, particularly in developing countries. South Africa (SA), as a

developing country, is also facing the problem of failure of WWTPs to produce effluent of high

quality (Dungeni and Momba, 2010; Naidoo and Olaniran, 2014; Sroka et al. 2013).   In actual

fact, in South Africa WWTPs have been identified as one of the major polluters of water

sources (Dungeni and Momba, 2010; CSIR, 2010).

To date, there are relatively few studies in South Africa that have looked at the prevalence and

efficiency of wastewater treatment plants in removing *Cryptosporidium* oocysts and *Giardia*

cysts. Thus this current study aimed at investigating the prevalence of *Cryptosporidium* oocysts

and *Giardia* cysts in raw influent, through treatment stages, and in final treated effluent of two

South African WWTPs over a period of 12 months. Furthermore, the removal efficiency of

Cryptosporidium oocysts and *Giardia* cysts by the two WWTPs was evaluated. Results

obtained in this study will be useful to water utilities as it can assist in the evaluation of WWTPs

as polluters of the receiving water bodies and also in establishing *Cryptosporidium* and *Giardia*

risk status in water sources (Sigudu et al. 2014).

2. MATERIALS AND METHODS

Study sites

Two wastewater treatment plants (WWTPs) were selected as sampling sites for this study.

These wastewater treatment plants were referred to as WWTP 1 and WWTP 2 and they are

located in two South African Provinces namely Gauteng and the Free State. Table 1

summarises the characteristics of the two WWTPs.

Table 1: Characteristics of the sampling sites

WWTP	Wastewater source	WWTP capacity (ML/day)	Treatment processes
WWTP 1	Domestic, Industries Abattoir	36	Coarse and fine steel filters, Grit remover, Settling tanks, Biological nutrient removal; Settling tanks, Chlorine dosage
WWTP 2	Domestic	2.1	Course steel filters, Sedimentation (Settling ponds),Trickling Filters; Sedimentation; Chlorine dosage,

140 *Note: WWTP1- has two raw influent receiving points. One that receives wastewater from industries and abattoirs*

141 *and other that receives from domestic.*

Sample collection for *Cryptosporidium* oocysts and *Giardia* cysts analysis

Grab wastewater samples were collected at different treatment stages, namely: raw influent,

primary treated effluent, secondary treated effluent and the final treated effluent over a period

of 12 months (February 2014 to January 2015) from WWTP1 and WWTP2. For the first six

months of the study, samples were taken twice a month (second and fourth week of each

month), and in the last six months sampling was done once every month. Wastewater samples

were collected using 500 mL bottles for *E. coli* analysis, 1 L sampling bottles for

Cryptosporidium and *Giardia* analysis in raw influent and 10 L sampling bottles

Cryptosporidium and *Giardia* analysis in primary effluent, secondary effluent and final treated

effluent samples.

Sample analysis

153 The United States Environmental Protection Agency method 1693 was used for detecting and

154 enumerating *Cryptosporidium* oocysts and *Giardia* cysts in the collected wastewater samples

155 (USEPA, 2014).

Sample concentration

Treated effluent samples (10 L) were filtered using Pall Envirochek^{TM} High Volume (HV filter)

filters at a flow rate of 2 L min^{-1}. Laureth 12 elution buffer and wrist actions shakers were used

to elute the filters. The eluates were transferred into 250 mL conical tubes. centrifuged using a

Sorvall RC3 centrifuge at 1500 xG for 15 minutes. After centrifugation, the pellet was used to

detect *Cryptosporidium* oocysts and *Giardia* cysts using the immuno-magnetic separation.

Raw influent samples were directly poured into the 250 mL conical tubes and centrifuged.

Immunomagnetic separation

The IMS procedure was performed as described in USEPA 1693 (USEPA, 2014. Briefly,

kaolin (0.75 g) was added into the LT tubes containing SL-buffer-A and B and the pellet, for

raw water samples. One hundred microliters of anti-*Cryptosporidium* and 100 µL of anti-

Giardia Dynabeads were added to the LT tubes and mixed by rotating using at 18 rpm for 1

hour. The LT tubes were placed in a magnetic particle concentrator (MPC) and gently mixed

by tilting at 90˚ angle for 2 minutes. The supernatant was decanted and the LT tubes were

removed from the MPC. One millilitre of 1X buffer A was added to the LT tubes to collect the

beads. The suspension containing bead-oocysts/cysts complex was transferred into 1.5 mL

Eppendorf tubes. These tubes were placed in MPC and rocked for 1 minute. The supernatant

was gently removed through aspiration. The magnets were removed, 50 µL of 0.1 N

hydrochloric acid (HCl) was added to the tubes and the tube was vortexed for 20 seconds. The

tubes were allowed to stand for 10 minutes, the vortexed for 20 seconds and the magnets were

inserted. The tubes were allowed to stand undisturbed for few seconds. The beads collected at

the back of the tubes and the acidified suspension was transferred into microscope slides

containing 10 µL of 1.0 N of Sodium hydroxide (NaOH). This procedure was repeated twice

for each tube and transferred into the same microscope slide.

Staining procedure (***Immunofluorescence assay***)

The slides were dried up and fixed using methanol. Twenty five microliters of fluorescein

isothiocyanate anti-*Cryptosporidium* and 25 µL of fluorescein isothiocyanate of anti-*Giardia*

monoclonal antibodies (FITC-mAb) stains were added onto the slides and incubated at 37 ±

2˚C for 45 minutes. The excess FITC-Mab was aspirated. Fifty microliters of 4'6-diamidino-

2-phenyindole (DAPI) (50 µL) was added onto the slides and allowed to stand at room

temperature for 2 min then the excess was aspirated. Fifty microliters of PBS was added onto

the slides and allowed to stand at room temperature for 2 min then the excess was aspirated.

The slides were allowed to dry. Diazabicyclooctane (DABCO) mounting medium (8 µL) was

used to mount the slides and they were sealed. An Olympus epiflourescence microscope

containing the blue filter block was used to examine the stained slides, and to detect and

enumerate FITC-mAb labelled *Cryptosporidium* oocysts and *Giardia* cysts at 40x

magnification. DAPI stain characteristics were examined at 40x magnification with the UV

filter.

3. RESULTS

3.1 Prevalence and concentrations of *Cryptosporidium oocysts* and *Giardia cysts* in raw influents and final treated effluents

Table 2 provide the results of the prevalence of *Cryptosporidium* oocysts and *Giardia* cysts through the treatment process in WWTP1. A total of 90 samples (36 raw influents, 18 primary treated effluents, 18 secondary treated effluents and 18 final treated effluents) were collected and analysed in WWTP1 over a period of one year. According to these results both *Cryptosporidium* and *Giardia* were detected in 99% (89/90) of the samples collected from WWTP1. As shown in Table 2, *Cryptosporidium* and *Giardia* were detected in 100% (n=36) of raw influent samples. The same result was obtained in primary treated effluent, where both *Cryptosporidium* and *Giardia* was detected in 100% (n=18) of the samples collected. In secondary treated effluent, a total of 18 samples were collected and *Cryptosporidium* was detected in 100% (n=18) of the samples, whilst *Giardia* was detected in 94% (n=18) of the samples. Visa versa results were obtained in the final treated effluent samples, where *Cryptosporidium* was detected in 94% of the samples, whereas *Giardia* was detected in 100% of the samples.

Also shown in table 2 are the results of the prevalence of *Cryptosporidium* oocysts and *Giardia* cysts through the treatment process in WWTP2. A total of 72 samples (18 raw influents, 18 primary treated effluents, 18 secondary treated effluents and 18 final treated effluents) were collected and analysed in WWTP2 over a period of one year. According to these results *Cryptosporidium* was detected in 97% (70/90) of the samples collected, whilst *Giardia* was detected in 99% (71/72) of the samples collected from WWTP1. As illustrated by Table 2, *Crytposporium* and *Giardia* were detected in 100% (n=18) of raw influent samples. The same result was obtained in primary treated effluent, where both *Cryptosporidium* and *Giardia* was detected in 100% (n=18) of the samples collected. In secondary treated effluent, a total of 18 samples were collected and *Cryptosporidium* was detected in 94% (n=18) of the sample, whereas *Giardia* was detected in 100% (n=18) of the samples. In the final treated effluent samples. Both *Cryptosporidium* and *Giardia* where detected 94% (n=18) of the samples collected.

Figure 1 and figure 2 illustrates the mean concentration of both *Cryptosporidium* oocysts and *Giardia* cysts in different treatment stages over 12 months period of the study. According to the results, mean concentration of both *Cryptosporidium* oocysts and *Giardia* cysts were

227   considerable higher in the raw influent treatment stage. Reduction in mean concentration for

228   both *Cryptosporidium* and *Giardia* was observed through the treatment stages with the lowest

229   values obtained in the final treated effluent. WWTP2 had the highest mean concentration for

230   *Crytposporium* (9134 oocysts/ 10 L), whilst for WWTP2 high mean concentration for *Giardia*

231   (13572 cysts/ 10 L).

241 **Table 2:** Prevalence; mean, max (maximum), min (minimum) and the percentage number of samples positive for *Cryptosporidium* and *Giardia*

242 in raw influent and the final treated effluent from the two WWTPs.

Figure 1: Mean Concentrations of *Cryptosporidium* and *Giardia* in raw influent, primary treated effluent, secondary treated effluent and the final treated effluent at WWPT1 (A) and WWTP2 (B).

Table 3: Mean % removal efficiency of *Cryptosporidium* and *Giardia* in WWTP 1 and WTTP 2

4. DISCUSSION

This study aimed at determining the prevalence of *Cryptosporidium* and *Giardia* in two South African WWTPs. Using method 1693, *Cryptosporidium* oocysts and *Giardia* cysts were isolated, identified and enumerated in different treatment stages (raw influent, primary treated effluent, secondary treated effluent and final treated effluent) in both WWTPs. These results suggest that *Cryptosporidium* and *Giardia* are one of the predominant pollutants in wastewater treated by the two WWTPs. Moreover, detection of *Cryptosporidium* oocysts and *Giardia* cysts in the final treated effluent indicated that the two WWTPs were inefficient in removing either *Cryptosporidium* oocysts or *Giardia* cysts in the wastewater.

Both *Cryptosporidium* and *Giardia* were detected consistently in all treatment stages (raw influent, primary effluent, secondary effluent and the final treated effluent) at both WWTPs. In raw influent, results obtained revealed that both *Cryptosporidium* oocysts and *Giardia* cysts were consistently detected in all raw influents samples (100% samples) analysed throughout the year. These results indicate a high prevalence of *Cryptosporidium* and *Giardia* in the wastewater treated by the two wastewater treatment plants. The occurrence of *Cryptosporidium* and *Giardia* obtained in this study was higher than that reported by Dungeni and Momba (2010), where *Cryptosporidium* oocysts and *Giardia* cysts were isolated in 50% and 78% of raw influent samples. Robertson and co-workers (2006) also detected *Cryptosporidium* and *Giardia* in 93% and 80% of raw wastewater samples. Also, it is likely that percentages of *Cryptosporidium* and *Giardia* reported in several studies are an underestimation, due to the challenge of low recovery efficiency of the method used. In a study carried out in north-eastern Spain, *Cryptosporidium* was detected in 100% of raw influent wastewater samples (Montemayor et al. 2005). Cheng and co-workers (2009) also obtained both *Cryptosporidium* and *Giardia* in 100% of the samples analysed.

Concentrations of *Cryptosporidium* oocysts and *Giardia* cysts varied throughout the study
period with highest concentrations of 60120 oocysts/10L and the lowest concentration of 32
cysts/10L. According to McCuin and Clancy (2006), high variations in concentrations oocysts
could be due to several factors such as the 83 levels of infections in the contributing
community, the severity of the infection in the community and the size of the community. In
the current study industries and abattoirs were also contributors of wastewater. Therefore,
levels of *Cryptosporidium* oocysts and *Giardia* cysts in the wastewater discharged by the
domestic, industries and abattoirs contributed to the concentration variations. Similar trends
where *Cryptosporidium* and *Giardia* concentrations varied between sampling occasions were
obtained in a study by Robertson and co-workers (2000). According to Robertson and co-
workers (2000), concentrations of *Cryptosporidium* and *Giardia* detected in raw influent are
determined by the contributors such as the infected animals and humans in the community
served by the wastewater treatment plants (Robertson et al. 2000). The mean concentration of
Giardia (13572 cysts/10L) was higher than *Cryptosporidium* concentration in WWTP 1
(P=0.01). This phenomenon could indicate that there is a lower prevalence of *Cryptosporidium*
in the wastewater contributors (abattoirs, human and industries). On the other hand, results on
concentrations of these two protozoans should be interpreted with caution, considering the fact
that the method used in this study had low recoveries in the raw influent matrix. Similar results
were obtained in a study by Castro-Hermida and co-workers (2010), where mean concentration
of *Giardia* was higher than mean concentration of *Cryptosporidium* (Castro-Hermida et al
2010). In WWTP 2, the mean concentration of *Cryptosporidium* (9134 oocysts/10L) was
higher than *Giardia* concentration. WWTP 2 treated human associated waste mostly.
Therefore, the high mean concentration of *Cryptosporidium* could indicate that
Cryptosporidium infections were common in the communities served by the WWTP.

Throughout the treatment process, both *Cryptosporidium* and *Giardia* were detected through
the period of the study. According to the results, mean concentration of both *Cryptosporidium*
oocysts and *Giardia* cysts were considerable higher in the raw influent treatment stage.
Reduction in mean concentration for both *Cryptosporidium* and *Giardia* was observed through
the treatment stages with the lowest values obtained in the final treated effluent. This trend in
concentrations of both protozoans was also observed in study by Cheng and co-workers (2009),
where high concentrations of *Cryptosporidium* and *Giardia* were observed in raw influents and
low concentrations were obtained in the final treated effluent. Mean removal efficiency ranged
from 96.5 to 97.7% for both *Cryptosporidium* oocysts and *Giardia* cysts in both WWTPs.

According to Robertson et al (2000), several factors play a role in the removal of parasites by
WWTPs. These factors include parasite properties and the properties of the WWTP.

Both WWTPs had high percentage removal efficiency. However, the fact that *Cryptosporidium*
and *Giardia* were detected in the final treated effluents discharged by the WWTPs indicated
the risk of contamination of the receiving water bodies. This phenomenon is of major concern
if the receiving water bodies are being used for domestic purposes by communities living
downstream of the WWTPs (Lim et al. 2007). The percentage removal results obtained in the
study were similar to those obtained by Dungeni and Momba (2010) and Montemayor et al
(2005). Parker et al (1993) obtained a percentage removal efficiency of 80-98% for
Cryptosporidium oocysts and 98-99% for *Giardia* cysts. Dungeni and Momba (2010) obtained
a similar percentage removal efficiency ranging from 67.4% to 98.26% and from 86.81 to
99.96% for *Cryptosporidium* oocysts and *Giardia* cysts respectively.

Results obtained revealed the presence of *Cryptosporidium* oocysts and *Giardia* cysts in final
treated wastewater effluent in both WWTPs. In WWTP 1, *Cryptosporidium* oocysts were
isolated in most of the treated effluent samples (94%, 0 - 454 *Cryptosporidium* oocysts/10L
and 100%, 1 - 1032 *Giardia* cysts/10L). *Cryptosporidium* and *Giardia* were also detected in
most treated effluent samples collected from WWTP 2 (94%, 0-1955 *Cryptosporidium*
oocysts/10L and 94%, 0-1214 *Giardia* cysts/10L). A significant difference between
concentrations of *Cryptosporidium* and *Giardia* were observed in WWTP 1 (P = 0.02).
However, for WWTP 2, there were no significant difference observed between
Cryptosporidium and *Giardia*. Moreover, *Cryptosporidium* oocysts and *Giardia* cysts
concentrations reported might be an underestimation, since the recovery efficiency data in (not
given in this paper) were low and the method was inhibited by unknown matrix components.
Nevertheless, detection of *Cryptosporidium* oocysts and *Giardia* cysts in treated effluents from
the two WWTPs suggested that the treatment processes used in these wastewater treatment
plants were not effective in removing these two protozoans. The prevalence of
Cryptosporidium and *Giardia* reported in the study were comparable to the results obtained by
Montemayor and co-workers (2005), where *Cryptosporidium* oocysts and *Giardia* cysts were
detected in 100% of wastewater treated effluent from two WWTPs (Montemayor et al. 2005).
The prevalence and concentrations of *Cryptosporidium* and *Giardia* measured in this study
were higher than those reported in a study conducted by Dungeni and Momba (2010) in
selected South African WWTPs. This could due to differences in the prevalence of

Cryptosporidium and *Giardia* in the wastewater contributors in different areas and also the
differences in the methodology.

5. CONCLUSION

The findings of this study revealed that *Cryptosporidium* and *Giardia* were prevalent
throughout the treatment process and in the final treated effluent from the two WWTPs. This
is an indication that these two WWTPs could pollute the receiving water sources in terms of
Cryptosporidium and *Giardia*. Pollution of surface water with *Cryptosporidium* and *Giardia*
parasites could have serious public health risks when the water is directly used for drinking
purposes, irrigation and for recreational purposes. Hence, this study identified a need for
Cryptosporidium and *Giardia* to be included in the monitoring programmes for treated
wastewater effluent. Also, to ensure that the limited South African surface water resources are
protected from being polluted by WWTPs, enhancement of the performance of South African
WWTPs to remove these protozoan parasites needs serious attention. This process of
enhancement will start with the process of determining the main challenges that are hindering
the performance of the WWTPs.

5. REFERENCES

1.   Abdel-Raouf, N., Al-Homaidan, A. A. and Ibraheem, I. B. M. (2012) Microalgae and
wastewater treatment. Saudi Journal of Biological Sciences Volume 19 pp. 257-275.

2.   APHA (American Public Health Association) (1995) Standard methods for
examination of water and wastewater. American Water Works, Water Environment Federation.

3.   Bukhari, Z. and Clancy, J. (2000) Development of method for Cryptosporidium
analysis in wastewater. Presented at WISA Biennial Conference, Sun City, South Africa.

4.   Carmena, D., Aguinagalde, X., Zigorraga, C., Fernandez-Crespo, J.C. and Ocio, J. A
(2007) Presence of Giardia cysts and Cryptosporidium oocysts in drinking water supplies in
northern Spain. Journal of Applied Microbiology Volume 102 pp.619-629.

5.   Carmena, D. (2010) Waterborne transmission of Cryptosporidium and Giardia:
detection, surveillance and implications for public health. Current Research, Technology, and

Education Topics in Applied Microbiology and Microbial Technology. A. Mendez-Vilas
Edition.

6.   Castro-Hermida, J. A., Garcia-Presedo, J., Gonzalez-Warleta, M. and Mezo, M. (2010)
Cryptosporidium and Giardia detection in water bodies of Galicia, Spain. Water Research
Volume 44 pp. 5887-5896

7.   Cheng, H. A., Lucy, F. E., Graczyk, T. K., Broaders, M. A., Taman., L. and Connolly,
M. (2009) Fate of Cryptosporidium parvum and Cryptosporidium hormonis oocysts and
Giardia duodenalis cysts during secondary wastewater treatments. Parasitology Research. DOI
10. 1007/s00436-009-1440-y.

8.   Council for Scientific and Industrial Research (CSIR) (2010) A CSIR perspective on
water in South Africa-2010. CSIR Report No- CSIR/NRE/PW/IR/2011/0012/A. ISBN: 978-0-
79885595-2.

9.   Dungeni, M. and Momba M., N., B. (2010) The abundance of Cryptosporidium and
Giardia in treated effluent produced by four wastewater treatment plants in the Gauteng
province of South Africa. Water South Africa Volume 36 Number 4 pp. 425-432.

10.   DWAF (Department of Water Affairs and Forestry) (2013) Revision of general
authorisation in terms of section 39 of NWA, 1998. Government Gazette, Act No. 36 of 1998.

11.   Fayer, R., Morgan, U. and Upton, S., J. (2000) Epidemiology of Cryptosporidium:
transmission, detection and identification. International Journal of Parasitology Volume 30 pp.
1305-1322.

12.   Health Canada (2012). Guidelines for Canadian Drinking Water Quality: Guideline
Technical Document — Enteric Protozoa: Giardia and Cryptosporidium. Water, Air and
Climate Change Bureau, Healthy Environments and Consumer Safety Branch, Health Canada,
Ottawa, Ontario. (Catalogue No H129-23/2013E-PDF)

13.   Kfir, R., Hilner, C. du Preez, M. and Bateman, B. (1995) Studies in the prevalence of
Giardia cysts and Cryptosporidium oocysts in South Africa water. Water Science &
Technology 31 (5).435-438.

14.   Kuhn, R. C., Rock, C. M. and Oshima, K. H. (2002) Effects of pH and immuno-
magnetic separation of Cryptosporidium oocysts from concentrated water samples. Applied
and Environmental Microbiology Volume 68 Number 4 pp. 2066-2070.

15.   Lim, Y. A. L., Wan Hafiz, W. I. and Nissaparton, V. (2007) Reduction of
Cryptosporidium and Giardia by sewage treatment processes. Tropical Biomedicine Volume
24 Number 1 pp. 95 -104.

16.   McCuin, R., M. and Clancy, J., L. (2006) Occurrence of Cryptosporidium oocysts in
US wastewaters. Journal of Water and Health Volume 4 Number 4 pp. 437-451.

17.   Montemayor, M., Valero, F., Jofre, J. and Lucena, F. (2005) Occurrence of
Cryptosporidium spp. oocysts in raw and treated sewage and river water in north-eastern Spain.
Journal of Applied Microbiology Volume 99 pp. 1455-1462.

18.   Naidoo, S. and Olaniran, A., O. (2014) Treated wastewater effluent as a microbial
pollution of surface water sources. International Journal of Environmental Research and Public
Health Volume11 pp. 249-270.

19.   Nikiema, J., Figoli, A., Weissenbancher, N., Langergraber, G., Marrot, B. and Moulin
P. (2013) Wastewater treatment practises in Africa-experiences from seven countries.
Sustainable Sanitation Practice Issue14 pp. 26-34.

20.   Parker, J. F. W., Smith, H. V. and Girdwood, R. W. A. (1993) Survey of Loch Lomond
to assess the occurrence and prevalence of Cryptosporidium sp. Oocysts and their likely on
human health. Foundation for Water Research. Report No. FR0409.

21.   Putignani, L., Menichella, D. (2010) Global distribution, public health and clinical
impact of the protozoan pathogen Cryptosporidium. Interdisciplinary Perspective on Infectious
Diseases Volume 12 pp.1-39.

22.   Quintero-Betancourt, W., Gennaccaro, T. M., Scott, T. M. and Rose, J. B. (2003)
Assessment of the methods for detection of infectious Cryptosporidium oocysts and Giardia
cysts in reclaimed effluents. Applied and Environmental Microbiology Volume 69 pp. 5380-
5388.

23.   Robertson, L. J., Paton, C. A., Campbell, A. T., Smith, P. G., Jackson, M. H., Gilmour,
R. A., Black, S. E., Stevenson, D. A. and Smith, H., V. (2000) Giardia cysts and

Cryptosporidium oocysts at sewage treatment works in Scotland, UK. Water Research Volume
34 Number 8 pp. 2310-2322.

24. Robertson, L. J., Hermansen, L. and Gjerde, B. K. (2006) Occurrence of
Cryptosporidium oocysts and Giardia cysts in sewage in Norway. Applied and Environmental
Microbiology Volume 27 Number 8 pp. 5297-5303.

25. Samie, A. and Ntekele, P. (2013) genotypic detection and evaluation of the removal
efficiency of Giardia duodenalis at municipal wastewater treatment plants in Northern South
Africa. Tropical Medicine Volume 31 Number 1 pp. 122-133.

26. Sigudu, M. V., du Preez, H., H. and Retief, F. (2014) Application of a basic monitoring
strategy for Cryptosporidium and Giardia in drinking water. Water South Africa Volume 40
Number 2 pp. 297-312.

27. Sroka, J., Stojecki, K., Zdybel, J., Karamon, J., Cencek, T. and Dutkiewicz, J. (2013)
Occurrence of Cryptosporidium oocysts and Giardia cysts in effluent from sewage treatment
plant from eastern Poland. Annals of Agriculture and Environmental Medicine Special issue 1
pp. 57-62.

28. Teklehaimanot, G. Z., Genthe, B., Kamika, I. and Momba, M. N. B. (2015) Prevalence
of enteropathogenic bacteria in treated effluents and receiving water bodies and their potential
health risks. Science of the Total Environment pp. 441-449.

29. United States Environmental Protection (USEPA). (2012) Method 1623.1:
Cryptosporidium and Giardia in water by filtration/IMS/FA http://water.epa.gov.drink.

30. Yakub, G. P. and Stadterman-knauer, K. L. (2000) Evaluation of the immunomagnetic
separation for recovery of Cryptosporidium parvum and Giardia duodenalis from high-iron
matrices. Applied and Environmental Microbiology Volume 66 Number 8 pp. 36

Table 2: Prevalence; mean, max (maximum), min (minimum) and the percentage number of samples positive for *Cryptosporidium* and *Giardia*

in raw influent and the final treated effluent from the two WWTPs.

Sample source	Treatment stages	*Cryptosporidium* oocysts/10L					*Giardia* cysts/10L				
		Mean	n	Min	Max	% of positive	Mean	n	Min	Max	% of positive
WWTP 1	Raw influent	4623	36	560	13920	100	13572	36	318	48000	100
	Primary treated effluent	1062	18	76	3720	100	6775	18	106	52800	100
	Secondary treated effluent	150	18	2	448	100	360	18	0	1164	94
	Final treated effluent	116	18	0	454	94	308	18	1	1032	100
WWTP 2	Raw influent	9134	18	40	60120	100	8013	18	32	31200	100
	Primary treated effluent	873	18	9	3150	100	693	18	6	2214	100
	Secondary treated effluent	474	18	0	1578	94	262	18	5	1400	100
	Final treated effluent	323	18	0	1955	94	190	18	0	1214	94

Figure 1: Mean Concentrations of *Cryptosporidium* and *Giardia* in raw influent, primary

treated effluent, secondary treated effluent and the final treated effluent at WWPT1 (A) and

WWTP2 (B).

1 **Table 3:** Mean % removal efficiency of *Cryptosporidium* and *Giardia* in WWTP 1 and WTTP

2 2

WWTP	% Removal efficiency	
	Cryptosporidium	*Giardia*
WWTP 1	97.5	97.7
WWTP 2	96.5	97.6