Transient Non-local Interactions Dominate the Dynamics of Measles

Virus N_{TAIL}

Lillian Otteson[1,2&], Gabor Nagy[3&], John Kunkel[1,2], Gerdenis Kodis[1,2], Lars V. Bock[3],

Christophe Bignon[5], Sonia Longhi[5], Wenwei Zheng[4*], Helmut Grubmüller[3*], Andrea C.

Vaiana[3,6*], and Sara M. Vaiana[1,2*]

[1]Center for Biological Physics, Arizona State University, Tempe, AZ, USA,

[2]Department of Physics, Arizona State University, Tempe, AZ 85287, USA,

[3]Department of Theoretical and Computational Biophysics, Max Planck Institute for

Multidisciplinary Sciences, Göttingen, Germany,

[4]College of Integrative Sciences and Arts, Arizona State University, Mesa, AZ 85212, USA

[5]Aix Marseille Univ, CNRS, AFMB, UMR 7257, Marseille, France

[6]Present address: Department of Chemistry and Chemical Biology, University of New Mexico,

Albuquerque, NM 87131, USA

[&]These authors have contributed equally

*To whom correspondence may be addressed

Abstract.

The RNA genome of measles virus is encapsidated by the nucleoprotein within a helical nucleocapsid that serves as template for both transcription and replication. The intrinsically disordered domain of the nucleoprotein (N_{TAIL}), partly protruding outward from the nucleocapsid, is essential for binding the polymerase complex responsible for viral transcription and replication. As for many IDPs, binding of N_{TAIL} occurs through a short molecular recognition element (MoRE) that folds upon binding, with the majority of N_{TAIL} remaining disordered. Though N_{TAIL} regions far from the MoRE influence the binding affinity, interactions between them and the MoRE have not been investigated in depth. Using an integrated approach, relying on photo-induced electron transfer (PET) experiments between tryptophan and cysteine pairs placed at different positions in the protein under varying salt and pH conditions, combined with simulations and analytical models, we identified transient interactions between two disordered regions distant in sequence, which dominate N_{TAIL} dynamics, and regulate the conformational preferences of both the MoRE and the entire N_{TAIL} domain. Co-evolutionary analysis corroborates our findings and suggests an important functional role for the same intramolecular interactions. We propose mechanisms by which these non-local interactions may regulate binding to the measles phosphoprotein, polymerase recruitment, and ultimately viral transcription and replication. Our findings may be extended to other IDPs, where non-local intra-protein interactions affect the conformational preferences of intermolecular binding sites.

Introduction

The measles virus (MeV) is a non-segmented, single-stranded, negative sense RNA virus, in which the viral genome is wrapped within a helical nucleocapsid, made of thousands of repeats of the nucleoprotein (N) [1]. This structure is shared among all Paramyxo- and Pneumoviruses, [2]

which include common human viruses (e.g., MeV, mumps, respiratory syncytial virus,

metapneumovirus, and other para-influenza viruses) and animal viruses (e.g., Sendai virus,

Newcastle disease virus and rinderpest virus), as well as zoonotic biosafety level 4 agents of

great concern to public health (e.g., Hendra and Nipah viruses, respectively HeV and NiV)[2].

The N protein not only provides the interaction with the RNA and holds the nucleocapsid in its

helical structure, but it is the substrate used for viral transcription and replication. Viral

transcription and replication in Paramyxoviruses are ensured by a complex formed by the large-

polymerase (L) and the phosphoprotein (P), where the latter enables tethering L to the

nucleoprotein-RNA (N-RNA) template for transcription and replication[2,3].

In MeV, N consists of a folded 400 amino-acid (a.a.) domain (N_{CORE}) which binds the RNA

and forms the structured helical nucleocapsid, and of an intrinsically disordered 125 aa. C-

terminal domain (N_{TAIL}), which partly protrudes radially outward from the helical nucleocapsid

structure[4–7]. The intrinsically disordered N_{TAIL} domain binds to the folded X domain (P_{XD}) of

the phosphoprotein (P) and is essential for MeV transcription and replication. P in turn, binds

as a tetramer to both L and N_{CORE}[6,8–11]. A similar structure and binding of N_{TAIL} to P_{XD} are found

in NiV and HeV[3,12]. Binding of N_{TAIL} to P_{XD} is thought to assist in positioning the L

appropriately on the nucleocapsid to read the RNA, while allowing efficient transcription re-

initiation at intergenic regions[13] and hence progression from one reading frame to the next. It

may also assist in maintaining contact between L and the nucleocapsid template, preventing it

from leaving before transcribing all the viral genome[11,14]. In addition, the presence of N_{TAIL}

loosens the nucleocapsid helical structure, possibly facilitating transcription and replication by

L[11].

Because of its crucial role in transcription and replication, binding of MeV N_{TAIL} to P_{XD} has

been extensively studied[15–21]. It occurs through coupled folding and binding of a short

molecular recognition region (MoRE) of 18 a.a. (grey in Fig. 1)[21]. Upon binding, the MoRE

folds into an α-helix, forming a four-helix bundle with three helices from P_{XD} (PDB: 1t6o[8,9]).

However, the majority of N_{TAIL} (i.e., the remaining 109 a.a. on the two sides of the MoRE)

remains disordered in the bound complex[21]. The N_{TAIL}-P_{XD} complex is structurally similar in

MeV, NiV and HeV[3,12]. While the regions outside the MoRE (from here on referred to as

"flanking regions") do not fold, or engage in stable binding interactions[16,17], they modulate

binding affinities, as indicated by the binding affinity increase in truncated variants of N_{TAIL}

containing the intact MoRE[22,23]. This increase might be due to a pure excluded volume

(entropic) repulsion of the flanking regions, as well as to potential interactions involving the

flanking regions of N_{TAIL} (either at the intra- or inter-protein level)[22,24,25].

For N_{TAIL} free in solution, the secondary structure propensity of the MoRE region has been

characterized by NMR on full length N_{TAIL}[7], and by all-atom simulation studies of the MoRE

fragment[19,26]. In contrast, we know very little about potential interactions between residues that

are distant in sequence (non-local interactions), despite their particular importance for N_{TAIL}

binding to P_{XD}[24,27] and, more generally, for IDP function. Large-scale reconfiguration dynamics

can regulate IDP binding mechanisms, rates, and affinities; in fact, examples of this are found

even in simple models of proteins undergoing two-state conformational transitions[28]. For N_{TAIL},

we expect interactions involving the regions flanking the MoRE on either side to confer binding

specificity to the IDP by dictating large-scale reconfiguration dynamics. In addition, non-local

interactions potentially act as allosteric regulators in IDPs by stabilizing certain conformational

states. The resulting shift in the population of states within the ensemble can lead to allosteric

regulation, as described in the ensemble allosteric model for IDPs[29,30].

Probing non-local intra-chain interactions is therefore essential for a molecular understanding

of IDP dynamics, binding, and function in general[22,31–33]. Direct experimental probing of such

interactions is particularly difficult, due to their transient nature, and the fact that IDPs undergo

large conformational changes on sub-microsecond time scales. Atomistic simulations of IDPs,

on the other hand, are equally challenging due to the wide conformational space sampled by

IDPs. To overcome these experimental and theoretical challenges, we leveraged the high

sensitivity of IDP dynamics to non-local interactions. We experimentally quantified the

dynamics of full-length N_{TAIL} in solution, by probing intra-molecular contact formation times,

using photo-induced electron transfer (PET) between a tryptophan (W) and a cysteine (C)

placed in the sequence[34–37]. Using this technique, we measured contact formation rates along

the well-defined C-W distance reaction coordinate, probing large-scale backbone motions on

the hundreds of nanoseconds to microsecond internal reorganization time scale[36], which is most

relevant to IDP function[38]. With minimal perturbation to the sequence, compared to introducing

large prosthetic dyes, we achieved a high sensitivity at short spatial distances. To obtain a

structural interpretation and a causal description, we compared the measurements to

corresponding observables derived from both simulations and analytical models. Our integrated

approach allowed us to identify key intra-protein interactions that extend beyond the MoRE

region, and play an important role in N_{TAIL} dynamics and likely also in function.

Results

Photo-induced electron transfer reveals the heteropolymeric nature of N_{TAIL} dynamics

To characterize the dynamics of N_{TAIL} in solution, we probed intra-molecular contact formation,

using photo-induced electron transfer (PET) between tryptophan (W) and cysteine (C) residues

at different positions in the sequence.[35–37] After exciting W to the triplet state using a

nanosecond UV laser pulse, we monitored the triplet state population in time, via transient

absorption measurements. As N_{TAIL} undergoes spontaneous conformational changes, W can

come into contact with C and PET can occur. During PET, the excited-state (triplet) electron is

transferred from W to C, forming a W radical cation which eventually relaxes to the ground

state. The relaxation time due to PET, τ_{CW} , therefore provides information on how often and

quickly C and W come into contact after W excitation. If the C-W contact formation rate is of

the same order of magnitude, or faster than the natural decay rate of W, τ_{CW} can be obtained

from the short time behavior of the triplet state relaxation curves (Supporting Fig. S1).

Figure 1. *Results of photo-induced electron transfer (PET) experiments at T=20 °C, on N$_{TAIL}$ variants containing a*

unique C-W pair. A) N$_{TAIL}$ sequence, showing charged residues at pH 7.6, location of Y to W, and S to C substitutions,

and location of MoRE, which folds into a helix upon binding to P$_{XD}$. B) Net charge per residue calculated using a window

size of six amino acids. C) W and C positions. Each variant contains a unique C-W pair. The C-W distance in each

variant is indicated by orange dashed lines. C3W2 and C4W2 variants encompass the MoRE transient helix. D) C-W

relaxation time τ_{CW}, due to PET as a function of C-W sequence separation. Experimental conditions: pH 7.6 and 150

mM NaCl (black circles); pH 4.0 and 150 mM NaCl (red squares); pH 7.6 and 500mM NaCl (green triangles). Dashed

line: expected scaling of τ_{CW}, from a SAW-v homopolymer model at pH 7.6 (black). Inset: Comparison of N$_{TAIL}$ circular

dichroism spectra at pH 7.6 (black) and pH 4 (red).

The wild-type N_{TAIL} sequence (Fig. 1A) does not contain native W or C residues. To probe the N_{TAIL} conformational dynamics, we generated five N_{TAIL} variants (PET variants), each bearing a unique C-W pair, to probe the five distances indicated in Fig. 1C. To minimize perturbation of the chemical properties, we substituted native tyrosine (Y) residues with W, and native serines (S) with C residues. To determine the natural decay rate of excited W at the two positions, we also generated two variants with the respective W substitution, but lacking any C substitution. The Supporting Information contains the sequences of all seven N_{TAIL} variants (Table S1) and details about their production and purification (Supporting Methods S1).

Relaxation times τ_{CW} were estimated from multiple, independent sample preparations for each N_{TAIL} variant, and a series of repeated measurements to account for W and C photodamage during the measurements (Supporting Methods S2). All measured relaxation curves and fitted functions to obtain τ_{CW} are shown in Fig S3-S16, with the fit results summarized in Table S2.

We first performed measurements with the five PET variants of N_{TAIL} under near physiological conditions (i.e. 15mM Tris, 150mM NaCl, 1mM TCEP, pH 7.6, in the following referred to as "reference conditions"). Figure 1D shows the τ_{CW} (black circles) estimated from these measurements as a function of the W-C sequence separation (|i-j|).

To test whether the estimated τ_{CW} can be explained based on C-W sequence separation alone, we compared them to relaxation times expected from a simple homopolymer model (Fig. 1D, dashed line), where the only non-local interactions between the residues are excluded volume interactions. In this model, both the physical distance in space between C and W at positions i and j as well as the corresponding relaxation time τ_{ij} scale with the sequence separation |i-j|.[39] The scaling relation for relaxation times $\tau_{ij} = \tau_r |i - j|^{\nu_\tau}$ depends on a prefactor τ_r and an exponent ν_τ. The exponent ν_τ = 1.69 (corresponding to a Flory scaling exponent ν of 0.52) was estimated from small-angle X-ray scattering (SAXS) measurements of N_{TAIL}[6] based on a

homopolymer model (SAW-ν model[40], Supporting Methods S4). The prefactor $\tau_r = 0.011$ μs

was obtained by fitting to the estimated τ_{CW} (Fig. 1D) from PET experiments.

The general trend of the estimated τ_{CW} at physiological pH (Fig. 1D, black circles) follows the

simple homopolymer scaling (dashed curve), however some contacts form faster and others

slower than expected. This dynamic heterogeneity (deviation from homopolymer model)

observed along the sequence is most striking when comparing the C3W2 ($|i-j| = 36$) and C4W2

($|i-j| = 30$) variants for which the C-W pairs encompass the MoRE region. These variants share

the W position and differ by just six residues in sequence separation between W and C (Figs.

1A-1C). According to the homopolymer model, such a difference in sequence separation would

result in a τ_{CW} increase by a factor of 1.3. In contrast, the τ_{CW} estimated from relaxation curves

increases by a factor of 2.8. This considerable deviation from the homopolymer behavior is

primarily due to the measured 7.6 μs relaxation time for C3W2 that is substantially longer than

the 4.7 μs homopolymer estimate. The fact that this anomalously slow relaxation time is not

reproduced by the homopolymer model indicates that the N_{TAIL} dynamics near the MoRE region

is considerably affected by more complex heteropolymeric interactions, likely secondary

structure formation, electrostatic interactions between charged residues, or solvent-induced

interactions.

Charged amino acids partially account for N_{TAIL} dynamic heterogeneity

To test whether interactions between charged residues or secondary structure formation causes

the dynamic heterogeneity, we performed additional experiments at low pH and increased salt

concentrations, i.e. under conditions that should modulate electrostatic interactions.

The N_{TAIL} sequence contains 40 charged residues (13 Asp, 10 Glu, 14 Arg and 3 Lys), often

clustered together (Fig. 1B). If electrostatic interactions between charged residues were

markedly contributing to the observed dynamic heterogeneity, we would expect that screening

the interactions by increasing the salt concentration would decrease or eliminate the

heterogeneity. Indeed, increasing the salt concentration from 150mM (Fig. 1D, black circles)

to 500mM (green triangles) results in a τ_{CW} decrease from 7.6 μs to 5.8 μs for C3W2, and an

increase from 3.2 μs to 3.9 μs for C3W1, bringing both data points close to the homopolymer

curve. This suggests that electrostatic interactions are a major contributor to the observed

dynamic heterogeneity of N_{TAIL}.

Next, we investigated why electrostatic interactions accelerate relaxation for the C3W1 variant,

but slow down relaxation for the C3W2 variant. We hypothesized that changes in contact

dynamics are caused by the non-uniform distribution of charged amino acids (charge

patterning[41,42]) along the N_{TAIL} sequence. However, the increased electrostatic screening at

higher salt concentrations could also affect the stability of the transient helix in the MoRE

region, and thereby modulate the C3W2 relaxation time. To address these two possibilities, we

performed coarse-grained molecular dynamics (CG) simulations that allowed us to explicitly

include both charge patterning and the secondary structure contributions.

Figure 2. *Coarse-grained (CG) simulation of N_{TAIL}. Differences between the mean pairwise distances ($R_{i,j}$) of the CG simulations at the reference condition (0.15 M salt and pH 7.6) and simulations A) at high salt concentration (0.5 M salt, pH 7.6) or C) low pH (0.15 M salt pH 4.0), respectively. The diagonals show the charged amino acid positions with positively charged amino acids in blue and negatively charged amino acids in magenta. The MoRE is highlighted in grey. The full amino acid sequence is provided in Fig 1A. B) Ratios of C-W relaxation times due to PET, calculated from CG simulations (bars) at different conditions (high salt: green, or low pH: red), relative to the reference condition. For comparison, experimental ratios are shown where available (black squares). D) C-W relaxation times for all five N_{TAIL} PET variants estimated from CG simulations at different conditions (reference: black, high salt: green, low pH: red). The blue triangles denote the relaxation times for the C3W2 and C4W2 variants after varying the MoRE helix content (helix- 35%, helix+: 60%). All CG relaxation times were scaled by a factor, chosen so that the same homopolymer fitting function (dashed line) of Fig 1D fit both experimental and CG data.*

For these simulations we used a modified version of the recently developed residue-based HPS model[43] for IDPs. The CG model was adjusted to reproduce the experimental radius of gyration of N_{TAIL} obtained from SAXS[6] and the helical propensity of the MoRE region from NMR[7]. The effects of varying salt concentrations in the CG model were described by Debye-Hückel electrostatic screening[44]. Additional details on the CG simulations are provided in Supporting Methods S5.

To determine the effect of increased salt concentration (150 mM to 500 mM), we computed the

mean distance for each pair of amino acids i and j from the CG simulations at both salt

concentrations. Fig. 2A shows the difference of these means, highlighting the

compaction/expansion of each N_{TAIL} segment. In the simulations, the N-terminal part of N_{TAIL}

up to C3 expands at higher salt concentrations (red areas), whereas the C-terminal part becomes

more compact (blue areas).

The positively and negatively charged amino acids are well-blended in the N-terminal region,

with a close to zero net charge. The N-terminus therefore behaves like a polyampholyte, where

interactions between oppositely charged residues result in compact conformations and

screening of these electrostatic interactions at higher salt concentrations results in more

expanded conformations. This expansion in turn increases the average distance between C3 and

W1, leading to a slower expected relaxation time at high salt concentration. In contrast, the C-

terminal region has more negatively charged amino acids than positively charged amino acids

under physiological conditions. This imbalance leads to net repulsive electrostatic interactions

that keep this region expanded. Increasing the electrostatic screening at high salt concentrations

reduces the repulsion, which leads to the compaction of the C-terminus of N_{TAIL}, a decrease of

the mean C3-W2 distance, and an expected decrease in the C3W2 relaxation time.

To determine if the salt-induced changes in relaxation times can be fully described by changes

in inter-residue distances due to charge patterning alone, we estimated the expected ratio of

relaxation times (τ/τ_0) from the CG simulations at 500 mM and 150 mM NaCl concentrations,

respectively. For these calculations, we assumed reaction-limited quenching kinetics, in which

case, the relaxation time is inversely proportional to the probability of W and C being at a

reactive distance. We used a cutoff distance of 1 nm to determine the probability of reactive

conformations (further details in Supporting Methods S5). Figure 2B shows the τ/τ_0 ratios

upon increasing the salt concentration for each variant (green bars). A ratio larger than one

indicates an increase of the relaxation time, exclusively due to the expansion of that segment.

Comparison with experimental ratios (Fig. 2B, black squares) shows that the CG simulations

accurately predict both the increase in C3W1 relaxation times and the decrease in C3W2

relaxation times, which indicates that charge patterning suffices to explain the salt-induced

changes in N_{TAIL} relaxation times. It also corroborates the hypothesis that electrostatic

interactions between charged residues are a key factor in the observed dynamic heterogeneity

in N_{TAIL}.

To further test the effect of electrostatic interactions on the dynamic heterogeneity of N_{TAIL}, we

measured C-W relaxation times, upon decreasing the pH from 7.6 to 4.0, while maintaining a

near-physiological salt concentration (150 mM) (Fig. 1D, red squares). In contrast to increasing

the salt concentration which screens all electrostatic interactions, decreasing the pH only alters

the charge on specific residues that titrate in the 7.6-4.0 pH range, specifically negatively

charged Asp and Glu may become neutral, and neutral His become positively charged. If

electrostatic interactions involving these charged residues were the main contributor to the

dynamic heterogeneity observed at physiological pH, we would expect a reduction in

heterogeneity when decreasing the pH to 4.0. In particular, we expect the N-terminal part,

which acts as a balanced polyampholyte at pH 7.6, to become positively charged and therefore

to expand at pH 4.0, even beyond what was observed at high salt concentrations. This expansion

should then increase the PET relaxation times for the C1W1 and C2W1 variants. Conversely,

we expect the negatively charged C-terminal part to become more neutral, resulting in more

compact conformations that reduce the C3W2 and C4W2 relaxation times.

To capture the effects of lower pH in CG simulations, the charges of the residues that titrate

between 7.6 and 4.0 were adjusted to match the average level of protonation at the simulated

pH. To that aim, the charges assigned at pH 7.6 (-1.0 for Asp and Glu and +0 for His residues)

were changed to partial charges at pH 4.0 (-0.69 for Asp, -0.36 for Glu and +0.99 for His).

Further details on charge assignments are provided in Supporting Methods S5. If electrostatic

interactions involving specific charged residues drive the dynamic heterogeneity at

physiological pH, lowering the pH to 4.0 should reduce this heterogeneity. The N-terminal

region, a balanced polyampholyte at pH 7.6, is expected to become positively charged and

expand, while the negatively charged C-terminal region should become more neutral and

compact. The CG simulations with simple fixed-charge adjustment at low pH are consistent

with the expected behaviors described above, i.e. the N terminal region expands and the C

terminal region becomes more compact (Fig. 2C), which is also reflected in the calculated

relative relaxation times (red bars Fig. 2B).

The C-W relaxation times of all five PET variants estimated from measurements at pH 4.0 are

closer to the homopolymer curve (Fig. 1D, red squares) than those at pH 7.6. Lowering the pH

also reduces the observed dynamic heterogeneity between the C3W2 and C4W2 variants, with

the ratio of relaxation times reduced from 2.8 at pH 7.6 to 1.5 at pH 4.0, which is closer to the

homopolymer prediction of 1.3. However, in contrast to the reduction of the C-W relaxation

time expected from compaction based on charge patterning and predicted by CG simulation,

the measured relaxation time for C4W2 increases slightly upon lowering the pH (Fig. 1D).

In Fig. 2B the ratios of relaxation times at pH 4.0 and pH 7.6 estimated from the CG simulations

(red bars) were compared to those estimated from PET experiments (black squares). The CG

model correctly captures the trend of C1W1 and C3W2, but not that of C4W2 and C3W1.

Further, the measured increase of τ_{CW} for C4W2 is not only contrary to the CG simulation

results, but also contrary to the expectation based on charge patterning, suggesting that other

unaccounted factors influence the relaxation times. In the following, we will consider three

factors that can affect relaxation times in a pH dependent manner: the MoRE helicity, W to C

electron transfer efficiency, and residue specific changes in protonation equilibria.

pH-dependence of MoRE helix content. The CG model suggests limited difference in helical

propensity upon changing pH (Supporting Fig. S20). We further measured and compared the

far-ultraviolet circular dichroism (CD) spectra of the N_{TAIL} variant Y518W (W2) at pH 7.6 (Fig

1C, inset, black curve) and 4.0 (red curve). CD spectroscopy in this wavelength range of 180-

260 nm is highly sensitive to the presence of α-helices and the two spectra appear to be very

close to each other, indicating that the helical content of N_{TAIL} is not affected significantly by

pH. In addition, analysis of the measured spectra by the SESCA software[45] (Supporting

Methods S13) predicts at most 8% global helix content, which is compatible with the NMR

estimates used to set the MoRE helicity in CG simulations. Together, these results suggest the

MoRE helix content is not affected by the change in pH.

Changes in electron transfer efficiency. Because PET involves formation of W radical

cations, the pH may directly affect the electron transfer process in solution, independent of pH-

induced changes in the protein conformational ensemble and dynamics. To address this

possibility, we performed a bimolecular quenching study between N-acetyl-tryptophan-amide

(NATA) and cysteine molecules, freely diffusing in solution, at pH 7.6 and 4.0 (Supporting

Methods S2). In these experiments the bimolecular quenching rate was determined by

measuring the W triplet relaxation times in samples at fixed NATA concentration and

increasing cysteine concentrations. These measurements showed that the C-W bimolecular

quenching rate (determined from the slopes of Stern-Vollmer plots) at pH 7.6 and 4.0 are very

similar (Fig. S19). This result shows that the efficiency of electron transfer from the excited

state W to C, under the conditions of our PET experiments is not directly affected by pH, and

the pH dependence of τ_{CW} in N_{TAIL} is due to changes in the protein structure and dynamics.

Residue-specific protonation. N_{TAIL} is rich in Asp and Glu, amino acids that reach protonation

equilibrium close to pH 4.0. Under these conditions, the protonation state of individual acidic

residues can be influenced, e.g., by the spatial vicinity of similarly or oppositely charged

residues, resulting in a dynamic, conformation-dependent charge distribution.[46] This charge

distribution would also allow electrostatic interactions to stabilize certain conformational states,

increase or decrease the observed τ_{CW} for individual N_{TAIL} variants compared to those predicted

by our fixed-charge CG simulations.

Dynamic heterogeneity at the N_{TAIL} C-terminus

Although dynamic protonation may explain deviations from CG and homopolymer models, this

effect is an unlikely explanation for the dynamic heterogeneity between C3W2 and C4W2

under reference conditions at pH 7.6. To address this heterogeneity with CG simulations, we

assumed these processes are reaction limited, and that τ_p the proportionality between C-W

contact probabilities ($P_{r_{ij}<r_0}$) and corresponding τ_{CW} relaxation times are the same for all PET

variants, and thus $\tau_{ij} = \tau_p/P_{r_{ij}<r_0}$. Under these assumptions, it is possible to compare

relaxation times of different C-W pairs obtained from CG simulations relative to one another

and address the dynamic heterogeneity at the N_{TAIL} C-terminus.

To this aim, we fitted a homopolymer model to these CG-derived relaxation times at the

reference condition, and chose τ_p such that the scaling equation is identical to the one obtained

from experimental measurements in Fig. 1D. The resulting relaxation times from CG

simulations for all three conditions (reference, high salt, and low pH) are shown in Fig. 2D. The

figure shows that all CG-derived relaxation times at the reference condition (black circles)

follow homopolymer scaling (dashed black line), with only small deviations for individual CW

pairs. Although the CG simulations accurately predict many of the changes upon increasing salt

concentrations or lowering the pH, they do not show either the considerable slowdown for the

C3W2 relaxation time, or the pronounced dynamic heterogeneity measured at pH 7.6 between

C3W2 and C4W2 (ratio 2.8). In fact, the CG-derived τ_{C3W2}/τ_{C4W2} ratio of 1.6 is close to the

1.3 ratio predicted by homopolymer models, suggesting that the CG simulations do not capture

the main cause of the dynamic heterogeneity observed in PET measurements.

An additional reason why the CG simulations may not have captured the C3W2/C4W2 dynamic

heterogeneity is that we underestimated or overestimated the MoRE helix propensity (~45%)

under the conditions of PET experiments. This can alter the CG-derived heterogeneity if the

MoRE helix in its folded state affects C3W2 and C4W2 relaxation differently. To test this

hypothesis, we performed additional CG simulations in which we altered dihedral potentials

that control the MoRE helicity and computed the τ_{CW} for C3W2 and C4W2 variants at high

(60%) and low (35%) helix contents (blue triangles on Fig. 2D). Relaxation times from these

simulations show that considerable changes in the helix content can indeed shift the relaxation

times obtained from CG simulations, but this shift affects both C3W2 and C4W2 similarly.

Therefore, even a significant error in the imposed MoRE helicity would have only a small

(±0.1) effect on τ_{C3W2}/τ_{C4W2} ratio, certainly not sufficient to explain the dynamic

heterogeneity observed in PET experiments.

Overall, our results show that N_{TAIL} dynamics is heteropolymeric in nature, especially in the

region encompassing the MoRE, with electrostatic interactions playing an important role. It

appears that the dynamic heterogeneity of N_{TAIL} cannot be explained by a simple polymer

scaling model (Fig. 1D) or by CG models including helix formation within the MoRE and

explicit interactions between charged amino acids (Fig. 2). These observations suggest that the

cause of the dynamic heterogeneity is either side-chain specific transient interactions, such as

salt-bridges and hydrogen bonds[47], or interactions between the protein and solvent. To study

the effects of these interactions on N_{TAIL} dynamics in more detail, we turned to all-atom

explicit-solvent molecular dynamics simulations.

Disordered N_{TAIL} is characterized by a conformational ensemble with distinct states

In addition to specific interactions between amino acids and protein-solvent interactions, all-atom simulations include the backbone and side-chain dynamics required to accurately compute the contact formation rates. The relaxation times τ_{CW} estimated from these contact formation rates can be compared to τ_{CW} derived from PET experiments. However, this increased level of detail comes with two drawbacks; considerably higher computational costs that limit conformational sampling and the need for a larger number of accurate interaction parameters (force field). IDP simulations are particularly sensitive to variations in the used force field[48–51], and force field accuracy for IDPs often varies in a system-dependent manner[52].

To maximize conformational sampling, we limited all-atom simulations to the reference conditions (pH 7.6 at 150 mM NaCl concentration) and focused on the C3W2 and C4W2 variants, which showed large dynamic heterogeneity (Fig. 1D). Further, to ensure the reliability of all-atom simulations, we first validated eight force fields, characterized by different combinations of protein and water parameters, against independent experimental SAXS, CD, and NMR[6,16] data on the wild-type (WT) N_{TAIL} sequence (shown in Supporting Table S3). To evaluate these force fields, we first performed simulations of WT N_{TAIL} using each force field, and then compared the resulting trajectories to the SAXS data, which reports on protein compactness, using the software CRYSOL[53]. Next, we compared the trajectories to CD and NMR chemical shift measurements, which report on secondary structure propensities, using SESCA[45], and SPARTA+[54]. For further details of the force field validation, see Supporting Methods S8.

The two force fields that reproduced the WT validation data most accurately were the CHARMM36m (C36M)[51] force field with optimal point charge (OPC) water model[55], and Amber99SB-disp (A99SB-d) force field[49]. Both of these force fields produced extended N_{TAIL} ensembles with transiently forming helices in the MoRE region, that were in good agreement

with the available experimental data (Supporting Table S3), and suggested helix propensies

similar to those reported in computational studies of MoRE region[19,26] (Supporting Methods

S23).

The two best force fields were used to perform longer all-atom simulations of the C3W2 and

C4W2 variants. As a second force field validation step (Supporting Table S4), the variant

trajectories were tested against PET relaxation measurements of the same variants, and the CD

spectrum of the W2 variant (Fig. S21B). Simulations of both force fields estimated C3W2 and

C4W2 relaxation times showing considerable dynamic heterogeneity. However, the C36M-

OPC force field was the only parameter set that reproduced measurements of both WT N_{TAIL}

and the two variants within the experimental and computational uncertainty, including the two

measured W relaxation curves (Supporting Fig. S22). Thus C36M-OPC trajectories were used

for all subsequent analysis. The relaxation times τ_{CW} estimated from these simulations were

2.0 ± 0.5 µs for C4W2 and 5.3 ± 3.3 µs for C3W2, in good agreement with the corresponding

measured values of 2.3 µs and 7.6 µs. Although the uncertainty on the C3W2 relaxation time

is large due to limited sampling, the ratio of the relaxation times estimated from the simulations

is 2.7 ± 1.8 close to the experimentally determined ratio of 2.8. Together, these results suggest

that the all-atom simulations accurately describe the N_{TAIL} dynamics causing the C3W2/C4W2

dynamic heterogeneity.

To study the origin of the dynamic heterogeneity in detail, we computed the free energy

landscape of the N_{TAIL} variants along the C-W distances for positions C3W2 and C4W2.

Variants C3W2 and C4W2 only differ in the exchange of an oxygen and a sulfur atom between

two nearby residues at position 482 (C3) and 488 (C4), respectively. We therefore assumed that

the free-energy landscape of the two variants is similar and combined their trajectories for

computing it. The C-W distances were defined as the distance between the gamma sulphur of

C (or gamma oxygen of S) and the nearest heavy atom of the W side-chain ring. These serve as

intuitive reaction coordinates for describing C-W contact dynamics in the N_{TAIL} C-terminal

region.

Unexpectedly, the computed free-energy landscape (Fig. 3) is characterized by four local

minima corresponding to four N_{TAIL} conformational states (S1 to S4). Approximately 60% of

the sampled N_{TAIL} conformations are within the respective energy wells, including almost all

conformations with C-W contacts necessary for PET. S1 corresponds to the conformational

state around the absolute free-energy minimum and is occupied for approximately 24% of the

simulation time. S2 and S3 are less populated conformational states, occupied for 14% and 8%

of the simulation time, respectively. S4 is the least populated state, with approximately 2% of

conformations. The landscape shows only small barriers (< 1 kT) separating S1, S2, and S3,

and a low free-energy barrier between S1 and S4 (< 2.0 kT). This shallow and accessible free

energy landscape suggests nearly-free diffusion between states, which is in line with previous

SAXS results suggesting N_{TAIL} to be a pre-molten globule with residual structure[6].

Figure 3. Conformational states revealed by all-atom MD simulations. The free-energy landscape of sampled N_{TAIL}

The four conformational states influence C-W relaxation times as well. PET cannot occur in

the two most populated states (S1 and S2), because the distances between electron acceptors

C4 or C3 and electron donor W2 are more than 2 nm. In state S3, C4-W2 distances below 0.4

nm are observed, allowing PET in the C4W2 variant, while in state S4, C3-W2 distances below

0.4 nm are reached, which allows PET in the C3W2 variant. Further, from the most populated

S1 state the free-energy barrier that N_{TAIL} has to cross to reach S4 is about 0.9 kT larger than

the barrier to reach S3. This suggests that S3 is more easily accessible than S4, with an

approximately 2.5-fold difference between the respective transition rates. This result is

consistent with and provides a possible explanation for the measured 2.8 ratio between the

C3W2 and C4W2 relaxation times.

Non-local N_{TAIL} interactions explain dynamic heterogeneity

To determine the molecular basis of the observed dynamic heterogeneity, we investigated

which specific interactions stabilize the four conformational states identified above. To this

aim, Fig. 4 shows intramolecular contact probability maps for each conformational state, with

backbone (C_α) contacts shown in yellow to green. Because both the pH and salt concentration

dependence of PET experiments and CG simulations indicated an important role of electrostatic

interactions in N_{TAIL} dynamics, we also calculated contact probability maps for salt bridges

between charged amino acids shown in purple to blue. For the sake of clarity when discussing

the contacts, we defined five regions along the N_{TAIL} sequence (Fig. 4 bottom, color-coded and

labelled A-E). Regions A-C are subdivisions of the polyampholyte N-terminal part identified

in our CG simulations, regions D and E constitute the negatively charged C-terminal part, with

the MoRE located in region D.

Figure 4. Contact maps and representative conformations in N_{TAIL} conformational states. For each conformational state

/

The contact maps (Fig. 4, right) clearly show that the four states are stabilized by distinct sets

of local (within a region) and non-local (between regions) interactions, although some

interactions occur in several conformational states. The maps also provide information on the

secondary and tertiary structure preferences of N_{TAIL} in these conformational states. Helical

secondary structure elements and tight turns appear on contact maps as continuous sets of

interactions, parallel and close (within 5 residues) to the diagonal. β-strands, β-hairpins or

longer loops appear as sets of interactions that are further away and either parallel or

perpendicular to the diagonal. Disordered regions typically have few or sporadic interactions.

The four conformational states show markedly different structures (Fig. 4, left), including the

MoRE in region D. In state S1, region D is largely non-helical but forms several transient local

salt bridges. Several non-local interactions (encircled in green) may prevent donor-acceptor

contacts for both variants (C3W2 and C4W2). S1 conformations are also stabilized by salt

bridges formed between regions B and E. Salt bridges involving K441, R444 (region B) and

E507 and D511 (region E) are particularly frequent and probably play an important role in

stabilizing this conformational state.

State S2 includes conformations where region D is non-helical and conformations with a folded

α-helix (residues 492-502). The helical structures appear on the contact map as an interaction

patch close to the diagonal of region D (encircled in orange). Non-local interactions in S2

appear to be less stable than in S1 and include contacts between regions A and C, C and D, as

well as B and D. Stable interactions in S2 also involve two salt bridges between R413 and D476

(regions A and C) and K441 and E449 (local in region B). Many sporadic interactions (both

backbone contacts and salt bridges) were observed within the N-terminus (between regions A

and B). Taken together, the interactions suggest that S2 consists of multiple conformational

substates and most interactions are linking central regions (B and C) with both termini (A, D,

and E).

Compared to S1 and S2, the conformations of S3 comprise fewer non-local interactions and are

instead characterized by two very stable local interaction sites. These local interactions

correspond to a shorter helix between residues 492-497 present in all S3 conformations and to

a stable hairpin or zipper in region B, stabilized by a salt bridge (K441-E449) observed

previously in S2. Further, there are two sets of notable non-local interactions. First, a patch of

interactions between regions D and E folds the C-terminus into a loop, stabilized by the salt

bridge between R490 and D520. These interactions also bring the electron donor W2 (518) very

close to the potential acceptor at position C4 (488), thus promoting electron transfer in the

C4W2 variant. At the same time, the zipper in region B brings the N-terminal close to regions

C and D, where a salt bridge between K405 and D484 may prevent electron transfer to position

C3 (482) in the C3W2 variant.

The S4 state allows electron transfer to position C3 (482) in the C3W2 variant. It is structurally

very different from states S1-S3. The most defining structural feature of this state is a long loop

reminiscent of an anti-parallel β-sheet stretching over the entire C-terminus (regions D and E).

This loop is stabilized by interactions between regions C and E including a salt bridge between

R463 and D513. The N-terminus of N_{TAIL} in this conformational state has only weaker

interactions, mostly limited to contacts between regions A and C as well as between regions B

and C.

To determine if frequent contacts observed in one of the four conformational states (Fig. 4) are

specific to that state and if they are correlated with other contacts, including C-W contact

formation for the C3W2 and C4W2 variants, we applied normalized pointwise mutual

information analysis (NPMI, see Supporting Methods S22).

The NPMI analysis revealed that many salt bridges between regions B, D, and E occur

selectively in their respective conformational states. These stable salt bridges often co-occur

with clusters of backbone contacts (encircled in Fig. 4), possibly nucleating the interaction

clusters that stabilize the states.

Further, all frequent contacts observed in the S1 state were anti-correlated with C-W contact

formation (Table S5), indicating that the specific interactions that stabilize the S1 state would

increase the expected relaxation times compared to the flat free-energy landscape of a

homopolymer model. In contrast, all frequent contacts of states S3 and S4 are positively

correlated with C-W contact formation for variants C4W2 and C3W2, respectively. Thus,

interactions in these two states selectively reduce the relaxation times of either C4W2 or C3W2

variants, contributing to the dynamic heterogeneity observed in PET experiments.

Finally, the same charged residues (e.g., K441, R497, and D501) in regions B, D, and E form

salt-bridges with different residues in different conformational states (Table S5). Switching

between these sets of interactions is necessary to transition between the most populated

conformational states, and therefore likely define the free-energy barriers between the states

(Fig. 3). Further, a conformational switch based on these competing intramolecular N_{TAIL}

interactions of region B may have a functional role, by regulating the helical propensity of the

MoRE in region D, which in turn modulates affinity for the measles phosphoprotein.

Functional relevance of non-local interactions is supported by coevolutionary analysis

Having identified the most important interactions that dictate N_{TAIL} structure and dynamics (Fig. 4), we asked whether they also have a functional role. To this aim, we performed a coevolutionary analysis of N_{TAIL} using EVcouplings[56], which identifies and quantifies correlated mutations of residue pairs within the sequence of homologous proteins from different species or strains, and which often indicates pairwise-interactions of functional importance. A total of 12,072 sequences similar to N_{TAIL} have been aligned and selected for this analysis (Supporting Methods S21).

Most non-local residue pairs with positive EVcouplings scores connect regions B and D (Fig. S28), suggesting evolutionarily conserved interactions between these two regions. Here we specifically highlight the four non-local residue pairs with the highest EVcoupling scores (Fig. S28, top left, Table S5). These pairs include hydrophobic contacts involving three leucine residues (L491, L496 and L498) interacting with S443 and A447, and one salt bridge between R444 and D493. Correlated mutations of residues in the distant regions B and D suggest a biologically relevant, functional interaction between these two regions of N_{TAIL}. Given that both L491 and L498 are located in the MoRE region (in region D) and are directly involved in binding to P_{XD}, it is possible that their interactions with residues of region B serve the function of regulating binding. We note that the contact analysis of our simulations does not identify these specific coevolving residue pairs as frequent interactions. However, all-atom simulations independently identify non-local interactions between regions B and D (Fig. 4, S2 state), including A447 contacts with P_{XD} binding residues L498 and M501. These differences between all-atom simulations and the co-evolution analysis may be due to insufficient sampling or the lack of biological context in all-atom simulations (i.e. the absence of the N_{CORE} domain or P_{XD} itself).

Discussion

To understand the role of disordered regions flanking the MoRE on either side in N_{TAIL} dynamics and conformational preferences, we probed non-local interactions of the full-length protein in solution, combining photo-induced electron transfer experiments, analytical polymer models, molecular dynamics simulations, as well as coevoluionary analysis. With PET experiments, we measured and compared contact formation times between tryptophan and cysteine residue-pairs introduced in different regions of the N_{TAIL} sequence, under varying solution conditions. The experiments revealed a pronounced dynamic heterogeneity within the protein (Fig. 1) under physiological conditions, especially at the N_{TAIL} C-terminal region, close to the MoRE, which is involved in binding to P_{XD}. PET experiments also showed a reduced heterogeneity under conditions that weaken electrostatic interactions, such as increased salt concentration or lower pH, pointing to an important role of electrostatic interactions between charged residues in the dynamics of N_{TAIL}. Based on our molecular dynamics simulations, we attribute this heterogeneity to salt-bridge induced non-local interactions between specific regions of N_{TAIL}.

Interpretation of the general role of electrostatics

This finding is supported and explained in structural terms by our simulations (both coarse-grained and all-atom). Even without including atomic-level details, our CG simulations suggest that when increasing the electrostatic screening or changing the charge of acidic residues to emulate pH related protonation, the N-terminal region expands while the C-terminal region compacts. In addition, CG simulations performed with different helical propensities in the MoRE indicate that changes in helicity of the MoRE do not give rise to the dynamic heterogeneity within the N_{TAIL} C-terminal region (i.e. C3W2 and C4W2). These results indicate that side-chain specific transient interactions, not described by the CG model, contribute to the large difference between the C3W2 and C4W2 relaxation times.

Specific non-local interactions dictate N$_{TAIL}$ conformation and dynamics

Our all-atom simulations, performed with a carefully validated force-field, describe interactions

with atomic-level detail. These simulations reproduce the dynamic heterogeneity of C-W pairs

encompassing the N$_{TAIL}$ MoRE (C3W2 compared to C4W2). We found that, while N$_{TAIL}$ is

clearly disordered, a few key non-local interactions stabilize four distinct conformational states

which in total represent 48% of the structural ensemble. The two most populated states do not

allow PET for either C3W2 or C4W2 variants, while the third and fourth states allow PET for

the C4W2 and C3W2 variants, respectively. The free energy landscape of N$_{TAIL}$ (Fig. 3) reveals

rapid interconversions between the three most populated states. In contrast, structural

rearrangements required to reach the C3W2 contact state are slower due to a small, though

comparatively higher free-energy barrier (~4.4 kJ/mol) under physiological conditions. This

free-energy barrier and the lower equilibrium population of the C3W2 contact state contributes

to the dynamic heterogeneity of the N$_{TAIL}$ C-terminal region.

Analysis of the state-specific contacts (Fig. 4) reveals that competing clusters of interactions

stabilize the four conformational states. These interaction clusters involve (and are possibly

nucleated by) stable salt bridges between charged amino acid side chains. Further, focusing on

the two most populated conformational states S1 and S2, our all-atom simulations show that

specific charged residues in region B (residues 435-451) can form either non-local salt bridges

with C-terminal regions D-E (484-525, close to the MoRE), or local salt bridges with other

region B residues. Switching between these sets of interactions (as sketched in Fig. 5) is

necessary for N$_{TAIL}$ to transition between the two dominant states, affecting its large-scale

dynamics.

Figure 5. *Schematic representation of the two major N$_{TAIL}$ conformational states (S1 and S2). Different regions of N$_{TAIL}$*
are color-coded according to Figure 4. Non-local interactions between the regions are represented as dashed lines.
Key interactions between regions are highlighted by orange shading. The helical binding site of the phosphoprotein X-
domain (P$_{XD}$) in region D is marked by an orange arrow.

Co-evolutionary analysis independently shows that correlated mutations across regions were

only found between regions B and D. Given the distance in sequence between these two regions,

this not only corroborates the presence of non-local interactions between region B and D, but

also suggests that they play a functionally important role.

In summary, our simulations and coevolutionary analysis independently highlighted region B

of N$_{TAIL}$ as a potential allosteric interaction partner to region D, which includes the binding site

(MoRE) for the phosphoprotein X-domain (P$_{XD}$). Additionally, fluorescence-based binding

kinetics experiments on truncated N$_{TAIL}$ variants[23], indicated that intramolecular interactions of

N$_{TAIL}$ involving residues 435-451 (region B) have a significant impact on P$_{XD}$ binding in vitro,

despite being located far away in sequence from the MoRE.

Identified interactions in the context of N$_{TAIL}$ to P$_{XD}$ binding

We hypothesize that the intramolecular N$_{TAIL}$ interactions observed in our simulations not only

govern the N$_{TAIL}$ dynamics but also play a role in regulating P$_{XD}$ binding to the MoRE.

Mutational studies by Bignon *et al.*[57] have shown that the helix content of the MoRE correlates

with the binding affinity, and that N-terminal truncation of N_{TAIL} increases its affinity for

P_{XD}[22,24,27]. Similarly, Gruet *et al.*[27] have identified mutations in several regions of the N_{TAIL}

sequence, on either side of the MoRE, which increase P_{XD} binding affinity. The crystallographic

structure of the N_{TAIL} MoRE bound to P_{XD}[9] (PDB: 1t6o, Fig. 6B) shows that the binding

interface consists mainly of a hydrophobic edge along the MoRE helix (region D), between

residues S491 and M501 (hereafter D1-site, shown in dark green), which interacts via

intermolecular contacts with the $\alpha2$-$\alpha3$ face of the P_{XD} triple helix (Fig. 6A). The other side of

the MoRE helix is a hydrophilic edge formed by charged and polar residues (D2-site, dark red)

between Q486 and R497. The N_{TAIL}-P_{XD} complex is further stabilized by intermolecular

interactions involving R497 and salt bridges formed by D487 and R490 which are located at

the beginning of the MoRE helix between the two edges (D3-site, light green)[27]. Many of these

key amino acids are involved in intramolecular N_{TAIL} interactions in one or more

conformational states, as shown by the contact probabilities obtained from all-atom simulations

(Fig. 4).

Based on these interactions, we propose two independent mechanisms that provide a possible

explanation for the observed changes in P_{XD} binding affinity upon N_{TAIL} mutation or truncation.

Figure. 6. *Speculated biological context of non-local N$_{TAIL}$ interactions. (**A**) N$_{TAIL}$ MoRE residues in region D constituting the hydrophobic P$_{XD}$-binding interface are highlighted (D1 site, dark green) along with charged residues possibly involved in intramolecular N$_{TAIL}$ interactions (D2 site, dark red), and residues possibly involved in both intra- and intermolecular interaction (D3 site, light green). (**B**) Cartoon representation of the P$_{XD}$-MoRE complex based on the available crystal structure (PDB: 1t6o). (**C**) During the initial steps of the virus replication the MeV nucleocapsid protein (N, blue) recruits the large polymerase (L, green), by binding to the phosphoprotein (P, orange). Here N$_{TAIL}$ is extended. (**D**) When Brownian motion allows L to approach the RNA (red) in the nucleocapsid groove, N$_{TAIL}$ adopts a hairpin conformation, which promotes intramolecular interactions between the MoRE and the N-terminal region of N$_{TAIL}$. (**E**) Once the contact between nucleocapsid and L is established, additional complexes can be formed between free P$_{XD}$ and downstream N monomers. Further, intramolecular N$_{TAIL}$ interactions help release bound P to allow polymerase progression, by weakening the binding to P$_{XD}$ when L is nearby. Numbers (1-4) on P$_{XD}$ denote which step individual domains are in a "cartwheeling" L progression mechanism. In panels C, D, and E, only 1-3 N$_{TAIL}$ domains have been depicted for the sake of clarity. The RNA molecule is shown as a solid line on its first helix turn around the nucleocapsid to indicate that it is more solvent exposed, and as a dashed line on downstream turns to highlight that it is partially buried in the RNA-binding ridge.*

Direct competition for MoRE: In all-atom simulations, the D1-site residues (Fig. 6A, dark green) form intramolecular interactions with residues of regions B and C in state S2 as well as with residues of region E in state S3 (see Supporting Table S7), both of which are states with considerable helical propensity of the MoRE region. Assuming that a folded MoRE helix is a

necessary part of the biologically active N_{TAIL} conformation, these intramolecular interactions

would directly compete with the intermolecular interactions with P_{XD}.

First, the intramolecular interactions would reduce the free helix population of the MoRE,

thereby decreasing the binding rate of P_{XD}. Second, the intramolecular interactions may also

displace P_{XD} from the binding site without exposing the hydrophobic D1 site to water, thereby

lowering the P_{XD} unbinding activation barrier and increasing the unbinding rate. Both effects

would regulate MoRE-P_{XD} binding by reducing the binding affinity, and explain why the

binding affinity increases when flanking regions of the MoRE are removed[27,57].

Indirect allosteric regulation: Helical conformations of the N_{TAIL} MoRE appear to have higher

affinity for P_{XD}[57]. Therefore, changing the population of helical MoRE conformations via

intramolecular interactions would also affect the MoRE-P_{XD} binding affinity. This mechanism

is similar to the ensemble allosteric model for IDPs described by White *et al*[29]. Specifically,

intramolecular interactions that stabilize non-helical MoRE conformations, such as the contacts

seen in state S1, would decrease the population of helical conformations, reducing both N_{TAIL}

binding affinity and the rate of binding to P_{XD}. The observed contacts (Fig. 4) suggest that the

ratio of states S1 and S2 is controlled by the competing interactions of region B, which acts as

a conformational switch. Specifically, in state S1, the interactions between regions B and E

stabilize a loop-like, non-helical conformation of the N_{TAIL} C-terminus, which is likely binding-

incompetent towards P_{XD}. In contrast, in state S2, region B residues engage in local interactions,

allowing the N_{TAIL} C-terminus to form the high-affinity helical structure in the MoRE at region

D.

Potential functional role of N_{TAIL} intramolecular interactions in viral replication

Non-local interactions between regions B and D/E of N_{TAIL} may also play a functional role in

RNA polymerization and viral replication. In the following we present three established

mechanisms where intramolecular interactions between the B and D regions of N_{TAIL} may

contribute to optimizing the binding kinetics for viral replication.

Tether shortening: Initial polymerase complex recruitment likely involves the formation of one

or more disordered tethers between the polymerase complex and the nucleocapsid. By limiting

the volume available for diffusion, tethering increases the local concentration of the polymerase

complex around the nucleocapsid, thereby increasing the rate of transcription/replication

initiation. There are two experimentally established tethers between N and P. The first tether

consists of the disordered N-terminal tail of P (1-304, P_{TAIL}) binding to two opposite sides of

N_{CORE}, and its primary function appears to be the prevention of nucleocapsid association in the

absence of the viral RNA[58]. The second, tether is formed by regions B and C of N_{TAIL} and the

C-terminal domain (P_{LOOP}, 376-459) of P, through coupled binding and folding between the

MoRE of N and P_{XD} (Fig. 6C). This tether regulates L progression on the nucleocapsid.[59] The

two tethers may coexist and in fact work in concert to form N-P condensates that serve as the

basis of viral factories within the host cell[60,61], and to securely attach the polymerase complex

to the nucleocapsid during transcription/replication initiation[11].

During replication initiation, it is preferable to have long tethers to recruit polymerase

complexes from a larger volume around the nucleocapsid, but later, when L has to be guided to

the 3' end of the RNA, shorter tether lengths may be preferable. Intramolecular N_{TAIL}

interactions between the MoRE D2-site and region B would shorten the tether by 45 amino

acids (Fig. 6D), further reducing the volume available to the polymerase complex, and increase

the rate of initiation of genome transcription and/or replication.

P_{XD} unbinding during polymerase progression: Sourimant *et al.*[10] carried out recombinant

minigenome bioactivity assays found that polymerase progression is slowed down considerably

by a deletion of regions B and C of N_{TAIL} and proposed a mechanism for polymerase

progression. This mechanism is based on a newly discovered binding site on N_{CORE} for the α1-

$\alpha 2$ face of P_{XD}, at which compensatory point mutations could weaken P_{XD}-N_{CORE} binding and

restore bioactivity. Briefly, the mechanism suggests that the L-P polymerase complex is

tethered to the nucleocapsid by forming the N_{TAIL} MoRE– P_{XD} complex. Then, the P_{XD}-MoRE

complex binds the recently discovered N_{CORE} site. P_{XD} binding to this N_{CORE} site likely

facilitates P_{XD}-MoRE unbinding, possibly after compaction of the N_{TAIL} loop to allow

polymerase passage. P_{XD} unbinding in turn allows subsequent binding to MoRE of the

downstream N monomer, and polymerase translocation.

The mechanism proposed by Sourimant *et al.* provides previously unprecedented molecular

insight into the measles virus replication. However, it mostly considers the deleted regions of

N_{TAIL} as passive linkers or "roadblocks" for L, and it does not give an in-depth explanation on

how weakening the N_{CORE}-P_{XD} binding affinity compensates for reduced bioactivity caused by

the deleted N_{TAIL} segments. Our current results and previous binding efficiency studies on

truncated N_{TAIL} constructs[22] suggest that non-local interactions between MoRE and N_{TAIL}

regions B and C can in fact actively compete with P_{XD} for its binding site and may do so in the

context of a MoRE-P_{XD}-N_{CORE} ternary complex as well. This implies that deleting regions B

and C does not just shorten N_{TAIL} but also eliminates intramolecular interactions that promote

MoRE-P_{XD} unbinding. This would explain why the deletion hampers viral replication, why

mutations at the N_{CORE} interface compensate for the deletion, and provides a possible reason

for region B residues to co-evolve with the MoRE (Table S5). In addition, based on the

available cryo-EM structure of the nucleocapsid[1] (PDB: 4uft), region B would emerge from the

RNA binding ridge close to the N_{CORE} binding site, with sufficient space for region B to fold

into a loop and interact with the MoRE region (Fig. S24).

Polymerase progression by cartwheeling: Our findings are also compatible with a modified

version of the cartwheeling mechanism[62] for polymerase progression (Fig. 6E). Here, P is

recruited (Fig. 6E, step 1) by N_{TAIL} and bound to the N_{CORE} binding site (step 2) of the same

monomer as described by Sourimant *et al.*[10]. However, in this mechanism P_{XD} remains bound

to the same N monomer until the L polymerase approaches. Then, due to steric hindrance from

L or conformational changes at the N-N interface, N_{TAIL} folds back onto the nucleocapsid (step

3) in a conformation that allows region B to weaken P_{XD}-MoRE interactions. This, in turn,

allows P_{XD} to unbind and be transferred to its binding site on L (step 4) as L progresses.

Subsequently, P_{XD} is either released to the solvent or directly binds to the N_{TAIL} of a downstream

N monomer in the next cartwheeling cycle.

As discussed above, evidence suggests intramolecular N_{TAIL} interactions weaken P_{XD} binding.

Tight binding to P_{XD} is beneficial for tethering the viral polymerase to the nucleocapsid during

polymerase recruitment and progression, and thereby ensures that the transcription and

replication of the genome is completed. However, unbinding of individual P_{XD}-N_{TAIL}

complexes is required for polymerase progression during viral replication[63]. Therefore, MeV

and related viruses may be at an evolutionary disadvantage if the binding between P_{XD} and

N_{TAIL} is so tight that it limits the replication rate, as supported by mutational studies of P_{XD}[64]. It

is thus tempting to hypothesize that a possible functional role of non-local intra-protein N_{TAIL}

interactions is to provide a conformation-dependent optimization of P_{XD} binding affinity.

Conclusions

Our combined experimental and simulation approach shows that N_{TAIL} is clearly disordered,

contains very little secondary structure, and shows nearly diffusive dynamics. However, it still

displays some structural features which cause its dynamics to deviate significantly from that of

a homopolymer. In particular, we identified several key transient interactions between

disordered regions distant in sequence as the main reason for this deviation. These interactions

affect the overall conformation and dynamics of N_{TAIL} in solution. Interestingly, similar

interactions also emerge from our independent co-evolutionary analysis, corroborating our

simulation results and suggesting that they not only govern the conformational dynamics of the

essential N_{TAIL} domain, but also are functionally important. We therefore propose possible

mechanisms by which these non-local interactions regulate binding to the phosphoprotein X

domain (P_{XD}), and consequently, recruitment and progression of the polymerase complex onto

the nucleocapsid template. Our results for N_{TAIL} corroborate the importance of flanking regions

in IDPs that bind via short molecular recognition elements (MoREs)[64], and suggest a regulatory

role of regions that may be far in the sequence from MoREs. These non-local interactions within

N_{TAIL} can regulate binding *via* an ensemble allosteric model[29,30] without requiring the binding

of a third molecule. Further studies targeting the N_{TAIL}-P_{XD} interaction may provide additional

insight into the underlying mechanisms and effects of these interactions.

It is plausible that similar mechanisms are also at work for other IDPs that share sequence

characteristics, such as charge patterning, with N_{TAIL}. Specifically, non-local intra-protein

interactions in IDPs may regulate the conformational preferences and dynamics of both the

MoRE and of the entire protein, either via direct competition or indirectly by shifting the

population of free-energy minima.

Methods

Protein expression, purification, and sample preparation.

Wild type (WT) N_{TAIL} and the seven variants listed in Supporting Table S1, were expressed in

E. coli, and further purified as described in detail in Supporting Methods S1. Briefly, after

expression of tagged protein in *E. coli*, cell pellets were resuspended in 8 M urea, 50 mM Tris

pH8, 0.3 M NaCl and frozen. After thawing, the solution was sonicated, spun, and tagged

proteins were purified using Ni-Sepharose fast flow beads (Cytiva). After dialyzing the eluent

in 50 mM Tris pH8, 0.3 M NaCl, TEV protease was used to remove tags from the desired

protein. TEV protease as well as uncut tagged proteins were eventually removed using Ni-

sepharose beads. Purified N_{TAIL} proteins, free of histidine tags, were dialyzed against 15 mM

Tris, 150 mM NaCl, 1 mM TCEP buffer at neutral pH, checked using SDS-PAGE, UV

absorption, and far-UV circular dichroism (CD), and frozen for shipping. After shipping,

samples were further purified via HPLC (semipreparative Vydac C18 column) and lyophilized.

Before PET experiments, the lyophilized protein was dissolved directly into filtered buffers (for

reference conditions: 15 mM Tris, 150 mM NaCl, 1 mM TCEP pH 7.6; for low pH

measurements: 20 mM NaAc, 150 mM NaCl, 1 mM TCEP pH 4.0; for high salt concentrations:

15 mM Tris, 500 mM NaCl, 1mM TCEP pH 7.6). Protein concentration was adjusted to ~100

μM as determined by UV-absorbance at 280 nm (extinction coefficient 6990 cm^{-1} M^{-1}). Samples

(~350 μL) were placed in 5x10x30(h) mm Spectrocell quartz gas tight cuvettes with screwcap

(equipped with Teflon coated silicon membrane), and bubbled with USP grade nitrous oxide

for at least one hour, to reduce the concentration of dissolved oxygen (a quencher of excited

tryptophan triplet state), and to introduce solvated electron scavenger (to prevent reactions with

electrons, produced by water decomposition under UV laser pulses).

Photo-induced electron transfer (PET) experiments.

To probe intra-molecular contact formation, we used a technique based on photo-induced

electron transfer (PET) between a tryptophan (W) and a cysteine (C)[34–36] placed at different

positions within the sequence (sequences in Supporting Table S1). Details are described in

Supporting Methods S2. Briefly, we used a homebuilt nanosecond transient absorption

apparatus to excite the W to the triplet state, and to monitor the excited state population as a

function of time. When, C comes into contact with W (within Van der Waals distance) due to

stochastic collisions, an excited-state electron is transferred from the triplet state of W to C. As

explained in Supporting Methods S2, and in Sizemore et al. 2015[36], the measured relaxation

time is related to the intra-molecular contact formation time between W and C in the protein.

The contribution to the triplet state relaxation times due to C-W quenching via electron transfer,

are reported in Fig. 1D. These were obtained by fitting W triplet state relaxation curves as

illustrated in Supporting Figs. S1 and S2, and taking into account the natural lifetime of the

triplet state in the absence of C, measured on single W variants (see Supporting Methods S2).

Each value is the result of globally fitting multiple, repeated measurements, under given

solution conditions. All raw data and their fits are shown in Supporting Figs. S3 through S16.

The corresponding fitting parameters, obtained for each variant under each solution condition,

are reported in Supporting Table S2. Solution conditions used in PET experiments of Fig. 1

were, for reference conditions: 15 mM Tris, 150 mM NaCl, 1 mM TCEP pH 7.6; for low pH

measurements: 20 mM NaAc, 150 mM NaCl, 1 mM TCEP pH 4.0; for high salt concentrations:

15 mM Tris, 500 mM NaCl, 1mM TCEP pH 7.6.

Circular Dichroism

Circular Dichroism (CD) spectra reported in Fig. 1D inset were measured as described in

Supporting Methods S3. To improve signal to noise, and access low wavelengths which are

particularly important for the interpretation of IDP spectra, at each pH we carried out

measurements for a series of protein concentrations and combined them. Measurements at high

salt concentration were not possible due to the high absorbance of these samples below 210nm,

a range which is most important for the interpretation of IDP spectra. Solution conditions used

for spectra of Fig. 1 were: 10mM NaPO$_4$, 150 mM, NaF for pH 7.6 measurements, and 20 mM

NaAc, 150 mM NaF for pH 4.0 measurements.

Coarse-grained molecular dynamics

We applied a coarse-grained model based on the original HPS model[43] to N$_{TAIL}$. Each amino

acid is represented by a bead with charge (+1, 0, -1) and hydropathy. There are three types of

interactions in the HPS model: bonded interactions, electrostatic interactions, short-range

pairwise interactions. The electrostatic interactions are modeled using a Coulombic term with

Debye-Hückel[43] electrostatic screening to account for salt concentration. The short-range

pairwise potential accounts for both protein-protein and protein-solvent interactions, which was

optimized using the experimental radius of gyration of N$_{TAIL}$[6]. We further added additional

terms for angle and dihedral preference so that the secondary structure of the MoRE region can

be captured (see Supporting Methods S5). The HOOMD-Blue software v2.9.3[65] together with the azplugins (https://github.com/mphowardlab/azplugins) were used for running the molecular dynamics simulations. All simulations were run using a Langevin thermostat with a friction coefficient of 0.01 ps^{-1}, a time step of 10 fs and a temperature of 298 K for 5 µs. The first 0.5 µs were dumped for equilibration.

All atom molecular dynamics

We performed all molecular dynamics (MD) simulations on three full length N_{TAIL} variants. The wild type (WT) N_{TAIL} (401-525) simulations were performed including an N-terminal hexa-histidine (His6) tag, while the latter was not included in the case of variants C3W2 (i.e. S482C-Y518W) and C4W2 (i.e. S488C-Y518W) to match the available experimental data as much as possible. All simulations were performed using the GROMACS 2019 simulation package[66]. Each N_{TAIL} variant was simulated using two different force fields (CHARMM36m[51] and AMBER99SB-disp[49]) and at least three replicas per force field started from different initial conformations. The WT N_{TAIL} simulations were performed only with the CHARMM36m force field. For the Amber force field validation, we used previously published wild type N_{TAIL} simulation trajectories[26]. The simulations of the two variants in Amber was carried out using force field parameters adapted to the GROMACS simulation package. Both the WT trajectories and the force field parameters were kindly provided by Piana et al. Each simulation was carried out using periodic boundary conditions in a box filled with explicit solvent molecules consisting of either optimal point charge (OPC[55], for CHARMM36m) or a modified four point-charge (TIP4P[50] for AMBER99SB-disp) water models as well as Na+ and Cl- ions corresponding to an ion concentration of 150 mM. The total size of simulated systems was approximately 120,000 to 200,000 atoms. Total simulation time was more than 40 µs per variant with conformations recorded after every nanosecond. Hydrogen bond vibrations were constrained using virtual atom sites to enable a 4 fs time step during simulations. All simulations were kept at 1 atm pressure and 298 K temperature. Further details on the MD simulation parameters and

system preparation are provided in the Supporting Methods S7 and S9. Analysis methods related to the comparison to experimental data, calculation of mutual information, contact maps, and free-energy landscapes are described in the Supporting Methods S11, S19-21.

Data availability.

The implementation of the coarse-grained HPS model can be downloaded from https://github.com/wzhenglab/ntail. Other data supporting the findings of this study are available from the corresponding authors upon reasonable request.

Acknowledgements

L.O, J.K., G.K., S.L. and S.M.V. acknowledge the support from the National Institutes of Health (R01GM120537). A.C.V., H.G., L.V.B. and G.N. were supported by the Max Planck Society and by the German Science Foundation (DFG), Excellence Strategy Grant MBExC 2067/1; computer time was provided by the Max Planck Computing and Data Facility. G.N. was additionally supported by the Alexander von Humboldt-Foundation. A.C.V. was additionally supported by the European Union's Horizon 2020 Framework Programme for Research and Innovation, Specific Grant Agreement No. 945539 (Human Brain Project SGA3). W.Z. acknowledges the support from the National Institutes of Health (R35GM146814) and the research computing at Arizona State University.

References.

1. Gutsche, I. *et al.* Near-atomic cryo-EM structure of the helical measles virus nucleocapsid. *Science (1979)* **348**, 704 (2015).

2. Bloyet, L. M. The nucleocapsid of paramyxoviruses: Structure and function of an encapsidated template. *Viruses* **13**, 2465 (2021).

3. Longhi, S., Bloyet, L. M., Gianni, S. & Gerlier, D. How order and disorder within paramyxoviral nucleoproteins and phosphoproteins orchestrate the molecular interplay of transcription and replication. *Cellular and Molecular Life Sciences* **74**, 3091 (2017).

4. Karlin, D., Ferron, F., Canard, B. & Longhi, S. Structural disorder and modular organization in Paramyxovirinae N and P. *Journal of General Virology* **84**, 3239 (2003).

5. Kingston, R. L., Baase, W. A. & Gay, L. S. Characterization of Nucleocapsid Binding by the Measles Virus and Mumps Virus Phosphoproteins. *J Virol* **78**, 8630 (2004).

6. Longhi, S. *et al.* The C-terminal domain of the measles virus nucleoprotein is intrinsically disordered and folds upon binding to the C-terminal moiety of the phosphoprotein. *Journal of Biological Chemistry* **278**, 18638–18648 (2003).

7. Jensen, M. R. *et al.* Intrinsic disorder in measles virus nucleocapsids. *Proc Natl Acad Sci U S A* **108**, 9839–9844 (2011).

8. Johansson, K. *et al.* Crystal Structure of the Measles Virus Phosphoprotein Domain Responsible for the Induced Folding of the C-terminal Domain of the Nucleoprotein. *Journal of Biological Chemistry* **278**, 44567 (2003).

9. Kingston, R. L., Hamel, D. J., Gay, L. S., Dahlquist, F. W. & Matthews, B. W. Structural basis for the attachment of a paramyxoviral polymerase to its template. *Proc Natl Acad Sci U S A* **101**, 8301–8306 (2004).

10. Sourimant, J., Thakkar, V. D., Cox, R. M. & Plemper, R. K. Viral evolution identifies a regulatory interface between paramyxovirus polymerase complex and nucleocapsid that controls replication dynamics. *Sci Adv* **6**, eaaz1590 (2020).

11. Guseva, S., Milles, S., Blackledge, M. & Ruigrok, R. W. H. The nucleoprotein and phosphoprotein of measles virus. *Front Microbiol* **10**, 1832 (2019).

12. Gondelaud, F. *et al.* Functional benefit of structural disorder for the replication of measles, Nipah and Hendra viruses. *Essays Biochem* **66**, 915 (2022).

13. Bloyet, L. M. *et al.* Modulation of Re-initiation of Measles Virus Transcription at Intergenic Regions by PXD to NTAIL Binding Strength. *PLoS Pathog* **12**, e1006058 (2016).

14. Cox, R. & Plemper, R. K. The paramyxovirus polymerase complex as a target for next-generation anti-paramyxovirus therapeutics. *Frontiers in Microbiology* vol. 6 459 Preprint at https://doi.org/10.3389/fmicb.2015.00459 (2015).

15. Huang, W., Ravikumar, K. M., Chance, M. R. & Yang, S. Quantitative mapping of protein structure by hydroxyl radical footprinting-mediated structural mass spectrometry: A protection factor analysis. *Biophys J* **108**, 107–115 (2015).

16. Gely, S. *et al.* Solution structure of the C-terminal X domain of the measles virus phosphoprotein and interaction with the intrinsically disordered C-terminal domain of the nucleoprotein. *Journal of Molecular Recognition* **23**, 435 (2010).

17. Belle, V. *et al.* Mapping α-helical induced folding within the intrinsically disordered C-terminal domain of the measles virus nucleoprotein by site-directed spin-labeling EPR spectroscopy. *Proteins: Structure, Function and Genetics* **73**, 973 (2008).

18. Kavalenka, A. *et al.* Conformational analysis of the partially disordered measles virus N TAIL-XD complex by SDSL EPR spectroscopy. *Biophys J* **98**, 1055 (2010).

19. Wang, Y. *et al.* Multiscaled exploration of coupled folding and binding of an intrinsically disordered molecular recognition element in measles èirus nucleoprotein. *Proc Natl Acad Sci U S A* **110**, E3743 (2013).

20. Dosnon, M. *et al.* Demonstration of a folding after binding mechanism in the recognition between the measles virus NTAIL and X domains. *ACS Chem Biol* **10**, 795 (2015).

21. Bourhis, J.-M. *et al.* The intrinsically disordered C-terminal domain of the measles virus nucleoprotein interacts with the C-terminal domain of the phosphoprotein via two distinct sites and remains predominantly unfolded. *Protein Science* **14**, 1975 (2005).

22. Gruet, A. *et al.* Fuzzy regions in an intrinsically disordered protein impair protein-protein interactions. *FEBS Journal* **283**, 576–594 (2016).

23. Bignon, C., Troilo, F., Gianni, S. & Longhi, S. Modulation of measles virus NTAIL interactions through fuzziness and sequence features of disordered binding sites. *Biomolecules* **9**, 8 (2019).

24. Troilo, F., Bonetti, D., Bignon, C., Longhi, S. & Gianni, S. Understanding Intramolecular Crosstalk in an Intrinsically Disordered Protein. *ACS Chem Biol* **14**, 337 (2019).

25. Mukhopadhyay, R. & Hoh, J. H. AFM force measurements on microtubule-associated proteins: The projection domain exerts a long-range repulsive force. *FEBS Lett* **505**, 374 (2001).

26. Robustelli, P., Piana, S. & Shaw, D. E. Mechanism of Coupled Folding-upon-Binding of an Intrinsically Disordered Protein. *J Am Chem Soc* **142**, 11092–11101 (2020).

27. Gruet, A. *et al.* Dissecting partner recognition by an intrinsically disordered protein using descriptive random mutagenesis. *J Mol Biol* **425**, 3495 (2013).

28. Cai, L. & Zhou, H. X. Theory and simulation on the kinetics of protein-ligand binding coupled to conformational change. *Journal of Chemical Physics* **134**, 105101 (2011).

29. White, J. T., Li, J., Grasso, E., Wrabl, J. O. & Hilser, V. J. Ensemble allosteric model: Energetic frustration within the intrinsically disordered glucocorticoid receptor. *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**, 20170175 (2018).

30. Berlow, R. B., Dyson, H. J. & Wright, P. E. Hypersensitive termination of the hypoxic response by a disordered protein switch. *Nature* **543**, 447 (2017).

31. Chowdhury, A., Nettels, D. & Schuler, B. Interaction Dynamics of Intrinsically Disordered Proteins from Single-Molecule Spectroscopy. *Annu Rev Biophys* **52**, 433–462 (2023).

32. Staby, L. *et al.* Flanking Disorder of the Folded αα-Hub Domain from Radical Induced Cell Death1 Affects Transcription Factor Binding by Ensemble Redistribution: Flanking disorder tunes the RCD1-RST ensemble. *J Mol Biol* **433**, 167320 (2021).

33. Peng, Y. *et al.* A Metastable Contact and Structural Disorder in the Estrogen Receptor Transactivation Domain. *Structure* **27**, 229–240 (2019).

34. Buscaglia, M., Schuler, B., Lapidus, L. J., Eaton, W. A. & Hofrichter, J. Kinetics of intramolecular contact formation in a denatured protein. *J Mol Biol* **332**, (2003).

35. Vaiana, S., Best, R. B., Yau, W.-M., Eaton, W. A. & Hofrichter, J. Evidence for a partially structured state of the amylin monomer. *Biophys. J.* **97**, 2948–2957 (2009).

36. Sizemore, S. M., Cope, S. M., Roy, A., Ghirlanda, G. & Vaiana, S. M. Slow Internal Dynamics and Charge Expansion in the Disordered Protein CGRP: A Comparison with Amylin. *Biophys J* **109**, 1038 (2015).

37. Lapidus, L. J., Eaton, W. A. & Hofrichter, J. Measuring the rate of intramolecular contact formation in polypeptides. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 7220–7225 (2000).

38. Schuler, B. & Hofmann, H. Single-molecule spectroscopy of protein folding dynamics—expanding scope and timescales. *Curr. Opin. Struct. Biol.* **23**, 36–47 (2013).

39. Flory, P. J. *Principles of Polymer Chemistry. Principles of Polymer Chemistry* (Cornell University Press, 1953).

40. Zheng, W. *et al.* Inferring properties of disordered chains from FRET transfer efficiencies. *Journal of Chemical Physics* **148**, 123329 (2018).

41. Sawle, L. & Ghosh, K. A theoretical method to compute sequence dependent configurational properties in charged polymers and proteins. *J. Chem. Phys.* **143**, 85101 (2015).

42. Das, R. K. & Pappu, R. V. Conformations of intrinsically disordered proteins are influenced by linear sequence distributions of oppositely charged residues. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 13392–13397 (2013).

43. Dignon, G. L., Zheng, W., Kim, Y. C., Best, R. B. & Mittal, J. Sequence determinants of protein phase behavior from a coarse-grained model. *PLoS Comput. Biol.* **14**, e1005941 (2018).

44. Debye, P. & Hückel, E. De la theorie des electrolytes. I. abaissement du point de congelation et phenomenes associes. *Physikalische Zeitschrift* **24**, 185–206 (1923).

45. Nagy, G., Igaev, M., Jones, N. C., Hoffmann, S. V. & Grubmüller, H. SESCA: Predicting Circular Dichroism Spectra from Protein Molecular Structures. *J Chem Theory Comput* **15**, 5087 (2019).

46. Fossat, M. J. & Pappu, R. V. Q-Canonical monte carlo sampling for modeling the linkage between charge regulation and conformational equilibria of peptides. *Journal of Physical Chemistry B* **123**, 6952 (2019).

47. Zheng, W. *et al.* Molecular Details of Protein Condensates Probed by Microsecond Long Atomistic Simulations. *Journal of Physical Chemistry B* **124**, 11671 (2020).

48. Best, R. B., Zheng, W. & Mittal, J. Balanced protein-water interactions improve properties of disordered proteins and non-specific protein association. *J. Chem. Theor. Comput.* **10**, 5113–5124 (2014).

49. Robustelli, P., Piana, S. & Shaw, D. E. Developing a molecular dynamics force field for both folded and disordered protein states. *Proc. Natl. Acad. Sci. U.S.A.* **115**, E4758–E4766 (2018).

50. Piana, S., Donchev, A. G., Robustelli, P. & Shaw, D. E. Water dispersion interactions strongly influence simulated structural properties of disordered protein states. *J. Phys. Chem. B* **119**, 5113–5123 (2015).

51. Huang, J. *et al.* CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nat Methods* **14**, 71 (2017).

52. Henriques, J. & Skepö, M. Molecular dynamics simulations of intrinsically disordered proteins: On the accuracy of the TIP4P-D water model and the representativeness of protein disorder models. *J Chem Theory Comput* **12**, 3407–3415 (2016).

53. Svergun, D., Barberato, C. & Koch, M. H. J. CRYSOL - a program to evaluate X-ray solution scattering of biological molecules from atomic coordinates. *J. Appl. Cryst.* **28**, 768–773 (1995).

54. Shen, Y. & Bax, A. SPARTA+: a modest improvement in empirical NMR chemical shift prediction by means of an artificial neural network. *J. Biomol. NMR* **48**, 13–22 (2010).

55. Shabane, P. S., Izadi, S. & Onufriev, A. V. General Purpose Water Model Can Improve Atomistic Simulations of Intrinsically Disordered Proteins. *J Chem Theory Comput* **15**, 2620 (2019).

56. Hopf, T. A. *et al.* The EVcouplings Python framework for coevolutionary sequence analysis. *Bioinformatics* **35**, 1582–1584 (2019).

57. Bignon, C., Troilo, F., Gianni, S. & Longhi, S. Partner-Mediated Polymorphism of an Intrinsically Disordered Protein. *J Mol Biol* **430**, 2493–2507 (2018).

58. Milles, S. *et al.* An ultraweak interaction in the intrinsically disordered replication machinery is essential for measles virus function. *Sci Adv* **4**, eaat7778 (2018).

59. Krumm, S. A., Takeda, M. & Plemper, R. K. The measles virus nucleocapsid protein tail domain is dispensable for viral polymerase recruitment and activity. *Journal of Biological Chemistry* **288**, 29943 (2013).

60. Guseva, S. *et al.* Measles virus nucleo- and phosphoproteins form liquid-like phase-separated compartments that promote nucleocapsid assembly. *Sci Adv* **6**, eaaz7095 (2020).

61. Zhou, Y., Su, J. M., Samuel, C. E. & Ma, D. Measles Virus Forms Inclusion Bodies with Properties of Liquid Organelles. *J Virol* **93**, e00948-19 (2019).

62. Kolakofsky, D., Le Mercier, P., Iseni, F. & Garcin, D. Viral RNA polymerase scanning and the gymnastics of Sendai virus RNA synthesis. *Virology* **318**, 463 (2004).

63. Brunel, J. *et al.* Sequence of Events in Measles Virus Replication: Role of Phosphoprotein-Nucleocapsid Interactions. *J Virol* **88**, 10851 (2014).

64. Bugge, K. *et al.* Interactions by Disorder – A Matter of Context. *Front Mol Biosci* **7**, 110 (2020).

65. Anderson, J. A., Glaser, J. & Glotzer, S. C. HOOMD-blue: A Python package for high-performance molecular dynamics and hard particle Monte Carlo simulations. *Comput Mater Sci* **173**, 109363 (2020).

66. Abraham, M. J. *et al.* Gromacs: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **1**, 19 (2015).

47