1

Patterns of Ongoing Thought in the Real World and Their Links to Mental Health and

Well-Being

Bridget Mulholland[1*], Louis Chitiz[1], Raven Wallace[1], Brontë Mckeown[2], Michael Milham[3,4],

Arno Klein[5], Robert Leech[6], Elizabeth Jefferies[7], Giulia Poerio[8], Jeremy Stewart[1], Jeffery

Wammes[1], Jonathan Smallwood[1]

[1]Department of Psychology, Queen's University, Kingston, Ontario, Canada

[2]National Institute of Teaching, London, England

[3]Center for the Developing Brain, Child Mind Institute, New York City, New York, United
States of America

[4]Nathan S. Kline Institute for Psychiatric Research, Orangeburg, New York, United States of
America

[5]Child Mind Institute, New York City, New York, United States of America

[6]Department of Neuroimaging, King's College, England

[7]Department of Psychology, University of York, York, England

[8]School of Psychology, University of Sussex, Falmer, Brighton, England

[*]Corresponding author
E-mail: 15bbm@queensu.ca

Abstract

The thoughts we experience in daily life have implications for our mental health and

well-being. However, it is often difficult to measure thought patterns outside of laboratory

conditions due to concerns about the voracity of measurements taken in daily life. To address

this gap in the literature, our study set out to measure patterns of thought as they occur in daily

life and assess the robustness of these measures and their associations with trait measurements of

mental health and well-being. A sample of undergraduate participants completed multi-

dimensional experience sampling surveys eight times per day for five days as they went around

their normal lives. Principal component analysis reduced these data to identify the dimensions

that explained the patterns of thought reported by our participants. We used linear modelling to

map how these thought patterns related to the activities taking place at the time of the probe,

highlighting good consistency within the sample, as well as substantial overlap with prior work.

Multiple regression was used to examine associations between patterns of ongoing thought and

aspects of mental health and well-being, highlighting a pattern of "Intrusive Distraction" that had

a positive association with anxiety, and a negative association with social well-being. Notably,

this thought pattern tended to be most prevalent in solo activities and was relatively suppressed

when interacting with other people (either in person or virtually). Our study, therefore, highlights

the use of multi-dimensional experience sampling as a tool to understand how ongoing thoughts

in daily life impact on our mental health and well-being, and establishes the important role social

connectedness plays in the etiology of intrusive thinking.

Introduction

A fundamental goal of psychology is to understand how the thoughts we experience in daily life relate to our productivity, mental health, and well-being. Contemporary work indicates a link between patterns of ongoing thought, as determined by experience sampling, and various aspects of health and well-being (e.g., [1, 2]). However, research on human thought has historically relied on observations made in controlled laboratory contexts such as brain imaging (e.g., [3]) or behavioral laboratories (e.g., [4]) because of the assumption that the robustness of these measures is improved when they are gathered in controlled conditions. However, recent work in the laboratory and in daily life has emphasized that patterns of thought are intimately related to the context in which they emerge [5]. Since the tasks people perform in the lab are unlikely to perfectly match the activities people perform in daily life, the role that context plays in influencing thought patterns may be one reason why links between traits and patterns often do not generalize from the laboratory to daily life [6].

The influence that context has on thought patterns has been established both within the laboratory and in daily life (for a review see [7]). For example, Konu and colleagues [8] used a battery of laboratory-based activities, including cognitive and attentional tasks, videos, and audiobooks to establish that the patterns of thought people engaged in changed as a function of the tasks people engage in. In a similar vein, Mulholland and colleagues [9] established that in daily life, the thought patterns participants reported varied with the activities people engaged in. There were also similarities in how the ongoing context influenced the patterns that participants reported. In both studies, patterns of thought with (i) unpleasant, intrusive, and distracting features were identified that dominated low-demanding situations in both the lab and daily life, (ii) thoughts with social and episodic features were important in social situations in daily life and

tasks relying on social cognition in the lab, and (iii) patterns of detailed focus on a task were

important during homework and while working in daily life, and were present during demanding

laboratory task such as those depending on working memory. These prior lab and daily life

studies [8, 9] took advantage of a novel approach to measuring cognition known as multi-

dimensional experience sampling (mDES, [10]). This approach asks participants to rate their

experience along a number of dimensions (e.g. how detailed they are, whether are they intrusive,

whether they involve other people, etc.…) on multiple occasions. These data are then often

decomposed into a set of underlying dimensions that describe the patterns of answers that the

participants provided. This dimension derived from this decomposition provides a 'thought

space' in which different experiential moments can be located [11-13]).

Emerging evidence suggests that one reason why mDES can characterize contextual

influences on ongoing thought is because it provides high precision descriptions of individuals'

experiences. For example, states identified using mDES are associated with brain activity during

attention tasks, where patterns off-task thought with episodic features are related to greater

activity within the medial prefrontal cortex, a region of the default mode network [14]. In

contrast, patterns of detailed focus relating to a task are linked to greater activity in the

frontoparietal network [12]. mDES is also sensitive to changes in brain activity emerging during

movie watching [11], highlighting that patterns of intrusive distraction are related to moments

during a film when frontoparietal activity is reduced [11], a pattern consistent with the notion

that intrusive thought reflects a failure to control cognition [15]. Recently, we demonstrated that

mDES is sufficiently sensitive to variation in thought patterns to the extent that it can be used to

build a fully generative model of the mapping between thought patterns and brain activity during

task states [13].

Current Study

Together, these prior studies demonstrate that mDES provides a powerful tool for mapping

thinking patterns in the lab and daily life. Thus, our goal was to understand how the thought

patterns mDES reveals in daily life relate to measures of mental health and well-being. Prior

studies have already used experience sampling in daily life and in the laboratory to highlight

important links between patterns of ongoing thought and factors that impact mental health and

well-being, such as depression [16, 17], anxiety [18], and obsessional thinking [19]. The goal of

the current study was to leverage the precision mapping of experience sampling offered by

mDES to understand the links between thought patterns in daily life and mental health and well-

being. However, before examining links to mental health and well-being, we first investigated

the reliability of the thought patterns produced using mDES. We took advantage of the fact that

the thought patterns established by mDES are sensitive to the activities being performed in daily

life [9] and examined whether the mappings between thought patterns and activities are

consistent within the current dataset and also with those seen in our prior study [9]. Once we

established the reliability of mDES as a tool for mapping thought patterns in daily life, we

examined how the thought patterns established by our approach related to aspects of mental

health and well-being as measured by a battery of questionnaires.

Participant Population

    A total of 261 participants (women = 227, men = 29, non-binary or similar gender identity = 4, prefer not to say = 1; M = 21.44; SD = 6.16; range = 17-52; note, two ages were recorded incorrectly and therefore not included and two additional ages were absent) were included in this study. This study was granted ethics clearance by the Queen's University General Research Ethics Board. Participants were recruited between January 9th, 2023, and April 10[th], 2023, through the Queen's University Psychology Participant Pool. Eligible participants were Queen's University students enrolled in designated first- and second-year psychology courses. Participants gave informed, written consent through electronic documentation prior to taking part in any research activities and were awarded two course credits and fully debriefed upon the completion of their participation.

Procedure

    Participants completed demographic questionnaires (i.e., age, gender, gender identity, sex assigned at birth, language(s) spoken most often at home, country of primary place of residence, program, and year of study). Participants also completed evidence-based health and wellness questionnaires, including the Quick Inventory of Depressive Symptomatology self-report (QIDS-SR_{16}) [20], the Anxiety Sensitivity Index (ASI) [21], the Autism Spectrum Quotient (AQ) [22] to assess for autistic traits, the Adult ADHD Self-Report Scale (ASRS) [23] to asses for ADHD traits, and the Montgomery Asberg Depression Rating Scale self-report (MADRS-S) [24] to measure severity of depression symptomatology and the Overall Anxiety Severity and Impairment Scale (OASIS) [25] to measure severity of anxiety symptomatology, the MOS 36-item short form survey (SF-36) [26], the Sheehan Disability Scale (SDS) [27] to assess

functional impairment, and the World Health Organization Quality-of-Life Brief Version

(WHOQOL-BREF) [28] to quantify quality-of-life. These questionnaires were selected based on

a widespread, general approach to health and wellness, rather than clinical diagnosis. For the

purpose of this study, our analysis was limited to the AQ, ASRS, MADRS, OASIS, and

WHOQOL-BREF questionnaires which were deemed the most relevant indicators of mental

health and well-being. Once participants completed these questionnaires, they were notified via

Samply Research, a mobile application, to complete mDES questionnaires eight times daily for

five consecutive days between 7:00 am and 11:00 pm. Each questionnaire was randomly

delivered within this 16-hour time window, with a minimum of 30 minutes in between each

notification.

Multi-dimensional Experience Sampling

Participants received notifications on their phones through the Samply Research mobile

application. All responses were made in reference to their thoughts, feelings, environment,

location, and activities immediately before receiving the notification. This study used a 16-

question mDES battery that has been used in prior studies [11, 13] (see Table 1). The mDES

questions were always asked first and delivered in a random order. Participants then rated their

stress on a 1-to-10 Likert scale. For the purpose of this study, responses to the stress question

were not analyzed. Participants then answered questions about their physical and virtual social

environment (Table 2). Finally, participants also indicated the type of physical location they

were in (Table 3) and their primary activity (Table 4). The activity list was developed from the

day reconstruction method [29] and modified based on the options in our prior studies [9, 30].

 Analysis

 Data and Code Availability Statement

All custom code used to prepare data for analysis and figure development is available

online at https://github.com/ThinCLabQueens and https://github.com/BridgMul10. Anonymized

data has been uploaded to a publicly accessible database, Mendeley Data, and is available online

at https://doi.org/10.17632/gfhwg3p8r6.1.

Principal Component Analysis: mDES Questions

As in our prior studies (e.g. [8, 9, 12, 14, 30]), common patterns of thought (Figure 1)

were identified by applying principal component analysis (PCA) using Varimax rotation to the

data generated from responses to the 16 mDES questions (Table 1) using IBM SPSS Statistics

(version 29). PCA was applied at the observation level and included 6776 observations. The

loadings from the five components with an eigenvalue > 1 were retained for further analysis (see

Table 5 and Figure 1).

Reliability Analysis: mDES Question Data

Component reliability analysis was conducted in IBM SPSS Statistics (version 29).

mDES thought dimension data was randomly shuffled and divided into two halves (n = 3445

probes per sample). PCA (varimax) was applied to each subset separately and Pearson

correlations were run to compare the component loadings generated from each subset with the

overall solution. The higher the correlation between the two halves of the data to the overall

sample, the more consistent the dimensional structure within the overall sample was.

Linear Mixed Modeling: Generating "Thought-Activity" Mappings

To analyze contextual distributions of thought in relation to activities, we conducted a

series of linear mixed models (LMMs), one with each thought component as the dependent

variable and activity as the independent variable, to examine whether patterns of thought varied

across activity categories. Restricted maximum likelihood (REML) was used as the estimation

method and a variance components model was used as the covariance type. Participants were

included as a random intercept. The parameter estimates for each activity in each model were

saved for the eventual generation of activity word clouds that describe the experiential features

that contribute to each dimension (Figure 2). This analysis is identical to that found in

Mulholland et al. [9]. The parameter estimates were also saved to be used in reliability analysis

(Figure 3).

Reliability Analysis: Within Data Set Consistency of Thought-Activity Mappings

202   Primary activity data reliability analysis was conducted in IBM SPSS Statistics (version

29). As before, component scores were split into two subsets ($n = 3445$ probes per subset). We

conducted a series of LMMs, with thought components for each subset as the dependent variable

and activity as the independent variable. Restricted maximum likelihood (REML) was used as

the estimation method and a variance component model was used as the covariance type.

Participants were included as a random intercept. Pearson correlations were run on the parameter

estimates generated from each subset for each activity. This analysis allowed us to estimate

whether the results of our whole sample LMM are generalizable to sub-samples of the data.

Reliability Analysis: Comparing Thought-Activity Mappings Across Studies

211   Reliability analysis was conducted in IBM SPSS Statistics (version 29). The LMM

parameter estimates from Mulholland et al. [31] contained a sample of $n = 1451$ probes, and the

LMM parameter estimates from our current study contained a sample of $n = 6889$ probes.

Pearson correlations were run on the parameter estimates for overlapping activity from each

dataset. This analysis allowed us to estimate whether the results of our whole sample LMM in

the current data are generalizable to those seen in the prior dataset.

Linear Mixed Modeling: Social Environment Data

To analyze contextual distributions of thought in relation to socialization, we conducted a

series of LMMs, one with each thought component as the dependent variable and social

environment as the independent variable (i.e., physical environment with people, physical

environment with a pet, or virtual environment with people were all options as independent

variables). These analyses allowed us to examine whether patterns of thought varied

meaningfully across social environments. Restricted maximum likelihood (REML) was used as

the estimation method, and a variance components model was used as the covariance type.

Participants were included as a random intercept.

Multiple Regression: Linking mDES to Traits

To analyze whether mDES questions can differentiate individuals based on underlying

traits, we performed a series of multiple regressions with the average PCA component scores for

each individual in the overall "Thought Space" as the dependent variable and trait questionnaires

as explanatory variables. Trait questionnaires included in this analysis include the AQ, ASRS,

MADRS, OASIS, and WHOQOL-BREF, all of which were z-scored. The WHOQOL-BREF

questionnaire was subdivided into four categories based on scoring domains: physical health,

psychological health, social relationships, and environment. mDES questionnaire data was

reduced to a single score for each participant.

 Results

Patterns of Ongoing Thought

First, mean scores for each thought dimension measured were calculated and are shown

in the bar plots in Figure 1. Next, as is standard in our laboratory work (e.g. [8, 12, 14, 30]) the

mDES data was decomposed using PCA (varimax rotation) to reveal the dimensions that best

described the patterns of ongoing thought reported by the participants. Based on eigenvalue > 1,

five components were selected for further analysis (see Figure 1 for scree plot). PCA loadings

(Table 5) from the five components were used to generate thought word clouds (Figure 1).

Components were named based on mDES thought dimensions that dominated their composition.

Component 1 (21.5% of variance) was labelled "Detailed Task Focus" because loadings were

high for "detailed," "deliberate," "problem," and "task" (Figure 1). Component 2 (13.6% of

variance) was labelled "Intrusive Distraction" because loadings were high for "(negative)

emotion," "intrusive," and "distracting" (Figure 1). Component 3 (8.6% of variance) was

labelled "Episodic Social Cognition" because loadings were high for dimensions such as

"knowledge," "person," and "past" (Figure 1). Component 4 (6.6% of variance) was labelled

"Future Problem-Solving" because loadings were high for dimensions such as "future," "self,"

and "problem" [as in problem solving] (Figure 1). Component 5 (6.3% of variance) was labelled

"Sensory Engagement" because loadings were high for dimensions such as "images," and

"sounds" (Figure 1). Please note that these terms are used for convenience to summarize the

features that characterize each component. They do not constitute the only labels which could be

applied to these patterns.

The Within Dataset Consistency of the Dimensional Structure of Ongoing Experience in Daily Life

Note. Upper horizontal panel: (Left) Bar graph describing the mean mDES score for each dimension. Error bars

represent 99% confidence intervals (CIs). (Right) Scree plot generated from PCA of mDES questionnaire data.

Lower horizontal panel: Word clouds describing the experiential features that contribute the most to the dimensions

of experience as described by PCA (varimax rotation). Larger fonts describe features with more importance on the

dimension (i.e., stronger loadings) and the font colour denotes direction of the loading (i.e., warm colours relate to

positive loadings). For the purpose of exposition, these components are named based on their strongest features

(from left to right): "Detailed Task Focus," "Intrusive Distraction," "Episodic Social," "Future Problem-Solving,"

and "Sensory Engagement." Scatter plots below show the split half reliability of these dimensions (Y-axis: Subset 1;

X-axis: Subset 2). Left vertical panel: Illustrations of different activities in daily life.

Component Reliability

To understand the robustness of the dimensions within our sample, we conducted a split-

half reliability analysis. In this analysis, the data was divided into two random samples (n = 3445

probes per sample) and then examined to determine how the components generated in each half

of the data related to each other. We used the robustness of the solutions across PCAs with 3-, 4-,

and 5-component solutions as a complementary method to determine the best solution for the

entire sample. The mean correlation for the set of the homologous pairs from each solution was

calculated with a higher score reflecting the most reproducible components. The 3-, 4-, and 5-

component solutions all produced highly reliable results, with average homologue similarities

scores of $r = .996$ (3-component solution, range $r = .994$-$.995$, Figure 1), $r = .93525$ (4-

component solution, range $r = .869$-$.978$, Figure 1), and $r = 0.9772$ (5-component solution,

range $r = .957$-$.991$, Figure 1). Since the similarity was high for 3-, 4-, and 5-component

solutions, we focused our analysis on the 5-component solution.

Thought-Activity Mappings

Having identified the stability of the dimensions identified within this dataset, we

examined if mDES provides a reliable way to map cognition in daily life onto activities that are

being performed. To this end, we ran a series of LMMs in which the activities are the

explanatory variables and the mDES scores for each dimension from the common "Thought-

Space" are the dependent variables. In each case we found a significant association between

reported patterns of thought and activities ("Detailed Task Focus" ($F(21, 6639.50) = 74.08$, p

$<.001$); "Intrusive Distraction" ($F(21, 6591.47) = 10.21$, $p <.001$); "Episodic Social Cognition"

($F(21, 6632.08) = 19.08$, $p <.001$); "Future Problem-Solving" ($F(21, 6648.67) = 18.26$, p

$<.001$); "Sensory Engagement" ($F(21, 6559.04) = 16.82$, $p <.001$)). To visualize these thought-

activity mappings we generated a set of word clouds based on the estimated marginal means for

reported activities in each component (Figure 2). "Detailed Task Focus" was high when doing

homework and working (paid or volunteer) and lowest when napping or resting, watching TV,

and doing nothing or waiting. "Intrusive Distraction" was highest when doing nothing or

waiting, napping or resting, and doing homework, and lowest during physical leisure or sports,

and shopping. "Episodic Social" features were highest when talking on the phone, messaging by

phone/device, video-calling, walking the dog, talking in person, traveling or commuting, and

physical leisure or sports, and lowest when watching TV, napping or resting, using a computer or

electronic device, and doing homework. "Future Problem-Solving" was higher during hygiene

activities, texting, commuting, phone-call, exercise, and shopping, and lowest when watching

TV, reading, and doing homework. Lastly, "Sensory Engagement" had high loadings when

watching TV, listening to music, playing sports, and using a computer, and low loadings when

doing homework, talking on the phone, and working.

 Figure 2

Consistency of Thought-Activity Mappings

Note. First row: "Thought Patterns." Words represent PCA (varimax) loadings for mDES data and LMM loadings

for primary activities. Larger fonts are features with more importance (i.e., higher loadings) and colour denotes

features with a similar loading on the dimensions (i.e., warm colours relate to positive loadings). See Table 5-10 for

specific component loadings. Second row: "Activity Loadings." In these word clouds the large the font the stronger

this activity loads on that thought component and activities with a similar colour load in the same direction.

Third row: Scatter plots showing the within study consistency of the thought-activity mappings. (Y-axis: Activity

loadings from Subset 1; X-axis: Activity Loadings from Subset 2). Fourth row: Scatter plots showing the between

study consistency of the thought-activity mappings (Y-axis: Activity loadings from Mulholland et al. [7]; X-axis:

Activity loadings in the current dataset).

Next, we investigated the robustness of the thought-activity mappings produced by

mDES. Our first analysis examined the consistency of their mappings within the current data by

performing a split-half reliability analysis with current data. We created two subsets of our data

(n = 3445 probes per subset). In each case, we found a significant association between each of

the reported thought patterns and activities in each subset ("Detailed Task Focus" subset 1 ($F(21,$

$3284.10) = 36.96, p <.001$) and subset 2 ($F(21, 3333.340) = 36.101, p <.001$); "Intrusive

Distraction" subset 1 ($F(21, 3274.05) = 4.63, p <.001$) and subset 2 ($F(21, 3285.57) = 6.53, p$

$<.001$); "Episodic Social Cognition" subset 1 ($F(21, 3323.14) = 8.92, p <.001$) and subset 2

($F(21, 3271.06) = 10.32, p <.001$); "Future Problem-Solving" ($F(21, 3285.56) = 10.02, p <.001$)

and subset 2 ($F(21, 3294.73) = 9.06, p <.001$); "Sensory Engagement" subset 1 ($F(21, 3242.37)$

$= 9.69, p <.001$) and subset 2 ($F(21, 3239.06) = 8.13, p <.001$)).

To understand whether the thought-activity mappings in daily life were similar for each

subset of the data the loadings of each activity on each component each component was

correlated (see Methods). Correlations between the LMM activity estimated marginal means in

each subset ranged from $r = .60$-$.91$, with the highest reproducibility for the "Detail Task Focus"

component and the lowest for the "Sensory Engagement" component. This shows a reasonably

high degree of consistency in the feature loadings for each dimension of experience within the

current sample.

Next, we examined how the thought-activity mappings from the current study map onto

those from the homologous components from Mulholland et al. [9]. To this end, we used Pearson

Correlation to compare the activity mappings from our prior study to the mappings found in our

omnibus sample. Mulholland et al. [9] did not include a "Sensory Engagement" component

(because the modality questions did not include sounds), so only four components were

compared. This analysis identified correlations with a range of $r = .67$-$.86$, with the strongest

mappings for "Detailed Task Focus" and the lowest for "Future Problem-Solving." This analysis

shows a reasonably high degree of overlap between the thought activity mappings as described in

these data and those seen in our prior study [9].

One feature of our analysis is that it allows us to understand experience within each

activity as a linear combination of different thought components which can be represented

graphically by an activities location in a multivariate "Thought Space." To visualize the data this

way, the estimated marginal means for each activity derived from the LMMs and plotted against

each mDES components for the common "Thought-Space." We present these data a three-

dimensional space constructed using the three most reliable components (i.e., "Detailed Task

Focus," "Intrusive Distraction," and "Episodic Social Cognition") as the dimensions. We also

present a two-dimensional space that describes the relationship between activities in terms of

their weighting for "Future Problem-Solving" and "Sensory Engagement." These are both

presented in (, which show how certain activities occupy extreme values on multiple

components. For example, "homework" is high on both the "Detailed Task Focus" and

"Intrusive Distraction" components but low on "Episodic Social Cognition," "Future Problem-

Solving," and "Sensory Engagement."

 Figure 3

The Multivariate Landscape of Thought-Activity Mappings in Daily Life

Note. The thought-activity mapping revealed by our analysis are presented in two- and three-dimensional spaces to

provide the opportunity to visualize the multivariate nature of the relationships. Left image: Thought-activity

mappings in a three-dimensional "Thought Space." Each point is an activity. X-axis is the activity loadings for

"Detailed Task Focus", the Y-axis is the activity loading on "Intrusive Distraction," and the Z-axis are the activity

loadings for "Episodic Social Cognition." Right image: Thought-activity mappings in a two-dimensional thought

space. The Y-axis is the activity loading on "Sensory Engagement" and the X-axis as the activity loadings on

"Future Problem-Solving." Lower panel: This shows the feature loadings for each dimension of the thought space in

the form of word clouds. As described previously, larger fonts are activities with more importance (i.e., higher

loadings on a dimension) and words in a similar colour denotes direction (i.e., warm colours relate to positive

loadings).

Thought Dimensions and Social Environments

Another way to examine the robustness of mDES as a tool for mapping cognition in daily

life by measuring the similarity in how the thought patterns map onto social environments in

daily life. Prior studies (e.g. [9]) have found that being alone compared to being with other

people have important implications for a persons' thought patterns, in particular facilitating

patterns of thought with social and episodic features. We ran a series of LMMs in which

different social environments (Table 2) were the explanatory variable and the mDES scores from

the common "Thought Space" the dependent variables. We found a significant association

between reported patterns of thought and the physical social environments for each component.

"Detailed Task Focus" ($F(3, 6721.87) = 16.06$, $p <.001$) was lowest when with people and

interacting with them ($M = -.08$, 95% CI [-.08, -.13]) and highest when around people but not

interacting with them ($M = .14$, 95% CI [.08,-.21]). "Intrusive Distraction" ($F(3, 6656.39) =$

37.54, $p <.001$) was lowest when with people and interacting ($M = -.15$, 95% CI [-.22, -.08]) and

highest when alone ($M = .11$, 95% CI [.05,.18]). "Episodic Social Cognition" ($F(3, 6699.28) =$

56.29, $p <.001$) was highest when around people and interacting ($M = .22$, 95% CI [.16,.28]) and

lowest when around people but not interacting ($M = -.15$, 95% CI [-.08,-.22]). "Future Problem-

Solving" ($F(3, 6716.79) = 2.79$, $p = .039$) was lowest when around people and interacting with

them ($M = -0.46$, 95% CI [-.11,.02]) and highest when around people but not interacting ($M =$

$.04$, 95% CI [-.02,.15]). Finally, "Sensory Engagement" ($F(3, 6619.47) = 5.30$, $p = .001$) was

highest when alone ($M = .03$, 95% CI [-.08,.02]). A broadly similar pattern was found for our

analysis of the consequence of virtual environments. "Detailed Task Focus" ($F(3, 6705.72) =$

2.67, $p = .046$) was higher when around people and not interacting ($M = .05$, 95% CI [-.03,.14])

and lowest when alone ($M = -.03$, 95% CI [-.83,.02]). "Intrusive Distraction" ($F(3, 6695.70) =$

11.53, $p <.001$) was least prevalent when participants were around people and interacting

virtually ($M = -.11$, 95% CI [-.19,-.03]) and highest when around people but not interacting

virtually ($M = .07$, 95% CI [-.02, .16]). "Episodic Social Cognition" ($F(3, 6715.54) = 39.37$, p

424 <.001) was highest when around people and interacting virtually ($M = .27$, 95% CI [.19,.34]) and

425 lowest when alone ($M = -.07$, 95% CI [.19,.33]). "Future Problem-Solving" ($F(3, 6709.60)$ 9.55,

426 $p <.001$) was highest when around people but not interacting ($M = .15$, 95% CI [.07,.22]) and

427 lowest when around people but not interacting virtually with another person ($M = -.03$, 95% CI [-

428 .12,.05]). Finally, "Sensory Engagement" ($F(3, 6663.16) = 2.89$, $p = .03$) was highest when alone

429 ($M = .01$, 95% CI [-.05,.08]) and lowest around people but not interacting virtually ($M = -.003$,

430 95% CI [-.09,.09]). These analyses replicate prior observations that found a mapping of social

431 cognition onto social situations (e.g. [9, 30]) but extends this to establish that such situations also

432 reduce intrusive features of cognition.

433 **Thought Dimensions and Traits**

434   Having established a high level of consistency between the thought-activity mappings

435 seen in the current study and prior studies [9, 30], the final goal of our study was to investigate

436 whether mDES dimensions can differentiate people based on traits associated with mental health

437 and well-being. To do so, we performed a series of multiple regressions with the average loading

438 of each individual on each dimension of the common "Thought Space" as the dependent

439 variables and their score on the trait questionnaires as explanatory variables. Results indicated

440 the location of each individual on the "Intrusive Distraction" dimension ($F(8, 252) = 8.35$, $p =$

441 <.001; see Table 11 for results for each component) could be predicted based on high levels of

442 anxiety (Beta = 0.23, $p = .003$) and low levels of social well-being (Beta = -0.17, $p =.010$). This

443 analysis, therefore, shows a clear mapping between patterns of "Intrusive Distraction" with

444 higher levels of anxiety and lower social well-being (Figure 4).

Figure 4

mDES Questionnaire Data and Trait Data Multiple Regression Results

Note. Left panel: Scatterplot with WHOQOL-BREF social relationships (z-scored) on the y-axis and "Intrusive

Distraction" PCA component scores on the x-axis. Right panel: Scatterplot with OASIS (z-scored) on the y-axis and

"Intrusive Distraction" PCA component scores on the x-axis. For both scatterplots, all values were mean z-scored

per participant.

Discussion

Our study sought to investigate the underlying relationships between patterns of reported

thought, behavior, and measures of mental health in daily life. First, we investigated the stability

of the patterns of thoughts reported in our study, establishing a reasonable range of consistency

for a five-dimensional solution. These dimensions were "Detailed Task Focus," "Intrusive

Distraction," "Episodic Social," "Future Problem-solving," and "Sensory Engagement." Second,

we examined the reproducibility of how activities in daily life map onto these dimensions. The

thought-activity mappings were both consistent within these data and were similar to the thought

patterns described by Mulholland et al. [9] (Figure 2), who administered mDES in daily life to a

different sample of participants. We also replicated patterns in prior studies that show, for

example, that patterns of "Episodic Social Cognition" are higher when individuals are interacting

either physically or virtually observed in both prior studies [9], a pattern of thought that was also

reduced during COVID lockdowns in the United Kingdom because of reductions in the

opportunities for socialization [30]. Taken together these analyses show that the "Thought

Spaces" generated by the application of mDES to daily life provide a reproducible measure

ongoing thought and that these organize the activities we engage in as we go about our daily

lives in a meaningful manner (see Figure 3). It is important to note that populations in both

studies are undergraduate students, so this reproducibility highlights that if mDES is employed

within a similar group of individuals, then it captures thought patterns that are reasonably

consistent in their profile and their associations with activities. Further studies are needed to

understand how thought-activity mappings vary across cultures. However, it is possible that the

observed thought patterns, as well as their links to activities or social contexts, could conceivably

change if they were measured in different cultures or age groups.

Links Between Thought Patterns in Daily Life and Psychological Well-Being

The main goal of the study was to understand the links between patterns of thought in

daily life and mental health. We found that patterns of thought that loaded heavily on "Intrusive

Distraction" were associated with greater levels of anxiety and less social support (Figure 4).

Importantly this thought pattern showed a reasonable degree of consistency in terms of how it

mapped onto activities within this sample ($r = .77$) and with our prior study ($r = .73$) highlighting

that this thought component has a reasonably consistent thought-activity mapping.

Contemporary perspectives that emphasize patterns of thought often have an important

social component (e.g., [32]). One important contribution of our study is that our analysis

suggests that higher levels of "Intrusive Distraction" are also linked to social processes. As well

as linking intrusive thought to the relatively low social well-being, our analysis also shows that

this pattern of thought tends to dominate solo activities (such as exercise, doing nothing, and

homework) and is relatively absent during social interactions, either virtually or in person.

Interestingly, despite intrusive thought being relatively high during exercise, this component was

relatively low during sports, providing further evidence that social contexts may reduce levels of

unpleasant intrusive experiences. Importantly, these data are consistent with prior studies that

shown links between anxiety and increased levels of mind-wandering (e.g., [33]), a self-

generated state that is likely to have distracting features [15, 34]. Altogether, therefore, these data

suggest that unpleasant intrusive thoughts may be promoted by loneliness and social isolation, a

situation that may promote personal worries, often known as current concerns [35]. It is possible

based on our study that interventions that increase opportunities for socialization may be an

important way to help individuals who suffer from anxiety.

Open Questions

As well as detailing links between thinking in daily life and well-being, our study also

raises a number of important open questions. For example, the consistency of the thought-

activity mappings suggests that in any given population there may be systematic relationships

between what people think and what they do in daily life. Based on these results, it may be

possible to develop an understanding of the activities in daily life that are most related to

different features of a person's thought content, a taxonomy of mappings between thoughts and

activities which, as a discipline, we currently lack. It is worth noting that the capacity for mDES

to map thoughts in the lab and in daily life [36] makes it a useful tool for this goal. The relative

ease with which we can administer mDES in daily life makes it possible to study a relatively

large and socio-culturally diverse population (e.g. older adults or individuals from a different

culture). Thus while our study shows that mDES is reliable, the specific results presented in this

study may not generalize to different cultures, ethnicities, or ages, our methodology of

smartphone sampling is free and easy to use, so future studies can address this gap by broadening

the scope of their sampling population in a reasonably accessible manner. It is also important to

note that there were a number of instances of incorrect and/or missing demographic data in our

data which should be considered a limitation of our current study.

It is also likely that the battery of questions we used in our study could be improved. In

our study, we used a combination of mDES questions that had been used previously in both lab-

based and daily-life-based settings, as well as new mDES questions based on previously

identified limitations. Mulholland et al. [9] noted that including a single modality question does

not allow participants to describe their conscious experience of an activity fully. In that study, a

component that was common while listening to music was dominated by thoughts with images as

a key feature (in that mDES battery there were no questions about sounds). In light of this

discovery, we decided to replace the modality probe with three questions relating to an

individual's sensory experience (i.e., images, words, and sounds). The inclusion of these three

questions most likely led to the discovery of the "Sensory Engagement" component in the

current study. It is important to note that a similar component was observed in the context of

movie watching (using the same set of items as was applied in the current study [11]) and was

associated with periods of films when brain activity was activated within sensory systems (both

auditory and visual). This demonstrates that it is possible to optimize the questions included in

mDES approaches that could, in the future, provide more accurate ways for participants to

describe their conscious experience. In the same vein, there are likely to be ways to improve the

measures of mental health we employed in this analysis. The health and wellness surveys were

selected based on a widespread, general approach to health and wellness, rather than clinical

diagnosis. In the future, more accurate or informative questionnaires could be selected that

would provide a better understanding of how our thoughts in daily life impact upon our mental

health.

547 **Table 1**

548 *mDES Questions*

Dimension	Question	Scale Low	Scale High
Task	My thoughts were focused on an external task or activity:	Not at all	Completely
Future	My thoughts involved future events:	Not at all	Completely
Past	My thoughts involved past events:	Not at all	Completely
Self	My thoughts involved myself:	Not at all	Completely
People	My thoughts involved other people:	Not at all	Completely
Emotion	The emotion of my thoughts was:	Negative	Positive
Images	My thoughts were in the form of images:	Not at all	Completely
Words	My images were in the form of words:	Not at all	Completely
Sounds	My thoughts were in the form of sounds:	Not at all	Completely
Detailed	My thoughts were detailed and specific:	Not at all	Completely
Deliberate	My thoughts were:	Spontaneous	Deliberate
Problem	I was thinking about solutions to problems or goals:	Not at all	Completely
Intrusive	My thoughts were intrusive:	Not at all	Completely
Knowledge	My thoughts contained information I already knew (e.g., knowledge or memories):	Not at all	Completely
Absorption	I was absorbed in the contents of my thoughts:	Not at all	Completely
Distracting	My thoughts were distracting me from what I was doing:	Not at all	Completely

549 *Note*. Participants rated statements on a 1-to-10 Likert scale.

552　**Table 2**

553　*Social Environment Questions*

Environment	Question	Environment Type
Physical	Were you alone, or physically with other people?	Alone
		Around people but not interacting with them
		Around people and interacting with them
	Where you alone, or physically with a pet?	Not with a pet
		Around a pet but not interacting with them
		Around a pet and interacting with them
Virtual	Were you alone, or virtually with other people?	Alone
		Around people but not interacting with them (e.g., reading messages but not replying, being on a video call but not participating, etc.)
		Around people and interacting with them (e.g., direct communication with another person by text, instant messaging, calling, or video calling, etc.)

556 **Table 3**

557 *Physical Environment Question*

List Type	Question	Location List
Location	Where were you?	Inside a home
		Inside a shop
		Inside a workplace
		Inside (other)
		Outside in a city or town
		Outside in nature
		Outside (other)

558 *Note*. If participants selected "Inside (other)," or "Outside (other)," they were asked to specify their location.

561   **Table 4**

562   *Primary Activity Question*

List Type	Question	Activity List
Activity	What were you doing?	Eating
		Homework
		Household chores
		Listening to music
		Napping or resting
		Nothing or waiting
		Personal exercise
		Personal hygiene care
		Physical leisure or sports
		Reading
		Shopping
		Talking in person
		Talking on the phone
		Video-calling
		Messaging by phone/device
		Traveling or commuting
		Using a computer or an electronic device
		Walking the dog
		Watching TV
		Working
		Other activity

Note. If participants selected "Other activity," they were asked to specify what they were doing. If participants

selected "Using a computer or an electronic device," they were asked to further specify their activity from a list of

alternative activities that included "Social media: Passive scrolling," "Social media: Active engagement,"

"Gaming," "Admin (e.g., banking, calendar, etc.)," and "Other." If participants selected "Other," they were asked to

specify what they were doing).

Table 5

PCA Loadings Generated from mDES Questions

Dimension	Detailed Task Focus	Intrusive Distraction	Episodic Social Cognition	Future Problem-Solving	Sensory Engagement
Task	.605	-.207	.017	-.033	.142
Future	.228	.068	.106	.797	.064
Past	-.021	.408	.636	.054	.169
Self	.003	.166	.227	.730	-.009
Person	-.079	-.004	.645	.253	.208
Emotion	.018	-.730	.115	.042	.351
Images	.017	.048	.032	.216	.742
Words	.435	.020	.452	-.092	-.264
Sounds	.051	.080	.127	-.177	.680
Detailed	.748	.038	.224	.102	.034
Deliberate	.702	-.109	-.059	.009	-0.137
Problem	.609	.146	.010	.412	-0.052
Intrusive	-.013	.710	.211	.137	0.207
Knowledge	.240	.059	.651	.141	-.002
Absorption	.560	.267	.084	.155	.230
Distracting	-.015	.689	.174	.220	.228

580 **Table 6**

581 *LMM Estimated Marginal Means for the "Detailed Task Focus" Component*

Primary Activity	Mean	Std. Error	Lower Bound 95% Confidence Interval	Upper Bound 95% Confidence Interval
Eating	-.198	0.044	-.284	-.112
Homework	.664	.033	.599	.729
Chores	-.017	.058	-.130	.096
Music	-.205	.062	-.326	-.083
Resting	-.686	.043	-.770	-.602
Nothing	-.344	.057	-.455	-.233
Exercise	.042	.080	-.114	.199
Hygiene	-.029	.060	-.146	.088
Sports	-.149	.127	-.398	.099
Reading	.281	.080	.124	.438
Shopping	-.042	.094	-.226	.143
Conversation	-.293	.043	-.377	-.209
Phone-call	.043	.102	-.157	.243
Video-call	.168	.089	-.007	.344
Texting	.025	.084	-.140	.189
Commuting	-.139	.077	-.291	.012
Computer	-.250	.042	-.333	-.167
Walking dog	-.005	.165	-.329	.318
TV	-.388	.042	-.470	-.306
Working	.544	.052	.441	.646
Other	.122	.043	.038	.206

588   **Table 7**

589   *LMM Estimated Marginal Means for the "Intrusive Distraction" Component*

Primary Activity	Mean	Std. Error	Lower Bound 95% Confidence Interval	Upper Bound 95% Confidence Interval
Eating	-.036	.048	-.130	.059
Homework	.232	.038	.157	.308
Chores	0.026	.062	-.147	.095
Music	.102	.066	-.028	.232
Resting	.199	.048	-.106	.292
Nothing	.316	.061	.197	.436
Exercise	-.156	.084	-.320	.009
Hygiene	-.160	.064	-.285	-.035
Sports	-.552	.131	-.810	-.295
Reading	-.079	.084	-.244	.086
Shopping	-.229	.098	-.422	-.037
Conversation	-.084	.047	-.177	.009
Phone-call	-.006	.106	-.214	.202
Video-call	-.031	.093	-.214	.152
Texting	.041	.088	-.131	.214
Commuting	-.133	.081	-.293	.026
Computer	.070	.047	-.022	.162
Walking dog	.063	.171	-.272	.397
TV	-.121	.047	-.213	-.030
Working	-.125	.057	-.236	-.014
Other	-.090	.043	-.183	.004

592 **Table 8**

593 *LMM Estimated Marginal Means for the "Episodic Social Cognition" Component*

Primary Activity	Mean	Std. Error	Lower Bound 95% Confidence Interval	Upper Bound 95% Confidence Interval
Eating	0.170	0.047	0.078	0.262
Homework	-0.178	0.036	-0.247	-0.108
Chores	0.057	0.062	-0.063	0.178
Music	0.070	0.066	-0.060	0.200
Resting	-0.194	0.046	-0.284	-0.104
Nothing	0.082	0.061	-0.036	0.201
Exercise	-0.089	0.085	-0.256	0.078
Hygiene	0.124	0.064	-0.001	0.249
Sports	0.195	0.135	-0.070	0.459
Reading	-0.171	0.085	-0.338	-0.003
Shopping	-0.050	0.100	-0.246	0.147
Conversation	0.405	0.046	0.315	0.494
Phone-call	0.610	0.109	0.397	0.823
Video-call	0.469	0.095	0.283	0.656
Texting	0.561	0.089	0.385	0.736
Commuting	0.202	0.083	0.040	0.364
Computer	-0.178	0.045	-0.267	-0.089
Walking dog	0.429	0.176	0.085	0.773
TV	-0.243	0.045	-0.331	-0.155
Working	0.136	0.056	0.026	0.245
Other	-0.058	0.046	-0.148	0.032

 Table 9

 LMM Estimated Marginal Means for the "Future Problem-Solving" Component

Primary Activity	Mean	Std. Error	Lower Bound 95% Confidence Interval	Upper Bound 95% Confidence Interval
Eating	.127	0.047	0.035	0.218
Homework	-0.196	0.035	-0.265	-0.128
Chores	0.229	0.062	0.108	0.349
Music	0.082	0.066	-0.048	0.213
Resting	0.066	0.046	-0.023	0.156
Nothing	0.232	0.061	0.113	0.351
Exercise	0.344	0.086	0.176	0.512
Hygiene	0.523	0.064	0.398	0.648
Sports	0.260	0.136	-0.007	0.528
Reading	-0.199	0.086	-0.367	-0.030
Shopping	0.330	0.101	0.132	0.529
Conversation	0.084	0.045	-0.005	0.174
Phone-call	0.373	0.110	0.158	0.588
Video-call	0.284	0.096	0.096	0.473
Texting	0.365	0.090	0.188	0.541
Commuting	0.384	0.083	0.221	0.546
Computer	-0.104	0.045	-0.192	-0.015
Walking dog	0.007	0.177	-0.340	0.355
TV	-0.374	0.045	-0.461	-0.286
Working	0.065	0.056	-0.044	0.175
Other	0.012	0.046	-0.078	0.101

601   **Table 10**

602   *LMM Estimated Marginal Means for the "Sensory Engagement" Component*

Primary Activity	Mean	Std. Error	Lower Bound 95% Confidence Interval	Upper Bound 95% Confidence Interval
Eating	-0.017	0.049	-0.113	0.080
Homework	-0.235	0.041	-0.315	-0.155
Chores	-0.041	0.061	-0.160	0.079
Music	0.272	0.065	0.145	0.400
Resting	-0.003	0.049	-0.098	0.092
Nothing	-0.043	0.060	-0.161	0.075
Exercise	0.007	0.081	-0.152	0.165
Hygiene	0.025	0.063	-0.098	0.148
Sports	0.227	0.125	-0.018	0.471
Reading	0.101	0.081	-0.059	0.260
Shopping	-0.007	0.094	-0.191	0.177
Conversation	0.032	0.048	-0.063	0.126
Phone-call	-0.186	0.101	-0.385	0.013
Video-call	0.085	0.090	-0.091	0.261
Texting	-0.127	0.085	-0.293	0.039
Commuting	-0.093	0.079	-0.248	0.061
Computer	0.217	0.048	0.123	0.311
Walking dog	-0.038	0.162	-0.355	0.279
TV	0.427	0.048	0.333	0.520
Working	-0.171	0.057	-0.282	-0.060
Other	0.047	0.049	-0.048	0.143

605 **Table 11**

606 *Mental Health and Well-Being Multiple Regression Results Per PCA Component Scores*

Component	Model	Sum of Squares	df	Mean Square	F	Sig.
Detailed Task	Regression	.660	8	.083	.444	.894
Focus	Residual	46.866	252	.186		
	Total	47.526	260			
Intrusive	Regression	15.071	8	1.884	8.348	<.001
Distraction	Residual	56.869	252	.226		
	Total	71.940	260			
Episodic Social	Regression	2.127	8	.266	1.085	.374
Cognition	Residual	61.773	252	.245		
	Total	63.900	260			
Future	Regression	3.129	8	.391	1.806	.076
Problem-	Residual	54.569	252	.217		
Solving	Total	57.698	260			
Sensory	Regression	1.248	8	.156	.415	.911
Engagement	Residual	94.767	252	.376		
	Total	96.015	260			

Acknowledgements

This research is supported by (i) a National Science and Engineering Research Council

(NSERC) graduate fellowship awarded to Bridget Mulholland, (ii) an award from the

Government of Canada's New Frontiers in Research Fund (NFRF) (grant ID: NFRF-2021-

00183) to Dr. Jonathan Smallwood and Dr. Jeffrey D. Wammes, and (iii) an NSERC Discovery

Grant awarded to Dr Jonathan Smallwood (Grant ID: 2023-03496).

References

1.   Asztalos, M., et al., *Specific associations between types of physical activity and components of*

mental health. Journal of science and medicine in sport, 2009. **12**(4): p. 468-474.

2.   Kucyi, A., et al., *Recent advances in the neuroscience of spontaneous and off-task thought:*

implications for mental health. Nature mental health, 2023. **1**(11): p. 827-840.

3.   Vatansever, D., et al., *Distinct patterns of thought mediate the link between brain functional*

connectomes and well-being. Network neuroscience, 2020. **4**(3): p. 637-657.

4.   Kane, M.J., et al., *For whom the mind wanders, and when: An experience-sampling study of*

working memory and executive control in daily life. Psychological science, 2007. **18**(7): p. 614-

621.

5.   Smallwood, J., et al., *The neural correlates of ongoing conscious thought.* Iscience, 2021. **24**(3).

6.   Kane, M.J., et al., *For whom the mind wanders, and when, varies across laboratory and daily-life*

settings. Psychological science, 2017. **28**(9): p. 1271-1289.

7.   Turnbull, A., et al., *Age-related changes in ongoing thought relate to external context and*

individual cognition. Consciousness and cognition, 2021. **96**: p. 103226.

8.   Konu, D., et al., *Exploring patterns of ongoing thought under naturalistic and conventional task-*

based conditions. Consciousness and cognition, 2021. **93**: p. 103139.

9.   Mulholland, B., et al., *Patterns of ongoing thought in the real world.* Consciousness and

cognition, 2023. **114**: p. 103530.

10.   Smallwood, J., et al., *Representing representation: Integration between the temporal lobe and the*

posterior cingulate influences the content and form of spontaneous thought. PloS one, 2016.

11(4): p. e0152272.

11.   Wallace, R.S., et al., *Mapping patterns of thought onto brain activity during movie-watching.*

elife, 2025. **13**: p. RP97731.

12.   Mckeown, B., et al., *Experience sampling reveals the role that covert goal states play in task-*

relevant behavior. Scientific reports, 2023. **13**(1): p. 21710.

667 13. McKeown, B., et al., *Self-reports map the landscape of task states derived from brain imaging.*

668   Communications Psychology, 2025. **3**(1): p. 8.

669 14. Konu, D., et al., *A role for the ventromedial prefrontal cortex in self-generated episodic social*

670   *cognition.* NeuroImage, 2020. **218**: p. 116977.

671 15. McVay, J.C. and M.J. Kane, *Does mind wandering reflect executive function or executive*

672   *failure? Comment on Smallwood and Schooler (2006) and Watkins (2008).* 2010.

673 16. Chaieb, L., C. Hoppe, and J. Fell, *Mind wandering and depression: A status report.* Neuroscience

674   & Biobehavioral Reviews, 2022. **133**: p. 104505.

675 17. Smallwood, J. and R.C. O'Connor, *Imprisoned by the past: unhappy moods lead to a*

676   *retrospective bias to mind wandering.* Cognition & emotion, 2011. **25**(8): p. 1481-1490.

677 18. Forster, S., et al., *Unraveling the anxious mind: anxiety, worry, and frontal engagement in*

678   *sustained attention versus off-task processing.* Cerebral Cortex, 2015. **25**(3): p. 609-618.

679 19. Gaynor, N. and L. Fitzgerald, *Mind-Wandering and Its Relationship With Psychological*

680   *Wellbeing and Obsessive-Compulsive Symptomatology in the Context of Covid-19.* Psychological

681   Reports, 2023: p. 00332941231203563.

682 20. Rush, A.J., et al., *The 16-Item Quick Inventory of Depressive Symptomatology (QIDS), clinician*

683   *rating (QIDS-C), and self-report (QIDS-SR): a psychometric evaluation in patients with chronic*

684   *major depression.* Biological psychiatry, 2003. **54**(5): p. 573-583.

685 21. Reiss, S., et al., *Anxiety sensitivity, anxiety frequency and the prediction of fearfulness.* Behaviour

686   research and therapy, 1986. **24**(1): p. 1-8.

687 22. Baron-Cohen, S., et al., *The autism-spectrum quotient (AQ): Evidence from asperger*

688   *syndrome/high-functioning autism, malesand females, scientists and mathematicians.* Journal of

689   autism and developmental disorders, 2001. **31**: p. 5-17.

690 23. Kessler, R.C., et al., *The World Health Organization Adult ADHD Self-Report Scale (ASRS): a*

691   *short screening scale for use in the general population.* Psychological medicine, 2005. **35**(2): p.

692   245-256.

693   24.   Montgomery, S.A. and M. Åsberg, *A new depression scale designed to be sensitive to change.*

694   The British journal of psychiatry, 1979. **134**(4): p. 382-389.

695   25.   Norman, S.B., et al., *Development and validation of an overall anxiety severity and impairment*

696   *scale (OASIS).* Depression and anxiety, 2006. **23**(4): p. 245-249.

697   26.   Ware, J. and C. Sherbourne, *SHERBOURNE CD. THE MOS 36 ITEM SHORTFORM HEALTH*

698   *SURVEY (SF-36).* 1992.

699   27.   Leon, A., et al., *Assessing impairment in patients with panic disorder: the Sheehan Disability*

700   *Scale.* Social psychiatry and psychiatric epidemiology, 1992. **27**: p. 78-82.

701   28.   Group, W., *Development of the World Health Organization WHOQOL-BREF quality of life*

702   *assessment.* Psychological medicine, 1998. **28**(3): p. 551-558.

703   29.   Kahneman, D., et al., *A survey method for characterizing daily life experience: The day*

704   *reconstruction method.* Science, 2004. **306**(5702): p. 1776-1780.

705   30.   Mckeown, B., et al., *The impact of social isolation and changes in work patterns on ongoing*

706   *thought during the first COVID-19 lockdown in the United Kingdom.* Proceedings of the National

707   Academy of Sciences, 2021. **118**(40): p. e2102565118.

708   31.   Vatansever, D., et al., *The devil is in the detail: exploring the intrinsic neural mechanisms that*

709   *link attention-deficit/hyperactivity disorder symptomatology to ongoing cognition.* Psychological

710   medicine, 2019. **49**(7): p. 1185-1194.

711   32.   Poerio, G.L. and J. Smallwood, *Daydreaming to navigate the social world: What we know, what*

712   *we don't know, and why it matters.* Social and Personality Psychology Compass, 2016. **10**(11): p.

713   605-618.

714   33.   Fell, J., L. Chaieb, and C. Hoppe, *Mind wandering in anxiety disorders: a status report.*

715   Neuroscience & Biobehavioral Reviews, 2023: p. 105432.

716   34.   Smallwood, J. and J.W. Schooler, *The science of mind wandering: Empirically navigating the*

717   *stream of consciousness.* Annual review of psychology, 2015. **66**(1): p. 487-518.

718   35.   Klinger, E. and W.M. Cox, *Handbook of motivational counseling: Concepts, approaches, and*

719   *assessment*. 2003: john wiley & sons.

720   36.   Chitiz, L., et al., *Mapping cognition across lab and daily life using experience-sampling.* 2023.

721