TITLE: Constructing Ensemble Gene Functional Networks Capturing Tissue/condition-specific

Co-expression from Unlabled Transcriptomic Data with TEA-GCN

AUTHORS: Peng Ken Lim[1*], Ruoxi Wang[1], Jenet Princy Antony Velankanni[1], Marek Mutwil[1*]

[1]School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive, Singapore,

637551, Singapore

*Corresponding author:

Marek Mutwil

School of Biological Sciences,

Nanyang Technological University,60 Nanyang Drive,

637551, Singapore,

Singapore

Email: mutwil@ntu.edu.sg

Peng Ken Lim

School of Biological Sciences,

Nanyang Technological University,60 Nanyang Drive,

637551, Singapore,

Singapore

Email: pengkenlim.sbs@gmail.com

Abstract

Gene co-expression networks (GCNs) generated from public transcriptomic datasets can elucidate the co-regulatory and co-functional relationships between genes, making GCNs an important tool to predict gene functions. However, current GCN construction methods are sensitive to the quality of the data, and the interpretability of the identified relationships between genes is still difficult. To address this, we present a novel method — Two-Tier Ensemble Aggregation (TEA-) GCN. TEA-GCN utilizes unsupervised partitioning of big transcriptomic datasets and three correlation coefficients to generate ensemble GCNs in a two-step aggregation process. We show that TEA-GCN outperforms in finding correct functional relationships between genes over the current state-of-the-art across three model species, and is able to not only capture condition/tissue-specific gene co-expression but explain them through the use of natural language processing (NLP). In addition, we found TEA-GCN to be especially performant in identifying relationships between transcription factors and their activation targets, making it effective in inferring gene regulatory networks. TEA-GCN is available at https://github.com/pengkenlim/TEA-GCN.

Introduction

Gene co-expression networks (GCNs) play a pivotal role in elucidating co-regulatory and co-functional relationships between genes[1–5]. This concept, rooted in the observation that functionally related genes often exhibit coordinated expression patterns, has proven invaluable in predicting gene function and uncovering modules of co-regulated genes[3–6]. The utility of GCNs to uncover biological insights becomes especially important in non-model species where experimental validation of gene function is scarce[6,7]. The rapidly increasing affordability of RNA-seq, coupled with the fast rate at which genomes are continuously published has led to an explosion of public transcriptomic data, particularly in non-model species[7–9], and offers an unprecedented opportunity to unravel the evolutionary dynamics of gene function across diverse organisms via comparative-GCN analysis[6,7,10,11].

However, the potential of GCNs is hindered by several challenges when working with public transcriptome data. The composition of transcriptome data can significantly impact the quality of GCNs and the context of biological insights embedded within them[1,1,12,13]. For example, the overrepresentation of certain sample types (e.g. tissues, conditions) within the dataset can selectively represent some tissue/condition-specific regulatory information while ignoring others[14]. Additionally, the size and heterogeneity of the transcriptome data can also have a huge impact on the resultant GCN. For example, dataset size (i.e. the number of samples) can directly affect the statistical power and therefore, the resolving power of GCNs to differentiate authentic transcriptional co-regulation of genes from spurious correlation of expression patterns[12]. On the other hand, GCN built from a highly heterogeneous dataset consisting of diverse sample types can excel in capturing ubiquitous gene modules but can fail to capture condition-specific regulatory relationships due to insufficient sample-type replication[14]. Thus, the variability of publicly available RNA-seq data of different species, in terms of sample quality, heterogeneity, and size poses a substantial hurdle for constructing performant GCNs. To address these challenges, many studies have utilized highly curated public transcriptomic datasets that are sample-balanced and/or context-specific to construct well-performing GCNs[15–19]. However, these methods rely on labor-intensive sample annotation that might not be available for some species.

Recent advancements in obtaining performant GCNs have seen the use of ensemble approaches that combine (aggregate) multiple GCNs to enrich biological information[20–22]. In one study, gene expression omnibus (GEO) datasets of *Vitis vinifera* (Grape) were used to generate unweighted GCNs and aggregate them to form an ensemble GCN[21]. Aside from producing a more

robust GCN, this approach has achieved some degree of sample balancing as coexpression was

determined from context-(GEO dataset)-specific GCNs rather than directly calculated across all

samples. However, this approach might not work for cases where the number of GEO datasets

is limited, even for species that possess numerous public RNA-seq accessions. Another approach

(henceforth referred to as the "Subbaging" method) combined weighted GCNs calculated from

random subsets of principal components of the gene expression matrix[16,17]. However, the

stochastic nature of this approach due to random subsampling, as well as the ambiguous meaning

of principal components, compromises the interpretability of the resultant ensemble GCN. In

addition, both approaches only use Pearson's correlation coefficient (PCC), which can fail to

consider non-linear co-expression relationships between genes[1,23].

In response to these challenges, we present a novel method — Two-Tier Ensemble Aggregation

(TEA-) GCN. The TEA-GCN framework leverages unsupervised partitioning of publicly derived

transcriptomic data and utilizes three correlation coefficients to generate ensemble CGNs in a

two-step aggregation process (Figure 1). Importantly, the TEA-GCN approach does not require

annotated samples for batch correction and/or sample balancing, making it readily applicable to

large public transcriptomic datasets with limited preprocessing and suitable for species with

scarce sample annotation. We evaluated the performance of the TEA-GCN in uncovering diverse

biological relationships between genes and demonstrated its superior performance over the state-

of-the-art "subbagging" ensemble GCN method[16,17] in *Saccharomyces cerevisiae* (*S. cerevisiae*;

Baker's yeast), *Arabidopsis thaliana (A. thaliana*; Thale cress*)* and *Homo sapiens* (*H. sapiens*;

Human*),* three model organisms of vastly different biological/organizational complexities and

evolutionary lineages. Using *A. thaliana* as a model, we also demonstrated the framework's ability

to capture condition/tissue-specific gene co-expression and characterized its flexibility to scale to

smaller transcriptomic datasets. In addition, we found TEA-GCN to be especially performant in

identifying relationships between transcription factors and their activation targets that can rival the

well-established GENIE3 algorithm[24]. Finally, we show that TEA-GCN enables explainability

through the use of natural language processing (NLP) to reveal the experimental contexts

underpinning the co-expression relationships. Put together, we present TEA-GCN as not only a

more robust method to generate ensemble GCNs than the current state-of-the-art but also a

landmark method to generate explainable ensemble GCNs. TEA-GCN is available at

https://github.com/pengkenlim/TEA-GCN .

Figure 1. Overview of Two-tier Ensemble Aggregation Gene Co-expression Network (TEA-GCN) framework and GCN evaluation methodology. A) The processing of data from an initial gene expression matrix to the final ensemble GCN is visualized using a toy model for which there are only three genes (A, B, C) in the genome. The gene expression matrix is first partitioned via k-means clustering at a clustering granularity (k=3) that has been determined using a combination of semi-supervision and/or internal clustering evaluation metrics. For each dataset partition, GCNs are generated based on three different correlation coefficients and aggregated into one ensemble GCN. Resultant GCNs from each partition are again aggregated to yield the final ensemble TEA-GCN. GCNs with coexpression strength represented by Pearson's correlation coefficient (PCC), Spearman's rank correlation coefficient (SCC), and Biweight midcorrelation (Bicor) are denoted by purple, brown, and blue edges, respectively. The node color of GCNs corresponds to the particular dataset partition they represent. **B)** We define three sets of positive edges connecting genes sharing different biological relationships. Positive Metabolic (Met) edges are made by pairing experimentally validated enzymes from the same

pathway. Edges connecting enzymes that catalyze the same reaction step are removed from positive Met edges. Positive Transcription factor (TF) edges are generated between genes and their corresponding activator TFs. Positive Gene Ontology (GO) edges are made from pairing genes that were annotated for the same biological process (BP) GO term at the IDA (inferred from direct assay) evidence level. Edges connecting genes annotated with the same molecular function (MF) GO term are removed from positive GO edges. **C)** To evaluate a given GCN, co-expression strengths of positive edge sets for each biological aspect are compared to co-expression strengths of randomly determined negative edges of the same size using the Area-under-curve (AUC) of the receiver operator characteristics curve (ROC) and precision-recall curve (PRC). Scores calculated for each positive-negative edge set pair represent a GCN's performance to identify different biologically significant relationships between genes and can be aggregated using harmonic mean to infer a GCN's overall performance.

Results

Overview of Two-tier Ensemble Aggregation Gene Co-expression Network (TEA-GCN) Framework

TEA-GCN relies on the Two-tier Ensemble Aggregation (TEA) process consisting of two tandem aggregation steps, Coefficient Aggregation and Partition Aggregation (Figure 1A). Coefficient Aggregation involves the calculation of different co-expression metrics (coefficients) between a given pair of genes and then unifying (aggregating) said metrics into one measure of co-expression strength (Figure 1A). We used the three most popular metrics: Pearson's correlation coefficient (PCC, capturing linear relationships), Spearman's rank correlation coefficient (SCC, capturing monotonic relationships)[1,25,26] and biweight midcorrelation[27] (Bicor), a coefficient employed in the widely used WGCNA R package[28] that is more resistant to outliers and noise as compared to PCC[27]. Although there have been several studies comparing the three coefficients, there is no current consensus as to which coefficient is the most ideal for generating robust GCNs as different coefficients have been shown to excel in detecting different biological relationships[26]. Therefore, to detect more types of complex yet biologically significant co-expression relationships between genes, the TEA process uses Coefficient Aggregation to derive co-expression based on all three coefficients.

Partition Aggregation aims to capture tissue/condition-specific GCNs that might be missed if all samples are used for the network construction. The tissue/condtion-specific networks have been shown to capture relevant biological relationships that only appear in certain spatiotemporal settings[15,18,19], or present in response to certain environmental conditions[29–31]. However, the sample composition of publicly available RNA-seq samples is expected to be biased towards certain sample types (e.g., easily attainable leaves in plants and certain popular mammalian cell lines), which might miss co-expression relationships between underrepresented samples. To address this, the TEA-GCN framework uses K-means clustering[32], an unsupervised clustering algorithm to partition the transcriptomic dataset into samples with similar transcriptome profiles[33] (Figure 1A). Then, the co-expression strengths between all gene pairs are aggregated across all partitions to detect tissue/condition-specific co-expression. Our approach combines the Coefficient and Partition aggregation in two steps. In the first step, Coefficient Aggregation combines the three coefficients in every partition and in the second step, Partition Aggregation combines the coefficients from the different partitions (Figure 1A).

To benchmark the TEA-GCN framework against the state of the art, we investigated

whether the obtained networks can predict biologically significant relationships between genes

(Figure 1B and C). To this end, we investigated whether the networks can recover experimentally-

verified metabolic networks (Met) (Figure 1B; green), transcription factor-target relationships (TF)

(Figure 1B; purple) and genes involved in the same biological process (GO) (Figure 1B; blue).

We obtained the experimentally verified relationships (positive edges) for Met, TF and GO

categories for the three species (see methods, Table S1 to S9). Since there is a tendency for co-

linearity between the estimated expression abundances of sequence-similar homologs due to

how multi-mapping reads are accounted for in pseudoalignment[34], edges that connect genes

annotated with the same molecular function were removed from GO positive edges to prevent

these edges from inflating performance scores (Figure 1B; blue). Similarly, we removed edges

that connect genes coding for enzymes that catalyze the same reaction steps from Met positive

edges. Using the co-expression strengths extracted for different sets of positive edges and

negative edges (representing genes with no functional relationship, obtained from sampling

random gene pairs), the Area-under-curves of the receiver operator characteristics curve

(AUROC) and precision-recall curve (AUPRC) performance scores were calculated for each

biological aspect (e.g. AUROC-Met and AUROC-Met; Figure 1C). We aggregated the scores with

harmonic mean to infer a GCN's overall performance (i.e., AUROC-HM and AUROC-HM; Figure

1C). To account for the random effect on performance scores due to the random selection of

negative edges, median scores from 100 evaluation iterations are reported.

Coefficient Aggregation and Partition Aggregation improves GCN performance

The TEA process uses different aggregation functions for Coefficient Aggregation and Partition

Aggregation. Coefficient Aggregation employs the Maximum (Max) function which outputs the

highest correlation magnitude reported by three different coefficients (PCC, SCC, bicor). In

contrast, Partition Aggregation employs the Rectified Average (RAvg) function to aggregate co-

expression strengths from different dataset partitions. The RAvg function takes in co-expression

strengths from every dataset partition and converts all negative co-expression strengths into zero

before averaging. Effectively, the RAvg function considers negative co-expression, which results

from the inverse correlation between gene expression, as the absence of co-expression (i.e. zero

co-expression).

The TEA process relies on k-means clustering[32] to generate dataset partitions from public

transcriptome datasets[33]. However, the optimal number of partitions (i.e. the number of

biologically relevant subsets of the data) is unknown. Thus, we explored a range of partitions (k)

for publicly available data for *S. cerevisiae*, *A. thaliana*, and *H. sapiens*. After removing samples

with poorly mapped reads (see methods), we obtained 150,096, 57,759, and 95,553 RNA-seq

samples for *S. cerevisiae*, *A. thaliana*, and *H. sapiens,* respectively (Table S10 to S12). To

investigate how partitioning granularity (k) affects the performance of the GCN produced by TEA,

we tested preselected *ks* ranging from 2 to 1000 (see methods; Table S13 to S15) (Figure 1C).

Across all three organisms, the performance (measured by AUROC-HM and AUPRC-HM)

of the TEA generally increases with partitioning granularity and saturates quickly over a large

window of k (Figure 2A; green line). The ensemble GCN inferred by the TEA can outperform GCN

obtained from calculating PCC from all samples (PCC GCN) at every k for *S. cerevisiae* and *A.*

thaliana (Figure 2A; blue dotted baseline). Although the outperformance of TEA for *H. sapiens* is

not observed for every k (low AUCPRC-HM at *ks* < 248 and > 902), TEA still shows good

performance for the majority (approx. 69%, 248 to 902 k) of the sampled *ks* (2 to 1000). More

importantly, the performance of TEA is stable over a wide range of k and does not decay

substantially even at high k, which suggests the robustness of TEA to resist over-partitioning

(Figure 2A). Thus, the effectiveness of TEA can be achieved by testing the different k values or

by using a sufficiently large k (identified using internal clustering metrics like silhouette coefficient)

that does not under-partition the public transcriptome dataset (Figure 1B).

To investigate whether aggregating different coefficients (i.e. Coefficient Aggregation)

improves the performance, we evaluated Partition Aggregation-only GCNs using only one of the

three coeffieicent (PCC, SCC, or Bicor) (Figure 2B) over the same preselected ks for each species

(Figure S1). Curiously, we found that the optimal k (k at which the sum of median AUROC-HM

and AUPRC-HM is highest) is dependent on the coefficient. For example, optimal ks for *A.*

thaliana Partition Aggregation-only GCNs were 62, 444 and 950 when PCC, SCC and Bicor were

used, respectively. As such, median performance scores of various Partition Aggregation-only

GCNs (PCC-RAvg, SCC-RAvg, and Bicor-RAvg) were compared to GCNs generated by the TEA

process (Max-RAvg) at their respective optimal ks (Figure 2C). We found that Max-RAvg GCNs

achieves median performance scores (AUROC-HM, AUPRC-HM) of (0.650, 0.633), (0.659,

0.670), and (0.554, 0.554) for *S. cerevisiae*, *A. thaliana*, and *H. sapiens* respectively (Figure 2C),

which is higher than the performance scores of (0.532, 0.529), (0.602, 0.622), and (0.506, 0.545)

of their corresponding PCC GCNs (Figure 2A). Crucially, we show that Max-RAvg outperforms

Partition Aggregation-only GCNs across all species in AUROC-HM and AUPRC-HM (Figure 2C,

heatmap section 2), which demonstrates that the combination of Coefficient Aggregation and

Partition Aggregation results in more performant GCNs, as compared to only using Partition

Aggregation alone.

The performance of the GCNs can be further improved by transforming the co-expression

strengths to rank-based metrics, such as Mutual Ranks (MR)[35]. GCNs based on MR-transformed

co-expression values of Max-RAvg (Max-RAvg-MR) further increased performance across all

biological aspects in the three species (Figure 2C). Finally, we also explored whether other

aggregation functions influence the network performance. To this end, we compared the

performances of GCNs generated by Max-Avg and Max-RAvg, which use Avg and RAvg

functions, respectively, for Partition Aggregation. The performance differences were minor before

MR transformation (Figure 2C, heatmap section 1). However, we found that RAvg for Partition

Aggregation yielded more performant GCNs after MR transformation (i.e. Max-Avg-MR and Max-

RAvg-MR) in *S. cerevisiae* and *A. thaliana*, while performance between the two variants remained

similar in *H. sapiens* (Figure 2C, heatmap section 2)*. Therefore, removing negative correlation

values improved the performance of the GCN and Max-RAvg-MR (Coefficient Aggregation →

Partition Aggregation → Mutual Rank transformation) was chosen as the most performant

approach for the rest of the study and designated as the TEA-GCN method.

Figure 2. Benchmarking performance of TEA-GCN across Yeast, Arabidopsis and Human. A) Performance of TEA-GCNs in *S. cerevisiae*, *A. thaliana* and *H. sapiens* across different levels of partitioning granularity (*k*). The colors of solid lines correspond to the median performance scores of TEA-GCNs across 100 evaluations using different aggregation functions for Partition aggregation. Shaded regions denote upper and lower quartiles of performance scores. The blue baseline corresponds to median performances of non-ensemble GCN using Pearson's correlation coefficient (PCC). **B)** Performance of the Partition aggregation-only GCN generated

241 from one correlation metric. For each dataset partition, GCNs were generated based on the selected metric and subsequently
242 aggregated across partitions. **C)** Comparison of median harmonic mean-aggregated performances and biological aspect-specific
243 performances between TEA-GCNs (section 1), TEA-GCNs with mutual rank transformation (section 2) and Partition aggregation-only
244 GCNs (section 3) across the three different model organisms.

TEA-GCN achieves better overall performance over the state-of-the-art

To benchmark the performance of the TEA-GCN (i.e. Max-RAvg-MR), we compare the performances of *S. cerevisiae*, *A. thaliana*, and *H. sapiens* GCNs generated by our approach to networks from two industry-leading databases, COXPRESdb[16] (https://coxpresdb.jp/download/) and ATTED-II[16] (https://atted.jp/download/). According to the authors, the *A. thaliana* ATTED-II GCN, *S. cerevisiae* COXPRESdb GCN, and *H. sapiens* COXPRESdb GCN were generated by first constructing gene expression matrices of selected public batch-corrected RNA-seq samples. The GCNs were subsequently calculated from subsampled principal components of the matrices, and aggregated into one GCN[16,17]. As expected, these "Subbaging" ensemble GCNs have a better overall performance than non-ensemble PCC GCNs across all three species (ΔAUROC-HM and ΔAUPRC-HM > 0.06-0.16) (Figure 3A-B; Table S16 to 18). However, GCNs based on our approach showed a substantially higher performance for nearly all metrics (Figure 3A, 3B, and S2; Table S16 to 18).

To better understand the differences between our method and the state-of-the-art, we compared the median AUROC and AUPRC scores and their respective curves for TF, Met and GO-BP biological aspects of TEA-GCN and ATTED-II/COXPRESdb GCNs. TEA-GCNs were consistently more performant in detecting the experimentally verified transcription factor-target regulatory relationships (TF) across the three species (Figure 3C and 3D; Table S16 to 18), especially in *A. thaliana* and *H. sapiens.* To exemplify, *A. thaliana* TEA-GCN has a TF performance (AUROC-TF, AUPRC-TF) of (0.747, 0.785) as opposed to (0.680, 0.715) in *A. thaliana* ATTED-II GCN, while *H. sapiens* TEA-GCN has a TF performance of (0.604, 0.626) as opposed to (0.542, 0.587) in *H. sapiens* COXPRESdb GCN (Figure 3C and 3D; Table S16 to 18). Conversely, the ATTED-II/COXPRESdb GCNs perform better in identifying enzymes involved in the same metabolic pathway for *S. cerevisiae* and *H. sapiens*. *A. thaliana* TEA-GCN has a slightly higher Met performance (AUROC-Met, AUPRC-Met of 0.726, 0.757) than *A. thaliana* ATTED-II (0.711,0.749), and better performance in identifying genes sharing the same GO-BP term in *S. cerevisiae* (0.700, 0.716 vs 0.673, 0.712) and *A.thaliana* (0.621,0.636 vs 0.599, 0.719) (Figure 3C and 3D; Table S16 to 18).

We speculate that the superior performance of TEA-GCN lies in its ability to detect condition/tissue-specific co-expression from individual dataset partitions. As such, the observed TF performance disparity between TEA-GCN and the "Subbaging" method is more pronounced in multicellular *A. thaliana* and *H. sapiens* than in unicellular *S. cerevisiae* (Figure 3C and 3D;

277   Table S16 to 18), could be attributed to the presence of more specialized transcriptional programs

278   in multicellular organisms than unicellular organisms[36,37].

279
Figure 3. Benchmarking TEA-GCN across three different model organisms against "Subbaging" ensemble GCNs. Distribution
of **A)** AUROC-HM performance scores and **B)** AUPRC-HM performance scores between Subbaging CGNs and TEA-GCNs across *S.*
cerevisiae, *A. thaliana* and *H. sapiens.* We report delta values of performance scores (i.e. Δ AUROC-HM and Δ AUPRC-HM) used to
compare GCNs on a unified axis. The delta values were calculated by subtracting performance scores with the median performance
score of baseline non-ensemble PCC network of respective organisms. Median performance scores of PCC (blue) networks,
Subbaging CGNs (orange) and TEA-GCNs (green) are indicated. **C)** Receiver operating characteristics curves (ROCs) and **D)**
Precision-recall curves (PRCs) correspond to median performance scores for each biological aspect and organism. Across all panels,
performance information relating to the PCC network, Subbaging CGN and TEA-GCN are represented in blue, orange and green,
respectively.

TEA-GCN captures co-expression between experimentally validated enzymes in _Arabidopsis thaliana_ metabolic pathways

To characterize how well the TEA-GCN framework captures co-expression between enzymes of the same pathway, we compared the co-expression strengths of Met positive edges in _A. thaliana_ TEA-GCN and ATTED-II GCN. We define Met positive edges as edges that connect experimentally validated enzymes within individual PlantCyc pathways[38]. For a fair comparison of co-expression strengths across the two CGNs, we used standardized co-expression strengths (Z(Co-exp.)), which measure the co-expression strengths of each edge relative to the co-expression strengths of all edges in the GCN (see methods). _A. thaliana_ TEA-GCN performed better in identifying Met positive edges over the ATTED-II GCN across all major classes (Figure 4A). The exception was the "Generation of Precursor Metabolites & Energy" class, where the performance between the two GCNs is nearly identical (Figure 4A; Table S19), irrespective of co-expression thresholds (i.e. Z(Co-exp.) ≥ 0.5 and Z(Co-exp.) ≥ 1). On the other hand, TEA-GCN excelled in the "Metabolic Clusters" and "Biosynthesis" classes, where the percentage of positive edges at both thresholds is more than 10% higher than in ATTED-II GCN (Figure 4A; Table S19). The "Generation of Precursor Metabolites & Energy" class consists of conserved pathways such as the TCA cycle, glycolysis, respiration, and photosynthesis that are ubiquitously expressed and not likely condition/tissue-specific, which could explain the less pronounced performance of our approach. Conversely, "Biosynthesis" and "Detoxification" classes, which tend to comprise specialized pathways that are lineage-specific and/or condition/tissue-specific[39–44], has more co-expressed edges in TEA-GCN than ATTED-II GCN at both thresholds. As such, this observation is compatible with our hypothesis that the good performance of TEA-GCN could be due to its ability to capture condition/tissue-specific co-expression.

Figure 4. Characterizing TEA-GCN's ability to capture metabolic pathways in *A. thaliana*. **A)** Proportion of Met positive edges

of different PlantCyc pathways that are co-expressed in *A. thaliana* TEA-GCN and ATTED-II GCN at two (>0.5 and >=1) co-expression
thresholds. Met positive edges connect experimentally validated enzymes of different reaction steps. Grey labeled plot corresponds
to first-level ontological classes and compares between major pathway categories. The blue labeled plot breaks down second-level
ontological classes within "Biosynthesis", of which third-level ontological classes consisting of "Aromatic Compound Biosynthesis",
"Cell Structure Biosynthesis", "Hormone and Signalling Molecule Biosynthesis", and "Secondary Metabolite Biosynthesis" were further
broken down in the brown, purple, yellow and pink plots, respectively. **B)** Co-expression between enzymes of "Glucosinolate
Biosynthesis from Tryptophan" PlantCyc pathway (PWY-601) in *A. thaliana* TEA-GCN and ATTED-II GCN. Nodes represent
experimentally validated enzymes according to PlantCyc and are labeled with gene symbols (if available) or gene identifiers. The
thickness of grey edges connecting nodes corresponds to standardized co-expression strengths between enzymes in the respective
GCNs. **C)** Reaction steps in Glucosinolate Biosynthesis from Tryptophan PWY-601. Colored blocks in the middle of reaction steps
annotate enzyme nodes in panel B based on the reaction step they participate in.

To better compare the *A. thaliana* GCNs of TEA-GCN and ATTED-II, we investigated the

different sub-classes within the "Biosynthesis" class. TEA-GCN identified at least 10% more co-

expressed edges for "Hormone and Signaling Molecule Biosynthesis", "Cyclitol Biosynthesis",

"Aromatic Compound Biosynthesis", "Cell Structure Biosynthesis", "Secondary Metabolite

Biosynthesis", "Amide, Amidine, Amine, and Polyamine Biosynthesis", at both co-expression

thresholds (Figure 4A, Blue). Impressively, we observed an extra 23% of positive met edges

belonging to "Hormone and Signaling Molecule Biosynthesis" for TEA-GCN (61.5%; 123 out of

200) compared toATTED-II GCN (38.5%; 77 out of 200) (Figure 4A, blue; Table S19) at the stricter

co-expression threshold of Z(Co-exp.) ≥ 1. This is largely attributed to the "Gibberellin and

Gibberellin Precursor Biosynthesis", and "Jasmonate Biosynthesis" (Figure 4A, yellow) that

collectively account for more than half of the edges belonging to "Hormone and Signaling

Molecule Biosynthesis" (Table S19). Given that the biosynthesis of hormones is known to be

highly localized to certain tissues at specific stages of development or environmental conditions[45–

47], we attribute the higher performance to TEA-GCN's ability to capture condition-specific co-

expression. Other notable classes where TEA-GCN excels include "Fatty Acid Derivative

Biosynthesis" (TEA-GCN: 55.97% and ATTED-II: 31.45%, strict threshold; 159 total edges),

"Flavonoid Biosynthesis" (TEA-GCN: 51.97% and ATTED-II: 30.71%, strict threshold; 127 total

edges), "Lignin Biosynthesis" (TEA-GCN: 67.48% and ATTED-II: 47.15%, strict threshold; 123

total edges) and "Glucosinolate Biosynthesis" (TEA-GCN: 86.74% and ATTED-II: 66.85%, strict

threshold; 181 total edges) (Figure 4A; Table S19).

To exemplify the utility of TEA-GCN in predicting pathway memberships of enzymes, we

compared the co-expression strengths between genes that code for experimentally validated

enzymes participating in the "Glucosinolate Biosynthesis from Tryptophan" PlantCyc pathway

(henceforth referred to by its corresponding PlantCyc pathway accession, PWY-601; Figure 4B

and C). Glucosinolates are a class of specialized natural compounds synthesized by plants in the

Capparales order that consists of many agriculturally valuable crops in the Brassicaceae family

like *Brassica oleracea* (*B. oleracea*; cabbage, kale, and broccoli), *Brassica juncea (B. juncea*;

chinese mustard) *Brassica napus* (*B. napus*; rapeseed) and *Brassica rapa* (*B. rapa*; turnip and

field mustard)[48–50]. Glucosinolates and their breakdown products possess a plethora of

bioactivities, some of which can be toxic (e.g. goitrogenic and genotoxic) and others beneficial

(e.g. anti-cancer)[48–50]. As such, there is a significant research interest in transgenically modifying

the level of specific glucosinolates in Brassicaceae food crops[51], which involves the the

elucidation of enzymes involved in the biosynthesis of glucosinolate pathways. The PWY-601

pathway converts L-tryptophan to indole glucosinolates such as Glucobrassicin (3-IMG),

Neoglucobrassicin (MIMG), and 4-methoxyglucobrassicin (4-MoIMG), which have anti-

inflammatory and anti-cancer activities in humans[48,52–54]. We show that experimentally validated

enzymes of PWY-601 tend to be co-expressed with more experimentally validated enzymes from

the same pathway in *A. thaliana* TEA-GCN than in ATTED-II GCN (Figure 4B), which

demonstrates the superior ability of TEA-GCN to predict enzymes involved in the said pathway.

Furthermore, genes coding for enzymes involved in the derivatization of 3-IMG to 4-MoIMG (i.e.

IGMT1, *IGMT2*, *CYP91A2*, *CYP81F3*, and *CYP81F2*) are notably less co-expressed with other

PWY-601 enzymes in ATTED-II GCN, as opposed to the more connected PWY-601 network of

TEA-GCN (Figure 4B and C). This scarce connection is especially prominent for *CYP91A2* and

IGMT2, which codes for monooxygenase and methyltransferase that catalyze the hydroxylation

of 3-IMG to 4-hydroxylglucobrassicin (4-OhIMG) and the methylation of 4-OhIMG to 4-MoIMG,

respectively[55]; *CYP91A2* is not co-expressed with any genes in PWY-601, while *IGMT2* is only

connected to *IGMT1* which codes for a methyltransferase catalyzing the same reaction step

(Figure 4B). To summarize, we have shown that TEA-GCN can better capture co-expression

between genes coding for enzymes within specialized metabolic pathways than the state-of-the-

art "Subbagging" method, possibly due to TEA-GCN's ability to detect tissue/conditional-specific

gene co-expression.

TEA-GCN can accurately reveal condition/tissue-specific gene regulatory networks

Our analysis revealed that TEA-GCN is particularly performant in revealing transcription factors and their gene targets (Figure 3C and D; TF aspect). While a large portion of TF positive edges is co-expressed in both GCNs (237 out of 673, 35.2%, Figure 5A), a substantial portion (143 out of 673, 21.2%; Figure 5A) was found only in TEA-GCN. Conversely, only 19 TF positive edges were uniquely co-expressed in ATTED-II GCN (2.8%; Figure 5A). Thus, TEA-GCN contains nearly all edges found in ATTED-II and can identify additional positive edges.

To further characterize the difference between the gene regulatory networks identified by TEA-GCN and ATTED-II, we analyzed transcription factors where TEA-GCN, but not ATTED-II, showed a high co-expression (Z(Co-exp.) >1.5) with their transactivation gene targets (Figure 5B). The identified transcription factor genes are known to be highly tissue/condition-specific in their expression, which further supports our hypothesis that TEA-GCN excels in detecting tissue/condition-specific co-expression. To exemplify, *JAGGED LATERAL ORGANS*'s (*JLO*) expression is specific to embryos[56], while *ABA INSENSITIVE 5*'s (*ABI5*) and *FUSCA3*'s (*FUS3*) expression are seed-specific[57,58]. On the other hand, *OBF-BINDING PROTEIN 2*'s (*OBP2*) expression is known to be induced by wounding and phytohormone Methyl jasmonate (MeJA)[59] , *NAC DOMAIN CONTAINING PROTEIN 19*'s (*ANAC019*) expression is induced by drought, high-salt stress and phytohormone abscisic acid[60], and *WRKY6*'s expression is induced by phosphate-starvation[61]. Curiously, *WRKY25* is found by ATTED-II GCN to be negatively co-expressed with its transactivation targets of *HEAT SHOCK FACTOR 4* (*HSF4*) and *HEAT SHOCK TRANSCRIPTION FACTOR B2A* (*HSFB2A*)[62] (Figure 5B). *WRKY25* is known to be involved in responses against a multitude of abiotic stresses such as heat, cold, osmotic, and high-salt conditions[62–65]. Given the multimodal functions of *WRKY25* and its participation in various abiotic responses, we speculate that its mode of action might be condition-specific, which might be the reason why its co-expression with *HSF4* and *HSFB2A* were not detected in ATTED-II GCN.

Next, to investigate whether the transcription factors are co-expressed with biologically relevant genes, we compared the functional enrichment of genes co-expressed with the selected transcription factors for *A. thaliana* TEA-GCN and ATTED-II GCN. To this end, we encapsulated the co-expression neighborhoods of transcription factors as lists of genes ranked by co-expression and used them as input for Gene Set Enrichment Analysis (GSEA)[66,67] using GO-BP gene sets (Figure 5C). Conceptually, if genes involved in a particular biological function are overrepresented in highly ranked genes (i.e. strongly co-expressed with the transcription factor),

the calculated enrichment score (ES) for the biological function will be large and statistically

significant (Figure 5C). The statistical significance of said overrepresentation can be determined

in the form of a null-hypothesis p-value (p_BH, after multiple-testing correction using the

Benjamini-Hochberg method) by comparing ESs calculated from permuted rankings (permuted

ESs) to the ES calculated from the original ranking[66], while the magnitude of overrepresentation

can be inferred from the Normalized Enrichment Score (NES) calculated as the fold-change of

ES from the mean of permuted ESs[66].

Convincingly, we show that the functional enrichment profiles of co-expression

neighborhoods of selected transcription factors are significant for many of their known biological

functions in TEA-GCN, but not for ATTED-II (Figure 5D and E, Figure S3). For example, we show

genes involved in "Hyperosmotic response" (GO:0006972), "Hyperosmotic salinity response"

(GO:0042538), "Response to salt" (GO:1902074), "Cellular response to salt stress"

(GO:0071472), "Response to salt stress" (GO:0009651), "Heat acclimation" (GO:0010286),

"Response to heat" (GO:0009408), "Response to cold" (GO:0009409) and "Response to reactive

oxygen species" (GO:0000302) are enriched (p_BH ≥ 0.05) in the *WRKY25* co-expression

neighborhood of TEA-GCN but not ATTED-II GCN (Figure 5D and E). Although both *WRKY25*

co-expression neighborhoods from *A. thaliana* TEA-GCN and ATTED-II GCN are enriched for

genes involved in "Response to osmotic stress" (GO:0006970), the former has a higher

magnitude of enrichment according to NES scores (Figure 5D and E). Thus, in addition to

showcasing the robust gene function prediction abilities of the TEA-GCN, we have exemplified its

potential to predict the multimodal functions of genes.

Figure 5. Characterization of transcription factor-target predictions in *A. thaliana* with TEA-GCN. A) Comparing co-expression of 673 TF positive edges in TEA-GCN and ATTED-II GCN. Venn diagram (top) shows common and unique TF positive edges that are co-expressed (*Z*(Co-exp.) ≥ 1) in *A. thaliana* TEA-GCN and ATTED-II GCN. Heatmap (bottom shows) co-expression strengths of TF positive edges in the two GCNs. Edges are sorted based on their co-expression in ATTED-II GCN. Color intensities correspond to co-expression strength (i.e. *Z*(co-exp.)). **B)** Co-expression between selected TFs and their known targets in TEA-GCN and ATTED-II GCN. TF and non-TF genes are represented by yellow diamond-shaped nodes and purple circular-shaped nodes, respectively. Edges with different colors and thicknesses represent co-expression strengths between TFs and their known targets. Grey chevrons within the edges indicate the direction of regulation between TFs and their targets. **C)** Functional enrichment of the co-expression neighborhood of TFs is determined using Gene Set Enrichment Analysis (GSEA). For a given TF, its co-expression neighborhood information is encapsulated by genes ranked by their co-expression with said TF and used as input for GSEA. Gene Ontology (GO) biological processes (BP) were used as gene sets. **D)** Dot plots comparing the enrichment statistics of selected GO terms in the neighborhood of *WRKY25* in TEA-GCN and ATTED-II GCN. GO terms were selected based on the known functions of *WRKY25*. Dot sizes for GO terms correspond to normalized enrichment scores (NES), while inverted triangles represent GO terms with negative NES scores. For GO terms with positive NES scores, the ax-axis indicates the statistical significance in the form of Benjamini-Hochberg-corrected p-values (p_BH). The red dotted lines denote the statistical significance threshold (p_BH ≥ 0.05). **E)** Running-sum profiles for enrichment scores and annotated gene rankings of selected GO terms between co-expression neighborhoods of *WRKY25* in TEA-GCN and ATTED-II GCN. For panels D and E, colors of dots, running sum profiles, and annotated gene rankings correspond to the different selected GO terms.

TEA-GCN remains performant on smaller datasets

So far, we showed that the TEA-GCN framework performs well on large transcriptome datasets of *S. cerevisiae* (n = 150,096), *A. thaliana* (n = 57,759), and *H. sapiens* (n = 95,553). Given that available public RNA-seq data might be less abundant for some species, we investigated TEA-GCN perforamnce on smaller datasets. To this end, we generated GCNs for downsampled datasets consisting of 5,000 samples and 500 samples that were randomly selected (without replacement) from the *A. thaliana* public transcriptome dataset. We show that while the overall performance of the *A. thaliana* TEA-GCNs markedly decreases as the datasets become smaller, the *A. thaliana* TEA-GCN at n = 500 is still substantially more performant (AUROC-HM: 0.649, AUPRC-HM: 0.677; Figure 6A; Table S20) than the non-ensemble PCC GCN (AUROC-HM: 0.602, AUPRC-HM: 0.622; Figure 3A and B; Table S17) generated from the complete *A. thaliana* public transcriptome dataset (n = 57,759). Crucially, we show that *A. thaliana* TEA-GCN at n = 5,000 (AUROC-HM: 0.669, AUPRC-HM: 0.699; Figure 6A; Table S20) can still outperform ATTED-II GCN (AUROC-HM: 0.659, AUPRC-HM: 0.690; Figure 6A; Table S17), despite using a downsampled dataset that is more than one order of magnitude smaller than the complete dataset. This outperformance is made more impressive considering that ATTED-II GCN is generated via the "subbaging" method using 14,741 curated, sample-balanced, and batch-corrected RNA-seq samples (according to https://atted.jp/download/), as opposed to *A. thaliana* TEA-GCN at n = 5,000 which uses 5,000 non-batch corrected RNA-seq samples that have been randomly selected from the public transcriptomic dataset. Thus, we have demonstrated that the performance of TEA-GCN is not only translatable to smaller datasets, but can likely outperform the "subbaging" methods, which require the time-consuming and complex annotation and balancing of samples.

Figure 6. Scalability of TEA-GCN to smaller datasets and its GRN inference performance compared to GENIE3. A) Distribution
of AUROC-HM performance scores and AUPRC-HM performance scores between ATTED-II GCNs and *A. thaliana* TEA-GCN
constructed from the complete transcriptomic dataset (*n* = 57,759) and randomly downsampled datasets (*n* = 5,000 and *n* = 500), that
were sampled 100 number of times. **B)** AUROC-GRN and AUPRC-GRN calculation relies on GRN positive edges and GRN negative
edges. GRN positive edges connect transcription factors to the targets they activate. GRN positive edges are directional, differing
from TF positive edges, thus transcription factors targeting each other are connected bytwo separate positive edges. GRN negative
edges are generated by randomizing genes connected to transcription factors in GRN positive edges. **C)** Distribution of AUROC-GRN

480   and AUPRC-GRN scores of *A. thaliana* TEA-GCNs and GENIE3-GRNs constructed from the complete and downsampled
481   transcriptomic datasets. Non-ensemble PCC GCN is included as a comparison. Receiver operating characteristics curves (ROCs)
482   and Precision-recall curves (PRCs) correspond to median performance scores for each GCN.

TEA-GCN outperforms state-of-the-art in GRN inference

As demonstrated earlier, the TEA-GCN framework excels in revealing gene regulatory networks

(Figure 3C, 3D, and 5), which prompted us to compare its GRN-inference capabilities to GENIE3[24]

(ref), a well-established machine learning-based algorithm dedicated to inferring GRN from

transcriptomic data. Briefly, GRN-inference methods like GENIE3 calculate the propensities of

genes as regulatory targets for transcription factors defined by the user. Since GRNs are

directional (e.g. TF1 → TF2) and the aim is to predict TF - target gene pairs (rather than any gene-

gene pairs), we defined a new set of positive directional edges from experimentally verified GRNs

for *A. thaliana* (Figure 6B; Table S21; see methods). Similarly, the negative edges were defined

by randomizing target genes connected to edges in the GRN positive edge set (Figure 6B).

AUROC-GRN and AUPRC-GRN were calculated 100 times from propensities (for GENIE3) and

co-expression strengths (for TEA-GCN) of edges (Figure 6B; Table S22).

Comparison of TEA-GCN and GENIE3 revealed that TEA-GCN outperforms GENIE3

even when a smaller dataset comprising < 10% of samples is used (Figure 6C; Table S22).

Impressively, TEA-GCN at n = 500 (AUROC-GRN: 0.757, AUPRC-GRN: 0.777; Figure 6C; Table

S22) outperforms the GENIE3 GRN at n = 5,000 (AUROC-GRN: 0.711, AUPRC-GRN: 0.722;

Figure 6C; Table S22) in GRN inference despite a ten-fold difference in dataset size. We also

show that *A. thaliana* TEA-GCN generated using the complete dataset (n = 57,759) has the best

GRN-inference ability (AUROC-GRN: 0.812, AUPRC-GRN: 0.837; Figure 6C; Table S22). On the

other hand, the non-ensemble PCC GCN generated using the complete dataset (n = 57,759)

performed the worst (AUROC-GRN: 0.636, AUPRC-GRN: 0.653; Figure 6C; Table S22). Thus,

the excellent GRN-inference capabilities of TEA-GCNs demonstrate that our method can

outperform dedicated GRN algorithms, such as GENIE3.

Experimental contexts underpinning gene co-expression are discoverable in TEA-GCN

Despite their utility in identifying functional associations between genes, current network-based methods cannot explain which biological and experimental conditions (e.g., organ, tissue, treatment) underpin the observed co-expression between genes, making it difficult to understand in which conditions they are functionally related. This lack of explainability is common for machine-learning GRN methods such as GENIE3[24] and GRNBoost2[68], which are often considered "black boxes" due to the difficulty in understanding how the network edges are determined.

To investigate whether TEA-GCN provides explainable gene-gene relationships, we analyzed the datasets underpinning the condition/tissue-specific co-expression (Figure 7). To this end, we developed a Natural Language Processing (NLP) method that analyzes the metadata for each of the data partitions identified by k-means (Figure 7A). Specifically, we identified statistically overrepresented lemmas within the metadata of the *A. thaliana* RNA-seq samples (Table S14) in each dataset partition (Figure 7A; see methods). As a result, dataset partitions can be annotated by the overrepresented lemmas that might describe the experimental conditions and/or tissue of samples contained within partitions (Table S23). Next, the overrepresented lemmas of the partitions were used to annotate TEA-GCN edges by a statistical approach similar to GSEA. To achieve this, for each edge in TEA-GCN, we first ranked the underpinning co-expression strengths stemming from each the dataset partition (step immediately after Coefficient Aggregation).(Table S24). Then, we used these ranks to identify the overrepresented lemmas of dataset partitions with strongly observed co-expression (Figure 7B, see methods).

To test this approach, we applied this method on GRN positive edges that exhibited strong co-expression in *A. thaliana* TEA-GCN (i.e. $\mathbf{Z}$(Co-exp.) ≥ 1.5). The majority of identified edges (293 out of 400 edges; 73.25%; Table S25) could be annotated with at least one lemma, and these annotations largely correspond with the known functions of the gene pairs (Figure 7C; Table S23). To exemplify, we selected two well-characterized transcription factors, ABSSISIC ACID INSENSITIVE 5 (ABI5) and MALE STERILE 188 (MS188). Five positive edges of *ABI5* (connected to *GEA6*, *EM1*, *SDH2-3,* and *AT5G66400)* were annotated with the "ABA" co-expression context (Figure 7C; Table S26), which is a common acronym for the abscisic acid phytohormone that mediates important physiological processes such as seed dormancy maintenance and abiotic stress response[69–71]. This is consistent with the prominent regulatory role of ABI5 in ABA signaling[57,72,73]. Interestingly, *ABI5* edges connecting *AT5G66400, EM1*, and *GEA6* wereannotated with experimental contexts reflecting low-light/dark growth conditions (i.e.

"Shade", "Etiolate", "Darkness"; Figure 7C; Table S26), which represents novel information that
can be used to further dissect the regulatory wirings underpinning ABA signaling of ABI5. This is
supported by recent studies that have revealed darkness-specific regulatory players such as
COP1 E3 ligase[74] and phytochrome-interacting factors[75] in ABA-induced signaling by modulating
ABI5 activity. In another example, all three edges of *MS188* (connected to *FAR2*, *MYB35*, and
MS1) were annotated with experimental contexts related to floral tissues (i.e. "Flower",
"Flowering", "Inflorescence") which is congruent with MS188's known function in anther
developmental processes[76–78] (e.g. tapetum development, pollen exine formation, and pollen
callose development; Figure 7C; Table S26). Thus, TEA-GCN, when combined with NLP
approaches, produces interpretable edges that allow the elucidation of the experimental context,
further boosting gene function prediction approaches.

550
Figure 7. Annotating co-expression edges of *A. thaliana* TEA-GCN with experimental contexts. A) The metadata of RNA-seq
samples are broken down into lemmas using the "en_core_web_sm" natural language processing (NLP) model from spaCy. Lemmas
of RNA-seq samples belonging to the same dataset partition are then combined. For each partition, occurrences of different lemmas
are counted, and enriched lemmas were determined by comparing the observed lemma occurrences to permuted occurrences when

partition assignments of RNA-seq samples are randomly shuffled. As a result, dataset partitions can be annotated by overrepresented

lemmas that likely denote the experimental conditions or tissues which samples in the dataset partition were predominantly generated

from. **B)** Co-expression edges of TEA-GCN can be annotated with lemmas enriched with an approach similar to Gene-set Enrichment

Analysis (GSEA). The ensemble co-expression strength of a given TEA-GCN edge is first deconvoluted into co-expression strengths

obtained from the different dataset partitions, and these strengths are then used to rank dataset partitions. This facilitates the

calculation of enrichment scores as seen in GSEA. **C**) Lemma enrichment analysis of edges connecting *ABSCISIC ACID*

INSENSITIVE 5 (*ABI5*) and *MALE STERILE 188* (*MS188*) transcription factors to their known targets. Running-sum profiles for

enrichment scores and annotated partition rankings of the co-expression context lemmas detected for each of the presented edges

are shown. Line colors of running sum profiles correspond to colors of experimental context lemmas.

Discussion

Gene co-expression networks (GCNs) generated from transcriptomic datasets can elucidate the intricate co-regulatory and co-functional relationships between genes thus enabling gene function prediction. Consequently, GCNs have proven invaluable as a hypothesis generation tool to characterize genes in non-model species whose functions are hard to predict based on sequence similarity against characterized homologs in well-studied models (i.e., members of large gene families, orphan and lineage-specific genes/proteins)[79–83]. Due to the increased affordability of Next-generation Sequening (NGS), the number of species with transcriptomic data is rapidly increasing, which presents an unprecedented opportunity to unravel the evolutionary dynamics of gene function across diverse organisms via the construction of GCNs. However, current GCN construction methodologies [4,28,84,85] were developed for use on small in-house sequenced datasets or manually curated public RNA-seq samples and therefore, are not well equipped to handle and exploit the various complexities of public transcriptome data (e.g. high heterogeneity, sample imbalance, batch effects, poor sample quality). To address this unmet need, we developed TEA-GCN, an ensemble methodology centered around determining gene co-expression based on local co-expression observed in different dataset partitions. We show that TEA-GCN is superior to the widely-used method of calculating PCC across the whole dataset[28,84,86,87], and the state-of-the-art "Subbagging" method[16,17], across three model organisms of vastly different organizational/biological complexities and evolutionary lineages.

Through the characterization of co-expression edges, we show that TEA-GCN excels in capturing co-expression relationships that manifest in certain conditions and/or in specific tissues which enables the discovery of complex regulatory mechanisms that can be elusive. Inspired by the growing field of Explainable Artificial intelligence[88] (XAI) towards explaining the inner workings of complex machine-learning models, we demonstrated TEA-GCN's ability to explain experimental contexts underpinning gene co-expression through the use of NLP. This was possible by capitalizing on our method's ability for capturing condition/tissue-specific co-expression. Thus, we have showed that TEA-GCN can not only outperform more complex state-of-the-art methods, but also for the first time ever, report biologically significant information as experimental contexts that can supercharge gene function prediction. We made the TEA-GCN method available for use via https://github.com/pengkenlim/TEA-GCN as a easy to install and well documented standalone program for users to generate high quality GCNs from their own large transcriptomic datasets.

Methods

Generation and preprocessing of *S. cerevisiae*, *A. thaliana,* and *H. sapiens* public transcriptomic datasets

S. cerevisiae and *H. sapiens* mRNA sequences were downloaded from the NCBI RefSeq annotations of their reference genome assemblies (GCF_000146045.2 and GCF_000001405.40). Representative mRNA sequences of protein-coding genes (primary transcripts) for *S. cerevisiae* and *H. sapiens* were then obtained by selecting the longest transcript isoform for every protein-coding gene loci. Primary transcripts of *A. thaliana* were obtained by downloading the "Araport11_cdna_20220914_representative_gene_model.gz" file from TAIR database[89]. 6,612 *S. cerevisiae*, 19,379 *H. sapiens*, and 27,562 *A. thaliana* primary transcrtipts were used as pseudoalignment references for gene expression estimation of RNA-seq sample Illumina sequencing reads (read data) downloaded from ENA using Kallisto's quant functionality in single-end mode with default parameters (ref). Fastq files of 188,189, 66,048, and 142,828 RNA-seq samples were download and pseudoaligned for *S. cerevisiae, A. thaliana, and H. sapiens,* respectively. Transcripts per million (TPM) values of primary transcripts of RNA-seq samples estimated by Kallisto were used as gene expression abundances to construct gene expression matrices for the different species. RNA-seq samples with poor read alignment were identified based its %_pseudoaligned statistic (reported by Kallisto) below a certain threshold (5% for *H. sapiens*; 20% for *S. cerevisiae* and *A. thaliana*) as described[90–92] (Table S10 to 12). After removal RNA-seq samples with poor read alignment, gene expression matrices comprising of gene expression abundances from 150,096 (79.8%), 57,759 (87.5%), and 95,553 (66.9%) retained RNA-seq samples were designated as the public transcriptomic datasets for *S. cerevisiae, A. thaliana* and *H. sapiens,* respectively.

Dataset partitioning

TPM values of each RNA-seq sample in the gene expression matrices were standardized to unit variance (within-sample standardization) to control for batch effects at the sample level as described[33,90]. Next, the standardized gene expression values were subjected to dimension reduction using Principal Component Analysis (PCA; decomposition.IncrementalPCA function from scikit-learn version 1.3.0) where only the first 1,000 Principal components (PCs) were used as high-level features to embed RNA-seq samples in PC space (PC embeddings) that describes their relative transcriptome similarity between each other. PC embeddings of RNA-seq samples

626 were then used as features for k-means clustering (cluster.Kmeans function from scikit-learn
627 version 1.3.0) using k-means++ as initialization method using random seed of 42 (init = k-
628 means++, n_init = 5, max_iter = 1000, random_state = 42).

Preselection of partitioning granularities (*ks*) for optimal *k* determination

For each species's public transcriptomic dataset, at least 50 *k* values were preselected at fixed
intervals based on the silhouette coefficient, an internal clustering metric commonly used to
determine the clustering performance of unsupervised clustering algorithms like k-means[32]. To
ensure preselected *k*s were sampled at regular intervals across the *k* range from 2 to 1000, we
sectioned said range into *k* windows of length 19. For each *k* window, K-means clustering was
iteratively used to generate dataset partitions, and silhouette coefficients was calculated from the
resultant partition assignments. *k*s that produced the highest silhouette coefficient in each window
were selected (Table S13 to 13).

Data sources for establishing positive edges

S. cerevisiae TF positive edges are generated by downloading activation regulatory links from
the YEASTRACT+ database[93] (http://www.yeastract.com/) that fufill both"Documented" and
"Potential" criteria. According to the authors, "Documented " regulatory links are infered from
experimental transcription factor-binding evidence while "Potential" regulatory links connects
transcription factors with target genes that contains corresponding consensus sequences in
upstream promoter region[93]. *A. thaliana* TF positive edges are generated by downloading
activation regulatory links from the PlantTFDB database[94] (https://planttfdb.gao-lab.org/). *H.*
sapiens TF positive edges are generated by downloading activation regulatory links from
TRRUST version 2 database[95] (https://www.grnpedia.org/trrust/) and then eliminating regulatory
links that does not connect to any known *H. sapiens* transcription factors. *H. sapiens* transcription
factors are obtained from the HumanTFs database[96] (https://humantfs.ccbr.utoronto.ca/). GO
annotations for S. *cerevisiae*, *A. thaliana* and *H. sapiens* were downloaded from EBI Gene
Ontology Annotation (GOA) Database[97], The Arabidopsis Information Resource[89] (TAIR), and
Saccharomyces Genome Database[98] (SGD), respectively. Pathway annotations for S.
cerevisiae, *A. thaliana* and *H. sapiens* were webscraped from BioCyc[99], PlantCyc[38], and
HumanCyc[100] databases, respectively.

Correlation coefficients and mutual rank calculation

PCC, SCC, and Bicor were calculated based on previous publications detailing their use in measuring gene coexpression[1,27] (ref). In the event where correlation coefficients cannot be determined (e.g. expression values of one gene are all 0), correlation coefficients are set to 0. MR transformation of determined co-expression strengths of a GCN are calculated as described[35].

Aggregation functions

The mathematical description of the 3 ensemble aggregation functions, namely the Maximum (Max), Average (Avg), and Rectified Average(RAvg) are detailed below.

Let $V = [v_1, v_2, v_3, \ldots, v_n]$ and be a vector of co-expression strengths between a given pair of genes from different (n) GCNs. The Max aggregation method involves taking the maximum co-expression strength value across co-expression strengths in V and can be expressed as a function where:

$$Max(V) = max_i v_i$$

The Avg aggregation method involves taking the arithmetic mean of V and can be expressed as a function where:

$$Avg(V) = \frac{\sum_i v_i}{n}$$

The RAvg aggregation method involves first rectifying negative edges in V to zero. The arithmetic mean is then calculated from the resultant rectified vector, W. RAvg can be expressed as a function where:

$$W = \frac{|V| + V}{2}$$

$$RAvg(V) = \frac{\sum_i w_i}{n}$$

Standardizing co-expression strengths within GCNs

To compare the co-expression strengths of edges connecting the same gene pair across different GCNs, the co-expression strengths of edges were standardized before comparison. Because different CGNs do not necessarily contain the same genes and edges, the co-expression strengths of each GCN were standardized based on the mean and standard deviation of the co-

684　expression strengths from a subset of common edges shared between the GCNs. The

685　mathematical description of said standardization is detailed below.

686　　　　Let $X = [x_1, x_2, x_3 \ldots, x_N]$ be a vector of co-expression strengths of common edges in one

687　of the GCNs to be compared. The mean co-expression strength (μ) and standard deviation (σ) of

688　X can be calculated as follows:

690
$$\mu = \frac{\sum_i x_i}{N}$$

691
$$\sigma = \sqrt{\frac{\sum(X - \mu)}{N}}$$

692　　　　The co-expression strength of any given edge in the GCN (x_i) can then be standardized

693　as follows:

694
$$Z(x_i) = \frac{x_i - \mu}{\sigma}$$

For interpretability, **Z**(co-exp.) is used to refer to standardized co-expression strengths

throughout the text.

Inferring *A. thaliana* GRNs using GENIE3

The Python implementation of GENIE3[24] was used to infer GRNs used in this study with default

settings (i.e. ntrees = 1000, tree-method = Random Forest, K = square root method). A list of 324

A. thaliana transcription factors (Table S21) was obtained from PlantTFDB[94] and used as

regulators for GENIE3.

Downloading of *A. thaliana* RNA-seq metadata

The metadata of RNA-seq samples was downloaded from ENA via the ENA Browser API using

"sample_description", "dev_stage", "study_title", "sample_title", "tissue_lib" and "tissue_type" as

search fields (Table S13 to 15).

Identifying overrepresented lemmas in *A. thaliana* dataset partitions

The "en_core_web_sm" NLP model (version 3.7.1) from the SpaCy (version 3.7.1) Python

package was used to lemmatized metadata of *A. thaliana* RNA-seq samples. Before

lemmatization, the metadata of RNA-seq samples was preprocessed. First, metadata returned

from all search fields of every RNA-seq sample were concatenated. Next, stop words (e.g. about,

quite, the) were detected from the concatenated metadata using the token.is_stop attribute of

word tokens generated by the "en_core_web_sm" model. Digits and punctuations were also

removed from the concatenated metadata.

To obtain overrepresented lemmas from the different dataset partitions, metadata lemmas

of RNA-seq samples belonging to the same partition were grouped and combined to represent

lemmas of each dataset partition. The occurrence of lemmas in each dataset partition was then

counted and stored as observed lemma occurrences. Permuted lemma occurrences were

similarly obtained by repeating the counting process on dataset partition lemmas combined from

randomly shuffled sample-partition assignments. A null hypothesis p-value indicating the

overrepresentation significance of a particular lemma in a dataset partition can be mathematically

defined below.

Let $S = [s_1, s_2, s_3, \ldots s_n]$ be a vector of permuted occurrences of a lemma in a dataset

partition from n random shufflings of sample-partition assignments and o be the observed

occurrence. The p-value (P) can be calculated as follows:

725
$$P = \frac{1 + \sum_{i=1}^{n} I(s_i \geq o)}{n}$$

Where I is an indicator function that takes a value of 1 or 0 if the statement is true or false

respectively.

P-values determined for every lemma in each dataset partition were then subjected to

multiple-testing correction using the Benjamini-Hochberg method[101], yielding Benjamini-Hochberg

corrected p-values (p_BH). Based on 1000 permutations, 2,333 overrepresented lemmas (p_BH

≤ 0.05, o > 20) were identified for *A. thaliana* dataset partitions . After manual selection to remove

lemmas that are unlikely to contribute to biological insight (e.g. "arabidopsis", "bottle", "clean",

"device", etc.), 276 lemma labels remain (Table S23).

Enrichment-based analyses

GSEA of co-expression neighborhoods of selected *A. thaliana* transcription factors was

implemented by the GSEApy python package[67] (version 1.1.2). For each transcription factor,

genes were ranked based on co-expression strength. The ranks were centered and gene sets

were constructed using *A. thaliana* GO annotations (biological process catergory). Together, the

centered ranks and constructed gene sets were used as input for the prerank sub-command of

GSEApy (weight = 1). Annotating edges between co-expressed genes with experimental

context lemmas also uses the prerank sub-command of GSEApy in the same way. However,

centered partition rankings determined from ranking dataset partitions based on co-expression

strengths after Coefficient Aggregation were used in place of gene rankings. Instead of gene

sets, sets of dataset partitions for which lemma labels were assigned were used.

Data and code availability

TEA-GCN is available at https://github.com/pengkenlim/TEA-GCN for the scientific community to generate high-quality GCNs from their transcriptomic datasets. S. cerevisiae, A. thaliana, and H. sapiens TEA-GCNs generated by this study can be found at FigShare DOIs 10.6084/m9.figshare.26347018, 10.6084/m9.figshare.26347078, and 10.6084/m9.figshare.26347297, respectively. Transcriptomic datasets in the form of gene expression matrices used in this study can be found under 10.6084/m9.figshare.26346916, 10.6084/m9.figshare.26346964, and 10.6084/m9.figshare.263470105, respectively. Supplemental tables referenced in this study are available at FigShare DOI 10.6084/m9.figshare.26347507.

References

1. Usadel, B. *et al.* Co-expression tools for plant biology: opportunities for hypothesis generation and caveats. *Plant Cell Environ.* **32**, 1633–1651 (2009).

2. van Dam, S., Võsa, U., van der Graaf, A., Franke, L. & de Magalhães, J. P. Gene co-expression analysis for functional classification and gene-disease predictions. *Brief. Bioinform.* **19**, 575–592 (2018).

3. Lim, P. K., Zheng, X., Goh, J. C. & Mutwil, M. Exploiting plant transcriptomic databases: Resources, tools, and approaches. *Plant Commun.* **3**, 100323 (2022).

4. Rao, X. & Dixon, R. A. Co-expression networks for plant biology: why and how. *Acta Biochim. Biophys. Sin.* **51**, 981–988 (2019).

5. Aoki, K., Ogata, Y. & Shibata, D. Approaches for extracting practical information from gene co-expression networks in plant biology. *Plant Cell Physiol.* **48**, 381–390 (2007).

6. Rhee, S. Y. & Mutwil, M. Towards revealing the functions of all genes in plants. *Trends Plant Sci.* **19**, 212–221 (2014).

7. Julca, I., Tan, Q. W. & Mutwil, M. Toward kingdom-wide analyses of gene expression. *Trends Plant Sci.* **28**, 235–249 (2023).

8. Burgin, J. *et al.* The European Nucleotide Archive in 2022. *Nucleic Acids Res.* **51**, D121–

D125 (2023).

9. Moreno, P. *et al.* Expression Atlas update: gene and protein expression in multiple species.

Nucleic Acids Res. **50**, D129–D140 (2022).

10. Ruprecht, C., Vaid, N., Proost, S., Persson, S. & Mutwil, M. Beyond Genomics: Studying

Evolution with Gene Coexpression Networks. *Trends Plant Sci.* **22**, 298–307 (2017).

11. Ovens, K., Eames, B. F. & McQuillan, I. Comparative Analyses of Gene Co-expression

Networks: Implementations and Applications in the Study of Evolution. *Front. Genet.* **12**,

695399 (2021).

12. Ballouz, S., Verleyen, W. & Gillis, J. Guidance for RNA-seq co-expression network

construction and analysis: safety in numbers. *Bioinforma. Oxf. Engl.* **31**, 2123–2130 (2015).

13. Vandenbon, A. Evaluation of critical data processing steps for reliable prediction of gene co-

expression from large collections of RNA-seq data. *PloS One* **17**, e0263344 (2022).

14. Gupta, C. & Pereira, A. Recent advances in gene function prediction using context-specific

coexpression networks in plants. *F1000Research* **8**, F1000 Faculty Rev-153 (2019).

15. Pearce, S., Ferguson, A., King, J. & Wilson, Z. A. FlowerNet: a gene expression correlation

network for anther and pollen development. *Plant Physiol.* **167**, 1717–1730 (2015).

16. Obayashi, T., Kodate, S., Hibara, H., Kagaya, Y. & Kinoshita, K. COXPRESdb v8: an animal

gene coexpression database navigating from a global view to detailed investigations.

Nucleic Acids Res. **51**, D80–D87 (2023).

17. Obayashi, T., Hibara, H., Kagaya, Y., Aoki, Y. & Kinoshita, K. ATTED-II v11: A Plant Gene

Coexpression Database Using a Sample Balancing Technique by Subagging of Principal

Components. *Plant Cell Physiol.* **63**, 869–881 (2022).

18. Lin, H., Yu, J., Pearce, S. P., Zhang, D. & Wilson, Z. A. RiceAntherNet: a gene co-

expression network for identifying anther and pollen development genes. *Plant J. Cell Mol.*

Biol. **92**, 1076–1091 (2017).

19. Lancour, D. *et al.* Analysis of brain region-specific co-expression networks reveals clustering

of established and novel genes associated with Alzheimer disease. *Alzheimers Res. Ther.*

12, 103 (2020).

20. Liesecke, F. *et al.* Improved gene co-expression network quality through expression dataset

down-sampling and network aggregation. *Sci. Rep.* **9**, 14431 (2019).

21. Wong, D. C. J. Network aggregation improves gene function prediction of grapevine gene

co-expression networks. *Plant Mol. Biol.* **103**, 425–441 (2020).

22. Arshad, Z. & McDonald, J. F. A computational approach to generate highly conserved gene

co-expression networks with RNA-seq data. *STAR Protoc.* **3**, 101432 (2022).

23. Liesecke, F. *et al.* Ranking genome-wide correlation measurements improves microarray

and RNA-seq based global and targeted co-expression networks. *Sci. Rep.* **8**, 10885

(2018).

24. Huynh-Thu, V. A., Irrthum, A., Wehenkel, L. & Geurts, P. Inferring Regulatory Networks from

Expression Data Using Tree-Based Methods. *PLOS ONE* **5**, e12776 (2010).

25. Hou, J. *et al.* Distance correlation application to gene co-expression network analysis. *BMC*

Bioinformatics **23**, 81 (2022).

26. Song, L., Langfelder, P. & Horvath, S. Comparison of co-expression measures: mutual

information, correlation, and model based indices. *BMC Bioinformatics* **13**, 328 (2012).

27. Hardin, J., Mitani, A., Hicks, L. & VanKoten, B. A robust measure of correlation between two

genes on a microarray. *BMC Bioinformatics* **8**, 220 (2007).

28. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

29. Yue, L. *et al.* Gene co-expression network analysis of the heat-responsive core

transcriptome identifies hub genes in Brassica rapa. *Planta* **253**, 111 (2021).

30. Li, J.-X. *et al.* Investigation of hub gene associated with the infection of Staphylococcus

aureus via weighted gene co-expression network analysis. *BMC Microbiol.* **21**, 329 (2021).

31. Lv, L. *et al.* Gene co-expression network analysis to identify critical modules and candidate

genes of drought-resistance in wheat. *PloS One* **15**, e0236186 (2020).

32. Arthur, D. & Vassilvitskii, S. k-means++: the advantages of careful seeding. in *Proceedings*

of the eighteenth annual ACM-SIAM symposium on Discrete algorithms 1027–1035 (Society

for Industrial and Applied Mathematics, USA, 2007).

33. Zheng, X., Lim, P. K., Mutwil, M. & Wang, Y. A method for mining condition-specific co-

expressed genes in Camellia sinensis based on k-means clustering. *BMC Plant Biol.* **24**,

373 (2024).

34. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq

quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

35. Obayashi, T. & Kinoshita, K. Rank of correlation coefficient as a comparable measure for

biological significance of gene coexpression. *DNA Res. Int. J. Rapid Publ. Rep. Genes*

Genomes **16**, 249–260 (2009).

36. Aravind, L., Anantharaman, V. & Venancio, T. M. Apprehending multicellularity: regulatory

networks, genomics, and evolution. *Birth Defects Res. Part C Embryo Today Rev.* **87**, 143–

164 (2009).

37. Anantharaman, V., Iyer, L. M. & Aravind, L. Comparative Genomics of Protists: New Insights

into the Evolution of Eukaryotic Signal Transduction and Gene Regulation*. *Annu. Rev.*

Microbiol. **61**, 453–475 (2007).

38. Hawkins, C. *et al.* Plant Metabolic Network 15: A resource of genome-wide metabolism

databases for 126 plants and algae. *J. Integr. Plant Biol.* **63**, 1888–1905 (2021).

39. Delli-Ponti, R., Shivhare, D. & Mutwil, M. Using Gene Expression to Study Specialized

Metabolism-A Practical Guide. *Front. Plant Sci.* **11**, 625035 (2020).

40. Brockington, S. F. *et al.* Lineage-specific gene radiations underlie the evolution of novel

betalain pigmentation in Caryophyllales. *New Phytol.* **207**, 1170–1180 (2015).

41. Wink, M. Evolution of secondary metabolites from an ecological and molecular phylogenetic

perspective. *Phytochemistry* **64**, 3–19 (2003).

42. Lim, P. K., Julca, I. & Mutwil, M. Redesigning plant specialized metabolism with supervised

machine learning using publicly available reactome data. *Comput. Struct. Biotechnol. J.* **21**,

1639–1650 (2023).

43. Colinas, M. & Goossens, A. Combinatorial Transcriptional Control of Plant Specialized

Metabolism. *Trends Plant Sci.* **23**, 324–336 (2018).

44. Fang, C., Fernie, A. R. & Luo, J. Exploring the Diversity of Plant Metabolism. *Trends Plant*

Sci. **24**, 83–98 (2019).

45. Voß, U., Bishopp, A., Farcot, E. & Bennett, M. J. Modelling hormonal response and

development. *Trends Plant Sci.* **19**, 311–319 (2014).

46. Zhang, Y., Berman, A. & Shani, E. Plant Hormone Transport and Localization: Signaling

Molecules on the Move. *Annu. Rev. Plant Biol.* **74**, 453–479 (2023).

47. Verma, V., Ravindran, P. & Kumar, P. P. Plant hormone-mediated regulation of stress

responses. *BMC Plant Biol.* **16**, 86 (2016).

48. Prieto, M. A., López, C. J. & Simal-Gandara, J. Glucosinolates: Molecular structure,

breakdown, genetic, bioavailability, properties and healthy and adverse effects. *Adv. Food*

Nutr. Res. **90**, 305–350 (2019).

49. Barba, F. J. *et al.* Bioavailability of Glucosinolates and Their Breakdown Products: Impact of

Processing. *Front. Nutr.* **3**, 24 (2016).

50. Johnson, I. T. Glucosinolates: bioavailability and importance to health. *Int. J. Vitam. Nutr.*

Res. Int. Z. Vitam.- Ernahrungsforschung J. Int. Vitaminol. Nutr. **72**, 26–31 (2002).

51. Sønderby, I. E., Geu-Flores, F. & Halkier, B. A. Biosynthesis of glucosinolates--gene

discovery and beyond. *Trends Plant Sci.* **15**, 283–290 (2010).

52. Paul, S., Geng, C.-A., Yang, T.-H., Yang, Y.-P. & Chen, J.-J. Phytochemical and Health-

Beneficial Progress of Turnip (Brassica rapa). *J. Food Sci.* **84**, 19–30 (2019).

53. Higdon, J. V., Delage, B., Williams, D. E. & Dashwood, R. H. Cruciferous Vegetables and

Human Cancer Risk: Epidemiologic Evidence and Mechanistic Basis. *Pharmacol. Res. Off.*

J. Ital. Pharmacol. Soc. **55**, 224–236 (2007).

54. Latté, K. P., Appel, K.-E. & Lampen, A. Health benefits and possible risks of broccoli – An

overview. *Food Chem. Toxicol.* **49**, 3287–3309 (2011).

55. Pfalz, M. *et al.* Metabolic engineering in Nicotiana benthamiana reveals key enzyme

functions in Arabidopsis indole glucosinolate modification. *Plant Cell* **23**, 716–729 (2011).

56. Bureau, M. & Simon, R. JLO regulates embryo patterning and organ initiation by controlling

auxin transport. *Plant Signal. Behav.* **3**, 145–147 (2008).

57. Collin, A., Daszkowska-Golec, A. & Szarejko, I. Updates on the Role of ABSCISIC ACID

INSENSITIVE 5 (ABI5) and ABSCISIC ACID-RESPONSIVE ELEMENT BINDING

FACTORs (ABFs) in ABA Signaling in Different Developmental Stages in Plants. *Cells* **10**,

1996 (2021).

58. Moreno-Risueno, M. A. *et al.* FUSCA3 from barley unveils a common transcriptional

regulation of seed-specific genes between cereals and Arabidopsis. *Plant J. Cell Mol. Biol.*

53, 882–894 (2008).

59. Skirycz, A. *et al.* DOF transcription factor AtDof1.1 (OBP2) is part of a regulatory network

controlling glucosinolate biosynthesis in Arabidopsis. *Plant J. Cell Mol. Biol.* **47**, 10–24

(2006).

60. Podzimska-Sroka, D., O'Shea, C., Gregersen, P. L. & Skriver, K. NAC Transcription Factors

in Senescence: From Molecular Structure to Function in Crops. *Plants Basel Switz.* **4**, 412–

448 (2015).

61. Chen, Y.-F. *et al.* The WRKY6 transcription factor modulates PHOSPHATE1 expression in

response to low Pi stress in Arabidopsis. *Plant Cell* **21**, 3554–3566 (2009).

62. Li, S., Fu, Q., Huang, W. & Yu, D. Functional analysis of an Arabidopsis transcription factor

WRKY25 in heat stress. *Plant Cell Rep.* **28**, 683–693 (2009).

63. Doll, J. *et al.* Arabidopsis thaliana WRKY25 Transcription Factor Mediates Oxidative Stress

Tolerance and Regulates Senescence in a Redox-Dependent Manner. *Front. Plant Sci.* **10**,

1734 (2019).

64. Jiang, Y. & Deyholos, M. K. Functional characterization of Arabidopsis NaCl-inducible

WRKY25 and WRKY33 transcription factors in abiotic stresses. *Plant Mol. Biol.* **69**, 91–105

(2009).

65. Li, S., Fu, Q., Chen, L., Huang, W. & Yu, D. Arabidopsis thaliana WRKY25, WRKY26, and

WRKY33 coordinate induction of plant thermotolerance. *Planta* **233**, 1237–1252 (2011).

66. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for

interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–

15550 (2005).

67. Fang, Z., Liu, X. & Peltz, G. GSEApy: a comprehensive package for performing gene set

enrichment analysis in Python. *Bioinforma. Oxf. Engl.* **39**, btac757 (2023).

68. Moerman, T. *et al.* GRNBoost2 and Arboreto: efficient and scalable inference of gene

regulatory networks. *Bioinforma. Oxf. Engl.* **35**, 2159–2161 (2019).

69. Kishor, P. B. K., Tiozon, R. N., Fernie, A. R. & Sreenivasulu, N. Abscisic acid and its role in

the modulation of plant growth, development, and yield stability. *Trends Plant Sci.* **27**, 1283–

1295 (2022).

70. Bulgakov, V. P., Wu, H.-C. & Jinn, T.-L. Coordination of ABA and Chaperone Signaling in

Plant Stress Responses. *Trends Plant Sci.* **24**, 636–651 (2019).

71. Nambara, E. & Marion-Poll, A. ABA action and interactions in seeds. *Trends Plant Sci.* **8**,

213–217 (2003).

72. Lopez-Molina, L., Mongrand, S. & Chua, N. H. A postgermination developmental arrest

checkpoint is mediated by abscisic acid and requires the ABI5 transcription factor in

Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 4782–4787 (2001).

73. Pollo-Rodríguez, F., Sánchez-Vicente, I. & Lorenzo, O. The turnover of ABI5 by scaffold

proteins to attenuate ABA signaling. *J. Exp. Bot.* **75**, 3749–3753 (2024).

74. Peng, J. *et al.* COP1 positively regulates ABA signaling during Arabidopsis seedling growth

in darkness by mediating ABA-induced ABI5 accumulation. *Plant Cell* **34**, 2286–2308

(2022).

75. Qi, L. *et al.* PHYTOCHROME-INTERACTING FACTORS Interact with the ABA Receptors

PYL8 and PYL9 to Orchestrate ABA Signaling in Darkness. *Mol. Plant* **13**, 414–430 (2020).

76. Xiong, S.-X. *et al.* The transcription factors MS188 and AMS form a complex to activate the

expression of CYP703A2 for sporopollenin biosynthesis in Arabidopsis thaliana. *Plant J.*

Cell Mol. Biol. **88**, 936–946 (2016).

77. Wang, K. *et al.* The Regulation of Sporopollenin Biosynthesis Genes for Rapid Pollen Wall

Formation. *Plant Physiol.* **178**, 283–294 (2018).

78. Zhang, Z.-B. *et al.* Transcription factor AtMYB103 is required for anther development by

regulating tapetum development, callose dissolution and exine formation in Arabidopsis.

Plant J. Cell Mol. Biol. **52**, 528–538 (2007).

79. Gerlt, J. A., Babbitt, P. C., Jacobson, M. P. & Almo, S. C. Divergent evolution in enolase

superfamily: strategies for assigning functions. *J. Biol. Chem.* **287**, 29–34 (2012).

80. Arendsee, Z. W., Li, L. & Wurtele, E. S. Coming of age: orphan genes in plants. *Trends*

Plant Sci. **19**, 698–708 (2014).

81. Jeffery, C. J. Current successes and remaining challenges in protein function prediction.

Front. Bioinforma. **3**, 1222182 (2023).

82. Jacobson, M. P., Kalyanaraman, C., Zhao, S. & Tian, B. Leveraging structure for enzyme

function prediction: methods, opportunities and challenges. *Trends Biochem. Sci.* **39**, 363–

371 (2014).

83. Jeffery, C. J. Enzymes, pseudoenzymes, and moonlighting proteins: diversity of function in

protein superfamilies. *FEBS J.* **287**, 4141–4149 (2020).

84. Tzfadia, O. *et al.* CoExpNetViz: Comparative Co-Expression Networks Construction and

Visualization Tool. *Front. Plant Sci.* **6**, 1194 (2015).

85. Zainal-Abidin, R.-A., Harun, S., Vengatharajuloo, V., Tamizi, A.-A. & Samsulrizal, N. H.

Gene Co-Expression Network Tools and Databases for Crop Improvement. *Plants Basel*

Switz. **11**, 1625 (2022).

86. Proost, S. & Mutwil, M. CoNekT: an open-source framework for comparative genomic and

transcriptomic network analyses. *Nucleic Acids Res.* **46**, W133–W140 (2018).

87. De Bodt, S., Hollunder, J., Nelissen, H., Meulemeester, N. & Inzé, D. CORNET 2.0:

integrating plant coexpression, protein-protein interactions, regulatory interactions, gene

associations and functional annotations. *New Phytol.* **195**, 707–720 (2012).

88. A., S. & R., S. A systematic review of Explainable Artificial Intelligence models and

applications: Recent developments and future trends. *Decis. Anal. J.* **7**, 100230 (2023).

89. Garcia-Hernandez, M. *et al.* TAIR: a resource for integrated Arabidopsis data. *Funct. Integr.*

Genomics **2**, 239–253 (2002).

90. Lim, P. K., Wang, R. & Mutwil, M. LSTrAP-denovo: Automated Generation of Transcriptome

Atlases for Eukaryotic Species Without Genomes. *Physiol. Plant.* **176**, e14407 (2024).

91. Lim, P. K. *et al.* Bacteria.guru: Comparative Transcriptomics and Co-Expression Database

for Bacterial Pathogens. *J. Mol. Biol.* **434**, 167380 (2022).

92. Villanueva, E. M. F. *et al.* Protist.guru: A Comparative Transcriptomics Database for

Protists. *J. Mol. Biol.* **434**, 167502 (2022).

93. Monteiro, P. T. *et al.* YEASTRACT+: a portal for cross-species comparative genomics of

transcription regulation in yeasts. *Nucleic Acids Res.* **48**, D642–D649 (2020).

94. Jin, J. *et al.* PlantTFDB 4.0: toward a central hub for transcription factors and regulatory

interactions in plants. *Nucleic Acids Res.* **45**, D1040–D1045 (2017).

95. Han, H. *et al.* TRRUST v2: an expanded reference database of human and mouse

transcriptional regulatory interactions. *Nucleic Acids Res.* **46**, D380–D386 (2018).

96. Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650–665 (2018).

97. Camon, E. *et al.* The Gene Ontology Annotation (GOA) Database: sharing knowledge in

Uniprot with Gene Ontology. *Nucleic Acids Res.* **32**, D262–D266 (2004).

98. Wong, E. D. *et al.* Saccharomyces genome database update: server architecture, pan-

genome nomenclature, and external resources. *Genetics* **224**, iyac191 (2023).

99. Karp, P. D. *et al.* The BioCyc collection of microbial genomes and metabolic pathways.

Brief. Bioinform. **20**, 1085–1093 (2019).

100.   Romero, P. *et al.* Computational prediction of human metabolic pathways from the

complete human genome. *Genome Biol.* **6**, R2 (2005).

101.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300 (1995)

Acknowledgement

The authors wish to thank SingAREN (Singapore Advanced Research and Education Network) and Mr. Soh Hwee Jin, Melvin from the High Performance Computing Centre (HPCC) of Nanyang Technological University (NTU) for providing the network infrastructure used for downloading of public sequencing data. The authors also thank Asst. Prof. Jarkko Tapani Salojarvi and Asst. Prof. Goh Wen Bin Wilson from the School of Biological Sciences and Lee Kong Chian School of Medicine, respectively, of NTU for their input in the project. Lastly, the authors thank all members of the Mutwil Lab for their support in the project and comments on the manuscript. P.K.L is supported by the NTU Research Scholarship (RSS). M.M is supported by Singaporean Ministry of Education grant MOE-T2EP30122-0017.

Author information

Authors and Affiliations

School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive, Singapore, 637551, Singapore

Peng ken Lim, Ruoxi Wang, Jenet Princy Antony Velankanni and Marek Mutwil

Contributions

P.K.L and M.M conceived the project and wrote the manuscript. P.K.L developed and programmed code for the method. P.K.L and R.W performed data analysis and generated figures. J.P.A.V helped with the construction of edges. P.K.L. was supervised by M.M, while R.W and J.P.A were supervised by P.K.L. M.M secured funding for the project.

Corresponding authors

Correspondence to Marek Mutwil or Peng Ken Lim.

Supplemental Figures

1014

Figure S1. Overall performances of Partition Aggregation-only ensemble GCNs using different coefficients across different levels of partitioning granularity (k). The colors of solid lines correspond to the median AUCROC-HM and AUCPRC-HM scores of GCNs across 100 evaluations using aggregation functions RAvg or Avg for Partition Aggregation at different ks. Shaded regions of the same color mapping denote upper and lower quartiles of performance scores. The orange line denotes optimal k at which the RAvg Partition Aggregation-only GCNs has the highest sum of median AUCROC-HM and AUCPRC-HM scores.

Figure S2. Comparison of TEA-GCN against "Subagging" ensemble GCNs in terms of performance across different biological aspects in every species. Blue, orange, and green boxes represents performance scores measured for non-ensemble PCC GCNs, "Subagging" GCNs, and TEA-GCNs respectively in each species. Asterisks represents different levels of statistical significance (Two-sample student's T-test) calculated between score distributions of "Subagging" GCNs and TEA-GCNs (NS: P> 0.05, one asterisk: 0.01 ≤ P ≤ 0.05, three asterisks: P ≤ 0.001

1026

1027

Figure S3. Comparing *A. thaliana* TEA-GCN and ATTED-II GCN co-expression neighbourhoods of selected transcription
factors in terms of their functional enrichment profiles. GO terms were selected based on the known functions of the selected
transcription factors. Dot sizes for GO terms correspond to normalized enrichment scores (NES) while inverted triangles represent
CO terms with negative NES scores. For GO terms with positive NES scores, statistical significance in the form of Benjamini-
Hochberg-corrected p-values (p_BH) can be inferred from dot placement on the x-axis. The red dotted lines denote the statistical
significance threshold (p_BH ≥ 0.05).

A

Dataset Partitioning

Expression matrix

K-means clustering of samples

Divide dataset into partitions

Samples

Genes

Two-tier Ensemble Aggregation (TEA)

PCC   SCC   Bicor

Partition 1

Partition 2

Partition 3

Coefficient Aggregation

Partition Aggregaton

TEA-GCN

Calculate correlation between genes in every partition using different coefficients

B

Metabolic Pathway *X*

Compound 1

C   A

Compound 2

B

Compound 3

Met Positive Edges

A —— B

A —✕— C

B —— C

Transcription Factor

H

I

J

TF Positive Edges

H —— I

H —— J

BP-GO: *Y*

E   F

G

MF-GO: *Z*

GO Positive Edges

E —— F

E —— G

F —✕— G

C

Gene Co-expression Network (GCN)

Extract co-exp.

−ve Edges (Randomly Determined)

Extract co-exp.

+ve Edge sets

Split

−ve Edge sets

Calculate performace scores

Edge frequency

AUROC-Met AUPRC-Met

Co-expression strength

Edge frequency

AUROC-TF AUPRC-TF

Co-expression strength

Edge frequency

AUROC-GO AUPRC-GO

Co-expression strength

Harmonic mean

AUROC-HM AUPRC-HM