PA28γ promotes the malignant progression of tumor by elevating mitochondrial function via C1QBP

Jiongke Wang[1#], Yujie Shi[1, 2#], Ying Wang[1], Yingqiang Shen[1], Huan Liu[3], Silu Sun[1], Yimei Wang[1], Xikun Zhou[3], Yu Zhou[1], Xin Zeng[1*], Jing Li[1*], Qianming Chen[1]

[1] State Key Laboratory of Oral Diseases & National Center for Stomatology & National Clinical Research Center for Oral Diseases & Research Unit of Oral Carcinogenesis and Management & Chinese Academy of Medical Sciences, West China Hospital of Stomatology, Sichuan University, Chengdu 610041, Sichuan, China

[2] Department of Stomatology, The First Affiliated Hospital with Nanjing Medical University, Nanjing 210029, Jiangsu, China

[3] State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University and Collaborative Innovation Center for Biotherapy, Chengdu 610041, P. R. China.

#These authors contributed equally to this article.

*Correspondence and requests for materials should be addressed to:

Jing Li

State Key Laboratory of Oral Diseases & National Center for Stomatology & National Clinical Research Center for Oral Diseases & Research Unit of Oral Carcinogenesis and Management & Chinese Academy of Medical Sciences, West China Hospital of Stomatology, Sichuan University, Chengdu 610041, Sichuan, China

Email: lijing1984@scu.edu.cn

Phone: 8615828142267

Fax: 8602885501484

Xin Zeng

State Key Laboratory of Oral Diseases & National Center for Stomatology & National Clinical Research Center for Oral Diseases & Research Unit of Oral Carcinogenesis and Management & Chinese Academy of Medical Sciences, West China Hospital of Stomatology, Sichuan University, Chengdu 610041, Sichuan, China

Email: zengxin22@163.com

Phone: 8615928080862

31   Fax: 8602885503480

PA28γ promotes the malignant progression of tumor by elevating mitochondrial function via C1QBP

Abstract

Proteasome activator 28γ (PA28γ) plays a critical role in malignant progression of various tumors, however, its role and regulation are not well understood. Here, using oral squamous cell carcinoma (OSCC) as main research model, we discovered that PA28γ interacted with complement 1q binding protein (C1QBP), which is dependent on the N-terminus of C1QBP rather than the known functional domain (amino acids 168-213). Notably, we found that PA28γ can enhance C1QBP protein stability in OSCC. Functionally, PA28γ contributes to the malignant progression of OSCC by affecting mitochondrial morphology and oxidative phosphorylation (OXPHOS) through C1QBP in vitro and vivo. Mechanically, PA28γ upregulate the expression of optic atrophy 1 (OPA1), mitofusin 2 (MFN1), mitofusin 2 (MFN2) and the mitochondrial respiratory complex by C1QBP. Moreover, in a clinical cohort of OSCC patients, PA28γ was positively correlated with C1QBP expression and negatively correlated with prognosis. Therefore, C1QBP is also a potential target for the treatment and prognosis of cancer.

Keywords: oral squamous cell carcinoma, PA28γ, C1QBP, protein–protein interactions, mitochondria, oxidative phosphorylation

Introduction

Proteasome activator 28γ (PA28γ), also known as REGγ or PSME3, is a well-known non-ATP-dependent proteasome regulator. PA28γ is overexpressed in some malignant tumors, such as thyroid, colon and breast cancer and is significantly associated with poor prognosis (Mao, Liu, Li, & Luo, 2008; Stadtmueller & Hill, 2011). By regulating the stability of key components within various complex signaling pathways, PA28γ impacts the biological behavior of tumor cells. Our previous studies revealed that the expression level of PA28γ in oral squamous cell carcinoma (OSCC) cancer nest tissues is positively correlated with patient prognosis (J. Li et al., 2015; S. Liu et al., 2018). However, its role and regulation are not well understood.

OSCC is the most common oral and maxillofacial malignant tumor(Siegel, Miller, Wagle, & Jemal, 2023; Sung et al., 2021). The pathogenic factors of OSCC include smoking, drinking and viral infection, such as human papilloma virus (HPV). Therapies for OSCC include surgery, radiotherapy, chemotherapy and immunotherapy. Despite the rapid development of imaging, surgery, radiotherapy and immunotherapy in recent years, along with the emphasis on personalized treatment and multidisciplinary cooperation in the treatment of OSCC, the overall survival rate of OSCC patients has not significantly improved (Peres et al., 2019). The main reasons for unsatisfactory treatment efficacy include invasion of surrounding tissue, early lymph node metastasis and a tendency toward recurrence (Lindemann, Takahashi, Patel, Osman, & Myers, 2018). Metabolic reprogramming is a crucial characteristic of tumors, closely associated with malignant behaviors such as tumor growth, invasion, metastasis, and immune escape (Ohshima & Morii, 2021). Current research on metabolic reprogramming in OSCC primarily focused on how fibroblasts regulate mitochondria to promote malignant tumor progression (X. Li et al., 2022; Zhang et al., 2020). Understanding the molecular events that lead to the occurrence and development of OSCC can aid in developing new tumor-targeted therapies, which hold significant clinical application potential.

The complement 1q binding protein (C1QBP) is a crucial protein for maintaining mitochondrial function, particularly in mitochondrial oxidative phosphorylation (OXPHOS) (Ghebrehiwet, Geisbrecht, Xu, Savitt, & Peerschke, 2019; Q. Wang et al.,

2022). OXPHOS is essential for the production of adenosine triphosphate (ATP),

which is necessary for tumor development. C1QBP has been shown to play a

significant role in cancer progression, as influencing tumor growth, invasion and

metastasis (Bai et al., 2019; Hou, Lu, Wang, & Yang, 2022; Vendramin et al., 2018).

Due to these roles, C1QBP presents a promising therapeutic target for various tumors,

including melanoma, breast cancer, and colorectal cancer (Matsumoto & Bay, 2021).

In OSCC, enhanced mitochondrial OXPHOS function might also be linked to

malignant tumor progression (Vyas et al., 2021; Xiao et al., 2021; Zhu, Liu, Wu, &

Wang, 2021), although the underlying mechanism remain unclear.

In this study, we discovered that PA28γ interacts with C1QBP and that PA28γ can

stabilize C1QBP. This interaction enhances OXPHOS and promotes the growth,

migration and invasion of OSCC cells. Additionally, the expression of PA28γ and

C1QBP is increased and positively correlated in OSCC. Therefore, PA28γ and C1QBP

are potential targets for the treatment and prognosis of cancer.

99 Results

100 *PA28γ interacts with and stabilizes C1QBP*

To explore how PA28γ promotes OSCC progression, we conducted a gene–gene

interaction analysis using the GeneMANIA database. This analysis identified C1QBP

as a significant gene within the interaction network, with the protein-coding gene of

PA28γ also playing a key role in the C1QBP interaction network (Fig. 1A and

Appendix Fig. 1A). Subsequently, pull-down and coimmunoprecipitation analysis

demonstrated that both endogenous and exogenous PA28γ and C1QBP could bind to

each other (Fig. 1B, C and Appendix Fig. 1 B, C). Then, we developed four OSCC

cell lines with stable overexpression of PA28γ using a Flag-PA28γ-expressing

lentivirus and observed that the protein level of C1QBP was higher in these cell lines

compared to control cells (Fig. 1D). Notably, PA28γ upregulated C1QBP protein

levels in a dose-dependent manner (Fig. 1E). To determine whether the upregulated

C1QBP was due to increased stability, cells were pretreated with cycloheximide

(CHX) to inhibit protein synthesis. This treatment minimally affected C1QBP

degradation in the presence of PA28γ (Fig. 1F, G). Furthermore, pull-down analysis

revealed that the PA28γ-C1QBP interaction depends on the N-terminus of C1QBP,

rather than the known functional domain (amino acids 168-213) that bind to
mitochondrial antiviral proteins (Peerschke et al., 2020; Xu, Xiao, Liu, Ren, & Gu,
2009) (Fig. 1H, I). Therefore, our results illustrated that PA28γ could interacts with
C1QBP, in a manner dependent on the N-terminus of C1QBP, and that PA28γ could
stabilizes C1QBP.

PA28γ and C1QBP colocalize in mitochondria and affect mitochondrial functions

To further investigate the physical interaction between PA28γ and C1QBP, we
conducted an immunofluorescence (IF) assay in two OSCC cell lines, as shown these
two proteins were colocalized in the mitochondria (Fig. 2A). Remarkably,
transmission electron microscopy (TEM) images revealed fewer mitochondrial
vacuoles and higher mitochondrial ridge density in PA28γ-overexpressing cells
compared to control cells (Fig. 2B and Appendix Fig. 2A). IF analysis also showed
increased mitochondrial lengths and areas in the PA28γ-overexpressing cells (Fig. 2C,
D). Following that, we conducted a Cell Mito Stress Test to measure the oxygen
consumption rate (OCR) (Fig. 2E, F). The results indicated significantly higher basal
respiration, maximal OCRs and ATP production were significantly greater in
PA28γ-overexpressing cells compared to control cells (Fig. 2G-I and Appendix Fig.
2B-D). Considering that mitochondria are the primary site for ROS generation and
that ROS play a crucial role in carcinogenesis, we measured ROS levels in OSCC
cells. Consistently, ROS levels were significantly higher in PA28γ-overexpressing
cells (Fig. 2J and Appendix Fig. 2E). In addition, experiments conducted in PA28γ-sh
OSCC cells yielded opposite results, further confirming these conclusions (Appendix
Fig. 2F-K).

To demonstrate the carcinogenic abilities of PA28γ and C1QBP, we injected
4MOSC2 cells, a mouse OSCC line, with or without Pa28γ overexpression under the
surface of the tongue in C57BL/6 mice (Appendix Fig. 3A). After 10 days of normal
feeding, the tumors in the Pa28γ-overexpressing groups were significantly larger than
those in the control group. Furthermore, via immunohistochemical (IHC) staining
revealed that that PA28γ overexpression upregulate C1QBP in vivo (Fig. 3A-C).
Consistent with the in vitro experiments, ATP production and ROS generation were
significantly increased (Fig. 3D, E). Additionally, analysis of OSCC xenograft tumor
tissue in nude mice revealed that PA28γ's regulation of C1QBP in OSCC cells is
independent of the immune system (Appendix Fig. 3B, C). In addition, we established

stable Pa28γ-silenced B16 cells, a mouse melanoma cell line, and found that the level

of C1qbp protein was decreased in Pa28γ-silenced B16 cells (Appendix Fig. 3D).

Cells with or without Pa28γ silencing were grafted into the flanks of Pa28γ $^{ko/ko}$ and

Pa28γ $^{wt/wt}$ C57BL/6 mice by subcutaneous injection (Appendix Fig. 3E). Remarkably,

the tumors derived from the silenced Pa28γ cells in Pa28γ $^{ko/ko}$ mice were smaller than

those in the other groups (Fig. 3F-H). The trend of ATP and ROS was consistent with

tumor size (Fig. 3I, J), and the protein levels of C1qbp in the tumors were also

reduced in the Pa28γ-silenced groups (Fig. 3K).

Collectively, these data suggest that PA28γ, which co-localizes with C1QBP in

mitochondria, is involved in regulating mitochondrial morphology and function.

PA28γ regulates mitochondrial OXPHOS by upregulating C1QBP

Next, we explored the protein levels of the mitochondrial respiratory chain complex

and other mitochondrial functional proteins. We found that the levels of complex I and

IV proteins, as well as OPA1, MFN1 and MFN2 proteins, were upregulated in

PA28γ-overexpressing OSCC cells (Fig. 4A, B and Appendix Fig. 4A, B), These

increases could be reversed by C1QBP silencing in PA28γ-overexpressing OSCC

cells (Fig. 4C).

Consistent with this, the OCRs of C1QBP-silenced PA28γ-overexpressing OSCC

cells were markedly lower than those of control cells (Fig. 4D and Appendix Fig.

4C-E). Subsequently, we constructed a series of experiments to detect the biological

behavior of control and PA28γ-overexpressing OSCC cells with or without C1QBP

silencing. C1QBP knockdown significantly attenuated the migration, invasive and

proliferation capabilities previously augmented by PA28γ overexpression (Fig. 4E-H).

These data suggest that PA28γ enhances mitochondrial OXPHOS function through

C1QBP.

C1QBP expression was positively associated with PA28γ expression and was associated with a worse prognosis in patients with OSCC or SKCM

To determine the correlation between PA28γ and C1QBP, IHC staining was performed

in the clinical cohort of oral mucosa carcinogenesis. The results showed that both

C1QBP and PA28γ were notably upregulated during the progression of oral mucosa

carcinogenesis (Fig. 5A, B and Appendix Fig. 5A), and the levels of PA28γ and

C1QBP were positively related (Fig. 5C). Similarly, both C1QBP and PA28γ were

upregulated in metastatic OSCC tissues (Fig. 5D, E and Appendix Fig. 5B), and the

levels of PA28γ and C1QBP were consistently positively correlated in OSCC cohort

(Fig. 5F). Notably, consistent with our previous findings on PA28γ (J. Li et al., 2015),

C1QBP or combining PA28γ and C1QBP could be a negative predator in our

multicenter OSCC clinical cohort (Fig. 5G, H), TCGA HNSC database (Fig. 5I, J) and

TCGA SKCM database (Appendix Fig. 5C, D).

Discussion

Metabolic reprogramming is one of the hallmarks of malignant tumors and is related

to the malignant biological behavior of tumors (Faubert, Solmonson, & DeBerardinis,

2020; Hanahan, 2022; Tsai et al., 2023; Xia et al., 2021). The metabolic phenotype of

tumor cells not only differs from that of normal cells but also dynamically changes

during tumor progression. OSCC is a common head and neck malignancy that is still a

growing global health problem (Johnson et al., 2020). The development of OSCC is a

complex and multifaceted process in which metabolic reprogramming appears to be

important (B. Liu, Si, Wei, Zhang, & Chen, 2023). However, the mechanism of

metabolic reprogramming remains unclear. Excitingly, we found solid evidence that

PA28γ interacts with and stabilizes C1QBP. Aberrantly accumulated C1QBP enhances

the function of mitochondrial OXPHOS and leads to the production of additional ATP

and ROS by activating the expression and function of OPA1, MNF1, MFN2 and

mitochondrial respiratory chain complex proteins. This process results in resulting in

mitochondrial fusion and malignant tumor progression (Fig. 6). Our study highlights

the underlying mechanism by which PA28γ participates in the regulation of OXPHOS

by upregulating C1QBP in OSCC.

C1QBP is also called gClqR, p32, p33 and HABP1, can be distributed inside cells,

on the cell surface, and can also be secreted extracellularly (Egusquiza-Alvarez &

Robles-Flores, 2022). Within cells, C1QBP is primarily located in the mitochondria.

The key functions of mitochondrial C1QBP include maintaining OXPHOS, supplying

energy to cells, and achieving homeostasis of mitochondrial metabolism

(Ghebrehiwet et al., 2019; Q. Wang et al., 2022). C1QBP is highly expressed in cells

that require a large amount of energy, such as the heart, skeletal muscle, testis, ovary,

small intestine and colon (Feichtinger et al., 2017). Sufficient energy is essential for

cancer development, and C1QBP promotes malignant behaviors such as tumor

invasion and metastasis by enhancing OXPHOS function in various cancers
(Raschdorf et al., 2021; Q. Wang et al., 2022). However, there are no reports on the
expression and functions of C1QBP in OSCC. In our study, we found solid evidence
that PA28γ interacts with and stabilizes C1QBP. As a proteasome activator, PA28γ
usually affects the development of tumors by regulating the stability of various
important proteins. For example, PA28γ can promote the degradation of p53, p21
leading to cancer progression (J. Liu et al., 2010). However, the function of
mitochondrial PA28γ may be differ from that of nuclear PA28γ (Mao et al., 2008).
PA28γ interacts with C1QBP and stabilizes C1QBP at the protein level, suggesting
that the binding sites of PA28γ and C1QBP may mask the specific post-translational
modification sites of C1QBP and inhibit its degradation. For example, the
transcriptional coactivator p300 interacts with Smad7 by acetylating two lysine
residues in its N-terminus, which stabilize Smad7 and protect it from TGFβ-induced
degradation (Grönroos, Hellman, Heldin, & Ericsson, 2002). Overall, the specific
mechanism involved in the interaction between PA28γ and C1QBP still needs further
exploration.

Considering that C1QBP is a vital protein for maintaining mitochondrial
metabolism (Tian et al., 2023), we further investigated the function underlying the
malignant progression of OSCC through the interaction between PA28γ and C1QBP
in vitro and in vivo. PA28γ and C1QBP colocalized in the mitochondria, and the
stabilization of C1QBP by PA28γ enhanced mitochondrial morphology and OXPHOS.
This enhancement is crucial for ATP production, and ATP is the primary energy
currency of cellular metabolism. Therefore, PA28γ's regulation of OXPHOS directly
impacts cellular energy metabolism. PA28γ has been reported to activate the
mTORC1 signaling pathway in hepatocellular carcinoma cells to promote glycolysis
and inhibit OXPHOS (Yao et al., 2021). These findings contrast with our study,
potentially due to organ heterogeneity or the complexity of metabolic reprogramming.

In the tissue microenvironment of oral lichen planus, a potentially malignant oral
disorder, PA28γ in epithelial cells can regulate T cell differentiation (Y. Wang et al.,
2024), while PA28γ in CAFs is involved in the crosstalk between stromal cells and
tumor cells (Z. Li et al., 2024). This suggests that PA28γ may interact with the tumor
immune microenvironment. Our phenotypically similar OSCC xenograft tumor
models in both immunocompetent and nude mice indicate that the regulation of

mitochondrial OXPHOS by PA28γ localized in tumor cell mitochondria does not
entirely depend on the immune system (Z. Wang et al., 2019).

Furthermore, we confirmed that PA28γ regulates C1QBP and influences
mitochondrial morphology and function by enhancing the expression of OPA1, MFN1,
MFN2 and the mitochondrial respiratory complex. Mitochondrial fusion, crucial for
oxidative metabolism and cell proliferation, is regulated by the large GTPases MFN1
and MFN2, which fuse with the outer mitochondrial membrane, and OPA1, which
fuses with the inner mitochondrial membrane (Westermann, 2010). OPA1 is an
essential inner mitochondrial membrane protein with multiple functions, including
regulating mitochondrial fusion, crest morphology, mtDNA stability, and interacting
with the mitochondrial respiratory chain to regulate OXPHOS (Del Dotto et al.,
2018).

Finally, we analyzed the expression patterns and significance of PA28γ and
C1QBP in a clinical cohort. PA28γ and C1QBP were negatively correlated with
prognosis in the OSCC clinical cohort. Notably, some studies have shown that
OXPHOS is upregulated in various tumors, with some advanced tumors preferring
OXPHOS metabolically (Ohshima & Morii, 2021; Qiu, Li, & Zhang, 2023).
OXPHOS activity is associated with the recurrence of OSCC (Noh et al., 2023).
Additionally, the enhancement of OXPHOS and ATP production by ROS
proto-oncogene 1, which localizes to mitochondria in OSCC, promotes the OSCC
invasion(Noh et al., 2023; Grimm et al., 2014). These findings are consistent with our
results. Our study not only explain the mechanisms and signaling networks underlying
the oncogenic potential of PA28γ but also contribute to understanding the possible
mechanisms involved in OSCC metabolic reprogramming.

In summary, we found that PA28γ could interact with C1QBP and stabilize
C1QBP. This enhances the function of OXPHOS, leading to increased production of
ATP and ROS by activating the expression and function of OPA1, MNF1, MFN2 and
the mitochondrial respiratory chain complex. This process promotes mitochondrial
fusion and malignant tumor progression. The expression of PA28γ and C1QBP was
increased and positively correlated in OSCC. High expression of PA28γ and C1QBP
was correlated with poor prognosis in OSCC patients, indicating that both factors
could serve as potential targets for the treatment and prognosis of OSCC.

Materials and methods

Patients and Follow-up

Our study included 4 cohorts. The first cohort included 8 normal controls, 13 patients with oral potentially malignant disorder (OPMD), and 45 patients with OSCC (Appendix Table 1). These patients visited the West China Hospital of Stomatology in 2021. The normal tissues were obtained from patients who underwent maxillofacial plastic surgery. The second cohort for survival analysis included 295 patients diagnosed with primary OSCC tumors who visited West China Hospital of Stomatology, Peking University Hospital of Stomatology and Guangdong Provincial Stomatological Hospital from 2005 to 2009 (Appendix Table 2). None of the patients had cancer in other organs. These patients received regular follow-up for 1-133 months after the operation. The protocol of this study was approved by the ethics committee of the West China Hospital of Stomatology (Approval number: WCHSIRB-D-2020-046). The third cohort, comprising 518 patients with head and neck squamous cell carcinoma (HNSCC), was obtained from The Cancer Genome Atlas (TCGA) database (Appendix Table 3). The last cohort, comprising 458 patients with skin cutaneous melanoma (SKCM), was obtained from TCGA database (Appendix Table 4).

Laboratory Experiments

All animal studies were approved by the Animal Care and Use Committee for the State Key Laboratory of Oral Diseases (Approval number: WCHSIRB-D-2022-032) in compliance with the Guide for the updated Animal Research: Reporting of In Vivo Experiments (ARRIVE) 2.0 guidelines. Other methods are detailed in the *Supplemental Materials and Methods*, including cell culture and cell-related experiments, animal models, mitochondrial function detection, western blot/immunoprecipitation assays, hematoxylin and eosin (H&E) staining, and immunohistochemistry.

Statistical Analysis

The data were analyzed using GraphPad Prism software version 8.0. The statistical data are expressed as the mean ± standard deviation (SD). Chi-square and Fisher exact

tests, Spearman correlation tests, Wilcoxon rank-sum tests, one-way ANOVA and

Student's t tests were used to analyze the data. The Kaplan–Meier method was used

for survival analysis, and the log-rank test was used to evaluate the prognostic value

of C1QBP in OSCC patients. The data are shown as the mean ± standard error, and a

value of $P<0.05$ was considered to indicate statistical significance: $P<0.05*$,

$P<0.01**$, $P<0.001***$, and $P<0.0001****$. Each experiment was repeated 3 times.

References

Bai, Y., Wang, W., Li, S., Zhan, J., Li, H., Zhao, M., . . . Wang, J. (2019). **C1QBP Promotes Homologous Recombination by Stabilizing MRE11 and Controlling the Assembly and Activation of MRE11/RAD50/NBS1 Complex**. *Mol Cell, 75*(6): 1299-1314.e1296. doi:10.1016/j.molcel.2019.06.023

Del Dotto, V., Fogazza, M., Lenaers, G., Rugolo, M., Carelli, V., & Zanna, C. (2018). **OPA1: How much do we know to approach therapy?** *Pharmacol Res, 131*: 199-210. doi:10.1016/j.phrs.2018.02.018

Egusquiza-Alvarez, C. A., & Robles-Flores, M. (2022). **An approach to p32/gC1qR/HABP1: a multifunctional protein with an essential role in cancer**. *J Cancer Res Clin Oncol, 148*(8): 1831-1854. doi:10.1007/s00432-022-04001-5

Faubert, B., Solmonson, A., & DeBerardinis, R. J. (2020). **Metabolic reprogramming and cancer progression**. *Science, 368*(6487): eaaw5473.. doi:10.1126/science.aaw5473

Feichtinger, R. G., Oláhová, M., Kishita, Y., Garone, C., Kremer, L. S., Yagi, M., . . . Prokisch, H. (2017). **Biallelic C1QBP Mutations Cause Severe Neonatal-, Childhood-, or Later-Onset Cardiomyopathy Associated with Combined Respiratory-Chain Deficiencies**. *The American Journal of Human Genetics, 101*(4): 525-538. doi:10.1016/j.ajhg.2017.08.015

Ghebrehiwet, B., Geisbrecht, B. V., Xu, X., Savitt, A. G., & Peerschke, E. I. B. (2019). **The C1q Receptors: Focus on gC1qR/p33 (C1qBP, p32, HABP-1)**. *Semin Immunol, 45*: 101338. doi:10.1016/j.smim.2019.101338

Grimm, M., Cetindis, M., Lehmann, M., Biegner, T., Munz, A., Teriete, P., . . . Reinert, S. (2014). **Association of cancer metabolism-related proteins with oral carcinogenesis – indications for chemoprevention and metabolic sensitizing of oral squamous cell carcinoma?** *Journal of Translational Medicine, 12*(1): 208. doi:10.1186/1479-5876-12-208

Grönroos, E., Hellman, U., Heldin, C. H., & Ericsson, J. (2002). **Control of Smad7 stability by competition between acetylation and ubiquitination**. *Mol Cell, 10*(3): 483-493. doi:10.1016/s1097-2765(02)00639-1

Hanahan, D. (2022). **Hallmarks of Cancer: New Dimensions**. *Cancer Discov, 12*(1): 31-46. doi:10.1158/2159-8290.cd-21-1059

Hou, G., Lu, Z., Wang, Z., & Yang, X. (2022). **The Mitochondrial Protein C1QBP Promotes Hepatocellular Carcinoma Progression by Enhancing Cell Survival, Migration and Invasion**. *J Cancer, 13*(8): 2477-2489. doi:10.7150/jca.69379

Johnson, D. E., Burtness, B., Leemans, C. R., Lui, V. W. Y., Bauman, J. E., & Grandis, J. R. (2020). **Head and neck squamous cell carcinoma**. *Nat Rev Dis Primers, 6*(1): 92. doi:10.1038/s41572-020-00224-3

Li, J., Feng, X., Sun, C., Zeng, X., Xie, L., Xu, H., . . . Chen, Q. (2015). **Associations between proteasomal activator PA28γ and outcome of oral squamous cell carcinoma: Evidence from cohort studies and functional analyses**. *EBioMedicine, 2*(8): 851-858. doi:10.1016/j.ebiom.2015.07.004

Li, X., Jiang, E., Zhao, H., Chen, Y., Xu, Y., Feng, C., . . . Shang, Z. (2022). **Glycometabolic reprogramming-mediated proangiogenic phenotype enhancement of cancer-associated fibroblasts in oral squamous cell carcinoma: role of PGC-1α/PFKFB3 axis**. *Br J Cancer, 127*(3): 449-461. doi:10.1038/s41416-022-01818-2

Li, Z., Sun, S., Wang, Y., Hua, Y., Liu, M., Zhou, Y., . . . Li, J. (2024). **PA28γ coordinates the cross-talk between cancer-associated fibroblasts and tumor cells to promote OSCC progression via HDAC1/E2F3/IGF2 signaling**. *Cancer Lett, 594*:216962. doi:10.1016/j.canlet.2024.216962.

Lindemann, A., Takahashi, H., Patel, A. A., Osman, A. A., & Myers, J. N. (2018). **Targeting the DNA Damage Response in OSCC with TP53 Mutations**. *J Dent Res, 97*(6): 635-644. doi:10.1177/0022034518759068

Liu, B., Si, W., Wei, B., Zhang, X., & Chen, P. (2023). **PTP4A1 promotes oral squamous cell**

carcinoma (OSCC) metastasis through altered mitochondrial metabolic reprogramming. *Cell Death Discov,* **9(1)**: 360. doi:10.1038/s41420-023-01657-x

Liu, J., Yu, G., Zhao, Y., Zhao, D., Wang, Y., Wang, L., . . . Li, X. (2010). **REGgamma modulates p53 activity by regulating its cellular localization**. *J Cell Sci,* **123**: 4076-4084. doi:10.1242/jcs.067405

Liu, S., Liu, D., Zeng, X., Wang, J., Liu, J., Cheng, J., . . . Chen, Q. (2018). **PA28γ acts as a dual regulator of IL-6 and CCL2 and contributes to tumor angiogenesis in oral squamous cell carcinoma**. *Cancer Lett,* **428**: 192-200. doi:10.1016/j.canlet.2018.04.024

Mao, I., Liu, J., Li, X., & Luo, H. (2008). **REGgamma, a proteasome activator and beyond?** *Cell Mol Life Sci,* **65(24)**: 3971-3980. doi:10.1007/s00018-008-8291-z

Matsumoto, K., & Bay, B. H. (2021). **Role of C1QBP/p32 and its Therapeutic Potential in Breast Carcinoma and other Cancers**. *Curr Med Chem,* **28**(25): 5048-5065. doi:10.2174/0929867328666201231124038

Noh, J. K., Woo, S. R., Kong, M., Lee, M. K., Lee, J. W., Lee, Y. C., . . . Eun, Y. G. (2023). **Gene signature predicting recurrence in oral squamous cell carcinoma is characterized by increased oxidative phosphorylation**. *Mol Oncol,* **17(1)**: 134-149. doi:10.1002/1878-0261.13328

Ohshima, K., & Morii, E. (2021). **Metabolic Reprogramming of Cancer Cells during Tumor Progression and Metastasis**. *Metabolites,* **11(1)**: 28. doi:10.3390/metabo11010028

Peerschke, E., Stier, K., Li, X., Kandov, E., de Stanchina, E., Chang, Q., . . . Adusumilli, P. S. (2020). **gC1qR/HABP1/p32 Is a Potential New Therapeutic Target Against Mesothelioma**. *Front Oncol,* **10**: 1413. doi:10.3389/fonc.2020.01413

Peres, M. A., Macpherson, L. M. D., Weyant, R. J., Daly, B., Venturelli, R., Mathur, M. R., . . . Watt, R. G. (2019). **Oral diseases: a global public health challenge**. *Lancet,* **394(10194)**: 249-260. doi:10.1016/s0140-6736(19)31146-8

Qiu, X., Li, Y., & Zhang, Z. (2023). **Crosstalk between oxidative phosphorylation and immune escape in cancer: a new concept of therapeutic targets selection**. *Cell Oncol (Dordr),* **46(4)**: 847-865. doi:10.1007/s13402-023-00801-0

Raschdorf, A., Sünderhauf, A., Skibbe, K., Ghebrehiwet, B., Peerschke, E. I., Sina, C., & Derer, S. (2021). **Heterozygous P32/C1QBP/HABP1 Polymorphism rs56014026 Reduces Mitochondrial Oxidative Phosphorylation and Is Expressed in Low-grade Colorectal Carcinomas**. *Frontiers in Oncology,* **10**: 631592. doi:10.3389/fonc.2020.631592

Siegel, R. L., Miller, K. D., Wagle, N. S., & Jemal, A. (2023). **Cancer statistics, 2023**. *CA Cancer J Clin,* **73(1)**: 17-48. doi:10.3322/caac.21763

Stadtmueller, B. M., & Hill, C. P. (2011). **Proteasome Activators**. *Molecular Cell,* **41(1)**: 8-19. doi:10.1016/j.molcel.2010.12.020

Sung, H., Ferlay, J., Siegel, R. L., Laversanne, M., Soerjomataram, I., Jemal, A., & Bray, F. (2021). **Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries**. *CA Cancer J Clin,* **71(3)**: 209-249. doi:10.3322/caac.21660

Tian, H., Chai, D., Wang, G., Wang, Q., Sun, N., Jiang, G., . . . Zheng, J. (2023). **Mitochondrial C1QBP is essential for T cell antitumor function by maintaining mitochondrial plasticity and metabolic fitness**. *Cancer Immunol Immunother,* **72(7)**: 2151-2168. doi:10.1007/s00262-023-03407-5

Tsai, C. H., Chuang, Y. M., Li, X., Yu, Y. R., Tzeng, S. F., Teoh, S. T., . . . Ho, P. C. (2023). **Immunoediting instructs tumor metabolic reprogramming to support immune evasion**. *Cell Metab,* **35(1)**: 118-133.e117. doi:10.1016/j.cmet.2022.12.003

Vendramin, R., Verheyden, Y., Ishikawa, H., Goedert, L., Nicolas, E., Saraf, K., . . . Leucci, E. (2018). **SAMMSON fosters cancer cell fitness by concertedly enhancing mitochondrial and cytosolic translation**. *Nat Struct Mol Biol,* **25(11)**: 1035-1046. doi:10.1038/s41594-018-0143-4

Vyas, A., Harbison, R. A., Faden, D. L., Kubik, M., Palmer, D., Zhang, Q., . . . Duvvuri, U. (2021).
Recurrent Human Papillomavirus-Related Head and Neck Cancer Undergoes Metabolic
Reprogramming and Is Driven by Oxidative Phosphorylation. *Clin Cancer Res,* **27(22)**:
6250-6264. doi:10.1158/1078-0432.ccr-20-4789

Wang, Q., Chai, D., Sobhani, N., Sun, N., Neeli, P., Zheng, J., & Tian, H. (2022). **C1QBP regulates**
mitochondrial plasticity to impact tumor progression and antitumor immune response.
Front Physiol, **13**: 1012112. doi:10.3389/fphys.2022.1012112

Wang, Y., Zhang, Q., Deng, X., Wang, Y., Tian, X., Zhang, S., . . . Li, J. (2024). **PA28γ induces**
dendritic cell maturation and activates T-cell immune responses in oral lichen planus.
MedComm, **5(5)**: e561. doi:10.1002/mco2.561

Wang, Z., Wu, V. H., Allevato, M. M., Gilardi, M., He, Y., Luis Callejas-Valera, J., . . . Gutkind, J. S.
(2019). **Syngeneic animal models of tobacco-associated oral cancer reveal the activity of**
in situ anti-CTLA-4. *Nat Commun,* **10(1)**: 5546. doi:10.1038/s41467-019-13471-0

Westermann, B. (2010). **Mitochondrial fusion and fission in cell life and death**. *Nature Reviews*
Molecular Cell Biology, **11(12)**: 872-884. doi:10.1038/nrm3013

Xia, L., Oyang, L., Lin, J., Tan, S., Han, Y., Wu, N., . . . Liao, Q. (2021). **The cancer metabolic**
reprogramming and immune response. *Mol Cancer,* **20(1)**: 28.
doi:10.1186/s12943-021-01316-8

Xiao, L., Hu, Q., Peng, Y., Zheng, K., Zhang, T., Yang, L., . . . Liu, Y. (2021). **TRAP1 suppresses oral**
squamous cell carcinoma progression by reducing oxidative phosphorylation metabolism
of Cancer-associated fibroblasts. *BMC Cancer,* **21(1)**: 1329.
doi:10.1186/s12885-021-09049-z

Xu, L., Xiao, N., Liu, F., Ren, H., & Gu, J. (2009). **Inhibition of RIG-I and MDA5-dependent**
antiviral response by gC1qR at mitochondria. *Proc Natl Acad Sci USA,* **106(5)**: 1530-1535.
doi:10.1073/pnas.0811029106

Yao, L., Xuan, Y., Zhang, H., Yang, B., Ma, X., Wang, T., . . . Zhang, B. (2021). **Reciprocal**
REGγ-mTORC1 regulation promotes glycolytic metabolism in hepatocellular carcinoma.
Oncogene, **40(3)**: 677-692. doi:10.1038/s41388-020-01558-8

Zhang, Z., Gao, Z., Rajthala, S., Sapkota, D., Dongre, H., Parajuli, H., . . . Liang, X. (2020). **Metabolic**
reprogramming of normal oral fibroblasts correlated with increased glycolytic
metabolism of oral squamous cell carcinoma and precedes their activation into
carcinoma associated fibroblasts. *Cell Mol Life Sci,* **77(6)**: 1115-1133.
doi:10.1007/s00018-019-03209-y

Zhu, Z., Liu, Y., Wu, D., & Wang, H. (2021). **Association Between Mitochondrial DNA Copy**
Number and Head and Neck Squamous Cell Carcinoma: A Systematic Review and
Dose-Response Meta-Analysis. *Med Sci Monit,* **27**: e928327. doi:10.12659/msm.928327

Acknowledgments

This work was supported by grants from the National Natural Science Foundation of China (82072999 and 82273320 to J. Li, U19A2005 and 82270986 to X. Zeng, 82201074 to J. Wang), the Province Natural Science Foundation of Sichuan (23NSFSC0145 to J. Li), and the Innovation Research Project of Sichuan University (2022SCUH0029 to J. Li).

Author contributions

J. Wang, Y. Shi, X. Zeng, J. Li contributed to conception and design, data acquisition, analysis, and interpretation, drafted and critically revised the manuscript; Y. Wang, Y. Shen, H Liu contributed to data acquisition; S. Sun, Y. Wang contributed to data analysis and interpretation; X. Zhou, Q. Chen contributed to conception and design, critically revised the manuscript. All the authors gave final approval and agreed to be accountable for all aspects of the work.

Conflict of interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

                           Figure legends

Fig. 1 The interaction between PA28γ and C1QBP. **(A)** The gene network of PA28γ

and C1QBP in the GeneMANIA database. **(B, C)** The interaction between endogenous

PA28γ and C1QBP in HSC-3 cells was verified via immunoprecipitation. **(D)** Western

blot analysis of C1QBP in four OSCC cell lines with PA28γ-overexpressing. **(E)**

Western blot analysis of C1QBP in 293T cells transfected with increasing doses of

Flag-PA28γ. **(F)** 293T cells transfected with Flag-C1QBP with or without Myc-PA28γ

were treated with CHX (100 μg/ml) for the indicated periods of time. **(G)**

Quantification of Flag-C1QBP levels relative to tubulin levels is shown (the data are

representative of 1 experiment with 3 independent biological replicates, *P<0.05). **(H)**

Full-length C1QBP and truncation with deletion of functional domains. **(I)** Pull-down

of 293T cells transfected with Myc-PA28γ and full-length Flag-C1QBP or truncation

mutants of functional domains for 36h.

Fig. 2 PA28γ and C1QBP colocalize in mitochondria and affect mitochondrial

functions *in vitro*. **(A)** Confocal image of IF in two OSCC cell lines. **(B)** TEM

images of PA28γ-overexpressing and control UM1 cells. **(C)** Representative confocal

images of mitochondria in two OSCC cell lines. **(D)** The area and mean branch length

of mitochondria in two OSCC cell lines were measured by ImageJ (the data are

presented as the mean ± SD of 3 independent experiments; **P<0.01, ***P<0.001,

and ****P<0.0001). **(E, F)** OCRs of PA28γ-overexpressing and control UM1 and

HN12 cells were plotted using a Cell Mito Stress Test Kit (the data are presented as

the means ± SDs of 3 independent experiments). **(G-I)** Basal OCRs, maximal OCRs

and ATP production of PA28γ-overexpressing and control UM1 cells measured by the

Cell Mito Stress Test (the data are presented as the means ± SDs of 3 independent

experiments, **P<0.01). **(J)** ROS generation in PA28γ-overexpressing and control

UM1 cells (the data are presented as the means ± SDs of 3 independent experiments;

* P<0.05).

Fig. 3 PA28γ affect mitochondrial functions *in vivo*. **(A)** Representative images of

H&E staining and IHC staining of C1qbp of tongue sections from mice (n=3). **(B)**

Quantification of the sectional area in tumors from the Pa28γ-overexpressing and

control groups (n=3; the data are presented as the means ± SDs of 3 samples;

P<0.05). **(C)** Comparison of the immunoreactive scores (IRSs) of C1qbp antibody

staining in the Pa28γ-overexpressing and control groups (n=3, the data are presented

as the means ± SDs; *P<0.05). **(D, E)** Quantification of ATP production and ROS

levels in tumors from the Pa28γ-overexpressing and control groups (n=3; the data are

presented as the means ± SDs of 3 samples; *P<0.05, ***P<0.001, ****P<0.0001).

(F) Images of the tumors in different groups at the endpoint (n=6). **(G-J)** The volume,

weight, ATP and ROS of tumors in different groups at the endpoint (n=6; the data are

presented as the means ± SDs of 3 independent experiments; *P<0.05, **P<0.01,

***P<0.001, ****P<0.0001). **(K)** Western blot analysis of tumors in different groups.

Fig. 4 PA28γ regulates mitochondrial OXPHOS and cellular biological behavior

via C1QBP. (A, B) Western blot analysis of PA28γ-overexpressing and control UM1

and 4MOSC2 cells. **(C)** Western blot analysis of PA28γ-overexpressing OSCC cells

transfected with siNC or siC1QBP. **(D)** OCRs of C1QBP-silenced and control

PA28γ-overexpressing UM1 and 4MOSC2 cells (the data are presented as the means

± SDs of 3 independent experiments). **(E-H)** Cell migration, invasion and

proliferation in control, C1QBP-silenced, PA28γ-overexpressing and

PA28γ-overexpressing + C1QBP-silenced UM1 cells (the data are presented as the

means ± SDs of 3 independent experiments; *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001).

Fig. 5 The correlation between PA28γ and C1QBP in the carcinogenesis and

development of OSCC. **(A)** Representative IHC staining of PA28γ and C1QBP in

normal (n=8), OPMD (n=13) and OSCC (n=45) samples. **(B)** Comparison of the

immunoreactive scores (IRSs) of C1QBP between the normal, OPMD and OSCC

groups (the data are presented as the means ± SDs; *P<0.05, **P<0.01,

****P<0.0001). **(C)** Spearman correlation analysis was used to test the correlation

between PA28γ and C1QBP in normal, OPMD and OSCC tissues (P<0.0001,

r=0.8227). **(D)** Representative IHC staining of PA28γ and C1QBP in nonmetastatic

(n=27) and metastatic (n=18) OSCC patients. **(E)** Comparison of the IRSs of C1QBP

in the nonmetastatic and metastatic OSCC groups (the data are presented as the means

± SDs; *P<0.05). **(F)** Spearman correlation analysis was used to test the correlation

between PA28γ and C1QBP in OSCC tissues (P<0.0001, *r*=0.6977). **(G)** Kaplan–

Meier analysis of the protein expression of C1QBP in our multicenter OSCC clinical

cohort (n=295, *P*=0.0480). **(H)** Kaplan–Meier analysis of both low or high protein expression of C1QBP and PA28γ in our multicenter OSCC clinical cohort (n=295, *P*=0.0100). **(I)** Kaplan–Meier analysis of the protein expression of C1QBP in TCGA HNSC database (n=259, *P*=0.025). **(J)** Kaplan–Meier analysis of both low or high protein expression of C1QBP and PA28γ in TCGA HNSC database (n=259, *P*=0.033).

Fig. 6 Molecular mechanism through which PA28γ interacts with C1QBP in the malignant progression of tumor. PA28γ can interacts with and stabilize C1QBP, which can activate the expression and function of OPA1, MNF1, MFN2 and mitochondrial respiratory chain complex proteins, resulting in enhanced mitochondrial OXPHOS, mitochondrial fusion and malignant tumor progression.