Δ133p53α and Δ160p53α isoforms of the tumor suppressor protein p53 exert dominant-

negative effect primarily by co-aggregation

Liuqun Zhao[1], Tanel Punga[2], Suparna Sanyal[1*]

[1]Department of Cell and Molecular Biology, Uppsala University, Biomedical Center, SE-75124,

Uppsala, Sweden

[2]Department of Medical Biochemistry and Microbiology, Uppsala University, Biomedical Center, SE-

75123, Uppsala, Sweden

[*]Corresponding author e-mail: suparna.sanyal@icm.uu.se

Abstract

p53 is a tumor suppressor protein with multiple isoforms with shared or specific functions. However,

two of its isoforms, Δ133p53α and Δ160p53α, with large N-terminal deletions, can cause cancer.

These isoforms exert a dominant-negative effect on full-length p53 (FLp53), although the precise

molecular mechanisms are unknown. Here, we investigate the mechanisms of action of Δ133p53α

and Δ160p53α isoforms using chromatin immunoprecipitation, luciferase expression, subcellular

fractionation, immunofluorescence assays, and apoptotic caspase activity assay. Our study elucidates

that these DNA-binding deficient p53 isoforms form hetero-tetrameric complexes with FLp53 and

disrupt FLp53's DNA binding and transcriptional activities when present in a higher proportion than

FLp53 in the tetramer. However, these structurally unstable isoforms promote vigorous protein

aggregation involving FLp53, disrupting its structure and sequestering it in the cytoplasmic and

nuclear aggregates, thereby limiting its availability to function as a transcription activator protein.

Thus, co-aggregation of Δ133p53α and Δ160p53α with FLp53, rather than hetero-tetramerization, is

likely the primary factor contributing to their dominant-negative effect. Modulating the stability and

aggregation of p53 isoforms could be a novel strategy for cancer therapy.

Keywords: p53, Δ133p53α, Δ160p53α, dominant-negative effect, protein aggregation

Introduction

The tumor suppressor protein p53 is an essential transcription activator protein modulating the expression of multiple target genes. By regulating various cellular processes such as cell cycle arrest, apoptosis, senescence, DNA repair, and angiogenesis, p53 prevents cancer development[1, 2]. The intricate p53 pathway is stimulated by various stress stimuli, including DNA damage and oncogene activation, which lead to the stabilization and activation of the p53 protein[3]. Stabilized p53 acts as a transcription activator protein as it can directly bind to its response elements (REs) and recruit different transcription co-activator proteins at various gene promoters[4]. One of the best-known p53 targets is the p21 (also known as CIP1, WAF1) gene promoter, which contains two p53REs and is efficiently activated by the p53 protein[5, 6]. The encoded p21 protein blocks cell cycle progression and thereby executes tumor suppressor activity of the p53 protein[7, 8].

The TP53 gene produces at least twelve distinct p53 isoforms: p53 (α, β, γ), Δ40p53 (α, β, γ), Δ133p53 (α, β, γ), and Δ160p53 (α, β, γ). It has been shown that the p53 isoforms contribute to the regulation of the p53 pathway by regulating the expression of p53 target genes[9, 10, 11]. Notably, p53 isoforms exhibit both overlapping and distinct functions compared to canonical p53[11, 12, 13] and have been reported to promote tumor progression in various cancers[11, 14]. The p53 isoforms are generated through alternative promoter usage, alternative splicing, and alternative transcription starting site usage[9, 15]. These isoforms differ from the full-length (FL) p53 (canonical p53, p53α) by lacking variable segments of the N-terminus (N-terminal isoforms: Δ40, Δ133, and Δ160) and possessing alternative C-terminus (C-terminal isoforms - α, β, and γ). The FLp53 protein consists of four primary functional domains: transactivation domains (TAD-1 and TAD-2), DNA-binding core domain (DBD), nuclear localization sequence (NLS), and the oligomerization domain (OD)[16]. Native FLp53 typically functions as a tetrameric protein, where the individual monomers bind cooperatively to DNA[17, 18]. The C-terminal truncated isoforms (β/γ-isoforms) lacking the OD have lower transactivation activity because of reduced tetramerization efficiency[19, 20, 21, 22]. In contrast, the N-terminal truncated isoforms (α-isoforms), which retain the entire C-terminal OD, can form various complexes with FLp53 and exert dominant-negative effects on p53's transcriptional activity[23, 24, 25]. This study focuses on two large N-terminal deleted p53 isoforms: Δ133p53α and Δ160p53α, the latter has been largely unexplored. For convenience and simplicity, we have written Δ133p53 and Δ160p53 to represent the α isoforms

(Δ133p53α and Δ160p53α) throughout this manuscript. The Δ133p53 protein lacks both TADs and a

part of the first conserved cysteine box of the DBD, whereas the Δ160p53 protein lacks both TADs

and the entire first conserved cysteine box of the DBD[9, 26].

The Δ133p53 isoform exhibit complex biological functions, with both oncogenic and non-oncogenic

potentials. Recent studies demonstrate the non-oncogenic yet context-dependent role of the Δ133p53

isoform in cancer development. Δ133p53 expression has been reported to correlate with improved

survival in patients with TP53 mutations[27, 28], where it promotes cell survival in a non-oncogenic

manner[29, 30], especially under low oxidative stress[31]. Alternatively, other recent evidences emphasize

the notable oncogenic functions of Δ133p53 as it can inhibit p53-dependent apoptosis by directly

interacting with the FLp53 [23, 32]. The oncogenic function of the newly identified Δ160p53 isoform is

less known, although it is associated with p53 mutation-driven tumorigenesis[33] and in melanoma cells'

aggressiveness[34]. Whether or not the Δ160p53 isoform also impedes FLp53 function in a similar way

as Δ133p53 is an open question. However, these p53 isoforms can certainly compromise p53-

mediated tumor suppression by interfering with FLp53 binding to target genes such as p21 and miR-

34a[24, 25] by dominant-negative effect, the exact mechanism is not known.

Protein aggregation has become a central focus of modern biology research and has documented

implications in various diseases, including cancer[35, 36, 37]. Protein aggregates can be of different types

ranging from amorphous aggregates to highly structured amyloid or fibrillar aggregates, each with

different physiological implications. In the case of p53, whether protein aggregation, and in particular,

co-aggregation with large N-terminal deletion isoforms, plays a mechanistic role in its inactivation is

yet underexplored. Interestingly, the Δ133p53β isoform has been shown to aggregate in several

human cancer cell lines[38]. Additionally, the Δ40p53α isoform exhibits a high aggregation tendency in

endometrial cancer cells[39]. Although no direct evidence exists for Δ160p53 yet, these findings imply

that p53 isoform aggregation may play a major role in their mechanisms of actions.

The precise molecular mechanisms involved in the biological function of the Δ133p53 and the

Δ160p53 proteins, especially how they exert dominant-negative effects on FLp53, remain poorly

understood. In this study, we have analyzed the DNA-binding, transcriptional activity, and

oligomerization capacity of the human FLp53, Δ133p53, and Δ160p53 proteins. Our data show that

the Δ133p53 and Δ160p53 proteins are deficient in binding to p53-regulated promoters and inducing

expression of the p53 target genes. Moreover, the Δ133p53 and Δ160p53 isoforms can form hetero-

oligomers with FLp53 and inhibit FLp53 function in a concentration-dependent manner. Two potential

mechanisms for their dominant-negative effects have been suggested: (i) the formation of hetero-

tetramers with FLp53, inhibiting its DNA-binding and transcriptional activity, and (ii) the aggregation

property of the isoforms leading to co-aggregation of the FLp53 protein. Interestingly, while hetero-

tetramerization partially attenuates p53 activity, the co-aggregation of FLp53 with Δ133p53 and

Δ160p53 emerges as the predominant factor contributing to the dominant-negative effect.

Results

Δ133p53 and Δ160p53 proteins exhibit structural destabilization, exposure of hydrophobic

surfaces, and high propensity to aggregation.

The DBD, also known as the core domain of the FLp53 protein (human p53, residues 94-292) has a

structure with an aggregation-prone sequence (residues 251–257)[40] (Figure 1A). The core domain

contains a hydrophobic core of β-sandwich, which consists of two antiparallel β-sheets with four

strands (S1, S3, S8, and S5) and five strands (S6, S7, S4, S9, and S10) respectively[41]

(Supplementary Figure 1A). The DNA-binding surface is constituted by two large loops (L2 and L3)

and a loop–sheet–helix (LSH) motif, which comprises loop L1, a β-sheet formed by the β-hairpin S2–

S2′, the C-terminal residues of strand S10, and the α-helix H2[42] (Supplementary Figure 1A).

Compared to FLp53, the N-terminal deletion isoforms Δ133p53 and Δ160p53 lack the entire TADs

(Figure 1A), which are crucial for recruiting transcription co-activator proteins[43]. In addition, the

Δ133p53 protein is missing the β-strand S1 in the β-sandwich and a part of the LSH motif (L1, S2, and

the N-terminal residue 132 of the strand S2′) in the DBD (Supplementary Figure 1A). The Δ160p53

protein, on the other hand, lacks a part of the β-sandwich (β-strands S1, S3, and N-terminal residues

156-159 of S4) and a part of the LSH motif (L1 and β-hairpin S2–S2′) (Supplementary Figure 1A).

Therefore, the N-terminal deletion affects the structural integrity of both the β-sandwich and the LSH

motif in the core domain.

To assess the structural impact of the N-terminal deletions in FLp53, we analyzed the contact

information derived from the core domain of FLp53 (PDB ID: 3KMD) structure[44]. We observed that

both Δ133p53 and Δ160p53 lack several crucial contacts between the N- and C-terminal regions in

FLp53 (Figure 1B). For Δ133p53, the contact between the L1 and S2 segments in the N-terminal

region and the α-helix H2 in the C-terminal region within the LSH motif is disrupted (Figure 1B, insert

1), suggesting a potential impairment of the stabilizing of DNA-binding interface. For Δ160p53, in

addition to the missing contact between L1, S2 and H2, there are also no contacts between the S2′

segment in the N-terminal region and the S10 and H2 segments in the C-terminal region within the

LSH motif (Figure 1B, insert 2), as well as between S4 and S7 or S9 in the β-sandwich core domain

(Figure 1B, inserts 3 and 4). These contacts are crucial for maintaining the structural integrity and

folding of the core domain. Therefore, both Δ133p53 and Δ160p53 will likely have a partially or fully

destabilized core domain that may impair their DNA-binding ability.

We predicted the monomer structure of Δ133p53 and Δ160p53 by Alphafold2[45] (Supplementary

Figure 1B). Hydrophobic surface analysis revealed that Δ133p53 and Δ160p53 exposed four highly

hydrophobic residues (L252, I254, I255, and L257) within the aggregation-prone sequence, while

FLp53 exposes only one residue (L252) (Figure 1C). This increased hydrophobicity may induce the

aggregation of Δ133p53 and Δ160p53. Moreover, Δ133p53 and Δ160p53 have a partial or complete

structural loss of the β-hairpin S2–S2′ (Supplementary Figure 1A), which restricts access to the

hydrophobic core[46]. This structural alteration results in an enhanced exposure of the hydrophobic core

of the β-sandwich domain compared to FLp53 (Figure 1C). This exposure potentially facilitates the

propensity for protein aggregation, as suggested by previous studies[46, 47] and our *in vitro* aggregation

assay with purified FLp53, Δ133p53 and Δ160p53 proteins (Supplementary Figure 2). In summary,

these findings suggest that the large N-terminal deletion in Δ133p53 and Δ160p53 would compromise

the structural stability of FLp53 and induce aggregation, with Δ160p53 being more severely affected

among the two.

Δ133p53 and Δ160p53 form complexes with FLp53.

To investigate the oligomerization properties of Δ133p53 and Δ160p53, which retain the OD known to

facilitate p53 tetramerization[48], we co-expressed the FLAG-tagged p53 proteins as the baits along

with the V5-tagged FLp53, Δ133p53, and Δ160p53 proteins as preys. The V5-tagged FLp53,

Δ133p53, and Δ160p53 proteins specifically interacted with their corresponding FLAG-tagged

proteins (Figure 2A, lanes 1–6), demonstrating their capacity to form homo-oligomers (Figure 2B). In

addition, interactions between FLp53-FLAG/Δ133p53-V5, FLp53-FLAG/Δ160p53-V5, Δ133p53-

FLAG/Δ160p53-V5, and Δ160p53-FLAG/Δ133p53-V5 were found (Figure 2A, lanes 7–10), suggesting

that they can also form hetero-oligomers (Figure 2B). These findings are consistent with previous

studies indicating that Δ40p53 can hetero-oligomerize with FLp53[49, 50]. To rule out the possibility that

the observed interactions between FLp53 and its isoforms Δ133p53 and Δ160p53 were artifacts

caused by the FLAG and V5 antibody epitope tags, we co-expressed FLAG-tagged FLp53 with

untagged Δ133p53 and Δ160p53. Immunoprecipitation assays demonstrated that FLAG-tagged

FLp53 could still interact with the untagged Δ133p53 and Δ160p53 isoforms (Supplementary Figure 3,

lanes 3 and 4), confirming the formation of hetero-oligomers between FLp53 and its isoforms. These

findings demonstrate that Δ133p53 and Δ160p53 can oligomerize with FLp53 and with each other.

Δ133p53 and Δ160p53 negatively influence FLp53's DNA-binding activity.

Previous studies have shown that the p53 lacking the TADs and the PRD (ΔNp53DBD, residues 94-

393) retain full DNA-binding capacity as the FLp53[51, 52]. In contrast, the Δ133p53 (residues 133-393)

and Δ160p53 (residues 160-393) isoforms additionally lack part of the DBD (residues 94-292). This

deficiency raises questions about the impact of partial DBD deletion on the DNA-binding activity of

p53. To address this question, we performed a chromatin immunoprecipitation (ChIP) assay to

measure the DNA-binding activity of FLp53 and its isoforms Δ133p53 and Δ160p53 to several known

p53 targeted promoters controlling proteins regulating cycle arrest (p21), protein stability (MDM2), and

apoptosis induction (BAX and PUMA) (Figure 3A). Previous studies have classified p53 DNA-binding

affinities. Accordingly, the p21 promoter has the highest affinity, the MDM2 promoter has moderate

affinity, whereas the BAX and PUMA promoters show relatively low p53-binding affinity[53, 54, 55].

Consistent with these findings, our results revealed distinct binding affinities of the FLp53 protein to

these target gene promoters (Figure 3B). Notably, the Δ133p53 and Δ160p53 isoforms demonstrated

a significant loss of DNA-binding activity at these promoters (Figure 3B).

We subsequently hypothesized that Δ133p53 and Δ160p53 might compromise the DNA-binding

ability of FLp53 through hetero-oligomer formation (Figure 2). To study this, we co-transfected FLp53

along with either Δ133p53 or Δ160p53 expressing plasmid and measured FLp53's binding to the

aforementioned promoters. The presence of Δ133p53 or Δ160p53 significantly reduced the binding of

FLp53 to the p21 promoter, which has the highest affinity for FLp53, while having minimal impact on

its interaction with the other promoters (Figure 3C). These findings suggest that Δ133p53 and

Δ160p53 isoforms may selectively impair the ability of FLp53 to bind to high-affinity target promoters.

Δ133p53 and Δ160p53 impede FLp53-mediated transcription activation.

The p53-mediated transcription activation depends on its binding ability to specific p53REs at the target gene promoters[56, 57]. To evaluate the influence of the FLp53 protein interaction with its isoforms, Δ133p53 and Δ160p53, on transcription, we employed a luciferase reporter assay with the p21, MDM2, BAX, and PUMA promoters fused to a firefly luciferase reporter gene. As expected, all four promoters were responsive to FLp53-mediated transcriptional activation (Figure 4A-D). Notably, FLp53 activation of the PUMA promoter was relatively low, possibly due to the lower binding affinity of FLp53 to the PUMA promoter (Figure 3B). In contrast, neither Δ133p53 nor Δ160p53 activated any tested promoters (Figure 4A-D), indicating a lack of functional capability to stimulate transcription from these promoters.

In order to assess how Δ133p53 and Δ160p53 influence FLp53-mediated transcription on different p53-responsive promoters, we co-expressed FLp53 along with increasing amounts of Δ133p53 or Δ160p53 in H1299 cells. We found that co-expression of FLp53 with Δ133p53 or Δ160p53 at a 1:1 ratio did not significantly alter the activation of the p21 and MDM2 promoters compared to FLp53 alone (Figure 4A and B). This result suggests that at an equal protein expression ratio (Figure 2A), these isoforms did not exert a dominant-negative effect on the FLp53 at these promoters. However, a strong reduction in BAX promoter activation was evident under the same experimental condition (Figure 4C), suggesting that the isoforms can disrupt FLp53 function on the BAX promoter. Interestingly, both isoforms enhanced the activation of the PUMA promoter when added with FLp53 in a 1:1 ratio (Figure 4D).

The elevated Δ133p53 protein modulates p53 target genes such as miR-34a and p21, facilitating cancer development[24, 58]. To mimic conditions where isoforms are upregulated relative to FLp53, we increased the ratios to 1:5 and 1:10. Adjusting the ratio of FLp53 to isoforms to 1:5 revealed that higher isoform expression levels significantly impaired FLp53-mediated activation of the p21, MDM2, and BAX promoters (Figure 4A-C). Interestingly, Δ133p53 at a 1:5 ratio had almost no impact on PUMA promoter activity, whereas Δ160p53 reduced it compared to FLp53 alone (Figure 4D). These results indicate that higher expression ratios of Δ133p53 and Δ160p53 relative to FLp53 might be required to exert dominant-negative effects. To test this hypothesis, we escalated the ratio of FLp53 to isoforms to 1:10. As expected, the activity of all four promoters decreased significantly at this ratio

(Figure 4A-D). Notably, Δ160p53 showed a more potent inhibitory effect than Δ133p53 at the 1:5 ratio

on all promoters except for the p21 promoter, where their impacts were similar (Figure 4E-H).

However, at the 1:10 ratio, Δ133p53 and Δ160p53 had similar effects on all transactivation except for

the MDM2 promoter (Figure 4E-H). This differential impact underscores the variability in how p53

isoforms influence FLp53 function across different promoters and expression ratios. Collectively,

these findings demonstrate that isoforms with impaired DNA-binding capability can indeed exert

dominant-negative effects on the function of FLp53, with the magnitude of this impact being promoter-

specific and dependent on the relative expression levels of the proteins.

The Δ133p53 and Δ160p53 proteins affect the subcellular localization and aggregation of

FLp53.

p53 exerts its transcriptional activity in the nucleus, and its appropriate subcellular localization is

crucial for regulating p53 functions[59]. Immunoblotting analysis with isolated nuclear and cytoplasmic

fractions revealed distinct subcellular localization patterns for the FLp53 and its isoforms Δ133p53

and Δ160p53 (Figure 5A and B). The cytoplasm-to-nucleus ratio of Δ133p53 and Δ160p53 was

approximately 1.5-fold higher than that of FLp53 (Figure 5B). These results suggest that Δ133p53 and

Δ160p53 exhibit a greater tendency for cytoplasmic localization.

Immunofluorescence analysis of the cells expressing FLp53 and the two isoforms confirmed the

above-observed protein localization pattern (Figure 5C). The FLp53 protein was predominantly

localized in the nucleus, with minimal to no fluorescence signal detected in the cytoplasm (Figure 5C,

panel 1 and Supplementary Figure 4, panels 1 and 2). In contrast, the Δ133p53 and Δ160p53

proteins exhibited reduced nuclear staining and enhanced cytoplasmic staining relative to FLp53, with

Δ160p53 showing more pronounced cytoplasmic staining (Figure 5C). Notably, Δ133p53 and

Δ160p53 demonstrated a heterogeneous distribution in the nucleus and cytoplasm (Figure 5C, panels

2 and 3). Furthermore, both Δ133p53 and Δ160p53 exhibited predominantly cytoplasmic punctate

staining, particularly Δ160p53, suggesting the formation of protein aggregates (Figure 5C, panels 2

and 3).

Given that p53 aggregation has been found in numerous tumor cells[60, 61, 62, 63], it is important to

explore the effects of Δ133p53 or Δ160p53 isoforms on the aggregation of FLp53. To investigate the

aggregation property of the p53 isoforms, we performed a subcellular fractionation of the H1299 cells

expressing FLp53 along with Δ133p53 or Δ160p53 at 1:5 ratio. Notably, our fractionation approach

allows us to analyze soluble cytoplasmic and nuclear fractions as well as insoluble nuclear fraction.

Immunoblotting of the soluble fraction showed that an excess of Δ133p53 and Δ160p53 isoforms

reduced the levels of FLp53 in both the cytoplasm and nucleus, with a notable reduction in the

nucleus (Figure 6A and C). Conversely, insoluble nuclear FLp53 increased significantly upon co-

expression with the p53 isoforms, with Δ160p53 showing a more prominent effect (Figure 6B and D,

Supplementary Figure 5).

To exclude the possibility that FLAG or V5 tags contribute to protein aggregation, we also

conducted subcellular fractionation of H1299 cells expressing FLAG-tagged FLp53 along with

untagged Δ133p53 or Δ160p53 at a 1:5 ratio. The results showed (Supplementary Figure 6) a similar

distribution of FLp53 across cytoplasmic, nuclear, and insoluble nuclear fractions as in the case of

tagged Δ133p53 or Δ160p53 (Figure 6A to D). Notably, the aggregation of untagged Δ133p53 or

Δ160p53 markedly promoted the aggregation of FLAG-tagged FLp53 (Supplementary Figure 6B and

D), demonstrating that the antibody epitope tags themselves do not contribute to protein aggregation.

These results collectively demonstrate that Δ133p53 and Δ160p53 isoforms strongly induce FLp53

aggregation, and the decrease in soluble FLp53 levels could be attributed to its aggregation when co-

expressed with isoforms Δ133p53 and Δ160p53.

We complemented the immunoblotting assays with immunofluorescence assays. Our results

clearly show the co-localization of FLp53 with the isoforms Δ133p53 and Δ160p53. More importantly,

we find co-aggregation of FLp53 with the two isoforms forming punctate spots in the cytoplasm and

nucleus (Figure 6E, panels 2 and 3 and Supplementary Figure 4, panels 3 and 4). Furthermore, more

prominent subnuclear foci of FLp53 formed in the cells co-expressing the Δ160p53 protein (Figure 6E,

panel 3 and Supplementary Figure 4, panel 4), which were detected in the Δ133p53 protein

expressing H1299 cells. Thus, these results imply that Δ133p53 and Δ160p53 isoforms efficiently

induce co-aggregation of FLp53, potentially influencing its function in cancer cells.

The Δ133p53 and Δ160p53 proteins block pro-apoptotic function of FLp53.

One of the physiological read-outs of FLp53 is its ability to induce apoptotic cell death[64]. To

investigate the effects of p53 isoforms Δ133p53 and Δ160p53 on FLp53-induced apoptosis, we

measured caspase-3 and -7 activities in H1299 cells expressing different p53 isoforms (Figure 7).

Caspase activation is a key biochemical event in apoptosis, with the activation of effector caspases

(caspase-3 and -7) ultimately leading to apoptosis[65]. The caspase-3 and -7 activities induced by

FLp53 expression was approximately 2.5 times higher than that of the control vector (Figure 7). Co-

expression of FLp53 and the isoforms Δ133p53 or Δ160p53 at a ratio of 1: 5 significantly diminished

the apoptotic activity of FLp53 (Figure 7). This result aligns well with our reporter gene assay, which

demonstrated that elevated expression of Δ133p53 and Δ160p53 impaired the expression of

apoptosis-inducing genes BAX and PUMA (Figure 4G and H). Moreover, a reduction in the apoptotic

activity of FLp53 was observed irrespective of whether Δ133p53 or Δ160p53 protein was expressed

with or without a FLAG tag (Figure 7). This result, therefore, also suggests that the FLAG tag does not

affect the apoptotic activity or other physiological functions of FLp53 and its isoforms. Overall, the

overexpression of p53 isoforms Δ133p53 and Δ160p53 significantly attenuates FLp53-induced

apoptosis, independent of the protein tagging with the FLAG antibody epitope.

Discussion

The inactivation of p53 is one of the most common events in cancer cells[66]. Although functionally

compromised p53 mutations are prevalent in more than half of human cancers[67, 68], specific p53

isoforms have recently been associated with tumorigenesis in various cancer types[11, 14]. The aberrant

expression of Δ133p53 has been found in colorectal cancer[24, 69] and lung cancer[58]. In comparison, the

oncogenic role of the Δ160p53 isoform in human cancers is less explored[11], although links have been

established between the expression of Δ160p53 isoform and melanoma cells[34]. The mechanisms of

the oncogenic potential of p53 isoforms have attracted attention from researchers[14, 38, 39, 70, 71, 72]. A

theory of the dominant-negative effect has been introduced, but its exact details have remained

unknown.

The ChIP and luciferase activity assays show that the isoforms Δ133p53 and Δ160p53 are

defective in their ability to bind DNA and activate transcription (Figure 3B and Figure 4A-D). A possible

reason for the diminished DNA-binding function of these isoforms is that their core domains are

defective (Figure 1B). The Δ133p53 and Δ160p53 isoforms lack approximately 20% and 33% of the

DBD, which hinders their ability to bind to p53REs. We find that co-expression of Δ133p53 and

Δ160p53 with FLp53 exerts an inhibitory effect on FLp53's DNA-binding ability (Figure 3C). However,

the degree to which Δ133p53 and Δ160p53 impair FLp53 function is influenced by both the

expression levels of these isoforms and the specific promoters of the target genes (Figure 4). For the

p21 and MDM2 promoters, equal expression of FLp53 and the two isoforms didn't show much effect.

However, higher expression levels of the isoforms gradually inhibited FLp53's activity to induce

transcription from those promoters (Figure 4A and B). Notably, the PUMA promoter behaved

differently, whereby 1:1 co-expression of FLp53 with Δ133p53 or Δ160p53 significantly increased its

activity (Figure 4D). We speculate that this isoform expression level may activate p21 induction

pathways independent of FLp53[73], potentially facilitating the interaction between FLp53 and p21,

which is required for optimal activation of PUMA promoter activity[74]. These results underscore a

complex regulatory network where FLp53 and the isoform expression levels, target gene specificity,

and transcriptional activities modulate p53-mediated cellular responses.

Our co-expression data clearly indicate that the isoforms Δ133p53 and Δ160p53 exert a dominant-

negative effect on FLp53 function (Figures 3, 4 and 7). Since both of these p53 isoforms are defective

in DNA binding and they possess an intact C-terminal OD, it is not unexpected that they form different

species of homo- and hetero-tetramers with FLp53. In accordance with that, the immunoprecipitation

assays using FLAG-tagged FLp53 and V5-tagged/untagged Δ133p53 or Δ160p53 demonstrate a

potential direct interaction between these isoforms and FLp53 (Figure 2A, lanes 7 and 8 and

Supplementary Figure 3, lanes 3 and 4). However, in what ratio the p53 isoforms impose inhibition to

FLp53 function is worth investigating. We employed a theoretical model of p53 hetero-tetramer activity

in transcriptional activation from p21, MDM2, BAX, and PUMA promoters by varying relative

expression levels of the isoforms[75] (Supplementary Figure 7). We assumed that the transfection level

was a good proxy for the expression level. Interestingly, the experimental inhibition curves with

Δ133p53 or Δ160p53 derived from the luciferase reporter assay were all above the theoretical

inhibition curves for two isoform molecules per tetramer (Supplementary Figure 7B-E), suggesting

that at least three isoform molecules per tetramer are required to abolish the transcriptional activity of

FLp53. This is similar to an earlier report with Δ40p53 isoform, which was shown to exert a dominant-

negative effect at a 3:1 ratio with FLp53[50].

Interestingly, similar to Δ40p53[39], the Δ133p53, and Δ160p53 proteins can aggregate in the

cytoplasm (Figures 5C and 6E). Due to cytoplasmic aggregation and a stronger tendency for

cytoplasmic localization (Figure 5), only a limited amount of these isoforms can successfully be

imported into the nucleus successfully. Additionally, the isoforms tend to aggregate within the nucleus

(Figure 6B and E). Consequently, the ratio of these isoforms to FLp53 diminishes in the nucleus in

contrast to their expression levels. This decrease reduces the likelihood of forming hetero-tetramers,

in which the isoforms will be in higher proportion than FLp53. Thus, it seems unlikely that hetero-

tetramerization of Δ133p53 and Δ160p53 isoforms with FLp53 would be the main reason for their

dominant-negative effect.

Next we investigated whether aggregation of the Δ133p53 and Δ160p53 isoforms contributes to

their functional loss and dominant-negative effect on FLp53. Our immunofluorescence and

immunoprecipitation analyses of H1299 cells expressing these isoforms show a higher aggregation

propensity of the two isoforms than FLp53 (Figure 5C and Supplementary Figure 2). The aggregation

likely results from the exposure of hydrophobic sequences in the isoforms (Figure 1C). In addition, the

absence of N-terminal TADs in Δ133p53 and Δ160p53 isoforms may prevent their MDM2-mediated

ubiquitin-proteasome degradation[66, 76]. This may, in turn, lead to higher accumulation of Δ133p53 and

Δ160p53 isoforms in the cell, which, combined with an unstable core domain, promotes their

aggregation. Moreover, the observed cytoplasmic isoform aggregates may reflect autophagy-related

degradation, as suggested by the co-localization of Δ133p53 with autophagy substrate p62/SQSTM1

and autophagosome component LC3B[77]. Overexpression of these aggregation-prone proteins could

induce endoplasmic reticulum stress and activate autophagy[78]. Interestingly, we also observed

nuclear aggregation of these isoforms (Figure 6B and E and Supplementary Figure 6B), suggesting

that distinct mechanisms, such as intrinsic properties of the isoforms, may govern their localization

and behavior within the nucleus. This dual localization underscores the complexity of Δ133p53 and

Δ160p53 behavior in cellular systems.

More interestingly, our immunofluorescence and immunoprecipitation assays demonstrate large co-

aggregation of FLp53 with Δ133p53 and Δ160p53 isoforms in the cytoplasm as well as in the nucleus

when expressed together (Figure 6E and Supplementary Figure 4, panels 3 and 4). The predominant

cytoplasmic aggregation would sequester FLp53 in the cytoplasm. Moreover, it may destabilize

FLp53's conformation, potentially masking the nuclear localization sequence (NLS) and preventing its

nuclear import[79]. Additionally, it may disrupt the interaction of FLp53 with MDM2, thereby inhibiting its

MDM2-induced degradation and enhancing its accumulation[80], which would drive it more to the

aggregation pathway. All these processes involving FLp53 aggregation would result in reduction of

the functional FLp53 in the cell and more importantly, in the nucleus. Cytoplasmic aggregation would

also limit the isoforms from nuclear import, reducing the possibility of hetero-tetramerization.

Therefore, we propose that the co-aggregation of Δ133p53 and Δ160p53 with FLp53 is the primary

factor for their dominant-negative effect. This mechanism is consistent with the model where structural

mutations, such as R175H and R282W in FLp53, induce co-aggregation and exert a dominant-

negative effect[40].

In our study, we primarily utilized an overexpression strategy involving FLAG/V5-tagged proteins to

investigate the effects of p53 isoforms Δ133p53 and Δ160p53 on the function of FLp53. To address

concerns regarding potential overexpression artifacts, we performed the co-immunoprecipitation

(Supplementary Figure 6) and caspase-3 and -7 activity (Figure 7) experiments with untagged

Δ133p53 and Δ160p53. In both experimental systems, the untagged proteins behaved very similarly

to the FLAG/V5 antibody epitope-containing proteins (Figures 6 and 7 and Supplementary Figure 6).

Hence, the C-terminal tagging of FLp53 or its isoforms does not alter the biochemical and

physiological functions of these proteins.

We propose a model to explain how the p53 isoforms Δ133p53 and Δ160p53 inhibit FLp53 activity

and exert a dominant-negative effect on it (Figure 8A). Two mechanisms are involved: a subsidiary

mechanism in which Δ133p53 and Δ160p53 isoforms form hetero-tetramers with FLp53 impairing its

DNA-binding and transcriptional activities, and a predominant mechanism in which these isoforms

promote co-aggregation with FLp53 and abolish its transcriptional activities by disrupting its structure,

regulation and degradation pathways. Since elevated levels of Δ133p53 and Δ160p53 relative to

FLp53 attenuate its transcriptional activity (Figure 8B, left) and overexpression of FLp53 relative to the

isoforms allows to regain it (Figure 8B, right), the tumor-suppressive efficacy of p53 must depend on

the balanced expression of FLp53 and its isoforms, highlighting the importance of their relative

abundance in cancer cells. Based on our results and recent related publications[38, 39] we propose that

co-aggregation, rather than hetero-tetramerization, is the primary mechanism for the dominant-

negative effect exerted by the p53 isoforms with large N-terminal truncations.

Methods

Cell culture, plasmids, and plasmid transfections. Human non-small cell lung carcinoma cells

H1299 (p53-null) were obtained from ATCC and cultured in Dulbecco's modified Eagle's medium

(DMEM) supplemented with 10% fetal bovine serum (FBS, Thermo Fisher Scientific) and penicillin-

streptomycin solution (PEST; Thermo Fisher Scientific) at 37 °C in a 5% CO_2 incubator. The plasmid

pcDNA3.1-FLp53-FLAG, encoding human full-length p53 (amino acids 1-393), was synthesized by

GenScript (Rijswijk, Netherlands). A flexible linker $(GGGGG)_1$ was introduced between the FLp53

coding sequence and the C-terminal FLAG epitope tag. Subsequently, this fusion construct was

inserted into the pcDNA3.1 vector at the NheI and EcoRI restriction enzyme sites. The plasmids

encoding N-terminally truncated p53 proteins (Δ133p53 and Δ160p53) were created by deleting the

N-terminal 132 and 159 amino acids from the FLp53 sequence, respectively. To create plasmids

expressing C-terminal V5-tagged proteins (FLp53, Δ133p53, and Δ160p53), the respective DNA

sequences were amplified by PCR from the pcDNA3.1-FLp53-FLAG template and inserted into the

EcoRI and XhoI sites of the pcDNA3.1-V5. The plasmids encoding untagged p53 proteins were

constructed by inserting the respective DNA sequences into the NheI and EcoRI restriction sites of

the pcDNA3.1 empty vector. Luciferase reporter plasmids, WWP/p21-Luc (Plasmid #16451)[6], pGL3-

MDM2-Luc (Plasmid #32365)[81], and PUMA Frag1-Luc (Plasmid #16591)[82] were obtained from

Addgene. For the construction of the BAX-Luc reporter plasmid, the p53 binding site in the BAX

promoter was PCR amplified from HepG2 genomic DNA kindly provided by Dr. Gang Pan (Uppsala

University) and cloned into the pGL3-Basic plasmid (Promega) using XhoI and HindIII sites. The PCR

primers used for cloning are listed in Supplementary Table 1. All plasmid transfections were performed

with jetPRIME® (Polyplus) transfection reagent according to the manufacturer's protocol.

Immunoprecipitation and Western blotting. H1299 cells were seeded in a 6-well plate and

transfected with 2 μg of various plasmids. The total amount of plasmids (2 μg/well) was kept constant

in all samples by adding empty plasmid pcDNA3.1. To investigate p53 oligomerization, cells were co-

transfected with equal amounts (1 μg each) of FLAG- and V5-tagged plasmids. After 24 h post-

transfection (hpt), cells were harvested and lysed in RIPA buffer (50 mM Tris-HCl pH 7.4, 150 mM

NaCl, 1% Triton X-100, 1% sodium deoxycholate, 0.1% SDS, and 1 mM EDTA) supplemented with

the Halt Protease Inhibitor Cocktail (Thermo Fischer Scientific) as previously described[83]. The whole

cell lysates were incubated for 30 min on ice and centrifuged at 13,000 × g for 15 min at 4°C. A

fraction of the cleared cell lysates was reserved as the input sample; the remaining lysates were used

for immunoprecipitation with anti-FLAG M2 magnetic beads (Sigma, Cat No. M8823) overnight at 4°C.

The beads were washed with 4 ×1 ml of RIPA buffer. Proteins were then eluted from the beads using

2 × Laemmli Sample buffer, separated on a 10% SDS-PAGE, transferred onto nitrocellulose

membranes (Bio-Rad), and blocked with 5% nonfat dry milk in TBS containing 0.1% Tween-20

(TBST) for 1 h at room temperature. The blocked membranes were incubated overnight at 4°C with

the anti-FLAG (Sigma, Cat No. F2555) and anti-V5 (Sigma, Cat No. V8012) primary antibodies in a

blocking buffer overnight at 4°C. Proteins were detected with the IRDye 680RD Goat anti-Rabbit IgG

and IRDye 800CW Donkey anti-Mouse IgG secondary antibodies (Li-COR Biosciences). The Western

blots were visualized using an Odyssey CLX imaging system (Li-COR Biosciences).

Chromatin immunoprecipitation (ChIP) assays. H1299 cells were seeded in 10 cm cell culture

dishes and transfected with 5 μg of either pcDNA3.1-FLp53-FLAG, pcDNA3.1-Δ133p53-FLAG,

pcDNA-Δ160p53-FLAG or an empty vector pcDNA3.1 for 24 h. To investigate the impacts of isoforms

Δ133p53 and Δ160p53 on the DNA-binding activity of FLp53, H1299 cells were co-transfected with

2.5 μg of pcDNA3.1-FLp53-FLAG along with either 2.5 μg of pcDNA3.1-Δ133p53-V5, pcDNA3.1-

Δ160p53-V5 or an equal amount of empty vector pcDNA3.1. The ChIP assays were performed as

described previously[84], with minor modification. Briefly, cells were fixed with 1% formaldehyde in PBS

for 10 min at room temperature, followed by lysis in 250 μl of lysis buffer (50 mM Tris-HCl pH 8.0, 10

mM EDTA, 1% SDS) supplemented with Halt Protease Inhibitor Cocktail. Lysates were then

sonicated for 15 cycles (30 s ON/30 s OFF) using the Bioruptor Pico (Diagenode). The resulting

soluble cell lysates were diluted with dilution buffer (16.7 mM Tris-HCl pH 8.0, 167 mM NaCl, 1.2 mM

EDTA, 0.01% SDS, 1.1% Triton X-100) and immunoprecipitated overnight at 4°C with the anti-FLAG

M2 magnetic beads. After a series of washes, protein-DNA complexes were eluted, and the crosslinks

were reversed through incubation with RNase A (Thermo Fischer Scientific) at 37°C for 30 min,

followed by proteinase K (Invitrogen) digestion at 65°C overnight. The enrichment of specific DNA

fragments was evaluated with qPCR analysis using HOT FIREPol EvaGreen qPCR Supermix (Solis

Biodyne). All qPCR reactions were done in triplicate, employing the QuantStudio™ 6 Flex Real-Time

PCR System (Thermo Fisher Scientific). The relative fold enrichments were calculated after

normalizing to the input values and considering the pcDNA3.1 transfected samples as 1. Primer

sequences used for qPCR analysis are listed in Supplementary Table 1.

Luciferase assay. H1299 cells were seeded in 24-well plates and transfected in triplicate with

25 ng/well of luciferase reporter plasmids and various p53-expressing plasmids. Co-transfection ratios

for FLp53 and its isoforms Δ133p53 or Δ160p53, were set at 1:1 (12.5 ng/well of FLp53 to 12.5

ng/well of isoform), 1:5 (12.5 ng/well of FLp53 to 62.5 ng/well of isoform), and 1:10 (12.5 ng/well of

FLp53 to125 ng/well of isoform). The total plasmid amount (300 ng/well) was kept consistent across

the various groups by supplementing with the empty pcDNA3.1 plasmid. Cells were harvested 24 hpt

and lysed in a Passive Lysis Buffer (Promega). The cell lysate was centrifuged at 13,000 × g for 5 min

at 4°C, and 20 µl of the lysate was analyzed in triplicates using a firefly luciferase substrate on Infinite

200 PRO plate reader (TECAN)[85].The luciferase activity of the FLp53 and its isoforms transfected

samples was normalized against that of samples transfected with the control vector pcDNA3.1 to

obtain relative reporter activity. Results represent the mean values and standard deviations of three

independent experimental replicates. The data was analyzed using Origin 2018 software (Origin Lab

Corporation, USA).

Subcellular fractionation. The subcellular fractionation assay was carried out using NE-PER

Nuclear and Cytoplasmic Extraction Reagents (Thermo Fischer Scientific), with minor modifications.

In brief, H1299 cells were seeded in 10 cm cell culture dishes and transfected with 0.8 µg of the

following plasmids: pcDNA3.1-FLp53-FLAG, pcDNA3.1-Δ133p53-FLAG, pcDNA-Δ160p53-FLAG, and

the control vector pcDNA3.1. To investigate the effects of overexpressing p53 isoforms on the

subcellular localization of FLp53, co-transfection was performed using FLAG-tagged FLp53 and V5-

tagged Δ133p53 or Δ160p53 at a 1:5 ratio (0.8 µg to 4.2 µg). Cells were transfected with 5 µg of the

empty vector pcDNA3.1 as a control. About 24 hpt, cells were collected by trypsinization,

resuspended in ice-cold CER I buffer containing Halt Protease Inhibitor Cocktail and incubated for 10

min on ice. Subsequently, ice-cold CER II buffer was added according to the manufacturer's

instructions. The cytoplasmic extracts were isolated by centrifugation at 14,000 × g for 5 min at 4°C.

The nuclear pellet was then resolved in ice-cold RIPA buffer supplemented with Halt Protease

Inhibitor Cocktail, subjected to vortex for 15 s every 10 min for a total of 40 min, and centrifuged at

14,000 × g for 10 min to obtain the soluble nuclear fraction. The insoluble nuclear pellet (i.e., insoluble

nuclear fraction) was resuspended in RIPA buffer, an appropriate volume of 2 × Laemmli Sample

buffer was added, and the samples were heated at 95°C until the pellet fully dissolved. For Western

blot experiments, equal volumes of each subcellular fraction were separated by SDS-PAGE, and the

nitrocellulose membranes were probed with the respective antibodies. Histone H3 (nuclear marker)

and α-Tubulin (cytoplasmic marker) antibodies were used to verify the purity of the nuclear and

cytoplasmic fractions. All primary antibodies: anti-FLAG (Sigma, Cat No. F2555), anti-V5 (Sigma, Cat

No. V8012 and Cat No. AB3792), anti-Histone H3 (Abcam, Cat No. ab1791), and anti-α-Tubulin

(Sigma, Cat No. T6199) were incubated on membranes overnight at 4°C. The Western blots were

visualized using an Odyssey CLX imaging system, and the proteins were quantified using Image

Studio Version 5.5 (Li-COR).

Immunofluorescence analysis. H1299 cells were seeded on glass coverslips in 24-well plates and

transfected with 250 ng/well of either pcDNA3.1-FLp53-FLAG, pcDNA3.1-Δ133p53-FLAG, pcDNA-

Δ160p53-FLAG or the control vector pcDNA3.1. To investigate whether an excess of p53 isoforms

could influence the subcellular localization of FLp53, H1299 cells were co-transfected with 250 ng/well

of pcDNA3.1-FLp53-FLAG along with 1000 ng/well of pcDNA3.1-Δ133p53-V5, pcDNA3.1-Δ160p53-

V5, or pcDNA3.1. About 24 hpt, cells were fixed with 4% paraformaldehyde in PBS, followed by

permeabilization with 0.2% Triton X-100 in PBS for 15 min at room temperature as described

previously[86]. Fixed cells on coverslips were incubated overnight at 4°C with primary antibodies anti-

FLAG (Sigma, Cat No 1804, 1:1000 dilution), and anti-V5 (Sigma, Cat No AB3792, 1:1000 dilution).

Subsequently, coverslips were incubated with the Alexa Fluor 488 goat anti-mouse IgG (Invitrogen,

Cat No. A11001, 1:2000 dilution) and/or Alexa Fluor 594 chicken anti-rabbit IgG (H+L) (Invitrogen, Cat

No. A21442, 1:2000 dilution) secondary antibodies for 30 min at room temperature. The coverslips

were mounted with ProLong Diamond Antifade Mountant (Invitrogen, Cat No. P36961), and nuclei

were counterstained using DAPI. Subcellular localization was visualized using a Nikon Eclipse 90i

microscope (Nikon, Japan). Images were processed using ImageJ software[87] and subsequently

arranged by Inkscape.

Caspase-3/7 activity assay. Caspase-3 and -7 activities were measured with a Caspase-Glo® 3/7

assay kit (Promega, Madison, USA) according to the manufacturer's protocol. Briefly, H1299 cells

were seeded (5000 cells/well) into a white solid bottom 96-well plate and transfected in triplicate with

180 ng/well of plasmids expressing FLAG-tagged or untagged FLp53, Δ133p53, Δ160p53, or the

empty vector control pcDNA3.1. To evaluate the effects of overexpression of isoforms on the

apoptotic activity of FLp53, H1299 cells were co-transfected with plasmids encoding FLp53 (30

ng/well) with the control plasmid pcDNA3.1 (150 ng/well), or plasmids expressing isoforms Δ133p53

and Δ160p53 (150 ng/well), at a 1:5 ratio. After 28 hpt, caspase-3/7 substrate was added to each well

and incubated for 90 min at room temperature. Luminescence was recorded using an Infinite 200

PRO plate reader (TECAN). The background luminescence associated with the cell culture and assay

reagent (blank reaction) was subtracted from the experimental values.

Acknowledgments

501 We thank Dr. Zamaneh Hajikhezri for her help with the immunofluorescence experiments and Dr.

502 Gang Pan for reagents. This research was funded by grants from the Swedish Research Council

503 (2023-05237, 2018-05946 and 2018-05498), from Wenner-Gren Foundation (UPD2023-0185) and

504 from the Knut and Alice Wallenberg Foundation (KAW 2017.0055) to S.S.

Author contribution

506 S.S. conceived the idea of the project; L.Z. and T.P. performed experiments; L.Z., T.P., and S.S.

507 analyzed the data; L.Z. wrote the manuscript and prepared Figures. All authors reviewed, edited, and

508 approved the manuscript.

Competing interests

510 The authors declare no competing interests.

References

512 1. Levine AJ, Oren M. The first 30 years of p53: growing ever more complex. *Nature reviews*
513 *Cancer* **9**, 749-758 (2009).
514

515 2. Vogelstein B, Lane D, Levine AJ. Surfing the p53 network. *Nature* **408**, 307-310 (2000).
516

517 3. Pflaum J, Schlosser S, Müller M. p53 Family and Cellular Stress Responses in Cancer.
518 *Frontiers in oncology* **4**, 285 (2014).
519

520 4. Sullivan KD, Galbraith MD, Andrysik Z, Espinosa JM. Mechanisms of transcriptional
521 regulation by p53. *Cell death and differentiation* **25**, 133-143 (2018).
522

523 5. Macleod KF*, et al.* p53-dependent and independent expression of p21 during cell growth,
524 differentiation, and DNA damage. *Genes & development* **9**, 935-944 (1995).
525

526   6.   el-Deiry WS, *et al.* WAF1, a potential mediator of p53 tumor suppression. *Cell* **75**, 817-825
527   (1993).
528

529   7.   Shamloo B, Usluer S. p21 in Cancer Research. *Cancers* **11**, (2019).
530

531   8.   Indeglia A, Murphy ME. Elucidating the chain of command: our current understanding of
532   critical target genes for p53-mediated tumor suppression. *Critical reviews in biochemistry and*
533   *molecular biology*, 1-11 (2024).
534

535   9.   Joruiz SM, Bourdon JC. p53 Isoforms: Key Regulators of the Cell Fate Decision. *Cold Spring*
536   *Harbor perspectives in medicine* **6**, (2016).
537

538   10.   Khoury MP, Bourdon JC. p53 Isoforms: An Intracellular Microprocessor? *Genes & cancer* **2**,
539   453-465 (2011).
540

541   11.   Zhao L, Sanyal S. p53 Isoforms as Cancer Biomarkers and Therapeutic Targets. *Cancers* **14**,
542   3145 (2022).
543

544   12.   Mehta S*, et al.* Adaptive homeostasis and the p53 isoform network. *EMBO reports*, e53085
545   (2021).
546

547   13.   Pal A, Ghosh PK, Das S. The "LINC" between Δ40p53-miRNA Axis in the Regulation of
548   Cellular Homeostasis. *Molecular and cellular biology* **43**, 335-353 (2023).
549

550   14.   Vieler M, Sanyal S. p53 Isoforms and Their Implications in Cancer. *Cancers* **10**, 288 (2018).
551

552   15.   Sharathchandra A, Katoch A, Das S. IRES mediated translational regulation of p53 isoforms.
553   *Wiley interdisciplinary reviews RNA* **5**, 131-139 (2014).
554

555   16.   Khoury MP, Bourdon JC. The isoforms of the p53 protein. *Cold Spring Harbor perspectives in*
556   *biology* **2**, a000927 (2010).
557

558   17.   Jeffrey PD, Gorina S, Pavletich NP. Crystal structure of the tetramerization domain of the p53
559   tumor suppressor at 1.7 angstroms. *Science (New York, NY)* **267**, 1498-1502 (1995).
560

561   18.   Weinberg RL, Veprintsev DB, Fersht AR. Cooperative binding of tetrameric p53 to DNA.
562   *Journal of molecular biology* **341**, 1145-1159 (2004).
563

564   19.   Jänicke RU, Graupner V, Budach W, Essmann F. The do's and don'ts of p53 isoforms.
565   *Biological chemistry* **390**, 951-963 (2009).
566

567   20.   Wang P*, et al.* p53 domains: structure, oligomerization, and transformation. *Molecular and*
568   *cellular biology* **14**, 5182-5191 (1994).
569

570 21. Wang Y, Schwedes JF, Parks D, Mann K, Tegtmeyer P. Interaction of p53 with its consensus
571   DNA-binding site. *Molecular and cellular biology* **15**, 2157-2165 (1995).
572

573 22. Kim H*, et al.* p53 requires an intact C-terminal domain for DNA binding and transactivation.
574   *Journal of molecular biology* **415**, 843-854 (2012).
575

576 23. Bourdon JC*, et al.* p53 isoforms can regulate p53 transcriptional activity. *Genes &*
577   *development* **19**, 2122-2137 (2005).
578

579 24. Fujita K*, et al.* p53 isoforms Delta133p53 and p53beta are endogenous regulators of
580   replicative cellular senescence. *Nature cell biology* **11**, 1135-1142 (2009).
581

582 25. Horikawa I*, et al.* Δ133p53 represses p53-inducible senescence genes and enhances the
583   generation of human induced pluripotent stem cells. *Cell death and differentiation* **24**, 1017-
584   1028 (2017).
585

586 26. Pavletich NP, Chambers KA, Pabo CO. The DNA-binding domain of p53 contains the four
587   conserved regions and the major mutation hot spots. *Genes & development* **7**, 2556-2564
588   (1993).
589

590 27. Hofstetter G*, et al.* Δ133p53 is an independent prognostic marker in p53 mutant advanced
591   serous ovarian cancer. *British journal of cancer* **105**, 1593-1599 (2011).
592

593 28. Bischof K*, et al.* Influence of p53 Isoform Expression on Survival in High-Grade Serous
594   Ovarian Cancers. *Scientific reports* **9**, 5244 (2019).
595

596 29. Gong L*, et al.* p53 isoform Δ113p53/Δ133p53 promotes DNA double-strand break repair to
597   protect cell from death and senescence in response to DNA damage. *Cell research* **25**, 351-
598   369 (2015).
599

600 30. Gong L*, et al.* p53 isoform Δ133p53 promotes efficiency of induced pluripotent stem cells and
601   ensures genomic integrity during reprogramming. *Scientific reports* **6**, 37281 (2016).
602

603 31. Gong L, Pan X, Yuan ZM, Peng J, Chen J. p53 coordinates with Δ133p53 isoform to promote
604   cell survival under low-level oxidative stress. *Journal of molecular cell biology* **8**, 88-90 (2016).
605

606 32. Aoubala M*, et al.* p53 directly transactivates Δ133p53α, regulating cell fate outcome in
607   response to DNA damage. *Cell death and differentiation* **18**, 248-258 (2011).
608

609 33. Candeias MM, Hagiwara M, Matsuda M. Cancer-specific mutations in p53 induce the
610   translation of Δ160p53 promoting tumorigenesis. *EMBO reports* **17**, 1542-1551 (2016).
611

612 34. Tadijan A*, et al.* Altered Expression of Shorter p53 Family Isoforms Can Impact Melanoma
613   Aggressiveness. *Cancers* **13**, (2021).

615　35.　Petronilho EC, *et al.* Oncogenic p53 triggers amyloid aggregation of p63 and p73 liquid
616　　　droplets. *Communications chemistry* **7**, 207 (2024).
617

618　36.　Forget KJ, Tremblay G, Roucou X. p53 Aggregates penetrate cells and induce the co-
619　　　aggregation of intracellular p53. *PloS one* **8**, e69242 (2013).
620

621　37.　Farmer KM, Ghag G, Puangmalai N, Montalbano M, Bhatt N, Kayed R. P53 aggregation,
622　　　interactions with tau, and impaired DNA damage response in Alzheimer's disease. *Acta*
623　　　*neuropathologica communications* **8**, 132 (2020).
624

625　38.　Arsic N, *et al.* Δ133p53β isoform pro-invasive activity is regulated through an aggregation-
626　　　dependent mechanism in cancer cells. *Nature communications* **12**, 5463 (2021).
627

628　39.　Melo Dos Santos N, *et al.* Loss of the p53 transactivation domain results in high amyloid
629　　　aggregation of the Δ40p53 isoform in endometrial carcinoma cells. *The Journal of biological*
630　　　*chemistry* **294**, 9430-9439 (2019).
631

632　40.　Xu J, *et al.* Gain of function of mutant p53 by coaggregation with multiple tumor suppressors.
633　　　*Nature chemical biology* **7**, 285-295 (2011).
634

635　41.　Cho Y, Gorina S, Jeffrey PD, Pavletich NP. Crystal structure of a p53 tumor suppressor-DNA
636　　　complex: understanding tumorigenic mutations. *Science (New York, NY)* **265**, 346-355
637　　　(1994).
638

639　42.　Joerger AC, Allen MD, Fersht AR. Crystal structure of a superstable mutant of human p53
640　　　core domain. Insights into the mechanism of rescuing oncogenic mutations. *The Journal of*
641　　　*biological chemistry* **279**, 1291-1296 (2004).
642

643　43.　Raj N, Attardi LD. The Transactivation Domains of the p53 Protein. *Cold Spring Harbor*
644　　　*perspectives in medicine* **7**, (2017).
645

646　44.　Chen Y, Dey R, Chen L. Crystal structure of the p53 core domain bound to a full consensus
647　　　site as a self-assembled tetramer. *Structure (London, England : 1993)* **18**, 246-256 (2010).
648

649　45.　Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold: making
650　　　protein folding accessible to all. *Nature methods* **19**, 679-682 (2022).
651

652　46.　Cañadillas JM, Tidow H, Freund SM, Rutherford TJ, Ang HC, Fersht AR. Solution structure of
653　　　p53 core domain: structural basis for its instability. *Proceedings of the National Academy of*
654　　　*Sciences of the United States of America* **103**, 2109-2114 (2006).
655

47.　Olotu FA, Soliman MES. From mutational inactivation to aberrant gain-of-function: Unraveling
the structural basis of mutant p53 oncogenic transition. *Journal of cellular biochemistry* **119**,

2646-2652 (2018).

48.   Clore GM, *et al.* Refined solution structure of the oligomerization domain of the tumour
suppressor p53. *Nature structural biology* **2**, 321-333 (1995).

49.   Courtois S, *et al.* DeltaN-p53, a natural isoform of p53 lacking the first transactivation domain,
counteracts growth suppression by wild-type p53. *Oncogene* **21**, 6722-6728 (2002).

50.   Hafsi H, Santos-Silva D, Courtois-Cox S, Hainaut P. Effects of Δ40p53, an isoform of p53
lacking the N-terminus, on transactivation capacity of the tumor suppressor protein p53. *BMC*
cancer **13**, 134 (2013).

51.   Petty TJ, *et al.* An induced fit mechanism regulates p53 DNA binding kinetics to confer
sequence specificity. *The EMBO journal* **30**, 2167-2176 (2011).

52.   Emamzadah S, Tropia L, Vincenti I, Falquet B, Halazonetis TD. Reversal of the DNA-binding-
induced loop L1 conformational switch in an engineered human p53 protein. *Journal of*
molecular biology **426**, 936-944 (2014).

53.   Weinberg RL, Veprintsev DB, Bycroft M, Fersht AR. Comparative binding of p53 to its
promoter and DNA recognition elements. *Journal of molecular biology* **348**, 589-596 (2005).

54.   Purvis JE, Karhohs KW, Mock C, Batchelor E, Loewer A, Lahav G. p53 dynamics control cell
fate. *Science (New York, NY)* **336**, 1440-1444 (2012).

55.   Park SY, Jeong MS, Jang SB. In vitro binding properties of tumor suppressor p53 with PUMA
and NOXA. *Biochemical and biophysical research communications* **420**, 350-356 (2012).

56.   Menendez D, *et al.* Diverse stresses dramatically alter genome-wide p53 binding and
transactivation landscape in human cancer cells. *Nucleic acids research* **41**, 7286-7301
(2013).

57.   Beckerman R, Prives C. Transcriptional regulation by p53. *Cold Spring Harbor perspectives in*
biology **2**, a000935 (2010).

58.   Fragou A, *et al.* Increased Δ133p53 mRNA in lung carcinoma corresponds with reduction of
p21 expression. *Molecular medicine reports* **15**, 1455-1460 (2017).

59.   Stommel JM, Marchenko ND, Jimenez GS, Moll UM, Hope TJ, Wahl GM. A leucine-rich
nuclear export signal in the p53 tetramerization domain: regulation of subcellular localization
and p53 activity by NES masking. *The EMBO journal* **18**, 1660-1672 (1999).

60.   Costa DC, de Oliveira GA, Cino EA, Soares IN, Rangel LP, Silva JL. Aggregation and Prion-
Like Properties of Misfolded Tumor Suppressors: Is Cancer a Prion Disease? *Cold Spring*

Harbor perspectives in biology **8**, (2016).

61.   de Oliveira GA, Rangel LP, Costa DC, Silva JL. Misfolding, Aggregation, and Disordered

Segments in c-Abl and p53 in Human Cancer. *Frontiers in oncology* **5**, 97 (2015).

62.   Levy CB*, et al.* Co-localization of mutant p53 and amyloid-like protein aggregates in breast

tumors. *The international journal of biochemistry & cell biology* **43**, 60-64 (2011).

63.   Li J, Guo M, Chen L, Chen Z, Fu Y, Chen Y. Amyloid aggregates induced by the p53-R280T

mutation lead to loss of p53 function in nasopharyngeal carcinoma. *Cell death & disease* **15**,

35 (2024).

64.   Aubrey BJ, Kelly GL, Janic A, Herold MJ, Strasser A. How does p53 induce apoptosis and

how does this relate to p53-mediated tumour suppression? *Cell death and differentiation* **25**,

104-113 (2018).

65.   Ghorbani N, Yaghubi R, Davoodi J, Pahlavan S. How does caspases regulation play role in

cell decisions? apoptosis and beyond. *Molecular and cellular biochemistry* **479**, 1599-1613

(2024).

66.   Kussie PH*, et al.* Structure of the MDM2 oncoprotein bound to the p53 tumor suppressor

transactivation domain. *Science (New York, NY)* **274**, 948-953 (1996).

67.   Hollstein M, Sidransky D, Vogelstein B, Harris CC. p53 mutations in human cancers. *Science*

(New York, NY) **253**, 49-53 (1991).

68.   Bykov VJN, Eriksson SE, Bianchi J, Wiman KG. Targeting mutant p53 for efficient cancer

therapy. *Nature reviews Cancer* **18**, 89-102 (2018).

69.   Campbell H*, et al.* Δ133p53 isoform promotes tumour invasion and metastasis via interleukin-

6 activation of JAK-STAT and RhoA-ROCK signalling. *Nature communications* **9**, 254 (2018).

70.   Gong L, Chen J. Δ113p53/Δ133p53 converts P53 from a repressor to a promoter of DNA

double-stand break repair. *Molecular & cellular oncology* **3**, e1033587 (2016).

71.   Mondal AM*, et al.* p53 isoforms regulate aging- and tumor-associated replicative senescence

in T lymphocytes. *The Journal of clinical investigation* **123**, 5247-5257 (2013).

72.   Lei J*, et al.* Conformational stability and dynamics of the cancer-associated isoform

Δ133p53β are modulated by p53 peptides and p53-specific DNA. *FASEB journal : official*

publication of the Federation of American Societies for Experimental Biology **33**, 4225-4235

(2019).

73.   Karimian A, Ahmadi Y, Yousefi B. Multiple functions of p21 in cell cycle, apoptosis and

transcriptional regulation after DNA damage. *DNA repair* **42**, 63-71 (2016).

74.   Kim U, Kim KS, Park JK, Um HD. Involvement of the p53/p21 complex in p53-dependent
gene expression. *Biochemical and biophysical research communications* **621**, 151-156
(2022).

75.   Chan WM, Siu WY, Lau A, Poon RY. How many mutant p53 molecules are needed to
inactivate a tetramer? *Molecular and cellular biology* **24**, 3536-3551 (2004).

76.   Chen J, Marechal V, Levine AJ. Mapping of the p53 and mdm-2 interaction domains.
Molecular and cellular biology **13**, 4107-4114 (1993).

77.   Horikawa I*, et al.* Autophagic degradation of the inhibitory p53 isoform Δ133p53α as a
regulatory mechanism for p53-mediated senescence. *Nature communications* **5**, 4706 (2014).

78.   Lee H*, et al.* IRE1 plays an essential role in ER stress-mediated aggregation of mutant
huntingtin via the inhibition of autophagy flux. *Human molecular genetics* **21**, 101-114 (2012).

79.   Marine JC. p53 stabilization: the importance of nuclear import. *Cell death and differentiation*
17, 191-192 (2010).

80.   Moll UM, Petrenko O. The MDM2-p53 interaction. *Molecular cancer research : MCR* **1**, 1001-
1008 (2003).

81.   Chang CJ, Freeman DJ, Wu H. PTEN regulates Mdm2 expression through the P1 promoter.
The Journal of biological chemistry **279**, 29841-29848 (2004).

82.   Yu J, Zhang L, Hwang PM, Kinzler KW, Vogelstein B. PUMA induces the rapid apoptosis of
colorectal cancer cells. *Molecular cell* **7**, 673-682 (2001).

83.   Inturi R, Thaduri S, Punga T. Adenovirus precursor pVII protein stability is regulated by its
propeptide sequence. *PloS one* **8**, e80617 (2013).

84.   Punga T, Bühler M. Long intronic GAA repeats causing Friedreich ataxia impede transcription
elongation. *EMBO molecular medicine* **2**, 120-129 (2010).

85.   Schubert E*, et al.* Complex regulation of mitochondrial signaling by human adenovirus minor
capsid protein VI. *Journal of virology* **98**, e0035624 (2024).

86.   Kases K*, et al.* The RNA-binding protein ZC3H11A interacts with the nuclear poly(A)-binding
protein PABPN1 and alters polyadenylation of viral transcripts. *The Journal of biological*
chemistry **299**, 104959 (2023).

87.   Schindelin J*, et al.* Fiji: an open-source platform for biological-image analysis. *Nature*

methods **9**, 676-682 (2012).

88.   Vehlow C*, et al.* CMView: interactive contact map visualization and analysis. *Bioinformatics*

(Oxford, England) **27**, 1573-1574 (2011).

89.   Meng EC*, et al.* UCSF ChimeraX: Tools for Structure Building and Analysis. *Protein science :*

a publication of the Protein Society, e4792 (2023).

90.   Goddard TD*, et al.* UCSF ChimeraX: Meeting modern challenges in visualization and

analysis. *Protein science : a publication of the Protein Society* **27**, 14-25 (2018).

 Figure 1. The Δ133p53 and Δ160p53 proteins exhibit structural destabilization.

(A) Schematic domain structures of full-length (FL) p53 and its isoforms. TAD, transactivation domain; PRD, proline-rich domain; DBD, DNA-binding domain; NLS, nuclear localization sequence; OD, oligomerization domain; CTD, C-terminal domain. Δ133p53 lacks the entire TAD and approximately 20% of the DBD. Δ160p53 lacks the entire TAD and approximately 33% of the DBD. An aggregation region ILTIITL (residues 251-257) is indicated in the DBD domain. **(B)** Contact density map (C-α 8Å) of the core domain of FLp53 (PDB ID: 3KMD) using CWView[88]. The residues 94-132 and 133-159 were colored magenta and green, respectively. Higher densities are shown in red, and low densities are shown in blue. Among these contacts, four higher-density contacts were selected and shown in oval circles on the map. The corresponding secondary structures of the contacts between the deleted region (blue) and the region (red) in the loop-sheet-helix (LSH) motif or the β-sandwich are shown on the right. **(C)** Comparison analysis of surface hydrophobicity of the core domain in FLp53 (PDB ID: 3KMD) and its isoforms Δ133p53 and Δ160p53 by ChimeraX[89, 90]. The exposed hydrophobic residues in the aggregation region ILTIITL (residues 251-257) are labeled. The color scale of hydrophobicity is indicated with the maximum and minimum values in dark cyan and dark goldenrod, respectively. The β-hairpin S2–S2′ was shown in an oval circle.

Figure 2. Complex formation among FLp53 and its isoforms Δ133p53 and Δ160p53.

(A) Co-immunoprecipitation of FLp53-FLAG, Δ133p53-FLAG, and Δ160p53-FLAG along with the

FLp53-V5, Δ133p53-V5, and Δ160p53-V5 proteins in H1299 cells. The complexes were isolated with

anti-FLAG antibody-coupled magnetic beads. Western blot analysis was done with the anti-FLAG and

anti-V5 antibodies. The merged image corresponds to the overlay between the FLAG and V5 antibody

signals. Input corresponds to 5% of the whole cell lysate. **(B)** A summary of FLp53, Δ133p53, and

Δ160p53 oligomerization based on data shown in panel A.

Figure 3. Δ133p53 and Δ160p53 negatively influence FLp53's DNA-binding activity.

(A) Schematic representation of the promoters of p53 target genes (p21, MDM2, BAX, and PUMA).

An arrow denotes the transcription start site (TSS). The DNA sequences of p53 response elements

(REs) are shown, with uppercase and lowercase letters indicating matched and mismatched bases,

respectively, in relation to the canonical p53 RE sequence (RRRCWWGYYY). Paired arrows highlight

the regions subjected to quantitative PCR (qPCR) amplification. Specifically, the distal (5') p53 RE

within the p21 gene promoter was analyzed. **(B)** Relative DNA-binding of the FLp53-FLAG, Δ133p53-

FLAG, and Δ160p53-FLAG proteins to p53 target genes (p21, MDM2, BAX, and PUMA) in H1299

cells. **(C)** Relative DNA-binding of the FLp53-FLAG protein to the p53-target gene promoters in the

presence of the V5-tagged protein Δ133p53 or Δ160p53 at a 1: 1 ratio. ChIP-qPCR assay data are

shown as relative enrichment of promoter sequences of the target genes after normalization to the

control, pcDNA3.1 plasmid transfected cells. Data are represented as the mean of technical triplicates

± standard deviation (SD), **$P < 0.01$ (Student's t-test).

Figure 4. Impact of Δ133p53 and Δ160p53 on FLp53-activated transactivation.

(A-D) H1299 cells were transfected with luciferase reporter plasmids driven by p21 (A), MDM2 (B),

BAX (C), and PUMA (D) promoters, along with vectors expressing FLp53, Δ133p53, or Δ160p53. To

evaluate the influence of the isoforms on FLp53's transactivation capability, co-expression was

performed at ratios of 1:1, 1:5, and 1:10 relative to FLp53. Basal promoter activity was determined by

transfecting cells with the empty vector, pcDNA3.1. Relative promoter activity is shown after

normalization to the pcDNA3.1-treated sample activity. Data represent mean values ± standard

deviation (SD) (n=3). *$P < 0.05$; **$P < 0.01$ (Student's t-test). **(E-H)** The transcriptional activity of

FLp53 on the p21 (E), MDM2 (F), BAX (G), and PUMA (H) promoters was inhibited by Δ133p53 and

Δ160p53. Inhibition curve fitting was performed using the exponential function ExpDec1 in Origin

2018 software.

Figure 5. Subcellular distribution of the FLp53, Δ133p53 and Δ160p53 proteins.

(A) Western blot analysis of subcellular fractionation of H1299 cells transfected with the FLAG-

tagged FLp53, Δ133p53, or Δ160p53 proteins. Representative immunoblots indicate the presence of

these proteins in different cellular fractions. α-Tubulin (cytoplasmic marker) and Histone H3 (nuclear

marker) were used to ensure fraction integrity. **(B)** Quantitative bar graph summarizing the cytoplasm-

to-nucleus ratio of protein levels. FLAG-tagged protein levels were normalized to their respective

fractionation markers: α-Tubulin for the cytoplasmic fraction and histone H3 for the nuclear fraction.

The relative ratios of Δ133p53 and Δ160p53 protein levels are displayed, with the FLp53 cytoplasm-

to-nucleus ratio set as 1. **(C)** Immunofluorescence analysis of H1299 cells expressing FLAG-tagged

FLp53, Δ133p53 and Δ160p53. Cell nuclei are visualized with DAPI (blue). The punctate spots (white

arrows) indicating protein aggregates are marked. Scale bar, 50 μm.

Figure 6. Induction of FLp53 aggregation by Δ133p53 and Δ160p53 isoforms.

(A-B) Western blot analysis of soluble cytoplasmic and nuclear subcellular fractions **(A)** and insoluble

nuclear fraction **(B)**. Biochemical fractionation of H1299 cells transfected with the FLAG-tagged

FLp53 and V5-tagged Δ133p53 or Δ160p53 at a 1:5 ratio. The aggregated monomers of V5-tagged

Δ133p53 and Δ160p53 (indicated by white arrows), as well as the higher molecular weight

aggregates, are shown. This figure shows one representative experiment; n=3. **(C)** Quantitative bar

graph summarizing the levels of the soluble FLp53-FLAG protein in the cytoplasmic and nuclear

fractions. The FLp53-FLAG protein levels were normalized to the corresponding fractionation marker

(α-Tubulin or histone H3). Relative accumulation of the FLp53-FLAG in Δ133p53-V5 or Δ160p53-V5

expressing cells is shown after considering the FLp53-FLAG protein expressing sample as 1. **(D)**

Quantitative bar graph summarizing the levels of the FLp53-FLAG protein in the insoluble nuclear

fraction relative to the histone H3. Data normalization as in panel C, data represent mean values ±

standard deviation (SD). **$P < 0.01$ (Student's t-test). **(E)** Immunofluorescence analysis of H1299 cells

transfected with FLp53-FLAG alone or in combination with V5-tagged isoforms Δ133p53 and

Δ160p53 expressing plasmids at a concentration ratio of 1:5. Cell nuclei were visualized with DAPI

(blue). The co-aggregation of isoforms with FLp53 in the cytoplasm (white arrows) and nucleus (bright

green arrows) are indicated. Scale bar, 50 μm.

Figure 7. Overexpression of Δ133p53 and Δ160p53 block pro-apoptotic activity of FLp53.
Caspase-3 and -7 activities were detected in H1299 cells co-transfected with plasmid expressing
FLp53 or the isoforms Δ133p53 and Δ160p53 at a 1:5 ratio. Experiments were performed with the
plasmids expressing FLAG-tagged and untagged FLp53, Δ133p53 and Δ160p53. The relative
activities of caspase-3 and -7 were normalized to those of the control vector (transfected only with the
pcDNA3.1 plasmid). Data represent mean values ± standard deviation (SD). **$P < 0.01$ (Student's t-
test).

Figure 8. (A) Schematic representation of the underlying mechanisms by which Δ133p53 and Δ160p53 exert dominant-negative effect on FLp53.

(A) When expressed alone, FLp53 demonstrates strong transcriptional activity, but when expressed together with the Δ133p53 or Δ160p53 isoforms, FLp53 activity gets impaired due to the dominant-negative effect exerted by the isoforms. The Δ133p53 or Δ160p53 isoforms either attenuate FLp53's transcription activity by hetero-tetramerization in a ratio higher than 1:1, or abolish it completely by aggregation. Since Δ133p53 and Δ160p53 exhibit much higher propensity for aggregation than FLp53, they sequester FLp53 in the cytoplasmic aggregates, leading to reduction in the amount of functional FLp53 in the nucleus. Aggregation also limits the availability of these isoforms in the nucleus. Thus, co-aggregation is the primary reason for dominant negative effect of the Δ133p53 or Δ160p53 isoforms on FLp53. It is important to note that the numbers of tetramers and aggregates depicted do not represent the actual quantities present in the cytoplasm and nucleus. **(B)** The balance between the expression level of the FLp53 and its isoforms Δ133p53 and Δ160p53 regulates FLp53 activity. Increased levels of Δ133p53 and Δ160p53 relative to FLp53, lead to stronger dominant-negative effect due to co-aggregation and hetero-tetramerization, resulting in the inhibition of apoptosis. Increased levels of FLp53 compared to the isoforms Δ133p53 and Δ160p53 in cancer cells restore the function of FLp53, leading to the induction of apoptosis.