V-ATPase Disassembly at the Yeast Lysosome-Like Vacuole Is a Phenotypic Driver of Lysosome Dysfunction in Replicative Aging

Fiza Hashmi and Patricia M. Kane

Department of Biochemistry and Molecular Biology
SUNY Upstate Medical University, Syracuse, NY USA

Address correspondence to: Patricia M. Kane
                           kanepm@upstate.edu

Running title: V-ATPase disassembly in replicative aging

Declines in lysosomal acidification and function with aging are observed in organisms

ranging from yeast to humans. V-ATPases play a central role in organelle acidification and V-

ATPase activity is regulated by reversible disassembly in many different settings. Using the yeast

Saccharomyces cerevisiae as a replicative aging model, we demonstrate that V-ATPases

disassemble into their V_1 and V_0 subcomplexes in aging cells, with release of V_1 subunit C

(Vma5) from the lysosome-like vacuole into the cytosol. Disassembly is observed after ≥ 5 cell

divisions and results in overall vacuole alkalinization. Caloric restriction, an established

mechanism for reversing many age-related outcomes, prevents V-ATPase disassembly in older

cells and preserves vacuolar pH homeostasis. Reversible disassembly is controlled in part by the

activity of two opposing and conserved factors, the RAVE complex and Oxr1. The RAVE

complex promotes V-ATPase assembly and a *rav1Δ* mutant shortens replicative lifespan; Oxr1

promotes disassembly and an *oxr1Δ* mutation extends lifespan. Importantly, the level of Rav2, a

key subunit of the RAVE complex, declines in aged cells. These data indicate that reduced V-

ATPase assembly contributes to the loss of lysosome acidification with age, which affects

replicative lifespan.

20 Keywords: Aging, lysosomes, proton pumps, Saccharomyces cerevisiae, caloric restriction

INTRODUCTION

It is well-documented that lysosomal acidification is compromised with age across many organisms (Nixon, 2020). Reduced lysosomal acidification has multiple downstream functional consequences. Lysosomal hydrolases represent a major arm of the cellular proteostatic machinery and operate optimally at the acidic pH, so cargo degradation is compromised at higher pH (Vilchez, Saez, & Dillin, 2014). Lysosomes are the terminal compartment for multiple autophagy pathways, so clearance of autophagic cargoes and recycling of nutrients, both critical in aging cells, is slowed (Hansen, Rubinsztein, & Walker, 2018; Kaushik et al., 2021). Iron and other heavy metals are sequestered and buffered in the acidic lysosomes; loss of sequestration can induce both oxidative stress (Diab & Kane, 2013; Kurz, Terman, Gustafsson, & Brunk, 2008) and deficiency in mitochondrial iron-sulfur proteins (Chen et al., 2020). Reduced lysosomal storage can also create toxic imbalances in amino acids such as cysteine that contribute to loss of mitochondrial function (C. E. Hughes et al., 2020). Recent work has highlighted the central role of the lysosome in nutritional signaling and many aspects of this signaling are linked to acidification (Perera & Zoncu, 2016). It is clear that loss of lysosomal acidification can impact many processes associated with age-related functional decline, but the mechanisms behind increases in lysosomal pH are not fully understood.

The highly conserved vacuolar H^+- ATPase (V-ATPase) acidifies the lumen of lysosomes and lysosome-like vacuoles, as well as endosomes and the late Golgi apparatus, in all eukaryotes (Collins & Forgac, 2020). V-ATPases are multi-subunit protein complexes that couple ATP hydrolysis to proton pumping into organelle lumens. The V-ATPase consists of two subcomplexes: a peripheral V_1 subcomplex oriented toward the cytosol that is responsible for ATP hydrolysis connected to a membrane-embedded V_0 subcomplex containing the proton pore.

V-ATPase subunit sequences are conserved across eukaryotes and recent V-ATPase structures

indicate very strong structural similarity between yeast and mammalian V-ATPases (Oot &

Wilkens, 2020).

V-ATPase activity is highly regulated and responsive to multiple environmental

conditions. Reversible disassembly is a versatile mechanism of V-ATPase regulation that fine-

tunes the activity of the proton pump to meet cellular demands (Collins & Forgac, 2020;

Jaskolka, Winkley, & Kane, 2021; Wilkens, Khan, Knight, & Oot, 2023). In reversible

disassembly, the V_1 subcomplex is released from the V_0 subcomplex inhibiting both ATP

hydrolysis and proton pumping (Kane, 1995; Sumner et al., 1995). V_1 subunit C is dissociated

from both subcomplexes and also becomes cytosolic during disassembly (Kane, 1995).

Disassembly is post-translational and rapidly reversible (Kane, 1995). It was first observed in the

yeast *S. cerevisiae* and the tobacco hornworm *M. sexta* upon acute glucose deprivation and was

reversed by glucose replenishment (Kane, 1995; Sumner et al., 1995). Since that time, it has

become clear that reversible disassembly occurs in many different settings and in response to

diverse signals. For example, unlike yeast cells, most mammalian cells appear to promote V-

ATPase reassembly under conditions of nutrient deprivation and mTOR inhibition, possibly as a

means of promoting lysosomal proteolysis and nutrient recycling (Ratto et al., 2022; Stransky &

Forgac, 2015). In neurons, V-ATPases are reversibly disassembled as part of each synaptic

vesicle cycle (Bodzeta, Kahms, & Klingauf, 2017). In cardiomyocytes, lipid overload can

promote V-ATPase disassembly in endosomes, ultimately contributing to the long-term insulin

resistance (Liu et al., 2017). Reversible disassembly of V-ATPases can also be manipulated by

both host cells and pathogens to prevent or facilitate infection (Kohio & Adamson, 2013).

Several mechanisms for loss of lysosomal acidification with age have been proposed. In

yeast, the long-lived plasma membrane proton pump, Pma1, accumulates in mother cells, and it

has been proposed that increased proton export through Pma1 disrupts the balance of Pma1 and

V-ATPase activities and compromises organelle acidification (Henderson, Hughes, &

Gottschling, 2014). Early loss of vacuolar acidification in yeast has been correlated with

defective mitochondrial morphology and function. Both the deacidification and mitochondrial

morphology phenotypes are suppressed by overexpression of the V-ATPase catalytic subunit,

VMA1, or an ER-localized assembly factor, *VPH2*, suggesting a possible deficiency in these

factors with age (A. L. Hughes & Gottschling, 2012). In *C. elegans*, stability of the *VMA1*

transcript is controlled by a microRNA, miR-1, which can globally control lysosomal

acidification (Schiffer et al., 2021). In several systems, mRNA expression for one or more V-

ATPase subunit genes has been reported to decrease with age (Ghavidel et al., 2018), but in most

cases protein levels have not been assessed. All of these mechanisms could contribute to loss of

vacuolar/lysosomal acidification. Wilms et al. (Wilms et al., 2017) demonstrated that the mTOR

effector Sch9 promotes V-ATPase assembly and extends yeast chronological lifespan.

However, despite the importance of reversible disassembly in regulation of V-ATPase

activity, V-ATPase assembly state during aging has not been explored extensively. Here we show

that V-ATPase assembly does change with age in a yeast replicative aging model. Increased V-

ATPase disassembly in older cells is accompanied by decreased vacuolar acidification that does

not appear to stem from reduced V-ATPase subunit levels. Instead, we provide evidence that

reduced activity of the RAVE (Regulator of H^+-ATPase of Vacuolar and Endosomal membranes)

assembly complex may give rise to net V-ATPase disassembly and increased lysosomal pH with

age.

RESULTS

V-ATPases are more disassembled and vacuoles more alkaline after ~5 cell divisions.

Given the evidence that vacuoles and lysosomes are less acidic in older cells, we hypothesized that V-ATPase assembly and activity might also be changing with age. Replicative aging in the yeast *S. cerevisiae* is a widely accepted model for aging (He, Zhou, & Kennedy, 2018). Briefly, yeast cells divide asymmetrically with each cell division giving rise to a new, "rejuvenated" daughter cell from an established mother cell. Each cell division leaves a bud scar on the mother, allowing visual assessment of age. The yeast V_1C subunit (Vma5) is released from both V_1 and V_0 during V-ATPase disassembly, so we first visualized Vma5-GFP localization in a mixed age population of cells (Figure 1). (Note that vacuoles are visible as indentations under DIC (differential interference contrast) optics on the left of the images.) In parallel, we monitored replicative age of each cell by staining with calcofluor white, which labels the bud scars on mother cells. As shown in Figure 1a, Vma5-GFP is tightly localized to the vacuolar membrane in cells with few or no bud scars. In contrast, in older cells with more bud scars, Vma5-GFP exhibited a marked decrease in fluorescence at the vacuolar membrane and a notable increase in cytosolic fluorescence. In order to compare Vma5-GFP localization between cells, we quantitated maximum fluorescence from a line scan across each cell. As shown in Figure 1b, the line scan from a young cell has prominent peaks corresponding to the edges of the vacuole with a high maximum fluorescence signal, while the older cells have less prominent peaks. We conducted the same analysis across populations of cells of mixed age, normalized to the maximum fluorescence signal of young cells and binned the results by the number of bud scars. As shown at the left of Figure 1c, young cells, defined as having less than 5 bud scars, displayed Vma5-GFP localization at the vacuolar membrane. However, the normalized

maximum fluorescence signal, representative of vacuole localization, decreases significantly in

cells with five bud scars or more. This early onset of V-ATPase disassembly aligns with previous

reports indicating changes in lysosomal pH early in replicative aging (A. L. Hughes &

Gottschling, 2012).

Vma5 is a V_1 subunit that bridges the V_1 and V_0 subcomplexes of the V-ATPase. The V_1

subcomplex also contains three copies of Vma2, and the V_0 subcomplex contains the largest

subunit, Vph1, which comprises part of the proton pore (Figure 2a). Although Vma5 appears to

be the subunit that is released most completely from the membrane by reversible disassembly

(Tabke et al., 2014), the rest of the V_1 sector also dissociates from the vacuolar membrane. We

assessed the cellular distribution of Vma2-GFP, a V_1 subunit, and Vph1-GFP, a V_0 subunit

(Figure 2b). As shown in Figure 2b, there was less membrane-bound Vma2-GFP in older cells

relative to younger cells as assessed by line scans as in Figure 1. However, vacuolar Vph1-GFP

levels were the same between old and young cells (Figure 2c). These results are consistent with

disassembly of the V-ATPase as cells age.

Although comparable levels of Vph1 at the vacuole suggests that expression of V_0

subunits and V_0 assembly are intact, V_1 subunits could become cytosolic because of reduced V_1

subunit levels in older cells. To address this question, we isolated populations old and young

yeast cells by biotinylating the cell walls in a mixed age population, allowing growth to continue

for several generations, and then obtaining "old" cells by biotin-streptavidin affinity

chromatography (Jin, Cao, & Liu, 2021). Daughter cells that emerged after biotinylation cannot

bind to magnetic streptavidin beads and represent the "young" population. The age distribution

was determined by counting bud scars in each population and binning by the number of buds per

cell (Figure 2d). When prepared by this method, the population of old cells peaks at 20-24 bud

scars, while there was a median value of 0-4 bud scars in the young population. Cell lysates were

prepared from each population and analyzed by SDS-PAGE and immunoblotting. As shown in

Figures 2e and f, there is no significant difference in the cellular levels of V_1 subunits Vma1,

Vma2, and Vma5 between young and old populations. These results indicate that the cytosolic

populations of V_1 subunits arise from disassembly of the V-ATPase, rather than inability to

assemble because of lack of V-ATPase subunits.

Reversible disassembly of V-ATPases is employed in a number of contexts to provide

dynamic regulation of the complex in response to changing cellular conditions. Disassembled V_1

and V_0 subcomplexes lack ATPase and proton transport activity and ATP-driven proton pumping

is restored upon reassembly. In order to test whether the lower levels of assembled V-ATPases in

old cells result in reduced capacity for acidification, we measured the response of young and old

cells to an acute glucose deprivation (an abrupt shift to 0% glucose), which promotes

disassembly of the yeast V-ATPase, followed by readdition of glucose to a 2% final

concentration, which promotes reassembly and reactivation of the complex (Kane, 1995).

Young and old yeast cells obtained as described above were loaded with the ratiometric

pH sensor BCECF-AM (2',7'-Bis-(2-carboxyethyl)-5-(and-6)-carboxyfluorescein acetoxy methyl

ester) which localizes to the vacuole in yeast cells (Diakov, Tarsio, & Kane, 2013). Both young

and old cells were shifted to medium with no glucose for ~30 minutes. Fluorescence of cell

suspensions was then monitored continuously (Figure 3a), and glucose was added at the

indicated time. This assay revealed clear age-dependent differences in the vacuolar pH response

to glucose stimulation. Young cells exhibit a rapid drop in vacuolar pH upon glucose addition.

This drop was previously shown to be V-ATPase-dependent and to correlate with V-ATPase

assembly (Martinez-Munoz & Kane, 2008). In old cells, however, there was a smaller pH

decrease after glucose addition, indicating a more alkaline vacuole and consistent with the lower

levels of V-ATPase assembly observed by microscopy during growth in glucose-replete

conditions in Figures 1 and 2. Quantitative analysis of pH at defined time points across

biological replicates (Figure 3b) indicates that vacuoles in old cells are significantly more

alkaline than their younger counterparts at each of the indicated time points. These results

suggest an age-related alteration in vacuolar pH regulation through the inability of V-ATPase to

reassemble. As a result, vacuoles in old cells display a more alkaline pH than vacuoles in young

cells.

Caloric restriction restores V-ATPase assembly and vacuolar acidification in older cells.

Caloric restriction (CR) is defined as a reduction in caloric intake in the presence of

adequate nutrition (Longo & Anderson, 2022). CR promotes both cellular health and longevity.

Extensive research, conducted in diverse model organisms ranging from yeast to worms, flies,

and rodents, suggests that CR can have significant anti-aging effects and promote overall health

(Longo & Anderson, 2022). In the context of the yeast *S. cerevisiae*, adjusting the concentration

of glucose in the growth medium from the 2% used above to 0.5% is a common way to induce

CR. This treatment does not significantly reduce growth rate over several cell divisions

(Supporting information, Figure 1).

As shown in Figure 4, CR reverses the V-ATPase disassembly in older cells. Notably, the

V_1 subunits Vma5-GFP (Figure 4a) and Vma2-GFP (Figure 4b) are recruited to the vacuolar

membrane in both young and old cells, and there is no significant difference in fluorescence at

the vacuolar membrane with age. (Fluorescence intensities of Vph1-GFP continue to be similar

between old and young cells (Figure 4c).) This reversal suggests that CR extends V-ATPase

assembly beyond the replicative age of ≥5 bud scars when cells grown in higher glucose begin to

show disassembly.

We hypothesized that given the improvement in V-ATPase assembly, vacuolar pH in old

cells might also be restored. We grew cells under CR conditions, loaded the vacuoles with

BCECF-AM and monitored vacuolar pH before and after addition of glucose as described above.

Figure 4d demonstrates that the glucose-stimulated decrease in vacuolar pH, which was

compromised in older cells grown in 2% glucose (Figure 3), was restored to the level of young

cells in cells after growth under CR conditions. This observation indicates that CR has a direct

impact on both V-ATPase assembly and vacuolar acidification in aging cells. In addition, it

further highlights the potential connection between V-ATPase assembly, vacuolar acidification

and aging.

Regulators of V-ATPase assembly state affect replicative lifespan.

If reversible disassembly of the V-ATPase plays a central role in aging of yeast cells, we

hypothesized that the cellular factors that regulate V-ATPase assembly might also affect

replicative lifespan. The RAVE complex plays a crucial role in regulation of the V-ATPase by

reversible disassembly (Jaskolka, Winkley, et al., 2021; Seol, Shevchenko, & Deshaies, 2001). It

consists of three subunits: Rav1, Rav2, and Skp1. The RAVE complex is required for reassembly

of V-ATPase complexes disassembled by glucose deprivation. In mutants lacking Rav1 or Rav2,

V-ATPases are predominantly disassembled into V_1 and V_0 subcomplexes and vacuolar

acidification is lost (Seol et al., 2001; Smardon, Tarsio, & Kane, 2002). The RAVE complex

associates with V_1 subcomplexes in the cytosol. Although V_1 subunit C is also released from the

vacuolar membrane, very little is associated with cytosolic RAVE-V_1 (Jaskolka, Tarsio,

Smardon, Khan, & Kane, 2021). In contrast, recent studies suggest that Oxr1, a protein originally

associated with resistance to oxidative stress, promotes disassembly of V-ATPases (Khan et al.,

2022; Khan & Wilkens, 2024; Klossel et al., 2024) suggesting that RAVE and Oxr1 have

opposing effects on the assembly state of V-ATPases as diagrammed in Figure 5a.

To further explore the functional significance of these assembly regulators in aging cells,

we examined the replicative lifespans of deletion mutants lacking Rav1 or Oxr1 (Figure 5b).

Replicative lifespan was measured on YEPD, pH 5 plates, conditions that are optimal for growth

of *rav1Δ* strains. Deletion of Rav1 shortened replicative lifespan (median 17 cell divisions, n=

30) by 26% relative to wild-type cells (median 23 cell divisions, n= 30). In contrast, the deletion

of Oxr1 extended replicative lifespan by 47.8% over wild-type (median 34 cell divisions, n=30).

These results reinforce the significance of V-ATPase assembly in replicative aging and suggest

that RAVE pro-assembly activity (disrupted in the *rav1Δ* mutant) and Oxr1 anti-assembly

activity (lost in *oxr1Δ*) may be central determinants of lifespan.

Given these results, we asked whether there were differences in levels of RAVE subunits

or Oxr1 between young and old cells. We isolated young and old populations of cells containing

myc13-tagged Rav1 or Rav2 by biotinylation and streptavidin magnetic separation as described

above (Figure 2), then assessed the levels of the tagged proteins in cell lysates. Although there is

no significant difference in Rav1-myc13 levels between young and old cells (Figure 6a), we

consistently observed a significant decrease in protein levels of Rav2-myc13 in older cells

(Figure 6b). We also assessed expression of *RAV2* in young and old cells by quantitative PCR but

observed no significant difference in mRNA levels (Figure 6c). Because Rav2 is required for

RAVE complex function in promoting V-ATPase assembly (Seol et al., 2001; Smardon et al.,

2002), these results suggest that partial loss of RAVE function in old cells could contribute to

reduced V-ATPase assembly and replicative aging.

To further explore the basis of the extended replicative lifespan in *oxr1Δ* cells, we

observed Vma5-GFP localization in a population of *oxr1Δ* cells of mixed age. As shown in

Figure 6d, *oxr1Δ* mutants localize Vma5-GFP to the vacuole even in older cells with ≥5 bud

scars. In order to quantitate this effect, we again binned cells by the number of bud scars as

described above, and compared Vma5-GFP localization to localization in young daughter cells

(Figure 6e). In contrast to wild-type cells, there is no significant difference in Vma5-GFP

localization in the *oxr1Δ* mutant until cells have divided 25 or more times. We also isolated

young and old populations from an HA-tagged Oxr1 strain. As shown in Figure 6f, the levels of

Oxr1 do not change with age. However, loss of RAVE function could ultimately favor Oxr1-

induced disassembly.

Intriguingly, these results suggest that the *oxr1Δ* mutation mirrors the effects of CR on V-

ATPase assembly and longevity. The results also indicate that direct manipulation of V-ATPase

assembly by mutation of critical assembly factors can affect replicative lifespan. Specifically,

promoting V-ATPase assembly through the RAVE complex appears to be important for

preserving replicative lifespan, while the opposing effects of Oxr1 on the V-ATPase tend to

shorten lifespan. We hypothesized that restoration of V-ATPase assembly in older cells grown

under CR conditions (Figure 4) might be supported by restoration of Rav2 levels. To test this, we

isolated young and old populations from Rav2-myc13 tagged cells grown under CR conditions.

Under these conditions, V-ATPase assembly and function restored in older cells (Figure 4) and as

shown in Figure 6g, Rav2 levels are also restored.

DISCUSSION

The results described here establish V-ATPase disassembly as an important factor in the reduced vacuolar acidification observed in aging yeast cells. We demonstrate an increase in V-ATPase disassembly at relatively early replicative ages (5-9 cell divisions) similar to the age at which Hughes and Gottschling first observed compromised vacuolar acidification (A. L. Hughes & Gottschling, 2012). We show that promoting assembly of the V-ATPase through CR can restore V-ATPase assembly and vacuolar pH in older cells and that deletion of Oxr1, a negative regulator of V-ATPase assembly, can extend replicative lifespan. Taken together, these data support V-ATPase disassembly as a significant age-related factor behind reduced vacuolar and lysosomal acidification and the associated declines in function. This mechanism does not necessarily conflict with those proposed previously. If V-ATPases are more disassembled in older cells, cells may be even less able to tolerate an imbalance between Pma1 activity at the plasma membrane and V-ATPase activity at the vacuole (Henderson et al., 2014). We see no difference in protein levels of the core V_1 subunits Vma1 and Vma2, or Vma5, between young and old cells. The presence of similar levels of Vph1 in vacuoles of young and old cells suggests that V_0 subunits are also expressed at similar levels, since V_0 assembly occurs in the ER and reduced levels of any V_0 subunit reduces V_0 subcomplex levels at the vacuole (Kane, Kuehn, Howald-Stevenson, & Stevens, 1992). The data presented here suggest a post-transcriptional regulatory mechanism, but increased levels of a V_0 assembly factor like Vph2 (A. L. Hughes & Gottschling, 2012) might still promote assembly and improve acidification.

Many questions remain to be investigated in the future. The reduction in Rav2 protein levels with aging appears to be post-transcriptional but we do not yet know whether this arises from reduced translation or increased degradation of the protein. Perhaps more importantly, even

though reduced levels of Rav2 levels could help explain increased V-ATPase disassembly, we do

not know whether the Rav2 reduction occurs as a result of some general vulnerability in aging

cells or is deliberately programmed (Gladyshev et al., 2021). The significant increase in lifespan

and prolonged V-ATPase assembly in *oxr1Δ* cells seems to argue against any growth advantage

from increased V-ATPase disassembly in older cells, at least at the single cell level.

Many aging pathways are tightly linked to nutritional signaling, and reversible

disassembly is often driven by nutritional signaling pathways. Given the rapid disassembly of the

V-ATPase upon acute glucose deprivation (Kane, 1995), it was initially surprising that growth

under CR conditions suppresses V-ATPase disassembly with age. However, we previously

showed that the acute disassembly response required glucose concentrations well below 0.5%

(Parra & Kane, 1998). In mammalian cells, V-ATPase assembly generally increases in response

to nutrient deprivation (Ratto et al., 2022; Stransky & Forgac, 2015). CR could mimic this effect

in aging cells, preserving lysosomal acidification and function. Signals involved in reversible

disassembly of the V-ATPase are incompletely understood. However, it is intriguing that in yeast,

RAVE appears to play a central role in glucose signaling during acute glucose deprivation.

RAVE is released from the vacuolar membrane upon acute glucose deprivation and recruited

back to the membrane upon glucose restoration, even in the absence of V_1 subunit C and the V_1

subcomplex (Jaskolka, Winkley, et al., 2021). The RAVE complex appears to be a major

determinant of V-ATPase assembly state in multiple situations, including aging.

Reduced V-ATPase assembly could certainly be a factor in the age-related decline in

lysosomal acidification and function in higher eukaryotes. Reversible disassembly actively

occurs in higher eukaryotes including mammalian cells, so the apparatus for age-induced

assembly regulation is available. Rabconnectin-3 complexes of higher eukaryotes are the

functional homologues of the yeast RAVE complex (Ratto et al., 2022; Yan, Denef, &

Schupbach, 2009). Oxr1 belongs to a family of TLDc proteins that are also found in mammals,

and several of these proteins have been shown to bind to mammalian V-ATPases (Eaton, Brown,

& Merkulova, 2021; Wilkens et al., 2023). These data suggest that the core elements for

controlled V-ATPase disassembly during aging are present in other cells. Here, we observed V-

ATPase disassembly in a yeast replicative aging model, which is most comparable to mammalian

cell types that continue to divide, such as adult stem cells (He et al., 2018). However, V-ATPase

activity is also critical in the yeast chronological aging model, which is more analogous to long-

lived, non-dividing mammalian cells. In this model increased V-ATPase assembly has been

associated with longevity (Wilms et al., 2017). Taken together, these data suggest that V-ATPase

assembly state is linked to multiple aging models and could easily play a role in aging in higher

eukaryotes.

METHODS

Yeast strains and plasmids

All strains analyzed were in the BY4741 (*MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*) or BY4742

(*MATα his3Δ1 leu2Δ0 lys1Δ ura3Δ0*) background. BY4742 strains containing Vma5-GFP::*HIS3*

and Vph1-GFP::*HIS3* were constructed as part of the genome-wide GFP-tagging project (Huh et

al., 2003) and purchased from Thermo Fisher. *VMA2* was C-terminally tagged with GFP by PCR

amplification from pFA6a-GFP-KanMX6 and genomic integration (Longtine et al., 1998). The

strain containing *oxr1Δ::kanMX* and Vma5-GFP::*HIS3* was obtained by crossing BY4741

oxr1Δ::kanMX from the haploid deletion collection and BY4742 Vma5-GFP::*HIS3*, sporulating

the diploid, and obtaining spores with the desired genotype by tetrad dissection. The pRS316

Oxr1-HA plasmid (Khan & Wilkens, 2024) was transformed into *oxr1Δ* and transformants were selected for on SC medium (fully supplemented minimal medium) lacking uracil. Rav1-myc13::*kanMX* and Rav2-myc13::*kanMX* strains were described previously (Smardon et al., 2002). The tagged alleles were PCR amplified from the original strains and integrated into BY4741.

Fluorescence microscopy

Strains expressing GFP-tagged Vma5, Vma2, or Vph1 were grown to log phase in SC medium containing either 2% or 0.5% glucose overnight, pelleted by centrifugation, then suspended in fresh medium containing 2% or 0.5% glucose and grown for an additional 2 hours. Cells were stained with calcofluor white (CW) by diluting a 1 mg/ml CW stock to a final concentration of 10 µg/ml with cells, 5 min prior to imaging. Cells were visualized with a 100x oil (NA 1.4) objective on a Zeiss Imager.Z1 fluorescence microscope with a Hamamastu CCD camera and AxioVision software. Cells were viewed through differential interference contrast (DIC) optics or fluorescence was visualized using a DAPI filter set for CW and a GFP filter set for GFP-tagged subunits. Bud scars were counted from CW staining; in mixed age populations, cells with 5 or more bud scars were designated as old and those with less than 5 as young. To obtain a more precise count of bud scars for binning, cells were visualized on multiple focal planes. GFP fluorescence was then determined for each of the binned ranges. Images of GFP-tagged proteins were captured then processed in FIJI. To assess vacuolar localization, a line was drawn across a cell and through the vacuolar membrane. From this line scan, a plot illustrating fluorescence intensity along the line was generated, with peaks indicating areas of elevated fluorescence at the vacuolar membrane. By quantifying the peak intensity (maximum fluorescence), we quantitated the vacuolar localization for each GFP-tagged subunit. Each

biological replicate corresponds to a distinct culture of yeast cells. Maximum fluorescence was

quantitated for at least 20 young and 20 old cells per biological replicate. When cells were

binned by age, at least 20 cells per biological replicate were counted for each bin. To normalize

maximum fluorescence across replicates, the maximum fluorescence from line scans of young

and old cells in each biological replicate was averaged and then divided by the average of the

young cells in that replicate. The normalized fluorescence for each biological replicate was

plotted, along with average intensities across replicates +/- s.e.m. (standard error of the mean).

Statistical significance was determined by t-test for the young and old cell comparisons and by

ANOVA for the binned samples. In order to show the range of values for the young cells across

experiments, the values for the young cells in each biological replicate were averaged and the

values for the individual replicates were divided by the average and shown as points on each

graph.

Age Enrichment

Age enrichment was performed as described by Jin et al. (Jin et al., 2021). Cells were collected

from a 50 mL fresh overnight culture in YEP (yeast extract-peptone medium) supplemented with

2% or 0.5% glucose to an OD_{600} of 1.0 and washed twice with cold phosphate-buffered saline

(PBS), pH 7.4. Cells were pelleted by centrifugation and washed three times in cold sterile PBS,

then labeled with 1.6 mg/ml EZ-Link Sulfo-NHS-LC-Biotin (Pierce) at room temperature for 30

min with gentle agitation. After labeling, the cells were washed three times with cold PBS, pH

8.0, to remove free biotin, and resuspended in YEP supplemented with 2% or 0.5% glucose for

growth overnight. After 16 hours, cells were pelleted by centrifugation and resuspended in 35 ml

of cold PBS, pH 7.4, mixed with 250 µl of a magnetic streptavidin bead suspension (Pierce), and

incubated for 60 min at 4°C. The mixture, in a 50 mL conical tube, was loaded onto a magnetic

separation column (Permagen) at 4°C to separate biotinylated cells and allow unbound cells to

settle, and supernatant was removed gently by pipetting. Magnetically separated cells were

subsequently washed three times by resuspending in 35 mL PBS, pH 7.4 supplemented with 2%

or 0.5% glucose, repeating magnetic separation, and discarding the supernatant. After washing,

cells were resuspended in 200 mL of YEP or SC supplemented with 2% or 0.5% glucose and

allowed to grow for an additional 4 hours before obtaining the final "old" mother cells and

"young" daughter cells. Cells were loaded on the magnetic separation column as described above

and daughter cells obtained from the supernatant were pelleted by centrifugation to obtain a

concentrated population of young cells. After the final wash, the magnetic beads were

resuspended in 1 ml PBS 7.4 and centrifuged at 4000g to concentrate the old population. The two

populations were then stored at -80°C for further biochemical analysis or used immediately for

pH measurements (see below).

Whole Cell Lysates and Immunoblots

Cell pellets from age-enriched young and old cell populations were resuspended in hot

cracking buffer (8 M urea, 5% SDS, 1 mM EDTA, 50 mM Tris-HCl, pH 6.8) and glass beads.

The mixture was vortexed for 10 sec and incubated at 95 °C for 30 sec repeatedly for a total of 5

min. Cellular debris was pelleted by centrifugation at $16,000 \times g$ for 2 min, and supernatants

containing whole-cell extracts were used immediately or stored at −80 °C until use.

After determination of protein concentrations by Bradford assay, equal concentrations of

protein for each sample were separated by SDS-PAGE and transferred to a nitrocellulose

membrane. Blots were blocked for 1 hr in TBST (20 mM Tris-HCl pH 7.5, 150 mM NaCl,

0.05% Tween-20) plus 5% nonfat milk before incubating overnight with primary antibodies at 4

°C with agitation. Primary antibodies (all used at a 1:500 dilution) included mouse monoclonal

antibodies: 7A2 against Vma5, 10D7 against Vph1, 8B1 against Vma1, and 13D11 against Vma2

(Kane et al., 1992). In addition, anti-myc monoclonal 9E10 (Santa Cruz Biotechnology), anti-HA

monoclonal (BioLegend), and anti-GAPDH (Proteintech) antibodies were purchased and used at

1:500, 1:500, and 1:10000, respectively. After washing three times with TBST buffer, HRP-

conjugated anti-mouse secondary antibody (Bio-Rad) was added at a final dilution of 1:2000 and

incubated for 60 min at room temperature. The blot was washed again, incubated with Bio-Rad

Clarity Western ECL substrate and imaged in an Azure Sapphire FL Biomolecular Imager.

Images were quantified using FIJI. Molecular mass markers were included on every blot. In

images of blots, the mass of the marker nearest in size to the protein is indicated.

Vacuolar pH Measurements

Vacuolar pH was measured using the ratiometric fluorescent dye BCECF-AM (Invitrogen) as

described previously(Diakov et al., 2013). Age-enriched populations of cells were loaded with

BCECF-AM in YEP supplemented with 2% or 0.5% glucose. After washing with YEP media to

remove dye, cells were resuspended in YEP, deprived of glucose, and incubated on rotator for 30

min. For fluorescence measurement, 20 µl of cell suspension was diluted into 3 ml 1 mm MES

pH 5.0 buffer, and fluorescence intensity at excitation wavelengths 450 and 495 nm and emission

wavelength 535 nm and was monitored continuously in a Horiba Jovin Yvon Spectrafluor Max

fluorometer with temperature maintained at 30°C. The fluorescence ratio for each sample was

calibrated for each strain in every experiment by clamping the pH to a range of values from 5.0

to 7.0 as described, and the resulting calibration curve was used to convert the experimental

fluorescence ratios to vacuolar pH.

Replicative Life Span

Replicative life span assays were performed on YEP, 2% glucose plates buffered to pH 5.

Daughter cells were sequentially removed by micromanipulation (Steffen, Kennedy, &

Kaeberlein, 2009). Survival curves are pooled data from experiment-matched controls. The

number of divisions for 30 mother cells were scored for each curve. p values for replicative life

span survival curve comparisons were calculated with a Cox regression model. Kaplan-Meier

survival curves were plotted with GraphPad Prism.

RT-PCR of RAV2 from young and old cells

RNA was extracted from young and old cells, obtained by age enrichment as described above,

using the NEB Monarch Total RNA Miniprep Kit . RT-PCR was conducted using the NEB Luna

Universal One-Step RT-qPCR Kit and performed on a Bio-Rad CFX384 Touch System. Data

analysis was conducted using CFX Maestro Software to determine expression levels.

Acknowledgements: This work was supported by NIH R35 GM145256 to P.M.K. The authors

thank M. Murad Khan and Dr. Stephan Wilkens for sharing the *oxr1Δ* strain and Oxr1-HA

plasmid, and Dr. Xin Jie Chen for helpful discussions and a critical reading of the manuscript.

Author contributions: F.H. performed experiments, analyzed data, prepared figures, and wrote

the first draft of the manuscript; P.M.K. obtained funding for the project, analyzed data, prepared

figures, and contributed to writing of the manuscript.

Conflict of interest: The authors declare that they have no conflicts of interest with the contents

of this article.

Data availability: The data that support the findings of this study are openly available in

Upstate.figshare.com at https://upstate.figshare.com. DOI: 10.58120/upstate.26023660

REFERENCES

Bodzeta, A., Kahms, M., & Klingauf, J. (2017). The Presynaptic v-ATPase Reversibly Disassembles and Thereby Modulates Exocytosis but Is Not Part of the Fusion Machinery. *Cell Rep, 20*(6), 1348-1359. doi:10.1016/j.celrep.2017.07.040

Chen, K. L., Ven, T. N., Crane, M. M., Brunner, M. L. C., Pun, A. K., Helget, K. L., . . . Wasko, B. M. (2020). Loss of vacuolar acidity results in iron-sulfur cluster defects and divergent homeostatic responses during aging in Saccharomyces cerevisiae. *Geroscience, 42*(2), 749-764. doi:10.1007/s11357-020-00159-3

Collins, M. P., & Forgac, M. (2020). Regulation and function of V-ATPases in physiology and disease. *Biochim Biophys Acta Biomembr, 1862*(12), 183341. doi:10.1016/j.bbamem.2020.183341

Diab, H. I., & Kane, P. M. (2013). Loss of vacuolar H+-ATPase (V-ATPase) activity in yeast generates an iron deprivation signal that is moderated by induction of the peroxiredoxin TSA2. *Journal of Biological Chemistry, 288*(16), 11366-11377. doi:M112.419259 [pii] 10.1074/jbc.M112.419259

Diakov, T. T., Tarsio, M., & Kane, P. M. (2013). Measurement of vacuolar and cytosolic pH in vivo in yeast cell suspensions. *J Vis Exp, Apr 19*(74). doi:10.3791/50261

Eaton, A. F., Brown, D., & Merkulova, M. (2021). The evolutionary conserved TLDc domain defines a new class of (H(+))V-ATPase interacting proteins. *Sci Rep, 11*(1), 22654. doi:10.1038/s41598-021-01809-y

Ghavidel, A., Baxi, K., Prusinkiewicz, M., Swan, C., Belak, Z. R., Eskiw, C. H., . . . Harkness, T. A. (2018). Rapid Nuclear Exclusion of Hcm1 in Aging Saccharomyces cerevisiae Leads to Vacuolar Alkalization and Replicative Senescence. *G3 (Bethesda), 8*(5), 1579-1592. doi:10.1534/g3.118.200161

Gladyshev, V. N., Kritchevsky, S. B., Clarke, S. G., Cuervo, A. M., Fiehn, O., de Magalhaes, J. P., . . . Cummings, S. R. (2021). Molecular Damage in Aging. *Nat Aging, 1*(12), 1096-1106. doi:10.1038/s43587-021-00150-3

Hansen, M., Rubinsztein, D. C., & Walker, D. W. (2018). Autophagy as a promoter of longevity: insights from model organisms. *Nat Rev Mol Cell Biol, 19*(9), 579-593. doi:10.1038/s41580-018-0033-y

He, C., Zhou, C., & Kennedy, B. K. (2018). The yeast replicative aging model. *Biochim Biophys Acta Mol Basis Dis, 1864*(9 Pt A), 2690-2696. doi:10.1016/j.bbadis.2018.02.023

Henderson, K. A., Hughes, A. L., & Gottschling, D. E. (2014). Mother-daughter asymmetry of pH underlies aging and rejuvenation in yeast. *Elife, 3*, e03504. doi:10.7554/eLife.03504

Hughes, A. L., & Gottschling, D. E. (2012). An early age increase in vacuolar pH limits mitochondrial function and lifespan in yeast. *Nature, 492*(7428), 261-265. doi:nature11654 [pii] 10.1038/nature11654

Hughes, C. E., Coody, T. K., Jeong, M. Y., Berg, J. A., Winge, D. R., & Hughes, A. L. (2020). Cysteine Toxicity Drives Age-Related Mitochondrial Decline by Altering Iron Homeostasis. *Cell, 180*(2), 296-310 e218. doi:10.1016/j.cell.2019.12.035

Huh, W. K., Falvo, J. V., Gerke, L. C., Carroll, A. S., Howson, R. W., Weissman, J. S., & O'Shea, E. K. (2003). Global analysis of protein localization in budding yeast. *Nature, 425*(6959), 686-691. Retrieved from

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citati
on&list_uids=14562095
Jaskolka, M. C., Tarsio, M., Smardon, A. M., Khan, M. M., & Kane, P. M. (2021). Defining steps
in RAVE-catalyzed V-ATPase assembly using purified RAVE and V-ATPase
subcomplexes. *J Biol Chem, 296*, 100703. doi:10.1016/j.jbc.2021.100703
Jaskolka, M. C., Winkley, S. R., & Kane, P. M. (2021). RAVE and Rabconnectin-3 Complexes as
Signal Dependent Regulators of Organelle Acidification. *Front Cell Dev Biol, 9*, 698190.
doi:10.3389/fcell.2021.698190
Jin, X., Cao, X., & Liu, B. (2021). Isolation of Aged Yeast Cells Using Biotin-Streptavidin Affinity
Purification. *Methods Mol Biol, 2196*, 223-228. doi:10.1007/978-1-0716-0868-5_17
Kane, P. M. (1995). Disassembly and reassembly of the yeast vacuolar H(+)-ATPase in vivo. *J
Biol Chem, 270*(28), 17025-17032.
Kane, P. M., Kuehn, M. C., Howald-Stevenson, I., & Stevens, T. H. (1992). Assembly and targeting
of peripheral and integral membrane subunits of the yeast vacuolar H(+)-ATPase. *J Biol
Chem, 267*(1), 447-454.
Kaushik, S., Tasset, I., Arias, E., Pampliega, O., Wong, E., Martinez-Vicente, M., & Cuervo, A. M.
(2021). Autophagy and the hallmarks of aging. *Ageing Res Rev, 72*, 101468.
doi:10.1016/j.arr.2021.101468
Khan, M. M., Lee, S., Couoh-Cardel, S., Oot, R. A., Kim, H., Wilkens, S., & Roh, S. H. (2022).
Oxidative stress protein Oxr1 promotes V-ATPase holoenzyme disassembly in catalytic
activity-independent manner. *EMBO J, 41*(3), e109360. doi:10.15252/embj.2021109360
Khan, M. M., & Wilkens, S. (2024). Molecular mechanism of Oxr1p mediated disassembly of
yeast V-ATPase. *EMBO Rep*. doi:10.1038/s44319-024-00126-5
Klossel, S., Zhu, Y., Amado, L., Bisinski, D. D., Ruta, J., Liu, F., & Gonzalez Montoro, A. (2024).
Yeast TLDc domain proteins regulate assembly state and subcellular localization of the V-
ATPase. *EMBO J*. doi:10.1038/s44318-024-00097-2
Kohio, H. P., & Adamson, A. L. (2013). Glycolytic control of vacuolar-type ATPase activity: a
mechanism to regulate influenza viral infection. *Virology, 444*(1-2), 301-309. doi:S0042-
6822(13)00396-6 [pii]
10.1016/j.virol.2013.06.026
Kurz, T., Terman, A., Gustafsson, B., & Brunk, U. T. (2008). Lysosomes in iron metabolism,
ageing and apoptosis. *Histochem Cell Biol, 129*(4), 389-406. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citati
on&list_uids=18259769
Liu, Y., Steinbusch, L. K. M., Nabben, M., Kapsokalyvas, D., van Zandvoort, M., Schonleitner, P.,
. . . Luiken, J. (2017). Palmitate-Induced Vacuolar-Type H(+)-ATPase Inhibition Feeds
Forward Into Insulin Resistance and Contractile Dysfunction. *Diabetes, 66*(6), 1521-1534.
doi:10.2337/db16-0727
Longo, V. D., & Anderson, R. M. (2022). Nutrition, longevity and disease: From molecular
mechanisms to interventions. *Cell, 185*(9), 1455-1470. doi:10.1016/j.cell.2022.04.002
Longtine, M. S., McKenzie, A., 3rd, Demarini, D. J., Shah, N. G., Wach, A., Brachat, A., . . .
Pringle, J. R. (1998). Additional modules for versatile and economical PCR-based gene
deletion and modification in Saccharomyces cerevisiae. *Yeast, 14*(10), 953-961.
Martinez-Munoz, G. A., & Kane, P. (2008). Vacuolar and plasma membrane proton pumps
collaborate to achieve cytosolic pH homeostasis in yeast. *J Biol Chem, 283*(29), 20309-
20319. doi:M710470200 [pii]

10.1074/jbc.M710470200

Nixon, R. A. (2020). The aging lysosome: An essential catalyst for late-onset neurodegenerative
diseases. *Biochim Biophys Acta Proteins Proteom, 1868*(9), 140443.
doi:10.1016/j.bbapap.2020.140443

Oot, R. A., & Wilkens, S. (2020). A "Sugar-Coated" Proton Pump Comes into Focus: High-
Resolution Structure of a Human V-ATPase. *Mol Cell, 80*(3), 379-380.
doi:10.1016/j.molcel.2020.10.020

Parra, K. J., & Kane, P. M. (1998). Reversible association between the V1 and V0 domains of yeast
vacuolar H+-ATPase is an unconventional glucose-induced effect. *Mol Cell Biol, 18*(12),
7064-7074. Retrieved from http://mcb.asm.org/cgi/content/full/18/12/7064

Perera, R. M., & Zoncu, R. (2016). The Lysosome as a Regulatory Hub. *Annu Rev Cell Dev Biol,*
32, 223-253. doi:10.1146/annurev-cellbio-111315-125125

Ratto, E., Chowdhury, S. R., Siefert, N. S., Schneider, M., Wittmann, M., Helm, D., & Palm, W.
(2022). Direct control of lysosomal catabolic activity by mTORC1 through regulation of
V-ATPase assembly. *Nat Commun, 13*(1), 4848. doi:10.1038/s41467-022-32515-6

Schiffer, I., Gerisch, B., Kawamura, K., Laboy, R., Hewitt, J., Denzel, M. S., . . . Antebi, A. (2021).
miR-1 coordinately regulates lysosomal v-ATPase and biogenesis to impact proteotoxicity
and muscle function during aging. *Elife, 10*. doi:10.7554/eLife.66768

Seol, J. H., Shevchenko, A., & Deshaies, R. J. (2001). Skp1 forms multiple protein complexes,
including RAVE, a regulator of V-ATPase assembly. *Nat Cell Biol, 3*(4), 384-391.

Smardon, A. M., Tarsio, M., & Kane, P. M. (2002). The RAVE complex is essential for stable
assembly of the yeast V-ATPase. *J Biol Chem, 277*(16), 13831-13839.

Steffen, K. K., Kennedy, B. K., & Kaeberlein, M. (2009). Measuring replicative life span in the
budding yeast. *J Vis Exp*(28). doi:10.3791/1209

Stransky, L. A., & Forgac, M. (2015). Amino Acid Availability Modulates Vacuolar H+-ATPase
Assembly. *Journal of Biological Chemistry, 290*(45), 27360-27369. doi:M115.659128 [pii]
10.1074/jbc.M115.659128

Sumner, J. P., Dow, J. A., Earley, F. G., Klein, U., Jager, D., & Wieczorek, H. (1995). Regulation
of plasma membrane V-ATPase activity by dissociation of peripheral subunits. *J Biol*
Chem, 270(10), 5649-5653.

Tabke, K., Albertmelcher, A., Vitavska, O., Huss, M., Schmitz, H. P., & Wieczorek, H. (2014).
Reversible disassembly of the yeast V-ATPase revisited under in vivo conditions.
Biochemical Journal, 462(1), 185-197. doi:BJ20131293 [pii]
10.1042/BJ20131293

Vilchez, D., Saez, I., & Dillin, A. (2014). The role of protein clearance mechanisms in organismal
ageing and age-related diseases. *Nat Commun, 5*, 5659. doi:10.1038/ncomms6659

Wilkens, S., Khan, M. M., Knight, K., & Oot, R. A. (2023). Tender love and disassembly: How a
TLDc domain protein breaks the V-ATPase. *Bioessays, 45*(7), e2200251.
doi:10.1002/bies.202200251

Wilms, T., Swinnen, E., Eskes, E., Dolz-Edo, L., Uwineza, A., Van Essche, R., . . . Winderickx, J.
(2017). The yeast protein kinase Sch9 adjusts V-ATPase assembly/disassembly to control
pH homeostasis and longevity in response to glucose availability. *PLoS Genet, 13*(6),
e1006835. doi:10.1371/journal.pgen.1006835

Yan, Y., Denef, N., & Schupbach, T. (2009). The vacuolar proton pump, V-ATPase, is required for
notch signaling and endosomal trafficking in Drosophila. *Dev Cell, 17*(3), 387-402.
doi:S1534-5807(09)00260-3 [pii]

10.1016/j.devcel.2009.07.001
661

 FIGURE LEGENDS

Figure 1: V-ATPases are more disassembled in yeast cells of older replicative age. (a)

BY4742 cells expressing Vma5-GFP grown in SC containing 2% glucose. Differential

interference contrast microscopy (DIC) was used to visualize vacuoles. Bud scars are stained

with calcofluor white (CW) to determine replicative age. **(b)** Left panel provides representative

example of quantitative measurements using line scans. The young cell is from the dashed white

box in the Vma5-GFP image of **Figure 1a** and the old cell is from the dashed red box. Plot

profiles are superimposed for the young cell (black) and old cell (red). **(c)** Quantitation of

maximum fluorescent intensity across five biological replicates, after normalization to the

average intensity in the youngest bin for each replicate. Each biological replicate (dot) represents

at least 20 cells and bars represent the mean +/- s.e.m. Significance was calculated by ordinary

one-way ANOVA. "Old" for subsequent experiments (without age enrichment) is categorized at

≥ 5 bud scars. **** represents a p-value <0.0001.

Figure 2: Age-enriched populations of cells have comparable levels of V-ATPase subunits.

(a) Diagram of V-ATPase showing the relative positions of Vma5, Vma2, and Vph1 (Image was

prepared with Biorender.com). **(b)** BY4742 cells expressing Vma2-GFP (V_1B) grow in SC with

2% glucose. CW was used to visualize bud scars. The CW images for young cells were

overexposed relative to those for the old cells in order to visualize the low level staining in cells

with few bud scars. Normalized maximum fluorescence was obtained through line scan

quantitation using FIJI as in **Figure 1**. Means +/- s.e.m. of three biological replicates are shown;

each replicate is represented by a dot. Significance was calculated by unpaired Student's *t* test,

*** p value=0.0009 . **(c)** Cells expressing Vph1-GFP were visualized and analyzed as described

in **2b**. **(d)** After biotin-streptavidin age enrichment, bud scars were counted for 100 cells in

young and old populations, binned by bud count, and the age distribution plotted. **(e)** Lysates

were prepared from age-enriched populations defined as in **1c** and equal protein concentrations

were separated by SDS-PAGE and examined by immunoblot for V-ATPase protein levels in

young (Y) vs. old (O) cell populations defined as in **1c**. **(f)** Band intensities were quantified using

FIJI, ratios of V-ATPase subunit levels to the GAPDH internal control were calculated, and ratios

were normalized to the young population for each biological replicate. Significance calculated by

unpaired Student's *t* test. Data are presented as mean (horizontal bars) ± s.e.m. (whiskers) of

three biological replicates. n.s.= not significant.

Figure 3: Vacuolar pH is more alkaline in old cells. (a) Vacuolar pH responses were measured

for wild-type BY4742 age-enriched young and old populations as described in Methods.

Glucose-deprived cultures were loaded with BCECF-AM. Fluorescence intensity values were

collected every 10 sec at excitation wavelengths 450 and 490 nm and emission wavelength 535

nm, and glucose was added to a final concentration of 2% after 3 min. The ratio of fluorescence

signals from the two excitation wavelengths was calculated and converted to pH via a calibration

curve. **(b)** Fluorescence measurements at 1 min. (before glucose addition), 5 min (2 min. after

glucose addition), and 8 min. (5 min after glucose addition). Calculated pH measurements are

presented as mean (horizontal bars) ± s.e.m. (whiskers) of three biological replicates. * indicates

a p value of 0.02, ** represents a p value of 0.001.

Figure 4: Caloric restriction (0.5% glucose) restores V-ATPase assembly and vacuolar pH

in old cells. (a), (b), (c) Strains used in **Figures 1** and **2** show recruitment of Vma5-GFP **4a** and

Vma2-GFP **4b** to the vacuolar membrane after growth in SC with 0.5% glucose while Vph1-GFP

4**c** remains unchanged. Maximum fluorescence was measured and normalized as described in

Methods and **Figure 1**. CW was visualized in young and old cells as described in **Figure 2b**. No

significant differences between young and old cells was observed as calculated by unpaired

Student's *t* test. **(d)**Young and old cell populations were obtained by biotin-streptavidin age-

enrichment from cells grown in YEP supplemented with 0.5% glucose, and vacuolar pH

responses were measured. Data were collected and analyzed as in **Figure 3b**. Quantitated data

are presented as mean (horizontal bars) ± s.e.m. (whiskers) of three biological replicates.

Figure 5: Effects of *rav1Δ* and *oxr1Δ* on replicative lifespan. (a) Schematic of reversible

disassembly highlighting the roles of the RAVE complex and Oxr1 (Image was prepared with

Biorender.com). **(b)** Kaplan-Meier curves comparing the replicative lifespan (RLS) of *rav1Δ*

(green), *oxr1Δ* (red), and wild-type cells (black). Median number of replicative generations is

shown in parentheses for each strain, and the difference between *rav1Δ* and *oxr1Δ* median values

and wild-type are expressed as %. Deletion of *OXR1* significantly increases yeast RLS

(p<0.001), and deletion of RAVE component *RAV1* shortens RLS (p < 0.01).

Figure 6: Rav2 level is reduced during replicative aging, but restored by CR. (a) Analysis of

Rav1 levels in young and old cells. Age-enriched populations were obtained from BY4741 cells

containing Rav1-myc$_{13}$ (young (Y) versus old (O) cells), and cell lysates were prepared,

separated, and quantitated as described in Methods and **Figure 2**. Normalized data are presented

as mean (horizontal bars) ± s.e.m. (whiskers) of three biological replicates. ns=not significant **(b)**

Immunoblot analysis of BY4741 Rav2-myc$_{13}$ from young and old cells prepared as in **6a**.

Normalized data are presented as mean (horizontal bars) ± s.e.m. (whiskers) of three biological

replicates. P-value 0.0094. **(c)** Quantitative RT-PCR comparing expression of *RAV2* mRNA

between young and old cells. **(d)** Wild-type and *oxr1Δ* cells containing Vma5-GFP were grown

in SC containing 2% glucose. DIC used to visualize vacuoles and CW used to visualize bud scars

as described in **Figure 2b**. **(e)** BY4741 *oxr1*Δ cells were binned by the number of bud scars and

normalized maximal fluorescence quantitated as described in **Figure 1**. *** indicates a P value

of 0.001 in comparison to the youngest bin; other bins are not significantly different from the

youngest bin. **(f)** Immunoblot analysis of a BY4741 *oxr1*Δ strain expressing Oxr1-HA from a

low copy plasmid. Young and old cell populations were isolated as in **6a**. Data are presented as

mean (horizontal bars) ± s.e.m. (whiskers) of three biological replicates. **(g)** Immunoblot analysis

comparing Rav2-myc$_{13}$ levels in cells grown in YEP supplemented with 2% glucose and 0.5%

glucose (CR conditions). Samples were prepared and analyzed as in **6b**. Quantification of data

are presented as mean (horizontal bars) ± s.e.m. (whiskers) of three biological replicates. ***

indicates a P value of 0.0001, n.s.= not significant.

Supporting information, Figure 1: Cells show similar growth rates over 12 hours in 2%

and 0.5% glucose. BY4742 cells at log phase were diluted into YEP containing either 2% or

0.5% glucose as indicated. OD$_{600}$ was measured every hour for 12 hours.

a.

| DIC | Calcofluor White | Vma5-GFP |

b.

Young

CW

GFP

Old

CW

GFP

c.

**Vma5-GFP
Vacuole Localization**

Figure 1

Figure 2

Figure 3

Figure 4

a.

V$_1$

V$_o$

ACTIVE

Oxr1

RAVE

V$_1$

V$_o$

INACTIVE

Vma5

b.

rav1Δ (17, -26.1%)

WT (23)

oxr1Δ (34, +47.8%)

% Survival

Generation

Figure 5

Figure 6