V-ATPase Disassembly at the Yeast Lysosome-Like Vacuole Is a Phenotypic Driver of Lysosome Dysfunction in Replicative Aging

Fiza Hashmi and Patricia M. Kane

Department of Biochemistry and Molecular Biology
SUNY Upstate Medical University, Syracuse, NY USA

Address correspondence to: Patricia M. Kane
kanepm@upstate.edu

Running title: V-ATPase disassembly in replicative aging

Declines in lysosomal acidification and function with aging are observed in organisms

ranging from yeast to humans. V-ATPases play a central role in organelle acidification and V-

ATPase activity is regulated by reversible disassembly in many different settings. Using the yeast

Saccharomyces cerevisiae as a replicative aging model, we demonstrate that V-ATPases

disassemble into their V_1 and V_0 subcomplexes in aging cells, with release of V_1 subunit C

(Vma5) from the lysosome-like vacuole into the cytosol. Disassembly is observed after ≥ 5 cell

divisions and results in overall vacuole alkalinization. Caloric restriction, an established

mechanism for reversing many age-related outcomes, prevents V-ATPase disassembly in older

cells and preserves vacuolar pH homeostasis. Reversible disassembly is controlled in part by the

activity of two opposing and conserved factors, the RAVE complex and Oxr1. The RAVE

complex promotes V-ATPase assembly and a *rav1Δ* mutant shortens replicative lifespan; Oxr1

promotes disassembly and an *oxr1Δ* mutation extends lifespan. Importantly, the level of Rav2, a

subunit of the RAVE complex, declines in aged cells, and Rav2 overexpression delays V-ATPase

disassembly with age. These data indicate that reduced V-ATPase assembly contributes to the

loss of lysosome acidification with age, which affects replicative lifespan.

Keywords: Aging, lysosomes, proton pumps, Saccharomyces cerevisiae, caloric restriction

INTRODUCTION

Lysosomal acidification is compromised with age in many organisms (Nixon, 2020).

Reduced acidification has multiple downstream functional consequences. Lysosomal hydrolases

operate optimally at the acidic pH, so cargo degradation is compromised at higher pH (Vilchez,

Saez, & Dillin, 2014). Lysosomes are terminal compartments for autophagy pathways, so

clearance of autophagic cargoes and recycling of nutrients, both critical in aging cells, is slowed

(Hansen, Rubinsztein, & Walker, 2018; Kaushik et al., 2021). Iron and other heavy metals are

sequestered in the acidic lysosomes; loss of sequestration can induce both oxidative stress (Diab

& Kane, 2013) and deficiency in mitochondrial iron-sulfur proteins (Chen et al., 2020). Reduced

lysosomal storage can also create toxic imbalances in amino acids such as cysteine that

contribute to loss of mitochondrial function (C. E. Hughes et al., 2020). Recent work highlights

the central role of the lysosome in nutritional signaling; many aspects of this signaling are linked

to acidification (Perera & Zoncu, 2016). Loss of lysosomal acidification impacts many processes

associated with age-related functional decline, but the mechanisms behind increased lysosomal

pH are not fully understood.

The highly conserved vacuolar H^+- ATPase (V-ATPase) acidifies lumens of lysosomes

and lysosome-like vacuoles, as well as endosomes and the late Golgi apparatus (Collins &

Forgac, 2020). V-ATPases are multi-subunit protein complexes that couple ATP hydrolysis to

proton pumping into organelle lumens. The V-ATPase consists of two subcomplexes: a

peripheral V_1 subcomplex that is responsible for ATP hydrolysis connected to a membrane-

embedded V_0 subcomplex containing the proton pore. V-ATPase subunit sequences are

conserved across eukaryotes and recent V-ATPase structures indicate very strong structural

similarity between yeast and mammalian V-ATPases (Oot & Wilkens, 2020).

V-ATPase activity is highly regulated and responsive to environmental conditions.

Reversible disassembly is a versatile mechanism of V-ATPase regulation that fine-tunes the

activity of the proton pump to meet cellular demands (Collins & Forgac, 2020; Jaskolka,

Winkley, & Kane, 2021). In reversible disassembly, the V_1 subcomplex is released from the V_0

subcomplex inhibiting both ATP hydrolysis and proton pumping (Kane, 1995; Sumner et al.,

1995). V_1 subunit C dissociates from both subcomplexes and becomes cytosolic during

disassembly (Kane, 1995). Disassembly is post-translational and rapidly reversible (Kane,

1995). It was first observed in the yeast *S. cerevisiae* and the tobacco hornworm *M. sexta* upon

acute glucose deprivation and was reversed by glucose replenishment (Kane, 1995; Sumner et

al., 1995). It has since become clear that reversible disassembly occurs widely and in response to

diverse signals. For example, many cells appear to promote V-ATPase reassembly under

conditions of nutrient deprivation and mTOR inhibition, possibly as a means of promoting

lysosomal proteolysis and nutrient recycling (Ratto et al., 2022). In neurons, V-ATPases are

reversibly disassembled as part of each synaptic vesicle cycle (Bodzeta, Kahms, & Klingauf,

2017). In cardiomyocytes, lipid overload can promote V-ATPase disassembly in endosomes,

ultimately contributing to long-term insulin resistance (Liu et al., 2017). Reversible disassembly

can also be manipulated by both host cells and pathogens to prevent or facilitate infection (Kohio

& Adamson, 2013).

The RAVE (Regulator of Acidification of Vacuoles and Endosomes) complex and TLDc

(Tre2/Bub2/Cdc16 LysM domain catalytic) protein Oxr1 regulate reversible disassembly of the

V-ATPase (Jaskolka, Winkley, et al., 2021; Khan et al., 2022; Klossel et al., 2024; Seol,

Shevchenko, & Deshaies, 2001). The RAVE complex consists of three subunits: Rav1, Rav2, and

Skp1 and is required for reassembly of V-ATPase complexes disassembled by glucose

deprivation. In mutants lacking Rav1 or Rav2, V-ATPases are disassembled into V_1 and V_0

subcomplexes and vacuolar acidification is lost (Jaskolka, Winkley, et al., 2021). The role of

Skp1 in RAVE is less clear and it may be dispensable for RAVE function (Jaskolka, Winkley, et

al., 2021).   The RAVE complex associates with V_1 subcomplexes in the cytosol (Jaskolka,

Tarsio, Smardon, Khan, & Kane, 2021). In contrast, recent studies suggest that Oxr1, a member

of the TLDc family of proteins, promotes disassembly of V-ATPases. Mutants lacking Oxr1

maintain higher levels of assembled V-ATPases in the absence of glucose (Khan et al., 2022;

Khan & Wilkens, 2024; Klossel et al., 2024).

Several mechanisms for loss of lysosomal acidification with age have been proposed. In

yeast, the long-lived plasma membrane proton pump, Pma1, accumulates in mother cells, and it

has been proposed that increased proton export through Pma1 disrupts the balance of Pma1 and

V-ATPase activities and compromises organelle acidification (Devare et al., 2020; Henderson,

Hughes, & Gottschling, 2014). Early loss of vacuolar acidification in yeast has been correlated

with defective mitochondrial morphology and function. Both vacuolar deacidification and

mitochondrial morphology phenotypes are suppressed by overexpression of the V-ATPase

catalytic subunit, *VMA1*, or an ER-localized assembly factor, *VPH2 (VMA12)*, suggesting a

possible deficiency in these factors with age (A. L. Hughes & Gottschling, 2012). In *C. elegans*,

stability of the *VMA1* transcript is controlled by a microRNA, miR-1, which can globally control

lysosomal acidification (Schiffer et al., 2021). Wilms et al. (Wilms et al., 2017) demonstrated

that deletion of the mTOR effector Sch9 promotes V-ATPase assembly and extends yeast

chronological lifespan (CLS). All of these mechanisms could contribute to loss of

vacuolar/lysosomal acidification.

However, despite the importance of reversible disassembly in V-ATPase regulation, V-

ATPase assembly state during aging has not been explored extensively. Here we show that V-

ATPase assembly changes with age in a yeast replicative aging model. Increased V-ATPase

disassembly in older cells is accompanied by decreased vacuolar acidification that does not

appear to stem from reduced V-ATPase subunit levels. Instead, we provide evidence that reduced

activity of the RAVE (Regulator of H^+-ATPase of Vacuolar and Endosomal membranes)

assembly complex may give rise to net V-ATPase disassembly and increased lysosomal pH with

age.

RESULTS

V-ATPases are more disassembled and vacuoles more alkaline after ~5 cell divisions.

Given the evidence that vacuoles and lysosomes are less acidic in older cells, we

hypothesized that V-ATPase assembly and activity might also change with age. Replicative aging

in the yeast *S. cerevisiae* is a widely accepted model for aging (He, Zhou, & Kennedy, 2018).

Yeast cells divide asymmetrically with each cell division giving rise to a new, "rejuvenated"

daughter cell from an established mother cell. Each cell division leaves a bud scar on the mother,

allowing visual assessment of age. The yeast V_1C subunit (Vma5) is released from both V_1 and

V_0 during V-ATPase disassembly, so we first visualized Vma5-GFP localization in a mixed age

population of cells (Figure 1). (Vacuoles are visible as indentations under DIC (differential

interference contrast) optics.) In parallel, we monitored replicative age of each cell by staining

with calcofluor white (CW) which labels the bud scars on mother cells. As shown in Figure 1a,

Vma5-GFP is tightly localized to the vacuolar membrane in cells with few or no bud scars. In

contrast, in older cells with more bud scars, Vma5-GFP exhibited a marked decrease in

fluorescence at the vacuolar membrane and a notable increase in cytosolic fluorescence. To

compare Vma5-GFP localization between cells, we quantitated maximum fluorescence from a

line scan across each cell. As shown in Figure 1b, a line scan from a young cell has prominent

peaks corresponding to edges of the vacuole with a high maximum fluorescence signal, while

older cells have less prominent peaks. We conducted the same analysis across populations of

cells of mixed age, normalized to the maximum fluorescence signal of young cells and binned

the results by the number of bud scars. As shown in Figure 1c, young cells, defined as having

less than 5 bud scars, displayed Vma5-GFP localization at the vacuolar membrane. However, the

normalized maximum fluorescence signal, representative of vacuole localization, decreases

significantly in cells with five bud scars or more. This early onset of V-ATPase disassembly

aligns with previous reports indicating declines in lysosomal pH early in replicative aging (A. L.

Hughes & Gottschling, 2012).

Vma5 is a V_1 subunit that bridges the V_1 and V_0 subcomplexes of the V-ATPase. The V_1

subcomplex also contains Vma2, and the V_0 subcomplex contains Vph1, which comprises part of

the proton pore (Figure 2a). Vma5 appears to be released most completely from the membrane

by reversible disassembly (Tabke et al., 2014), but the rest of V_1 also dissociates from the

vacuolar membrane. We assessed the cellular distribution of Vma2-GFP and Vph1-GFP. There

was less membrane-bound Vma2-GFP in older cells relative to younger cells (Figure 2b), when

assessed as in Figure 1. However, vacuolar Vph1-GFP levels were the same between old and

young cells (Figure 2c). These results are consistent with disassembly of the V-ATPase as cells

age.

Although comparable levels of Vph1 at the vacuole suggests that expression of V_0

subunits and V_0 assembly are intact, V_1 subunits could become cytosolic because of reduced V_1

subunit levels in older cells. To address this question, we isolated populations old and young

yeast cells by biotinylating the cell walls in a mixed age population, allowing growth to continue

for several generations, and then obtaining "old" cells by biotin-streptavidin affinity

chromatography (Jin, Cao, & Liu, 2021). Daughter cells that emerged after biotinylation cannot

bind to magnetic streptavidin beads and represent the "young" population. The age distribution

was determined by counting bud scars in each population and binning by the number of buds per

cell (Figure 2d). When prepared by this method, the population of old cells peaks at 20-24 bud

scars, while there was a median value of 0-4 bud scars in the young population. Cell lysates were

prepared from each population and analyzed by SDS-PAGE and immunoblotting. As shown in

Figures 2e and f, there is no significant difference in the cellular levels of V_1 subunits Vma1,

Vma2, and Vma5 between young and old populations. These results indicate that the cytosolic

populations of V_1 subunits arise from disassembly of the V-ATPase, rather than inability to

assemble because of lack of V-ATPase subunits.

Reversible disassembly of V-ATPases is employed in a number of contexts to provide

dynamic regulation of the complex in response to changing cellular conditions. Disassembled V_1

and V_0 subcomplexes lack ATPase and proton transport activity and ATP-driven proton pumping

is restored upon reassembly. In order to test whether the lower levels of assembled V-ATPases in

old cells result in reduced capacity for acidification, we measured the response of young and old

cells to an acute glucose deprivation (an abrupt shift to 0% glucose), which promotes

disassembly of the yeast V-ATPase, followed by readdition of glucose to a 2% final

concentration, which promotes reassembly and reactivation of the complex (Kane, 1995).

Young and old yeast cells obtained as described above were loaded with the ratiometric

pH sensor BCECF-AM (2',7'-Bis-(2-carboxyethyl)-5-(and-6)-carboxyfluorescein acetoxy methyl

ester) which localizes to the vacuole in yeast cells (Diakov, Tarsio, & Kane, 2013). Both young

and old cells were shifted to medium with no glucose for ~30 minutes. Fluorescence of cell

suspensions was then monitored continuously (Figure 3a) and glucose was added at the indicated

time. This assay revealed clear age-dependent differences in the vacuolar pH response to glucose

stimulation. Young cells exhibit a rapid drop in vacuolar pH upon glucose addition. This drop

was previously shown to be V-ATPase-dependent and to correlate with V-ATPase assembly

(Martinez-Munoz & Kane, 2008). In old cells, however, there was a smaller pH decrease after

glucose addition, indicating a more alkaline vacuole and consistent with the lower levels of V-

ATPase assembly observed by microscopy during growth in glucose-replete media in Figures 1

and 2. Quantitative analysis of pH at defined time points across biological replicates (Figure 3b)

indicates that vacuoles in old cells are significantly more alkaline than their younger counterparts

at each of the indicated time points. These results suggest an age-related alteration in vacuolar

pH regulation through the inability of V-ATPase to reassemble. As a result, vacuoles in old cells

display a more alkaline pH than vacuoles in young cells.

Caloric restriction restores V-ATPase assembly and vacuolar acidification in older cells.

Caloric restriction (CR) is defined as a reduction in caloric intake in the presence of

adequate nutrition (Longo & Anderson, 2022). CR promotes both cellular health and longevity.

Extensive research in organisms ranging from yeast to worms, flies, and rodents suggests that

CR can have significant anti-aging effects and promote overall health (Longo & Anderson,

2022). For *S. cerevisiae*, adjusting the concentration of glucose in the growth medium from the

2% used above to 0.5% is a common way to induce CR. This treatment does not significantly

reduce growth rate over several cell divisions (Supporting information Figure 1a).

As shown in Figure 4, CR reverses the V-ATPase disassembly in older cells. Notably, the

V_1 subunits Vma5-GFP (Figure 4a) and Vma2-GFP (Figure 4b) are recruited to the vacuolar

membrane in both young and old cells, and there is no significant difference in fluorescence at

the vacuolar membrane with age. Fluorescence intensities of Vph1-GFP continue to be similar

between old and young cells (Figure 4c). This suggests that CR extends V-ATPase assembly

beyond the replicative age of ≥ 5 bud scars when cells grown in higher glucose begin to show

disassembly.

We hypothesized that vacuolar pH in old cells might also be restored. We grew cells

under CR conditions and monitored vacuolar pH before and after addition of glucose as

described above. Figure 4d demonstrates that the glucose-stimulated decrease in vacuolar pH,

which was compromised in older cells grown in 2% glucose (Figure 3), was restored to the level

of young cells in cells after growth under CR conditions. This observation indicates that CR has

a direct impact on both V-ATPase assembly and vacuolar acidification in aging cells. In addition,

it further highlights the potential connection between V-ATPase assembly, vacuolar acidification

and aging.

These results point to nutritional signaling as an important determinant of V-ATPase

assembly and activity in aging cells. Bond and Forgac demonstrated that V-ATPase assembly

was preserved under low glucose conditions in an *ira2Δ* mutant. Deletion of *IRA2*, or its

paralogue *IRA1*, activates the Ras/protein kinase A (PKA) pathway (Bond & Forgac, 2008). We

hypothesized that V-ATPase assembly might be preserved in older cells, mimicking the effects of

CR, in an *ira2Δ* mutant. We visualized Vma5-GFP localization and as shown in Supporting

Information Figure 1b, the *ira2Δ* cells do maintain V-ATPase assembly in older cells. Therefore,

we determined replicative lifespan (RLS) for the *ira2Δ* mutant by daughter dissection

(Supporting Information Figure 1c).   The median lifespan of the *ira2Δ* mutant is significantly

shorter than that of wild-type cells.   These data are consistent with evidence that upregulation of

the Ras/protein kinase A pathway shortens lifespan (Longo & Anderson, 2022) and suggest that

although there is improved V-ATPase assembly in older cells in the *ira2Δ* mutant, it is not

sufficient to balance effects on other targets of the pathway in aging cells.

Regulators of V-ATPase assembly affect replicative lifespan.

We next sought to manipulate V-ATPase assembly more directly in order to assess the

significance of maintaining V-ATPase assembly and activity in aging cells. We hypothesized

that the RAVE complex and Oxr1, cellular factors that promote V-ATPase assembly and

disassembly as shown in Figure 5a, might also affect RLS.

To explore the functional significance of these assembly regulators in aging cells, we

examined the RLS of deletion mutants lacking Rav1 or Oxr1 (Figure 5b). RLS was measured at

pH 5, conditions that are optimal for growth of *rav1Δ* strains. Deletion of Rav1 shortened RLS

(median 17 cell divisions, n= 30) by 26% relative to wild-type cells (median 23 cell divisions, n=

30). In contrast, the *oxr1Δ* mutation extended RLS by 47.8% over wild-type (median 34 cell

divisions, n=30). These results reinforce the significance of V-ATPase assembly in RLS and

suggest that the RAVE pro-assembly activity (disrupted in the *rav1Δ* mutant) and Oxr1 anti-

assembly activity (lost in *oxr1Δ*) may be central determinants of lifespan.

We next asked whether loss of RAVE and Oxr1 affects chronological aging of yeast cells.

Replicative aging assesses the number of times a mother cells can divide, while chronological

aging assesses cell survival as nutrients are depleted over time. Wild-type cells and *rav1Δ*,

oxr1Δ, and *vma1Δ* mutants were grown to saturation in medium buffered to pH 5.5 (Murakami et

al., 2012; Wilms et al., 2017). (*VMA1* encodes the catalytic subunit of the V-ATPase and the

vma1Δ mutant has no V-ATPase activity.). At the indicated times, aliquots were removed from

each culture, cell density was determined, and a defined number of cells was plated on fresh

medium.   The percentage of each cell type that formed colonies (c.f.u.) on fresh plates at each

time is shown in Figure 5c. Even in buffered medium, which improves the survival of *vma*

mutants (Wilms et al., 2017), less than half of the *vma1Δ* and *rav1Δ* mutant cells are able to form

colonies after 2 weeks. In contrast, almost half of the *oxr1Δ* cells form colonies at 4 weeks, and

wild-type cells have an intermediate phenotype. The percentage of cells staining with propidium

iodide, an indication of cell permeabilization and death, increases as colony formation declines

(Figure 5d). However, it is notable that even after 5 weeks, only ~60% of *oxr1Δ* cells stain with

propidium iodide. These results complement the replicative aging data and indicate that

mutations that preserve V-ATPase assembly can significantly increase lifespan, while mutations

that compromise activity and/or assembly decrease lifespan.   Yeast encode a second isoform of

the V_0 a-subunit, Stv1, which localizes to the Golgi in wild-type cells.   Stv1-containing V-

ATPases do not require RAVE for assembly (Smardon et al., 2014), and Stv1 has been shown to

relocalize to the vacuole in an *oxr1Δ* mutant (Klossel et al., 2024).   In order to test whether the

lifespan extension in *oxr1Δ* cells arises from Stv1-containing V-ATPases at the vacuole, we

constructed a *stv1Δoxr1Δ* double mutant strain and assessed its RLS and V-ATPase assembly. As

shown in Supporting Information Figure 2a, the double mutant strain retains the long lifespan of

the *oxr1Δ* relative to wild-type, suggesting that vacuolar Stv1 is not responsible for this effect.

Rather, V-ATPases containing the Vph1 isoform account for the assembly in older cells of the

oxr1Δ mutant (Supplemental Information Figure 2b).

Given the results with these deletion mutants, we asked whether there were differences in

levels of the RAVE subunits or Oxr1 between young and old cells. We isolated young and old

populations of cells containing functional myc13-tagged Rav1 or FLAG-tagged Rav2 by

biotinylation and streptavidin magnetic separation as described above (Figure 2), then assessed

the levels of the tagged proteins in cell lysates. Although there is no significant difference in

Rav1-myc13 levels between young and old cells (Figure 6a), we consistently observed a

significant decrease in the level of Rav2-FLAG in older cells (Figure 6b). We assessed

expression of *RAV2* in young and old cells by quantitative PCR but observed no difference in

mRNA levels (Supporting Information Figure 2c). Because Rav2 is required for RAVE complex

function in promoting V-ATPase assembly (Seol et al., 2001; Smardon, Tarsio, & Kane, 2002),

these results suggest that partial loss of RAVE function in old cells could contribute to reduced

V-ATPase assembly and replicative aging.   We transformed the cells containing a chromosomal

copy of Rav2-FLAG with an additional copy of Rav2-FLAG on a low-copy plasmid to

determine whether we could increase Rav2 levels and improve V-ATPase assembly in older cells.

As shown in Figure 6b, addition of the plasmid increases Rav2-FLAG levels in old cells. We

compared Vma5-GFP localization with and without the Rav2-FLAG plasmid in a mixed age

population in Figure 6c. In cells overexpressing Rav2-FLAG, Vma5-GFP is at higher levels in

the vacuoles of cells with ≥5 bud scars. The quantitation and binning by bud scar number in

Figure 6d confirms that there is no significant decline in vacuolar Vma5-GFP in cells with 5-9

bud scars and only in cells with ≥ 15 bud scars does vacuolar fluorescence reach the level in

"old" wild-type cells (Figure 1).   These data indicate that increasing Rav2-FLAG levels can help

preserve V-ATPase assembly in older cells.

We hypothesized that restoration of V-ATPase assembly in older cells grown under CR

conditions (Figure 4) might be supported by restoration of Rav2 levels. To test this, we isolated

young and old populations from Rav2-myc13 tagged cells grown under CR conditions. Under

these conditions, V-ATPase assembly and function is restored in older cells (Figure 4) and Rav2

levels are also restored (Figure 6e).

Alterations in Oxr1 levels with age could also affect V-ATPase assembly. We isolated

young and old populations from a strain containing HA-tagged Oxr1 and found that the levels of

Oxr1-HA in whole cell lysates do not change with age (Figure 6f). To further explore the basis of

the extended RLS in *oxr1Δ* cells, we observed Vma5-GFP localization in a population of *oxr1Δ*

cells of mixed age. As shown in Figure 6g, h, *oxr1Δ* mutants localize Vma5-GFP to the vacuole

even in older cells with ≥5 bud scars. To quantitate this effect, we again binned cells by the

number of bud scars as described above, and compared Vma5-GFP localization to localization in

young daughter cells (Figure 6h). In contrast to wild-type cells, there is no significant difference

in Vma5-GFP localization in the *oxr1Δ* mutant until cells have divided 25 or more times.

Intriguingly, these results suggest that the *oxr1Δ* mutation mirrors the effects of CR on V-

ATPase assembly and longevity. The results also indicate that direct manipulation of V-ATPase

assembly by mutation of critical assembly factors can affect replicative and chronological

lifespan, both of which are linked to cellular pH homeostasis (Murakami et al., 2012).

Specifically, promoting V-ATPase assembly through the RAVE complex appears to be important

for preserving lifespan, while the opposing effects of Oxr1 on the V-ATPase tend to shorten

lifespan.

DISCUSSION

These results establish V-ATPase disassembly as an important factor in reduced vacuolar

acidification in aging cells. We demonstrate an increase in V-ATPase disassembly at relatively

early replicative ages (5-9 cell divisions) similar to the age at which Hughes and Gottschling first

observed compromised vacuolar acidification (A. L. Hughes & Gottschling, 2012). We show that

promoting assembly of the V-ATPase through CR can restore V-ATPase assembly and vacuolar

pH in older cells. Deletion of Oxr1, a negative regulator of V-ATPase assembly, can extend both

replicative and chronological lifespan. In contrast, mutation of the RAVE complex shortens

lifespan and levels of the Rav2 subunit decline with age. Taken together, these data support V-

ATPase disassembly as a significant age-related factor behind reduced vacuolar and lysosomal

acidification and the associated declines in function.

This mechanism does not necessarily conflict with those proposed previously. If V-

ATPases are more disassembled in older cells, cells may be less able to tolerate an imbalance

between Pma1 activity at the plasma membrane and V-ATPase activity at the vacuole

(Henderson et al., 2014). Okreglak et al. (Okreglak et al., 2023) recently visualized oscillations

of vacuolar pH in mother cells that were coordinated with cell cycle. Oscillations cease upon V-

ATPase inhibition, but it is not clear whether reversible disassembly of the V-ATPase is required.

It is possible that this oscillatory behavior is lost in older mother cells, leading to loss of their

ability to restore vacuolar pH after cell division. Notably, we see no difference in protein levels

of V_1 subunits Vma1, Vma2, and Vma5, between young and old cells. The similar levels of

Vph1 in vacuoles of young and old cells suggest that V_0 subunits are present at similar levels,

since V_0 assembly occurs in the ER and reduced levels of any V_0 subunit reduces V_0 subcomplex

levels at the vacuole. The data presented here suggest a post-transcriptional regulatory

mechanism, but increased levels of a biosynthetic assembly factor (A. L. Hughes & Gottschling,

2012) could promote assembly and improve acidification.

Many questions remain to be investigated. The reduction in Rav2 protein levels with age

appears to be post-transcriptional but we do not yet know whether this arises from reduced

translation or increased degradation. It is intriguing in young cells, addition of an extra copy of

Rav2-FLAG did not increase levels of the protein in Figure 6b, suggesting the Rav2 subunit

levels are controlled. Perhaps more importantly, even though reduced levels of Rav2 levels helps

explain increased V-ATPase disassembly and addition of extra Rav2 improves assembly, we do

not know whether the Rav2 reduction occurs as a result of some general vulnerability in aging

cells or is deliberately programmed (Gladyshev et al., 2021). The significant increase in lifespan

and prolonged V-ATPase assembly in *oxr1Δ* cells seems to argue against any growth advantage

from increased V-ATPase disassembly in older cells, at least at the single cell level.

Many aging pathways are tightly linked to nutritional signaling, and reversible

disassembly is often driven by nutritional signaling pathways. Given the rapid disassembly of the

V-ATPase upon acute glucose deprivation (Kane, 1995), it was initially surprising that growth

under CR conditions suppresses V-ATPase disassembly with age. However, we previously

showed that the acute disassembly response required glucose concentrations well below 0.5%

(Parra & Kane, 1998). Activation of the PKA pathway in the *ira2Δ* mutant (Supporting

Information Figure 1b) promotes V-ATPase assembly in older cells, but also results in a decrease

in RLS. In mammalian cells, V-ATPase assembly generally increases in response to nutrient

deprivation or inhibition of mTOR function (Ratto et al., 2022). CR could mimic this effect in

aging cells, preserving lysosomal acidification and function. Deletion of the mTOR effector

SCH9 extends RLS and CLS and promotes V-ATPase assembly (Wilms et al., 2017), similar to

the effect of *oxr1Δ* mutation.   Signals involved in reversible disassembly of the V-ATPase are

incompletely understood and vary between physiological settings (Collins & Forgac, 2020). The

RAVE complex responds to nutritional signals associated with reversible disassembly with

release from the vacuolar membrane upon acute glucose deprivation and recruitment to the

membrane upon glucose restoration, even in the absence of V_1 subunit C and the V_1 subcomplex

(Jaskolka, Winkley, et al., 2021). The molecular mechanism of Oxr1-induced V-ATPase

assembly is understood (Khan & Wilkens, 2024), but its response to cellular signaling is still

being explored.

Reduced V-ATPase assembly could be a factor in the age-related decline in lysosomal

acidification and function in higher eukaryotes. Reversible disassembly occurs in higher

eukaryotes including mammalian cells. Rabconnectin-3 complexes of higher eukaryotes are

functional homologues of the yeast RAVE complex (Jaskolka, Winkley, et al., 2021). Oxr1

belongs to a family of TLDc proteins and several of these proteins have been shown to bind to

mammalian V-ATPases and induce disassembly (Oot & Wilkens, 2024). These data suggest that

the core elements for controlled V-ATPase disassembly during aging are present in other cells.

We observed V-ATPase disassembly in a yeast replicative aging model, which is most

comparable to mammalian cell types that continue to divide, such as adult stem cells (He et al.,

2018), and observed similar effects of *rav1Δ* and *oxr1Δ* mutations in chronological aging, which

is more analogous to long-lived, non-dividing mammalian cells. Taken together, these data

suggest that V-ATPase assembly state is linked to different types of aging and could easily play a

role in aging in higher eukaryotes.

METHODS

Yeast strains and plasmids

All strains analyzed were in the BY4741 (*MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*) or BY4742

(*MATα his3Δ1 leu2Δ0 lys1Δ ura3Δ0*) background. BY4742 strains containing Vma5-GFP::*HIS3*

and Vph1-GFP::*HIS3* were constructed as part of the genome-wide GFP-tagging project (Huh et

al., 2003). The *rav1Δ::kanMX, oxr1Δ::kanMX, vma1Δ::kanMX*, and *ira2Δ::kanMX* mutant

strains in the BY4741 background were purchased as part of a haploid deletion collection.

VMA2 was C-terminally tagged with GFP by PCR amplification from pFA6a-GFP-KanMX6 and

genomic integration (Longtine et al., 1998). The *oxr1Δ::kanMX* Vma5-GFP::*HIS3* strain was

obtained by crossing BY4741 *oxr1Δ::kanMX* and BY4742 Vma5-GFP::*HIS3* and obtaining

spores with the desired genotype by tetrad dissection. The *stv1Δ::natMX* mutation was

introduced by crossing this strain to a *stv1Δ::natMX* haploid and sporulating. The pRS316 Oxr1-

HA plasmid (Khan & Wilkens, 2024) was transformed into *oxr1Δ::kanMX* and transformants

were selected on medium lacking uracil. Rav1-myc13::*kanMX*, Rav2-myc13::*kanMX,* and Rav2-

FLAG::*kanMX* strains were constructed as described previously (Jaskolka, Tarsio, et al., 2021;

Smardon et al., 2002). The tagged alleles were PCR amplified from the original strains and

integrated into BY4741.

Fluorescence microscopy

Strains expressing GFP-tagged Vma5, Vma2, or Vph1 were grown to log phase in

synthetic complete (SC) medium containing either 2% or 0.5% glucose overnight, pelleted by

centrifugation, then suspended in fresh medium containing 2% or 0.5% glucose and grown for an

additional 2 hours. Cells were stained by addition of CW (final concentration 10 μg/ml) 5 min

prior to imaging. Cells were visualized with a 100x oil (NA 1.4) objective on a Zeiss Imager.Z1

fluorescence microscope with a Hamamastu CCD camera and AxioVision software. Cells were

viewed through DIC optics and fluorescence was imaged using a DAPI filter set (for CW) and a

GFP filter set. Bud scars were counted from CW staining; in mixed age populations, cells with 5

or more bud scars were designated as old and those with less than 5 as young. To obtain a more

precise count of bud scars for binning, cells were visualized on multiple focal planes. GFP

fluorescence images were captured for cells in each binned range then processed in FIJI. To

assess vacuolar localization, a line was drawn across a cell and through the vacuolar membrane.

From this line scan, a plot illustrating fluorescence intensity along the line was generated, with

peaks indicating areas of elevated fluorescence at the vacuolar membrane. The peak intensity

(maximum fluorescence) was quantitated as a measure of vacuolar localization. Each biological

replicate corresponds to a distinct culture of yeast cells. Maximum fluorescence was quantitated

for at least 20 young and 20 old cells per biological replicate. In cells were binned by age, at

least 20 cells per biological replicate were counted for each bin. Maximum fluorescence from

line scans of young and old cells in each biological replicate were averaged and then divided by

the average for young cells in that replicate to normalize. Normalized fluorescence for each

biological replicate was plotted, along with average intensities across replicates +/- s.e.m.

(standard error of the mean). Statistical significance was determined by t-test for the young and

old cell comparisons and by ANOVA for binned samples. To show the range of values for the

young cells across experiments, the values for young cells across biological replicates were

averaged; individual replicates were divided by this average are shown as points on each graph.

Age Enrichment

Age enrichment was performed as described by Jin et al. (Jin et al., 2021). Cells were collected

from a 50 mL fresh overnight culture in YEP (yeast extract-peptone medium) supplemented with

2% or 0.5% glucose to an OD_{600} of 1.0 and washed twice with cold phosphate-buffered saline

(PBS), pH 7.4. Cells were pelleted by centrifugation and washed three times in cold sterile PBS,

then labeled with 1.6 mg/ml EZ-Link Sulfo-NHS-LC-Biotin (Pierce) at room temperature for 30

min with gentle agitation. After labeling, the cells were washed three times with cold PBS, pH

8.0, to remove free biotin, and resuspended in YEP medium supplemented with 2% or 0.5%

glucose for growth overnight. After 16 hours, cells were pelleted by centrifugation and

resuspended in 35 ml of cold PBS, pH 7.4, mixed with 250 μl of a magnetic streptavidin bead

suspension (Pierce), and incubated for 60 min at 4°C. The mixture, in a 50 mL conical tube, was

loaded onto a magnetic separation column (Permagen) at 4°C to separate biotinylated cells.

Unbound (young) cells were removed gently by pipetting. Magnetically separated cells were

subsequently washed three times by resuspending in 35 mL PBS, pH 7.4 supplemented with 2%

or 0.5% glucose, repeating magnetic separation, and discarding the supernatant. After washing,

cells were resuspended in 200 mL of YEP or SC containing 2% or 0.5% glucose and allowed to

grow for an additional 4 hours before obtaining the final "old" mother cells and "young"

daughter cells. Cells were loaded on the magnetic separation column as described above and

daughter cells obtained from the supernatant were pelleted by centrifugation to obtain a

concentrated population of young cells. After the final wash, the magnetic beads were

resuspended in 1 ml PBS, pH 7.4 and centrifuged to concentrate the old population. The two

populations were then used immediately for pH measurements or stored at -80°C for further

biochemical analysis.

Whole Cell Lysates and Immunoblots

Cell pellets from age-enriched young and old cell populations were resuspended in

cracking buffer (8 M urea, 5% SDS, 1 mM EDTA, 50 mM Tris-HCl, pH 6.8) at 95°C and lysed

with glass beads. Cellular debris was pelleted by centrifugation at $16,000 \times g$ for 2 min, and

supernatants containing whole-cell extracts were used immediately or stored at -80 °C until use.

After determination of protein concentrations by Bradford assay, equal amounts of

protein were separated by SDS-PAGE and transferred to a nitrocellulose membrane. Blots were

blocked, then incubated overnight with primary antibodies at 4°C. Primary antibodies (all at a

1:500 dilution) included mouse monoclonal antibodies 7A2 against Vma5, 10D7 against Vph1,

8B1 against Vma1, and 13D11 against Vma2 (Kane, Kuehn, Howald-Stevenson, & Stevens,

1992). In addition, anti-myc monoclonal 9E10 (Santa Cruz Biotechnology), anti-HA monoclonal

(BioLegend), anti-FLAG (Sigma) and anti-GAPDH (Proteintech) antibodies were used at 1:500,

1:500, 1:1000, and 1:10,000 dilution, respectively. HRP-conjugated anti-mouse secondary

antibody (Bio-Rad) was added at a final dilution of 1:2000 and incubated for 60 min at room

temperature. The blot was washed, incubated with Bio-Rad Clarity Western ECL substrate and

imaged in an Azure Sapphire FL Biomolecular Imager. Images were quantified using FIJI.

Molecular mass markers were included on every blot. In blot images, the nearest molecular mass

marker is indicated.

Vacuolar pH Measurements

Vacuolar pH was measured using BCECF-AM (Invitrogen) as described previously (Diakov et

al., 2013). Age-enriched populations of cells were loaded with BCECF-AM in YEP

supplemented with 2% or 0.5% glucose. After washing with YEP media to remove dye, cells

were resuspended in YEP, deprived of glucose, and incubated for 30 min. For fluorescence

measurement, 20 µl of cell suspension was diluted into 3 ml 1 mM 2-(N-

morpholino)ethanesulfonic acid (MES) pH 5.0 buffer, and fluorescence intensity at excitation

wavelengths 450 and 495 nm and emission wavelength 535 nm and was monitored continuously

in a Horiba Jovin Yvon Spectrafluor Max fluorometer at 30°C. The fluorescence ratio for each

sample was calibrated in every experiment by clamping pH in the range of 5.0 to 7.0 and

constructing a calibration curve.

Replicative Life Span

RLS was determined on YEP, 2% glucose plates buffered to pH 5. Daughter cells were

sequentially removed by micromanipulation (Steffen, Kennedy, & Kaeberlein, 2009). Survival

curves are pooled data from experiment-matched controls. The number of divisions for 26-30

mother cells were scored for each curve. p-values for replicative life span survival curve

comparisons were calculated with a Cox regression model. Kaplan-Meier survival curves were

plotted with GraphPad Prism.

Chronological lifespan analysis

CLS was assessed as described (Wilms et al., 2017), with the following modifications. Cells

were pre-cultured to the stationary phase in SC containing 2% glucose. Stationary-phase cells

were then diluted to an OD_{600} of 0.1 in fresh medium and grown for 48 hours; the end of this

time is designated as time zero. Cells were aged in SC containing 2% glucose adjusted to pH 5.5

with 500 mM MES. Cell viability and clonogenicity (colony forming units; CFUs) were

analyzed on days 7, 14, 21, 28, and 35. Cell death was assessed by propidium iodide staining

(Sigma) and visualized by fluorescence microscopy. For CFU analysis, cell density was

measured using a hemocytometer, and 250 cells were plated onto YPD plates. After incubation,

CFUs were counted using FIJI, and the results expressed as the percentage of total cells plated.

Statistical significance was assessed by ANOVA with multiple sequence comparisons (GraphPad

Prism).

RT-PCR of RAV2 from young and old cells

RNA was extracted from young and old cells, obtained by age enrichment as described above,

using the NEB Monarch Total RNA Miniprep Kit. RT-PCR was conducted using the NEB Luna

Universal One-Step RT-qPCR Kit and performed on a Bio-Rad CFX384 Touch System. Data

analysis was conducted using CFX Maestro Software.

Acknowledgements: This work was supported by NIH R35 GM145256 to P.M.K. The authors

thank M. Murad Khan and Dr. Stephan Wilkens for sharing the *oxr1Δ* strain and Oxr1-HA

plasmid, and Dr. Xin Jie Chen for helpful discussions and a critical reading of the manuscript.

Illustrations in Figures 2a and 5a were prepared using Biorender.com.

Author contributions: F.H. performed experiments, analyzed data, prepared figures, and wrote

the first draft of the manuscript; P.M.K. obtained funding for the project, analyzed data, prepared

figures, and contributed to writing of the manuscript.

Conflict of interest: The authors declare that they have no conflicts of interest with the contents

of this article.

Data availability: The data that support the findings of this study are available at

Upstate.figshare.com. https://upstate.figshare.com. DOI: 10.58120/upstate.27902313

FIGURE LEGENDS

Figure 1: V-ATPases are more disassembled in yeast cells of older replicative age. (a) Vma5-

GFP cells grown in SC containing 2% glucose. DIC was used to visualize vacuoles. Bud scars

are stained with CW to determine replicative age. **(b)** Left panel provides representative example

of quantitative measurements using line scans. The young and old cells are from the dashed

white and red boxes, respectively, in the Vma5-GFP image of **Figure 1a**. Plot profiles are

superimposed for the young (black) and old (red) cells. **(c)** Quantitation of maximum fluorescent

intensity across five biological replicates, after normalization to the average intensity in the

youngest bin for each replicate. Each biological replicate (dot) represents at least 20 cells and

bars represent the mean +/- s.e.m. "Old" for subsequent experiments (without age enrichment) is

categorized as ≥5 bud scars. **** represents a p-value <0.0001 calculated by ordinary one-way

ANOVA.

Figure 2: Age-enriched populations of cells have comparable levels of V-ATPase subunits.

(a) Diagram of V-ATPase showing the positions of Vma5, Vma2, and Vph1. **(b)** Vma2-GFP cells

(V_1B) grown in SC with 2% glucose. CW was used to visualize bud scars. The CW images for

young cells were overexposed relative to those for the old cells in order to visualize the low level

staining in young cells. Normalized maximum fluorescence was obtained as in **Figure 1**. Means

+/- s.e.m. of three biological replicates are shown; each replicate is represented by a dot.

Significance was assessed by unpaired Student's *t* test, *** p value=0.0009 . **(c)** Vph1-GFP cells

were visualized and analyzed as described in **2b**. **(d)** After biotin-streptavidin age enrichment,

bud scars were counted for 100 cells in young and old populations, then binned by bud count. **(e)**

Lysates from age-enriched populations defined as in **1c** were separated by SDS-PAGE.

Immunoblots for the indicated proteins in young (Y) vs. old (O) cell populations as defined as in

1c. **(f)** Band intensities were quantified using FIJI, ratios of subunit levels to the GAPDH control

were calculated and normalized to the young population for each biological replicate. Data are

presented as mean (horizontal bars) ± s.e.m. (whiskers) of three biological replicates. n.s.= not

significant by unpaired Student's *t* test.

Figure 3: Vacuolar pH is more alkaline in old cells. (a) Vacuolar pH responses were measured

for wild-type BY4742 age-enriched young and old populations as described in Methods.

Glucose-deprived cultures were loaded with BCECF-AM. Fluorescence intensity values were

collected every 10 sec and glucose was added to a final concentration of 2% after 3 min. The

ratio of fluorescence signals was converted to pH as described in Methods. **(b)** pH measurements

at 1 min (before glucose addition), 5 min (2 min after glucose addition), and 8 min (5 min after

addition) are presented as mean (horizontal bars) ± s.e.m. (whiskers) of three biological

replicates. * = p value of 0.02, ** p value <0.001.

Figure 4: Caloric restriction restores V-ATPase assembly and vacuolar pH in old cells. (a-c)

Strains used in **Figures 1** and **2** show recruitment of Vma5-GFP (**a**) and Vma2-GFP **(b)** to the

vacuolar membrane after growth in SC with 0.5% glucose. Vph1-GFP (**c**) remains unchanged.

Normalized maximum fluorescence was quantitated as described in Methods. No significant

differences between young and old cells were observed by unpaired Student's *t* test. Designation

of young and old cells by CW staining was as described in **Figure 1**. **(d)** Young and old cell

populations were obtained by biotin-streptavidin age-enrichment from cells grown in YEP

supplemented with 0.5% glucose, and vacuolar pH responses were measured. Data were

collected and analyzed as in **Figure 3b**. Data are presented as mean (horizontal bars) ± s.e.m.

(whiskers) of three biological replicates.

Figure 5: Effects of *rav1Δ* and *oxr1Δ* on RLS and CLS. (a) Schematic of reversible

disassembly. **(b)** Kaplan-Meier curves comparing the RLS of *rav1Δ* (green), *oxr1Δ* (red), and

wild-type cells (black). Median number of replicative generations is shown in parentheses, with

the % difference between median values for *rav1Δ* and *oxr1Δ* vs. wild-type. Deletion of *OXR1*

increases RLS (p<0.001), and deletion of RAVE component *RAV1* shortens RLS (p < 0.01). **(c,**

d) Plots of colony-forming units **(c)** and percentage of cells stained with propidium iodide **(d)** vs.

time. Each point is the mean of three biological replicates. Strains included are defined to the

right of **(c)**. Statistical analysis of **(c)** indicates a p-value <0.0001 for all mutants in comparison

to wild-type. In **(d)**, *vma1Δ* and *oxr1Δ* differed from wild-type with a p-value <0.0001, and

rav1Δ with a p-value= 0.0007.

Figure 6: RAVE subunit and Oxr1 levels in aging cells. (a) Rav1 levels in young and old cells.

Age-enriched populations were obtained from WT cells containing Rav1-myc$_{13}$ (young (Y)

versus old (O) cells). Cell lysates were prepared and analyzed as in **Figure 2**. Normalized data

are presented as mean (horizontal bars) ± s.e.m. (whiskers) of three biological replicates. ns=not

significant. **(b)** Rav2-FLAG levels from young and old cells prepared as in **6a** from haploid cells

containing only a chromosomally integrated (int.) Rav2-FLAG or cells with both integrated

Rav2-FLAG and low copy Rav2-FLAG plasmid (+plasmid). Normalized data are presented as

mean (horizontal bars) ± s.e.m. (whiskers) of three biological replicates. ****, p-value<0.0001,

ns= not significant. **(c)** Rav2-FLAG Vma5-GFP cells, without (int.) or with the Rav2-FLAG

plasmid (+ plasmid). CW is used to visualize bud scars as in **Figure 1**; white stars on the Vma5-

GFP images indicate newborn cells with no bud scars. Images representative of cells with 0-4,

5-9, and 10-14 bud scars are shown. **(d)** Rav2-FLAG cells + plasmid were binned by the number

of bud scars and normalized maximal fluorescence quantitated as in **Figure 1**. **, p-value<0.01

and *, p-value<0.05 for the indicated comparisons; n.s.=not significant. **(e)** Immunoblot analysis

comparing Rav2-myc$_{13}$ levels in cells grown in YEP with 2% glucose or 0.5% glucose (CR

conditions). Samples were prepared, analyzed, and presented as in **6a**. ***, p- value <0.001,

n.s.= not significant. **(f)** Immunoblot of lysates from *oxr1Δ* strain expressing Oxr1-HA from a

low copy plasmid. Young and old cells were isolated as in **6a**. Data are presented as mean

(horizontal bars) ± s.e.m. (whiskers) of three biological replicates. **(g)** Wild-type and *oxr1Δ* cells

containing Vma5-GFP grown in SC with 2% glucose. DIC and CW were used to visualize

vacuoles and bud scars as in **Figure 2b**. **(h)** *oxr1Δ* cells were binned by the number of bud scars

and normalized maximal fluorescence quantitated as in **Figure 1**. ***, p-value <0.001 in

comparison to the youngest bin; there is no other significant difference.

REFERENCES

Bodzeta, A., Kahms, M., & Klingauf, J. (2017). The Presynaptic v-ATPase Reversibly Disassembles and Thereby Modulates Exocytosis but Is Not Part of the Fusion Machinery. *Cell Rep, 20*(6), 1348-1359. doi:10.1016/j.celrep.2017.07.040

Bond, S., & Forgac, M. (2008). The Ras/cAMP/protein kinase A pathway regulates glucose-dependent assembly of the vacuolar (H+)-ATPase in yeast. *J Biol Chem, 283*(52), 36513-36521. doi:M805232200 [pii]10.1074/jbc.M805232200

Chen, K. L., Ven, T. N., Crane, M. M., Brunner, M. L. C., Pun, A. K., Helget, K. L., . . . Wasko, B. M. (2020). Loss of vacuolar acidity results in iron-sulfur cluster defects and divergent homeostatic responses during aging in Saccharomyces cerevisiae. *Geroscience, 42*(2), 749-764. doi:10.1007/s11357-020-00159-3

Collins, M. P., & Forgac, M. (2020). Regulation and function of V-ATPases in physiology and disease. *Biochim Biophys Acta Biomembr, 1862*(12), 183341. doi:10.1016/j.bbamem.2020.183341

Devare, M. N., Kim, Y. H., Jung, J., Kang, W. K., Kwon, K. S., & Kim, J. Y. (2020). TORC1 signaling regulates cytoplasmic pH through Sir2 in yeast. *Aging Cell, 19*(6), e13151. doi:10.1111/acel.13151

Diab, H. I., & Kane, P. M. (2013). Loss of vacuolar H+-ATPase (V-ATPase) activity in yeast generates an iron deprivation signal that is moderated by induction of the peroxiredoxin TSA2. *Journal of Biological Chemistry, 288*(16), 11366-11377. doi:M112.419259 [pii] 10.1074/jbc.M112.419259

Diakov, T. T., Tarsio, M., & Kane, P. M. (2013). Measurement of vacuolar and cytosolic pH in vivo in yeast cell suspensions. *J Vis Exp, Apr 19*(74). doi:10.3791/50261

Gladyshev, V. N., Kritchevsky, S. B., Clarke, S. G., Cuervo, A. M., Fiehn, O., de Magalhaes, J. P., . . . Cummings, S. R. (2021). Molecular Damage in Aging. *Nat Aging, 1*(12), 1096-1106. doi:10.1038/s43587-021-00150-3

Hansen, M., Rubinsztein, D. C., & Walker, D. W. (2018). Autophagy as a promoter of longevity: insights from model organisms. *Nat Rev Mol Cell Biol, 19*(9), 579-593. doi:10.1038/s41580-018-0033-y

He, C., Zhou, C., & Kennedy, B. K. (2018). The yeast replicative aging model. *Biochim Biophys Acta Mol Basis Dis, 1864*(9 Pt A), 2690-2696. doi:10.1016/j.bbadis.2018.02.023

Henderson, K. A., Hughes, A. L., & Gottschling, D. E. (2014). Mother-daughter asymmetry of pH underlies aging and rejuvenation in yeast. *Elife, 3*, e03504. doi:10.7554/eLife.03504

Hughes, A. L., & Gottschling, D. E. (2012). An early age increase in vacuolar pH limits mitochondrial function and lifespan in yeast. *Nature, 492*(7428), 261-265. doi:nature11654 [pii]10.1038/nature11654

Hughes, C. E., Coody, T. K., Jeong, M. Y., Berg, J. A., Winge, D. R., & Hughes, A. L. (2020). Cysteine Toxicity Drives Age-Related Mitochondrial Decline by Altering Iron Homeostasis. *Cell, 180*(2), 296-310 e218. doi:10.1016/j.cell.2019.12.035

Huh, W. K., Falvo, J. V., Gerke, L. C., Carroll, A. S., Howson, R. W., Weissman, J. S., & O'Shea, E. K. (2003). Global analysis of protein localization in budding yeast. *Nature, 425*(6959), 686-691.

Jaskolka, M. C., Tarsio, M., Smardon, A. M., Khan, M. M., & Kane, P. M. (2021). Defining steps in RAVE-catalyzed V-ATPase assembly using purified RAVE and V-ATPase subcomplexes. *J Biol Chem, 296*, 100703. doi:10.1016/j.jbc.2021.100703

Jaskolka, M. C., Winkley, S. R., & Kane, P. M. (2021). RAVE and Rabconnectin-3 Complexes as Signal Dependent Regulators of Organelle Acidification. *Front Cell Dev Biol, 9*, 698190. doi:10.3389/fcell.2021.698190

Jin, X., Cao, X., & Liu, B. (2021). Isolation of Aged Yeast Cells Using Biotin-Streptavidin Affinity Purification. *Methods Mol Biol, 2196*, 223-228. doi:10.1007/978-1-0716-0868-5_17

Kane, P. M. (1995). Disassembly and reassembly of the yeast vacuolar H(+)-ATPase in vivo. *J Biol Chem, 270*(28), 17025-17032.

Kane, P. M., Kuehn, M. C., Howald-Stevenson, I., & Stevens, T. H. (1992). Assembly and targeting of peripheral and integral membrane subunits of the yeast vacuolar H(+)-ATPase. *J Biol Chem, 267*(1), 447-454.

Kaushik, S., Tasset, I., Arias, E., Pampliega, O., Wong, E., Martinez-Vicente, M., & Cuervo, A. M. (2021). Autophagy and the hallmarks of aging. *Ageing Res Rev, 72*, 101468. doi:10.1016/j.arr.2021.101468

Khan, M. M., Lee, S., Couoh-Cardel, S., Oot, R. A., Kim, H., Wilkens, S., & Roh, S. H. (2022). Oxidative stress protein Oxr1 promotes V-ATPase holoenzyme disassembly in catalytic activity-independent manner. *EMBO J, 41*(3), e109360. doi:10.15252/embj.2021109360

Khan, M. M., & Wilkens, S. (2024). Molecular mechanism of Oxr1p mediated disassembly of yeast V-ATPase. *EMBO Rep*. doi:10.1038/s44319-024-00126-5

Klossel, S., Zhu, Y., Amado, L., Bisinski, D. D., Ruta, J., Liu, F., & Gonzalez Montoro, A. (2024). Yeast TLDc domain proteins regulate assembly state and subcellular localization of the V-ATPase. *EMBO J*. doi:10.1038/s44318-024-00097-2

Kohio, H. P., & Adamson, A. L. (2013). Glycolytic control of vacuolar-type ATPase activity: a mechanism to regulate influenza viral infection. *Virology, 444*(1-2), 301-309. doi:S0042-6822(13)00396-6 [pii]

10.1016/j.virol.2013.06.026

Liu, Y., Steinbusch, L. K. M., Nabben, M., Kapsokalyvas, D., van Zandvoort, M., Schonleitner, P., . . . Luiken, J. (2017). Palmitate-Induced Vacuolar-Type H(+)-ATPase Inhibition Feeds Forward Into Insulin Resistance and Contractile Dysfunction. *Diabetes, 66*(6), 1521-1534. doi:10.2337/db16-0727

Longo, V. D., & Anderson, R. M. (2022). Nutrition, longevity and disease: From molecular mechanisms to interventions. *Cell, 185*(9), 1455-1470. doi:10.1016/j.cell.2022.04.002

Longtine, M. S., McKenzie, A., 3rd, Demarini, D. J., Shah, N. G., Wach, A., Brachat, A., . . . Pringle, J. R. (1998). Additional modules for versatile and economical PCR-based gene deletion and modification in Saccharomyces cerevisiae. *Yeast, 14*(10), 953-961.

Martinez-Munoz, G. A., & Kane, P. (2008). Vacuolar and plasma membrane proton pumps collaborate to achieve cytosolic pH homeostasis in yeast. *J Biol Chem, 283*(29), 20309-20319. doi:M710470200 [pii]

10.1074/jbc.M710470200

Murakami, C., Delaney, J. R., Chou, A., Carr, D., Schleit, J., Sutphin, G. L., . . . Kaeberlein, M. (2012). pH neutralization protects against reduction in replicative lifespan following chronological aging in yeast. *Cell Cycle, 11*(16), 3087-3096. doi:10.4161/cc.21465

Nixon, R. A. (2020). The aging lysosome: An essential catalyst for late-onset neurodegenerative diseases. *Biochim Biophys Acta Proteins Proteom, 1868*(9), 140443. doi:10.1016/j.bbapap.2020.140443

Okreglak, V., Ling, R., Ingaramo, M., Thayer, N. H., Millett-Sikking, A., & Gottschling, D. E. (2023). Cell cycle-linked vacuolar pH dynamics regulate amino acid homeostasis and cell growth. *Nat Metab, 5*(10), 1803-1819. doi:10.1038/s42255-023-00872-1

Oot, R. A., & Wilkens, S. (2020). A "Sugar-Coated" Proton Pump Comes into Focus: High-Resolution Structure of a Human V-ATPase. *Mol Cell, 80*(3), 379-380. doi:10.1016/j.molcel.2020.10.020

Oot, R. A., & Wilkens, S. (2024). Human V-ATPase function is positively and negatively regulated by TLDc proteins. *Structure, 32*(7), 989-1000 e1006. doi:10.1016/j.str.2024.03.009

Parra, K. J., & Kane, P. M. (1998). Reversible association between the V1 and V0 domains of yeast vacuolar H+-ATPase is an unconventional glucose-induced effect. *Mol Cell Biol, 18*(12), 7064-7074. Retrieved from http://mcb.asm.org/cgi/content/full/18/12/7064

Perera, R. M., & Zoncu, R. (2016). The Lysosome as a Regulatory Hub. *Annu Rev Cell Dev Biol, 32*, 223-253. doi:10.1146/annurev-cellbio-111315-125125

Ratto, E., Chowdhury, S. R., Siefert, N. S., Schneider, M., Wittmann, M., Helm, D., & Palm, W. (2022). Direct control of lysosomal catabolic activity by mTORC1 through regulation of V-ATPase assembly. *Nat Commun, 13*(1), 4848. doi:10.1038/s41467-022-32515-6

Schiffer, I., Gerisch, B., Kawamura, K., Laboy, R., Hewitt, J., Denzel, M. S., . . . Antebi, A. (2021). miR-1 coordinately regulates lysosomal v-ATPase and biogenesis to impact proteotoxicity and muscle function during aging. *Elife, 10*. doi:10.7554/eLife.66768

Seol, J. H., Shevchenko, A., & Deshaies, R. J. (2001). Skp1 forms multiple protein complexes, including RAVE, a regulator of V-ATPase assembly. *Nat Cell Biol, 3*(4), 384-391.

Smardon, A. M., Diab, H. I., Tarsio, M., Diakov, T. T., Nasab, N. D., West, R. W., & Kane, P. M. (2014). The RAVE complex is an isoform-specific V-ATPase assembly factor in yeast. *Mol Biol Cell, 25*(3), 356-367. doi:mbc.E13-05-0231 [pii]
10.1091/mbc.E13-05-0231

Smardon, A. M., Tarsio, M., & Kane, P. M. (2002). The RAVE complex is essential for stable assembly of the yeast V-ATPase. *J Biol Chem, 277*(16), 13831-13839.

Steffen, K. K., Kennedy, B. K., & Kaeberlein, M. (2009). Measuring replicative life span in the budding yeast. *J Vis Exp*(28). doi:10.3791/1209

Sumner, J. P., Dow, J. A., Earley, F. G., Klein, U., Jager, D., & Wieczorek, H. (1995). Regulation of plasma membrane V-ATPase activity by dissociation of peripheral subunits. *J Biol Chem, 270*(10), 5649-5653.

Tabke, K., Albertmelcher, A., Vitavska, O., Huss, M., Schmitz, H. P., & Wieczorek, H. (2014). Reversible disassembly of the yeast V-ATPase revisited under in vivo conditions. *Biochemical Journal, 462*(1), 185-197. doi:BJ20131293 [pii]
10.1042/BJ20131293

Vilchez, D., Saez, I., & Dillin, A. (2014). The role of protein clearance mechanisms in organismal ageing and age-related diseases. *Nat Commun, 5*, 5659. doi:10.1038/ncomms6659

Wilms, T., Swinnen, E., Eskes, E., Dolz-Edo, L., Uwineza, A., Van Essche, R., . . . Winderickx, J. (2017). The yeast protein kinase Sch9 adjusts V-ATPase assembly/disassembly to control pH homeostasis and longevity in response to glucose availability. *PLoS Genet, 13*(6), e1006835. doi:10.1371/journal.pgen.1006835

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6