Title: Reduced perceptual error monitoring is a biomarker of autism

Authors**:** Nathaniel J. Zuk[1,2*], Jaroslav Y. Slobodskoy-Plusnin[1], Yarden Weiss[1],

VishnuPriya Sampath[1], Kira M. Düsterwald[3], Athena Akrami[4§], Merav Ahissar[1§]

Affiliations: 1) Edmond & Lily Safra Center for Brain Sciences, Hebrew University,

Jerusalem, Israel; 2) NTU Psychology, Nottingham Trent University, Nottingham, United

Kingdom; 3) Gatsby Computational Neuroscience Unit, University College London,

London, United Kingdom; 4) Sainsbury Wellcome Centre, University College London,

London, United Kingdom

§These authors jointly supervised this work

Corresponding author: Nathaniel J. Zuk, NTU Psychology, Nottingham Trent University,

Nottingham, United Kingdom (nathaniel.zuk@ntu.ac.uk)

Abstract:

Can reduced flexibility in autistics' behavior be a reliable classifier of autism? We asked this question in the context of pitch discrimination, which is performed with similar accuracy by autistic and non-autistic participants, matched for cognitive skills. We found that three reliable, uncorrelated characteristics quantified reduced behavioral flexibility. Most non-autistic participants showed a clear feedback-related negativity (FRN) in their EEG when incorrect, larger EEG responses during feedback following difficult versus easy trials, and increased perceptual bias following correct and easy trials versus difficult ones. All three measures were weaker in autistic participants. To assess reliability, we replicated all three measures in an experiment approximately a year later. These measures were not correlated within participants, indicating that several mechanisms, associated with anterior cingulate cortex, are atypical in autism. Together, these three measures provide group classification with above 80% accuracy, showing promise for their utility as a neurocognitive biomarker of autism.

Significance statement: A characteristic of autism is cognitive "slowness" in updating expectations in new contexts. Our study shows that this cognitive difference is related to a reduced use of feedback, linked to atypical activity in a brain area involved in error detection and behavioral adjustment. Using EEG, we found that autistic individuals have weaker feedback-related signals and do not adjust their behavior as effectively as non-autistic individuals. Together, these neurocognitive signals could serve as a biomarker for autism. This is the first time such brain responses have been measured in a perceptual task requiring continuous adjustments. Understanding these brain mechanisms can inform strategies to enhance learning and adaptability in autism, providing new insights into the neural basis of behavioral flexibility.

Introduction

People with autism spectrum disorder (ASD) are known to be less flexible in updating their motor and perceptual performance to changes in the environment (Van de Cruys et al., 2014; Lieder et al., 2019). For example, when people perform a perceptual discrimination task they implicitly learn the distribution of the stimuli they experience and form predictions regarding the upcoming stimuli (Ashourian and Loewenstein, 2011). These predictions are updated on each trial (Akrami et al., 2018; Lieder et al., 2019), but in ASD the rate of this update is reduced (Lieder et al., 2019; Sapey-Triomphe et al., 2023; Noel et al., 2024). Other studies have demonstrated slower updating in tasks requiring continuous monitoring of errors (South et al., 2012; D'Cruz et al., 2016; Vishne et al., 2021; Kasten et al., 2023), which could also relate to the slower rate at which they update their internal representation of the external world.

A separate line of evidence using EEG suggests atypical impact of both external and internal feedback. Reduced sensitivity to internal feedback is manifested when participants detect that they have made an erred response in fast motor tasks (such as the flanker task; Gehring et al., 1993, 1995, 2018), which is smaller in autism (Vlamings et al., 2008; Sokhadze et al., 2010; South et al., 2010; Hüpen et al., 2016) (but see Henderson et al., 2006; Groen et al., 2008). Similarly, reduced sensitivity to external feedback, called the feedback-related negativity (FRN) is manifested as an error in the expected outcome (Miltner et al., 1997). The FRN was often studied in the context of success and failure in guessing random rewards (Gehring and Willoughby, 2002; Holroyd and Coles, 2002; Bellebaum and Daum, 2008; Larson et al., 2011) rather than in evaluating the outcome of reward-maximization strategies in perceptual decision making paradigms. In autism, such a study design has yielded mixed results (Groen et al., 2008; Larson et al., 2011; Bellebaum et al., 2014) (for review see Hüpen et al., 2016).

The anterior cingulate cortex (ACC) is important for evaluating both predictions and actions, integrating internal and external information (Falkenstein et al., 1991; Gehring et al., 1993; Dehaene et al., 1994; Carter et al., 1998; Botvinick et al., 1999, 2004; Beckmann et al., 2009; Stevens et al., 2011), including updating predictions (Behrens et al., 2007), although its specific role is not well understood. Indeed, ACC was found to be atypical in ASD (Thakkar et al., 2008; Sapey-Triomphe et al., 2023), in both connectivity (Simms et

al., 2009; Cauda et al., 2011; Postema et al., 2019) and in its pattern of activation (Chan et al., 2011) particularly when processing perceptual errors (Sapey-Triomphe et al., 2021, 2023). This is consistent with recent findings on under-weighting of sensory observation and prediction error in frontal cortices across several mouse models of autism (Noel et al., 2024).

Together, the above observations suggest that online monitoring of both motor and sensory performance is reduced in ASD. We now asked whether all these mechanisms manifest in a single task and whether their measures are reliably reduced in autism. We used a single, simple perceptual task and aimed to measure all three aspects of monitoring: perceptual updates, internal feedback, and external feedback. We found that all forms of performance monitoring are reliably reduced in ASD. Moreover, all these measures, while uncorrelated within participant groups, were replicated in a follow-up study 1-2 years later and were sufficient when combined to classify non-autistic and ASD participants with over 80% accuracy. These results show that reduced performance monitoring, which can be reliably measured behaviorally and neurophysiologically, provides a biomarker for autism.

Methods

All experimental methods were approved by the European Research Council and the Ethics Committee of the Hebrew University.

Study 1

Participants

35 non-autistic and 34 ASD individuals participated in this experiment. One ASD participant was rejected from further analyses because of poor performance which suggested that they did not understand the task, leaving 33 ASD included in our analyses. All autistic individuals had a prior formal diagnosis (by a psychologist, neurologist, or psychiatrist), and participation in our study did not require changes in medication. Except for one participant in each group, all participants had at most four years of musical training. The 35 non-autistics and 33 ASD participants that were included in subsequent analyses were also matched for age, gender, and reasoning scores, measured by Block Design, a standardized test of visuospatial reasoning (Wechsler, 2008) (see Table 1). The Block

Design scores (population average is scaled to 10 and standard deviation to 1.5) indicate high cognitive skills in both groups. By contrast, and as expected, there was a significant difference in the scores of the self-reported autism questionnaire, AQ50 (Autism Quotient questionnaire; (Baron-Cohen et al., 2001)), with an effect size of Cohen's $d = 1.10$.

Table 1: Demographic information for the participants included in the experiment. For age, Block Design, and AQ50 the mean is specified for each group, with the standard deviation in parentheses. ** $p < 0.01$ based on a Student's t-test.

	Non-autistics	ASD	t-statistic	p-value
Number of participants	35	33		
Number of females	5	6		
Age (years)	25.9 (3.7)	25.4 (5.3)	0.493	0.624
Block design (scaled score)	12.5 (2.2)	12.9 (3.4)	-0.384	0.702
AQ50**	50.6 (15.8)	74.2 (24.6)	-3.269	0.002

Experiment design

Experiments were coded using PsychToolbox in Matlab (Mathworks). In each trial, participants were sequentially presented with two tones 50 ms in duration with 5 ms onset and offset sine ramps. Tone frequencies ranged from 800 to 1250 Hz and were sampled so that both tones in a pair were either above or below 1000 Hz (as in Jaffe-Dax et al., 2017). The ratio of the two tone frequencies ranged from 1.01 to 1.25 (0.17 to 3.86 semitones). Tone onsets were separated by 2 seconds, resulting in a working memory delay (inter-stimulus-interval) of 1.95 seconds. After the second tone, participants had to specify which tone had a higher pitch. Upon response, participants were immediately presented with visual feedback for 400 ms: "Correct" in green if they got the correct answer, or "Incorrect" in red if they were wrong. Participants had 1300 ms to respond after the onset of the second tone, and if they did not respond within this interval, then they saw "Miss" in red on screen for 400 ms. This "time out" interval is not stressful, as the median response time across all

participants was 646 ms with a range of 398 ms to 923 ms. The next trial was always

presented four seconds after the first tone onset of the previous trial (see Figure 1A),

irrespective of the speed of the participant's response. The inter-trial interval (2 s after the

second tone) was set to be equal to the within-trial inter-stimulus interval (2 s) to reduce

the effect of within-trial adaptation, which might differ between the groups as it relates to

implicit learning (Jaffe-Dax et al., 2017; Regev et al., 2021). The sounds were presented

over Sennheiser HD 25 headphones at a comfortable sound level.

Before starting the formal experiment, participants completed a training set of 10-30

trials, presented in blocks of 10 trials with a break in between. To make the task easier,

during these training trials the frequency range of the tones was increased to 600 – 1600

Hz, and the tone ratios were randomly selected between 1.35 – 1.4 (equivalent to 5.20 –

5.83 semitones). Participants proceeded to the main experiment upon reaching 8/10 trials

correct in a block, or upon completing the three 10-trial training blocks, even if this

threshold performance was not attained (this was the case for 7/33 ASD participants and

5/35 non-autistics).   In the main experiment, participants performed 121 trials without

breaks.

Behavioral analysis

Recent trials biased the decision of participants. This bias can be viewed as a

contraction of the perceptual memory of the first tone in the trial (whose representation is

noisier by the time the second tone is presented) towards the expected frequency, namely

the geometric mean of the tone frequencies presented first in the previous trial.

Consequently, in trials where the first tone's frequency is closer than the second tone's

frequency to geometric mean of the previous trial performance improves (as representation

of the first tone becomes more distant from that of the second tone in the trial, "bias+"

trials). By contrast, in trials where the first tone is farther from the previous trial's

frequencies, the representation of the distance between the tones decreases, hampering

performance ("bias-" trials). Previous work has shown that people with autism are less

biased by the recent trial (Lieder et al., 2019).

Performance was assessed by calculating the proportion of correct trials, ignoring

misses. To quantify the effect of recent history on choice bias, we split trials into "bias+"

and "bias-" trials and calculated the difference in performance between these sets of trials.

To understand the effects of external feedback ("Correct" or "Incorrect" trials) and trial

difficulty on subsequent behavior, we further grouped trials based on the participant's

accuracy (Correct/Incorrect) and the trial difficulty (Easy/Hard trials). For trial difficulty,

easy versus hard trials were identified with respect to the median frequency difference

which was 1.242 semitones. Trials with smaller frequency difference were tagged "hard"

whereas trials with larger frequency difference were tagged "easy". We asked how implicit

strategies differed between trials that followed easy ones and trials that followed difficult

ones.

EEG recording and preprocessing

Biosemi ActiveTwo with a 32-channel layout was used to record EEG while

participants performed the task. Additionally, external channels were recorded bilaterally

on the mastoids, to the outside of each eye, above each eye, and on the nose (7 external

electrodes total). The EEG data was digitized at 512 Hz.

All preprocessing was done in Matlab. After recording, the EEG data was filtered

using a 4th order zero-phase Butterworth filter with -3 dB cutoff frequencies of 1-30 Hz.

Eyeblinks and eye motion artifacts were isolated using fastICA (Hyvärinen and Oja, 2000)

applied to the 32 EEG channels as well as the externals and then removed. The data was

re-referenced to the average of the mastoid channels. If one of the mastoid channel's

variance exceeded the median plus two times the interquartile range of the main 32 EEG

channels (not including the externals), the mastoid channel was deemed too noisy and only

the other mastoid channel was used for reference. In two non-autistic participants, both

mastoid channels exceeded this range, and we used the average of neighboring P7 and P8

electrodes instead.

Calculating evoked responses

All analyses of the EEG were done using channel Cz referenced to the mastoids.

Event-related potentials (ERP) to the stimuli and the button presses (the participant's

response in the task) were first calculated individually for each participant. ERPs to the

stimuli were baselined to the interval of -100 to 0 ms before the stimulus, and ERPs to the

button presses were baselined -400 to -200 ms; we chose a different interval before the
button press to avoid a negativity that appeared between -200 to 0 ms in some participants.
To avoid the influence of outlier EEG signals on the evoked responses, all ERPs were
calculated using a spline-based median of the evoked responses (see Zuk et al., 2021 for
information on the spline transform). First, each EEG segment was transformed into a basis
of 173 cubic splines which enforced a smooth representation of the EEG segments with a
-3 dB cutoff frequency around 40 Hz. Second, the median of the basis spline weights across
trials was calculated. Then, the basis spline weights were transformed back into the time
domain for interpretation, plotting, and further analysis. These steps were also used when
calculating the median ERP across participants. For the shaded regions in the plots of the
ERP (e.g. Figure 1B), a bootstrap resampling procedure with replacement was done across
participants and repeated 200 times to calculate the 95% confidence interval of the median
ERP. Analyses of individual participant's ERP magnitudes are based on the minima or
maxima of the median ERPs in the time domain.
202
203   <u>Study 2</u>
204   Study 2 was a replication of Study 1, and the same software and hardware were
205   used for administering this experiment. 51 participants (26 non-autistics and 25 ASD) took
206   part in this replication. Of these, 18 non-autistics and 18 ASD had previously participated
207   in Study 1. The time interval between Study 1 and Study 2 was 1-2 years.
208
Group classification
To explore group-level classification, we trained a linear support vector machine
(SVM) with second-degree polynomial features using scikit-learn (Python). We chose this
model architecture because it performed consistently well across a range of data splits (80%
for training, 20% for testing) and performed at least as well as alternative approaches,
including random forest, logistic regression, decision trees, and more complex SVMs with
radial basis function (RBF) kernels. The feature set included behavioural recency bias after
correct and easy trials, the ERP peak-to-peak amplitude difference between easy and hard
trials, and FRN magnitude. The preprocessing pipeline involved z-scoring and polynomial
feature expansion (degree = 2, excluding the bias term). The regularisation parameter (C),

which controls the trade-off between margin size and classification error, was tuned using a grid search over logarithmically spaced values between 0.01 and 100, with 5-fold cross-validation on the training set. Model performance was evaluated using accuracy and area under the ROC curve (AUC) on a held-out test set, and statistical significance was assessed with 1000-iteration permutation testing. For the main analysis, we performed 20 stratified random 80/20 train–test splits and reported average test accuracy and AUC across seeds. We then examined a representative model using the most common hyperparameter (C=0.1) with model performance (accuracy) closest to the median performance, and we used this model to assess the contribution of each feature. Additional evaluations tested generalization across participants (training on unique individuals and testing on non-unique, or overlapping, individuals).

Feature contribution was assessed by inspecting the weights associated with individual features and calculating the amount of impact each weight had on the data. In general, linear SVM classifiers are trained to choose the weights w associated with each feature or interaction feature so that the decision function $f(x)=w^\mathsf{T}x+b$ best classifies positive values of $f(x)$ to one class and negative values to the other, where x is a vector for each sample containing feature values and b is a (constant) bias term (here, non-autistic and ASD are the positive and negative classes respectively). By isolating just $w^\mathsf{T}x$ for each feature, we can compare the feature's contribution to the classifier. A feature that is a good discriminator (and has low class overlap) should transform data from a non-autistic participant to a positive value and an ASD participant to a negative value. We plot these feature contributions using a kernel density estimator (KDE) in Figure 7B-D.

Results

Event-related potential (ERP) and behavioral responses to stimuli are similar for people with and without ASD

Participants performed two-tone pitch discrimination, during which we recorded EEG. On each trial, two pure tones with different frequencies were presented serially. Participants were asked to determine which of the two tones had a higher pitch (Figure 1A, in the example the second tone frequency is "higher"). Upon responding, they were immediately presented with visual feedback: "Correct" in green text or "Incorrect" in red

text. Since we were interested in the ability to track performance, which might depend on performance level, we aimed to attain similar success rates for the two groups. We used a simple temporal protocol, with two-second intervals both between stimuli and between trials, in order to minimize possible effects of within-trial neural adaptation that may differ in neurodivergent groups (Jaffe-Dax et al., 2017). With these intervals, the peaks of the three main EEG response components (N1 between 70-130 ms, P2 between 150-230 ms, and P3 between 250-350 ms) to the two tones (f1 and f2) were similar in the two groups (Rank-sum test: 1^{st} N1, $z = -0.52$, $p = 0.606$; 2^{nd} N1, $z = -0.11$, $p = 0.912$; 1^{st} P2, $z = 0.18$, $p = 0.854$; 2^{nd} P2, $z = -0.43$, $p = 0.668$; 1^{st} P3, $z = -1.88$, $p = 0.061$; 2^{nd} P3, $z = -0.87$, $p = 0.384$). We found difference in N1 magnitudes between the tones, indicating no significant within-trial adaptation (signed-rank test: Non-autistics, $z = -0.87$, $p = 0.385$; ASD, $z = -0.85$, $p = 0.396$). P2 showed a weak, though significant, adaptation in non-autistics (Non-autistics, $z = -2.23$, $p = 0.023$), which was not significantly different from the absence of adaptation in ASD (ASD, $z = -0.96$, $p = 0.339$; rank-sum test for the difference between groups: $z = -0.92$, $p = 0.357$). As expected, both groups had a P3 component following the second tone (Figure 1B), which characterizes time points where a categorical decision is made ("which tone was higher?"; Nahum et al., 2010; Jaffe-Dax et al., 2017). The P3 magnitude was similar for the two groups (rank-sum test: $z = 0.209$, $p = 0.835$). Behaviorally, both groups had similar median success rates in the task (Figure 1C; Non-autistics = 71.9%, ASD = 70.0%; rank-sum test: $z = 0.76$, $p = 0.447$), though the cross-participant distribution was broader in the ASD group.

Figure 1: Non-autistics and ASD show similar performance and similar ERP responses to the tones in each trial. (A) A schematic diagram of a trial's temporal structure. Participants performed a two-tone discrimination task where they had to determine if the pitch of the second tone was higher than that of the first. Visual feedback ("Correct" or "Incorrect") was given immediately upon response in every trial. (B) Grand average response (median of participants' median ERPs, see Methods) for electrode Cz. Time zero denotes the onset of the first tone on each trial. Thick lines show the median across participants' median ERPs. Shaded regions show the 95% confidence interval of the median using a bootstrap resampling across participants. Non-autistics are plotted in blue (top) and ASD in green (bottom). Dashed vertical lines mark the onset of each tone. The three main response components, N1, P2 and P3 are marked on the top plot. (C) Distributions of participants' performance accuracy, in each group. Each dot denotes the percent correct of an individual participant, the bars indicate the median in each group. Median performance was similar in the two groups (no significant difference in a rank-sum test).

The Feedback-Related Negativity (FRN) is significantly weaker in ASD

Under similar task performance and EEG responses to tones in the two groups, we next asked whether the response to external feedback is also similar. The visual feedback was provided immediately (within 20 ms) after the participant's button press. To quantify the EEG response to the feedback, we aligned our analyses to participants' button press. We rejected three ASD participants who performed extremely well in the experiment and thus had too few incorrect answers to get a reliable FRN response (< 6 trials incorrect; 3 best performers, see Figure 1C where these participants performed better than all non-

autistics). The EEG signal following "incorrect" feedback is known as the "feedback-
related negativity" (FRN) (Miltner et al., 1997). The topography of activity at the time of
the negativity (e.g. inset in Figure 2B) has been source-localized in previous studies to
ACC (Dehaene et al., 1994; Miltner et al., 1997). Figure 2A plots samples of 12 single
participants, six with ASD (right) and six non-autistics (left), matched for the rank order
of their FRN with respect to their group. It illustrates that the majority of non-autistic
participants (sampled here by 1st to 26th in descending FRN magnitude) have a clear and
large FRN, whereas most ASD (also sampled here by 1st to 26th) do not.

To quantify the magnitude of the FRN, we determined, for each participant, the
minimal response between 200-350 ms after the participant's button press (Figure 2C).
This time window captured the span of the negativity for all participants (dashed vertical
lines in Figure 2A, median ERP in Figure 2B). FRN magnitude was then quantified as the
difference between the minimum in incorrect and the minimum in correct trials. Figure 2C
indicates that in both groups the minimum for the evoked response to incorrect trials was
more negative than to positive trials (signed-rank test: Non-autistic, $z = 5.16$, $p < 0.001$;
ASD, $z = 3.16$, $p = 0.002$). However, this was significantly smaller in ASD participants
than non-autistics (Figure 2D, rank-sum test: $z = -3.56$, $p < 0.001$). Importantly, the effect
size of this group difference was large (Cohen's $d = 0.90$). For reference, the effect size
(Cohen's d) of AQ, a questionnaire specifically designed to detect ASD traits (Baron-
Cohen et al., 2001), was 1.10 for our participants, comparable to the effect size of the FRN
measure. We did not find a correlation within each group between AQ score and FRN
magnitude (Non-autistics, $\rho = -0.274$, $p = 0.117$; ASD, $\rho = -0.062$, $p = 0.733$), perhaps
because FRN magnitude also depends on performance levels (in non-autistics it was shown
that "incorrect" feedback yields a larger error- or feedback-related negativity when the task
is easier (Miltner et al., 1997) and when errors are more unexpected (Gehring et al., 1993;
Holroyd and Coles, 2002)).

To highlight this difference at the single participant level, figure 2D on the right
shows a histogram of the differences between incorrect and correct evoked responses in
each group. It shows that by using discriminant boundary of -2.5 µV (close to the median
magnitude for ASD participants, plotted in Figure 2D), only a minority of the ASD

participants (13/30) show an FRN magnitude lying below this boundary, whereas nearly

all (33/35) of the non-autistics have magnitudes below this boundary.

Figure 2: ASD participants' EEG activity in response to external feedback (Feedback-Related Negativity, FRN) is weaker than that of non-autistics'. (A) Single participants' FRN, defined as the difference between (minimal) responses following "Correct" (blue) and "Incorrect" (red) feedback within 200-350 ms after their behavioral response (range marked by the dashed lines). FRNs were ranked according to these differences. We plotted responses of six participants from each group, whose ranked magnitude of FRNs were matched with respect to their group (non-autistics on the left, ASD on the right). These ranks are denoted on the right (1st being largest). While all six non-autistics had an FRN, only two of the six plotted autistics' responses manifest a clear FRN. (B) Median ERP across participants, shaded error bars indicate the bootstrapped 95% confidence intervals of the median. The topography of the difference in voltage between Incorrect and Correct trials averaged within this time interval is shown above each plot. (C) Quantification of the FRN magnitude as the difference between the ERP minimum for incorrect and correct trials, within the range 200-350 ms as indicated in A and B, plotted separately for each participant. At the group level both

groups show a significant FRN. (D) Individual dots show the difference in magnitude between correct and
incorrect, and the horizontal bars denote the median of each group. The group difference in FRN is highly
significant (*** = p < 0.001). On the left, each symbol denotes a single participant. Note that more than half
of the ASD participants (above the green line) are distributed symmetrically around zero difference, meaning
that they have no clear FRN. By contrast, only 2/35 neurotypicals do not show a clear FRN (upper-most blue
dots in the plot). The histogram on the right also shows the distribution of FRN in each group with 2.5 μV
bins. It illustrates the efficiency of FRN magnitude as an indicator of group; a plausible discrimination
boundary of -2.5 μV is shown as a red dashed line.
348
Only non-autistics' EEG responses are modulated by task difficulty

External feedback is often processed in tandem with one's confidence in their
decision (Fleming and Daw, 2017; Grogan et al., 2023). When participants are more
confident in their knowledge of task-relevant information they are more likely to use that
information (Samaha et al., 2019). This is a form of internal feedback that can also
modulate the participants' behavior in the task, but does it affect their EEG responses to
feedback?

As a proxy for participants' assurance of their perceptual decision, we separated the
trials into "Hard" and "Easy" (Figure 3A) using a median of the frequency difference in
the trial as the threshold (the threshold of 1.242 semitones was the median frequency
difference across all trials and participants). We reasoned that in easy trials, participants
are more confident in their decision. Indeed, non-autistics' EEG responses were modulated
by task difficulty (Figure 3B), such that the amplitude was larger following hard trials. We
used the peak-to-peak difference in the evoked response, calculated as the difference
between the minimum in 0-150 ms following the participant's button press and the
maximum in 300-450 ms following the button press (see the dashed lines in Figure 3B) to
quantify the magnitude of this difference (Figure 3C). We found that, in non-autistics, this
difference was significantly larger in hard trials compared to easy ones (signed-rank: $z = -$
2.69, $p = 0.007$), suggesting that participants' effort following difficult discriminations
modulated the EEG response to feedback. ASD participants, however, did not show any
difference ($z = 1.51$, $p = 0.131$). The group difference was significant (rank-sum: $z = -2.98$,
$p = 0.028$; Cohen's $d = 0.574$).

Our results indicate that both internal feedback and external feedback significantly
modulate the EEG response in non-autistics but not in ASD, showing that ASD participants
have weaker processing of both types of feedback.
374

375
376

Figure 3: Weaker effects of internal feedback were observed in ASD participants compared to non-autistics. (A) Diagram showing the distinction between easy and hard trials. Easy trials have a larger frequency difference than the median difference across all possible trials (1.242 semitones), while hard trials have frequency differences smaller than this difference. (B) Median evoked responses across participants, time-aligned to the behavioral response, plotted separately for hard (purple) and easy (orange) trials. (C) The peak-to-peak difference in the evoked EEG responses to hard compared to easy trials is significantly larger in non-autistic participants. Note that, here, ASD participants with excellent performance (also excluded from Figure 2) were also rejected to remain consistent with the other measures used later for autism/non-autism classification. (D) Individual dots show the difference in peak-to-peak magnitude between easy and hard trials. The group difference is significant (rank-sum z = -2.98, p = 0.028, Cohen's d = 0.574). * p < 0.05, *** p < 0.001.

Only non-autistics' perceptual bias is modulated online by task difficulty and success

The modulation of non-autistics' EEG response by task difficulty suggests that it may also have a behavioral impact. We therefore examined how external and internal feedback affect participants' response tendencies in subsequent trials. Previous work had shown that participants' behavioral responses in a two-interval forced choice task are substantially biased by previous trials' statistics (Ashourian and Loewenstein, 2011; Raviv et al., 2012, 2014; Akrami et al., 2018; Lieder et al., 2019). Within a Bayesian framework, it was proposed that since the mental representation of the first tone is noisier, it is contracted more towards an implicit prior (predicted frequency), which is formed by previous trials' frequencies (Ashourian and Loewenstein, 2011) (but see Boboeva et al., 2024). Because of this, when the first tone is closer to this prior, contraction improves performance (consequently termed bias+ trials), whereas when it is farther, performance deteriorates (bias-) (Raviv et al., 2012; Akrami et al., 2018) (Figure 4A).

We calculated the recency bias, namely the bias of the first tone towards the most recent trial's average frequency (Lieder et al., 2019). The bias is the difference in success rate between bias+ trials (where this contraction improves performance) and bias- trials (where performance decreases) (Ashourian and Loewenstein, 2011; Raviv et al., 2012). In line with previous work (Lieder et al., 2019), the bias for ASD participants was on average smaller than non-autistics (rank-sum test: z = 2.39, p = 0.017). Next, we divided these trials based on the difficulty of the previous trial (easy versus hard, based on the threshold of 1.242 semitones) and based on the participant's accuracy in the previous trial (correct or incorrect) (Figure 4B). All participants were included for previously correct trials. For incorrect trials, we excluded participants who did not have sufficient trials in at least one of the conditions; 32 non-autistics and 28 ASD participants were included.

We first looked at the effect of the previous trial's difficulty (internal feedback) on performance, separately for correct and incorrect trials (external feedback). Importantly, we found that non-autistic participants showed a larger recency bias when the previous trial was easy compared to when it was hard (signed-rank test: z = -3.01, p = 0.003). In contrast, ASD participants did not show this effect on bias in correct trials (z = 1.02, p = 0.309), and the bias difference between easy and hard trials when the previous trial was easy was significantly weaker than non-autistics' (z = 2.88, p = 0.004). The group difference was

particularly driven by a difference in bias when participants were previously correct in an
easy trial, where non-autistics showed a larger bias than ASD ($z = 2.72$, $p = 0.006$, Cohen's
$d = 0.752$).

Together, the behavioral group differences suggest that the magnitude of the
recency bias and the participant's subsequent decisions are affected by the difficulty and
accuracy in the previous trial. When non-autistic participants are confident in the stimulus
information from the previous trial (after easy) and their confidence has been externally
validated (after correct), the effect on the perceptual prior is stronger on their subsequent
decisions. ASD participants have weaker processing of internal feedback, and their
decisions were unaffected by the accuracy or internal reliability of their response.
Importantly, this observation is in line with previous studies reporting reduced online
flexibility in updating perceptual priors based on recent events (Sapey-Triomphe et al.,
2021). But while studies assessed stimulus statistics, here we assessed the impact of
certainty and show that the effect on recency bias is reliable and effective.

Figure 4: Only non-autistics show a larger recency bias following correct and easy trials. (A) A schematic illustration of recency bias towards recent trial's frequencies in perceptual decisions. In bias+ trials the first tone's frequency is closer to the mean frequency of the previous trial,, whereas in bias- the second tone is closer. Better performance occurs in "bias+" trials versus "bias-" trials though they differ only in their preceding trial's frequency. (B) Diagram illustrating the main question of how internal feedback (Hard / Easy) and external feedback (Correct / Incorrect) in the previous trial affect the participant's choices in the current trial. (C) When non-autistic participants were previously correct yet more confident (easy trials), there was significantly larger recency bias in the subsequent trial. This effect was not found for the ASD group. (D) A closer comparison of the bias difference between groups when the previous trial was both easy and correct, plotted identically to figures 2D and 3D. The group difference was significant (p = 0.006, Cohen's d = 0.752). ** p < 0.01

 A mini control experiment – Real feedback is important
450
Since an FRN response to perceptual performance has not been studied before to
the best of our knowledge, and to better understand its nature and the temporal dynamics
of the response, we administered a small control experiment with three non-autistic
participants and three ASD participants. The non-autistic participants were selected
because they had a large FRN in the previous experiment and were available for the follow
up, and the ASD participants were selected based on willingness to participate in a follow-
up session. All participants did three blocks in a random order, where each block had a
different type of feedback: True feedback was identical to the previous experiment (as in
Figure 1A); Delayed feedback showed true feedback, but the feedback was delayed by 200
ms relative to the participant's response; and Random feedback showed "Correct" or
"Incorrect" randomly after the participant's response. In the latter condition, participants
were told that the feedback would be random and uninformative, but they were still
instructed to look at the feedback during the task.

Our first test compared true and delayed feedback in non-autistics; for participants
who showed a reliable FRN, was the ERP timed to the button press or the feedback? When
the feedback was delayed, both the correct response and the incorrect response were also
delayed by 200 ms (Figure 5A), indicating that the ERP and FRN was timed to the external
feedback rather than to the response itself. Next, we compared the FRN to true and random
feedback, asking whether, in the absence of informative feedback, the FRN response
disappears. In non-autistics, randomizing the feedback causes the FRN to disappear (Figure
5B). Additionally, non-autistics' positivity 200 ms after a correct response also
disappeared, indicating that the signal was related to information in the feedback rather
than the visual input. Interestingly, the ASD participants had EEG responses to both correct
and incorrect feedback that did not appear to differ between the true feedback and random
feedback conditions, unlike the non-autistics (Figure 5B). While preliminary, this result
suggests that ASD participants may process feedback the same way irrespective of its
informativeness.

Taken together, the mini follow-up study indicates that the responses we observed
in non-autistics were time-locked to the feedback. Importantly, both the positivity in the

response to Correct feedback after 200 ms (Figure 5B) and the FRN reflected the
processing of errors in the task, rather than visual processing of the feedback.

Figure 5: A (mini) follow-up control experiment (3 participants from each group) for feedback timing
and validity. (A) Evoked responses to correct (blue) and incorrect (red) feedback presented immediately
after the participant's response (left), and delayed by 200ms (right), shown separately for the non-autistic
(top) and ASD participants (bottom). For non-ASD participants the FRN response was delayed by 200ms.
For autistic participants there is still little to no FRN. Interestingly, at a shorter time window than the FRN

there is a large negative response, which does not differ between correct and incorrect response, and suggests
some enhanced anticipation that should be studied further. The thin colored lines are evoked responses for
individual subjects, and the thick line is the average. Dashed lines label the peaks and troughs of the evoked
response to true feedback (on the left). The dashed lines on the right mark the same time points in A shifted
by 200 ms, to make clear the shift in the response. (B) Non-autistics (left) and ASD (right) evoked responses
after seeing "Correct" or "Incorrect" on screen when the feedback was true (blue, as in A) or randomized
(orange).

495
496
497   *Follow-up study replicates prior results*
498
In Study 1 we got impressive results for three aspects of online performance
monitoring. Given that the two-tone discrimination protocol is composed of 121 trials only,
we wanted to verify that these parameters are replicable and that a reliable classification
can be done based on these parameters. We therefore invited participants to return for a
follow-up assessment with the same protocol, approximately 1-2 years after participating
in Study 1. Overall, we had 51 participants in this study (25 non-autistics and 23 ASD),
and 36 of them were people who also participated in study 1, allowing assessment of within
participants' replicability. Two ASD participants had above 95% trials correct in
discrimination (as they did in Study 1) and hence were excluded from FRN calculations
(in both Study 1 and Study 2), leaving 18 non-autistics and 16 ASD who participated in
both studies. As in Study 1, overall performance was similar for the two groups (mean
percent correct in non-autistics = 79%, in ASD = 78%).

All three measures related to performance monitoring that were significantly
different between groups in Study 1 were also significantly different in Study 2 (Figure
6A-C): FRN magnitude was larger in non-autistics (Figure 6A, $z = -2.87$, $p = 0.004$,
Cohen's $d = 0.852$); ASDs' difference in ERP magnitude following easy and difficult trials
was smaller than non-autistics' (Figure 6B, $z = 2.18$, $p = 0.028$, Cohen's $d = 0.680$) and
ASDs' reduced flexibility, as manifested in their reduced effect of prior's magnitude
following easy trials, was smaller (Figure 6C, $z = 2.70$, $p = 0.006$, Cohen's $d = 0.883$).

Importantly, within people who participated in both studies, each of these measures
were correlated, even though more than a year passed between the two measures (Figure
6D-F; Pearson's correlations – FRN: Non-autistics, $r = 0.484$, $p = 0.042$; ASD, $r = 0.760$,
$p < 0.001$; Hard-Easy Peak-to-peak: Non-autistics, $r = 0.437$, $p = 0.070$; ASD, $r = 0.512$, p

= 0.042; Bias following easy and correct: Non-autistics, r = 0.540, p = 0.021; ASD, r =

0.767, p < 0.001). The replicability and correlation of three measures indicate that each of

these three measures reliably characterize the participants.

Figure 6 – In Study 2, all parameters were replicated and correlated within participants across

sessions. (A) Single participant distribution of FRN magnitude, plotted identically to Figure 2D (Cohen's d

= 0.852). (B) Single participant distribution of the difference between Hard and Easy peak-to-peak

magnitudes of the EEG following the participant's button press, plotted identically to Figure 3D (Cohen's d

= 0.680). (C) Single participants' distribution of magnitude of online bias updating (following easy and

correct trials), plotted identically to Figure 4D (Cohen's d = 0.883). (D) Within participant correlation of

these parameters between study 1 and study 2, for the subgroup that participated in both studies.

Additionally, we combined the participants for study 1 and study 2 (taking only

their first session; 42 non-autistics, 35 ASD) and examined if there were correlations

between these measures. Interestingly, we found no correlations between measures within

each group (Spearman's correlations, two-tailed, FRN vs peak-to-peak difference: non-

autistics, $\rho = 0.06$, $p = 0.727$; ASD, $\rho = -0.19$, $p = 0.262$; FRN vs bias after easy and correct:

non-autistics, $\rho = 0.12$, $p = 0.442$; ASD, $\rho = -0.12$, $p = 0.483$; peak-to-peak difference vs

bias after easy and correct: non-autistics, $\rho = 0.21$, $p = 0.176$; ASD, $\rho = 0.23$, $p = 0.197$).

This implies that the measures may represent distinct cognitive processes which are all
generally weaker in autistics.
544
Combining these neurocognitive measures classifies ASD participants
Given that the three performance-monitoring measures were both replicable across
sessions and were uncorrelated with each other, we reasoned that a classification model
based on all three could be both effective and informative, as it can point to the specific
performance monitoring characteristics that are atypical in each participant.
We trained a linear support vector machine (SVM) classifier using second-degree
polynomial features (for details see Methods). This model choice allows for linear decision
boundaries while capturing both main effects and pairwise interactions among features.
Over random train-test splits of all participants, the model achieved a median test accuracy
of 80.4% (±7.0%) and a mode (for 6/20 train-test splits) of 82.6%. Similarly, the median
receiver operating characteristic (ROC) area under the curve (AUC) was 0.86 (±0.06).
AUC represents the probability that a randomly chosen non-autistic participant's data will
be ranked higher by the model than a randomly chosen autistic participant's data, where an
AUC of 0.5 is chance level. An AUC above 0.8 is indicative of a good model classifier
(Hanley and McNeil, 1982).
Next, a representative model using the most common hyperparameter setting with
performance close to the median accuracy across all models was selected to examine
feature contributions to classification. This specific model achieved 82.6% test accuracy
and an AUC of 0.88 (Fig. 7B-D; see Methods). FRN and behavioral recency bias following
easy and correct trials contributed more to the separation of classes than the peak-to-peak
ERP magnitude, where separation can be understood based on the amount of overlap
between the density distributions for non-autistic and ASD groups. Interaction effects
between the peak-to-peak amplitude and FRN magnitude as well as between FRN
magnitude and bias were also contributive (data not shown).
Performance remained robust when training on participants' initial experience with
the experiment (i.e. all participants in study 1 and non-repeating participants in study 2)
and testing on individuals who had already participated once before (participants in study
2 who also completed study 1) with an accuracy of 77% and AUC of 0.85. Permutation

tests confirmed that classification performance was significantly above chance across all

data splits (p < 0 .001).

Overall, these results demonstrate that these three behavioral and

neurophysiological measures alone can identify an autistic participant with good accuracy,

implying that they are a promising biomarker of autism.

Figure 7: Group classification based on the three parameters of performance monitoring. (A) Receiver

operator characteristic (ROC) curve for the representative model (AUC is 0.88). (B-D) shows the contribution

of each main parameter to this classification model, with negative values favoring ASD classification, and

positive values favoring NT. The FRN magnitude difference (D) has the largest impact (i.e. largest effect

size) since there are fewer areas of overlap between the true classes and each is pushed away, while behavioral

bias (B) provides a smaller but still significant contribution. The ERP peak-to-peak amplitude (C) contributes

much less.

Discussion

In this study, non-autistics and autistic individuals performed a two-tone

discrimination task, exhibiting similar levels of performance. Yet autistics' flexibility

during performance was much reduced; while non-autistic individuals adapted their

reliance on perceptual predictions and their EEG responses to feedback and task difficulty,

people with autism did not. These neurocognitive measures were highly robust, but

interestingly they were not correlated with each other. Together, however, they provided a

reliable, theory-based, classifier of autism.

First, we found that error processing of external feedback, manifested by a large
EEG component termed the feedback-related negativity (FRN) is produced by almost all
non-autistics. By contrast, FRN was weaker or absent among most ASD participants.
Groups also differed in their processing of internal feedback: non-autistics had a larger
EEG response to hard trials (small frequency difference) compared with easy trials, while
no such difference was found in autistic participants. Behaviorally, non-autistics showed a
larger perceptual bias after correct and easy trials, indicating that estimates of reliability
and confidence affect one's subsequent perceptual prior (Samaha et al., 2019). ASD
participants did not show this trend, demonstrating less flexibility in updating their
perceptual prior that stems from reduced monitoring of their internal evaluation of
performance.

Since ACC is involved in both error monitoring and evaluating perceptual priors,
and since both characterize autism, we initially thought they may stem from a common
mechanism. However, ACC's function in cognition is multifaceted, responsible for
monitoring many pieces of information that inform subsequent actions. Indeed, ACC has
a complex parcellation (Bush et al., 2000), with different subregions functionally
connected to various areas of frontal, parietal, and limbic regions of the brain (Devinsky et
al., 1995; Margulies et al., 2007). Though all three of our measures have been associated
with ACC, the lack of correlation between them may not be surprising, as perceptual and
response bias may stem from different processing levels (Margulies et al., 2007; Fritsche
et al., 2017; Oesch et al., 2025). Taken together, our results suggest that different
atypicalities associated with ACC may contribute to autism spectrum.

The reduction in serial effects in perceptual bias that we found here links two
findings in the literature. One indicates that ASD participants use recent prior information
less than non-autistics in perceptual tasks. This is the case when inter-trial intervals are
short and require fast trial-to-trial updating (Lieder et al., 2019; Boboeva et al., 2024) or
when external statistics are volatile such that optimal inference yields large trial-to-trial
prediction updating (Lawson et al., 2017; Noel et al., 2024). Similarly, other studies have
indicated poor online error correction in motor tasks (Noel et al., 2021; Vishne et al., 2021;
Kasten et al., 2023) and perhaps reduced conflict resolution when fast responses are

required (South et al., 2010). Both are suggested to be mediated by an ACC pattern of
activation (Botvinick et al., 2001; Kolling et al., 2016a).

The weaker FRN response in the ASD group aligns with other human studies, using
functional imaging, showing weaker activation in autistics' ACC (Sapey-Triomphe et al.,
2023). ACC encodes the expectation of errors or conflicting information (Carter et al.,
1998; Botvinick et al., 1999; Kennerley et al., 2006; Brown and Braver, 2007) (for review
see also (Botvinick et al., 2001)) which is subsequently important for identifying changes
in task rules and the value of future choices (Kolling et al., 2016b; Takeuchi et al., 2022).
We now show that these ACC signals (as measured with the FRN and peak-to-peak
differences in the EEG response) are produced during error monitoring in perceptual
discrimination. This also affected the majority of the ASD participants (17/30 ASD
participants had magnitudes above -2.5 µV) and its effect size (Cohen's d = 0.90, see Figure
2) was comparable to that of the AQ self-report questionnaire (Baron-Cohen et al., 2001)
(d = 1.10, see Table 1), designed specifically to detect self-report social difficulties in
autism.

Previous EEG work had also shown that the error-related negativity in the absence
of external feedback (hence reflecting internal feedback) is weaker in ASD participants
compared to non-autistics (Vlamings et al., 2008; Sokhadze et al., 2010; South et al., 2010;
Hüpen et al., 2016). However, evidence for group differences in the FRN were mixed
(Hüpen et al., 2016). Importantly, much of this work had focused on the effect of feedback
on reward expectations (Groen et al., 2008; Larson et al., 2011; Bellebaum et al., 2014).
Specifically, these studies varied the probability that the participant could get a reward after
making a decision. Our results imply that group differences in error monitoring require the
formation of perceptual expectations rather than processing of the reward outcome itself.
In prior work, the actual percept generating the error could be dismissed in subsequent
trials because the probability of a reward was randomized (Larson et al., 2011), while in
our study it is well-established that participants use previous information to both learn
about the task rules (Lak et al., 2020; Pedrosa et al., 2023) and update their perceptual bias
in subsequent trials (Raviv et al., 2012; Akrami et al., 2018; Lieder et al., 2019).

By combining all three measures of reduced flexibility in updating implications of
outcomes, accurate classification of ASD versus non-autistic individuals (above 80%) was

possible, highlighting the potential utility of this multi-dimensional approach for
identifying neurocognitive signatures of autism. Importantly, the measures were also
reliably reproduced within participants though the studies were conducted more than a year
apart. Taken together, these results demonstrate the importance of several independent
components in error monitoring during a perceptual task and merge two lines of
observations in the ASD literature. It is notable that despite the variability within groups,
each of these three measures reliably differed between groups.

Several studies have addressed the question of biomarkers for ASD. They
compared motor characteristics during development (Wu et al., 2018), based on atypical
brain connectivity (Pagnozzi et al., 2018), altered event-related potentials (Sysoeva et al.,
2018; Mason et al., 2022), differences in neurochemical (Yizhar et al., 2011; Robertson et
al., 2016; Marotta et al., 2020) or immunological profiles (Xu et al., 2015; Masi et al.,
2017). However, the heterogeneity of ASD complicates biomarker development (Charman
et al., 2017; Frye et al., 2019; Jensen et al., 2022). Our study suggests that slower updating
of priors is a general trait. It is in line with a recent rodent study, which assessed three
genetic mouse models of ASD, and found that all underutilize adaptive statistical
regularities during decision-making, and frontal cortices underweight sensory observation
and prediction errors (Noel et al., 2024). Similarly, reduced online error monitoring and
use of perceptual priors was found before in several different paradigms (South et al., 2010;
Noel et al., 2021, 2022; Sapey-Triomphe et al., 2021, 2023), further supporting the
hypothesis that this is a prevailing characteristic.

Taken together, our findings open a promising avenue for developing objective,
reliable, neurocognitive markers for better understanding of autism and for measuring the
specific impact of intervention on behavioral monitoring in autism.

Acknowledgements

This work was funded by the Israel Science Foundation (grant 1731/24) and by the European Research Council (grant 833694). NZ was also funded by the ELSC-SWC Postdoctoral Fellowship, co-advised by MA and AA. YW was also funded by the ELSC PhD Fellowship. KD was funded by the Gatsby Charitable Foundation (GAT3850). AA was funded by the Gatsby Charitable Foundation (GAT3755) and the Wellcome Trust (219627/Z/19/Z).

Author contributions

NZ, AA, and MA designed the experiment. NZ, JS, YW, VS collected the data. NZ, JS, and KD wrote the analysis scripts and analyzed the data. NZ, JS, KD, AA, and MA drafted the manuscript, and all authors contributed to editing the manuscript.

Data availability

All EEG and behavioral data from the experiment are available here: <OSF link>. The code used to run the experiment and analyze the data can be found here: https://gitlab.com/auditory-predictive-coding/erp-adaptation. The code used to run the classification analysis can be found here: https://github.com/kiradust/asd-cat.

Declaration of interests

The authors report no competing interests.

References

Akrami A, Kopec CD, Diamond ME, Brody CD (2018) Posterior parietal cortex represents sensory history and mediates its effects on behaviour. Nature 554:368–372.

Ashourian P, Loewenstein Y (2011) Bayesian Inference Underlies the Contraction Bias in Delayed Comparison Tasks. PLOS ONE 6:e19551.

Baron-Cohen S, Wheelwright S, Skinner R, Martin J, Clubley E (2001) The Autism-Spectrum Quotient (AQ): Evidence from Asperger Syndrome/High-Functioning Autism, Males and Females, Scientists and Mathematicians. J Autism Dev Disord 31:5–17.

Beckmann M, Johansen-Berg H, Rushworth MFS (2009) Connectivity-Based
Parcellation of Human Cingulate Cortex and Its Relation to Functional
Specialization. J Neurosci 29:1175–1190.

Behrens TEJ, Woolrich MW, Walton ME, Rushworth MFS (2007) Learning the value of
information in an uncertain world. Nat Neurosci 10:1214–1221.

Bellebaum C, Brodmann K, Thoma P (2014) Active and observational reward learning in
adults with autism spectrum disorder: relationship with empathy in an atypical
sample. Cognitive Neuropsychiatry 19:205–225.

Bellebaum C, Daum I (2008) Learning-related changes in reward expectancy are
reflected in the feedback-related negativity. European Journal of Neuroscience
27:1823–1835.

Boboeva V, Pezzotta A, Clopath C, Akrami A (2024) Unifying network model links
recency and central tendency biases in working memory Sharpee TO, Colgin LL,
eds. eLife 12:RP86725.

Botvinick M, Nystrom LE, Fissell K, Carter CS, Cohen JD (1999) Conflict monitoring
versus selection-for-action in anterior cingulate cortex. Nature 402:179–181.

Botvinick MM, Braver TS, Barch DM, Carter CS, Cohen JD (2001) Conflict monitoring
and cognitive control. Psychological Review 108:624–652.

Botvinick MM, Cohen JD, Carter CS (2004) Conflict monitoring and anterior cingulate
cortex: an update. Trends in Cognitive Sciences 8:539–546.

Brown JW, Braver TS (2007) Risk prediction and aversion by anterior cingulate cortex.
Cognitive, Affective, & Behavioral Neuroscience 7:266–277.

Bush G, Luu P, Posner MI (2000) Cognitive and emotional influences in anterior
cingulate cortex. Trends in Cognitive Sciences 4:215–222.

Carter CS, Braver TS, Barch DM, Botvinick MM, Noll D, Cohen JD (1998) Anterior
Cingulate Cortex, Error Detection, and the Online Monitoring of Performance.
Science 280:747–749.

Cauda F, Geda E, Sacco K, D'Agata F, Duca S, Geminiani G, Keller R (2011) Grey
matter abnormality in autism spectrum disorder: an activation likelihood
estimation meta-analysis study. J Neurol Neurosurg Psychiatry 82:1304–1313.

Chan AS, Han YMY, Leung WW, Leung C, Wong VCN, Cheung M (2011)
Abnormalities in the anterior cingulate cortex associated with attentional and
inhibitory control deficits: A neurophysiological study on children with autism
spectrum disorders. Research in Autism Spectrum Disorders 5:254–266.

Charman T et al. (2017) The EU-AIMS Longitudinal European Autism Project (LEAP):
clinical characterisation. Molecular Autism 8:27.

D'Cruz A-M, Mosconi MW, Ragozzino ME, Cook EH, Sweeney JA (2016) Alterations
in the functional neural circuitry supporting flexible choice behavior in autism
spectrum disorders. Transl Psychiatry 6:e916–e916.

Dehaene S, Posner MI, Tucker DM (1994) Localization of a Neural System for Error
Detection and Compensation. Psychol Sci 5:303–305.

Devinsky O, Morrell MJ, Vogt BA (1995) Contributions of anterior cingulate cortex to
behaviour. Brain 118:279–306.

Falkenstein M, Hohnsbein J, Hoormann J, Blanke L (1991) Effects of crossmodal divided
attention on late ERP components. II. Error processing in choice reaction tasks.
Electroencephalography and Clinical Neurophysiology 78:447–455.

Fleming SM, Daw ND (2017) Self-evaluation of decision-making: A general Bayesian
framework for metacognitive computation. Psychological Review 124:91–114.

Fritsche M, Mostert P, de Lange FP (2017) Opposite Effects of Recent History on
Perception and Decision. Current Biology 27:590–595.

Frye RE, Vassall S, Kaur G, Lewis C, Karim M, Rossignol D (2019) Emerging
biomarkers in autism spectrum disorder: a systematic review. Annals of
Translational Medicine 7:792–792.

Gehring W, Coles M, Meyer D, Donchin E (1995) A brain potential manifestation of
error-related processing. Electroencephalogr Clin Neurophysiol Suppl 44:261–
272.

Gehring WJ, Goss B, Coles MGH, Meyer DE, Donchin E (1993) A Neural System for
Error Detection and Compensation. Psychol Sci 4:385–390.

Gehring WJ, Goss B, Coles MGH, Meyer DE, Donchin E (2018) The Error-Related
Negativity. Perspect Psychol Sci 13:200–204.

Gehring WJ, Willoughby AR (2002) The Medial Frontal Cortex and the Rapid
Processing of Monetary Gains and Losses. Science 295:2279–2282.

Groen Y, Wijers AA, Mulder LJM, Waggeveld B, Minderaa RB, Althaus M (2008) Error
and feedback processing in children with ADHD and children with Autistic
Spectrum Disorder: An EEG event-related potential study. Clinical
Neurophysiology 119:2476–2493.

Grogan JP, Rys W, Kelly SP, O'Connell RG (2023) Confidence is predicted by pre- and
post-choice decision signal dynamics. Imaging Neuroscience Available at:
https://doi.org/10.1162/imag_a_00005 [Accessed August 10, 2023].

Hanley JA, McNeil BJ (1982) The meaning and use of the area under a receiver operating
characteristic (ROC) curve. Radiology 143:29–36.

Henderson H, Schwartz C, Mundy P, Burnette C, Sutton S, Zahka N, Pradella A (2006)
Response monitoring, the error-related negativity, and differences in social
behavior in autism. Brain and Cognition 61:96–109.

Holroyd CB, Coles MGH (2002) The neural basis of human error processing:
Reinforcement learning, dopamine, and the error-related negativity. Psychological
Review 109:679–709.

Hüpen P, Groen Y, Gaastra GF, Tucha L, Tucha O (2016) Performance monitoring in
autism spectrum disorders: A systematic literature review of event-related
potential studies. International Journal of Psychophysiology 102:33–46.

Hyvärinen A, Oja E (2000) Independent component analysis: algorithms and
applications. Neural Networks 13:411–430.

Jaffe-Dax S, Frenkel O, Ahissar M (2017) Dyslexics' faster decay of implicit memory for
sounds and words is manifested in their shorter neural adaptation King AJ, ed.
eLife 6:e20557.

Jensen AR, Lane AL, Werner BA, McLees SE, Fletcher TS, Frye RE (2022) Modern
Biomarkers for Autism Spectrum Disorder: Future Directions. Mol Diagn Ther
26:483–495.

Kasten K, Jacoby N, Ahissar M (2023) Poor synchronization yet adequate tempo-keeping
in adults with autism. Autism Research 16:1161–1173.

Kennerley SW, Walton ME, Behrens TEJ, Buckley MJ, Rushworth MFS (2006) Optimal
decision making and the anterior cingulate cortex. Nat Neurosci 9:940–947.

Kolling N, Behrens T, Wittmann M, Rushworth M (2016a) Multiple signals in anterior
cingulate cortex. Current Opinion in Neurobiology 37:36–43.

Kolling N, Wittmann MK, Behrens TEJ, Boorman ED, Mars RB, Rushworth MFS
(2016b) Value, search, persistence and model updating in anterior cingulate
cortex. Nat Neurosci 19:1280–1285.

Lak A, Hueske E, Hirokawa J, Masset P, Ott T, Urai AE, Donner TH, Carandini M,
Tonegawa S, Uchida N, Kepecs A (2020) Reinforcement biases subsequent
perceptual decisions when confidence is low, a widespread behavioral
phenomenon Salinas E, Frank MJ, Salinas E, Brody CD, Ding L, eds. eLife
9:e49834.

Larson MJ, South M, Krauskopf E, Clawson A, Crowley MJ (2011) Feedback and
reward processing in high-functioning autism. Psychiatry Research 187:198–203.

Lawson RP, Mathys C, Rees G (2017) Adults with autism overestimate the volatility of
the sensory environment. Nat Neurosci 20:1293–1299.

Lieder I, Adam V, Frenkel O, Jaffe-Dax S, Sahani M, Ahissar M (2019) Perceptual bias
reveals slow-updating in autism and fast-forgetting in dyslexia. Nat Neurosci
22:256–264.

Margulies DS, Kelly AMC, Uddin LQ, Biswal BB, Castellanos FX, Milham MP (2007)
Mapping the functional connectivity of anterior cingulate cortex. NeuroImage
37:579–588.

Marotta R, Risoleo MC, Messina G, Parisi L, Carotenuto M, Vetri L, Roccella M (2020)
The Neurochemistry of Autism. Brain Sciences 10:163.

Masi A, Glozier N, Dale R, Guastella AJ (2017) The Immune System, Cytokines, and
Biomarkers in Autism Spectrum Disorder. Neurosci Bull 33:194–204.

Mason L et al. (2022) Stratifying the autistic phenotype using electrophysiological
indices of social perception. Science Translational Medicine 14:eabf8987.

Miltner WHR, Braun CH, Coles MGH (1997) Event-Related Brain Potentials Following
Incorrect Feedback in a Time-Estimation Task: Evidence for a "Generic" Neural
System for Error Detection. Journal of Cognitive Neuroscience 9:788–798.

Nahum M, Daikhin L, Lubin Y, Cohen Y, Ahissar M (2010) From Comparison to
Classification: A Cortical Tool for Boosting Perception. J Neurosci 30:1128–
1136.

Noel J-P, Balzani E, Acerbi L, Benson J, Laboratory TIB, Savin C, Angelaki DE (2024)
A common computational and neural anomaly across mouse models of
autism. :2024.05.08.593232 Available at:
https://www.biorxiv.org/content/10.1101/2024.05.08.593232v1 [Accessed July
16, 2024].

Noel J-P, Shivkumar S, Dokka K, Haefner RM, Angelaki DE (2022) Aberrant causal
inference and presence of a compensatory mechanism in autism spectrum disorder
Yu X, Shinn-Cunningham BG, Beierholm U, eds. eLife 11:e71866.

Noel J-P, Zhang L-Q, Stocker AA, Angelaki DE (2021) Individuals with autism spectrum
disorder have altered visual encoding capacity. PLOS Biology 19:e3001215.

Oesch LT, Thomas MC, Sandberg D, Couto J, Churchland AK (2025) Anterior cingulate
cortex mixes retrospective cognitive signals and ongoing movement signatures
during decision-making. Available at:
https://www.biorxiv.org/content/10.1101/2025.04.11.648398v3 [Accessed August
12, 2025].

Pagnozzi AM, Conti E, Calderoni S, Fripp J, Rose SE (2018) A systematic review of
structural MRI biomarkers in autism spectrum disorder: A machine learning
perspective. International Journal of Developmental Neuroscience 71:68–82.

Pedrosa V, Menichini E, Pajot-Moric Q, Vincent P, Zhou L, Teachen L, Latham P,
Akrami A (2023) Humans, rats and mice show species-specific adaptations to
sensory statistics in categorisation behaviour. :2023.01.30.526119 Available at:
https://www.biorxiv.org/content/10.1101/2023.01.30.526119v1 [Accessed July
16, 2024].

Postema MC et al. (2019) Altered structural brain asymmetry in autism spectrum disorder
in a study of 54 datasets. Nat Commun 10:4958.

Raviv O, Ahissar M, Loewenstein Y (2012) How Recent History Affects Perception: The
Normative Approach and Its Heuristic Approximation. PLOS Computational
Biology 8:e1002731.

Raviv O, Lieder I, Loewenstein Y, Ahissar M (2014) Contradictory Behavioral Biases
Result from the Influence of Past Stimuli on Perception. PLOS Computational
Biology 10:e1003948.

Regev TI, Markusfeld G, Deouell LY, Nelken I (2021) Context Sensitivity across
Multiple Time scales with a Flexible Frequency Bandwidth. Cerebral Cortex
Available at: https://doi.org/10.1093/cercor/bhab200 [Accessed October 7, 2021].

Robertson CE, Ratai E-M, Kanwisher N (2016) Reduced GABAergic Action in the
Autistic Brain. Current Biology 26:80–85.

Samaha J, Switzky M, Postle BR (2019) Confidence boosts serial dependence in
orientation estimation. Journal of Vision 19:25.

Sapey-Triomphe L-A, Pattyn L, Weilnhammer V, Sterzer P, Wagemans J (2023) Neural
correlates of hierarchical predictive processes in autistic adults. Nat Commun
14:3640.

Sapey-Triomphe L-A, Timmermans L, Wagemans J (2021) Priors Bias Perceptual
Decisions in Autism, But Are Less Flexibly Adjusted to the Context. Autism
Research 14:1134–1146.

Simms ML, Kemper TL, Timbie CM, Bauman ML, Blatt GJ (2009) The anterior
cingulate cortex in autism: heterogeneity of qualitative and quantitative
cytoarchitectonic features suggests possible subgroups. Acta Neuropathol
118:673–684.

Sokhadze E, Baruth J, El-Baz A, Horrell T, Sokhadze G, Carroll T, MD AT, PhD LS,
MD MFC (2010) Impaired Error Monitoring and Correction Function in Autism.
Journal of Neurotherapy Available at:

https://www.tandfonline.com/doi/abs/10.1080/10874201003771561 [Accessed
April 25, 2024].

South M, Larson MJ, Krauskopf E, Clawson A (2010) Error processing in high-
functioning Autism Spectrum Disorders. Biological Psychology 85:242–251.

South M, Newton T, Chamberlain PD (2012) Delayed reversal learning and association
with repetitive behavior in autism spectrum disorders. Autism Research 5:398–
406.

Stevens FL, Hurley RA, Taber KH (2011) Anterior Cingulate Cortex: Unique Role in
Cognition and Emotion. JNP 23:121–125.

Sysoeva OV, Constantino JN, Anokhin AP (2018) Event-related potential (ERP)
correlates of face processing in verbal children with autism spectrum disorders
(ASD) and their first-degree relatives: a family study. Molecular Autism 9:41.

Takeuchi D, Roy D, Muralidhar S, Kawai T, Bari A, Lovett C, Sullivan HA, Wickersham
IR, Tonegawa S (2022) Cingulate-motor circuits update rule representations for
sequential choice decisions. Nat Commun 13:4545.

Thakkar KN, Polli FE, Joseph RM, Tuch DS, Hadjikhani N, Barton JJS, Manoach DS
(2008) Response monitoring, repetitive behaviour and anterior cingulate
abnormalities in autism spectrum disorders (ASD). Brain 131:2464–2478.

Van de Cruys S, Evers K, Van der Hallen R, Van Eylen L, Boets B, de-Wit L, Wagemans
J (2014) Precise minds in uncertain worlds: Predictive coding in autism.
Psychological Review 121:649–675.

Vishne G, Jacoby N, Malinovitch T, Epstein T, Frenkel O, Ahissar M (2021) Slow
update of internal representations impedes synchronization in autism. Nat
Commun 12:5439.

Vlamings PHJM, Jonkman LM, Hoeksma MR, Van Engeland H, Kemner C (2008)
Reduced error monitoring in children with autism spectrum disorder: an ERP
study. European Journal of Neuroscience 28:399–406.

Wechsler D (2008) Wechsler Adult Intelligence Scale--Fourth Edition.

Wu D, José JV, Nurnberger JI, Torres EB (2018) A Biomarker Characterizing
Neurodevelopment with applications in Autism. Sci Rep 8:614.

Xu N, Li X, Zhong Y (2015) Inflammatory Cytokines: Potential Biomarkers of
Immunologic Dysfunction in Autism Spectrum Disorders. Mediators of
Inflammation 2015:531518.

Yizhar O, Fenno LE, Prigge M, Schneider F, Davidson TJ, O'Shea DJ, Sohal VS,
Goshen I, Finkelstein J, Paz JT, Stehfest K, Fudim R, Ramakrishnan C,

Huguenard JR, Hegemann P, Deisseroth K (2011) Neocortical
excitation/inhibition balance in information processing and social dysfunction.
Nature 477:171–178.

Zuk NJ, Murphy JW, Reilly RB, Lalor EC (2021) Envelope reconstruction of speech and
music highlights stronger tracking of speech at low frequencies. PLOS
Computational Biology 17:e1009358.