Genetic Architecture and Analysis Practices of Circulating Metabolites in the NHLBI Trans-Omics for Precision Medicine (TOPMed) Program

Nannan Wang[1,29], Franklin P. Ockerman[2,29], Laura Y. Zhou[3,29], Megan L. Grove[4], Taryn Alkis[1], John Barnard[5], Russell P. Bowler[6], Clary B. Clish[7], Shinhye Chung[1], Emily Drzymalla[8], Anne M. Evans[9], Nora Franceschini[10], Robert E. Gerszten[11], Madeline G. Gillman[12], Scott R. Hutton[9], Rachel S. Kelly[13], Charles Kooperberg PhD[14], Martin G. Larson[15], Jessica Lasky-Su[16], Deborah A. Meyers[17], Prescott G. Woodruff[18], Alexander P. Reiner[19], Stephen S. Rich[20], Jerome I. Rotter[21], Edwin K. Silverman[22], Vasan S. Ramachandran[23], Scott T. Weiss[24], Kari E. Wong[9], Alexis C. Wood[25], Lang Wu[26], NHLBI Trans-Omics for Precision Medicine (TOPMed) Consortium, Ronit Yarden[27], Thomas W. Blackwell[28], Albert V. Smith[28], Han Chen[1,30], Laura M. Raffield[12,30], Bing Yu[1,30,31]

[1]Department of Epidemiology, The University of Texas Health Science Center at Houston, Houston, TX, USA,
[2]Department of Biostatistics, University of North Carolina, Chapel Hill, NC, 27599, USA,
[3]Department of Biostatistics and Health Data Science, Indiana University School of Medicine, Indianapolis, IN, USA,
[4]Human Genetics Center, Department of Epidemiology, School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX, USA,
[5]Department of Inflammation and Immunity, Cleveland Clinic Lerner Research Institute, Cleveland, OH, USA,
[6]Department of Medicine, National Jewish Health, Denver, CO, USA,
[7]Division of Cardiovascular Medicine, Broad Institute of MIT and Harvard, Cambridge, MA, USA,
[8]Department of Epidemiology, University of North Carolina, Chapel Hill, NC, USA,
[9]Metabolon, Inc, Morrisville, NC, USA,
[10]Department of Epidemiology, University of North Carolina, Chapel Hill, NC, USA,
[11]Division of Cardiovascular Medicine, Broad Institute of MIT and Harvard, MA, USA,
[12]Department of Genetics, University of North Carolina, Chapel Hill, NC, USA,
[13]Channing Division of Network Medicine, Brigham and Women's Hospital, 181 Longwood Ave, Boston, MA, USA,
[14]Division of Public Health Sciences, Fred Hutchinson Cancer Center, Seattle, WA, USA,
[15]Department of Biostatistics, Boston University, Boston, MA, USA,
[16]Channing Division of Network Medicine, Brigham and Women's Hospital and Harvard Medical School, 181 Longwood Ave, Boston, MA, USA,
[17]Department of Medicine, University of Arizona College of Medicine, Tucson, AZ, USA,
[18]Department of Medicine, University of California, San Francisco, CA, USA,

[19]Department of Epidemiology, University of Washington, Seattle, WA, USA,

[20]Center for Public Health Genomics, Department of Public Health Sciences, University of Virginia School of Medicine, Charlottesville, VA, USA,

[21]Department of Pediatrics, The Institute for Translational Genomics and Population Sciences, The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance, CA, USA,

[22]Channing Division of Network Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA,

[23]Department of Quantitative and Qualitative Health Sciences, School of Public Health, University of Texas Health Science Center at San Antonio, San Antonio, TX, USA,

[24]Channing Division of Network Medicine, Harvard Medical School, Boston, MA, USA,

[25]USDA/ARS Children's Nutrition Center, Baylor College of Medicine, Houston, TX, USA,

[26]Cancer Epidemiology Division, Population Sciences in the Pacific Program, University of Hawaiʻi Cancer Center, University of Hawaiʻi at Mānoa, Honolulu, HI, USA,

[27]Epidemiology Branch, Division of Cardiovascular Sciences, National Heart, Lung, and Blood Institute, Bethesda, MD, 20817.

[28]Department of Biostatistics, University of Michigan, Ann Arbor, MI, 48109, USA.

[29]These authors contributed equally, Nannan Wang, Franklin P. Ockerman, and Laura Y. Zhou.

[30]Senior authors.

[31]Corresponding author.

Correspondence:
Bing Yu, PhD
Department of Epidemiology and Human Genetics Center, School of Public Health,
University of Texas Health Science Center at Houston, Houston, TX
1200 Pressler St., Suite E407, Houston, TX 77030
bing.yu@uth.tmc.edu

Abstract

Circulating metabolite levels partly reflect the state of human health and diseases, and can be impacted by genetic determinants. Hundreds of loci associated with circulating metabolites have been identified; however, most findings focus on predominantly European ancestry or single study analyses. Leveraging the rich metabolomics resources generated by the NHLBI Trans-Omics for Precision Medicine (TOPMed) Program, we harmonized and accessibly cataloged 1,729 circulating metabolites among 25,058 ancestrally-diverse samples. We provided recommendations for outlier and imputation handling to process metabolite data, as well as a general analytical framework. We further performed a pooled analysis following our practical recommendations and discovered 1,778 independent loci associated with 667 metabolites. Among 108 novel locus-metabolite pairs, we detected not only novel loci within previously implicated metabolite associated genes, but also novel genes (such as *GAB3* and *VSIG4* located in the X chromosome) that have putative roles in metabolic regulation. In the sex-stratified analysis, we revealed 85 independent locus-metabolite pairs with evidence of sexual dimorphism, including well-known metabolic genes such as *FADS2*, *D2HGDH*, *SUGP1*, *UTG2B17*, strongly supporting the importance of exploring sex difference in the human metabolome. Taken together, our study depicted the genetic contribution to circulating metabolite levels, providing additional insight into the understanding of human health.

Main

Metabolites, small molecules produced during metabolism, play a pivotal role in cellular and physiological processes. They are widely used as clinical biomarkers, as their observed levels in biofluids can reflect physiological and pathophysiological processes. Many metabolites are highly heritable[1,2], indicating a strong genetic influence. Investigation of genetic associations with metabolite levels may help improve our understanding of the pathways involved in diseases and health. Recently, hundreds of metabolite quantitative trait loci (metQTLs) have been reported from genome-wide association studies[3–8]. However, most studies are focused on one population (predominantly of European ancestry), a single study or a single metabolomic platform. Additional work integrating metabolite and genetic information across diverse studies and populations to broaden scope for metQTL discovery and improve generalizability is required.

The National Heart Lung and Blood Institute (NHLBI) Trans-Omics for Precision Medicine (TOPMed) program aims to provide information that allows the personalization of disease treatments based on an individual's unique genes and environment. TOPMed is generating a rich resource of metabolomic data across multiple studies at two core laboratories. The inherent complexity of metabolite analysis and the diversity of methodologies employed across different laboratories underscore the importance of the harmonization of metabolites. Further, systematically cataloging metabolites can provide scientifically rigorous information for future studies for elucidating complex metabolic networks and unraveling their implications in health, diseases, and environmental interaction. Our study collected and harmonized the circulating metabolites measured from eight studies at the following two TOPMed Centralized Omics Resource Core (CORE) centers: 1.) Baylor-UTHealth Metabolomics Center with Metabolon, Inc. and 2.) Broad-Beth Israel Deaconess Medical Center; hereafter referred to as the Global Discovery Panel and Broad-BIDMC Panel, respectively. l The resulting metabolite catalog is publicly available on the TOPMed website (Fig. 1) and as a manuscript supplemental table.

Given the broad spectrum of metabolomic data from multiple studies released by TOPMed, investigators face challenges in pre-processing and integrating data across multiple studies. Consistent protocols for pre-processing and integration are integral to maximize the potential of this resource. Here we utilized TOPMed metabolomics and whole-genome sequencing (WGS) data from multiple studies to provide practical recommendations for metabolite data pre-processing and to identify novel metQTLs across platforms and studies. We compared the impacts of various processing steps, i.e., outlier handling, missingness imputation, as well as study pooled or stratified analysis strategies, using age and sex as example analysis phenotypes (Fig. 1).

WGS based associations with the metabolome have been reported[9–11]. Most previous studies focused on predominantly European ancestry studies. Some existing studies are focused on one population (predominantly of European ancestry)[3,6,9,11], or a single study[8]. For studies including multiple studies, meta-analysis is mostly used to account for study-specific covariates and overcome data sharing and privacy issues[4]. Moreover, research suggests that strong

differences may exist between males and females in the incidence or severity of diseases, metabolism, and reaction to medications[12-14]. However, sex-stratified analysis on metQTLs is still limited. Following the above practice recommendations for pre-processing data, we conducted association analyses using WGS to investigate the associations with 1,135 metabolites from eight studies profiled across two metabolomic platforms. We also systematically assessed sex differences in the genetic regulation of the circulating metabolites (Fig. 1).

Results

Catalog of circulating metabolites

We analyzed metabolite data from 25,058 participants from case-control and population-based studies, including 15,648 participants from three studies profiled on the Global Discovery Panel (i.e., Women's Health Initiative [WHI], The Subpopulations and Intermediate Outcomes in Chronic Obstructive Pulmonary Disease Study [SPIROMICS], The Genetic Epidemiology of Chronic Obstructive Pulmonary Disease [COPDGene]), and 9,426 participants from five studies profiled on the Broad-BIDMC panel (i.e., Women's Health Initiative [WHI], Framingham Heart Study [FHS], Childhood Asthma Management Program [CAMP], Genetic Epidemiology of Asthma in Costa Rica [CRA], and Multi-Ethnic Study of Atherosclerosis [MESA]). Table 1 summarizes the demographic characteristics and number of participants from each study with available plasma metabolites funded through TOPMed, and participants included for WGS analysis. Overall, samples with metabolites profiled by Broad-BIDMC from WHI had the oldest mean age of 80.0 years with standard deviations (SD) being 7.5, while samples from CAMP had the youngest mean age of 13.0 years with SD being 2.2. All samples with metabolites profiled by the Global Discovery Panel had similar mean ages, ranging from 63.2 years to 66.3 years. Across the eight studies, we identified 1,729 named metabolites, which are summarized in Fig. 2. There were 366 metabolites identified on both the Broad-BIDMC and the Global Discovery Panels. RefMet annotations showed that named metabolites mostly belonged to organic acids (14%), fatty acyls (21%), glycerolipids (11%), glycerophospholipids (7%), sterol lipids (7%), and nucleic acids (4%). Additional information about studiesis described in Supplementary Materials. The catalog is provided in Supplementary Table 1, including the HMDB ID, metabolite names based on the latest Metabolon metabolite library and Broad metabolite library, RefMet name, and annotations provided by Metabolomics Workbench.

Fig. 1 Workflow for comparison of outlier, imputation, and QTL analysis methods. Gold boxes indicate recommended outlier management, imputation, and QTL analysis methods. Abbreviations:

CAMP = Childhood Asthma Management Program; CRA = Genetic Epidemiology of Asthma in Costa Rica; MESA = Multi-Ethnic Study of Atherosclerosis; COPDGene = Genetic Epidemiology of COPD; SPIROMICS = SubPopulations and InteRmediate Outcome Measures in COPD Study; WHI = Women's Health Initiative; INT = inverse normal transform. Created with Biorender.com. (*) The WHI study has data on both metabolomics platforms. Note that the BCM/UTH core utilized the Global Discovery Panel.

Table 1 Study samples with available plasma metabolite data from TOPMed.

Study Name	Study Abbreviation	dbGaP ID	Study Samples*	N with Metabolomics	Female (%)	Age (mean ± sd)
Broad-BIDMC Panel						
The Childhood Asthma Management Program	CAMP	phs001726	Overall	960	350 (36)	12.96 (2.15)
			WGS	787	315 (40)	13.01 (2.14)
The Genetic Epidemiology of Asthma in Costa Rica	CRA	phs000988	Overall	2,040	918 (45)	16.38 (14.40)
			WGS	1,758	806 (46)	16.78 (14.67)
Framingham Heart Study	FHS	phs000974	Overall	3,025	1,629 (54)	61.74 (15.15)
			WGS	3,021	1,626 (54)	61.76 (15.15)
Multi-Ethnic Study of Atherosclerosis **	MESA	phs001416	Overall	2,000	1,064 (53)	64.98 (10.72)
			WGS	998	530 (53)	60.22 (9.68)
Women's Health Initiative	WHI	phs001237	Overall	1,400	1,400 (100)	80.04 (7.54)
			WGS	1,024	1,024 (100)	81.69 (6.33)

Metabolon Global Discovery Panel						
Genetic Epidemiology of COPD	COPDGene	phs000951 phs000946	Overall	9,218	4,431 (50)	65.93 (9.04)
			WGS	6,302	3,121 (50)	64.70 (8.86)
SubPopulations and InteRmediate Outcome Measures in COPD Study	SPIROMICS	phs001927	Overall	3,417	1,602 (47)	63.23 (9.10)
			WGS	1,924	888 (46)	63.32 (9.13)
Women's Health Initiative	WHI	phs001237	Overall	2,998	2,998 (100)	66.33 (9.68)
			WGS	545	545 (100)	63.65 (7.61)

*Study samples in the row of "Overall" have available metabolite data, and those in the row of "WGS" have meta-data (i.e. age and sex) and existing WGS and were included for subsequent analyses. Average age and percent females in Overall study samples was calculated regardless of missing, WGS study samples had complete age and sex.

**The overall sample size in MESA included participants from visit 1 and visit 5. Only samples from visit 1 were included for WGS analysis.

Fig.2 Named metabolites identified across two metabolomic cores, Broad-BIDIMC panel and the Metabolon Global Discovery Panel, and eight studies. A) Venn diagram of named metabolites measured between platforms; B) Refmet super classes that the named metabolites belong to; and C) the number of named metabolites measured by each study.

Comparison of outlier strategies

For the comparison of outlier and imputation methods, we used pooled analyses to assess associations with age and sex or metQTLs (for imputation methods) to reduce computational burden, and in concordance with prior work using TOPMed generated data[10], combining data across study sites. However, all outlier handling strategies and missing data imputation (for example, deciding what value is considered to be minimum or median) was done within studies. To compare the outlier methods in the pooled data, we used complete-case analysis (per-metabolite), by removing missing values (distribution of missingness by study summarized in Supplementary Table 2). For this analysis, we removed any related individuals and removed any metabolites that were only measured in one study(Supplementary Table 4). Fig. 3A-C compares the parameter estimates of age using removal of values more than six standard deviations away from the mean metabolite value (6SD), rank-based inverse normal

transformation (INT), and winsorization outlier management methods. The same comparison for metabolite-sex associations is provided in Supplementary Figure 1. Agreement of significance of the parameter estimates is indicated by color, using Bonferroni adjusted *p*-values. All three outlier strategies have very similar estimates for age and sex (minimum correlation > 0.999). Likewise, test statistics for age-metabolite and sex-metabolite associations are highly concordant between outlier strategies (Supplementary Figure 2, Supplementary Table 3). Due to the advantages of normalizing the data, notably inclusion of extreme values which may be biologically plausible as opposed to dropping subjects altogether, and favorable statistical properties for parametric modeling methods, we continued with missing imputation using INT for the outlier management.

Fig. 3 Comparison of age-metabolite associations for outlier handling and data pooling strategies. Parameter estimates for age-metabolite associations are compared for the three outlier handling strategies (A-C) and for the two data integration strategies (D).

Comparison of pooled vs. meta-analysis

Using INT transformation for outlier management and complete-case analysis, we compared pooled versus study-stratified meta-analysis for age and sex. As in the previous analysis, we removed related individuals and metabolites measured in only one study. Fig. 3D summarizes the effect estimates in the pooled analysis versus the meta-analysis for age for each metabolite and Supplementary Figure 1 summarizes the results for sex. Agreement of significance of the parameter estimates is indicated by color, using Bonferroni adjusted *p*-values. In both metabolite age and sex analyses, the parameter estimates and significance have high agreement (correlation 0.998 and > 0.999 for age and sex, respectively). Full association results are reported in Supplementary Table 5. Our results demonstrate that in most circumstances, pooled and meta-analysis, after inverse normalization by study, give similar results. Our results

recapitulate many previously reported associations with age and sex, such as the association of increased leucine (beta = 0.517, $p = 4.83\times10^{-202}$), carnitine (beta = 0.250, $p = 8.12\times10^{-48}$), and nicotinamide (beta = 0.129, $p = 6.31\times10^{-15}$) with age[15,16]. We also find higher levels of glycine in females (beta = -0.380, $p = 4.41\times10^{-109}$) and higher levels of creatinine in males (beta = 0.707, $p = 0$)[17].

Comparison of metabolite imputation strategies

Literature-validated metQTLs were selected from three large broad-spectrum metabolomics GWAS studies[3,11,18], allowing for a large number of assumed positive control signals for testing of eight missing data management strategies. Analysis was restricted to metabolites with > 1% missingness in at least one study. In the Broad-BIDMC Panel, 85 literature metQTLs were selected for 26 metabolites. In the Global Discovery Panel, 690 metQTLs were selected for 188 metabolites. Imputation methods, followed by rank-based inverse normal transformation, were applied per-study before conducting association analyses. metQTL analysis was performed with a mixed model approach (GMMAT), adjusting for age, sex, and study(for metabolites found in multiple studies).

Results for the number and percent of Bonferroni-corrected significant metQTLs detected for each imputation method, within bins of metabolite missingness, are reported in Table 2. Overall, a maximum of 53/85 metQTLs were validated in the Broad-BIDMC panel for the best-performing imputation methods (k-nearest neighbors (kNN), no imputation). In the Global Discovery Panel, a maximum of 512/690 QTLs were validated in the best-performing method (median), a 2.6% improvement over no imputation (499 metQTLs). Of particular interest is imputation quality for metabolites with the highest degree of missingness. Among metabolites with 15-50% missingness, we observed the highest number of QTL-metabolite associations for single value imputation and left-imputation methods (zero, half-min, median, Quantile Regression for Imputation of Left-Censored data (QRILC)[19] in the Metabolon dataset. However, the total number of significant associations was largely consistent between imputation methods. The distribution of test statistics for metQTL associations is provided in Supplementary Figure 3. A pairwise comparison of these test statistics between each imputation method reveals very similar test statistics for methods which do not assume missing not at random (MNAR) (Random Forest (RF)[20], Mechanism-Aware Imputation (MAI)[21], kNN), and similar test statistics for left imputation methods (zero, half-min, QRILC). Full association results are reported in Supplementary Table 6.

A comparison of test statistics for imputed metabolite-age (A) and imputed metabolite-sex (B) associations is reported in Supplementary Figure 4. While the distributions per-imputation method are highly similar overall, we observe greater magnitude in the test statistics for left-imputation methods (zero, half-min, QRILC) relative to methods which do not assume MNAR (RF, MAI, kNN). Full association results are reported in Supplementary Table 7.

Table 2 Comparison of quantitative trait loci results by missingness strategy. In the Broad-BIDMC Panel, 85 and 690 literature metQTLs were selected to validate the imputation methods for the Broad-BIDMC Panel and the Global Discovery Panel. The number and percent of Bonferroni-corrected significant metQTLs detected for each imputation method, within bins of metabolite missingness are reported.

Platform	Imputation	Missingness			
		All	0-5%	5-15%	15-50%
Broad-BIDMC Panel	None	53 (62.4%)	40 (72.7%)	7 (30.4%)	6 (85.7 %)
	Zero	52 (61.2%)	38 (69.1%)	9 (39.1%)	5 (71.4%)
	Half-min	52 (61.2%)	38 (69.1%)	9 (39.1%)	5 (71.4%)
	Median	52 (61.2%)	38 (69.1%)	7 (30.4%)	7 (100%)
	QRILC	50 (58.9%)	38 (69.1%)	9 (39.1%)	3 (42.9%)
	kNN	53 (62.4%)	38 (69.1%)	9 (39.1%)	6 (85.7 %)
	RF	51 (60%)	38 (69.1%)	6 (26.8%)	7 (100%)
	MAI	52 (61.2%)	39 (70.9%)	6 (26.8%)	7 (100%)
Global Discovery Panel	None	499 (72.3%)	270 (72.4%)	180 (75.6%)	49 (62.0%)
	Zero	508 (73.6%)	276 (74.0%)	182 (76.5%)	50 (63.3%)
	Half-min	508 (73.6%)	276 (74.0%)	182 (76.5%)	50 (63.3%)
	Median	512 (74.2%)	276 (74.0%)	183 (76.9%)	53 (67.1%)
	QRILC	509 (73.8%)	276 (74.0%)	183 (76.9%)	50 (63.3%)
	kNN	500 (72.4%)	271 (72.7%)	178 (74.8%)	51 (64.6%)
	RF	503 (72.9%)	274 (73.5%)	179 (75.2%)	50 (63.3%)
	MAI	496 (71.9%)	273 (73.2%)	176 (73.9%)	47 (59.5%)

Pooled metQTL scanning

We performed a pooled metQTL scanning in our studies, analyzing a total of 16,518,857 variants (Minor Allele Frequency; MAF>0.5%) on 22 autosomal chromosomes and the X chromosome for associations with 1,135 metabolites (Supplementary Figure 5). A total of 147,166 single variant-metabolite associations were identified to be significant ($P \leq 4.4 \times 10^{-11}$),

including 51,866 unique variants covering 667 metabolites. These associations were further grouped into 1,775 conditionally independent locus-metabolite pairs (993 loci, 667 metabolites; see Methods, Supplementary Figure 5).

To identify novel associations from these 1,775 conditionally independent locus-metabolite pairs, we further excluded 55 association pairs in unannotated metabolite pathways. Among the remaining 1,720 pairs,1,612 pairs were previously reported or already detected in available public databases (Supplementary Table 8) and 108 locus-metabolite pairs were novel, including 77 loci and 81 metabolites (Supplementary Table 9). Among 108 independent novel pairs, 82 pairs were available for replication (Methods), and 80% of loci-trait pairs (66/82) (including 51 unique loci and 51 metabolites) were successfully replicated ($P \leq 6.10 \times 10^{-4}$; Supplementary Figure 5). We then annotated these novel loci with their closest genes. Among them (for lead variants and all significant variants within 500 kb), 1% were classified as missense variants, 5% were in 5 prime UTR regions, 46% were in downstream and upstream regions and others were in non-coding regions (Supplementary Figure 6).

The circular plot in Figure. 4 showed the genomic location of significant associations with metabolites. The genes closest to the novel locus-metabolite pairs were labeled. Of the 77 loci from 108 independent novel association pairs, we observed that 20.8% of loci (16/77) were associated with multiple metabolites, either within the same super-pathways or across super-pathways. Given the one-to-many associations between genes and metabolites, we classified these loci into two categories (Figure.4; Supplementary Table 10). The first category includes 26 loci (labeled in red color) associated with both reported sub-pathways and unreported sub-pathways in our study. For example, for the known metabolic gene *CD36*, we reproduced a known association of rs3211938 with plasmalogen[9], and also discovered novel associations with sphingomyelins and ceramides. In addition, we detected rs1047891 located in gene *CPS1* that was associated with 22 known metabolites in previous studies[1,3,5–7,9,10,18,22–24]. Meanwhile, we discovered novel associations of *CPS1* with branched-chain fatty acids and partially characterized molecules (Supplementary Table 10). For loci in the second category (labeled in blue color), we only identified novel metabolite associations. For example, the missense variant rs148051057 located in the *RGN* gene was associated with gluconate level (replicated p-value 4.82×10^{-10} in Feofanova et al.[10]).

Interestingly, we identified 50 novel association pairs from chromosome X. The high percentage of novel pairs (46.3%, 50/108) annotated on chromosome X is in part due to the lack of replication using public databases, as chromosome X variants were not analyzed in many prior studies. The rs112701110 variant near the *MID1IP1* gene was not only associated with the well known guanidino and acetamido levels[10], but also had novel associations with leucine, isoleucine, and valine metabolism and methionine, cysteine, SAM, and taurine metabolism pathway metabolites, among others etc. Novel association between rs867460874 in the *VSIG4* with long chain monounsaturated fatty acid were identified (replication p-value 3.31×10^{-4}). In addition, seven novel loci located in the *GAB3* gene were associated with hemoglobin/porphyrin metabolism and fatty acid metabolism, among which six associations have been replicated (Supplementary Table 10).

Prior work suggested potential links between metQTLs and traits/diseases[25,26], we therefore performed genotype-phenotype scanning to understand their possible mechanisms (Fig. 5). The significant loci were not only detected in complex traits such as height, systolic blood pressure, high density lipoprotein, total cholesterol, but also overlapped in fat-free mass (including whole body fat-free mass, leg-free mass). Regarding the disease category, the top three enriched diseases are malabsorption/coeliac disease, peripheral vascular disease, and coronary artery disease. The metQTLs related to peripheral vascular disease are from one independent locus(with 14 significant variants) associated with diacylglycerol (DAG) level, including one missense variant rs3812316 located in the *MLXIPL* gene and rs13232120 located in the 3'-*UTR* of *TBL2* (Supplementary Table 11), indicating shared genetic determinants of DAG-protein kinase C (PKC) pathway and vascular function[27].

Sex-stratified metQTL scanning

For the sex-stratified metQTL scanning, 2,453 variant-metabolites associations were identified to show dimorphism by sex and 85 locus-metabolite pairs were identified to be conditional independent (Supplementary Figure 7; Supplementary Table 12). These independent sex-stratified loci were classified into three categories: 1) significant in female only (44.7%, 33/85); 2) significant in male only (29.4%, 25/85), and 3) significant in both male and female but with different effect sizes (31.8%, 27/85) (Figure. 6a-b; Supplementary Figure 8).

We observed that rs1047891 located in the *CPS1* gene had the most sex-stratified association pairs, including ten amino acid metabolites, seven lipid metabolites, and one metabolite from partially characterized molecules, peptides, and xenobiotics, separately (Figure. 6c). All of them show higher levels in females than males. Importantly, 12 associations were significant in females only, i.e., betaine, creatine, threonine, serine etc. Among the 20 sex-stratified associations on *CPS1*, 16 associations have been reported in Wittemans et al., 2019[28]. For example, consistent with Wittemans et al., 2019[28], the effect of the *CPS1* on isobutyrylglycine is nearly 5-fold stronger in females than males ($\hat{\beta}_{female} \pm se_{female} = 0.33 \pm 0.02$, $\hat{\beta}_{male} \pm se_{male} = 0.06 \pm 0.02$, $P_{diff} = 7.49 \times 10^{-16}$). The four metabolites associated with *CPS1* gene with dimorphism by sex were trans-2-hexenoylglycine, 2-Butenoylglycine, cis-3,4-methyleneheptanoylglycine, and 2R,3R-dihydroxybutyrate with 2-fold, 2-fold, 3-fold, and 7-fold stronger in females than males separately.

We observed additional 21 association pairs that had significant effects on females only (Supplementary Figure 8A). One example is the association of rs625978 located in *GSTP1* with N-acetylglycine level (Fig. 6d; $\hat{\beta}_{female} \pm se_{female} = -0.16 \pm 0.02$, $\hat{\beta}_{male} \pm se_{male} = -0.03 \pm 0.02$, $P_{diff} < 2.53 \times 10^{-4}$). We also identified associations for well-known metabolic genes such as *FADS2* (rs2526678 with 1,2-dillnoleoyl-GPC), *D2HGDH* (rs1106639 with 2-hydroxyglutarate), *SULT2A1* (rs2547235 with 5alpha-androstan-3beta), *ASMTL* (rs11553055 with gualacol sulfate), and *ACADL* (rs1509567 with nonanoylcarnitine) etc.

There were 25 association pairs that only showed significant effects in males (Supplementary Figure 8B). We observed the rs34707604 in *UTG2B17* associated with three metabolites of lipid androgenic steroids, including androsterone sulfate (Fig. 6f), 5alpha-androstan-3alpha,17beta-diol monosulfate (1), and androstenediol (3alpha, 17alpha) monosulfate (3). Its effect on androsterone sulfate was nearly 4-fold stronger in males than females ($\hat{\beta}_{female} \pm se_{female} = -0.07 \pm 0.03$, $\hat{\beta}_{male} \pm se_{male} = -0.30 \pm 0.03$, $P_{diff} < 5.3 \times 10^{-9}$). Similarly, there were male-specfic associations for well-known metabolic genes such as *PTPRN2* (rs73510583 with propyl 4-hydroxybenzoate sulfate), *UGT2B17* (rs34707604 with androsterone sulfate), *SUGP1* (rs8100204 with Cer 18:1;O2/24:0), *ABCC1* (rs2062541 with glutarylcarnitine (C5-DC)), *UGT3A1* (rs6889699 with 5alpha-pregnan-3beta).

There were four sex-stratified loci located on the X chromosome. The top significant signal was rs34805104 located in *TMLHE-SPRY3* associated with N6-trimethyllysine level, with the effect size estimate in males was nearly 2-fold of that in females ($\hat{\beta}_{female} \pm se_{female} = -0.21 \pm 0.02$, $\hat{\beta}_{male} \pm se_{male} = -0.43 \pm 0.03$, $P_{diff} = 2.02 \times 10^{-12}$).

Fig. 4 Circular plot illustrating the genomic location of significant associations with metabolites from metabolite pooled analyses. Metabolites were classified into circular bands, and each colored section represented one assigned major pathway class. Significant associations were indicated by black points. The genes closest to the novel locus-metabolites associations were labeled. The genes were classified into two categories: the gene names labeled in red indicated that there were both reported sub-pathways and novel sub-pathways associated with the gene in our study, and the gene names labeled in blue indicated that the gene was only associated with unreported sub-pathways.

Fig. 5 Phenome-wide associations for novel metabolite loci from the pooled analysis. Traits and diseases available in PhenoScanner (version 2) database were analyzed. The scanning was performed on 7,300 novel significant associated pairs (from 6,500 variants) detected in the pooled analysis (from 108 locus-metabolite pairs). For each category, only top 25 phenotypes were shown. The y-axis denoted the number of variant hits (with the logarithmic scale) from the PhenoScanner database.

Fig. 6 Significant associations identified from sex-stratified analyses. a, Scatter plot showing the female- and male-specific association P-values from sex-stratified analysis, the associations are significant for female only (top-left) or male only (bottom-right), **b**, Scatter plot showing significant associations in both males and females but with different effect size level. Each dot denoted one association pair labeled with the rsID and colored by the super pathway of the associated metabolite. Four association pairs with *P* values out of the range (i.e, 0-100)

were not shown in **b**), including rs1047891 associated with glycine, N-Acetylglylycine, gamma-Glutamylglycine, and rs7907555 with N2-acetyl,N6-dimethyl lysine. **c**, Sex-stratified per-allele effect size of rs1047891 (in *CPS1*) on metabolite levels. Metabolites not detected in Wittemans et al., 2019[28] were highlighted with the star symbol. **d-f** Miami-Plot showed available sex-specific association P-value and for trans-2-hexenoylglycine (**d**), cholesterol sulfate (**e**), and androsterone sulfate (**f**) from sex-stratified analyses. The genes mapped by proximity to the sex-specific associations were colored in green.

Discussion

In this study, we harmonized and cataloged circulating metabolites from eight TOPMed studies. We demonstrated one set of reasonable strategies for both genetic and epidemiological association analyses within the TOPMed program, concerning methods for handling outliers, missing data, and pooled versus meta-analysis. We expect these recommendations will prompt increasing use of the valuable TOPMed metabolomics resources. We also conducted a WGS study to detect genetic association with 1,135 circulating metabolites. Among 108 novel locus-metabolite pairs, we detected not only novel associations for known metabolic genes, but also novel genes that have putative roles in metabolic regulation based on prior literature. The sex-stratified analysis also provided a systematic, large-scale examination of sex difference in the genetic regulation of the circulating metabolites.

Metabolite overlap across five metabolomic platforms (the Global Discovery Panel, the Broad-BIDMC Panel, Nightingale Health, Biocrates, and West Coast Metabolomics) has previously been assessed by COMETS[29]. Between the two major metabolite platforms, there were 1,155 metabolites measured by the Global Discovery Panel, 350 metabolites measured by Broad-BIDMC Panel, and 111 overlapping metabolites identified using names and identifiers from the Human Metabolome Database, Pubchem, and Chemspider. In our current assessment of metabolite overlap across eight TOPMed studies, we identified 1,729 total metabolites with known identities and 366 overlapping metabolites between the Global Discovery Panel and Broad-BIDMC Panel. The number of identified metabolites overlapping between these two popular metabolite platforms were more than tripled with the latest metabolite libraries used in TOPMed. Our current work has harmonized additional metabolites that previously may not have been measured or had a known identity, expanding the scope of metabolomic research.

With technological advances of liquid chromatography combined with mass spectrometry (LC-MS), the collection of metabolomic data has increased. However, LC-MS generated data have missingness. These missingness could be true missing (i.e. metabolite is absent from the biological samples, true concentration of zero), missing due to technical factors (such as co-elution), or could be the level is below the detection limit. There are two main approaches to handle missingness, (1) to impute the data and use statistical methods that require complete data or (2) use statistical methods that can account for missingness without imputation. In the first approach, previous studies have compared different methods to handle or impute the missingness. Kokla et al.[30] simulated different missingness mechanisms, i.e. MCAR, MNAR, and MAR, and compared imputation methods including zero, mean, minimum value, half

minimum, singular value decomposition, probabilistic principal component analysis, Bayesian principal component analysis, random forest, and k-nearest neighbors. In the MCAR and MAR scenarios, RF and KNN performed the best. For our analysis, assuming that previously reported QTLs from GWAS studies in non-TOPMed studies and age and sex associations were true positives, we found that half min performed similarly to RF or KNN. In the case where there is high missingness (15-50%), we found that KNN, RF, and MAI have the best performance. MAI is also the recommended method by the Metabolomics Workbench[31]. Other studies also concluded that RF tended to outperform other methods in performance[32]. A similar analysis conducted by Wei et al. 2018[19] found that RF imputation performed well for MAR and MCAR, but poorly when the data was MNAR. For MNAR, they recommended quantile regression for imputation of left-censored data (QRILC), which we also examined. In the second approach, imputation-free methods have been previously compared with standard imputation methods and the results indicated that no imputation-free method was consistently better[31]. We find very few meaningful differences across imputation methods for Broad-BIDMC panel, and some improvements with left-imputation methods for the Global Discovery Panel. There may be differences in proportion of missing data that is MAR versus MNAR between the two platforms, and we note that there are both many more named metabolites and many more named metabolites with high missingness for the Global Discovery Panel, which may impact our comparisons. In both the Global Discovery Panel and Broad-BIDMC Panel, our results suggest that simpler approaches may provide comparable performance to more complex methods (MAI, kNN, etc). Accounting for the computation burden and technical complexity, we have recommended the use of half minimum value for most TOPMed analyses; this approach was utilized in the genome-wide metabolite QTL associations provided here.

Multi-study and integrative metabolomic analyses are often conducted via a meta-analytic approach[33–36], where association studies are conducted per-study, and effect sizes or p-values are combined. Meta-analyses can account for study-specific considerations and overcome many privacy issues where data cannot be shared and pooled in one analysis location. However, meta-analyses may also have bias and lower power to detect individual-level interactions[37]. Particularly in the scope of rare-variant analysis, meta-analysis has many challenges. Pooling data has been suggested as an alternative approach to meta-analysis[38]. Pooled data increases the sample size, has greater flexibility to control confounders and avoid within-study assumptions. Pooling data allows for better ability to detect rare outcomes in an efficient manner, especially if the outcome is low in individual studies, for example when assessing associations of rare genetic variants. However, the pooling analysis requires the data to be homogenous and more post-pooling cleaning, formatting, and resources for analysis. Feofanova et al. 2023[10] pooled five studies, including MESA, and metabolomic platforms with up to 1666 circulating metabolites and performed whole-genome sequencing association analysis. They successfully identified 75 novel and replicated metabolite-genetic locus associations. Their pooled data identified novel rare-variant gene pairs, while effectively controlling for genomic inflation. Additionally, they were able to extend metabolite genetic association discovery to multi-ethnic populations[39]. We similarly show the feasibility and power of a study-pooled analytic approach with appropriate outlier management via inverse normalization by study prior to analysis.

In our pooled WGS scanning, 93.7% (1612/1720) of associations overlapped with findings from previously reported studies, demonstrating the high reliability of metQTLs. For example with association scanning on glycine, we identified four significant loci located in genes *CPS1*, *ALDH1L1*, *PSPH*, and *GCSH* (Supplementary Figure 9). Remarkably, all these four genes are essentially involved in the glycine metabolism pathway[28]. We were also able to reproduce well known signals from the sex-stratified analysis. For example, about 80% of metabolites showing female specific associations for rs1047891 locating in *CPS1* in our study have been detected in a previous study[28].

Among the 66 novel and replicated pairs in the pooled analysis, nearly half of these detected involved the X chromosome, which has not been included in many prior genetic analysis efforts. For example, five novel and replicated pairs were identified between the *GAB3* locus and bilirubin, biliverdin, cerotoylcarnitine (C26), palmitoylcarnitine (C16), and stearoylcarnitine (C18). Cerotoylcarnitine (C26), palmitoylcarnitine (C16), and stearoylcarnitine (C18) are acylcarnitines involved in fatty acid metabolism. The *GAB3* gene encodes the protein that binds to the growth factor receptor-bound protein 2 (*GRB2*). *GRB2* plays a central role in signaling by receptor protein-tyrosine kinases (RTKs)[40,41], which respond to the FGF receptor family to affect the fatty acid synthesis[42]. Our association scanning indicates the putative role of *GAB3* on controlling lipid metabolism by the mediation of RTK ligands. Another novel association of rs867460874 in the *VSIG4* locus with myristoleate (14:1n5) in the sub-pathway of long chain monounsaturated fatty acid was identified. *VSIG4*, a B7 family-related protein expressed by resting macrophages, is able to regulate pyruvate/acetyl-CoA conversion by activating the PI3K/Akt-STAT3 pathway[43]. Since pyruvate and acetyl-CoA are important intermediaries in the conversion of carbohydrates into fatty acids and cholesterol[44], this may reveal the putative link of the *VSIG4* gene on lipid metabolism.

The influence of sexual dimorphism is gaining mounting interest in human metabolomic research[45,46]. We detected that *GSTP1* only showed female-specific effects on N-acetylglycine level. This gene encodes glutathione S-transferase enzymes and plays an important role in glutathione metabolism and drug metabolism[47]. Polymorphisms in this gene have been reported to be associated with male infertility[48], supporting the potential sex-stratified role of *GSTP1* on metabolite level. *UGT2B17* only shows male-specific effects on lipid androgenic steroid levels. This gene encodes an enzyme for glucuronidation of androgens and their metabolites, which is a major source of estrogen[49,50]. Higher concentration of dehydroeplandrosterone sulfate in male participants has been reported[49], further supporting the potential sex-stratified role of *UTG2B17* on androsterone sulfate level. Furthermore, we detected well-known metabolic genes, such as *CPS1, FADS2, D2HGDH*, and *SUGP1*, showing sex-stratified effects on multiple metabolite levels, which requires further investigation. Taken together, our study strongly supports the importance of sex-stratified analysis for studying the underlying mechanisms of sex difference in human metabolomes.

TOPMed has a variety of strengths and weaknesses versus other large-scale efforts to profile the circulating metabolome. TOPMed is a programof many studies, with centralized

metabolomic cores, but with varying measured phenotypes and recruitment strategies. COMETS has a very large total sample size (>136,000 samples with metabolomics, from >47 studies) but due to widely varying study design and laboratory methods and data sharing restrictions, conducting pooled analyses is challenging as compared to the TOPMed program[29]. UK Biobank has generated metabolomics measures in an extremely large sample size, as have other biobanks such as Estonian Biobank and Finnish THL Biobank[51], on the targeted Nightingale platform. This platform has advantages, particularly in terms of scalability and cost, but covers a narrow slice of the circulating metabolome, mostly from amino acids and lipids[52]. Untargeted metabolomic platforms have been used in TOPMed, as well as the inclusion of many studies with diverse participants from across the lifecourse, which enables the capture of a variety of metabolites from different pathways for a comprehensive assessment of impact from human metabolomes. TOPMed data generation, including longitudinal data generation, is ongoing, and sample sizes in diverse TOPMed studies will continue to accrue. Pooled analysis makes it more feasible for certain types of analysis strategies (such as aggregate tests for rare variants, as in previous TOPMed work using study generated data[10]). Our results show a feasible strategy for conducting cross-study analyses for the two different platforms in TOPMed.

Future analysis will also work to integrate both TOPMed generated and study generated metabolomics data (as in Feofanova 2023[10]) to promote the discovery of metQTLs, from both single variant and aggregate rare variant tests. We will also work to incorporate our analytical recommendations and to extend our harmonization efforts to additional metabolomic samples funded by TOPMed (with metabolomic quantification for >90,000 samples, including longitudinal measurements, across 13 studies currently funded). Our current work has focused on named metabolites; additional insights may be gained by examining unnamed compounds, with genetic analysis potentially providing clues to metabolite identity[9,53,54]. Such work will further elucidate our understanding of the genetic and epidemiological determinants of the circulating metabolome in diverse US populations.

In summary, we here describe the first set of TOPMed generated metabolomics data, across two harmonized data generation platforms (the Global Discovery Panel and Broad-BIDMC Panel) and eight diverse studies. A metabolite catalog was created and released on the TOPMed website and as supplementary data with this publication. We evaluate the impact on results of multiple common analysis strategies for handling outlier and missing values, as well as comparing pooled versus meta-analysis strategies. We find that within study inverse normalization and half-minimum imputation for missing data provide reasonable and well-powered results in both pooled and meta-analyses. We utilize these strategies for cross-study metQTL analyses, identifying known and novel loci and validating the high quality of the metabolomics resources generated. TOPMed generated individual level data (Supplementary Table 14) and summary metQTL data (phs001974) are available for access through dbGaP, providing a resource for the scientific community to integrate metabolomics for understanding of common complex diseases.

Methods

Metabolite Data Acquisition

Metabolite data for eight studies were generated at the following two TOPMed Centralized Omics Resource Core (CORE) centers: 1.) the Baylor-UTHealth Metabolomics Center used the Metabolon, Inc. Global Discovery Panel to process COPDGene, SPIROMICS and WHI; and 2.) the Broad-Beth Israel Deaconess Medical Center (Broad-BIDMC) Panel was used in CAMP, CRA, FHS, MESA, and WHI. WHI had metabolites measured at both COREs, considered as two studies, WHI-Baylor-UTHealth and WHI-Broad-BIDMC. All studies used plasma derived from venous blood.

The Metabolon Global Discovery Panel is comprised of multiple chromatographic methods encompassing an ACQUITY reverse phase ultra-performance liquid chromatography tandem mass spectrometry (RP/UPLC-MS) (Waters; Milford, MA) and a Q Exactive accurate-mass liquid chromatography mass spectrometer (LC-MS) (Thermo Fisher Scientific; Waltham, MA) interfaced with a heated electrospray ionization (HESI-II) source, and Orbitrap mass analyzer operated at 35,000 mass resolution. Raw mass spectrometry data were analyzed using custom developed software (Metabolon, Inc.; Morrisville, NC) and compounds were identified by comparing them to library entries of purified standards or recurrent unknown entities. Metabolon maintains a library based on the retention time index (RI), mass to charge ratio (m/z), and chromatographic data established from >5,400 authenticated standards. Detailed methods were previously described[55–58].

Metabolomic profiles by the Broad-BIDMC Panel were obtained using liquid chromatography mass spectrometry (LC-MS) as previously described[59,60]. In brief, water-soluble, polar metabolites were assayed with a Nexera X2 U-HPLC (Shimadzu Corp; Marlborough, MA) using an Atlantis hydrophilic interaction LC column (Waters) coupled to a Q Exactive hybrid quadrupole Orbitrap mass spectrometer (Thermo Fisher Scientific). Raw data were visualized using TraceFinder 3.1 (Thermo Fisher Scientific) and Progenesis QI (Nonlinear Dynamics; Newcastle upon Tyne, United Kingdom). A 1290 Infinity LC system (Agilent Technologies) with a XBridge Amide column (Waters) was used in negative ionization or amide mode to measure organic acids, sugars, purines, pyrimidines, and other intermediary metabolites. The LC was coupled to a 6490 Triple Quad MS (Agilent Technologies) in multiple reaction monitoring (MRM) scanning mode, and MassHunter Quantitative Analysis software V10.1 (Agilent Technologies) was used for peak identification.

Metabolite Harmonization and Catalog

Study specific metabolite data was obtained through dbGaP from two metabolomic cores across eight studies from the TOPMed Program. Metabolite names and available identifiers, such as Human Metabolome Database (HMDB), KEGG, and chemical IDs (unique to the Global

Discovery Panel), were provided in metadata from each data generating core across studies. We used HMDB and metabolite names based on the latest Broad metabolite library, as well as manual check, to harmonize across studies profiled by the Broad Institute. Metabolites profiled by the Broad-BIDMC Panel were measured using multiple Broad methods, and we used a priority of C8-pos, HILIC-pos, C18-neg, and Amide (highest to lowest) to only keep unique metabolites. This priority was based on the metabolomic core recommendation, and the metabolite missing rates and CV values across Broad-BIDMC method studies. To harmonize across studies profiled by Metabolon Inc., we used Metabolon chemical IDs, HMDB, and metabolite names based on the latest Metabolon metabolite library. Metabolites provided by the Global Discovery Panel were based on internal platform priority. To harmonize metabolites across metabolomic cores, we used the HMDB and metabolite names provided by RefMet workbench. Duplicate metabolites were identified as having the same metabolite name or same HMDB, and only unique metabolites were retained based on method priority and missing rates. After harmonization, the available RefMet name provided by Metabolomics Workbench was assigned to each metabolite to produce a cross-platform and cross-study metabolite catalog. We note that our analyses here prioritize named metabolites only, unknown metabolites are also available in TOPMed datasets but are not categorized and assessed in the current work due to incomplete chemical structural information.

Pre-processing metabolites

Treatment of Metabolite Outliers. Investigators face several critical choices when pre-processing metabolomics data. Here we assessed choices regarding outlier treatment, multi-study analysis, and metabolite imputation, and made our recommendations for investigators using TOPMed metabolomics data. Higher values in the metabolite data are, in many cases, biologically plausible, however, they may be more difficult to handle in standard linear modeling analysis methods, and some may be spurious values. Thus, for handling outliers within study, three methods were considered: (1) removing observations more than 6 standard deviations away from the mean metabolite value (6SD), (2) rank-based inverse normal transformation (INT), and (3) winsorization, where outliers are replaced by the percentile value. In winsorization, for observations above the 95th percentile or below the 5th percentile, we replaced them with the 95th percentile and 5th percentile, respectively. Removing outliers in 6SD and winsorization removes values which may have a large impact on the effect estimates and power. However, the overall distribution of the metabolites remains similar. We also considered INT, which transforms the distribution of the metabolite to be normal. One advantage to this method is that it retains the outliers; however, real information that may be gained from skewness in the distribution is lost. One disadvantage of INT for metabolites with high missingness is that rank will be randomly selected for a high proportion of the data on the left side of the distribution. All three methods were initially applied within studies for each metabolite, removing missing observations. For comparability, we then apply INT on the pooled data across studies. To reduce computational burden, and in concordance with prior work using TOPMed study generated data, we compare the outlier methods using pooled data.

Pooled vs. Meta-Analysis. To address concerns regarding low statistical power, previous work has demonstrated the feasibility of using platform- and study- pooled metabolites to increase

sample size and maximize statistical power[38]. Data pooling is a method that also allows investigators to conduct research at a lower cost and computation burden. However, this method creates some challenges. For example, there will be technical variation across different metabolomic platforms and across different analysis centers. Additionally, missing metabolite values may vary in their distribution and source between studies. The target population for which the metabolites are measured will differ in various studies. These challenges need to be considered to produce high-quality harmonized data for pooled analyses.

Combining data across studies will increase the sample size and power. To assess the best method in TOPMed for such cross-study analysis, we compared a meta-analysis versus pooling across the studies. Due to the different platforms and studies, not all metabolites will be measured in every study. To compare meta-analysis with pooled analysis, we only considered metabolites that have measurements in 2 or more studies. We used complete-case with no missing imputation and compared the parameter estimate as well as the p-value in the meta-analysis versus pooled in using 6SD, INT, or winsorization. To provide a recommendation for analysis, we examined age and sex associations with metabolites. We removed related individuals. FHS was the only study with significant numbers of related individuals and using the pedigree file we removed individuals from the same family.

In a pooled analysis, for the i-th metabolite, we fit the age association model
$$Metabolite_{ij} = \beta_{0i} + \beta_{age,i}age_j + \beta_{study,i}study_j + \epsilon_{ij}$$
Where j denotes the subject with measurement of the i-th metabolite and $study_j$ is a categorical variable.

In a meta-analysis, we used the inverse variance-weighted (IVW) method to perform fixed effects meta-analysis. IVW calculates the weighted mean of the effect size estimates using the inverse variance of the individual studies as weights to summarize effect sizes from the independent studies. This model assumes that studies share a single true effect size. Thus, for i-th metabolite, we fit the age association model
$$Metabolite_{i_kj_k} = \beta_{0i_k} + \beta_{age,i_k}age_{j_k} + \epsilon_{i_kj_k}$$
To integrate multiple effect size estimates from multiple studies, we estimated $\hat{\beta}_{age,i} = \sum_k w_k \hat{\beta}_{age,i_k} / \sum_k w_k$, where w_k is the inverse variance of the age effect estimate. Similar models are fit for sex by replacing age_j with sex_j. Age at metabolite measure and sex were study-provided by participating studies.

Metabolite Imputation. Missing data is common in broad-spectrum metabolomics data. While the use of single value left imputation methods (such as half-min or zero) is common, several authors have claimed better performance of more sophisticated methods such as Random Forest [30]. We compared the following missing imputation strategies: no imputation, zero, half-min, median, Quantile Regression for Imputation of Left-Censored data (QRILC)[19], k-Nearest Neighbors (KNN)[61], Random Forest (RF)[20], and Mechanism-Aware Imputation

(MAI)[30]. Recommendations were decided based on number of known metQTLs (from prior GWAS in non-TOPMed studies) detected in targeted analysis (and strength of associations) with different missingness imputation strategies, as well as comparisons of number and strength of associations with age and sex, both for known associations and overall (with the assumption that most additional age and sex associations identified are plausible true positives, given the broad impact of both age and sex on the circulating metabolome[50,62–69]).

To assess the downstream effect of imputation methods on genetic association analyses, we first assembled a list of putative metabolite-variant associations based on three large-scale metabolite GWAS studies in studiesindependent from those included here in TOPMed[3,11,18]. After linking metabolites by HMDB, as described above, and filtering by missingness (1-50%) in our own datasets, we were able to identify 125 reported metQTLs for 28 metabolites present on the Broad-BIDMC Panel, and 913 QTLs for 204 metabolites present on the Global Discovery Panel. Treating these associations as "true positives", we assessed our ability to replicate these associations for each imputation strategy. Using the GMMAT pipeline on BioData Catalyst, as described below, we tested for metabolite-variant associations. After imputing metabolite levels and inverse-normalizing values within study sites, we pooled datasets within each platform and tested for genetic associations with adjustment for age, sex, and study(where applicable). To correct for sample relatedness and population structure, we used sparse genetic relationship matrices and the first 10 genetic principal components. After performing association testing for each imputation method within both metabolomics platforms, we assessed imputation strategies based on the number of literature-reported metQTLs identified as Bonferroni-significant and with the previously reported direction of effect.

Detection of Metabolite QTLs

Only the metabolites measured in at least two studies and missing rates <50% in these studies were included in the following QRLs analysis. Totally, there are 1,135 metabolites from 16,359 participants across eight studies , i.e., CAMP, CRA, FHS, MESA, COPDGene, SPIROMICS, WHI-Baylor-UTHealth and WHI-Broad-BIDMC. The sample size of female and male participants in each study are roughly balanced, except for WHI, which only has female participants. The sample size of each study was shown in Supplementary Table 13. Presence of ancestry diversity in participating studies based on similarity to reference panels has been previously described[70].

Whole Genome Sequencing. TOPMed freeze 10 WGS data was utilized throughout our analyses. Joint calling was performed across all freeze 10 studies by the TOPMed Informatics Research Center (IRC). Detailed calling, processing and quality control were previously described[70]. In brief, variant calling was completed using the GotCloud pipeline. Variant quality control was performed by calculating Mendelian consistency scores and by applying a support vector machine (SVM) classifier trained on known variant sites and Mendelian inconsistent variants. Sample QC measures included: concordance between annotated and inferred genetic sex, concordance between prior array genotype data and TOPMed WGS data, and pedigree checks. Further details regarding data processing, and quality control are described on the

TOPMed   website
(https://topmed.nhlbi.nih.gov/topmed-whole-genome-sequencing-methods-freeze-9) and in a
common document accompanying each TOPMed study's dbGaP accession. Variants located on
both autosomal and X chromosomes were considered in both sex-combined and sex-stratified
analyses. Variants with a MAF less than 0.5% in the overall sample were removed, resulting in
16,518,857 variants included in the following association analyses.

Single variant tests on the pooled TOPMed data. For sex-pooled analysis, a total of 16,359
individuals were analyzed, including 8,855 females and 7,500 males. Missing values on
metabolites were imputed using half the minimum values and inverse normal transformations
were performed by the study. We included age, sex, study center, and top eleven ancestry
principal components as fixed-effects covariates[10]. Study_center is the study the individual
belongs to, as well as a recruitment center for relevant studies. The sparse kinship matrix was
used as a random effect to account for sample relatedness. A two-stage procedure for rank
normalization was applied in genotype-metabolite association analysis[71]. For each of the 1,135
metabolites, we rank-normalized the residuals after regressing the metabolites on fixed-effects
covariates, and then used them as the phenotype in downstream null model fitting and
association tests, which was performed by GMMAT pipeline on BioData Catalyst (BDC). We
applied a Bonferroni-correction to define   genome-wide significance at P-value <
$5.0 \times 10^{-8}/1,135 = 4.4 \times 10^{-11}$, accounting for 1,135 analyzed metabolites

Conditional analysis on single variant-metabolite associations. Across the analyzed
genome, we defined metabolite-associated genetic loci (i.e. genomic windows) as those
containing all statistically significant variants within 500 kb from each other. To account for
linkage disequilibrium, we added 500 kbp up- and downstream as a buffer to the genetic loci. If
any of these loci overlapped, we merged them together into a single locus. In this way, we
identified 248 loci, containing 1,248 locus-metabolite pairs.

To identify independent leading variants for each of the locus-metabolite pairs, we conducted
conditional analysis by GMMAT. For each locus-metabolites pair, the rank-normalized residuals
after regressing the metabolite on fixed-effects covariates from single variant tests was used as
the phenotype in the downstream null model fitting. The genotypes of those identified top
associated variants from each locus together with the fixed-effects covariates were used as
covariates in the null model fitting. This conditioning process started with the top associated
variant, across the locus followed by a stepwise procedure of selecting additional variants, one
by one, according to their conditional P values. At the *t*-th step, we calculated the conditional P
values of all the remaining SNPs in the locus conditional on the SNPs that have been selected
in the previous steps. We then selected the SNP with minimum conditional P value that was less
than the cutoff P-value of 5x10[-8] and added it to the covariates in the next step. We repeated
this process until no SNPs can be selected into the model. Variants that were selected in each
step in the conditional analysis were considered conditionally independent associations.

Annotation of known and novel metQTL findings. To access which of the independent
variant-metabolite associations were novel, we compiled results from the Metabolomic GWAS
Server[1], TwinsUK Study[8], GWAS Catalog, GRASP Search, previous reports from our group[6,10]

and performed manual search through published papers to detect known loci that overlap with our findings. The significant locus-metabolite pair we identified were classified as known if any variant from a locus-metabolite pair was previously associated with any of the metabolites in its sub-pathway. Otherwise, the locus-metabolite pair was considered novel.

Replication of novel metQTLs. We performed replication analysis using metQTLs previously reported from among 16,359 multi-ethnic participants from five studies with WGS, including the Atherosclerosis Risk in Communities study (ARIC), Hispanic Community Health Study/Study of Latinos (HCHS/SOL), Cardiovascular Health Study (CHS), FHS and MESA[10]. Of note, a small proportion, 996 out of 16,539 participants (6%), overlapped with the samples from the current analysis. Out of 108 novel metQTLs, 82 pairs were available for replication in Feofanova et al. (2023)[10]. The novel pairs were replicated with either the reported variant or a variant within 500 kb regions for the same metabolite having $P \leq 6.10 \times 10^{-4}$. Sixty six novel metQTL loci-trait pairs were defined to be significantly replicated.

Categorization of novel metQTLs based on PhenoScanner. We further used PhenoScanner2 (citation) to determine whether the novel metQTLs were also associated with other traits/diseases to improve the function annotation. We got the 135 novel independent loci (from 108 locus-metabolite associated pairs; Fig. 1) and generated the input list by combining all significant variants associated with corresponding metabolites from each independent loci, leading to a total of 6,500 variants for PhenoScanner queries. The scanning was performed on database version 2 (till 12 Dec, 2023) with categories setting with "*mQTL*" and "*GWAS*" respectively. We further extracted the "*Disease*" related queries from the "*GWAS*" category by matching the "disease" string within traits. The queries from multiple studies were merged as one unique search.

Sex-stratified data analysis

For sex-stratified analysis, the single variant test was conducted on female individuals (8,855 individuals) and male individuals (7,500 individuals) separately following the same analytical steps. Biological sex was defined as "male" for study individuals with sex chromosomes XY and "female" for those with sex chromosomes XX; we acknowledge that such chromosomal make-up data should not be equated with self-identified gender, which was not consistently examined across included studies. We identified the sex-dimorphic associations using the following steps. We first selected the variants that were statistically significant in the sex-stratified analyses for either females or males, i.e., $P_{female} \leq 4.4 \times 10^{-11}$ or $P_{male} \leq 4.4 \times 10^{-11}$. Significant variants located within 500 kb from each other were merged into a single locus. To account for linkage disequilibrium, we added 500 kb up- and downstream as a buffer to the genetic loci. If any of these loci overlapped, we merged them together into a single locus. In total, 184 independent clumped loci were identified from sex-stratified analyses. For the lead variant of each independent locus, we then tested if the genetic effects were different

between females and males and computed P-values (P_{diff}). Sex-dimorphic associations were defined as loci with P_{diff} smaller than the Bonferroni corrected significance level $0.05/184 = 2.7\times10^{-4}$.

In each locus-metabolite pair, we conducted conditional analysis using GMMAT to identify the independent associated variants following the aforementioned steps. Variants that were both statistically significant in the sex-stratified analysis ($P_{female}\leq5.0\times10^{-8}$ or $P_{male}\leq5.0\times10^{-8}$) and $P_{diff}\leq2.7\times10^{-4}$ in the conditional analysis were considered conditionally independent, sex-dimorphic associations.

Acknowledgments

Molecular data for the Trans-Omics in Precision Medicine (TOPMed) program was supported by the National Heart, Lung and Blood Institute (NHLBI). See the TOPMed Omics Support section in Supplementary Materials for study specific omics support information. Core support including centralized genomic read mapping and genotype calling, along with variant quality metrics and filtering were provided by the TOPMed Informatics Research Center (3R01HL-117626-02S1; contract HHSN268201800002I). Core support including phenotype harmonization, data management, sample-identity QC, and general program coordination were provided by the TOPMed Data Coordinating Center (R01HL-120393; U01HL-120393; contract HHSN268201800001I). We gratefully acknowledge the studies and participants who provided biological samples and data for TOPMed. Additional study specific acknowledgments are included in Supplementary Materials.

NF is supported by NIH MD012765, DK117445, HL163972. RSK is supported by NHLBI: K01 HL146980. LW is supported by NHGRI/NIMHD U54HG013243. The work and BY is in part supported by R01HL168638 and the JLH Foundation.

Conflict of Interests

In the past three years, EKS received grant support from Bayer and Northpond Laboratories. LW provided consulting service to Pupil Bio Inc. and reviewed manuscripts for Gastroenterology Report, not related to this study, and received honorarium. LMR is a consultant for the TOPMed Administrative Coordinating Center (through Westat).

References

1. Shin, S.-Y. *et al.* An atlas of genetic influences on human blood metabolites. *Nat. Genet.* **46**, 543–550 (2014).

2. Vattikuti, S., Guo, J. & Chow, C. C. Heritability and genetic correlations explained by common SNPs for metabolic syndrome traits. *PLoS Genet.* **8**, e1002637 (2012).

3. Yin, X. *et al.* Genome-wide association studies of metabolites in Finnish men identify disease-relevant loci. *Nat. Commun.* **13**, 1644 (2022).

4. Hysi, P. G. *et al.* Metabolome Genome-Wide Association Study Identifies 74 Novel Genomic Regions Influencing Plasma Metabolites Levels. *Metabolites* **12**, 61 (2022).

5. Lotta, L. A. *et al.* A cross-platform approach identifies genetic regulators of human metabolism and health. *Nat. Genet.* **53**, 54–64 (2021).

6. Feofanova, E. V. *et al.* A Genome-wide Association Study Discovers 46 Loci of the Human Metabolome in the Hispanic Community Health Study/Study of Latinos. *Am. J. Hum. Genet.* **107**, 849–863 (2020).

7. Bar, N. *et al.* A reference map of potential determinants for the human serum metabolome. *Nature* **588**, 135–140 (2020).

8. Long, T. *et al.* Whole-genome sequencing identifies common-to-rare variants associated with human blood metabolites. *Nat. Genet.* **49**, 568–578 (2017).

9. Tahir, U. A. *et al.* Whole Genome Association Study of the Plasma Metabolome Identifies Metabolites Linked to Cardiometabolic Disease in Black Individuals. *Nat. Commun.* **13**, 4923 (2022).

10. Feofanova, E. V. *et al.* Whole-Genome Sequencing Analysis of Human Metabolome in Multi-Ethnic Populations. *Nat. Commun.* **14**, 3111 (2023).

11. Surendran, P. *et al.* Rare and common genetic determinants of metabolic individuality and their effects on human health. *Nat. Med.* **28**, 2321–2332 (2022).

12. Mauvais-Jarvis, F. *et al.* Sex and gender: modifiers of health, disease, and medicine. *Lancet Lond. Engl.* **396**, 565–582 (2020).

13. Mauvais-Jarvis, F. *et al.* Sex- and Gender-Based Pharmacological Response to Drugs. *Pharmacol. Rev.* **73**, 730–762 (2021).

14. Ramsey, J. M., Cooper, J. D., Penninx, B. W. J. H. & Bahn, S. Variation in serum biomarkers with sex and female hormonal status: implications for clinical tests. *Sci. Rep.* **6**, 26947 (2016).

15. Teruya, T., Chen, Y.-J., Kondoh, H., Fukuji, Y. & Yanagida, M. Whole-blood metabolomics of dementia patients reveal classes of disease-linked metabolites. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2022857118 (2021).

16. Chaleckis, R., Murakami, I., Takada, J., Kondoh, H. & Yanagida, M. Individual variability in human blood metabolites identifies age-related differences. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 4252–4259 (2016).

17. Deja, S. *et al.* Gender-Specific Metabolomics Approach to Kidney Cancer. *Metabolites* **11**, 767 (2021).

18. Chen, Y. *et al.* Genomic atlas of the plasma metabolome prioritizes metabolites implicated in human diseases. *Nat. Genet.* **55**, 44–53 (2023).

19. Wei, R. *et al.* Missing Value Imputation Approach for Mass Spectrometry-based Metabolomics Data. *Sci. Rep.* **8**, 663 (2018).

20. Stekhoven, D. J. & Bühlmann, P. MissForest—non-parametric missing value imputation for mixed-type data. *Bioinformatics* **28**, 112–118 (2012).

21. Dekermanjian, J. P., Shaddox, E., Nandy, D., Ghosh, D. & Kechris, K. Mechanism-aware imputation: a two-step approach in handling missing values in metabolomics. *BMC Bioinformatics* **23**, 179 (2022).

22. Schlosser, P. *et al.* Genetic studies of paired metabolomes reveal enzymatic and transport processes at the interface of plasma and urine. *Nat. Genet.* **55**, 995–1008 (2023).

23. Raffler, J. *et al.* Genome-Wide Association Study with Targeted and Non-targeted NMR Metabolomics Identifies 15 Novel Loci of Urinary Human Metabolic Individuality. *PLoS Genet.* **11**, e1005487 (2015).

24. Kettunen, J. *et al.* Genome-wide study for circulating metabolites identifies 62 loci and reveals novel systemic effects of LPA. *Nat. Commun.* **7**, 11122 (2016).

25. Wang, Z. *et al.* Genome-wide association study of metabolites in patients with coronary artery disease identified novel metabolite quantitative trait loci. *Clin. Transl. Med.* **11**, e290 (2021).

26. Luo, S. *et al.* Genome-wide association study of serum metabolites in the African American Study of Kidney Disease and Hypertension. *Kidney Int.* **100**, 430–439 (2021).

27. Noh, H. & King, G. L. The role of protein kinase C activation in diabetic nephropathy. *Kidney Int. Suppl.* S49-53 (2007) doi:10.1038/sj.ki.5002386.

28. Wittemans, L. B. L. *et al.* Assessing the causal association of glycine with risk of cardio-metabolic diseases. *Nat. Commun.* **10**, 1060 (2019).

29. Yu, B. *et al.* The Consortium of Metabolomics Studies (COMETS): Metabolomics in 47 Prospective Cohort Studies. *Am. J. Epidemiol.* **188**, 991–1012 (2019).

30. Kokla, M., Virtanen, J., Kolehmainen, M., Paananen, J. & Hanhineva, K. Random forest-based imputation outperforms other methods for imputing LC-MS metabolomics data: a comparative study. *BMC Bioinformatics* **20**, 492 (2019).

31. Taylor, S., Ponzini, M., Wilson, M. & Kim, K. Comparison of imputation and imputation-free methods for statistical analysis of mass spectrometry data with missing data. *Brief. Bioinform.* **23**, bbab353 (2022).

32. Wilson, M. D., Wilson, M. D., Taylor, S. L. & Kim, K. Imputation of Missing Values for Multi-Biospecimen Metabolomics Studies: Bias and Effects on Statistical Validity. *Metabolites* **12**, (2022).

33. Richardson, S., Tseng, G. C. & Sun, W. Statistical Methods in Integrative Genomics. *Annu. Rev. Stat. Its Appl.* **3**, 181–209 (2016).

34. Jendoubi, T. Approaches to Integrating Metabolomics and Multi-Omics Data: A Primer. *Metabolites* **11**, 184 (2021).

35. Robinson, O. *et al.* Metabolic profiles of socio-economic position: a multi-cohort analysis. *Int. J. Epidemiol.* **50**, 768–782 (2020).

36. Zheng, R., Michaëlsson, K., Fall, T., Elmståhl, S. & Lind, L. The metabolomic profiling of total fat and fat distribution in a multi-cohort study of women and men. *Sci. Rep.* **13**, 11129 (2023).

37. Burke, D. L., Ensor, J. & Riley, R. D. Meta-analysis using individual participant data: one-stage and two-stage approaches, and why they may differ. *Stat. Med.* **36**, 855–875 (2017).

38. Boedhoe, P. S. W. *et al.* An Empirical Comparison of Meta- and Mega-Analysis With Data From the ENIGMA Obsessive-Compulsive Disorder Working Group. *Front. Neuroinformatics* **12**, 102 (2019).

39. Han, X. *et al.* Integrating genetics and metabolomics from multi-ethnic and multi-fluid data reveals putative mechanisms for age-related macular degeneration. *Cell Rep. Med.* **4**, 101085 (2023).

40. Lowenstein, E. J. *et al.* The SH2 and SH3 domain-containing protein GRB2 links receptor tyrosine kinases to ras signaling. *Cell* **70**, 431–442 (1992).

41. Downward, J. The GRB2/Sem-5 adaptor protein. *FEBS Lett.* **338**, 113–117 (1994).

42. Zhao, M., Jung, Y., Jiang, Z. & Svensson, K. J. Regulation of Energy Metabolism by Receptor Tyrosine Kinase Ligands. *Front. Physiol.* **11**, 354 (2020).

43. Li, J. *et al.* VSIG4 inhibits proinflammatory macrophage activation by reprogramming mitochondrial pyruvate metabolism. *Nat. Commun.* **8**, 1322 (2017).

44. Berg, J. M., Tymoczko, J. L. & Stryer, L. *Biochemistry (Loose-Leaf)*. (Macmillan, 2007).

45. Costanzo, M. *et al.* Sex differences in the human metabolome. *Biol. Sex Differ.* **13**, 30 (2022).

46. Huang, Y. *et al.* Deciphering genetic causes for sex differences in human health through drug metabolism and transporter genes. *Nat. Commun.* **14**, 175 (2023).

47. Bocedi, A. *et al.* Glutathione Transferase P1-1 an Enzyme Useful in Biomedicine and as Biomarker in Clinical Practice and in Environmental Pollution. *Nutrients* **11**, 1741 (2019).

48. Xiong, D.-K., Chen, H.-H., Ding, X.-P., Zhang, S.-H. & Zhang, J.-H. Association of polymorphisms in glutathione S-transferase genes (GSTM1, GSTT1, GSTP1) with idiopathic azoospermia or oligospermia in Sichuan, China. *Asian J. Androl.* **17**, 481–486 (2015).

49. Krumsiek, J. *et al.* Gender-specific pathway differences in the human serum metabolome. *Metabolomics Off. J. Metabolomic Soc.* **11**, 1815–1833 (2015).

50. Mittelstrass, K. *et al.* Discovery of sexual dimorphisms in metabolic and genetic biomarkers. *PLoS Genet.* **7**, e1002215 (2011).

51. Borodulin, K. & Katri, S. FinHealth 2017 Study – Methods.

52. Ritchie, S. C. *et al.* Quality control and removal of technical variation of NMR metabolic biomarker data in \textasciitilde120,000 UK Biobank participants. *Sci. Data* **10**, 64 (2023).

53. Chen, Z.-Z. *et al.* Nontargeted and Targeted Metabolomic Profiling Reveals Novel Metabolite Biomarkers of Incident Diabetes in African Americans. *Diabetes* **71**, 2426–2437 (2022).

54. O'Sullivan, J. F. *et al.* Dimethylguanidino valeric acid is a marker of liver fat and predicts diabetes. *J. Clin. Invest.* **127**, 4394–4402 (2017).

55. Ford, L. *et al.* Clinical metabolomics for inborn errors of metabolism. *Adv. Clin. Chem.* **107**, 79–138 (2022).

56. Daley-Yates, P. *et al.* Inhaled glucocorticoid-induced metabolome changes in asthma. *Eur. J. Endocrinol.* **187**, 413–427 (2022).

57. Peng, H. *et al.* Identification of Metabolite Markers Associated with Kidney Function. *J. Immunol. Res.* **2022**, 6190333 (2022).

58. Benedetti, E. *et al.* Plasma metabolomics profiling of 580 patients from an Early Detection Research Network prostate cancer cohort. *Sci. Data* **10**, 830 (2023).

59. Benson, M. D. *et al.* Protein-metabolite association studies identify novel proteomic determinants of metabolite levels in human plasma. *Cell Metab.* **35**, 1646-1660.e3 (2023).

60. Tahir, U. A. *et al.* Metabolomic Profiles and Heart Failure Risk in Black Adults: Insights From the Jackson Heart Study. *Circ. Heart Fail.* **14**, e007275 (2021).

61. Troyanskaya, O. *et al.* Missing value estimation methods for DNA microarrays. *Bioinformatics* **17**, 520–525 (2001).

62. Panyard, D. J., Yu, B. & Snyder, M. P. The metabolomics of human aging: Advances, challenges, and opportunities. *Sci. Adv.* **8**, eadd6155 (2024).

63. Bell, J. A. *et al.* Sex differences in systemic metabolites at four life stages: cohort study with repeated metabolomics. *BMC Med.* **19**, 58 (2021).

64. Rist, M. J. *et al.* Metabolite patterns predicting sex and age in participants of the Karlsruhe Metabolomics and Nutrition (KarMeN) study. *PloS One* **12**, e0183228 (2017).

65. Verri Hernandes, V. *et al.* Age, Sex, Body Mass Index, Diet and Menopause Related Metabolites in a Large Homogeneous Alpine Cohort. *Metabolites* **12**, 205 (2022).

66. Wang, Q. *et al.* Metabolic characterization of menopause: cross-sectional and longitudinal evidence. *BMC Med.* **16**, 17 (2018).

67. Adav, S. S. & Wang, Y. Metabolomics Signatures of Aging: Recent Advances. *Aging Dis.* **12**, 646–661 (2021).

68. Hertel, J. *et al.* Measuring Biological Age via Metabonomics: The Metabolic Age Score. *J. Proteome Res.* **15**, 400–410 (2016).

69. Robinson, O. *et al.* Determinants of accelerated metabolomic and epigenetic aging in a UK cohort. *Aging Cell* **19**, e13149 (2020).

70. Taliun, D. *et al.* Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. *Nature* **590**, 290–299 (2021).

71. Sofer, T. *et al.* A fully adjusted two-stage procedure for rank-normalization in genetic association studies. *Genet. Epidemiol.* **43**, 263–275 (2019).