Diet of Andean leaf-eared mice (*Phyllotis*) living at extreme elevations on Atacama

volcanoes: insights from metagenomics, DNA metabarcoding, and stable isotopes

Claudio Quezada-Romegialli[1], Marcial Quiroga-Carmona[2,3,4], Guillermo D'Elía[2,3], Chris

Harrod[5,6,7], Jay F. Storz[4]

[1]Plataforma de Monitoreo Genómico y Ambiental, Departamento de Química, Facultad de

Ciencias, Universidad de Tarapacá, Arica, Chile

[2]Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de

Chile, Valdivia, Chile

[3]Colección de Mamíferos, Facultad de Ciencias, Universidad Austral de Chile, Campus Isla

Teja, Valdivia, Chile

[4]School of Biological Sciences, University of Nebraska, Lincoln, Nebraska, USA.

[5]Instituto de Ciencias Naturales Alexander von Humboldt, Universidad de Antofagasta,

Antofagasta, Chile

[6]Núcleo Milenio de Salmónidos Invasores Australes, INVASAL, Concepción, Chile

[7]Scottish Centre for Ecology and the Natural Environment, School of Biodiversity, One Health

and Veterinary Medicine, University of Glasgow

Correspondence:

Jay F. Storz
School of Biological Sciences
University of Nebraska
Lincoln, Nebraska, USA
E-mail: jstorz2@unl.edu

Abstract

On the flanks of >6000 m Andean volcanoes that tower over the Atacama Desert, leaf-eared mice (*Phyllotis vaccarum*) live at extreme elevations that surpass known vegetation limits. What the mice eat in these barren, hyperarid environments has been the subject of much speculation. According to the arthropod fallout hypothesis, sustenance is provided by windblown insects that accumulate in snowdrifts ('aolian deposits'). It is also possible that mice feed on saxicolous lichen or forms of cryptic vegetation that have yet to be discovered at such high elevations. We tested hypotheses about the diet of mice living at extreme elevations on Atacama volcanoes by combining metagenomic and DNA metabarcoding analyses of gut contents with stable-isotope analyses of mouse tissues. Genomic analyses of contents of the gastrointestinal tract of a live-captured mouse from the 6739 m summit of Volcán Llullaillaco revealed evidence for an opportunistic but purely herbivorous diet, including lichens. Although we found no evidence of animal DNA in gut contents of the summit mouse, stable isotope data indicate that mice native to elevations at or near vegetation limits (~5100 m) include a larger fraction of animal prey in their diet than mice from lower elevations. Some plant species detected in the gut contents of the summit mouse are known to exist at lower elevations at the base of the volcano and in the surrounding Altiplano, suggesting that such plants may occur at higher elevations beneath the snowpack or in other cryptic microhabitats.

Running title: Diet of Andean mice from extreme elevations

Keywords: aolian zone, Andes, high altitude, lichenivory, Llullaillaco, Puna de Atacama

1. INTRODUCTION

Extreme high-elevation surveys of small mammals in the Central Andes have yielded live captures of numerous specimens of the Andean leaf-eared mouse *Phyllotis vaccarum* at elevations at or above the elevational limits of vegetation (>5,000 m)(Storz et al., 2024). One specimen was captured at 6,739 m (22,100 feet) above sea level on the very summit of Volcán Llullaillaco, a stratovolcano in the Central Andes that straddles the Argentina-Chile border (Storz et al., 2020). This summit specimen far surpasses previous elevational records for wild mammals in the Andes and Himalaya. Documentation of active burrows of *P. vaccarum* at >6,100 m on the flanks of Llullaillaco and the discovery of desiccated cadavers ('mummies') of *Phyllotis* on the summits of Llullaillaco and several neighboring >6,000 m volcanoes confirm that these mice inhabit extreme elevations well above the apparent limits of vascular plants (Halloy, 1991; Steppan et al., 2022; Storz et al., 2023, 2024). Evidence that high-elevation mice are living in an apparently barren world of rock, ice, and snow prompts numerous questions, perhaps none more basic than: What are they eating? In the perennial winter conditions that prevail at elevations >6,000 m, the scarcity of food poses a special physiological challenge for small endotherms like mice because of the energetic demands of thermoregulation. Moreover, leaf-eared mice in the genus *Phyllotis* do not hibernate, so the energetic challenge of sustaining endothermy in cold, hypoxic conditions is especially acute (Storz & Scott 2019).

It has been suggested that windblown arthropods and/or vegetation could provide a source of food for animals living at elevations that exceed the limits at which green plants grow (the 'aolian zone') (Swan, 1961, 1992). According to this hypothesis, the transport of airborne nutrients from lower elevations sustains life on the upper reaches of a volcano like Llullaillaco similar to the way that the fallout of organic detritus from upper layers of the water column sustains life in the aphotic zone of the deep ocean. Aolian deposits of windblown arthropods along the lee edge of mountain summits and ridgelines can attract birds and other insectivorous animals that normally forage at much lower elevations (Antor, 1995; Spalding, 1979). If aolian deposits of windblown arthropods ('arthropod fallout') help sustain populations of *P. vaccarum* at extreme elevations on Atacama volcanoes, we would expect arthropods to constitute a much larger part of their diet than at lower elevations where the species is mainly granivorous and frugivorous (Bozinovic and Rosenmann 1988; López-Cortés et al. 2007; Sassi et al. 2017). On the Quinghai-Tibetan Plateau, high elevation plateau pikas (*Ochotona curzoniae*) exploit the feces of yak (*Bos grunniens*) as a food source (Speakman et al., 2021). If *Phyllotis* mice practice a similar form of interspecific coprophagy, feces from Andean camelids such as vicuña (*Lama vicugna*) and guanaco (*Lama guanicoe*) would provide the most readily available source.

Although neither vicuña nor guanaco typically spend much time above the elevational limit of

vegetation, both species are known to traverse mountain passes at elevations >5,500 m in the

Central Andes (J. Storz, personal observation). It is also possible that the mice feed on lichen

that grows on rock substrates (saxicolous lichen) or some cryptic form of vegetation that is not

currently known to occur at such extreme elevations.

Here we test the above-mentioned hypotheses by conducting metagenomic and

metabarcoding analyses of gut contents from the world-record specimen of *P. vaccarum* that

was live-captured on the summit of Llullaillaco at 6,739 m. Since the metagenomic approach

involves high-throughput sequencing of all DNA extracted from a sample without PCR

enrichment of specific markers, it is not biased by *a priori* expectations about which taxonomic

groups to expect and is therefore well-suited to dietary assessments of omnivorous species

(Chua et al., 2020). DNA metabarcoding complements the metagenomic approach and can be

used to estimate the diversity and relative abundance of different items in the diet (Deagle et al.,

2019; Stapleton et al., 2022).

To complement the metagenomic and metabarcoding analysis of the summit specimen,

we conducted a stable isotope analysis of liver samples from a larger sample of wild-caught

mice from a broad range of elevations in the Chilean Altiplano and Puna de Atacama (2,370-

6,739 m). We used stable isotope values of three key elements (carbon, nitrogen and sulfur)

from liver tissue to characterize the diet of *P. vaccarum* over a timespan of weeks to months. In

the livers of small mammals, isotopic half-lives are <1 week for both carbon ($\delta^{13}C$) and nitrogen

($\delta^{15}N$) and we expect a similar half-life for sulfur ($\delta^{34}S$). Examination of stable isotope values

permits inferences about several key components of trophic ecology.

Carbon stable isotopes ($\delta^{13}C$) reflect the relative consumption of food derived from

different sources of primary production (Cerling et al., 1997; DeNiro & Epstein, 1978). The

stable isotope of nitrogen is typically used as an indicator of consumer trophic position

(Vanderklift & Ponsard, 2003; Quezada-Romegialli et al., 2018) but can also be used to

discriminate between consumption of food from distinct habitats (Harrod et al., 2005). For

example, it should be possible to assess the extent to which mice rely on lichen at high

elevations. Lichens are typically very [15]N-depleted relative to terrestrial plants (Fogel et al.,

2008; Lee, Lim & Yoon, 2009; Pinho et al., 2017), and this holds true for lichens from high

elevations (Biazrov, 2012; Marris et al., 2019; Szpak et al., 2013) and volcanic fumeroles (Tozer

et al. 2005). If lichen forms an important part of the diet of *P. vaccarum* at high elevations, we

would expect to observe negative $\delta^{15}N$ values. Sulfur stable isotopes ($\delta^{34}S$) are also useful

indicators of consumer habitat use as values measured from plants and their consumers often

exhibit high levels of spatial variation across biogeochemical gradients (Krouse et al., 1991;

Nielsen et al., 1991), as might be expected along the flanks of an historically active volcano like

Llullaillaco.

By combining metagenomics, metabarcoding, and stable-isotope analyses, we tested

several hypotheses about the diet of mice living at extreme elevations. The arthropod fallout

hypothesis would be supported by the presence of DNA from insects or other arthropods that

could be blown upslope, and isotopic estimates of trophic position would be higher for mice

living at especially high elevations on the flanks or summit of the volcano in comparison with

those from lower elevations in the surrounding Altiplano. Interspecific coprophagy (or

scavenging) would be supported by the presence of DNA from vicuña, guanaco, or other co-

distributed mammals. Lichenivory would be supported by the presence of DNA from lichen-

associated fungi or green algae and especially low ^{15}N values in mice from high elevations. The

particular plants detected in the gut contents of the summit mouse may suggest that some plant

species actually occur at much higher elevations than currently assumed, but they may be

sparsely distributed in cryptic microhabitats (e.g., in rock crevices or under the snowpack). If

that is the case, then the diet of mice living at >6,000 m may include a subset of the same

plants that mice feed on at lower elevations.

2. MATERIALS AND METHODS

2.1 Sampling

We live-captured all mice using Sherman live traps and other methods described in Storz et al.,

(2020, 2024). We collected mice from a broad range of elevations in the Altiplano/Puna de

Atacama ecoregions, from 2,340 m in the Atacama Desert to the 6,739 m summit of Volcán

Llullaillaco (Figure 1). We sacrificed mice in the field, prepared them as museum specimens,

and preserved liver tissue in 95‰ ethanol as source material for the stable isotope analysis. For

the *P. vaccarum* specimen captured on the summit of Volcán Llullaillaco, we preserved the

entire gastrointestinal tract in ethanol as a source of DNA for metagenomic and metabarcoding

analyses. All mouse specimens are housed in the Colección de Mamíferos of the Universidad

Austral de Chile, Valdivia, Chile.

We collected all mice in accordance with permissions to JFS and GD from the following

Chilean government agencies: Servicio Agrícola y Ganadero (SAG, Resolución exenta #

6633/2020), Corporación Nacional Forestal (CONAF, Autorización # 171219), and Dirección

Nacional de Fronteras y Límites del Estado (DIFROL, Autorización de Expedición Científica

#68/2020). We handled all mice in accordance with protocols approved by the Institutional

Animal Care and Use Committee (IACUC) at the University of Nebraska (project ID: 1919).

2.2 Dissection of gastrointestinal tract

For the mouse captured on the summit of Volcán Llullaillaco (UACH8291), we extracted DNA

from contents of the stomach for metagenomic sequencing and DNA metabarcoding. We also

dissected the lower gastrointestinal tract into 13 adjoining sections, the cecum and 12

consecutive segments of the colon, ordered from the outlet of the cecum to the rectum (Figure

2), and we extracted DNA from the contents of each section for additional DNA metabarcoding

analysis. This approach allowed us to examine temporal changes in the mouse's diet, as

determined by gut passage times: the stomach and cecum contain food items ingested within a

few hours of its capture, while the colon sections and rectum potentially contain food ingested

within the previous two or three days. The metagenomic sequencing represents an unbiased

approach to characterize the stomach contents of the mouse while the metabarcoding analysis

is designed to test specific hypotheses about the animal's diet (arthropod fallout, interspecific

coprophagy, lichenivory, or herbivory at elevations that surpass assumed vegetation limits).

2.3 Metagenomic sequencing

We sent contents of the dissected sections of the gastrointestinal tract to Azenta Life Sciences

(South Plainfield, NJ, USA) for metagenomic analysis. Genomic DNA was isolated using the

NucleoMag DNA Microbiome Kit (Takara Bio, Shiga, Japan) and was quantified using a Qubit

2.0 Fluorometer (ThermoFisher Scientific, Waltham, MA, US). NEBNext® Ultra™ II DNA Library

Prep Kit for Illumina (New England Biolabs, Ipswich, MA, USA) was used for library preparation

following manufacturer's recommendations. Briefly, genomic DNA was fragmented by acoustic

shearing with a Covaris S220 instrument, followed by end-repair. Adapters were ligated after

adenylation of the 3'ends followed by enrichment by a limited cycle PCR. DNA libraries were

quantified using Qubit 2.0 Fluorometer and by real time PCR (Applied Biosystems, Carlsbad,

CA, USA). Sequencing libraries were sequenced on an Illumina HiSeq instrument using a 2 x

150bp Paired End (PE) configuration. Image analysis and base calling were conducted using

the HiSeq Control Software (HCS). Raw BCL files were converted to FastQ files and de-

multiplexed using bcl2fastq v.2.1.9 (Illumina), keeping only >Q30 reads with 150 bp in length. A

de novo approach was followed for assembling reads using Spades v3.10 (Bankevich et al.,

2012), with a minimum contig length of 1,000 bp and using the newly assembled genome of

Phyllotis vaccarum (Storz et al., 2023) as a reference genome to discard reads aligned to the

host. QUAST (Gurevich et al., 2013) was used to generate statistics and EMBOSS tools getorf

was used to find the open reading frames within the *de novo* assembled genome. BLAST+

(v.2.6.0) (Altschul et al., 1990) was used to query assembled contigs in the nucleotide database

of Genbank.

2.4　DNA metabarcoding analysis and primer selection

The stomach, cecum, and 12 consecutive segments of the colon were sent to MrDNA

(www.mrdnalab.com) for DNA extraction and metabarcoding analysis. We identified and

discarded false positives using extraction blanks and multiple PCR replicates to avoid noise

from spurious amplification (Taberlet et al., 2018, Table 1). We used the following primer pairs

for specific taxonomic groups (Table 1): for plants, we amplified (i) the P6 loop of the chloroplast

trnL (UAA) intron using primers P6-trnLF: 5'-GGG CAA TCC TGA GCC AA-3' and p6-trnLR: 5'-

CCA TTG AGT CTC TGC ACC TAT C-3' (Taberlet et al 2007), and (ii) the internal transcribed

spacer 2 (ITS2) of nuclear ribosomal DNA using primers S2F: 5'-

ATGCGATACTTGGTGTGAAT-3' and S2R: 5'- GACGCTTCTCCAGACTACAAT-3' (Chen et al

2010); for eukaryotic algae and cyanobacteria, we amplified domain V of the 23S plastid rRNA

gene using primers p23SrV_f1 5'-GGA CAG AAA GAC CCT ATG AA-3' and p23SrV_r1 5'-TCA

GCC TGT TAT CCC TAG AG-3' (Sherwood & Presting, 2007); for fungi (Ascomycota and

Basidiomycota), we amplified the internal transcribed spacer (ITS) of nuclear ribosomal DNA

using primers: ITS1-F 5'-CTT GGT CAT TTA GAG GAA GTA A-3' (Gardes & Bruns, 1993) and

5'-GCT GCG TTC TTC ATC GAT GC-3' (White et al 1990); for metazoans, we amplified the

cytochrome c oxidase subunit I using primers: mlCOIintF: 5'-GGW ACW GGW TGA ACW GTW

TAY CCY CC-3' (Leray et al., 2013) and jgHCO2198: 5'-TAI ACY TCI GGR TGI CCR AAR AAY

CA-3' (Geller et al., 2013); and for invertebrates, we amplified the cytochrome c oxidase subunit

I using primers: fwhF2: 5'-GGD ACW GGW TGA ACW GTW TAY CCH CC-3' (Vamos et al.,

2017) and EPTDr2n: 5'-CAA ACA AAT ARD GGT ATT CGD TY-3' (Leese et al., 2021).

2.5　Bioinformatic processing – metabarcoding

Reads were analyzed separately for each primer in R v 4.3.1 (R Core Team, 2023) using

RStudio 2023.12.1 (RStudio Team, 2023) in dada2 (Callahan *et al*. 2016), after filtering (maxEE

= 2, Q-scores >30; Edgar & Flyvbjerg, 2015) and discarding reads with Ns. Error model

calculation, read correction, and read merging and removal of chimeric sequences was

performed using default settings. Amplicon sequence variants (ASVs) identified by dada2 were

assigned to taxa of origin using BLAST+ (v.2.6.0) (Altschul et al., 1990) and the GenBank

database.

2.6 Stable isotope analysis

Ethanol-preserved liver tissues were rinsed in distilled water and freeze-dried for 48 h. Once

freeze-dried, samples were ground to a fine powder using a laboratory bead-beater. Ground

samples were weighed in 8 x 5 mm pressed standard weight tin capsules using a high precision

microbalance (repeatability = 0.0008 mg). Elemental percentages of carbon, nitrogen, sulfur,

and stable isotope ratios ($\delta^{13}C$, $\delta^{15}N$ and $\delta^{34}S$) were measured using a Pyrocube elemental

analyser (Elementar, Langenselbold, Germany) linked to a visION continuous-flow isotope ratio

mass spectrometer (Elementar, Langenselbold, Germany) at the Universidad de Antofagasta

Stable Isotope Facility (UASIF), Chile. Stable isotope ratios are expressed using δ notation and

are reported in units of per mil (‰) relative to the following standards: Vienna Pee Dee

Belemnite for carbon, air for nitrogen, and Vienna Canyon Diablo Troilite for sulfur. International

standards were used in each batch to provide a multi-point calibration using the ionOS software

package v4.1.005 (Elementar, Langenselbold, Germany). Certified reference material USGS40

and USGS41a were used for carbon and nitrogen and IAEA-SO-5, IAEA-SO-6 and IAEA-S2 for

sulfur. Repeated analysis of standards showed analytical errors (± 1 SD) of ± 0.04 ‰ for $\delta^{13}C$, ±

0.06 ‰ for $\delta^{15}N$, and ± 0.6 ‰ for $\delta^{34}S$. We used two calibration standards, a) sulfonamide

(Elementar, Germany) and b) an in-house standard (rainbow trout dorsal muscle) to correct for

instrument drift.

2.7 Statistical analyses – stable isotopes

Liver is commonly used as a lipid-storage organ in vertebrates and liver lipid content can vary

significantly among individuals according to variation in nutritional state and physiological

condition. Lipids formed through *de novo* biosynthesis are isotopically lighter in $\delta^{13}C$ values

compared to proteins and the dietary sources from which they were formed (DeNiro & Epstein,

1977). If the isotopic effect of these lipids is not accounted for, they can affect assessment of

consumer $\delta^{13}C$ values. Furthermore, as the livers analyzed here were preserved in ethanol,

they may have undergone some partial uncontrolled lipid extraction prior to analysis. Variation in

individual lipid content can affect comparisons of $\delta^{13}C$ values and it is therefore common to use

chemical treatments to remove lipids prior to stable isotope analyses. However, the chemical

treatment can affect estimated values of other stable isotopes from the same sample. Another

possible solution is to use an arithmetic correction that relies on a predictable relationship

between lipid content and the elemental ratio between carbon and nitrogen (C:N) in the sample

(Kiljunen et al., 2006; Logan et al., 2008). Javornik et al. (2019) found small effects of ethanol

storage on $\delta^{13}C$ values in mammalian liver but reported no preservation effects on $\delta^{15}N$ or $\delta^{34}S$

values. Javornik et al. (2019) reported that C:N values decreased after ethanol storage but

suggested that lipid-free ^{13}C values could be reliably estimated mathematically from the C:N

ratio. In our samples, liver C:N ratios varied considerably (range = 3.2-5.7, mean ± SD = 3.8 ±

0.5, n = 41). Liver C:N ratios were lower in mice captured at higher elevations (r = -0.36, n = 40,

P = 0.024). As there was also a negative relationship between C:N and $\delta^{13}C$ within samples

from the same collection locality, we estimated lipid-corrected $\delta^{13}C$ values using Equation 1a

from Logan et al. (2008), resulting in a mean (± SD) isotopic shift of 1.1 ± 0.6‰. All $\delta^{13}C$ data

that we report are lipid-corrected, but liver $\delta^{15}N$ and $\delta^{34}S$ data are shown without correction.

264   Collection sites spanned ~4400 m of elevation and therefore exhibited considerable

variation in vegetation cover and plant species composition. We therefore examined how stable

isotopes varied within the dataset by plotting each of the stable isotopes against elevation. We

then used PERMANOVA (non-parametric permutation-based equivalent of ANOVA) to examine

whether stable isotope values varied among capture sites. Since we collected a single individual

from site 7 (the summit of Llullaillaco), it was not included in these comparisons. Although

PERMANOVA is typically used for multivariate comparisons (MANOVA), it can also be used to

make robust univariate comparisons. Finally, to assess the ability of the stable isotope analysis

to assign mice to capture location, and to identify mismatches that may be indicative of recent

dispersal, we used canonical analysis of principal coordinates (CAP), a distance-based

equivalent of discriminant function analysis (Anderson & Willis, 2003). This approach uses

multivariate data (e.g., $\delta^{13}C$, $\delta^{15}N$ and $\delta^{34}S$ values) to discriminate between groups defined by

elevation of capture sites. This approach also allowed us to infer the possible origin of the

summit mouse from Volcán Llullaillaco (site 7). We grouped mice in bins based on their capture

elevation (2000-3000 m, 3000-4000 m, 4000-5000 m, and >5000 m) and we used $\delta^{13}C$, $\delta^{15}N$

and $\delta^{34}S$ as dependent variables. We used a leave-one-out classification approach to examine

relative classification success, and we then used the model to identify the elevational range that

provided the best match to values from the Llullaillaco summit mouse. The ability of the CAP

model to statistically discriminate between groups was estimated via permutation (n = 9999).

PERMANOVA and CAP were both run in the PERMANOVA+1 add on to PRIMER 7 (Anderson

et al., 2008; Clarke & Gorley, 2015).

285   We estimated the trophic position of *P. vaccarum* at each site using liver $\delta^{15}N$ values

with those of primary producers collected across a similar (but truncated) elevational range

(Díaz et al., 2016). This approach (Cabana and Rasmussen 1996) allows the indirect calculation

of consumer trophic position (TP): TP = λ + ($\delta^{15}N_{Consumer}$ − $\delta^{15}N_{Baseline}$)/TDF, where λ is the

trophic position of the baseline taxon, $\delta^{15}N_{Consumer}$ is the nitrogen isotopic value of mice at a

given site, $\delta^{15}N_{Baseline}$ is the nitrogen isotopic value of the baseline at that site, and TDF is the

mean ± SD nitrogen trophic enrichment factor (TDF) for mouse liver (here we use 4.3 ± 0.2 ‰

from Arneson and MacAvoy (2005)). We used plants as our baseline (λ =1) based on data from

Díaz et al. (2016), which were collected in the same region as our study, over an elevational

range of 2670-4480 m. Plant $\delta^{13}C$ and $\delta^{15}N$ values exhibited considerable variation across

elevations (Figure 6), and we placed plants into broad elevational intervals (2000-3000 m, 3000-

4000 m, 4000-5000 m, and >5000 m). We then used values from the closest elevational interval

to estimate mouse trophic position at each capture site using *tRophicPostion* 0.8.0 (Quezada-

Romegialli et al., 2018) in R 4.2.3 (López-Cortés et al., 2007; R Core Team, 2023). Briefly,

tRophicPosition uses a Bayesian approach to estimate trophic position for a population of

consumers while accounting for variation in consumer and baseline isotope values. For most

sites we use the *onebaseline* model (assuming a single baseline) but we used the *twoBaselines*

model for mice from site 2. This is because the plants from the 3000 – 4000 m interval showed

a bimodal distribution of $\delta^{13}C$ values, which indicates the presence of plants using different

photosynthetic pathways (e.g. C3, C4/CAM). Since these groups also showed evidence for a

non-normal distribution of $\delta^{15}N$ values, we used the twoBaselines full model, which also uses

baseline $\delta^{13}C$. For all model runs we used the following parameters: chains = 3, number of

adaptive iterations = 1 000, iterations = 20 000, burn-in = 1 000, thinning = 10. In case of the

summit mouse from Volcán Llullaillaco we developed an individual model to calculate trophic

position, as *tRophicPosition* v 0.8.0 currently provides only population-level estimates of TP.

This new model with a one baseline approach was implemented in *greta* (Golding, 2019) which

allows the calculation of TP at the individual level. We modelled the baseline for the summit

mouse as having a mean and standard deviation of $\delta^{15}N$ values of plants >5,000 m with a

normal distribution for the mean and a Cauchy distribution for the SD, with a location of plants

$\delta^{15}N$ SD a scale of 3 and truncated from 0 to infinite. In this analysis λ is 1, the TDF was

modelled as having a normal distribution with a mean of 4.3 and SD of 0.2 (Arneson &

MacAvoy, 2005). We calculated 10 000 samples, with a thinning of 10, 1 000 samples as

warmup and 16 chains.

Due to the selective retention of heavier isotopes during the assimilation of food,

consumers are typically isotopically 'heavier' than their food (DeNiro & Epstein, 1978; DeNiro &

Epstein, 1981). These diet-tissue shifts are referred to as trophic enrichment or trophic

discriminations, and are typically estimated in experimental settings. Arneson and MacAvoy

(2005) provided empirical estimates for trophic discrimination factors (TDFs) in liver from groups

of laboratory mice fed diets that differed in the origin of their protein and carbohydrate

components. In their control diet, where carbohydrates and proteins originated from the same

source, mean ± SD TDFs were 0.7 ± 0.3 ‰ for carbon ($\Delta^{13}C$), 4.3 ± 0.2 ‰ for nitrogen ($\Delta^{15}N$),

and -2.1 ± 0.1 ‰ for sulfur ($\Delta^{34}S$). As such, we expect mouse livers to have $\delta^{13}C$ and $\delta^{15}N$

values that are ~1% and ~4 % higher, respectively, than their long-term average, in combination

with $\delta^{34}S$ values ~2 % lower than the long-term average.

3. RESULTS AND DISCUSSION

3.1 Metagenomics

For the stomach DNA sample of the Llullaillaco summit mouse, we sequenced a total of

423,477,275 reads, yielding 127,043 Mbases, with 92.48% of reads ≥ q30 and a mean quality

score of 35.69. We assembled a total of 9,138 contigs ≥1,000 bp in length (21,188,348 bp), with

a maximum length of 103,345 bp. Out of 991 contigs that were identified at the order level and

above, the vast majority were assigned to super kingdom Bacteria (699 contigs), with

Proteobacteria (563 contigs) and Firmicutes (99) as the dominant phyla. Only 1.3% of contigs

(*n*=13) were assigned to plants (clade Streptophyta, class Magnoliopsida [= dicotyledons]), and

all were assigned to a single representative of the coca family, Erythroxylaceae (*Erythroxylum*

novagranatense). This shrub species is widely cultivated in South America because its leaves

are a rich source of the psychoactive alkaloid, cocaine. In the stomach contents of the summit

mouse, we detected no traces of DNA from arthropods nor from vicuña, guanaco, or other

potentially co-distributed Andean mammals.

3.2 Metabarcoding

We sequenced a total of 9,487,085 reads as part of the DNA metabarcoding analysis,

maintaining 3,196,911 non-chimeric reads after filtering, denoising, and merging reads for all

primer combinations and all samples (Table 1, Tables S1-S6). The P6 loop of the chloroplast

trnL (UAA) intron marker (Figure 3a,b) yielded an average of 56,507 ± 22,886 (SD) non-

chimeric processed reads for each of the dissected sections of the gastrointestinal tract.

Consistent with results of the metagenomic analysis, sequences assigned to the family

Erythroxylaceae (*Erythroxylum novagranatense*) predominated in samples from each section,

from the stomach to the C12 portion of the colon (Figure 3a). Sequences of *E. novagranatense*

represent 97.3% of the 886,078 sequences derived from all surveyed sections of the

gastrointestinal tract. In addition to representatives of Erythroxylaceae, we detected

representatives of Amaryllidaceae (0.4% of the total), Poaceae (0.3%), Pinaceae (0.3%), and

Malvaceae (0.1%), and Fabaceae and Juglandaceae, which together accounted for 0.05% of

the total (Figure 3b).

The internal transcribed spacer 2 (ITS2) marker yielded an average of 19,952 ± 7,762

non-chimeric reads for each section of the gastrointestinal tract, and Erythroxylaceae accounted

for 87.2% of the 279,336 total reads. Sequences from Pinaceae (6.1% of the total),

Amaryllidaceae (5.9%), Urticaceae (0.64%), Poaceae (0.15%) were also detected, along with

traces of Laureaceae, Fabaceae, Ranunculaceae and Hypericaceae, which together accounted

for 0.04% of total reads (Figure 3d).

The third marker that provided information for Streptophyta was domain V of the 23S

plastid rRNA gene which yielded 1,596 ± 671 reads on average for all sections of the

gastrointestinal tract. Again, Erythroxylaceae predominated, accounting for 98.2% of 22,352

total reads (Figure 3e). In addition to Erythroxylaceae and Amaryllidaceae (1.5% of total reads),

the next two most abundant taxa were Cyatheaceae and Brassicaceae, which together

accounted for <0.6% of total reads (Figure 3f).

For Fungi, the internal transcribed spacer (ITS) of the nuclear ribosomal DNA marker

detected a wide variety of Ascomycota and Basidiomycota orders and families (Figure 4) in the

414,990 read total, with an average of 29,642 ± 12,846 reads for each section of the

gastrointestinal tract. For Ascomycota, the most abundant families were Cladosporiacea (28.4%

of the total reads for this Phylum), Pleosporaceae (20.2% of the total), Sacharomycetaceae

(16.6% of the total) and Nectriaceae (14.1% of the total), while the remaining 24 families

represent 20.6% of the total, with Phaeococcomycetaceae and Parmeliaceae (lichen-associated

families) accounting for 1.6% of the total. For the Phylum Basidiocomycota, the orders

Agaricales (40.7% of total reads for this Phylum) and Polyporales (36.5% of the total) represent

the most abundant groups, whereas Psathyrellaceae (18.8% of the total), Polyporaceae (13.6%

of the total), Agaricaceae (12.0% of the total), and the families Meripilaceae, Fomitopsidaceae

and Hyphodermataceae (together accounting for 16.7% of the total) were the most abundant

groups across all sections of the gastrointestinal tract.

For Metazoans the marker cytochrome c oxidase subunit I identified 41 amplicon

sequence variants (ASV), all of which were assigned to *Phyllotis*; none were derived from

vicuña or guanaco. Nine ASV remain uncategorized. Finally, the marker cytochrome c oxidase

subunit I specifically developed for arthropods did not detect any ASV for the group.

3.3 Diet of the summit mouse

One of the most puzzling results of the metagenomic and metabarcoding analyses is the

predominance of Erythroxylaceae (the coca family) and Amaryllidaceae (the garlic family) in the

stomach, cecum, and all 12 of the independently analyzed sections of the colon (Figure 3). Two

representatives of Erythroxylaceae, *Erythroxylum argentinum* and *E. cuneifolia*, exist at

elevations below ~2000 m far to the east of Llullaillaco, but they do not occur in Andean desert

or dry puna habitats. Coca was widely used throughout the Incan empire and is still used in

indigenous Quechua and Aymara communities and, occasionally, by mountain climbers. Upon

summiting a particular peak, there is a custom (especially among Argentine climbers) of leaving

offerings to Pachamama, an Andean "Earth mother", Gaia-type deity. Such offerings are left at

the base of rock piles called 'apacheta' that serve as summit markers. A typical offering to

Pachamama is a sprinkling of coca leaves or a small bag of such leaves at the base of the

apacheta. This custom provides a ready explanation for the predominance of Erythroxylaceae in

the gut contents of the summit mouse, which had presumably encountered just such an offering

on the summit of Llullaillaco. The presence of garlic in the stomach contents of the summit

mouse has a similar explanation. In the Argentine province of Salta (where the western portion

of Llullaillaco is located), garlic is a traditional folk remedy for altitude sickness. Argentine

climbers are known to chew cloves of garlic during their ascent. As is the case with any

unchewed coca that climbers possess upon reaching the summit, it is also customary to leave

leftover cloves of garlic at the base of the summit apacheta. The predominance of both coca

and garlic in the gut contents of the Llullaillaco summit mouse suggest that climbers' offerings to

Pachamama on the summits of high Andean summits may sometimes serve as unintentional

offerings to opportunistic *Phyllotis* mice living in an extremely food-scarce environment.

Aside from Erthroxylaceae and Amaryllidaceae, we also detected DNA representative of

several plant families such as Fabaceae, Malvaceae, and Poaceae, that occur at high

elevations at the base of Volcán Llullaillaco and in the surrounding Altiplano (Arroyo et al., 1988,

1998; Luebert & Gajardo, 1999; Marticorena et al., 2004). Within Fabaceae, the herb *Astragalus*

pusillus was documented at elevations up to 4300 m on the flanks of Llullaillaco (Marticorena et

al., 2004). Within Malvaceae, several perennial herbs such as *Cristaria andicola*, *Nototriche*

auricoma, and *N. clandestina* occur at elevations between 4000-4500 m (Arroyo et al. 1998)

and *Cristaria andicola* was documented as the most abundant plant species in the diet of

Phyllotis at another altiplano study site in northern Chile (López-Cortés et al., 2007). Within

Poaceae (=Gramineae), bunch grasses in the genus *Calamagrostis* (recognized as *Deyeuxia* in

Arroyo et al. [1998] and Luebert & Gajardo [1999]) occur above 4000 m (Marticorena et al.,

2004) and are also known to be included in the diet of *Phyllotis* from the Chilean Altiplano

(López-Cortés et al. 2007). Although all of these plants seem plausible as potential sources of

food for the summit mouse, there are no records of vascular plants or other vegetation above

~5000 m on the flanks of Llullaillaco (Arroyo et al., 1988, 1998; Luebert & Gajardo, 1999;

Marticorena et al., 2004; Storz et al., 2024; Vimercati et al., 2019), although it is also true that

botanical surveys typically do not venture above such elevations, so we should be careful about

interpreting absence of evidence as evidence of absence.

Given that sequences representative of Fabaceae, Malvaceae, and Poaceae were

detected in the gut contents of the mouse captured at 6739 m, far above the apparent

elevational limits of those plant taxa, there are three possible explanations to consider: (1) plant

material is carried upslope by the wind and accumulates in sufficient quantities on the lee edge

of ridge lines and snowdrifts to provide a source of sustenance for high-elevation mice ('Aolian

deposits'; Antor, 1995; Spalding, 1979; Swan 1961, 1992); (2) the plants in question actually

occur at much higher elevations than previously thought (though they may be scarce and

cryptic); or (3) the mouse was not a full-time summit resident, but rather a transient sojourner

that had simply consumed the plant material at or near the base of the mountain some days

prior to its capture. The former two hypothesis cannot be rejected, since few systematic plant

surveys have been performed on Llullaillaco or other >6000 m volcanoes (Storz et al., 2024). In

assessing the plausibility of the third hypothesis, it is important to note that the ~1.6 km

elevational distance between the summit of Llullaillaco (6739 m) and the apparent vegetational

limit (~5000-5100 m; Storz et al., 2024) translates into a linear distance of ~5 km from any side

of the volcano. Summiting the volcano from the vegetation limit is roughly equivalent to a direct-

from-basecamp ascent, a feat that only the most elite mountain climbers could accomplish in a

single day. We cannot rule out the possibility that mice undergo upslope/downslope dispersal on

a seasonal basis, but such movements could certainly not occur on a daily basis. Moreover, in

addition to the live capture of the *P. vaccarum* specimen on the Llullailllaco summit at 6739 m,

video records and identification of active burrows of *P. vaccarum* between 6145 – 6205 m on

the same volcano, and the discovery of desiccated cadavers and skeletal remains of numerous

P. vaccarum on the summits of four different >6000 m volcanoes in the same mountain chain

(Halloy, 1991; Steppan et al., 2023; Storz et al., 2020, 2023, 2024) provide a consilience of

evidence suggesting that these extreme high-elevation mice are representative of resident

populations. We think it is more likely that potential food plants exist at higher elevations,

although they must be scarce and patchily distributed. The plausibility of this hypothesis is

supported by the surprising discovery of bryophytes growing in association with active volcanic

fumaroles near the summit of Volcán Socompa (Halloy, 1991), a 6051 m volcano located 47 km

northeast of Llullaillaco along the Argentina-Chile border.

Primer set ITS detected sequences from families of two lichen-associated fungi,

Phaeococcomycetaceae y Parmeliaceae (phylum Ascomycota), indicating that *Phyllotis* feeds

on saxicolous lichen, as has been documented in other arctic and alpine mammals during

periods of food scarcity (Conner, 1983; Seward, 2008; Richardson & Young, 1977). During the

Arctic Winter, cricetid rodents such as snow voles (*Chionomys nivalis*) and northern bog

lemmings (*Synaptomys borealis*) feed on tundra lichen (Richardson & Young, 1977). Likewise,

during Winter months in the high alpine, North American pikas (*Ochotona princeps*) feed on

lichens under the snowpack (Conner, 1983). Terricolous, arboreal, and saxicolous lichens are

an important component of the winter diet of Caribou (*Rangifer tarandus*) in the northern

Holarctic (Seaward, 2008) and arboreal lichens are an important component of the winter diet of

Yunnan snub-nosed monkeys (*Rhinopithecus bieti*) in montane coniferous forests at elevations

>4000 m (Kirkpatrick et al., 1998). Although lichens may serve as a seasonal or short-term

supplement to the normal diet of many mammals living in arctic and alpine environments, their

low nutritive value suggest that they are unlikely to represent a year-round dietary staple for

small mammals like *Phyllotis* that have high metabolic demands.

We found no strong support for the arthropod fallout hypothesis, as we did not detect

arthropod DNA in the gut contents of our summit mouse. In contrast to high-elevation pikas on

the Quinghai-Tibetan Plateau that feed on yak feces, we found no evidence for interspecific

coprophagy in our high-elevation *Phyllotis*, as indicated by the absence of metagenomic

sequence reads and *COI* barcodes matching vicuña, guanaco, or any other potentially co-

distributed mammals. The absence of such sequences also constitutes absence of evidence for

scavenging.

3.4 Stable isotope analysis

Mice showed considerable variation in values of all three of the stable isotopes examined

(Figure 5a-c, Table 2). Patterns included isotopic variation both among sites (indicating an

elevational effect) and within sites (indicating individual variation in foraging habits). In the total

dataset, $\delta^{13}C$ values of *P. vaccarum* (Figure 5a, Table 2) varied between -22.8 and -12.7 ‰,

with more ^{13}C-enriched values being recorded at lower elevations (<4000 m). However, mice

with relatively ^{13}C-depleted values were captured at both sites 1 and 2. There was strong

statistical support for inter-site differences in $\delta^{13}C$ (PERMANOVA: $PseudoF_{5,34} = 36.6$, $P_{9999\ perms}$

= 0.0001). *Post-hoc* comparisons indicated significant differences ($P < 0.05$) between $\delta^{13}C$

values for all sites apart from sites 3 and 4 ($t = 1.42$, $P = 0.13$), sites 3 and 6 ($t = 1.92$, $P = 0.05$),

and sites 4 and 5 ($t = 1.4$, $P = 0.14$). *Phyllotis vaccarum* liver $\delta^{15}N$ ranged between 6.2 and 22.8

‰ (Figure 5b, Table 2), with considerable variation among sites (PseudoF$_{5,34}$ = 39.3, P$_{9999\ perms}$ =

0.0001). Values were notably ^{15}N-enriched at site 1 but included one individual with relatively

low $\delta^{15}N$. *Post-hoc* comparisons showed overlap in $\delta^{15}N$ values at sites 2 and 6 ($t = 0.10$, $P =$

0.92), sites 3 and 4 ($t = 0.05$, $P = 0.97$), sites 3 and 6 ($t = 2.27$, $P = 0.05$), and sites 4 and 5 ($t =$

0.97, $P = 0.34$). In the cases of both $\delta^{13}C$ and $\delta^{15}N$, *P. vaccarum* showed a similar pattern of

^{13}C- and ^{15}N-enriched values at sites from lower elevations (< 4000 m) relative to individuals

captured between 4000 and 5000 m (Figure 6, Table 2). This contrasted with the pattern in $\delta^{34}S$

(Figure 5c, Table 2) where *P. vaccarum* showed less variation in general, with values ranging

between -2.5 and 2.4 ‰, but exhibited a positive shift between ^{34}S-depleted values at sites

<4000 m and ^{34}S-enriched values >4000 m. Values of $\delta^{34}S$ for mouse livers varied among sites

(PseudoF$_{5,34}$ = 25.3, P$_{9999\ perms}$ = 0.0001), although variation was lower than that observed for C

and N. *Post-hoc* comparisons indicated that $\delta^{34}S$ values were similar for mice captured from

sites 1 and 2 ($t = 0.93$, $P = 0.70$), sites 3 and 5 ($t = 1.39$, $P = 0.18$), sites 3 and 6 ($t = 0.67$, $P =$

0.52), sites 4 and 5 ($t = 1.35$, $P = 0.21$), sites 4 and 6 ($t = 1.8$, $P = 0.09$, and sites 5 and 6 ($t =$

0.84, $P = 0.45$).

Mice from the highest elevations did not exhibit negative $\delta^{15}N$ values, suggesting that

lichenivory is not especially common.

3.4.1 Assignment to capture elevations

As abiotic and biotic conditions (e.g. temperature, aridity, UV concentrations, plant nutrient

availability, soil organic content) change with elevation, so too do the biomass and community

composition of the primary producers (Díaz et al., 2019), with consequent changes in the

availability of food for consumers and isotopic shifts at the base of the food web (Díaz *et al.*

2016). Given the known elevational gradient in stable isotope values, we can expect that stable

isotope values will provide a means of identifying variation in habitat use among mice captured

at different elevations.

Analysis of the combined stable isotope dataset using CAP showed that mice could be

reliably assigned to broad elevational zones using individual $\delta^{13}C$, $\delta^{15}N$ and $\delta^{34}S$ values (Figure

5d) (CAP: Trace = 1.71, P = 0.0001). The leave-one-out classification cross validation (Table 3)

indicated that by using the three isotope ratios we could assign an individual mouse to a 1000 m

interval with ~85 ‰ success. The CAP model predicted that the 6739 m summit mouse from

Volcán Llullaillaco was isotopically most similar to mice captured from the 4000-5000 m interval

(Figure 5d). The CAP model also suggested that 4 other individuals had stable isotope values characteristic of elevational zones distinct from where they were captured (Table 3). Including the summit mouse, this indicates that ca. 15 ‰ of the *P. vaccarum* in the study area had stable isotope values suggestive of upslope or downslope dispersal.

3.4.2 Trophic Position

Modal estimates of *P. vaccarum* trophic position (TP) varied between capture sites (Table 4) and ranged from 1.9 at site 2 to 4.3 at site 1. The latter estimate is extremely high and reflects the very high $\delta^{15}N$ values from mice collected at site 1 (mean $\delta^{15}N$ = 19.2 ‰). Discounting the results from site 1, mouse trophic position estimates were generally similar across sites and were indicative of omnivory with modal values between 1.9 and 2.3 at sites 2 to 5. The modal estimate for mice at site 6 was slightly higher (TP = 3.4), but the credibility limits overlapped with those from all sites apart from site 2. TP values between 2-3 are indicative of an herbivorous diet that includes some animal prey. TP values >3, as seen at site 6, indicates a diet dominated by animal prey. The TP of the summit mouse from Volcán Llullaillaco was estimated as 2.2, quite close to that of mice from sites 3, 4 and 5.

4. CONCLUSIONS

A combination of metagenomic, metabarcoding, and stable isotope data provided new insights into the diet of *Phyllotis* mice living at extreme elevations that far surpass known vegetation limits. Stable isotope data revealed that *Phyllotis vaccarum* maintains a mainly omnivorous diet in all elevational zones, and elevational variation in diet reflects variation in vegetation composition and the extent to which the mice rely on animal prey. Estimates of trophic position based on isotopic data indicated that mice collected near apparent vegetation limits (~5100 m) on the flanks of Llullaillaco rely more heavily on animal prey than mice from lower elevations. Metagenomic and metabarcoding analyses of gut contents from the mouse from the summit of Llullaillaco (6739 m) revealed a strictly herbivorous diet. The absence of animal DNA suggests that mice at extreme elevations do not subsist on wind-blown arthropods or other animal material. The detection of DNA from lichen-associated fungi indicate that *Phyllotis* mice living above known vegetation limits may supplement their diet with saxicolous lichens, as observed for other arctic and alpine mammals during periods of food scarcity in the winter. However, measured $\delta^{15}N$ levels indicate that lichen is not an important dietary staple in mice native to any of the surveyed elevational zones. The metagenomic and metabarcoding data also produce a scientific conundrum: the gut contents of a mouse captured at 6739 m elevation contained DNA

from several families of native plants that are not known to occur above ~5000 m elevation.

Clearly we have more to learn about the elevational distributions of both the plants and the

mice. It is possible that some of the plants identified in the diet of the summit mouse exist at

higher elevations than previously supposed, but they exist beneath the snowpack or in other

cryptic microhabitats.

AUTHOR CONTRIBUTIONS

CQ-R, CH, and JFS designed the study, MQC, GD, and JFS performed the fieldwork, CQ-R and

CH performed data analysis, CQ-R, CH, and JFS wrote the initial draft of the manuscript, and all

authors read and approved it.

ACKNOWLEDGMENTS

We thank Mario Pérez-Mamani and Juan Carlos Briceño for assistance and companionship in

the field, José Urquizo for help with figures, and Cristina Dorador for logistical assistance.

FUNDING INFORMATION

This work was funded by grants to JFS from the National Institutes of Health (R01 HL159061),

National Science Foundation (OIA-1736249 and IOS-2114465), and National Geographic

Society (NGS-68495R-20) and a grant to GD from the Fondo Nacional de Desarrollo Científico

y Tecnológico (Fondecyt 1180366).

REFERENCES

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment

search tool. *Journal of Molecular Biology*, *215*(3), 403–410.

Anderson, M.J., Gorley, R.N. & Clarke, K.R. (2008) PERMANOVA+ for PRIMER: Guide to

Software and Statistical Methods. PRIMER-R, Plymouth, UK.

Anderson, M.J. & Willis, T.J. (2003) Canonical analysis of principal coordinates: A useful

method of constrained ordination for ecology. *Ecology*, *84*, 511-525.

Antor, R.J. (1995) The importance of arthropod fallout on snow patches for the foraging of high-

alpine birds. *Journal of Avian Biology*, *26*, 81-85.

Arneson, L.S. & MacAvoy, S.E. (2005) Carbon, nitrogen, and sulfur diet-tissue discrimination in

mouse tissues. *Canadian Journal of Zoology*, *83*, 989-995.

Arroyo, M.T.K., Castor, C., Marticorena, C., Muñoz, M., Cavieres, L., Matthei, O., Squeo, F.A.,

M. Grosjean, M., & Rodríguez, R. (1998) The flora of Llullaillaco National Park located in the

transitional winter-summer rainfall area of the northern Chilean Andes. *Gayana Botanica*, *55(2)*,

93-110.

Arroyo, M.T.K., Squeo, F.A., Armesto, J.J., Villagrán, C. (1988) Effects of aridity on plant

diversity in the northern Chilean Andes: Results of a natural experiment. *Annals of the Missouri*

Botanical Garden, *75(1)*, 55-78.

Bender, M.M. (1971) Variations in the 13C/12C ratios of plants in relation to the pathway of

photosynthetic carbon dioxide fixation. *Phytochemistry*, *10*, 1239-1244.

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M.,

Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler, G.,

Alekseyev, M. A., & Pevzner, P. A. (2012). SPAdes: A new genome assembly algorithm and its

applications to single-cell sequencing. *Journal of Computational Biology*, *19*(5), 455–477.

https://doi.org/10.1089/cmb.2012.0021

Biazrov, L.G. (2012) Stable nitrogen isotopes (d15N) in thalli of arid vagrant lichen

Xanthoparmelia camtschadalis across an altitudinal gradient of the Khangai Plateau of

Mongolia. *Contemporary Problems of Ecology*, *5*, 200-207.

Bozinovic, F., & Rosenmann, M. (1988) Comparative energetics of South American cricetid

rodents. *Comparative Biochemistry and Physiology A*, *91*, 195-202.

Cabana, G. & Rasmussen, J.B. (1996) Comparison of aquatic food chains using nitrogen

isotopes. *Proceedings of the National Academy of Science*, *93*, 10844-10847.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.

(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*,

13(7), 581–583. https://doi.org/10.1038/nmeth.3869

Cerling, T.E., Harris, J.M., MacFadden, B.J., Leakey, M.G., Quade, J., Eisenmann, V. &

Ehleringer, J.R. (1997) Global vegetation change through the Miocene/Pliocene boundary.

Nature, *389*, 153-158.

Chen, S., Yao, H., Han, J., Liu, C., Song, J., Shi, L., Zhu, Y., Ma, X., Gao, T., Pang, X., Luo, K.,

Li, Y., Li, X., Jia, X., Lin, Y., & Leon, C. (2010). Validation of the ITS2 region as a novel DNA

barcode for identifying medicinal plant species. *PLoS ONE*, *5*(1), 1–8.

https://doi.org/10.1371/journal.pone.0008613

Chua, P.Y.S., Crampton-Platt, A., Lammers, Y., Alsos, I.G., Boessenkool, S., & Bohmann, K.

(2021) Metagenomics: a viable tool for reconstructing herbivore diet. *Molecular Ecology*

Resources, *21*, 2249-2263.

Clarke, K.R. & Gorley, R.N. (2015) PRIMER v7: User Manual/Tutorial. PRIMER-E, Plymouth.

Conner, D.A. (1983) Seasonal changes in activity patterns and the adaptive value of haying in pikas (*Ochotona princeps*). *Canadian Journal of Zoology*, *61*, 411-416.

Dalerum, F. & Angerbjörn, A. (2005) Resolving temporal variation in vertebrate diets using naturally occurring stable isotopes. *Oecologia*, *144*, 647-658.

Deagle, B.E., Thomas, A.C., McInnes, J.C., Clarke, L.J., Vesterinen, E.J., Clare, E.L., Kartzinel, T.R., & Eveson, J.P. (2019) Counting with DNA in metabarcoding studies: How should we convert sequence reads to dietary data? *Molecular Ecology*, *28(2)*, 391-406.

DeNiro, M. & Epstein, S. (1977) Mechanism of carbon isotope fractionation associated with lipid synthesis. *Science*, *197*, 261-263.

DeNiro, M.J. & Epstein, S. (1978) Influence of diet on the distribution of carbon isotopes in animals. *Geochimica et Cosmochimica Acta*, *42*, 495-506.

DeNiro, M.J. & Epstein, S. (1981) Influence of diet on the distribution of nitrogen isotopes in animals. *Geochimica et Cosmochimica Acta*, *45*, 341-351.

Díaz, F.P., Frugone, M., Gutiérrez, R.A. & Latorre, C. (2016) Nitrogen cycling in an extreme hyperarid environment inferred from δ15N analyses of plants, soils and herbivore diet. *Scientific Reports*, *6*, 22226.

Díaz, F.P., Latorre, C., Carrasco-Puga, G., Wood, J.R., Wilmshurst, J.M., Soto, D.C., Cole, T.L. & Gutiérrez, R.A. (2019) Multiscale climate change impacts on plant diversity in the Atacama Desert. *Global Change Biology*, *25*, 1733-1745.

Edgar, R. C., & Flyvbjerg, H. (2015). Error filtering, pair assembly and error correction for next-generation sequencing reads. *Bioinformatics*, *31*(21), 3476–3482. https://doi.org/10.1093/bioinformatics/btv401

Fogel, M.L., Wooller, M.J., Cheeseman, J., Smallwood, B.J., Roberts, Q., Romero, I. & Meyers, M.J. (2008) Unusually negative nitrogen isotopic compositions d15N of mangroves and lichens in an oligotrophic, microbially-influenced ecosystem. *Biogeosciences*, *5*, 1693-1704.

Gardes, M., & Bruns, T. D. (1993). ITS primers with enhanced specificity for basidiomycetes - application to the identification of mycorrhizae and rusts. *Molecular Ecology*, *2*(2), 113–118. https://doi.org/10.1111/j.1365-294X.1993.tb00005.x

Geller, J., Meyer, C., Parker, M., & Hawk, H. (2013). Redesign of PCR primers for mitochondrial cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic surveys. *Molecular Ecology Resources*, *13*(5), 851–861.

Golding, N. (2019). greta: simple and scalable statistical modelling in R. *Journal of Open Source Software*, *4*(40), 1601. https://doi.org/10.21105/joss.01601

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: Quality assessment tool for

genome assemblies. *Bioinformatics*, *29*(8), 1072–1075.

https://doi.org/10.1093/bioinformatics/btt086

Halloy, S. (1991) Islands of life at 6000 m altitude: the environment of the highest autotrophic

communities on Earth (Socompa Volcano, Andes). *Arctic and Alpine Research*, *23(3)*, 247-262.

Harrod, C., Grey, J., McCarthy, T.K. & Morrissey, M. (2005) Stable isotope analyses provide

new insights into ecological plasticity in a mixohaline population of European eel. *Oecologia*,

144, 673-683.

Javornik, J., Hopkins, J.B., III, Zavadlav, S., Levanič, T., Lojen, S., Polak, T. & Jerina, K. (2019)

Effects of ethanol storage and lipids on stable isotope values in a large mammalian omnivore.

Journal of Mammalogy, *100*, 150-157.

Kiljunen, M., Grey, J., Sinisalo, T., Harrod, C., Immonen, H. & Jones, R.I. (2006) A revised

model for lipid-normalising d13C values from aquatic organisms, with implications for isotope

mixing models. *Journal of Applied Ecology*, *43*, 1213-1222.

Kirkpatrick, R.C., Long, Y.C., Zhong, T., Xia, L. (1998) Social organization and range use in the

Yunnan snub-nosed monkey *Rhinopithecus bieti*. *International Journal of Primatology*, *19*, 13-

51.

Krouse, H.R., Stewart, J.W.B. & Grinenko, V.A. (1991) Pedosphere and biosphere. *Stable

Isotopes in the Assessment of Natural and Anthropogenic Sulphur in the Environment* (eds H.R.

Krouse & V.A. Grinenko), pp. 267-306. John Wiley & Sons, Chichester, UK.

Lee, Y.I., Lim, H.S. & Yoon, H.I. (2009) Carbon and nitrogen isotope composition of vegetation

on King George Island, maritime Antarctic. *Polar Biology*, *32*, 1607-1615.

Leese, F., Sander, M., Buchner, D., Elbrecht, V., Haase, P., & Zizka, V. M. A. (2021). Improved

freshwater macroinvertebrate detection from environmental DNA through minimized nontarget

amplification. *Environmental DNA*, *3*(1), 261–276. https://doi.org/10.1002/edn3.177

Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., Boehm, J. T., &

Machida, R. J. (2013). A new versatile primer set targeting a short fragment of the mitochondrial

COI region for metabarcoding metazoan diversity: application for characterizing coral reef fish

gut contents. *Frontiers in Zoology*, *10*(1), 34. https://doi.org/10.1186/1742-9994-10-34

Logan, J.M., Jardine, T.D., Miller, T.J., Bunn, S.E., Cunjak, R.A. & Lutcavage, M.E. (2008) Lipid

corrections in carbon and nitrogen stable isotope analyses: comparison of chemical extraction

and modelling methods. *Journal of Animal Ecology*, *77*, 838-846.

López-Cortés, F., Cortés, A., Miranda, E. & Rau, J.R. (2007) Dietas de Abrothrix andinus,

Phyllotis xanthopygus (Rodentia) y *Lepus europaeus* (Lagomorpha) en un ambiente altoandino

de Chile. *Revista Chilena De Historia Natural*, *80*, 3-12.

Luebert, F., Gajardo, R. (1999) Adiciones y notas a la flora del Parque Nacional Llullaillaco

(Región de Antofagasta, Chile). *Noticiario Mensual del Museo Nacional de Historia Natural*, 3-6.

Marris, J., Hawke, D. & Glenny, D. (2019) Eating at high elevation: an herbivorous beetle from

alpine rock outcrops relies on ammonia-absorbing lichens. *Ecology*, *100*, e02598.

Marticorena, C., Pardo, V., Peñaloza, A., Negritto, M.A., Cavieres, L.A., & Parada, M. (2004)

Adiciones y notas a la flora del Parque Nacional Llullaillaco, II Región, Chile. *Gayana Botanica*,

61(2), 49-54.

Nielsen, H., Pilot, L., Grinenko, L.N., Grinenko, V.A., Lein, A.Y., Smith, J.W. & Pankina, R.G.

(1991) Lithospheric sources of sulphur. *Stable Isotopes in the Assessment of Natural and*

Anthropogenic Sulphur in the Environment (eds H.R. Krouse & V.A. Grinenko), pp. 65-132.

John Wiley & Sons, Chichester, UK.

Pinho, P., Barros, C., Augusto, S., Pereira, M.J., Máguas, C. & Branquinho, C. (2017) Using

nitrogen concentration and isotopic composition in lichens to spatially assess the relative

contribution of atmospheric nitrogen sources in complex landscapes. *Environmental Pollution*,

230, 632-638.

Pizzimenti, J.J. & De Salle, R. (1980) Dietary and morphometric variation in some Peruvian

rodent communities: the effect of feeding strategy on evolution. *Biological Journal of the*

Linnean Society, *13*, 263-285.

Quezada-Romegialli, C., Jackson, A.L., Hayden, B., Kahilainen, K.K., Lopes, C. & Harrod, C.

(2018) tRophicPosition, an R package for the Bayesian estimation of trophic position from

consumer stable isotope ratios. *Methods in Ecology and Evolution*, *9*, 1592-1599.

R Core Team (2023). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/.

RStudio Team (2023). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA.

http://www.rstudio.com/.

Richardson, D.H.S., & Young, C.M. (1977). *Lichens and vertebrates*. In: Seaward, MRD

(ed.), *Lichen Ecology*. Academic Press, London.

Ruperto, E.F., Menéndez, J., Taraborelli, P.A., Scattolón, F.O. & Sassi, P.L. (2023) Behavioral

responses of two small-sized rodents, *Phyllotis vaccarum* and *Abrothrix andina*, to energy

challenges of high-altitude habitats in the Andes Mountains. *Ethology*, *129*, 117-132.

Sassi, P.L., Cuevas, M.F., Menéndez, J. & Dacar, M.A. (2017) Feeding strategies of a small

mammal (*Phyllotis xanthopygus*, Rodentia Cricetidae) at diverse altitudes in the Central Andes,

Argentina. *Ethology, Ecology, & Evolution*, *29*, 351-366.

Seaward, M. R. D. (2008) Environmental role of lichens. In: Nash III, TH (ed.), *Lichen Biology,*

2nd Edition. Cambridge University Press, Cambridge.

Sherwood, A. R., & Presting, G. G. (2007). Universal primers amplify a 23S rDNA plastid marker

in eukaryotic algae and cyanobacteria. *Journal of Phycology*, *43*(3), 605–608.

https://doi.org/10.1111/j.1529-8817.2007.00341.x

Silva, S.I. (2005) Posiciones tróficas de pequeños mamíferos en Chile: una revisión. *Revista*

Chilena De Historia Natural, *78*, 589-599.

Spalding, J.B. (1979) The Aeolian ecology of White Mountain Peak, California: windblown insect

fauna. *Arctic and Alpine Research*, *11*, 83-94.

Speakman, J.R., Chi, Q., Oldakowski, L., Fu, H., Fletcher, Q.E., Hambly, C., Togo, J., Liu, X.,

Piertney, S.B., Wang, X., Zhang, L., Redman, P., Wang, L., Tang, G., Li, Y., Cui, J., Thomson,

P.J., Wang, Z., Glover, P., Robertson, O.C., Zhang, Y., & Wang, D. (2021) *Proceedings of the*

National Academy of Sciences USA, *118(30)*, e2100707118.

Stapleton, T.E., Weinstein, S.B., Greenhalgh, R., & Dearing, M. D. (2022) Successes and

limitations of quantitative diet metabarcoding in a small, herbivorous mammal. *Molecular*

Ecology Resources, *22 (7)*, 2573–2586. doi: 10.1111/1755-0998.13643

Steppan, S.J., Bowen, T., Bangs, M.R., Farson, M., Storz, J.F., Quiroga-Carmona, M., D'Elía, G.,

Vimercati, L., Dorador Ortz, C., Zimmerman, G., Schmidt, S.K. (2022) Evidence of a population

of leaf-eared mice (*Phyllotis vaccarum*) above 6000 m in the Andes and a survey of high-elevation

mammals. *Journal of Mammalogy*, *103*, 776-785.

Storz, J. F., Liphardt, S., Quiroga-Carmona, M., Bautista, N. M., Opazo, J. C., Wheeler, T. B.,

D'Elía, G., & Good, J. M. 2023. Genomic insights into the mystery of mouse mummies on the

summits of Atacama volcanoes. *Current Biology*, *33*, R1040-R1042.

Storz, J. F., Quiroga-Carmona, M., Liphardt, S., Herrera, N. D., Bautista, N. M., Opazo, J. C.,

Rico-Cernhorska, A., Salazar-Bravo, J., Good, J. M., & G. D'Elía. 2024. Extreme high-elevation

mammal surveys reveal unexpectedly high upper range limits of Andean mice. *American*

Naturalist, *203*, 726-735.

Storz, J. F., M. Quiroga-Carmona, J. C. Opazo, T. Bowen, M. Farson, S. J. Steppan, & G.

D'Elía. 2020. Discovery of the world's highest-dwelling mammal. *Proceedings of the National*

Academy of Sciences USA 117: 18169-18171.

Storz, J. F., & Scott, G. R. 2019. Life ascending: mechanism and process of physiological

adaptation to high-altitude hypoxia. *Annual Review of Ecology, Evolution, and Systematics, 50*,

22.1-22.24.

Swan, L.W. (1961) The ecology of the high Himalayas. *Scientific American*, *205*, 68-78.

Swan, L.W. (1992) The aolian biome. *BioScience*, *42(4)*, 262-270.

Szpak, P., White, C.D., Longstaffe, F.J., Millaire, J.-F. & Vásquez Sánchez, V.F. (2013) Carbon

and nitrogen isotopic survey of northern Peruvian plants: Baselines for paleodietary and

paleoecological studies. *PLoS One*, *8*, e53763.

Taberlet, P., Coissac, E., Pompanon, F., Gielly, L., Miquel, C., Valentini, A., Vermat, T.,

Corthier, G., Brochmann, C., & Willerslev, E. (2007). Power and limitations of the chloroplast

trnL (UAA) intron for plant DNA barcoding. *Nucleic Acids Research*, *35*(3).

https://doi.org/10.1093/nar/gkl938

Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). *Environmental DNA: For Biodiversity*

Research and Monitoring. Oxford University Press.

https://doi.org/10.1093/oso/9780198767220.001.0001

Thomas, S.M. & Crowther, T.W. (2015) Predicting rates of isotopic turnover across the animal

kingdom: a synthesis of existing data. *Journal of Animal Ecology*, *84*, 861-870.

Tozer, W.C., Hackell, D., Miers, D.B. & Silvester, W.B. (2005) Extreme isotopic depletion of

nitrogen in New Zealand lithophytes and epiphytes; the result of diffusive uptake of atmospheric

ammonia? *Oecologia*, *144*, 628-635.

Vamos, E., Elbrecht, V., & Leese, F. (2017). Short COI markers for freshwater

macroinvertebrate metabarcoding. *Metabarcoding and Metagenomics*, *1*, e14625.

https://doi.org/10.3897/mbmg.1.14625

Vanderklift, M. & Ponsard, S. (2003) Sources of variation in consumer-diet d15N enrichment: a

meta analysis. *Oecologia*, *136*, 169-182.

Vimercati, L., Solon, A.J., Krinsky, A., Arán, P., Porazinska, D.L., Darcy, J.L., Dorador, C., &

Schmidt, S.K. (2019) Nieves penitentes are a new hábitat for snow algae in one of the most

extreme high-elevation environments on Earth. *Arctic, Antarctic, and Alpine Research*, *51(1)*,

190-200. doi.org/10.1080/15230430.2019.1618115

White, T.J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal

ribosomal RNA Genes for phylogenetics. In W. T. Innis MA, Gelfand DH, Sninsky JJ (Ed.), *PCR*

- Protocols and Applications - A Laboratory Manual (Issue January, pp. 315–322). Academic

Press.

Table 1. Summary statistics for metabarcoding sequence reads for each of the dissected portions of the gastrointestinal tract of the *Phyllotis vaccarum* specimen captured on the summit of Volcán Llullaillaco (see Figure 2).

Sample	Input reads	Filtered	Denoised forward	Denoised reverse	Merged	Non-chimeric
Extraction blanks	2.009	1.173	1.150	1.142	13	13
Stomach	755.758	350.866	349.765	349.840	242.553	237.567
Cecum	742.138	321.525	320.278	319.642	229.464	226.427
C1	548.838	283.847	283.023	282.946	213.947	204.749
C2	745.252	370.547	369.633	369.303	271.283	261.741
C3	634.524	324.615	323.596	323.317	226.881	224.844
C4	551.463	254.049	253.346	253.147	206.205	203.786
C5	559.460	272.415	271.609	271.402	159.567	158.241
C6	624.732	301.630	300.660	300.059	193.624	188.859
C7	670.502	313.308	311.935	311.397	214.056	205.772
C8	760.648	397.214	395.953	395.543	287.349	282.642
C9	731.646	367.289	365.790	365.438	275.122	269.431
C10	658.965	318.646	317.186	316.856	215.254	212.705
C11	702.488	340.687	339.058	338.399	224.913	219.703
C12	798.662	390.642	388.897	388.447	304.740	300.431
TOTAL	**9.487.085**	**4.608.453**	**4.591.879**	**4.586.878**	**3.264.971**	**3.196.911**

Table 2. Summary statistics (n, mean ± SD) for *Phyllotis vaccarum* liver stable isotope analyses of carbon (values shown for δ¹³C and lipid-corrected δ¹³C), nitrogen (δ¹⁵N), and sulfur (δ³⁴S) and the elemental C:N ratio.

Site	Elevation (m)	n	$\delta^{13}C$ (‰)	Lipid corrected $\delta^{13}C$ (‰)	$\delta^{15}N$ (‰)	$\delta^{34}S$ (‰)	C:N
Site 1	2370	8	-18.3 (± 2.2)	-16.9 (±2.0)	19.2 (± 3.6)	-1.6 (± 0.8)	4.2 (± 0.8)
Site 2	3240	8	-16.2 (±1.8)	-14.9 (± 1.8)	10.1 (± 0.9)	-1.5 (± 0.7)	4.0 (± 0.6)
Site 3	4150	4	-22.6 (± 0.4)	-22 (± 0.2)	7.6 (± 0.8)	0.5 (± 1.3)	3.4 (± 0.1)
Site 4	4360	8	-22.8 (± 0.5)	-21.8 (± 0.2)	7.6 (± 1.0)	1.6 (±0.5)	3.8 (± 0.3)
Site 5	4620	5	-23.3 (±0.2)	-22.2 (± 0.4)	7.1 (± 0.1)	1.3 (± 0.1)	3.7 (± 0.2)
Site 6	5070	7	-22.0 (± 1.1)	-20.9 (± 0.9)	10.2 (± 2.1)	1.0 (±0.9)	3.8 (± 0.4)
Site 7	6739	1	-22.0 (–)	-21.5 (–)	7.0 (–)	2.0 (–)	3.3 (–)

Table 3. Results of the canonical analysis of principal coordinates (CAP) leave-one-out cross-validation to assess the ability of the model to assign individual mice to the 1000 m elevational zone in which they were captured.

Known capture altitude (m)	Predicted capture altitude (m)				‰ correctly classified
	2000-3000	3000-4000	4000-5000	5000-6000	
2000 – 3000	7	0	0	1	86
3000 – 4000	0	7	0	1	86
4000 – 5000	0	0	16	1	94
5000 – 6000	0	0	2	5	71

Table 4. Estimates of trophic position for mice captured in different elevational zones. Summary statistics are provided as modal TP (95 ‰ credibility intervals) and as mean TP ± SD.

	tRophicPosition model	Baseline data (Díaz et al., 2016)	Trophic position (mode, 95 ‰ credibility intervals)	Trophic position (mean ± SD)
Site 01	*oneBaseline*	C3 & CAM data combined from 2000 – 3000 m	[†]4.3 (3.4 – 5.1)	[*]4.3 ± 0.2
Site 02	*twoBaselines*	C3 & CAM/C4 data 3000 – 4000 m	1.9 (1.7 – 2.2)	1.9 ± 0.1
Site 03	*oneBaseline*	Plants >4000 m (all C3)	2.3 (1.8 – 2.9)	2.3 ± 0.3
Site 04	*oneBaseline*	Plants >4000 m (all C3)	2.3 (1.9 – 2.7)	2.3 ± 0.2
Site 05	*oneBaseline*	Plants >4000 m (all C3)	2.2 (1.8 – 2.5)	2.2 ± 0.2
Site 06	*oneBaseline*	Plants >4000 m (all C3)	2.9 (2.3 – 3.5)	2.9 ± 0.3
Summit mouse	*oneBaseline-greta*	Plants >4000 m (all C3)	2.2 (1.9 – 2.4)	2.2 ± 0.1

[†] $\delta^{15}N$ values of mice captured at Site 1 were unusually high and were markedly ^{15}N enriched relative to the putative baseline (see Figure 5b) and may be artefactual.

Figure Legends

Figure 1. Sampling of *Phyllotis vaccarum* across an elevational gradient in the Altiplano and Puna de Atacama of northern Chile, Región de Antofagasta. (a) Map of seven collection localities on the flanks of Volcán Lullaillaco and the surrounding Altiplano. (b) Elevational profile of sampling transect, with sampling localities 1 to 7 shown in ascending order of elevation, from 2370 m (site 1) to 6739 m (the summit of Volcán Lullaillaco, site 7). (c) Northwest face of Volcán Llullaillaco (24°43.21′S, 68°32.22′W). Photo was taken from a point ~15 km northwest of the summit. Photo: J.F. Storz.

Figure 2. Schematic figure of the dissected portions of the gastrointestinal tract of the mouse from the summit of Volcán Llullaillaco. Contents of the upper digestive tract (stomach) were analyzed via metagenomics and DNA metabarcoding. Contents of the lower digestive tract (separated into the cecum and 12 consecutive segments from the outlet of the cecum to the rectum) were analyzed via DNA metabarcoding.

Figure 3. Taxonomic composition of reads identified through metabarcoding for the stomach, cecum and 12 consecutive portions of the lower GI from the outlet of the cecum to the rectum. (A) Proportion of sequence reads per taxon for the marker *trn* and (B) proportion of total reads for the same marker with Erythroxylaceae excluded. (C) Proportion of sequence reads per taxon for the marker *ITS2* and (D) proportion of total reads for the same marker with Erythroxylaceae excluded. (E) Proportion of sequence reads per taxon for the marker *23S* and (F) proportion of total reads for the same marker with Erythroxylaceae excluded.

Figure 4. Taxonomic composition of Ascomycota (A) and Basidiomycota (B) at the Order and Family levels.

Figure 5. Variation in stable isotopes in livers of *Phyllotis vaccarum* sampled from different elevational zones: (A) $\delta^{13}C$, (B) $\delta^{15}N$, and (C) $\delta^{34}S$. (D) Results of multivariate CAP ordination based on Euclidean distances calculated from combined $\delta^{13}C$, $\delta^{15}N$ and $\delta^{34}S$ values.

Figure 6. Variation in plant $\delta^{13}C$ (A) and $\delta^{15}N$ (B) across different elevational zones. Values taken from Díaz et al. (2016). Note the elevational shift in $\delta^{13}C$ values showing dominance of C_4 plants at lower elevations, a mix of C3, C4 and CAM plants at mid-elevations and a shift to C3 plants at higher elevations. Plant $\delta^{15}N$ values were similar at lower and mid-elevations but were relatively ^{15}N depleted at higher elevations. These data were used to define isotopic baselines for estimates of trophic position of *P. vaccarum*.

(a)

CHILE

ARGENTINA

0 20 40 60 km

(b)

(c)

stomach

cecum

colon

Upper GI tract

Lower GI tract

1 2 3 4 5 6 7 8 9 10 11 12

A

B