Cerebellar systems consolidation driven by the temporal dynamics of Purkinje cell

excitability

Jewoo Seo[1,2], Yong Gyu Kim[1,2], Yong-Seok Lee [1,2,3*], Sang Jeong Kim[1,2,3*]

Author affiliations

[1]Department of Physiology, Seoul National University College of Medicine, Seoul 03080, Korea

[2]Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul 03080, Korea

[3]Memory Network Medical Research Center, Neuroscience Research Institute, Wide River Institute of Immunology, Seoul National University College of Medicine, Seoul 03080, Korea

*: Corresponding authors

Abstract

Systems consolidation, essential for long-term memory formation, orchestrates the

reorganization of newly encoded memories into subsequent neural circuitry. While the role of

synaptic mechanism in consolidation is well understood, the contribution of neuronal intrinsic

excitability (IE) remains relatively unexplored. Herein, we adopted a cerebellum-dependent

learning model, and manipulated IE of the sole output of the cerebellar cortex, Purkinje cells

(PCs), to corroborate the direct causality between neuronal IE and memory consolidation and

assessed its impact on intrinsic plasticity of the downstream neural circuits. Optogenetic

excitation disrupted consolidation but was effective only within the 90 min post-learning time

window. Notably, PC-IE temporarily weakened, but the effect faded beyond 90 minutes.

Furthermore, abnormally increased PC-IE abolished intrinsic plasticity in flocculus-targeting

neurons, a post-circuitry of cortical PCs, within the medial vestibular nucleus. These findings

thus emphasize the precise temporal dynamics of IE, highlighting it as a crucial mechanism

for systems consolidation.

Keywords

Cerebellum, Purkinje cells, flocculus-targeting neurons, intrinsic excitability, motor learning,

systems consolidation

Introduction

Systems memory consolidation is the process by which memory becomes permanent(1).

Memories initially encoded within the hippocampal entorhinal cortex are consolidated in the

neocortex for long-term storage(2). In the cerebellum, short-term memories(3-5) initially

acquired from motor learning and driven by sensory inputs from the external environment,

become mature and consolidated(3, 6, 7) through neural communication between GABAergic

inhibitory Purkinje cells (PCs)(8) in the flocculus, a lobe-like structure responsible for the

ocular reflex(9-12), and flocculus-targeting neurons (FTNs) in the medial vestibular

nuclei(13-17). Cerebellar systems consolidation accounts for the enhancement of

compensatory eye movements, such as the optokinetic reflex (OKR)(18, 19), associated with

various sensory modalities including visual inputs to ameliorate the stability of vision(20).

The canonical view of cerebellar learning and memory has long been centered on

bidirectional modifications in synaptic transmission, particularly the cellular models of long-

term potentiation and long-term depression, which are widely recognized as pivotal processes

underlying the formation and retention of memory(21-25). Empirical evidences suggest that

the attenuation of synaptic strength between parallel fibers and PCs plays a prominent role in

the consolidation of motor memory, allowing for the adaptation of cerebellum-dependent

behaviors(18, 19, 26, 27).

Recent studies have demonstrated that consolidation extends beyond synaptic

mechanisms and involves complementary non-synaptic mechanisms, such as intrinsic

excitability (IE), to achieve intact learning capabilities(28, 29). Its plasticity manifests

bidirectionally in response to a separate set of bidirectional modifications in synaptic

plasticity(30-32). This is indicative of homeostatic maintenance within the neural

circuitry(33) and   supports the notion that multiple mechanisms are involved in the

cerebellar memory system(6). Notably, OKR learning and eye-blink conditioning are

accompanied by the down-regulation and upregulation of the intrinsic excitability of PCs

(PC-IE), respectively(34, 35). The use of a transgenic mouse model with intact synaptic

transmission but deficient intrinsic plasticity resulted in impaired consolidation, despite

complete acquisition from learning(36, 37). In different regions such as the hippocampus,

increased cortical excitability can result from various tasks, including eye-blink and fear

conditioning(38, 39). Although numerous studies have established a correlation between the

level of intrinsic excitability and the quality of learning and memory, research exploring the

direct impact of neuronal IE on the formation of long-term memory remains limited. This

knowledge gap is particularly pronounced in the cerebellum, as the potential implications of

manipulating PC-IE on memory consolidation process remain unknown. In addition, although

evidence indicates fluctuations in PC-IE in a time-dependent manner following motor

learning(37), the temporal dependency of its contribution to consolidation remains unclear.

Therefore, it is imperative to establish causality between consolidation and PC-IE and to

revisit the timing of the high effectiveness of PC-IE for long-term storage.

Inter-regional communication between the cerebellar cortex and the nuclei underlies

systems consolidation(11, 40) and it has been suggested to be accompanied by PC-dependent

synaptic plasticity and intrinsic excitability of neurons in the MVN(37, 41). Therefore,

cortical PC is a plausible neural substrate for connecting two different brain regions.

However, the direct impact of changes in PC-IE on systems consolidation regarding the

intrinsic excitability of FTNs remain elusive.

In the present study, we hypothesized that PC-IE is essential for the maturation of motor

memories and the formation of neural links between the cortex and the nuclei during the

consolidation period. Through the application of neuron-specific optogenetic manipulations

in conjunction with the OKR paradigm, we present compelling evidence that PC-IE plays a

crucial determinant in facilitating memory consolidation, as we observed that upregulation of

cortical PC-IE within a specific time window result in impaired memory consolidation.

Moreover, we demonstrate that PC-IE manipulation compromises the intrinsic plasticity of

subsequent FTNs in the nucleus, thereby suggesting that the PC-FTN circuit may heavily

depend on PC-IE. Our findings provide a unique perspective on the non-synaptic mechanism

underlying the process of systems consolidation in the cerebellar circuit, which goes beyond

the traditional perspective of synaptic plasticity.

Results

Optogenetic manipulation of PC-IE increases spontaneous firing

Recent studies have demonstrated that the intrinsic excitability of PCs is strongly associated

with cerebellar motor learning(34, 35, 37), but yet, the actual causality between PC-IE and

motor memory consolidation through cell-type specific manipulation remains unexplored.

Hence, we devised an appropriate method for the selective manipulation of PC-IE in the

flocculus. To achieve this, we employed optogenetics using the stabilized step-function opsin

(SSFO)(42). It promptly elicits depolarization and stable excitatory state in neurons in

response to optostimulation. The experimental design involved the bilateral injection of

AAV1-Ef1a-DIO hChR2(C128S/D156A)-EYFP or AAV1-Ef1a-DIO EYFP into the flocculi

of Pcp2-cre (L7-Cre) mice, which allowed specific recombinase activity in the floccular PC

layer (Figure 1a). Following the surgery, an in-vitro whole-cell patch clamp recording of PCs

was conducted (Figure 1b). We validated that in response to 473 nm optostimulation, which

activates the opsin, the spontaneous firing of SSFO-expressing PCs was significantly elevated

(Figure 1c). Specifically, the average firing rate of SSFO-expressing PCs was robustly

enhanced, whereas EYFP-expressing PCs did not show any changes (Figure 1d).

Furthermore, upon 593 nm optostimulation, which deactivates the opsin, the elevated firing

rate of SSFO-expressing PCs reverted to the basal condition, and no change was detected in

EYFP-expressing PCs, as expected (Figure 1c, d). Moreover, our ex-vivo result indicated that

PCs of optostimulated SSFO-expressing mice following motor learning exhibited

significantly higher IE than that of EYFP-expressing mice (Supplementary Figure 1a, b).

These results establish the efficacy of the proposed approach for selectively manipulating PC-

IE during the consolidation period, providing a promising foundation for investigating the

causal relationship between PC-IE and cerebellar memory consolidation.

Optogenetic manipulation of PC-IE impairs memory consolidation

To understand the contribution of intrinsic excitability to memory consolidation, we

employed optogenetics during the behavioral assessment. Specifically, we utilized the OKR

protocol to train mice in adapting their ocular reflex in response to visual stimulation. To

manipulate the floccular PC excitability *in vivo* during the consolidation period, we

bilaterally implanted an optic cannula into the flocculi of mice that had undergone surgery for

SSFO/EYFP expression. One mouse underwent 2 days of training. On the first day, it was

trained for 50 min and placed back into the home-cage in a completely dark condition to rest

without any visual interruption (Figure 2a, top panel). During the training session, an animal

was subjected to continuous oscillation ($\pm 5°$ peak-to-peak motion at 0.5 Hz) of the

optokinetic screen designed by vertical stripe patterns (Figure 2b). Once training was

completed, bilateral 473 nm optostimulation was administered at 0 min post-learning to

activate the SSFO and another set of bilateral 593 nm optostimulation was provided at 24 hr

post-learning to deactivate the SSFO (Figure 2a, bottom panel). Proper training reduced the

retinal slip, which directly reflected the accomplishment of motor learning in both the SSFO

and EYFP groups (Figure 2c, pre- and post-learning). When mice were exposed to

continuous sinusoidal screen oscillation before training, the gain, a value representing how

closely the eye motion overlaps with the rotation of a screen, ranged between 0.22 and 0.58

(EYFP = 0.235-0.576; SSFO = 0.217-0.557); after training, the gains were increased ranging

between 0.38 and 0.84 (EYFP = 0.571-0.746; SSFO = 0.381-0.838) (Figure 2d). Improved

oculomotor performance was observed in both groups. After confirming the learning results,

we activated the SSFO during the initial phase of the consolidation period. Thus, we were

able to instantly disrupt the learning-induced depression of intrinsic excitability that is

reported to be promptly exhibited after cerebellar motor learning(34, 37). As a result, long-

term memory was normally formed in the EYFP group after 24 hr of learning, but was

abolished in the SSFO group due to aberrant conditions in the neural circuitry for

consolidation (Figure 2d). Specifically, the consolidation percentage of motor memory in the

SSFO group was significantly lower than that in control group (Figure 2e). To investigate the

potential disruption of memory consolidation attributed to the mere expression of SSFO, we

conducted an additional set of OKR learning using mice expressing SSFO. Subsequently,

these mice, which had previously showed impaired consolidation with optogenetic

manipulation, underwent for a recovery period of 2 to 3 weeks. Mice with relatively high

gains before the second set of learning, indicative of the remaining effects from the first

learning set, were excluded from the second assessment. When SSFO-expressing mice were

trained and subjected to a consolidation period without interference from optogenetic

manipulation, their long-term memory was steadily retained (Supplementary Figure 2a, b).

Therefore, these results reveal the importance of the proper expression of PC-IE in

facilitating memory consolidation.

PC-IE contributes to the formation of long-term memory during the early phase of the

consolidation period

In addition to OKR learning, long-term memories of other cerebellum-dependent behaviors,

such as the vestibulo-ocular reflex (VOR) and eye-blink conditioning are susceptible to

interference within specific periods of consolidation. When the intracortical activity was

disrupted between 45 and 90 min after eye-blink conditioning, the rabbits were unable to

properly express retained long-term memory(43). Similar to OKR, VOR requires the

cerebellum for visual stabilization when stimulated by vestibular inputs. After the cats were

trained for the VOR, they were subsequently exposed to a condition in which their vision was

being constantly interrupted during the first hour of the consolidation period. Following the

visual interruption, the cats exhibited reverted learning outcomes(44). In addition,

consolidation was prevented when G-protein coupled receptors that are involved in learning

and memory were pharmaceutically suppressed within 1hr after the learning period(45).

Building upon the findings from previous studies, the primary objective of this study was

to ascertain whether a critical time window exists during which PC-IE plays a pivotal and

highly effective role in memory consolidation. Given our observation that optogenetic

manipulation throughout the entire consolidation period resulted in the loss of long-term

memory, we implemented two distinct behavioral schemes involving different time points of

optogenetic manipulation. Some mice were given 473 nm optostimulation at 0 min post-

learning (Figure 3a, *Scheme #1*), and a break of approximately 2 to 3 weeks was provided to

ensure the extinction of previously acquired motor memory for a revised scheme of the next

experiment. Subsequently, they underwent a second trial of learning, with optostimulation at

90 min post-learning (Figure 3a, *Scheme #2*). The order of manipulation was counterbalanced

to eliminate any potential bias. Consequently, regardless of the behavioral scheme, the EYFP

group demonstrated robustly sustained long-term memory, as shown by the high

consolidation percentage, in both tests (Figure 3b, c). However, the SSFO group exhibited

varying outcomes based on the timing of the manipulation. When SSFO was activated to

increase PC-IE at 0 min post-learning, the previously enhanced gain levels were reversed,

resulting in a relatively low consolidation percentage (Figures 3d [left] and 3e). By contrast,

when the manipulation was administered 90 min after learning, the increased gain was

maintained, resulting in a high consolidation percentage (Figures 3d [right] and 3e). Outside

this critical time window, alterations in PC-IE did not significantly affect formation of long-

term memory, emphasizing the pivotal role of the timing of PC-IE contribution to memory

consolidation. Furthermore, we substantiated the significance of 90 min post-learning period

by optogenetically manipulating PC-IE exclusively within this initial timeframe. Despite any

interference, the EYFP group exhibited intact memory consolidation (Figures 3f [left] and

3g), but, as anticipated, the SSFO group was incapable of retaining long-term memory

(Figure 3f [right] and 3g), highlighting that the induction of aberrant PC-IE was powerful

enough to disturb the nature of learning and memory. These compelling results clearly

demonstrate that PC-IE exercises physiological leverage to facilitate memory consolidation

within 90 min post-learning period.

Dynamics of PC-IE within the critical temporal window following motor learning

Under general circumstances, PC-IE remains constant, but according to a previous study,

VOR learning induces the temporal adjustment of PC-IE throughout the consolidation period

in a bidirectional manner, which consists of decreased or restored levels of excitability

depending on the timeline(34, 37). Kim et al. (2020) found that OKR learning reduces PC-IE

immediately after the completion of motor learning(34). However, the fluctuation of

excitability at various time points has not been thoroughly elaborated. Therefore, it remains

unclear how PC-IE dynamics are exhibited after OKR learning, particularly within the

observed critical time window. To address this question, we investigated the underlying

cellular mechanisms that establish a critical time window. Once the mice were fully trained

for OKR, we prepared brain slices to perform whole-cell patch clamp recordings from

floccular PCs at various post-learning periods, which allowing us to explore the dynamics of

PC-IE during consolidation (Figure 4a). Some slices were acquired at 0 min post-learning,

while others were prepared at 90 min post-learning. To evaluate PC-IE, we provided current

injections from +0 pA to +900 pA (100 pA increment per injection, 500 ms long) in the

presence of NBQX (2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo[*f*]quinoxaline) and picrotoxin

to block excitatory and inhibitory synaptic inputs, respectively. The results demonstrated that

the mean firing rate of floccular PCs was significantly reduced compared with that of naïve

control conditions, but after 90 min post-learning, the rate recovered to that of the naïve

condition (Figure 4b, c). This indicates that the depression of PC-IE did not last for more than

90 min.

Furthermore, we analyzed the changes in action potential (AP) properties to explain the

depression of PC-IE. We measured four AP parameters: the rheobase current, AP threshold,

AP amplitude and afterhyperpolarization (AHP) amplitude. Among these parameters, only

the rheobase current exhibited significant changes. It was significantly increased at 0 min

post-learning, but lasted temporarily such that the increased current was reduced back to the

basal level at 90 min post-learning (Figures 4b, c). This implies that, within 90 min post-

learning period PC-IE decreased and required a larger current threshold to fire an AP. Beside

the rheobase current, other properties were not altered following OKR learning (Figure 4d-f).

To assess whether our optogenetic manipulation at 0 min post-learning induced an abnormal

condition, thereby blocking the learning-induced depression in PC-IE. The ex-vivo

results revealed that PCs of optostimulated SSFO-expressing mice, following motor learning,

exhibited significantly higher firing rates than those of EYFP-expressing mice (Supplementary

Figure 1a, b). This heightened activity directly signifies the abolished depression in PC-IE of the

SSFO group. Consequently, the loss of long-term memory can be attributed to the prevention of

learning-induced depression in PC-IE, a process anticipated to occur after learning.

Disrupted plasticity of intrinsic excitability of flocculus-targeting neurons in the MVN

Postsynaptic circuitry of cortical PCs is primarily organized by neurons in the MVN,

characterized by two distinct types: flocculus-targeting neurons (FTNs) and nonFTNs(13,

46). Among these types, FTNs, specifically, play an active role in facilitating memory

consolidation(11, 16, 17, 40), as they serve as the subsequent destinations for memory

transfer. The ability of FTNs to maintain and express long-term memory is supplemented by

both synaptic and intrinsic plasticities(41, 47). Previous studies reported that not only

intrinsic plasticity of PC was elicited following motor learning, but also a potentiation of

intrinsic excitability was induced in VN neurons(37). Therefore, we investigated whether

optogenetic manipulation of PC-IE directly influences the expression of intrinsic plasticity of

FTNs. Pcp2-cre (L7-Cre) mice underwent surgery for SSFO and EYFP injections into the

flocculus. These mice underwent OKR learning, followed by optostimulation to activate

SSFO for the entire period of consolidation, and eventually, acute slices were obtained at 24

hr post-learning period for whole-cell patch clamp recording (Figure 5a, top). Rich

expression of SSFO/EYFP in floccular PCs allowed for the morphological distinction of

FTNs and nonFTNs, indicated by axon terminals with multiple boutons en passant densely

surrounding neurons in MVN (Figures 5a [bottom] and 5b).

To further investigate whether learning-induced plasticity of intrinsic excitability occurs

in the FTNs, we measured the mean firing rates of the EYFP and SSFO groups under both

naïve and learned conditions. Comparisons revealed distinct patterns. In EYFP-expressing

mice, intact memory consolidation post-learning was accompanied by the potentiation of

intrinsic excitability in *in-vitro* (Figures 5c, d). Conversely, in SSFO-expressing mice, the

abnormal increase in PC-IE resulted in impaired memory consolidation and the absence of

intrinsic plasticity in FTNs (Figures 5e, f). Upon completion of the learning and subsequent

consolidation period, significant disparities in mean firing rates between EYFP-expressing

and SSFO-expressing FTNs were particularly evident during 100 pA and 150 pA stimulations

(Figure 5g).

Turning our attention toward nonFTN circuitry, which remains isolated from the cortical

learning circuit, we speculated that there would be no discernible learning-induced plasticity

in nonFTNs following motor learning. As anticipated, motor learning did not elicit intrinsic

plasticity in nonFTNs in either the EYFP or SSFO groups, as there was no significant

difference in the mean firing rates (Supplementary Figures 3a-d). Unlike FTNs, nonFTNs in

both the EYFP and SSFO groups showed comparable excitability following motor learning

(Supplementary Figure 3e). Overall, these data clearly demonstrate that the proper

maintenance of long-term memory involves the expression of intrinsic plasticities in both PCs

and FTNs, forming the foundation for systems consolidation, which, in turn requires inter-

regional communication.

Discussion

Conventional theories of learning and memory have extensively focused on bidirectional

changes in synaptic efficacy at multiple sites within the cerebellar circuitry(6). However, in

addition to these changes in synaptic function, recent evidence has indicated that cerebellar

motor learning entails a complex neurobiological process involving an intricate interplay of

both synaptic and intrinsic plasticity mechanisms. Despite this, the contribution of intrinsic

excitability, which reflects the ability of neurons to generate APs independent of synaptic

input(28), remains poorly understood. To address this knowledge gap, we employed a

comprehensive approach that integrates optogenetics, behavioral assays, and

electrophysiological recordings. Optogenetic manipulation of PC-IE resulted in an aberrant

increase in neuronal excitability and spontaneous firing, resulting in a discernible impairment

in the consolidation of motor memory. This finding strongly indicates a disruption in the

nature of cerebellar learning circuitry. As a result, it unequivocally highlights the essential

role played by PC-IE in the process of memory consolidation, thereby emphasizing the

reliance of the cerebellar learning system on a coordinated interplay of not just synaptic

plasticity but also intrinsic plasticity mechanisms. Notably, the level of PC-IE fluctuates

throughout the consolidation period following VOR. Interestingly, it has been only reported

that PC-IE reduces shortly after OKR(34, 37). Hence, we further conducted a detailed

examination of the dynamics of PC-IE following OKR at different time points, uncovering

enduring changes in excitability over an extended period after learning. A pronounced

decrease in PC-IE underscores an important temporal window that decisively influences the

achievement of memory consolidation. Furthermore, motor learning reshapes the cortical IE

and extensively modifies the post circuitry of floccular PCs, which are primarily organized by

FTNs in the MVN(13-17). Importantly, the abnormal expression of PC-IE not only hinders

memory consolidation but also prevents the induction of intrinsic plasticity of FTNs in the

nuclei. This suggests that inter-regional coupling, which is crucial for systems consolidation

between the cortex and the nuclei, may be facilitated by enduring changes in intrinsic

excitability. This concept aligns with the hypothesis that intrinsic plasticity may function as a

trigger or conductor of systems-level consolidation within the circuits involving the medial

prefrontal cortex-basolateral amygdala, and hippocampus(48, 49).

The involvement of intrinsic neuronal excitability in memory consolidation has been

often observed in other memory systems(6, 36, 38, 39, 50, 51). Interestingly, our data

revealed that cerebellar learning elicited depression in PC-IE, whereas neurons in the

amygdala exhibited increased neuronal excitability associated with fear learning(52). Hence,

this difference indicates that PC-IE in the cerebellar memory circuit displays distinctive

features compared with other types of neuronal IE, as observed in amygdala neurons.

Although the intrinsic excitation of neurons in other memory circuits promotes a stable

process of memory consolidation, in the cerebellar memory circuit, potentiation of the

intrinsic state of PCs interrupts the formation of long-term memory. Hence, optogenetic

manipulation of exciting the PCs in this study may have disturbed the incomparable nature of

the cerebellum.

This study demonstrated the time-limited impact of intrinsic excitability on memory

consolidation, specifically within the critical window of 90 minutes following OKR. This

temporal constraint indicates that the initial phase of consolidation is a critical determinant

for transforming short-term memory into long-term memory. The acute slice recording

demonstrated that PC-IE at 0 min post-learning period was significantly decreased compared

with that of the naïve condition. Remarkably, by the 90 min post-learning mark, the level of

PC-IE had returned to that observed in the naïve condition. Considering that PC firing serves

as the primary output of the cortex and encodes various types of information including

temporal coding(53), we suggest that PC-IE, a key component of the PC output, is a plausible

mechanism for determining the critical time window. Synaptic plasticity events underlie the

early phase of hippocampal systems consolidation(54). Moreover, although transient

fluctuations of hippocampal excitability have been suggested to conduct memory

consolidation(55), this study represents the first detailed exploration of the critical time

window in cerebellum-dependent memory consolidation concerning the dynamics of

neuronal intrinsic excitability.

It is widely recognized that memory consolidation requires de novo protein synthesis, and

the temporal aspect of protein synthesis is of paramount importance for long-term memory

formation(56, 57). Long-term contextual fear memory formation in the hippocampus, for

instance, relies on transcription factors and associated signaling cascades, including the cyclic

AMP response element binding protein and the Ras/Erk pathway, which are essential for the

synthesis of proteins(58-60). Intriguingly, cerebellum-dependent motor learning has also

been associated with certain modifications in protein expression. However, the specific

proteins that undergo changes in expression vary depending on whether they are correlated

with short- or long-term memory(61). Our data emphasize the presence of a critical time

window for memory consolidation regulated by intrinsic excitability. Nonetheless, we

acknowledge that we have not yet provided further insights into the molecular mechanisms

underlying this phenomenon. As such, the involvement of protein synthesis in establishing

this critical time window and fluctuations in PC-IE remain unexplored aspects of this process.

Similar to the cerebellar cortex-to-nuclei network, the hippocampus employs systems

consolidation to establish remote memories in the neocortex. Previous studies have suggested

that hippocampal and neocortical synaptic plasticity share common forms, including

molecular substrates and conditioned pathways(62). Moreover, hippocampal modifications,

such as conformational changes in dendritic spines and suppression of neuronal output,

directly influence cortical representations(63-65), but unlike studies on the hippocampus,

there is limited understanding of the neural properties underlying cerebellar systems

consolidation. Therefore, we further investigated the post-circuitry of floccular PCs that are

primarily organized by FTNs in the MVN(13-17). PC excitability plays an important role in

the construction of cerebellum-dependent long-term memory, and the intrinsic excitability of

FTNs in the MVN was found to be crucial. Disruption of PC-IE was also accompanied by the

deficient plasticity of the intrinsic excitability of FTNs, indicating on the direct links of

different regional intrinsic plasticities within the single memory circuit. Our data reveal that

the intrinsic plasticity of PCs provides guidance to drive the induction of intrinsic plasticity in

FTNs, which aligns with previous expectations that PC activity can affect the IE of FTNs(37,

41). Additionally, we extended our analysis to another prominent group of neurons in the

MVN, nonFTNs(13-17). As nonFTNs inputs are made directly from vestibular organs and

lack physical synapse with floccular PC axons(13-17), aberrations in PC-IE did not exert an

influence on the intrinsic excitability of nonFTNs, as expected.

PCs, are a rare and exclusive type of neuron found solely within the cerebellar cortex, and

have long been associated with motor coordination and movement(66, 67). Overall, this study

demonstrated that the ability to fine-tune firing patterns in a time-dependent manner by

modulating intrinsic excitability plays an essential role in memory consolidation. This

mechanism is crucial for the precise refinement of motor memory, prompting us to explore

the fundamental question of why PCs employ such sophisticated approaches. The cerebellum,

a highly complex and organized structure comprising of several layers of specifically

localized neurons, is capable of processing massive amounts of input with remarkable

efficiency(68). Despite the abundance of sensory inputs, only one pathway of cerebellar

cortical output exists: the PCs, which exercise a far-reaching influence on numerous regions

in the cerebral cortex to modify both motor and non-motor behaviors(69-71). Although PCs

in different cerebellar lobules exhibit no morphological differences, compelling evidence

demonstrates that each lobule serves specific roles by encoding different sensory

modalities(72-74); hence, PCs exhibit different firing patterns with distinct membrane

properties according to their location(75). In addition to the results of previous studies, the

results of this study led us to speculate that PCs may have evolved to establish specific

patterns of spike activity to conduct a diverse range of cerebellum-dependent tasks, thereby

optimizing the biological system and allowing for efficient performance. In our case, PC-IE

was tuned through fluctuations in excitability at different time points to conduct

consolidation.

The construction of computational models of neuronal pathways has proven to be a useful

tool for analyzing and understanding memory circuits(15, 76, 77). Based on our findings,

future studies may apply these insights to computational modeling and simulations to predict

the patterns of PC activity during different behavior assessments across multiple time points.

This approach has the potential to provide deeper insights into the intricate interplay between

biological and computational mechanisms governing the precise tuning of PC firing through

intrinsic excitability. Moreover, extending the application of these computational models to

broader memory systems across diverse brain regions offers an opportunity to expand our

understanding of memory consolidation processes beyond the cerebellum and a unified

understanding of memory as a fundamental cognitive process.

In summary, our investigation provides valuable insights into the complex nature of

cerebellar learning circuitry. This highlights the crucial role of intrinsic excitability in the

process of memory consolidation, particularly within the critical 90-minute time window, and

emphasizes its significance in facilitating memory transfer from the cortex to the nuclei.

These findings have potential implications for understanding the pathophysiology of various

disorders associated with cerebellar dysfunction(78-81). Overall, these molecules hold

promise for the development of targeted therapeutic interventions that address and ameliorate

these conditions.

Materials and methods

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Virus	pAAV-Ef1a-DIO hChR2(C128S/D156A)-EYFP (AAV1)	Addgene	RRID: Addgene_35503	Plasmid #35503
Virus	pAAV-Ef1a-DIO EYFP (AAV1)	Addgene	RRID: Addgene_27056	Plasmid #27056
Chemical compound, drug	Zoletil	Virvac		
Chemical compound, drug	Rompun	Bayer		
Chemical compound, drug	Dexamethasone	Samyang Phamaceutical		
Chemical compound, drug	Meloxicam	Boehringer Ingelheim		
Software, algorithm	Intrinsic VIEW Analysis			
Software, algorithm	VOG Analyzer			
Software, algorithm	PatchMaster	HEKA	RRID: SCR_000034	
Software, algorithm	Igor Pro	WaveMetrics	RRID: SCR_000325	
Software, algorithm	GraphPad Prism	GraphPad	RRID: SCR_002798	

Animals and stereotaxic surgery for virus injection and optic cannula implantation

The experimental procedures were approved by the Seoul National University Institutional

Animal Care and Use Committee and performed the guidelines of the National Institutes of

Health. In this study, 6- to 10-week-old *PCP2-Cre* mice (Jackson Laboratory, ME, USA), a

driver line with high selectivity for *Cre* activity in PCs, were anesthetized with intraperitoneal

injections of Zoletil/Rompun mixture (30 mg/kg and 10 mg/kg, respectively) and then they

were placed on a stereotaxic frame (Narishige, Tokyo, Japan) for virus injection. AAV-Ef1a-

DIO hChR2(C128S/D156A)-EYFP (SSFO) and AAV-EF1A-DIO EYFP were manually

injected at a rate of 3 nl/ sec into the bilateral flocculi using *Picopump*. The glass pipette was

set at a 34° angle and the targeted coordinates for the virus injection were at 5.48 mm

dorsoventral (DV) and ±0.10 mm mediolateral (ML). After the mice were fully recovered

(approximately 1 week) from the first surgery of virus injection, the second operation of optic

cannula implantation (Ø 1.25 mm Ceramic Ferrule, L=3.0 mm) near the cerebellar flocculus

and head fixation were conducted. Unlike the fiber optic cannula, the head fixation

components were hand-made (personalized design and production). Prior to the surgery, the

mice were anesthetized using intraperitoneal injection of a mixture of Zoletil 50 (Virbac,

15mg/kg) and xylazine (Rompun, Bayer, 15mg/kg). Again, the mice were placed on placed

on a stereotaxic frame (Narishige, Tokyo, Japan), and optic cannulae were implanted on both

sides of the flocculi at ±2.80 mm ML.

Confirmation of viral expression.

After the completion of all experiments, sampling was performed using cardiac perfusion.

The brain was then extracted and coronal sections were made at 30 µm intervals. Images

were acquired and processed using a confocal microscope (Zeiss LSM 7 MP, Carl Zeiss,

Jena, Germany) and Zen software (Zeiss). The anatomical location of virus expression was

confirmed by comparison with Paxinos mouse brain atlas(82)

Slice preparation and electrophysiology recording

A acute brain slice preparation and electrophysiological experiments were performed as

previously described(34). Here, 5 to 9-week-old mice were anesthetized using isoflurane and

decapitated. To acquire the records of floccular PCs, 250 μm thick coronal slices of the

cerebellar flocculus were obtained from mice using a vibratome (VT1200S, Leica). However,

to acquire FTNs data, 200 μm thick coronal slices of the cerebellar MVN were obtained. The

ice-cold cutting solution contained 75 mM sucrose, 75 mM of NaCl, 2.5 mM of KCl, 7 mM

of $MgCl_2$, 0.5 mM of $CaCl_2$, 1.25 mM of NaH_2PO_4, 26 mM of $NaHCO_3$, and 25 mM of

glucose, with bubbled 95% O_2 and 5% CO_2. The slices were immediately transferred to the

artificial cerebrospinal fluid (ACSF) containing 125 Mm of NaCl, 2.5 Mm of KCl, 1 mM of

$MgCl_2$, 2 mM of $CaCl_2$, 1.25 mM of NaH_2PO_4, 26 mM of $NaHCO_3$, and 10 mM of glucose,

with bubbled 95 % O_2 and 5 % CO_2. Subsequently, they were recovered at 32 ℃ for 30 min

and at room temperature for 1 hr. Brain slices were placed in a submerged chamber perfused

with the ACSF for at least 10 min before recording. Whole-cell recordings were performed at

29.5–30 ℃. We used recording pipettes (3–4 MΩ) filled with 9 mM of KCl, 10 mM of KOH,

120 mM of K-gluconate, 3.48 mM of $MgCl_2$, 10 mM of HEPES, 4 mM of NaCl, 4 mM of

Na_2ATP, 0.4 mM of Na_3GTP, and 17.5 mM of sucrose (pH 7.25). Electrophysiological data

were acquired using an EPC9 patch-clamp amplifier (HEKA Elektronik) and PatchMaster

software (HEKA Elektronik) at a sampling frequency of 20 kHz, and the signals were filtered

at 2 kHz. All electrophysiological recordings of the PCs were acquired in the flocculus,

which is a lobe-like structure of the cerebellum. The spontaneous firing rate was analyzed

using Igor Pro (WaveMetrics) and normalized to the average firing rate during the baseline

period. Whole-cell recordings of FTNs were acquired from the MVN that exhibited EYFP-

expressing *boutons en passant* from floccular PCs. Inhibitory synaptic inputs were

completely blocked using 100 mM of picrotoxin (Sigma Millipore) during PC recordings,

and strychnine (1 mM) was added to block the glycinergic input in FTN recordings. Patch

pipettes (3–4 MΩ) were pulled from borosilicate glass and filled with an internal solution (pH

7.25) containing 9 mM of KCl, 10 mM of KOH, 120 mM of K-gluconate, 3.48 mM of

MgCl2, 10 mM of HEPES, 4 mM of NaCl, 4 mM of Na2ATP, 0.4 mM of Na3GTP, and 17.5

mM of sucrose for testing in vitro recordings, ex vivo PC excitability, and ex vivo FTN

recordings. To examine the excitability of cerebellar PCs, we provided a series of square-

wised current injections that ranged from 100–900 pA with increments of 100 pA for 500 ms.

To investigate the excitability of FTNs in the MVN, we also provided current injections from

-150 pA to 150 pA with increments of 50 pA for 1000 ms.

Optokinetic reflex behavior test

First, three sessions of acclimation were conducted. During each session, the mouse was

restrained using a custom-made restrainer for 30 min with and without light for habituation to

the recording environment. Calibration was performed as described by Stahl et al. (2002. This

allowed the conversion of the pupil into eye rotation. Next, prior to oculomotor training, three

basal oculomotor performances including OKR, VOR in the dark (dVOR) and VOR, in the

light (lVOR), were measured. During the recording of eye movements, it was necessary to

control the pupil dilation that was induced by the absence of light in the dark condition. Thus,

physostigmine salicylate solution (eserine; Sigma Millipore) was given to mice under brief

isoflurane anesthesia. The concentration of eserine solution was increased from 0.1%, 0.15%

and 0.2% based on the pupil size. Basal oculomotor performances were described in gains.

For the actual learning protocol, we adopted OKR, and it consisted of five training sessions

(10 min per session), six checkup points and 24 h of consolidation period. During the training

sessions, the mice were visually stimulated with a sinusoidally rotating drum (±5 degrees).

Completely trained mice were placed back into their home cage, which was stored in the dark

condition until the last gain checkup points. To assess the impact of optogenetic manipulation

during the acquisition period, the training consisted of three sessions, each lasting 15 minutes.

Optogenetic manipulation during optokinetic reflex

For in-vivo optogenetic manipulation, SSFO-expressing mice were stimulated by 473nm blue

laser and 593nm yellow laser for activation and deactivation, respectively. Each

optostimulation was briefly given for 2 seconds at 10.5 mW. During the behavior assessment

to determine the critical time window of memory consolidation, both SSFO and EYFP-

expressing mice were subjected to optogenetic manipulation during the OKR (optokinetic

reflex) training. The training was conducted twice, with each manipulation occurring at a

different time point. Following the first training session, a break period of 2 to 3 weeks was

implemented to reset the learning conditions before the commencement of the second

training.

Gain analysis

Gains acquired from basal oculomotor performances and OKR learning calculated by a ratio

of evoked eye movements to the movement of screen or turn-table as visual or vestibular

stimulus, respectively. A custom-built LabView (National Instrument) analysis tool was used

for all of the calculations.

Statistical analysis

Graph plotting and statistical analysis were performed using GraphPad Prism (GraphPad

Software Inc, CA, USA). The hypothesis was tested by using paired and unpaired t-tests

between sample pairs using GraphPad Prism. Statistical significance was set at $p < 0.05$.

Asterisks denoted in the graph indicate the statistical significance. * indicates $p < 0.05$, ** <

0.01, *** < 0.001, and **** < 0.0001. The test name and statistical values are presented in

each figure legend. Data are shown as mean ± SEM and statistical evaluations were

performed using, independent t test, one-way repeated-measured (RM) ANOVA and two-

way RM ANOVA with post hoc Tukey's test and Sidak's test and were performed.

Author Details

Jewoo Seo

Department of Physiology, Seoul National University College of Medicine, Seoul,

Republic of Korea

Department of Biomedical Sciences, Seoul National University College of Medicine,

Seoul, Republic of Korea

Contribution: Conceptualization, Resources, Data curation, Software, Formal analysis,

Validation, Investigation, Visualization, Methodology, Writing – original draft, review and

editing

Competing interests: No competing interests declared

Yong Gyu Kim

Department of Physiology, Seoul National University College of Medicine, Seoul,

Republic of Korea

Department of Biomedical Sciences, Seoul National University College of Medicine,

Seoul, Republic of Korea

Contribution: Conceptualization, Data curation, Software, Formal analysis, Investigation

Methodology

Competing interests: No competing interests declared

Yong-Seok Lee

Department of Physiology, Seoul National University College of Medicine, Seoul,

Republic of Korea

Department of Biomedical Sciences, Seoul National University College of Medicine,

Seoul, Republic of Korea

Memory Network Medical Research Center, Neuroscience Research Institute, Wide River

Institute of Immunology, Seoul National University College of Medicine, Seoul, Korea

Contribution: Conceptualization, Supervision, Writing – review and editing

For correspondence: yongseok7@snu.ac.kr

Competing interests: No competing interests declared

Sang Jeong Kim

Department of Physiology, Seoul National University College of Medicine, Seoul,

Republic of Korea

Department of Biomedical Sciences, Seoul National University College of Medicine,

Seoul, Republic of Korea

Memory Network Medical Research Center, Neuroscience Research Institute, Wide River

Institute of Immunology, Seoul National University College of Medicine, Seoul, Korea

Contribution: Conceptualization, Supervision, Funding acquisition, Writing – review and

editing

For correspondence: sangjkim@snu.ac.kr

Competing interests: No competing interests declared

578   **Acknowledgments**

579   This work was supported by National Research Foundation of Korea (NRF) grants funded by

580   the Korean Government (MSIT) (NRF- 2018R1A5A2025964 and 2022M3E5E8017970 to

581   S.J. Kim).

Conflict of Interest

The authors declare that they have no conflicts of interests.

 Figure legends

Figure 1. Optogenetic manipulation of PC-IE increases spontaneous firing

a. Schematic diagram of the virus injection for the optogenetic manipulation of floccular

Purkinje cells (PCs). Bottom left, the virus is specifically expressed in PCs of the

flocculus (FL), which is located below the paraflocculus (PFL) (scale bar indicates 500

µm). Bottom right, the granule cell layer (GCL), PC layer (PCL), and molecular layer

(ML) are shown (scale bar indicates 150 µm).

b. Scheme for optogenetic activation/deactivation of the SSFO virus expressed in PCs using

a blue laser (473 nm)/yellow laser (593 nm).

c. Representative traces of spontaneous PC firing. Traces of both the EYFP and SSFO

groups are shown under three different conditions: before optostimulation, after 473 nm

optostimulation, and after 593 nm optostimulation (vertical scale bar indicates 20 mV and

horizontal scale bar represents 100 ms).

d. Spontaneous firing rate of SSFO-expressing PCs (n=7) and EYFP-expressing PCs (n=7).

The rate of SSFO-expressing PCs significantly increased and decreased following 473 nm

and 593 nm optostimulation, respectively (SSFO, ** $p = 0.0017$ and *** $p = 0.0003$; one-

way ANOVA Tukey's multiple comparisons).

The error bars indicate ±SEM.

Figure 2. Optogenetic manipulation of PC-IE impairs memory consolidation

a. Top, experimental scheme for the optokinetic reflex (OKR). On the first day after

learning was complete, 473 nm optostimulation (2 s, 10.5mW) was administered to

mouse via an optic cannula at 0 min of post-learning period. The mouse was then placed

in a cage to rest in complete darkness for 24 hr. On the second day, 593 nm

optostimulation (2s, 10.5 mW) was administered to the mouse, and the remaining gain

was measured. Bottom, schematic diagram showing the location of optic cannula

implantation, which is the flocculus of the cerebellum, and the settings of

optostimulation.

b. Illustration of the OKR behavioral test. During the learning period, visual stimulation was

provided by the horizontal rotation of a screen at a constant speed (±5 deg at 0.5 Hz) and

the table was stationary.

c. Representative traces of the screen (black sinusoidal curve) and eye movements (grey

sinusoidal curve). Traces of both the EYFP and SSFO groups were presented at three

different periods: pre (before learning), post (after learning) and +24 hr (post 24 hr). The

vertical scale bar represents 10 degrees per second and horizontal scale bar represents 0.5

second.

d. Gain changes in the EYFP and SSFO groups following learning. The gain values in both

groups (EYFP, n = 14 mice; SSFO, n = 16 mice) significantly increased after learning

(EYFP, **** $p < 0.0001$; SSFO, **** $p < 0.0001$; repeated-measure [RM] one-way

analysis of variance [ANOVA], Tukey's multiple comparisons). After 24 hr of

consolidation period, the EYFP group showed the robust maintenance of long-term

memory ($p = 0.4122$; RM one-way ANOVA Tukey's multiple comparisons), whereas the

SSFO group shows impaired consolidation (**** $p < 0.0001$; RM one-way ANOVA,

Tukey's multiple comparisons).

e. Comparison of the consolidation percentages between the EYFP and SSFO groups. The

consolidation percentage in the EYFP group is significantly higher than that in the SSFO

group (EYFP, n = 14 mice; SSFO, n = 16 mice; Unpaired t test, **** $p < 0.0001$).

The error bars indicate ±SEM.

Figure 3. PC-IE contributes to the formation of long-term memory during the early

phase of the consolidation period

a. Two different experimental schemes for OKR. *Scheme #1*, 50 min of learning was

conducted on the first day; once the learning was complete, 473 nm optostimulation (2 s,

10.5mW) was administered to mouse via an optic cannula at 0 min post-learning period.

The mouse was then placed back in a cage to rest in complete darkness for 24 hr. On the

second day, 593 nm optostimulation (2s, 10.5 mW) was given to a mouse, and remaining

gain was measured. *Scheme #2*, same behavior protocol for OKR was conducted as

scheme #1, but 473 nm optostimulation (2 s, 10.5mW) was given at 90 min post-learning

period.

b. OKR learning curves from 0 to 50 min and +24 hr of the EYFP group under two different

conditions. Left, 473 nm optostimulation (2 s, 10.5mW) was administered to the mouse

via an optic cannula at 0 min post-learning period. Right, the same treatment was

administered at 90 min post-learning period.

c. Comparison of consolidation percentages of EYFP-expressing mice under two different

conditions. The consolidation percentages of EYFP-expressing mice optostimulated at 0

min post-learning period are comparable to those of EYFP-expressing mice

optostimulated at 90 min post-learning period (0 min, n = 8; 90 min, n = 8; Paired t test, p

= 0.3248)

d. OKR learning curves from 0 to 50 min, and +24 hr of the SSFO group under two

different conditions. Left, 473 nm optostimulation (2 s, 10.5mW) administered a mouse

via an optic cannula at 0 min post-learning period. Right, the same treatment was

administered at 90 min post-learning period.

e. Comparison of consolidation percentages of SSFO-expressing mice under two different

conditions. The consolidation percentages of SSFO-expressing mice optostimulated at 90

min post-learning period are significantly higher than those of SSFO-expressing mice

optostimulated at 0 min post-learning period (0 min, n = 10; 90 min, n = 10; paired t-test,

*** $p < 0.0010$).

f. OKR learning curves from 0 to 50 min and +24 hr. Left, learning curve of the EYFP

group: 473 nm optostimulation (2 s, 10.5mW) was administered to the mouse via an optic

cannula at 0 min post-learning period, and 593 nm optostimulation (2s, 10.5 mW) was

administered at 90 min post-learning period. Right, learning curve of the SSFO group:

473 nm optostimulation (2 s, 10.5mW) was administered to the mouse via an optic

cannula at 0 min post-learning period, and 593 nm optostimulation (2 s, 10.5mW) was

administered at 90 min post-learning period.

g. Comparison of consolidation percentages of between the EYFP (n = 6) and SSFO (n = 6)

groups. The consolidation percentages of the SSFO group was significantly lower than

that of the EYFP group (unpaired t-test, **** $p < 0.0010$).

The error bars indicate ±SEM.

Figure 4. Dynamics of PC-IE within the critical temporal window following motor

learning

a. Top, the ex-vivo experimental scheme for OKR and whole-cell patch clamp recording.

After 50 min of learning, slice preparation for whole-cell patch clamp recording was

conducted at 0 min post-learning and 90 min post-learning periods. Bottom left, learning-

induced increase in the OKR gain is presented (mouse, n = 5; paired t -test, *** p =

0.004). Bottom right, coronal slices are acquired, and electrophysiological recordings are

conducted in the flocculus.

b. Raw traces of PC firing under 600 pA current injection under three different conditions:

control, at 0 min post-learning period, and at 90 min post-learning period.

c. The mean firing rate (Hz) is significantly decreased in the learned group at 0 min post-

learning compared with that in the control (naïve) and 90 min post-learning groups

(control, n = 14; 0 min, n = 14; 90 min, n = 10; two-way analysis of variance [ANOVA]

with Tukey's multiple comparisons, **** p < 0.0001).

d. OKR learning elicits significant increase in the rheobase current at 0 min post-learning

period (One-way ANOVA multiple comparisons, 0 min vs. 90 min, * p = 0.0336; 0 min

vs. Control, * p = 0.0187).

e. No significant differences in the AP threshold.

f. No significant differences in the AP amplitude.

g. No significant differences in the AHP amplitude.

The error bars indicate ±SEM.

Figure 5. Disrupted plasticity of intrinsic excitability of flocculus-targeting neurons in the MVN

a. Top, an ex-vivo experimental scheme for OKR and whole-cell patch clamp recording. After 50 min of learning, slice preparation for whole-cell patch clamp recording was conducted at 24 hr post-learning period. Bottom (i), coronal slices are acquired, and electrophysiological signals of FTNs in the medial vestibular nucleus (MVN) were recorded (scale bars indicate 500 µm). (ii), The virus expression (EYFP/SSFO) in the flocculus is validated (scale bars indicate 500 µm).

b. Top, an image of floccular PC terminals expressed in the MVN (scale bar indicates 25 µm). Bottom, an image of FTN with *boutons en passant* (scale bar indicates 20 µm).

c. Raw traces representing the excitability of EYFP-expressing FTNs at 100 pA and 150 pA current injections under learned (n = 6) and naïve (n = 7) conditions. The vertical scale bar represents 10 mV and the horizontal scale bar represents 200 ms.

d. Comparison of intrinsic excitability of EYFP-expressing FTNs under naïve and learned conditions. Excitability at 100 pA and 150 pA current injections is significantly higher in the learned group (n = 6) than in the naïve group (n = 7) (Two-way analysis of variance [ANOVA] with Sidak's multiple comparisons: at 100 pA, ** p = 0.0065; at 150 pA, ** p = 0.0025).

e. Raw traces representing the excitability of SSFO-expressing FTNs at 100 pA and 150 pA current injections under learned (n = 7) and naïve (n = 6) conditions. The vertical scale bar represents 10 mV and the horizontal scale bar represents 200 ms.

f.   Comparison of intrinsic excitability of SSFO-expressing FTNs under naïve and learned

conditions. The excitability of the learned (n = 7) and naïve (n = 6) groups is (n = 6)

comparable. They exhibit no significant difference.

g.   Comparisons of mean firing rates between the FTNs of the EYFP and SSFO groups after

learning. EYFP-expressing FTNs (n = 6) exhibit significantly higher mean firing rates

than SSFO-expressing FTNs (n = 7) at 100 pA and 150 pA current injections (two-way

ANOVA with Sidak's multiple comparisons: at 100 pA, ** $p = 0.0020$; at 150 pA, *** p

$= 0.0002$).

The error bars indicate ±SEM.

754　**Figure 1.**

755
756
757

758　**Figure 2.**

Figure 3.

    Figure 4.

784 **Figure 5**

References

1.   Wiltgen BJ, Tanaka KZ. Systems consolidation and the content of memory.

Neurobiol Learn Mem. 2013;106:365-71.

2.   Kitamura T, Ogawa SK, Roy DS, Okuyama T, Morrissey MD, Smith LM, et al.

Engrams and circuits crucial for systems consolidation of a memory. Science.

2017;356(6333):73-8.

3.   Attwell PJE, Cooke SF, Yeo CH. Cerebellar function in consolidation of a motor

memory. Neuron. 2002;34(6):1011-20.

4.   Gais S, Albouy G, Boly M, Dang-Vu TT, Darsaud A, Desseilles M, et al. Sleep

transforms the cerebral trace of declarative memories. P Natl Acad Sci USA.

2007;104(47):18778-83.

5.   Robertson EM. Skill learning: Putting procedural consolidation in context. Curr Biol.

2004;14(24):R1061-R3.

6.   Boyden ES, Katoh A, Raymond JL. Cerebellum-dependent learning: The role of

multiple plasticity mechanisms. Annu Rev Neurosci. 2004;27:581-609.

7.   De Zeeuw CI, Lisberger SG, Raymond JL. Diversity and dynamism in the

cerebellum (Dec, 10.1038/s41593-020-00782-5, 2020). Nat Neurosci. 2021;24(3):450-.

8.   Kayakabe M, Kakizaki T, Kaneko R, Sasaki A, Nakazato Y, Shibasaki K, et al.

Motor dysfunction in cerebellar Purkinje cell-specific vesicular GABA transporter knockout

mice. Front Cell Neurosci. 2014;7.

9.   Inoshita T, Hirano T. Occurrence of long-term depression in the cerebellar flocculus

during adaptation of optokinetic response. Elife. 2018;7:e36209.

10.   Chang HHV, Cook AA, Watt AJ, Cullen KE. Loss of Flocculus Purkinje Cell Firing

Precision Leads to Impaired Gaze Stabilization in a Mouse Model of Spinocerebellar Ataxia

Type 6 (SCA6). Cells-Basel. 2022;11(17).

11.   Shutoh F, Ohki M, Kitazawa H, Itohara S, Nagao S. Memory trace of motor learning

shifts transsynaptically from cerebellar cortex to nuclei for consolidation. Neuroscience.

2006;139(2):767-77.

12.   Okamoto T, Shirao T, Shutoh F, Suzuki T, Nagao S. Post-training cerebellar cortical

activity plays an important role for consolidation of memory of cerebellum-dependent motor

learning. Neurosci Lett. 2011;504(1):53-6.

13.   Sekirnjak C, du Lac S. Physiological and anatomical properties of mouse medial

vestibular nucleus neurons projecting to the oculomotor nucleus. J Neurophysiol.

2006;95(5):3012-23.

14.   Sekirnjak C, Vissel B, Bollinger J, Faulstich M, du Lac S. Purkinje cell Synapses

target physiologically unique brainstem neurons. J Neurosci. 2003;23(15):6392-8.

15.   Yamazaki T, Nagao S, Lennon W, Tanaka S. Modeling memory consolidation

during posttraining periods in cerebellovestibular learning. P Natl Acad Sci USA.

2015;112(11):3541-6.

16.   Matsuno H, Kudoh M, Watakabe A, Yamamori T, Shigemoto R, Nagao S.

Distribution and Structure of Synapses on Medial Vestibular Nuclear Neurons Targeted by

Cerebellar Flocculus Purkinje Cells and Vestibular Nerve in Mice: Light and Electron

Microscopy Studies. Plos One. 2016;11(10): e0164037.

17.   Shin M, Moghadam SH, Sekirnjak C, Bagnall MW, Kolkman KE, Jacobs R, et al.

Multiple Types of Cerebellar Target Neurons and Their Circuitry in the Vestibulo-ocular

Reflex. J Neurosci. 2011;31(30):10776-86.

18.   Schweigart G, Mergner T, Evdokimidis I, Morand S, Becker W. Gaze stabilization

by optokinetic reflex (OKR) and vestibulo-ocular reflex (VOR) during active head rotation in

man. Vision Res. 1997;37(12):1643-52.

19.   Cahill H, Nathans J. The Optokinetic Reflex as a Tool for Quantitative Analyses of

Nervous System Function in Mice: Application to Genetic and Drug-Induced Variation. Plos

One. 2008;3(4): e2055.

20.   Lisberger SG. Internal Models of Eye Movement in the Floccular Complex of the

Monkey Cerebellum. Neuroscience. 2009;162(3):763-76.

21.   Medina JF, Nores WL, Ohyama T, Mauk MD. Mechanisms of cerebellar learning

suggested by eyelid conditioning. Curr Opin Neurobiol. 2000;10(6):717-24.

22.   Kandel ER, Dudai Y, Mayford MR. The Molecular and Systems Biology of

Memory. Cell. 2014;157(1):163-86.

23.   Ito M. Cerebellar Control of the Vestibulo-Ocular Reflex - around the Flocculus

Hypothesis. Annu Rev Neurosci. 1982;5:275-96.

24.   De Zeeuw CI, Hansel C, Bian F, Koekkoek SKE, van Alphen AM, Linden DJ, et al.

Expression of a protein kinase C inhibitor in Purkinje cells blocks cerebellar LTD and

adaptation of the vestibulo-ocular reflex. Neuron. 1998;20(3):495-508.

25.   Hansel C, de Jeu M, Belmeguenai A, Houtman SH, Buitendijk GHS, Andreev D, et

al. alpha CaMKII is essential for cerebellar LTD and motor learning. Neuron.

2006;51(6):835-43.

26.   Ito M. Error detection and representation in the olivo-cerebellar system. Front Neural

Circuit. 2013;7.

27.   Hirano T. Long-term depression and other synaptic plasticity in the cerebellum. P

Jpn Acad B-Phys. 2013;89(5):183-95.

28.   Chen LX, Cummings KA, Mau W, Zaki Y, Dong Z, Rabinowitz S, et al. The role of

intrinsic excitability in the evolution of memory: Significance in memory allocation,

consolidation, and updating. Neurobiol Learn Mem. 2020;173:107266.

29.   Kim SJ, Linden DJ. Ubiquitous plasticity and memory storage. Neuron.

2007;56(4):582-92.

30.   Belmeguenai A, Hosy E, Bengtsson F, Pedroarena CM, Piochon C, Teuling E, et al.

Intrinsic Plasticity Complements Long-Term Potentiation in Parallel Fiber Input Gain Control

in Cerebellar Purkinje Cells. J Neurosci. 2010;30(41):13630-43.

31.   Shim HG, Jang DC, Lee J, Chung G, Lee S, Kim YG, et al. Long-Term Depression

of Intrinsic Excitability Accompanied by Synaptic Depression in Cerebellar Purkinje Cells. J

Neurosci. 2017;37(23):5659-69.

32.   Yang Z, Santamaria F. Purkinje cell intrinsic excitability increases after synaptic

long term depression. J Neurophysiol. 2016;116(3):1208-17.

33.   Turrigiano G. Homeostatic Synaptic Plasticity: Local and Global Mechanisms for

Stabilizing Neuronal Function. Csh Perspect Biol. 2012;4(1):a005736.

34.   Kim YG, Kim SJ. Decreased intrinsic excitability of cerebellar Purkinje cells

following optokinetic learning in mice. Mol Brain. 2020;13(1):136.

35.   Titley HK, Watkins GV, Lin C, Weiss C, McCarthy M, Disterhoft JF, et al. Intrinsic

Excitability Increase in Cerebellar Purkinje Cells after Delay Eye-Blink Conditioning in

Mice. J Neurosci. 2020;40(10):2038-46.

36.   Ryu C, Jang DC, Jung D, Kim YG, Shim HG, Ryu HH, et al. STIM1 Regulates

Somatic Ca2+ Signals and Intrinsic Firing Properties of Cerebellar Purkinje Neurons. J

Neurosci. 2017;37(37):8876-94.

37.   Jang DC, Shim HG, Kim SJ. Intrinsic Plasticity of Cerebellar Purkinje Cells

Contributes to Motor Memory Consolidation. J Neurosci. 2020;40(21):4145-57.

38.   Kuo AG, Lee G, Mckay BM, Disterhoft JF. Enhanced neuronal excitability in rat

CA1 pyramidal neurons following trace eyeblink conditioning acquisition is not due to

alterations in I-M. Neurobiol Learn Mem. 2008;89(2):125-33.

39.   Mckay BM, Matthews EA, Oliveira FA, Disterhoft JF. Intrinsic Neuronal

Excitability Is Reversibly Altered by a Single Experience in Fear Conditioning. J

Neurophysiol. 2009;102(5):2763-70.

40.   Kassardjian CD, Tan YF, Chung JYJ, Heskin R, Peterson MJ, Broussard DM. The

site of a motor memory shifts with consolidation. J Neurosci. 2005;25(35):7979-85.

41.   McElvain LE, Bagnall MW, Sakatos A, du Lac S. Bidirectional Plasticity Gated by

Hyperpolarization Controls the Gain of Postsynaptic Firing Responses at Central Vestibular

Nerve Synapses. Neuron. 2010;68(4):763-75.

42.   Yizhar O, Fenno LE, Prigge M, Schneider F, Davidson TJ, O'Shea DJ, et al.

Neocortical excitation/inhibition balance in information processing and social dysfunction.

Nature. 2011;477(7363):171-8.

43.   Cooke SF, Attwell PJE, Yeo CH. Temporal properties of cerebellar-dependent

memory consolidation. J Neurosci. 2004;24(12):2934-41.

44.   Titley HK, Heskin-Sweezie R, Chung JYJ, Kassardjian CD, Razik F, Broussard DM.

Rapid consolidation of motor memory in the vestibuloocular reflex. J Neurophysiol.

2007;98(6):3809-12.

45.   Steinmetz AB, Freeman JH. Cannabinoid modulation of memory consolidation

within the cerebellum. Neurobiol Learn Mem. 2016;136:228-35.

46.   Bagnall MW, Stevens RJ, du Lac S. Transgenic mouse lines subdivide medial

vestibular nucleus neurons into discrete, neurochemically distinct populations. J Neurosci.

2007;27(9):2318-30.

47.   Gittis AH, du Lac S. Intrinsic and synaptic plasticity in the vestibular system. Curr

Opin Neurobiol. 2006;16(4):385-90.

48.   Lisman J, Cooper K, Sehgal M, Silva AJ. Memory formation depends on both

synapse-specific modifications of synaptic strength and cell-specific increases in excitability.

Nat Neurosci. 2018;21(3):309-14.

49.   Song CH, Ehlers VL, Moyer JR. Trace Fear Conditioning Differentially Modulates

Intrinsic Excitability of Medial Prefrontal Cortex-Basolateral Complex of Amygdala

Projection Neurons in Infralimbic and Prelimbic Cortices. J Neurosci. 2015;35(39):13511-24.

50.   Daoudal G, Debanne D. Long-term plasticity of intrinsic excitability: learning rules

and mechanisms. Learn Mem. 2003;10(6):456-65.

51.   Zhang W, Linden DJ. The other side of the engram: experience-driven changes in

neuronal intrinsic excitability. Nat Rev Neurosci. 2003;4(11):885-900.

52. Zhou Y, Won J, Karlsson MG, Zhou M, Rogerson T, Balaji J, et al. CREB regulates

excitability and the allocation of memory to subsets of neurons in the amygdala. Nat

Neurosci. 2009;12(11):1438-43.

53. De Zeeuw CI, Hoebeek FE, Bosman LWJ, Schonewille M, Witter L, Koekkoek SK.

Spatiotemporal firing patterns in the cerebellum. Nat Rev Neurosci. 2011;12(6):327-44.

54. Goto A, Bota A, Miya K, Wang JB, Tsukamoto S, Jiang XZ, et al. Stepwise synaptic

plasticity events drive the early phase of memory consolidation. Science.

2021;374(6569):857-+.

55. Thompson LT, Moyer JR, Disterhoft JF. Transient changes in excitability of rabbit

CA3 neurons with a time course appropriate to support memory consolidation. J

Neurophysiol. 1996;76(3):1836-49.

56. Shrestha P, Klann E. Spatiotemporally resolved protein synthesis as a molecular

framework for memory consolidation. Trends Neurosci. 2022;45(4):297-311.

57. Shafik AM, Allen EG, Jin P. Epitranscriptomic dynamics in brain development and

disease. Mol Psychiatr. 2022;27(9):3633-46.

58. Wood MA, Kaplan MP, Park A, Blanchard EJ, Oliveira AMM, Lombardi TL, et al.

Transgenic mice expressing a truncated form of CREB-binding protein (CBP) exhibit deficits

in hippocampal synaptic plasticity and memory storage. Learn Memory. 2005;12(2):111-9.

59. Athos J, Impey S, Pineda VV, Chen X, Storm DR. Hippocampal CRE-mediated gene

expression is required for contextual memory formation. Nat Neurosci. 2002;5(11):1119-20.

60. Tian XJ, Gotoh T, Tsuji K, Lo EH, Huang S, Feig LA. Developmentally regulated

role for Ras-GRFs in coupling NMDA glutamate receptors to Ras, Erk and CREB. Embo J.

2004;23(7):1567-75.

61.   Kim YG, Woo J, Park J, Kim S, Lee YS, Kim Y, et al. Quantitative Proteomics

Reveals Distinct Molecular Signatures of Different Cerebellum-Dependent Learning

Paradigms. J Proteome Res. 2020;19(5):2011-25.

62.   Kirkwood A, Dudek SM, Gold JT, Aizenman CD, Bear MF. Common Forms of

Synaptic Plasticity in the Hippocampus and Neocortex Invitro. Science.

1993;260(5113):1518-21.

63.   Vetere G, Restivo L, Cole CJ, Ross PJ, Ammassari-Teule M, Josselyn SA, et al.

Spine growth in the anterior cingulate cortex is necessary for the consolidation of contextual

fear memory. P Natl Acad Sci USA. 2011;108(20):8456-60.

64.   Restivo L, Vetere G, Bontempi B, Ammassari-Teule M. The Formation of Recent

and Remote Memory Is Associated with Time-Dependent Formation of Dendritic Spines in

the Hippocampus and Anterior Cingulate Cortex. J Neurosci. 2009;29(25):8206-14.

65.   Tanaka KZ, Pevzner A, Hamidi AB, Nakazawa Y, Graham J, Wiltgen BJ. Cortical

Representations Are Reinstated by the Hippocampus during Memory Retrieval. Neuron.

2014;84(2):347-54.

66.   Thach WT. A role for the cerebellum in learning movement coordination. Neurobiol

Learn Mem. 1998;70(1-2):177-88.

67.   Hoogland TM, De Gruijl JR, Witter L, Canto CB, De Zeeuw CI. Role of

Synchronous Activation of Cerebellar Purkinje Cell Ensembles in Multi-joint Movement

Control. Curr Biol. 2015;25(9):1157-65.

68.   Voogd J, Glickstein M. The anatomy of the cerebellum. Trends Cogn Sci.

1998;2(9):307-13.

69.   Li C, Saliba NB, Martin H, Losurdo NA, Kolahdouzan K, Siddiqui R, et al. Purkinje

cell dopaminergic inputs to astrocytes regulate cerebellar-dependent behavior. Nat Commun.

2023;14(1):1613.

70.   Carta I, Chen CH, Schott AL, Dorizan S, Khodakhah K. Cerebellar modulation of the

reward circuitry and social behavior. Science. 2019;363(6424).

71.   Low AYT, Goldstein N, Gaunt JR, Huang KP, Zainolabidin N, Yip AKK, et al.

Reverse-translational identification of a cerebellar satiation network. Nature.

2021;600(7888):269-73.

72.   Barmack NH, Baughman RW, Eckenstein FP. Cholinergic Innervation of the

Cerebellum of the Rat by Secondary Vestibular Afferents. Ann Ny Acad Sci. 1992;656:566-

79.

73.   Barmack NH, Baughman RW, Errico P, Shojaku H. Vestibular Primary Afferent

Projection to the Cerebellum of the Rabbit. J Comp Neurol. 1993;327(4):521-34.

74.   Kim YS, Shin JH, Hall FS, Linden DJ. Dopamine Signaling Is Required for

Depolarization-Induced Slow Current in Cerebellar Purkinje Cells. J Neurosci.

2009;29(26):8530-8.

75.   Kim CH, Oh SH, Lee JH, Chang SO, Kim J, Kim SJ. Lobule-specific membrane

excitability of cerebellar Purkinje cells. J Physiol-London. 2012;590(2):273-88.

76.   Yamazaki T, Nagao S. A Computational Mechanism for Unified Gain and Timing

Control in the Cerebellum. Plos One. 2012;7(3):e33319.

77.   Rizza MF, Locatelli F, Masoli S, Sanchez-Ponce D, Munoz A, Prestori F, et al.

Stellate cell computational modeling predicts signal filtering in the molecular layer circuit of

cerebellum. Sci Rep-Uk. 2021;11(1):3873.

78.   Wu T, Hallett M. The cerebellum in Parkinson's disease. Brain. 2013;136:696-709.

79.   Shakiba A. The Role of the Cerebellum in Neurobiology of Psychiatric Disorders.

Neurol Clin. 2014;32(4):1105-+.

80.   Moreno-Rius J. Opioid addiction and the cerebellum. Neurosci Biobehav R.

2019;107:238-51.

81.   Bruchhage MMK, Bucci MP, Becker EBE. Cerebellar involvement in autism and

ADHD. Handb Clin Neurol. 2018;155:61-72.

82.   Keith B. J. Franklin GP. Paxinos and Franklin's The mouse brain in stereotaxic

coordinates: Academic Press; 2013.

1025