A comprehensive guide through the fold and stability of cytochrome c with implications for disease

Muhammad Abrar Yousaf[a], Anna Savoia[b], Annalisa Pastore[c*]

[a] *Department of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Verona, Italy*
[b] *Department of Engineering for Innovation Medicine, University of Verona, Verona, Italy*
[c] *Department of Clinical Neuroscience, King's College London, Denmark Hill Campus, London, United Kingdom*

[*]Corresponding author:
Annalisa Pastore
Department of Clinical Neuroscience, King's College London, Denmark Hill Campus, London, United Kingdom
Email: annalisa.1.pastore@kcl.ac.uk

Keywords: Cytochrome c; thrombocytopenia 4; mutations; peroxidase activity; molecular dynamics simulation

Abstract

Cytochrome c (Cyt-c), encoded by *CYCS* gene, is crucial for electron transport, peroxidase activity, and apoptosis. Mutations in *CYCS* can cause thrombocytopenia 4 (THC4), a disorder with low platelet counts. We have, for instance, recently described six Italian families with five different heterozygous missense *CYCS* variants. These mutations likely enhance peroxidase and apoptotic activities, yet mechanisms causing reduced platelet production and increased apoptosis are unclear. This study investigates all clinically important Cyt-c variants using an integrated bioinformatics approach. Our findings reveal that all variants are at evolutionarily conserved sites, likely disrupting Cyt-c function and contributing to disease phenotypes. Specific variants likely affect phosphorylation (T20I, V21G, Y49H), ubiquitination (G42S, A52T, A52V), and glycosylation (H27Y, N32H, L33V, G42S). Cyt-c variants may influence the function of interacting genes and proteins involved in apoptosis and mitochondrial function. Molecular dynamics simulations reveal significant structural differences from the wild-type, with mutants showing reduced stability and increased flexibility, particularly in the omega loops. These changes weaken hydrogen bonds and increase distances between key residues, opening the heme active site. This open conformation likely enhances accessibility to small molecules such as H_2O_2, thereby promoting peroxidase activity, which may enhance apoptosis and likely impact megakaryopoiesis and platelet homeostasis in THC4.

1   Introduction

Cytochrome C (Cyt-c) is an essential small heme-binding protein [1] that functions as an electron carrier in the electron transport chain by transferring electrons from complex III to complex IV of the cytochrome oxidase complex [2–4]. Cyt-c is a small globular protein of 105 residues in humans that contains a single covalently bound heme group attached to Cys15 and Cys18 of a CXXCH motif with two thioether bonds. His19 and Met81 are axial ligands of the six-coordinated heme iron. Heme ligation remains the same both in the ferric and ferrous forms of Cyt-c. The protein fold comprises five α-helices connected by three so-called for historic reasons Ω-loop regions, namely distal (residues 72-86), central (residues 41-58) and proximal (residues 13-29 and 28-40) Ω-loops [5–7]. These are non-regular protein structural motifs that consist of loops of six or more amino acid residues (Fetrow, 1995). Being highly conserved, Cyt-c is attached to the inner membrane of the mitochondrion to fulfil multiple important functions [8] which include a role in the intrinsic apoptotic pathway where Cyt-c initiates cell apoptosis as an antioxidant agent [9–11]. Because of its essential cellular role, it is not a surprise that Cyt-c is associated to pathology: heterozygous mutations of the *CYCS* gene on chromosome 7p15 are associated to autosomal dominant non-syndromic thrombocytopenia 4 (THC4, OMIM: 612004) [12], a disease originally described in a New Zealander family and, more recently, found also in a few other families all over the world [12–15]. Thrombocytopenia are a heterogeneous group of diseases caused by mutations in as many as at least twenty different genes [16]. Some of these diseases are rare such as it is the case for THC4. Despite the important roles of Cyt-c, it is interesting to notice that THC4 patients present only a mild phenotype that usually consists of a decreased number of platelets in circulating blood [12,13] caused by ectopic platelet release from bone marrow [12]. Platelets from THC4 patients are otherwise normal in size and morphology, and no increased bleeding tendency is usually described. While the low severity of this condition could suggest that a better understanding of the mechanisms causing it may be uninteresting, the need of being able to distinguish this condition from other more serious pathologies is essential for a correct diagnosis which may avoid mistreatment of the patients. There have in fact been cases of THC4 that were incorrectly treated by steroids and/or splenectomy, without any significant benefit [17].

Various Cyt-c variants have been so far identified that cause THC4 [12–15,18–20]. Amongst them, G42S and Y49H have been extensively studied and the crystal structures of some variants have been solved [21,22]. More recently, we have reported new variants and characterized the six families of the carriers for their symptoms [17]. By a clinical characterization of the 22 affected individuals, the largest series of THC4 patients ever reported, we showed that this disorder is characterized by mild-to-moderate thrombocytopenia, normal platelet size, but no additional clinical phenotypes associated with reduced platelet count. We also identified a significant correlation between the spatial region of Cyt-c affected by mutations and the extent of thrombocytopenia, which suggests different degrees of impairment of Cyt-c functions caused by different pathogenetic variants.

In the present study, we present a comprehensive bioinformatics analysis of all currently known variants of Cyt-c, examining the structural features these variants induce in the Cyt-c fold through various computational tools and molecular dynamics simulations. We also explore how the structural roles of certain residues correlate with observed phenotypes, proposing new hypotheses regarding protein function. Hence, this study investigates the impact of different Cyt-c variants on protein folding and stability, with the aim of elucidating the molecular mechanisms underlying enhanced peroxidase activity, increased apoptosis, and their effects on platelet production in THC4. These findings could offer new insights into the role of Cyt-c in the context of disease.

2 Methodology

2.1 Data collection and multiple sequence alignment

This study describes the functional effects of fourteen single amino acid variants (T20I, V21G, H27Y, N32H, L33V, G42S, Y49H, A52T, A52V, R92G, A97D, Y98H, L99V, and T103I) in human Cyt-c using bioinformatics approaches. These variants have been reported in patients with or without clinical characterization or identified through sequencing analysis (Table 1). The three-dimensional structure of Cyt-c (PDB ID: 1j3s) was retrieved from the RCSB PDB database (https://www.rcsb.org/) [23]. Notably, the amino acid numbering in our study is offset by +1 from that in the PDB file, reflecting different sequence annotation conventions. Protein amino acid sequences of *Homo sapiens* and various other species were obtained from the UniProt database (https://www.uniprot.org/) [24]. A multiple sequence alignment was performed using the Clustal Omega tool (https://www.ebi.ac.uk/jdispatcher/msa/clustalo) [25] to facilitate the evolutionary conservation analysis of protein residues across different species.

Table 1. THC4-associated Cyt-c variants analyzed in the present study.

Protein	References (with molecular and/or clinical characterization)	References (without clinical characterization)
T20I		[26]
V21G		[26]
H27Y	[27]	
N32H		[26]
L33V		[26]
G42S	[12,17]	[20,26,28]
Y49H	[13,17]	
A52T	[17]	[26,29]
A52V	[14]	
R92G	[17]	[30]
A97D		ClinVar (ISTH-SSC Genomics)
Y98H		[31]
L99V	[17]	[20]

2.2   Prediction of functional effect and pathogenicity of variants

The functional impacts of Cyt-c variants were assessed using SIFT (https://sift.bii.a-star.edu.sg/) [32]. SIFT predicts the effects of amino acid substitutions on protein function based on their position and nature, with scores ≤ 0.05 indicating deleterious changes. For the association of variants with the disease, the SuSPect tool (http://www.sbg.bio.ic.ac.uk/~suspect/) [33] was utilized. This tool uses sequence, structural, and systems biology features to predict the phenotypic effects of genetic variants. Trained on seventy-seven distinct features, SuSPect distinguishes between neutral and disease-causing variants, assigning scores ranging from 0 (neutral) to 100 (disease), with a cutoff score of 50 to differentiate between the two types of variants.

2.3   Prediction of molecular effects of variants

The MutPred2 platform (http://mutpred.mutdb.org/) [34] predicted molecular effects for Cyt-c variants, utilizing structural, functional and evolutionary data. This tool integrates PSI-BLAST, SIFT and Pfam for profiling, with TMHMM, MARCOIL and DisProt algorithms for predicting transmembrane helices, coiled-coil and disordered regions, respectively. This integration aids in the assessment of structural damage.

2.4   Prediction of post-translational modification

Post-translational modifications (PTMs) are crucial for predicting protein function. Methylation sites were identified using GPS-MSP (https://msp.biocuckoo.org/) [35], whereas phosphorylation sites on Serine (S), Threonine (T), and Tyrosine (Y) were predicted using NetPhos 3.1 (https://services.healthtech.dtu.dk/services/NetPhos-3.1/) [36], with scores above 0.5 indicating likely phosphorylation. UbiNet 2.0 (https://awi.cuhk.edu.cn/~ubinet/index.php) [37] facilitated the estimation of ubiquitylation sites, and NetOglyc 4.0 (https://services.healthtech.dtu.dk/services/NetOGlyc-4.0/) [38] was employed for glycosylation site prediction, considering sites with scores over 0.5 as probable glycosylation candidates.

2.5   Analysis of gene and protein interactions

Cyt-c mutations may have effects on the associated genes and proteins. The *CYCS* gene interactions were analyzed using GeneMANIA (https://genemania.org/) [39] to predict and understand gene-gene relationships comprehensively. Further, the STRING database (version 12.0) (https://string-db.org/) [40] was utilized to predict protein-protein interactions for Cyt-c, assessing the impact of variants on specific interacting proteins. A high confidence threshold of 0.7 was applied in the analysis to minimize the likelihood of false positives and negatives, ensuring that only interactions with substantial supporting evidence were considered.

2.6   Prediction of protein stability

Predicting the stability of mutant proteins is crucial for understanding their activity, function, and regulation. The impact of variants on protein stability was assessed by calculating the difference

in Gibbs Free Energy ($\Delta\Delta G$, in kcal/mol) between the wild-type (ΔG_{WT}) and mutant (ΔG_{MT}) proteins, where:

$$\Delta\Delta G = \Delta G_{WT} - \Delta G_{MT}$$

Positive $\Delta\Delta G$ values indicate an increase in stability due to mutations, whereas negative values denote destabilization. For these predictions, we employed both three-dimensional structural and sequence-based methods. DynaMut2 (https://biosig.lab.uq.edu.au/dynamut2/) [41] analyzed the impact of variants on protein dynamics and stability through normal mode analysis of protein three-dimensional structure. MUpro (https://mupro.proteomics.ics.uci.edu/) [42], a sequence-based predictor, utilized a support vector machine model to assess the effects of mutations. These complementary approaches offer a comprehensive understanding of the stability of mutant proteins.

2.7   Molecular dynamics simulation

Molecular dynamics (MD) simulations were performed to investigate the impact of variants on protein stability, flexibility, and dynamics, utilizing the GROMACS 2019.2 platform [43] on the WebGRO for Macromolecular Simulations webserver. Simulations employed the GROMOS96 43a1 force field for fully solvated systems. To ensure electroneutrality, sodium chloride was dissolved in SPC water molecules. The system pressure and temperature were maintained at 1 bar and 300 K, respectively, using the Berendsen weak coupling algorithm. The LINCS algorithm was employed to maintain all bonds at their equilibrium distances while allowing torsion and bending internal motions during the simulation [44]. The system underwent 5000 steps of energy minimization via the steepest descent algorithm to reach equilibrium, followed by a 50 ns (50000 ps) production run MD simulation. The number of frames per simulation was 1000. To compare the significant differences in MD simulation parameters between wild-type (WT) and mutant proteins, the Kruskal-Wallis test (non-parametric test) followed by the Dunn test (p-value < 0.05) was applied. The results were presented as mean $\pm$ standard deviation.

3   Results

3.1   Cyt-c and evolutionary conserved residues

The variants analyzed in this study are present in the proximal and central loops and the C-terminus (Fig. 1A). These mutated residues were assessed for evolutionary conservation through multiple sequence alignment. Amino acid positions critical for protein structure and/or function tend to be more evolutionarily conserved. Consequently, mutations at these conserved sites are usually more deleterious [45–47]. The analysis revealed that residues T20, V21, H27, N32, L33, G42, Y49, A52, R92, A97, Y98, L99, and T103 are conserved across species from diverse taxonomic groups

(Fig. 1B). This conservation suggests that the described variants T20I, V21G, H27Y, N32H, L33V, G42S, Y49H, A52T/V, R92G, A97D, Y98H, L99V, and T103I are likely deleterious.

3.2 Prediction of the functional effect and pathogenicity of variants

Analysis of the variants and their predicted impacts on protein function and association with diseases was computed utilizing three computational prediction tools, SIFT, and SuSPect (Fig. 1C). SIFT scores (<0.05) indicate severe effects of all variants on protein function. SuSPect scores further support these findings, categorizing each variant with a 'Disease' label, where high scores (>50) suggest a potential association with the disease. This agreement across different predictive platforms underscores the robustness of the predictions.

3.3 Mapping variants on protein structure

The three-dimensional structure of Cyt-c (Fig. 2A) shows the presence of Ω-loops with the heme, thereby providing insights into the protein's function in electron transport and apoptosis within cells. All Cyt-c variants were mapped onto the protein three-dimensional structure (Fig. 2B, C, D). The mutant structures of Cyt-c were generated using the SWISS-MODEL webserver (https://swissmodel.expasy.org/) [48] using the WT structure (PDB ID: 1j3s) as a template. Ramachandran plots were generated using PROCHECK tool [49] to evaluate the conformational integrity of both WT and mutant structures. These plots assess the phi and psi dihedral angles of the amino acid residues, key indicators of structural feasibility. In all mutant structures, the Ramachandran plot did not reveal any significant differences in the steric hindrance at the mutated residues compared to the WT (Supplementary data, Fig. S1), except for G42S (Fig. 2E). How the clinically important mutations alter the affected amino acids is shown in Supplementary Data (Fig. S2).

Buried residues are crucial for the structural stability of proteins because they contribute to the hydrophobic core and are like the pillars of the three-dimensional structure. Mutations that destabilize proteins can induce unfolding or misfolding, potentially affecting the functionality of Cyt-c. All examined amino acid residues examined in this study are buried within the protein structure. The T20I variant in the Cyt-c coil region (Fig. 2B), substituting a buried hydrophilic threonine with a larger hydrophobic isoleucine, likely compromises protein stability and disrupts the T20-N32 H-bonds [50]. This is due to the absence of the hydroxyl group in isoleucine, impeding the formation of the H-bond. The V21G variant in a coil region (Fig. 2B) introduces a cavity into the hydrophobic core by replacing a medium-sized valine with a smaller, more flexible glycine, potentially affecting protein stability (Fig. 2B). The critical role of H27 in a coil region (Fig. 2B) for maintaining protein stability is highlighted by the formation of a key H-bond with P45, which ensures the close association of the 20s and 40s Ω-loops [51]. Substituting histidine with the bulkier, uncharged and hydrophobic tyrosine in the H27Y variant may induce structural alterations that significantly reduce protein stability and disrupt crucial interactions. The replacement of N32 (Fig. 2B) with a bulkier histidine in the N32H variant could impair the T20-N32 H-bonding [50] and is likely to affect the protein structural integrity, leading to instability.

Additionally, the L33V variant at the turn (Fig. 2B) might create a cavity in the hydrophobic core due to valine smaller size.

G42, located in a protein loop, exhibits positive phi and psi values in the Ramachandran plot (Fig. 2E). Glycines, with their single hydrogen atom side chain, are symmetric and have a wider range of acceptable phi/psi backbone dihedral angles than other amino acids, allowing glycines to occupy also typically disallowed region of the Ramachandran plot. Replacing glycine with the bulkier serine could thus lead to steric clashes with adjacent residues due to glycine unique conformational flexibility, significantly altering the loop conformation [52] (Fig. 2E). Further, the G42S variant (in a coil region, Fig. 2C), within the hydrophobic core, introduces a polar and hydrophilic serine, potentially disrupting hydrophobic interactions, destabilizing the core, and affecting the structural integrity and protein function. Y49, located in a buried hydrophobic loop, is a large residue with a bulky phenyl ring and a hydroxyl group (Fig. 2C). Replacing Y49 with the medium-sized, imidazole-ring-containing H49 could destabilize the hydrophobic core by creating a cavity. The hydrophilic and basic nature of H49 might disrupt hydrophobic interactions and introduce pH-dependent changes, potentially impacting the protein structure and function. The A52T variant, replacing the nonpolar alanine located in a helix within the hydrophobic core (Fig. 2C) with the polar and hydrophilic threonine, could disturb hydrophobic interactions, affecting structure. Similarly, substitution to a larger valine in the A52V variant might affect the structure. R92, a large, polar and hydrophilic residue located in the core helix near the cytochrome c active site (Fig. 2C), is crucial for interactions with the heme and other buried hydrophilic residues. Mutating R92 to the much smaller glycine could disrupt these interactions and create a cavity, compromising the protein function and structure.

The A97D variant in the helix (Fig. 2D) may disrupt the protein hydrophobic interactions due to the introduction of the hydrophilic aspartate at position 97, replacing hydrophobic alanine. The Y98H (Fig. 2D) variant could compromise the protein structural integrity by creating a cavity and disrupting hydrophobic interactions within the core helix, as it involves replacing the large hydrophobic tyrosine with a medium-sized hydrophilic histidine. Similarly, the L99V variant, located in a helix within the hydrophobic core (Fig. 2D), is likely to form a cavity due to the substitution of leucine with the smaller valine. The T103I variant in a coil region (Fig. 2D) replaces the small, polar, hydrophilic threonine with a larger, nonpolar, hydrophobic isoleucine, potentially affecting local packing and hydrophilic interactions.

Collectively, these variants could significantly influence the protein by altering its structure and functionality.

3.4   Molecular effects of the mutations

We assessed the molecular effects of the mutations of the Cyt-c variants, which could lead to pathogenicity, by the MutPred2 software. Results demonstrate that these variants often result in modified protein interfaces, alterations in metal binding and stability and gains or losses of catalytic and allosteric sites. These alterations may be associated with shifts in post-translational

modifications, such as acetylation, methylation, phosphorylation and ubiquitylation, indicating significant effects on enzymatic activity and protein regulation (Supplementary data, Table S1).

3.5   Predicting the effects of PTMs

PTMs refer to the chemical modifications that proteins undergo after translation, including methylation, phosphorylation, ubiquitylation, glycosylation, acetylation, and several others. PTMs play a pivotal role in the regulation of protein structures and functions, proving essential across various pathways, including cell signaling and protein-protein interactions, as well as numerous biological processes [53,54]. Mutations at or near the site of these modifications could impact their function, stability, localization, and interaction with other molecules. Our analysis utilized a combination of bioinformatics tools to evaluate the potential effects of Cyt-c mutations on its PTMs, which are crucial for its functionality and regulatory mechanisms. The GPS-MSP tool suggests that native Cyt-c lacks methylation sites, indicating a minimal role of methylation in its regulation. NetPhos 3.1 predicted phosphorylation sites at threonine 20 (T20), tyrosine 47 (Y47), threonine 64 (T64), and threonine 79 (T79). Variants such as T20I, V21G, and Y49H, located at or near these sites, could potentially modify the phosphorylation pattern, thereby affecting the protein activity or its interactions with other cellular molecules. Particularly, the T20I variant would abolish the possibility of phosphorylation at this site, potentially influencing the protein function or stability. Ubiquitylation, a modification marking proteins for degradation, was predicted at lysine residues K40, K54, K80 and K87. Variants near these sites (G42S, A52T, A52V) may affect the ubiquitylation process, thus altering the protein degradation rate or its role in cellular signaling. Glycosylation, which influences protein folding, stability, and interactions, was predicted at threonine residues T29 and T41 by NetOGlyc4.0. Variants proximal to these residues (H27Y, N32H, L33V, G42S) could change the glycosylation pattern, potentially affecting the protein structure and its functional capabilities.

3.6   Effects of variants on interactions

Gene network analysis using the GeneMANIA tool revealed that the *CYCS* gene, encoding Cyt-c, engages with twenty distinct genes (Fig. 3A). Mutations in the gene product of *CYCS* could potentially disrupt these interactions. Specifically, *CYCS* was found to co-express with CYC1, VDAC1, VDAC3, and EPS15L1, maintaining similar expression levels across various conditions. It shares a domain only with CYC1. Additionally, *CYCS* demonstrated co-localization with BCL2L1, VDAC1, VDAC2, and CASP9, suggesting that these genes are expressed together in the same cellular locations or tissues. All genes within the same pathway were associated with apoptosis. *CYCS* displayed genetic interactions with BAX, BAK1, and DAP3; all these genes are key players in apoptosis. *CYCS* encodes Cyt-c, which, upon mitochondrial release, initiates apoptosome formation with APAF1 and CASP9, subsequently activating caspases such as CASP3 to drive apoptosis. Genetic interactions between *CYCS* and BAX, BAK1, and DAP3 demonstrate that alterations in *CYCS* function or expression can modulate the apoptotic effects mediated by these genes, underscoring the interconnected nature of the apoptotic pathway and the pivotal role of *CYCS* in regulating apoptosis.

On the protein level, analyses of protein-protein interactions elucidated the functional relationships among cellular proteins. The STRING database revealed that Cyt-c interacts with ten other proteins, suggesting that mutations in Cyt-c might disrupt their functions (Fig. 3B). Notably, Cyt-c was found to interact with QCRFS1, CYC1, CASP9, APAF1, MT-CYB, BCL2L1, CASP3, BCL2, CYB5B, and CYB5A, all with high confidence scores (Supplementary data, Table S2). These proteins are involved in the electron transport chain and apoptotic pathway. This analysis highlights the complex network of protein-protein interactions involving Cyt-c, reflecting its critical role in cellular processes.

3.7 Gibbs free energy-based protein stability analysis

We assessed the impacts of Cyt-c variants on protein stability by the DynaMut2 and MUpro software, using $\Delta\Delta G$ values which quantify the difference between the energy of the unfolded and folded states of a protein. A positive $\Delta\Delta G$ indicates that a mutant protein is thermodynamically more stable (requiring more energy to unfold) than the WT, whereas a negative $\Delta\Delta G$ suggests it is less stable (requiring less energy to unfold). V21G, H27Y, N32H, L33V, G42S, Y49H, A52T, A52V, R92G, A97D, Y98H, L99V and T103I variants were identified as destabilizing (negative $\Delta\Delta G$) by both prediction tools (Table 2). Notably, variants carrying the V21G, Y49H, R92G, Y98H and L99V mutations showed the most pronounced destabilizing effects (larger negative $\Delta\Delta G$), underscoring their significant impact on protein function.

Table 2. Gibbs free energy ($\Delta\Delta G$) values for Cyt-c variants computed through DynaMut2 and MUpro servers. Negative $\Delta\Delta G$ indicate destabilizing variants, while positive values suggest the variant as stabilizing.

Mutation	DynaMut2 ($\Delta\Delta G$, kcal/mol)	MUpro ($\Delta\Delta G$, kcal/mol)
T20I	0.51	–0.30
V21G	–2.87	–2.12
H27Y	1.12	–0.29
N32H	–0.81	–1.25
L33V	–2.2	–0.62
G42S	–0.68	–0.53
Y49H	–1.16	–1.22
A52T	–1.36	–0.86
A52V	–0.76	–0.58
R92G	–1.61	–1.10
A97D	–0.69	–0.77
Y98H	–1.76	–2.04
L99V	–1.48	–1.48
T103I	–0.17	–0.24

3.8 Dynamics of the wildtype and mutant proteins in aqueous solution

Finally, we performed MD simulations to evaluate the conformational dynamics of Cyt-c. We observed significant alterations in the structured regions of the protein, notably affecting the α-helical and loop regions. The analysis was conducted on WT Cyt-c and on all the mutant structures over 50 nanoseconds (ns), as structural and confirmational modifications in the protein have been reported in the initial period during protein unfolding [55,56]. That is why previous Cyt-c studies conducted MD simulation analyses till or less than 30 ns [56–59].

Over the MD simulation period, WT and mutant variants exhibited distinct alterations in secondary structure, indicating structural rearrangements. Snapshots of protein structure were taken every 12.5 ns to visualize structural changes (Supplementary data, Fig. S3, S4, S5). The structure of WT transitioned from 30.6% alpha helices, 3.3% 3-10 helices, 26.3% extended strands (part of a beta sheet), 0.6% isolated beta-bridges, 10.2% turns, 8.6% bends, and 20.4% coils to 38.8% alpha helices, 5.8% 3-10 helices, 3.9% isolated beta-bridges, 5.8% turns, 20.4% bends, and 25.2% coils. This change was characterized by increased coil and helix structures within the Ω-loops while eliminating extended strands, leading to a more compact and stable protein structure. During the simulation, all the variants, except L33V and Y98H, show a decrease in alpha-helical content and an increase in turns or bends conformation. Interestingly, variants Y49H, A97D, and L99V lead to the temporary development of new beta-sheets during the simulations. Helices with stable H-bonds provide stability, and reduced helical conformation indicates the instability of protein structure. Conversely, increased turns or bends conformation can interfere with the proper folding of the protein, leading to misfolded proteins that are often less stable.

3.9 Effect of the mutations on protein conformational stability

The root mean square deviation (RMSD) measures how much the structure deviates from a reference conformation, usually the starting structure, and is thus used as an indicator of stability. Higher RMSD values indicate reduced stability and lower values suggest greater stability [60]. WT Cyt-c exhibits the lowest RMSD, indicating stable conformation close to its initial structure (Fig. 4A). In contrast, T20I, V21G, H27Y, N32H and L33V mutant forms depict higher RMSD values, suggesting increased conformational fluctuations and reduced stability relative to the WT. Statistical analysis confirms that all variants significantly increase RMSD compared to WT, affecting Cyt-c stability (Fig. 4B). WT Cyt-c stability could be due to

increased helix and coil structures and decreased extended strands. Mutant (T20I, V21G, H27Y, N32H) destabilization could be associated with conformational changes such as decreased alpha-helix. Furthermore, T20I displays a significant increase in turns; V21G increases 3_10-helices, turns, and bends; H27Y shows higher turns and bends; and N32H enhances 3_10-helices and bends. L33V lower RMSD values relative to other mutants could be because the L33V mutant structure displays alpha-helical content relatively stable, but there is an increase in 3_10-helices and extended strands, indicating minor structural adjustments.

G42S, Y49H, A52T, A52V and R92G mutants (Fig. 4C, D) depicted significantly higher RMSD values than WT, suggesting lower stability, with G42S and R92G being particularly destabilizing. This instability is linked to structural changes: G42S displays a decrease in alpha-helix and an increase in turn content; Y49H shows decreased alpha-helix and increased 3_10-helices, turns, and bends; A52T exhibits decreased alpha-helix and increased turns and 3_10-helices; and A52V shows an increase in alpha-helix and a notable appearance of extended strands.

A97D, Y98H and T103I also showed significantly higher RMSD than WT (Fig. 4E, F), indicating destabilization, especially T103I. A97D increases bends and turns with reduced alpha-helical content; Y98H significantly increases in turns and bends; and T103I increases 3_10-helices and turns while reducing alpha-helical content, suggesting significant structural alterations. L99V increases turns with a decrease in alpha-helical content, indicating minor flexibility as displayed by non-significant differences with the WT ($p > 0.05$) (Fig. 4F).

3.10 Effect of the mutations on protein compactness and folding

The radius of gyration (Rg) measures protein compactness and structural stability, with lower Rg values indicating tighter packing and higher values suggesting more extended or unfolded structures. The WT protein exhibits a stable Rg, indicating consistent compactness and stability. In contrast, mutant proteins display varying Rg profiles, suggesting changes in compactness and stability due to variants (Fig. 5). Significantly higher Rg values for G42S, A52V, and R92G mutants indicate less compact structures (Fig. 5C, D). A52T shows a small decrease in Rg, while Y49H has comparable compactness to WT. Additionally, A97D, Y98H, L99V, and T103I exhibit significantly increased Rg compared to WT, suggesting less compact structures (Fig. 5E, F).

3.11 Effect of the mutations on protein flexibility

The root mean square fluctuation (RMSF) quantifies the flexibility of individual amino acid residues within a protein structure throughout a simulation. Higher RMSF values indicate increased residue movements, often linked to protein destabilization. Our results demonstrate that all mutated proteins show significantly higher flexibility than the WT protein (Fig. 6A, B, C). Analysis reveals the impact of variants across the Ω-loops: proximal (residues 13-40), central (residues 41-58), and distal (residues 72-86). Specifically, T20I and V21G increase flexibility in the proximal and central loops, respectively, while H27Y enhances flexibility across all Ω-loops, indicating a potential impact on protein dynamics. N32H and L33V significantly elevate RMSF in the proximal and central loops, with L33V also affecting the distal loop (Fig. 6D). The G42S variant displays heightened flexibility in all loops. Y49H shows increased RMSF in proximal and central regions, whereas A52T has a negligible effect on the proximal loop but reduces RMSF in central and distal loops. The A52V variant increases flexibility in proximal and central regions, with R92G also showing significant fluctuations across all loops and the N-terminus (Fig. 6E). The A97D, Y98H, and T103I variants significantly raise RMSF values in all loop regions, with T103I exhibiting the most pronounced fluctuations. Conversely, L99V increases flexibility solely in the proximal loop region (Fig. 6F).

3.12 H-bond network analysis suggests increased peroxidase activity

A hydrogen bond (H-bond) analysis compares the bonding patterns between the WT and mutant variants. The results reveal that the WT protein exhibits a higher number of H-bonds compared to most mutants, including T20I, V21G, N32H, G42S, A52T, R92G, A97D, Y98H, L99V, and T103I (Fig. 7A, B, C). This finding suggests that these variants potentially disrupt or alter the H-bonding network within the protein structure. However, the A52V mutant shows a less significant difference in the number of H-bonds compared to WT. Interestingly, mutants such as H27Y, L33V, and Y49H display a similar number of H-bonds as the WT protein, indicating that these variants may not significantly impact the bonding pattern (Fig. 7D, E, F).

In Cyt-c, Tyr68 forms a H-bond network with the axial ligand Met81, and conserved residues Asn53 and Thr79. Tyr68 is also involved in H-bonding interactions with the central and distal Ω-loops. This network involving Tyr68 is crucial for the conformational switch, orienting the protein towards the apoptotic pathway [61]. Consequently, distances between these residues were calculated (Fig. 8, Table 3).

The results indicate a loss of H-bonds in the T20I, V21G, N32H, G42S, Y49H, A52T, A52V, Y98H, L99V, and T103I mutants of Cyt-c with increased distances between H-bonded residues. This disruption of specific interactions may destabilize the active heme configuration by altering the positions of Met81, Asn53, and Thr79. Such structural changes could increase the accessibility of the heme catalytic site to small molecules like H_2O_2, thereby enhancing peroxidase activity and facilitating the initiation of the apoptosis pathway. In contrast, the H27Y, L33V, R92G, and A97D Cyt-c mutants exhibit considerably stronger H-bonds involving these critical residues compared to WT, suggesting an increased catalytic activity of the heme.

Table 3. Distances between H-bonded residues between Tyr68-Asn53, Tyr68-Thr79 and Tyr68-Met81 of WT and mutant Cyt-c after 50 ns MD simulation. Note that amino acid numbering differs by +1 from that of the PDB file (1j3s).

Protein	Tyr68 and Asn53 (nm)	Tyr68 and Thr79 (nm)	Tyr68 and Met81 (nm)
WT	1.31	0.96	0.88
T20I	1.30	1.04	1.00
V21G	1.42	1.47	1.41
H27Y	1.16	0.91	0.70
N32H	1.21	1.47	0.90
L33V	1.10	1.02	0.81
G42S	1.28	1.04	1.01
Y49H	1.53	1.19	1.16
A52T	1.25	0.98	1.16
A52V	1.20	1.03	0.97
R92G	1.14	1.23	0.87
A97D	1.04	0.93	0.61

Y98H	1.53	1.19	1.05
L99V	1.27	0.98	0.99
T103I	1.20	1.10	0.97

3.13 Analysis of active site residues for electron transfer efficiency and apoptosis

Optimal distances between residues are crucial for electron transfer around the heme, which is essential for the electron flow [62]. Variations in these distances impact electron transfer in WT and mutant Cyt-c proteins (Table 4). In WT Cyt-c, the heme group forms covalent bonds with Cys15 and Cys18, with axial ligands His19 and Met81. The variants T20I, V21G, N32H, L33V, G42S, Y49H, A52T, A52V, Y98H, L99V, and T103I increase distances between electron-transfer residues relative to WT, while H27Y and A97D variants decrease these distances. The R92G variant does not significantly alter these distances. The increased distances in certain mutants lead to an opening of the active site conformation, likely enhancing accessibility to small molecules like H_2O_2 and promoting peroxidase activity [62]. Specifically, T20I, V21G, N32H, L33V, Y49H, A52T, A52V, Y98H, L99V, and T103I variants induce significant structural changes, resulting in an open active conformation throughout the simulation. Conversely, minor distance changes in G42S and R92G mutants suggest that their structural integrity and heme active conformation closely resemble WT Cyt-c, enabling these mutants to retain electron transfer competency within the respiratory chain similar to WT.

Table 4. Distance between key amino acid residues of the active site environment of WT and mutant Cyt-c after 50 ns MD simulation. Note that amino acid numbering differs by +1 from that of the PDB file (1j3s).

Protein	Tyr68 and Cys15 (nm)	Tyr68 and Cys18 (nm)	Tyr68 and His19 (nm)	Tyr68 and Phe83 (nm)	Met81 and Phe83 (nm)
WT	1.29	1.22	1.04	0.98	0.66
T20I	1.71	1.62	1.46	0.63	0.61
V21G	1.05	1.27	1.1	0.99	0.7
H27Y	1.11	1.00	1.05	0.91	0.57
N32H	1.28	1.42	1.19	0.89	0.68
L33V	1.52	1.46	1.34	1.13	0.69
G42S	1.09	1.57	1.39	0.72	0.62
Y49H	1.36	1.49	1.29	1.03	0.56
A52T	1.76	1.49	1.29	1.03	0.56
A52V	1.41	1.61	1.54	0.85	0.70
R92G	1.19	1.29	1.11	0.86	0.57
A97D	1.07	1.06	1.16	0.93	0.72
Y98H	1.13	1.65	1.70	0.75	0.66
L99V	1.41	1.44	1.16	0.94	0.70
T103I	1.67	1.68	1.32	0.86	0.65

Takano et al. [63] reported that the electron transfer pathway from mitochondrial Cyt-c reductase to oxidized Cyt-c (Fe^{3+}) differs from the reduced form (Fe^{2+}) to cytochrome c oxidase. X-ray crystallography of WT Cyt-c revealed parallel orientations of Tyr75 with Trp60 and Tyr68 with Heme105. In the Cyt-c ferric state, electron transfer is proposed to begin at Tyr75, then to Trp60, with a subsequent transfer from Tyr68 to the heme, potentially via Met81 sulfur, to preserve electron neutrality. This triggers conformational adjustments, aligning Trp60 with Tyr68 to enable electron transfer between these electron-rich and electron-deficient sites. Therefore, the distances between Trp60-Tyr75 and Tyr68-Met81 are crucial for the electron transfer process. Post-simulation analysis shows that mutants T20I, V21G, H27Y, N32H, G42S, Y49H, A52T, A52V, A97D, Y98H, L99V, and T103I significantly increase the Trp60-Tyr75 distance, while T20I, V21G, G42S, Y49H, A52T, A52V, Y98H, L99V, and T103I also show an increase in the Tyr68-Met81 distance compared to WT (Table 5). These changes suggest a potential impact on electron transfer efficiency and apoptotic activity.

Table 5. Distance between Trp60 to Tyr75 and Tyr68 to Met81 residues of WT and mutant Cyt-c. Note that amino acid numbering differs by +1 from that of the PDB file (1j3s).

Protein	Trp60-Tyr75 (nm)	Tyr68-Met81 (nm)
WT	0.93	0.88
T20I	1.04	1.00
V21G	1.06	1.41
H27Y	1.09	0.70
N32H	1.03	0.90
L33V	0.80	0.81
G42S	1.05	1.01
Y49H	1.01	1.16
A52T	1.00	1.16
A52V	0.99	0.97
R92G	0.90	0.87
A97D	1.16	0.61
Y98H	1.20	1.05
L99V	1.00	0.99
T103I	1.05	0.97

4   Discussion

The interplay between increased peroxidase activity and apoptosis in Cyt-c mutations presents a complex mechanism contributing to the pathogenesis of THC4. Cyt-c, besides the mitochondrial respiratory chain, is crucial in the intrinsic apoptosis pathway [7,64]. For apoptosis, Cyt-c moves across the outer mitochondrial membrane into the cytosol to activate the caspase cascade and initiate the apoptotic activity. This process could be initiated through several stimuli, including

oxidative stress. It is believed that permeabilization of the outer mitochondrial membrane for Cyt-c release to the cytosol is induced by the peroxidase activity of Cyt-c in complex with cardiolipin, a phospholipid found in the mitochondrial inner membrane [65,66].

Post-translational modifications and mutations in Cyt-c can alter its structure and apoptotic function [67,68]. In this study, we propose that Cyt-c variants cause structural destabilization and increased flexibility, specifically through the loss of the intra-protein H-bond network and increased distance between important residues in the heme catalytic site. These changes increase H_2O_2 access, causing oxidative stress and enhancing peroxidase activity, which stimulates apoptosis. This increase in apoptotic activity, particularly within megakaryocytes, the progenitor cells for platelets, could lead to a reduction in platelet count, manifesting as THC4 [22,27]. The precise molecular mechanisms linking these enhanced activities to the clinical phenotype of THC4 remain an area of active investigation, with current hypotheses suggesting a disruption in the balance between platelet production and apoptosis.

Our study shows that all variants, except L99V, display higher RMSD values during MD simulation, indicating protein destabilization. Higher RMSD is associated with increased apoptosis [22]. Mutant structures show increased flexibility in the different Ω-loop structures, which could alter the heme iron coordination, particularly by disrupting the native Met81-Fe or Arg39-Fe bond promoting H_2O_2 access to the heme, thus increasing peroxidase activity [22,68–70]. Increased Ω-loop flexibility also impacts Cyt-c interaction with cardiolipin facilitating Cyt-c translocation to the cytosol during apoptosis [56]. This flexibility further enhances apoptosome activation, suggesting that Cyt-c and Apaf-1 interactions may follow an "induced fit" mechanism, particularly in variants with greater conformational flexibility [22]. Structural analyses show that certain highly conserved residues are key to the structure, function, folding, and stability of Cyt-c [71,72]. Conserved Tyr68 in the 60-70 helix extends its side chain into the heme pocket and forms H-bonds with Asn53, Thr79, and Met81 [59] which are important for electron transfer and apoptosis initiation as well as heme pocket stability [56,73–77]. Disruption of Tyr68-Asn53, Tyr68-Thr79, and Tyr68-Met81 H-bonds increases caspase activation, impacting peroxidase activity and apoptosis [59,74]. Our study found that the T20I, V21G, N32H, G42S, Y49H, A52T, A52V, Y98H, L99V, and T103I mutants affect H-bonds. This loss can destabilize the heme conformation by altering the orientation of Met81, Asn53, and Thr79, leading to an open active conformation. We further observed that the increased flexibility of the mutant proteins causes an increase in the distance between critical electron-transfer residues in the active site. These changes facilitate the access of small molecules like H_2O_2 to the heme catalytic site, promoting peroxidase activity and apoptosis [59]. The increased distance between the electron-transfer residues may also lead to the inability to transfer the electrons [78].

Variants H27Y, G42S, Y49H, A52T, A52V, R92G, and L99V have been previously described in the literature. H27Y, found in a Chinese family with mild THC4 and low platelet counts, can destabilize protein structure and impair function [27]. Normally, H27 forms a crucial H-bond with P45 for maintaining heme orientation and α-helices, affecting cardiolipin binding to Cyt-c and

apoptosis [79]. Replacing histidine with tyrosine may affect electron transport in the mitochondrial respiratory chain [27]. G42S shows enhanced peroxidase activity and proapoptotic effects by altering heme electronic structure and accelerating electron exchange [12,21] G42S and Y49H variants reduce respiratory levels and increase apoptosis in yeast and mouse models [13]. A52V increases peroxidase activity by destabilizing Cyt-c native conformation. G42S, Y49H, and A52V reduce overall and local stability compared to WT Cyt-c [80]. From a clinical perspective, patients with G42S and Y49H variants have reduced mean platelet counts but normal platelet volume and function, while the A52V family shows lower mean platelet counts with a platelet function defect [18]. G42S, Y49H, A52T, R92G, and L99V variants have been described in Italian families with low platelet counts, where A52T, and R92G variants showed effects on oxidative growth and respiratory activity in a yeast model [17].

To validate our bioinformatics findings, future studies could employ biochemical and molecular techniques to examine the effects of Cyt-c mutations on peroxidase activity and apoptosis. This process includes introducing mutations via site-directed mutagenesis, expressing and purifying the mutated Cyt-c protein, and measuring peroxidase activity using substrate oxidation assays. Apoptosis can be assessed through caspase activation assays, Cyt-c release assays, and cell viability tests. Additionally, structural changes can be investigated using X-ray crystallography or NMR spectroscopy, whereas temperature scans followed by circular dichroism and/or differential scanning calorimetry may give a quantitative measurement of the protein stability. These experiments may confirm our computational predictions and enhance our understanding of the role of Cyt-c mutations in THC4.

5   References

[1]   J.M. Stevens, D.A.I. Mavridou, R. Hamer, P. Kritsiligkou, A.D. Goddard, S.J. Ferguson, Cytochrome c biogenesis System I, FEBS J. 278 (2011) 4170–4178. https://doi.org/10.1111/j.1742-4658.2011.08376.x.

[2]   V.E. Kagan, V.A. Tyurin, J. Jiang, Y.Y. Tyurina, V.B. Ritov, A.A. Amoscato, A.N. Osipov, N.A. Belikova, A.A. Kapralov, V. Kini, I.I. Vlasova, Q. Zhao, M. Zou, P. Di, D.A. Svistunenko, I. V Kurnikov, G.G. Borisenko, Cytochrome c acts as a cardiolipin oxygenase required for release of proapoptotic factors, Nat. Chem. Biol. 1 (2005) 223–232. https://doi.org/10.1038/nchembio727.

[3]   H.A. Kalpage, J. Wan, P.T. Morse, M.P. Zurek, A.A. Turner, A. Khobeir, N. Yazdi, L. Hakim, J. Liu, A. Vaishnav, T.H. Sanderson, M.-A. Recanati, L.I. Grossman, I. Lee, B.F.P. Edwards, M. Hüttemann, Cytochrome c phosphorylation: Control of mitochondrial electron transport chain flux and apoptosis, Int. J. Biochem. Cell Biol. 121 (2020) 105704. https://doi.org/10.1016/j.biocel.2020.105704.

[4]   Y.-L.P. Ow, D.R. Green, Z. Hao, T.W. Mak, Cytochrome c: functions beyond respiration, Nat. Rev. Mol. Cell Biol. 9 (2008) 532–542. https://doi.org/10.1038/nrm2434.

[5]   G.R. Moore, G.W. Pettigrew, Cytochromes c, Springer Berlin Heidelberg, Berlin, Heidelberg, 1990. https://doi.org/10.1007/978-3-642-74536-2.

[6]   G.W. Bushnell, G. V. Louie, G.D. Brayer, High-resolution three-dimensional structure of horse heart cytochrome c, J. Mol. Biol. 214 (1990) 585–595. https://doi.org/10.1016/0022-2836(90)90200-6.

[7]   D. Alvarez-Paggi, L. Hannibal, M.A. Castro, S. Oviedo-Rouco, V. Demicheli, V. Tórtora, F. Tomasina, R. Radi, D.H. Murgida, Multifunctional Cytochrome c : Learning New Tricks from an Old Dog, Chem. Rev. 117 (2017) 13382–13460. https://doi.org/10.1021/acs.chemrev.7b00257.

[8]   R. Santucci, F. Sinibaldi, P. Cozza, F. Polticelli, L. Fiorucci, Cytochrome c: An extreme multifunctional protein with a key role in cell fate, Int. J. Biol. Macromol. 136 (2019) 1237–1246. https://doi.org/10.1016/j.ijbiomac.2019.06.180.

[9]   Y. Zhao, Z.-B. Wang, J.-X. Xu, Effect of cytochrome c on the generation and elimination of O2*- and H2O2 in mitochondria, J. Biol. Chem. 278 (2003) 2356–2360. https://doi.org/10.1074/jbc.M209681200.

[10]   L. Min, X. Jian-xing, Detoxifying function of cytochrome c against oxygen toxicity, Mitochondrion. 7 (2007) 13–16. https://doi.org/10.1016/j.mito.2006.11.011.

[11]   G.C. Brown, V. Borutaite, Regulation of apoptosis by the redox state of cytochrome c, Biochim. Biophys. Acta - Bioenerg. 1777 (2008) 877–881. https://doi.org/10.1016/j.bbabio.2008.03.024.

[12]   I.M. Morison, E.M. Cramer Bordé, E.J. Cheesman, P.L. Cheong, A.J. Holyoake, S. Fichelson, R.J. Weeks, A. Lo, S.M.K. Davies, S.M. Wilbanks, R.D. Fagerlund, M.W. Ludgate, F.M. da Silva Tatley, M.S.A. Coker, N.A. Bockett, G. Hughes, D.A. Pippig, M.P. Smith, C. Capron, E.C. Ledgerwood, A mutation of human cytochrome c enhances the intrinsic apoptotic pathway but causes only thrombocytopenia, Nat. Genet. 40 (2008) 387–389. https://doi.org/10.1038/ng.103.

[13]   D. De Rocco, C. Cerqua, P. Goffrini, G. Russo, A. Pastore, F. Meloni, E. Nicchia, C.T. Moraes, A. Pecci, L. Salviati, A. Savoia, Mutations of cytochrome c identified in patients with thrombocytopenia THC4 affect both apoptosis and cellular bioenergetics, Biochim. Biophys. Acta - Mol. Basis Dis. 1842 (2014) 269–274.

https://doi.org/10.1016/j.bbadis.2013.12.002.

[14] B. Johnson, G.C. Lowe, J. Futterer, M. Lordkipanidze, D. MacDonald, M.A. Simpson, I. Sanchez-Guiu, S. Drake, D. Bem, V. Leo, S.J. Fletcher, B. Dawood, J. Rivera, D. Allsup, T. Biss, P.H. Bolton-Maggs, P. Collins, N. Curry, C. Grimley, B. James, M. Makris, J. Motwani, S. Pavord, K. Talks, J. Thachil, J. Wilde, M. Williams, P. Harrison, P. Gissen, S. Mundell, A. Mumford, M.E. Daly, S.P. Watson, N. V. Morgan, Whole exome sequencing identifies genetic variants in inherited thrombocytopenia with secondary qualitative function defects, Haematologica. 101 (2016) 1170–1179. https://doi.org/10.3324/haematol.2016.146316.

[15] Y. Uchiyama, K. Yanagisawa, S. Kunishima, M. Shiina, Y. Ogawa, M. Nakashima, J. Hirato, E. Imagawa, A. Fujita, K. Hamanaka, S. Miyatake, S. Mitsuhashi, A. Takata, N. Miyake, K. Ogata, H. Handa, N. Matsumoto, T. Mizuguchi, A novel CYCS mutation in the α-helix of the CYCS C-terminal domain causes non-syndromic thrombocytopenia, Clin. Genet. 94 (2018) 548–553. https://doi.org/10.1111/cge.13423.

[16] C.L. Balduini, A. Savoia, Genetics of familial forms of thrombocytopenia, Hum. Genet. 131 (2012) 1821–1832. https://doi.org/10.1007/s00439-012-1215-x.

[17] A. Marzollo, S. Zampieri, S. Barozzi, M.A. Yousaf, J. Quartararo, D. De Rocco, M. Faleschini, C. Marconi, C.C. Berti, V. Bozzi, G. Russo, P. Giordano, P. Goffrini, S. Bresolin, A. Pastore, A. Savoia, A. Pecci, Thrombocytopenia 4 (<scp>THC4</scp>): Six novel families with mutations of the cytochrome c gene, Br. J. Haematol. (2024). https://doi.org/10.1111/bjh.19567.

[18] E.C. Ledgerwood, C. Dunstan-Harrison, L. Ong, I.M. Morison, CYCS gene variants associated with thrombocytopenia, Platelets. 30 (2019) 672–674. https://doi.org/10.1080/09537104.2018.1543866.

[19] L. Salviati, Cytochrome c Defects in Human Disease, in: Mitochondrial Dis., Springer International Publishing, Cham, 2021: pp. 191–200. https://doi.org/10.1007/978-3-030-70147-5_7.

[20] E. Turro, W.J. Astle, K. Megy, S. Gräf, D. Greene, O. Shamardina, H.L. Allen, A. Sanchis-Juan, M. Frontini, C. Thys, J. Stephens, R. Mapeta, O.S. Burren, K. Downes, M. Haimel, S. Tuna, S.V. V. Deevi, T.J. Aitman, D.L. Bennett, P. Calleja, K. Carss, M.J. Caulfield, P.F. Chinnery, P.H. Dixon, D.P. Gale, R. James, A. Koziell, M.A. Laffan, A.P. Levine, E.R. Maher, H.S. Markus, J. Morales, N.W. Morrell, A.D. Mumford, E. Ormondroyd, S. Rankin, A. Rendon, S. Richardson, I. Roberts, N.B.A. Roy, M.A. Saleem, K.G.C. Smith, H. Stark, R.Y.Y. Tan, A.C. Themistocleous, A.J. Thrasher, H. Watkins, A.R. Webster, M.R. Wilkins, C. Williamson, J. Whitworth, S. Humphray, D.R. Bentley, S. Abbs, L. Abulhoul, J. Adlard, M. Ahmed, T.J. Aitman, H. Alachkar, D.J. Allsup, J. Almeida-King, P. Ancliff, R. Antrobus, R. Armstrong, G. Arno, S. Ashford, W.J. Astle, A. Attwood, P. Aurora, C. Babbs, C. Bacchelli, T. Bakchoul, S. Banka, T. Bariana, J. Barwell, J. Batista, H.E. Baxendale, P.L. Beales, D.L. Bennett, D.R. Bentley, A. Bierzynska, T. Biss, M.A.K. Bitner-Glindzicz, G.C. Black, M. Bleda, I. Blesneac, D. Bockenhauer, H. Bogaard, C.J. Bourne, S. Boyce, J.R. Bradley, E. Bragin, G. Breen, P. Brennan, C. Brewer, M. Brown, A.C. Browning, M.J. Browning, R.J. Buchan, M.S. Buckland, T. Bueser, C.B. Diz, J. Burn, S.O. Burns, O.S. Burren, N. Burrows, P. Calleja, C. Campbell, G. Carr-White, K. Carss, R. Casey, M.J. Caulfield, J. Chambers, J. Chambers, M.M.Y. Chan, C. Cheah, F. Cheng, P.F. Chinnery, M. Chitre, M.T. Christian, C. Church, J. Clayton-Smith, M. Cleary, N.C. Brod, G. Coghlan, E. Colby, T.R.P. Cole, J. Collins, P.W. Collins, C. Colombo, C.J. Compton, R. Condliffe,

S. Cook, H.T. Cook, N. Cooper, P.A. Corris, A. Furnell, F. Cunningham, N.S. Curry, A.J. Cutler, M.J. Daniels, M. Dattani, L.C. Daugherty, J. Davis, A. De Soyza, S.V. V. Deevi, T. Dent, C. Deshpande, E.F. Dewhurst, P.H. Dixon, S. Douzgou, K. Downes, A.M. Drazyk, E. Drewe, D. Duarte, T. Dutt, J.D.M. Edgar, K. Edwards, W. Egner, M.N. Ekani, P. Elliott, W.N. Erber, M. Erwood, M.C. Estiu, D.G. Evans, G. Evans, T. Everington, M. Eyries, H. Fassihi, R. Favier, J. Findhammer, D. Fletcher, F.A. Flinter, R.A. Floto, T. Fowler, J. Fox, A.J. Frary, C.E. French, K. Freson, M. Frontini, D.P. Gale, H. Gall, V. Ganesan, M. Gattens, C. Geoghegan, T.S.A. Gerighty, A.G. Gharavi, S. Ghio, H.-A. Ghofrani, J.S.R. Gibbs, K. Gibson, K.C. Gilmour, B. Girerd, N.S. Gleadall, S. Goddard, D.B. Goldstein, K. Gomez, P. Gordins, D. Gosal, S. Gräf, J. Graham, L. Grassi, D. Greene, L. Greenhalgh, A. Greinacher, P. Gresele, P. Griffiths, S. Grigoriadou, R.J. Grocock, D. Grozeva, M. Gurnell, S. Hackett, C. Hadinnapola, W.M. Hague, R. Hague, M. Haimel, M. Hall, H.L. Hanson, E. Haque, K. Harkness, A.R. Harper, C.L. Harris, D. Hart, A. Hassan, G. Hayman, A. Henderson, A. Herwadkar, J. Hoffman, S. Holden, R. Horvath, H. Houlden, A.C. Houweling, L.S. Howard, F. Hu, G. Hudson, J. Hughes, A.P. Huissoon, M. Humbert, S. Humphray, S. Hunter, M. Hurles, M. Irving, L. Izatt, R. James, S.A. Johnson, S. Jolles, J. Jolley, D. Josifova, N. Jurkute, T. Karten, J. Karten, M.A. Kasanicki, H. Kazkaz, R. Kazmi, P. Kelleher, A.M. Kelly, W. Kelsall, C. Kempster, D.G. Kiely, N. Kingston, R. Klima, N. Koelling, M. Kostadima, G. Kovacs, A. Koziell, R. Kreuzhuber, T.W. Kuijpers, A. Kumar, D. Kumararatne, M.A. Kurian, M.A. Laffan, F. Lalloo, M. Lambert, H.L. Allen, A. Lawrie, D.M. Layton, N. Lench, C. Lentaigne, T. Lester, A.P. Levine, R. Linger, H. Longhurst, L.E. Lorenzo, E. Louka, P.A. Lyons, R.D. Machado, R. V. MacKenzie Ross, B. Madan, E.R. Maher, J. Maimaris, S. Malka, S. Mangles, R. Mapeta, K.J. Marchbank, S. Marks, H.S. Markus, H.-U. Marschall, A. Marshall, J. Martin, M. Mathias, E. Matthews, H. Maxwell, P. McAlinden, M.I. McCarthy, H. McKinney, A. McMahon, S. Meacham, A.J. Mead, I.M. Castello, K. Megy, S.G. Mehta, M. Michaelides, C. Millar, S.N. Mohammed, S. Moledina, D. Montani, A.T. Moore, J. Morales, N.W. Morrell, M. Mozere, K.W. Muir, A.D. Mumford, A.H. Nemeth, W.G. Newman, M. Newnham, S. Noorani, P. Nurden, J. O'Sullivan, S. Obaji, C. Odhams, S. Okoli, A. Olschewski, H. Olschewski, K.R. Ong, S.H. Oram, E. Ormondroyd, W.H. Ouwehand, C. Palles, S. Papadia, S.-M. Park, D. Parry, S. Patel, J. Paterson, A. Peacock, S.H. Pearce, J. Peden, K. Peerlinck, C.J. Penkett, J. Pepke-Zaba, R. Petersen, C. Pilkington, K.E.S. Poole, R. Prathalingam, B. Psaila, A. Pyle, R. Quinton, S. Rahman, S. Rankin, A. Rao, F.L. Raymond, P.J. Rayner-Matthews, C. Rees, A. Rendon, T. Renton, C.J. Rhodes, A.S.C. Rice, S. Richardson, A. Richter, L. Robert, I. Roberts, A. Rogers, S.J. Rose, R. Ross-Russell, C. Roughley, N.B.A. Roy, D.M. Ruddy, O. Sadeghi-Alavijeh, M.A. Saleem, N. Samani, C. Samarghitean, A. Sanchis-Juan, R.B. Sargur, R.N. Sarkany, S. Satchell, S. Savic, J.A. Sayer, G. Sayer, L. Scelsi, A.M. Schaefer, S. Schulman, R. Scott, M. Scully, C. Searle, W. Seeger, A. Sen, W.A.C. Sewell, D. Seyres, N. Shah, O. Shamardina, S.E. Shapiro, A.C. Shaw, P.J. Short, K. Sibson, L. Side, I. Simeoni, M.A. Simpson, M.C. Sims, S. Sivapalaratnam, D. Smedley, K.R. Smith, K.G.C. Smith, K. Snape, N. Soranzo, F. Soubrier, L. Southgate, O. Spasic-Boskovic, S. Staines, E. Staples, H. Stark, J. Stephens, C. Steward, K.E. Stirrups, A. Stuckey, J. Suntharalingam, E.M. Swietlik, P. Syrris, R.C. Tait, K. Talks, R.Y.Y. Tan, K. Tate, J.M. Taylor, J.C. Taylor, J.E. Thaventhiran, A.C. Themistocleous, E. Thomas, D. Thomas, M.J. Thomas, P. Thomas, K. Thomson, A.J. Thrasher, G. Threadgold, C. Thys, T. Tilly, M. Tischkowitz, C. Titterton, J.A. Todd, C.-H. Toh, B. Tolhuis, I.P. Tomlinson, M. Toshner, M. Traylor, C. Treacy, P. Treadaway, R.

Trembath, S. Tuna, W. Turek, E. Turro, P. Twiss, T. Vale, C. Van Geet, N. van Zuydam, M. Vandekuilen, A.M. Vandersteen, M. Vazquez-Lopez, J. von Ziegenweidt, A. Vonk Noordegraaf, A. Wagner, Q. Waisfisz, S.M. Walker, N. Walker, K. Walter, J.S. Ware, H. Watkins, C. Watt, A.R. Webster, L. Wedderburn, W. Wei, S.B. Welch, J. Wessels, S.K. Westbury, J.-P. Westwood, J. Wharton, D. Whitehorn, J. Whitworth, A.O.M. Wilkie, M.R. Wilkins, C. Williamson, B.T. Wilson, E.K.S. Wong, N. Wood, Y. Wood, C.G. Woods, E.R. Woodward, S.J. Wort, A. Worth, M. Wright, K. Yates, P.F.K. Yong, T. Young, P. Yu, P. Yu-Wai-Man, E. Zlamalova, N. Kingston, N. Walker, J.R. Bradley, S. Ashford, C.J. Penkett, K. Freson, K.E. Stirrups, F.L. Raymond, W.H. Ouwehand, Whole-genome sequencing of patients with rare diseases in a national health system, Nature. 583 (2020) 96–102. https://doi.org/10.1038/s41586-020-2434-2.

[21] M.D. Liptak, R.D. Fagerlund, E.C. Ledgerwood, S.M. Wilbanks, K.L. Bren, The Proapoptotic G41S Mutation to Human Cytochrome c Alters the Heme Electronic Structure and Increases the Electron Self-Exchange Rate, J. Am. Chem. Soc. 133 (2011) 1153–1155. https://doi.org/10.1021/ja106328k.

[22] M. Fellner, R. Parakra, K.O. McDonald, I. Kass, G.N.L. Jameson, S.M. Wilbanks, E.C. Ledgerwood, Altered structure and dynamics of pathogenic cytochrome c variants correlate with increased apoptotic activity, Biochem. J. 478 (2021) 669–684. https://doi.org/10.1042/BCJ20200793.

[23] H.M. Berman, J. Westbrook, Z. Feng, G. Gilliland, T.N. Bhat, H. Weissig, I.N. Shindyalov, P.E. Bourne, The Protein Data Bank, Nucleic Acids Res. 28 (2000) 235–242. https://doi.org/10.1093/nar/28.1.235.

[24] A. Bateman, UniProt: A worldwide hub of protein knowledge, Nucleic Acids Res. 47 (2019) D506–D515. https://doi.org/10.1093/nar/gky1049.

[25] F. Sievers, D.G. Higgins, Clustal Omega for making accurate alignments of many protein sequences, Protein Sci. 27 (2018) 135–145. https://doi.org/10.1002/pro.3290.

[26] K. Megy, K. Downes, M. Morel-Kopp, J.M. Bastida, S. Brooks, L. Bury, E. Leinoe, K. Gomez, N. V. Morgan, M. Othman, W.H. Ouwehand, J. Perez Botero, J. Rivera, H. Schulze, D. Trégouët, K. Freson, GoldVariants, a resource for sharing rare genetic variants detected in bleeding, thrombotic, and platelet disorders: Communication from the ISTH SSC Subcommittee on Genomics in Thrombosis and Hemostasis, J. Thromb. Haemost. 19 (2021) 2612–2617. https://doi.org/10.1111/jth.15459.

[27] F. Che, J. Zhao, Y. Zhao, Z. Wang, L. Zhang, Y. Yang, A Novel Heterozygous Pathogenic Variation in CYCS Gene Cause Autosomal Dominant Non-Syndromic Thrombocytopenia 4 in a Large Chinese Family, Front. Genet. 12 (2022). https://doi.org/10.3389/fgene.2021.783455.

[28] K. Downes, K. Megy, D. Duarte, M. Vries, J. Gebhart, S. Hofer, O. Shamardina, S.V. V. Deevi, J. Stephens, R. Mapeta, S. Tuna, N. Al Hasso, M.W. Besser, N. Cooper, L. Daugherty, N. Gleadall, D. Greene, M. Haimel, H. Martin, S. Papadia, S. Revel-Vilk, S. Sivapalaratnam, E. Symington, W. Thomas, C. Thys, A. Tolios, C.J. Penkett, W.H. Ouwehand, S. Abbs, M.A. Laffan, E. Turro, I. Simeoni, A.D. Mumford, Y.M.C. Henskens, I. Pabinger, K. Gomez, K. Freson, Diagnostic high-throughput sequencing of 2396 patients with bleeding, thrombotic, and platelet disorders, Blood. 134 (2019) 2082–2091. https://doi.org/10.1182/blood.2018891192.

[29] E. Gálvez, E. Vallespín, E.G. Arias-Salgado, C. Sánchez-Valdepeñas, Y. Giménez, S. Navarro, P. Río, M. Bogliolo, R. Pujol, M. Peiró, J. Nevado, J. Zubicaray, E. Sebastián, A.

Catalá, C. Beléndez, C. Díaz de Heredia, A. Galera, I. Badell, L. Madero, R. Perona, L. Sastre, J. Surrallés, J. Bueren, P. Lapunzina, J. Sevilla, Next-generation Sequencing in Bone Marrow Failure Syndromes and Isolated Cytopenias: Experience of the Spanish Network on Bone Marrow Failure Syndromes, HemaSphere. 5 (2021) e539. https://doi.org/10.1097/HS9.0000000000000539.

[30] O. Gilad, O. Dgany, S. Noy-Lotan, T. Krasnov, J. Yacobovich, R. Rabinowicz, T. Goldberg, A.A. Kuperman, A. Abu-Quider, H. Miskin, N. Kapelushnik, N. Mandel-Shorer, S. Shimony, D. Harlev, T. Ben-Ami, E. Adam, C. Levin, S. Aviner, R. Elhasid, S. Berger-Achituv, L. Chaitman-Yerushalmi, Y. Kodman, N. Oniashvilli, M. Hameiri-Grosman, S. Izraeli, H. Tamary, O. Steinberg-Shemer, Syndromes predisposing to leukemia are a major cause of inherited cytopenias in children, Haematologica. 107 (2022) 2081–2095. https://doi.org/10.3324/haematol.2021.280116.

[31] K. Giavi, S. Glentis, A. Bouchla, A. Apostolidou, N.M. Marinakis, A. Kattamis, E. Katsantoni, V. Pappa, A Novel Variant of the CYCS Gene Alters Apoptosis of Megakaryocytes in a Family with Thrombocytopenia, Blood. 142 (2023) 5409–5409. https://doi.org/10.1182/blood-2023-186060.

[32] P.C. Ng, S. Henikoff, SIFT: predicting amino acid changes that affect protein function, Nucleic Acids Res. 31 (2003) 3812–3814. https://doi.org/10.1093/NAR/GKG509.

[33] C.M. Yates, I. Filippis, L.A. Kelley, M.J.E. Sternberg, SuSPect: Enhanced prediction of single amino acid variant (SAV) phenotype using network features, J. Mol. Biol. 426 (2014) 2692–2701. https://doi.org/10.1016/j.jmb.2014.04.026.

[34] V. Pejaver, J. Urresti, J. Lugo-Martinez, K.A. Pagel, G.N. Lin, H.J. Nam, M. Mort, D.N. Cooper, J. Sebat, L.M. Iakoucheva, S.D. Mooney, P. Radivojac, Inferring the molecular and phenotypic impact of amino acid variants with MutPred2, Nat. Commun. 11 (2020). https://doi.org/10.1038/s41467-020-19669-x.

[35] W. Deng, Y. Wang, L. Ma, Y. Zhang, S. Ullah, Y. Xue, Computational prediction of methylation types of covalently modified lysine and arginine residues in proteins, Brief. Bioinform. (2016) bbw041. https://doi.org/10.1093/bib/bbw041.

[36] N. Blom, S. Gammeltoft, S. Brunak, Sequence and structure-based prediction of eukaryotic protein phosphorylation sites, J. Mol. Biol. 294 (1999) 1351–1362. https://doi.org/10.1006/jmbi.1999.3310.

[37] V.-N. Nguyen, K.-Y. Huang, J.T.-Y. Weng, K.R. Lai, T.-Y. Lee, UbiNet: an online resource for exploring the functional associations and regulatory networks of protein ubiquitylation, Database. 2016 (2016) baw054. https://doi.org/10.1093/database/baw054.

[38] C. Steentoft, S.Y. Vakhrushev, H.J. Joshi, Y. Kong, M.B. Vester-Christensen, K.T.B.G. Schjoldager, K. Lavrsen, S. Dabelsteen, N.B. Pedersen, L. Marcos-Silva, R. Gupta, E. Paul Bennett, U. Mandel, S. Brunak, H.H. Wandall, S.B. Levery, H. Clausen, Precision mapping of the human O-GalNAc glycoproteome through SimpleCell technology, EMBO J. 32 (2013) 1478–1488. https://doi.org/10.1038/emboj.2013.79.

[39] D. Warde-Farley, S.L. Donaldson, O. Comes, K. Zuberi, R. Badrawi, P. Chao, M. Franz, C. Grouios, F. Kazi, C.T. Lopes, A. Maitland, S. Mostafavi, J. Montojo, Q. Shao, G. Wright, G.D. Bader, Q. Morris, The GeneMANIA prediction server: biological network integration for gene prioritization and predicting gene function, Nucleic Acids Res. 38 (2010) W214–W220. https://doi.org/10.1093/nar/gkq537.

[40] D. Szklarczyk, A.L. Gable, K.C. Nastou, D. Lyon, R. Kirsch, S. Pyysalo, N.T. Doncheva, M. Legeay, T. Fang, P. Bork, L.J. Jensen, C. von Mering, The STRING database in 2021:

Customizable protein-protein networks, and functional characterization of user-uploaded gene/measurement sets, Nucleic Acids Res. 49 (2021) D605–D612. https://doi.org/10.1093/nar/gkaa1074.

[41] C.H.M. Rodrigues, D.E.V. Pires, D.B. Ascher, DynaMut2 : Assessing changes in stability and flexibility upon single and multiple point missense mutations, Protein Sci. 30 (2021) 60–69. https://doi.org/10.1002/pro.3942.

[42] J. Cheng, A. Randall, P. Baldi, Prediction of protein stability changes for single-site mutations using support vector machines, Proteins Struct. Funct. Bioinforma. 62 (2005) 1125–1132. https://doi.org/10.1002/prot.20810.

[43] M.J. Abraham, T. Murtola, R. Schulz, S. Páll, J.C. Smith, B. Hess, E. Lindah, Gromacs: High performance molecular simulations through multi-level parallelism from laptops to supercomputers, SoftwareX. 1–2 (2015) 19–25. https://doi.org/10.1016/J.SOFTX.2015.06.001.

[44] B. Hess, C. Kutzner, D. Van Der Spoel, E. Lindahl, GROMACS 4: algorithms for highly efficient, load-balanced, and scalable molecular simulation, J. Chem. Theory Comput. 4 (2008) 435–447. https://doi.org/10.1021/ct700301q.

[45] M. Landau, I. Mayrose, Y. Rosenberg, F. Glaser, E. Martz, T. Pupko, N. Ben-Tal, ConSurf 2005: the projection of evolutionary conservation scores of residues on protein structures, Nucleic Acids Res. 33 (2005) W299–W302. https://doi.org/10.1093/nar/gki370.

[46] G. Celniker, G. Nimrod, H. Ashkenazy, F. Glaser, E. Martz, I. Mayrose, T. Pupko, N. Ben-Tal, ConSurf: Using Evolutionary Data to Raise Testable Hypotheses about Protein Function, Isr. J. Chem. 53 (2013) 199–206. https://doi.org/10.1002/ijch.201200096.

[47] A.M. Lesk, C. Chothia, How different amino acid sequences determine similar protein structures: The structure and evolutionary dynamics of the globins, J. Mol. Biol. 136 (1980) 225–270. https://doi.org/10.1016/0022-2836(80)90373-3.

[48] A. Waterhouse, M. Bertoni, S. Bienert, G. Studer, G. Tauriello, R. Gumienny, F.T. Heer, T.A.P. De Beer, C. Rempfer, L. Bordoli, R. Lepore, T. Schwede, SWISS-MODEL: Homology modelling of protein structures and complexes, Nucleic Acids Res. 46 (2018) W296–W303. https://doi.org/10.1093/nar/gky427.

[49] R.A. Laskowski, M.W. MacArthur, D.S. Moss, J.M. Thornton, PROCHECK: a program to check the stereochemical quality of protein structures, J. Appl. Crystallogr. 26 (1993) 283–291. https://doi.org/10.1107/S0021889892009944.

[50] B. Nutho, S. Samsri, S. Pornsuwan, Structural Dynamics of the Precatalytic State of Human Cytochrome c upon T28C, G34C, and A50C Mutations: A Molecular Dynamics Simulation Perspective, ACS Omega. 8 (2023) 15229–15238. https://doi.org/10.1021/acsomega.3c00220.

[51] F. Sinibaldi, B.D. Howes, M.C. Piro, P. Caroppi, G. Mei, F. Ascoli, G. Smulevich, R. Santucci, Insights into the role of the histidines in the structure and stability of cytochrome c, JBIC J. Biol. Inorg. Chem. 11 (2006) 52–62. https://doi.org/10.1007/s00775-005-0057-6.

[52] S. Ittisoponpisan, S.A. Islam, T. Khanna, E. Alhuzimi, A. David, M.J.E. Sternberg, Can Predicted Protein 3D Structures Provide Reliable Insights into whether Missense Variants Are Disease Associated?, J. Mol. Biol. 431 (2019) 2197–2212. https://doi.org/10.1016/j.jmb.2019.04.009.

[53] K. Shinwari, H.M. Rehman, N. Xiao, L. Guojun, M.A. Khan, M.A. Bolkov, I.A. Tuzankina, V.A. Chereshnev, Novel high-risk missense mutations identification in FAT4 gene causing

Hennekam syndrome and Van Maldergem syndrome 2 through molecular dynamics simulation, Informatics Med. Unlocked. 37 (2023) 101160. https://doi.org/10.1016/j.imu.2023.101160.

[54] Y. Shiloh, Y. Ziv, The ATM protein kinase: regulating the cellular response to genotoxic stress, and more, Nat. Rev. Mol. Cell Biol. 14 (2013) 197–210. https://doi.org/10.1038/nrm3546.

[55] P. Brochette, C. Petit, M.P. Pileni, Cytochrome c in sodium bis(2-ethylhexyl) sulfosuccinate reverse micelles: structure and reactivity, J. Phys. Chem. 92 (1988) 3505–3511. https://doi.org/10.1021/j100323a037.

[56] G. Muneeswaran, S. Kartheeswaran, M. Pandiaraj, K. Muthukumar, M. Sankaralingam, S. Arunachalam, Investigation of structural dynamics of Thrombocytopenia Cargeeg mutants of human apoptotic cytochrome c : A molecular dynamics simulation approach, Biophys. Chem. 230 (2017) 117–126. https://doi.org/10.1016/j.bpc.2017.09.004.

[57] S. Abel, M. Waks, M. Marchi, Molecular dynamics simulations of cytochrome c unfolding in AOT reverse micelles: The first steps, Eur. Phys. J. E. 32 (2010) 399–409. https://doi.org/10.1140/epje/i2010-10635-x.

[58] G. Muneeswaran, S. Kartheeswaran, K. Muthukumar, C.D. Dharmaraj, C. Karunakaran, Comparative structural and conformational studies on H43R and W32F mutants of copper–zinc superoxide dismutase by molecular dynamics simulation, Biophys. Chem. 185 (2014) 70–78. https://doi.org/10.1016/j.bpc.2013.11.010.

[59] G. Muneeswaran, M. Pandiaraj, S. Kartheeswaran, M. Sankaralingam, K. Muthukumar, C. Karunakaran, Molecular dynamics simulation approach to explore atomistic molecular mechanism of peroxidase activity of apoptotic cytochrome c mutants, Informatics Med. Unlocked. 11 (2018) 51–60. https://doi.org/10.1016/j.imu.2018.04.003.

[60] A.K. Agrahari, P. Sneha, C. George Priya Doss, R. Siva, H. Zayed, A profound computational study to prioritize the disease-causing mutations in PRPS1 gene, Metab. Brain Dis. 33 (2018) 589–600. https://doi.org/10.1007/s11011-017-0121-2.

[61] Z.-P. Wang, X.-Z. Ding, J. Wang, Y.-M. Li, Double-edged sword in cells: chemical biology studies of the vital role of cytochrome c in the intrinsic pre-apoptotic mitochondria leakage pathway, RSC Adv. 5 (2015) 28258–28269. https://doi.org/10.1039/C4RA16856A.

[62] J. Zheng, S. Ye, T. Lu, T.M. Cotton, G. Chumanov, Circular dichroism and resonance Raman comparative studies of wild type cytochromec and F82H mutant, Biopolymers. 57 (2000) 77–84. https://doi.org/10.1002/(SICI)1097-0282(2000)57:2<77::AID-BIP4>3.0.CO;2-Y.

[63] T. Takano, O.B. Kallai, R. Swanson, R.E. Dickerson, The Structure of Ferrocytochrome c at 2.45 A Resolution, J. Biol. Chem. 248 (1973) 5234–5255. https://doi.org/10.1016/S0021-9258(19)43594-1.

[64] M. Hüttemann, P. Pecina, M. Rainbolt, T.H. Sanderson, V.E. Kagan, L. Samavati, J.W. Doan, I. Lee, The multiple functions of cytochrome c and their regulation in life and death decisions of the mammalian cell: From respiration to apoptosis, Mitochondrion. 11 (2011) 369–381. https://doi.org/10.1016/j.mito.2011.01.010.

[65] A.M. Firsov, E.A. Kotova, E.A. Korepanova, A.N. Osipov, Y.N. Antonenko, Peroxidative permeabilization of liposomes induced by cytochrome c/cardiolipin complex, Biochim. Biophys. Acta - Biomembr. 1848 (2015) 767–774. https://doi.org/10.1016/j.bbamem.2014.11.027.

[66] R. V. Chertkova, A.M. Firsov, N.A. Brazhe, E.I. Nikelshparg, Z. V. Bochkova, T. V.

Bryantseva, M.A. Semenova, A.A. Baizhumanov, E.A. Kotova, M.P. Kirpichnikov, G. V. Maksimov, Y.N. Antonenko, D.A. Dolgikh, Multiple Mutations in the Non-Ordered Red Ω-Loop Enhance the Membrane-Permeabilizing and Peroxidase-like Activity of Cytochrome c, Biomolecules. 12 (2022) 665. https://doi.org/10.3390/biom12050665.

[67] A. Guerra-Castellano, I. Márquez, G. Pérez-Mejías, A. Díaz-Quintana, M.A. De la Rosa, I. Díaz-Moreno, Post-Translational Modifications of Cytochrome c in Cell Life and Disease, Int. J. Mol. Sci. 21 (2020) 8483. https://doi.org/10.3390/ijms21228483.

[68] O.M. Deacon, R.W. White, G.R. Moore, M.T. Wilson, J.A.R. Worrall, Comparison of the structural dynamic and mitochondrial electron-transfer properties of the proapoptotic human cytochrome c variants, G41S, Y48H and A51V, J. Inorg. Biochem. 203 (2020) 110924. https://doi.org/10.1016/j.jinorgbio.2019.110924.

[69] S. Samsri, S. Pornsuwan, Influence of cysteine-directed mutations at the Ω-loops on peroxidase activity of human cytochrome c., Arch. Biochem. Biophys. 709 (2021) 108980. https://doi.org/10.1016/j.abb.2021.108980.

[70] A.I. Karsisiotis, O.M. Deacon, M.T. Wilson, C. Macdonald, T.M.A. Blumenschein, G.R. Moore, J.A.R. Worrall, Increased dynamics in the 40–57 Ω-loop of the G41S variant of human cytochrome c promote its pro-apoptotic conformation, Sci. Rep. 6 (2016) 30447. https://doi.org/10.1038/srep30447.

[71] S. Sperandio, I. de Belle, D.E. Bredesen, An alternative, nonapoptotic form of programmed cell death, Proc. Natl. Acad. Sci. 97 (2000) 14376–14381. https://doi.org/10.1073/pnas.97.26.14376.

[72] T. Ying, Z.-H. Wang, Y.-W. Lin, J. Xie, X. Tan, Z.-X. Huang, Tyrosine-67 in cytochrome c is a possible apoptotic trigger controlled by hydrogen bonds via a conformational transition, Chem. Commun. (2009) 4512. https://doi.org/10.1039/b904347k.

[73] T.L. Luntz, A. Schejter, E.A. Garber, E. Margoliash, Structural significance of an internal water molecule studied by site-directed mutagenesis of tyrosine-67 in rat cytochrome c., Proc. Natl. Acad. Sci. 86 (1989) 3524–3528. https://doi.org/10.1073/pnas.86.10.3524.

[74] E. Margoliash, S. Ferguson-Miller, J. Tulloss, C.H. Kang, B.A. Feinberg, D.L. Brautigan, M. Morrison, Separate Intramolecular Pathways for Reduction and Oxidation of Cytochrome c in Electron Transport Chain Reactions, Proc. Natl. Acad. Sci. 70 (1973) 3245–3249. https://doi.org/10.1073/pnas.70.11.3245.

[75] L.A. Abriata, A. Cassina, V. Tórtora, M. Marín, J.M. Souza, L. Castro, A.J. Vila, R. Radi, Nitration of Solvent-exposed Tyrosine 74 on Cytochrome c Triggers Heme Iron-Methionine 80 Bond Disruption, J. Biol. Chem. 284 (2009) 17–26. https://doi.org/10.1074/jbc.M807203200.

[76] D. Roccatano, I. Daidone, M.A. Ceruso, C. Bossa, A. Di Nola, Selective excitation of native fluctuations during thermal unfolding simulations: Horse heart cytochrome c as a case study, Biophys. J. 84 (2003) 1876–1883. https://doi.org/10.1016/S0006-3495(03)74995-9.

[77] W. Pulawski, S. Filipek, A. Zwolinska, A. Debinski, K. Krzysko, R. Garduño-Juárez, S. Viswanathan, V. Renugopalakrishnan, Low-temperature molecular dynamics simulations of horse heart cytochrome c and comparison with inelastic neutron scattering data, Eur. Biophys. J. 42 (2013) 291–300. https://doi.org/10.1007/s00249-012-0874-9.

[78] S.R. Singh, S. Prakash, V. Vasu, C. Karunakaran, Conformational flexibility decreased due to Y67F and F82H mutations in cytochrome c: Molecular dynamics simulation studies, J. Mol. Graph. Model. 28 (2009) 270–277. https://doi.org/10.1016/j.jmgm.2009.08.005.

[79] G. Balakrishnan, Y. Hu, T.G. Spiro, His26 Protonation in Cytochrome c Triggers

Microsecond β-Sheet Formation and Heme Exposure: Implications for Apoptosis, J. Am. Chem. Soc. 134 (2012) 19061–19069. https://doi.org/10.1021/ja307100a.

[80] H. Lei, B.E. Bowler, Naturally Occurring A51V Variant of Human Cytochrome c Destabilizes the Native State and Enhances Peroxidase Activity, J. Phys. Chem. B. 123 (2019) 8939–8953. https://doi.org/10.1021/acs.jpcb.9b05869.

Figure Legends

Fig. 1. Overall structure of Cyt-c and information on the position and conservation of the clinically important variants. (A) The Cyt-c variants, T20I, V21G, H27Y, N32H and L33V are located in the proximal loop, G42S, Y49H, A52T and A52V in the central loop and R92G, A97D, Y98H, L99V and T103I in the protein C-terminus. (B) Multiple sequence alignment analysis of Cyt-c with orthologues from different species including *Homo sapiens* (human), *Macaca mulatta* (rhesus monkey), *Mus musculus* (mouse), *Canis lupus familiaris* (dog), *Oryctolagus cuniculus* (European rabbit), *Sus scrofa* (wild boar/pig), *Gallus gallus* (chicken), *Xenopus tropicalis* (western clawed frog) and *Danio rerio* (zebrafish) belonging to diverse taxonomic groups. The alignment showed that all analyzed variants (T20I, V21G, H27Y, N32H, L33V, G42S, Y49H, A52T, A52V, R92G, A97D, Y98H, L99V and T103I) are at evolutionary conserved residues and are likely deleterious. (C) SIFT analysis shows that all variants affect protein function (score < 0.05) and SuSPect associates the variants with the disease (score > 50).

Fig. 2. Mapping the currently known clinically important variants on the structure of Cyt-c, and Ramachandran plot analysis. The protein structure is depicted as a grey ribbon, with heme in green. (A) Human Cyt-c (PDB ID: 1j3s) structure showing three Ω-loop regions: proximal (residues 13-40, magenta), central (residues 41-58, cyan) and distal (residues 72-86, red) Ω-loops. Amino acids investigated for variants are highlighted as follows: (B) T20 (magenta), V21 (blue), H27 (blue), N32 (magenta), L33 (blue); (C) G42 (magenta), Y49 (blue), A52 (blue), R92 (blue); (D) A97 (magenta), Y98 (blue), L99 (red), T103 (blue). Note that the residue numbering is offset by +1 relative to the PDB entry 1j3s. (E) In the Ramachandran plot, G42 is an exception, positioned in a typically disallowed region due to its small size and high flexibility. The G42S variant introduces steric hindrance due to larger and less flexible serine in a disallowed region.

Fig. 3. Visualization of gene and protein interactions. (A) Interactions of the *CYCS* gene with other genes, analyzed using the GeneMANIA tool. (B) The protein-protein interaction network of Cyt-c analyzed using the STRING tool. Cytc variants may affect the functions of interacting genes or proteins.

Fig. 4. Root Mean Square Deviation (RMSD) profiles of WT and mutant Cyt-c protein structures over a 50 ns (50000 ps) of MD simulation. (A) Fluctuations of RMSD values for WT and T20I, V21G, H27Y, N32H and L33V mutant proteins. (B) RMSD values of mutants T20I (0.38 ± 0.03 nm), V21G (0.36 ± 0.04 nm), H27Y (0.39 ± 0.0 nm), N32H (0.42 ± 0.05 nm) and L33V (0.33 ± 0.02 nm) were significantly higher than the WT (0.30 ± 0.03 nm) (****: $p < 0.0001$). (C) The mutant

proteins showed higher RMSD values (D) which were significant (****: p < 0.0001) compared to WT (0.30±0.03 nm). The mutant displayed mean±SD as G42S: 0.42±0.05 nm; Y49H: 0.34±0.04 nm; A52T: 0.37±0.03 nm; A52V: 0.41±0.04 nm; and R92G 0.45±0.06 nm. (E) Variations of RMSD values over simulation where (F) A97D (0.43±0.05 nm), Y98H (0.43±0.03 nm) and T103I (0.45±0.05 nm) displayed significantly higher RMSD values than the WT (0.30±0.03 nm) (****: p < 0.0001). However, L99V (0.03±0.02 nm) showed non-significant (ns) differences.

Fig. 5. Radius of gyration (Rg) analysis of WT and mutant Cyt-c protein structures over a 50 ns (50000 ps) of MD simulation. (A) Variants of Rg values for WT and T20I, V21G, H27Y, N32H and L33V mutant proteins. (B) Rg values of mutants T20I (1.29±0.02 nm), V21G (1.26±0.01 nm), H27Y (1.28±0.01 nm), N32H (1.27±0.02 nm) and L33V (1.27±0.01 nm) were significantly higher than the WT (1.25±0.01 nm) (****: p < 0.0001). (C) Rg profiles for G42S, Y49H, A52T, A52V and R92G (D) where G42S (1.25±0.01 nm), A52V (1.28±0.01 nm) and R92G (1.26±0.01 nm) were significant (****: p < 0.0001) compared to WT (1.25±0.01 nm). Y49H (1.25±0.01 nm) showed non-significant results while A52T (1.26±0.01 nm) displayed less significant differences (*: p = 0.03). (E) Rg variations over simulation where (F) A97D (1.27±0.01 nm), Y98H (1.25±0.02 nm), L99V (1.26±0.01 nm) and T103I (1.27±0.01 nm) displayed significantly higher Rg values than the WT (1.25±0.01 nm) (****: p < 0.0001).

Fig. 6. Root mean square fluctuation (RMSF) profiles of Cyt-c WT and mutant protein structures. (A) Mutants T20I (0.16±0.07 nm), V21G (0.18±0.06 nm), H27Y (0.19±0.07 nm), N32H (0.20±0.08 nm) and L33V (0.20±0.08 nm) displayed significantly higher RMSF values compared to the WT (0.16±0.07 nm) (****: p < 0.0001). (B) G42S (0.21±0.07 nm), Y49H (0.18±0.07 nm), A52T (0.15±0.05 nm), A52V (0.18±0.06 nm) and R92G (0.20±0.10 nm) exhibited significantly higher RMSF compared to the WT (****: p < 0.0001). (C) WT showed significantly lower RMSF values compared to A97D (0.23±0.09 nm), Y98H (0.20±0.07 nm), L99V (0.17±0.07 nm) and T103I (0.27±0.13 nm) (****: p < 0.0001). (D) Impact of variants on Ω-loop dynamics. T20I and V21G increased flexibility in proximal (residues 13-40) and central (residues 41-58) loops, respectively, while H27Y enhanced flexibility across all Ω-loops, suggesting altered protein dynamics. N32H and L33V variants notably elevated RMSF in proximal (residues 13-40) and central (residues 41-58) loops, with L33V affecting the distal (residues 72-86) loop as well. (E) G42S variant increased flexibility in all loops. Y49H showed increased RMSF in proximal and central regions; A52T minimally impacted the proximal loop but reduced RMSF in central and distal loops. A52V and R92G variants increased flexibility in proximal and central regions, with R92G exhibiting significant fluctuations across all loops and the N-terminus. (F) A97D, Y98H and T103I variants significantly increased RMSF values in all loop regions.

Fig. 7. Variations of H-bonds in Cyt-c WT and mutant structures. (A) WT structure (68.46±5.05) showed a significantly higher number of H-bonds than that of T20I (62.03±4.51), V21G (65.51±4.76) and N32H (66.66±4.60) (****: $p < 0.0001$). Differences of WT with H27Y and L33V were non-significant. (B) G42S (59.58±4.86), A52T (64.17±5.08). A53V (67.37±4.84) and R92G (60.50±4.74) showed significantly lower (****: $p < 0.0001$) compared to the WT while the difference with Y49H was non-significant. (C) A97D (64.06±4.73), Y98H (65.33±6.01), L99V (66.37±4.51) and T103I (62.15±4.70) displayed significantly lesser (****: $p < 0.0001$) than WT. Variation of H-bonds for WT and mutants (D) T20I, V21G, H27Y, N32H, L33V, (E) G42S, Y49H, A52T, A52V, R92G, (F) A97D, Y98H, L99V and T103I over the MD simulation of 50 ns (50000 ps).

Fig. 8. Distances between H-bonded residue: Tyr68-Asn53, Tyr68-Thr79 and Tyr68-Met81. Tyr68 forms a hydrogen bond network with the conserved residues Asn53 and Thr79 and the axial ligand Met81, which is crucial for maintaining the structural integrity and functional properties of Cyt-c. The data highlights the loss of H-bonds and increased distances in mutants such as T20I, V21G, N32H, G42S, Y49H, A52T, A52V, Y98H, L99V, and T103I, which may destabilize the heme configuration by altering the spatial arrangement of Asn53, Thr79, and Met81. This destabilization potentially increases the accessibility of the heme catalytic site to small molecules like H_2O_2, enhancing peroxidase activity and facilitating the initiation of apoptosis.

Figures

Fig. 1. Overall structure of Cyt-c and information on the position and conservation of the clinically important variants. (A) The Cyt-c variants, T20I, V21G, H27Y, N32H and L33V are located in the proximal loop, G42S, Y49H, A52T and A52V in the central loop and R92G, A97D, Y98H, L99V and T103I in the protein C-terminus. (B) Multiple sequence alignment analysis of Cyt-c with orthologues from different species including *Homo sapiens* (human), *Macaca mulatta* (rhesus monkey), *Mus musculus* (mouse), *Canis lupus familiaris* (dog), *Oryctolagus cuniculus* (European rabbit), *Sus scrofa* (wild boar/pig), *Gallus gallus* (chicken), *Xenopus tropicalis* (western clawed frog) and *Danio rerio* (zebrafish) belonging to diverse taxonomic groups. The alignment showed that all analyzed variants (T20I, V21G, H27Y, N32H, L33V, G42S, Y49H, A52T, A52V, R92G, A97D, Y98H, L99V and T103I) are at evolutionary conserved residues and are likely deleterious. (C) SIFT analysis shows that all variants affect protein function (score < 0.05) and SuSPect associates the variants with the disease (score > 50).

Fig. 2. Mapping the currently known clinically important variants on the structure of Cyt-c, and Ramachandran plot analysis. The protein structure is depicted as a grey ribbon, with heme in green. (A) Human Cyt-c (PDB ID: 1j3s) structure showing three Ω-loop regions: proximal (residues 13-40, magenta), central (residues 41-58, cyan) and distal (residues 72-86, red) Ω-loops, which make the large part of Cyt-c. These loops are important for both structural and functional reasons, including protein folding and conformational transition. Amino acids investigated for variants are highlighted as follows: (B) T20 (magenta), V21 (blue), H27 (blue), N32 (magenta), L33 (blue); (C) G42 (magenta), Y49 (blue), A52 (blue), R92 (blue); (D) A97 (magenta), Y98 (blue), L99 (red), T103 (blue). Note that the residue numbering is offset by +1 relative to the PDB entry 1j3s. (E) In the Ramachandran plot, G42 is an exception, positioned in a typically disallowed region due to its small size and high flexibility. The G42S variant introduces steric hindrance due to larger and less flexible serine in a disallowed region.

Fig. 3. Visualization of gene and protein interactions. (A) Interactions of the *CYCS* gene with other genes, analyzed using the GeneMANIA tool. (B) The protein-protein interaction network of Cyt-c analyzed using the STRING tool. Variants in these networks may affect the functions of interacting genes or proteins.

Fig. 4. Root Mean Square Deviation (RMSD) profiles of WT and mutant Cyt-c protein structures over a 50 ns (50000 ps) of MD simulation. (A) Fluctuations of RMSD values for WT and T20I, V21G, H27Y, N32H and L33V mutant proteins. (B) RMSD values of mutants T20I (0.38 ± 0.03 nm), V21G (0.36 ± 0.04 nm), H27Y (0.39 ± 0.0 nm), N32H (0.42 ± 0.05 nm) and L33V (0.33 ± 0.02 nm) were significantly higher than the WT (0.30 ± 0.03 nm) (****: $p < 0.0001$). (C) The mutant proteins showed higher RMSD values (D) which were significant (****: $p < 0.0001$) compared to WT (0.30 ± 0.03 nm). The mutant displayed mean$\pm$SD as G42S: 0.42 ± 0.05 nm; Y49H: 0.34 ± 0.04 nm; A52T: 0.37 ± 0.03 nm; A52V: 0.41 ± 0.04 nm; and R92G 0.45 ± 0.06 nm. (E) Variations of RMSD values over simulation where (F) A97D (0.43 ± 0.05 nm), Y98H (0.43 ± 0.03 nm) and T103I (0.45 ± 0.05 nm) displayed significantly higher RMSD values than the WT (0.30 ± 0.03 nm) (****: $p < 0.0001$). However, L99V (0.03 ± 0.02 nm) showed non-significant (ns) differences.

Fig. 5. Radius of gyration (Rg) analysis of WT and mutant Cyt-c protein structures over a 50 ns (50000 ps) of MD simulation. (A) Variants of Rg values for WT and T20I, V21G, H27Y, N32H and L33V mutant proteins. (B) Rg values of mutants T20I (1.29±0.02 nm), V21G (1.26±0.01 nm), H27Y (1.28±0.01 nm), N32H (1.27±0.02 nm) and L33V (1.27±0.01 nm) were significantly higher than the WT (1.25±0.01 nm) (****: $p < 0.0001$). (C) Rg profiles for G42S, Y49H, A52T, A52V and R92G (D) where G42S (1.25±0.01 nm), A52V (1.28±0.01 nm) and R92G (1.26±0.01 nm) were significant (****: $p < 0.0001$) compared to WT (1.25±0.01 nm). Y49H (1.25±0.01 nm) showed non-significant results while A52T (1.26±0.01 nm) displayed less significant differences (*: $p = 0.03$). (E) Rg variations over simulation where (F) A97D (1.27±0.01 nm), Y98H (1.25±0.02 nm), L99V (1.26±0.01 nm) and T103I (1.27±0.01 nm) displayed significantly higher Rg values than the WT (1.25±0.01 nm) (****: $p < 0.0001$).

Fig. 6. Root mean square fluctuation (RMSF) profiles of Cyt-c WT and mutant protein structures. (A) Mutants T20I (0.16±0.07 nm), V21G (0.18±0.06 nm), H27Y (0.19±0.07 nm), N32H (0.20±0.08 nm) and L33V (0.20±0.08 nm) displayed significantly higher RMSF values compared to the WT (0.16±0.07 nm) (****: p < 0.0001). (B) G42S (0.21±0.07 nm), Y49H (0.18±0.07 nm), A52T (0.15±0.05 nm), A52V (0.18±0.06 nm) and R92G (0.20±0.10 nm) exhibited significantly higher RMSF compared to the WT (****: p < 0.0001). (C) WT showed significantly lower RMSF values compared to A97D (0.23±0.09 nm), Y98H (0.20±0.07 nm), L99V (0.17±0.07 nm) and T103I (0.27±0.13 nm) (****: p < 0.0001). (D) Impact of variants on Ω-loop dynamics. T20I and V21G increased flexibility in proximal (residues 13-40) and central (residues 41-58) loops, respectively, while H27Y enhanced flexibility across all Ω-loops, suggesting altered protein dynamics. N32H and L33V variants notably elevated RMSF in proximal (residues 13-40) and central (residues 41-58) loops, with L33V affecting the distal (residues 72-86) loop as well. (E) G42S variant increased flexibility in all loops. Y49H showed increased RMSF in proximal and central regions;

A52T minimally impacted the proximal loop but reduced RMSF in central and distal loops. A52V and R92G variants increased flexibility in proximal and central regions, with R92G exhibiting significant fluctuations across all loops and the N-terminus. (F) A97D, Y98H and T103I variants significantly increased RMSF values in all loop regions.

Fig. 7. Variations of H-bonds in Cyt-c WT and mutant structures. (A) WT structure (68.46±5.05) showed a significantly higher number of H-bonds than that of T20I (62.03±4.51), V21G (65.51±4.76) and N32H (66.66±4.60) (****: p < 0.0001). Differences of WT with H27Y and L33V were non-significant. (B) G42S (59.58±4.86), A52T (64.17±5.08). A53V (67.37±4.84) and R92G (60.50±4.74) showed significantly lower (****: p < 0.0001) compared to the WT while the difference with Y49H was non-significant. (C) A97D (64.06±4.73), Y98H (65.33±6.01), L99V (66.37±4.51) and T103I (62.15±4.70) displayed significantly lesser (****: p < 0.0001) than WT. Variation of H-bonds for WT and mutants (D) T20I, V21G, H27Y, N32H, L33V, (E) G42S, Y49H, A52T, A52V, R92G, (F) A97D, Y98H, L99V and T103I over the MD simulation of 50 ns (50000 ps).

Fig. 8. Distances between H-bonded residue: Tyr68-Asn53, Tyr68-Thr79 and Tyr68-Met81. Tyr68 forms a hydrogen bond network with the conserved residues Asn53 and Thr79 and the axial ligand Met81, which is crucial for maintaining the structural integrity and functional properties of Cyt-c. The data highlights the loss of H-bonds and increased distances in mutants such as T20I, V21G, N32H, G42S, Y49H, A52T, A52V, Y98H, L99V, and T103I, which may destabilize the heme configuration by altering the spatial arrangement of Asn53, Thr79, and Met81. This destabilization potentially increases the accessibility of the heme catalytic site to small molecules like H_2O_2, enhancing peroxidase activity and facilitating the initiation of apoptosis.