Unraveling the Role of Ctla-4 in Intestinal Immune Homeostasis: Insights from a novel Zebrafish Model of Inflammatory Bowel Disease

Lulu Qin[a], Chongbin Hu[a], Qiong Zhao[a], Yong Wang[a], Dongdong Fan[a], Aifu Lin[a], Lixin Xiang[a*], Ye Chen[a,c*], Jianzhong Shao[a,b*]

Affiliations:

[a]College of Life Sciences, Key Laboratory for Cell and Gene Engineering of Zhejiang Province, Zhejiang University, Hangzhou, China.

[b]Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, Qingdao, China

[c]Department of Genetic and Metabolic Disease, the Children's Hospital, Zhejiang University School of Medicine, National Clinical Research Center for Child Health, Hangzhou 310052, China.

[*]Corresponding authors: shaojz@zju.edu.cn, yechency@zju.edu.cn, and xianglx@zju.edu.cn

Address correspondence and reprint requests to Prof. Jianzhong Shao, Prof. Ye Chen, and Assoc. Prof. Lixin Xiang, College of Life Sciences, Zhejiang University, 866 YuHangTang Road, Hangzhou 310058, China. Tel.: +86 (571) 88206582; Fax: +86 (571) 88206582.

E-mail addresses: shaojz@zju.edu.cn (J.-z.S.), yechency@zju.edu.cn (Y.C), and xianglx@zju.edu.cn (L.-x.X.)

Abstract

Inflammatory bowel disease (IBD) is a chronic and relapsing immune-mediated disorder characterized by intestinal inflammation and epithelial injury. The underlying causes of IBD are not fully understood, but genetic factors have implicated in genome-wide association studies, including CTLA-4, an essential negative regulator of T cell activation. However, establishing a direct link between CTLA-4 and IBD has been challenging due to the early lethality of CTLA-4 knockout mice. In this study, we identified zebrafish Ctla-4 homolog and investigated its role in maintaining intestinal immune homeostasis by generating a Ctla-4-deficient (*ctla-4$^{-/-}$*) zebrafish line. These mutant zebrafish exhibited reduced weight, along with impaired epithelial barrier integrity and lymphocytic infiltration in their intestines. Transcriptomics analysis revealed upregulation of inflammation-related genes, disturbing immune system homeostasis. Moreover, single-cell RNA-sequencing analysis indicated increased Th2 cells and interleukin 13 expression, along with decreased innate lymphoid cells and upregulated proinflammatory cytokines. Additionally, Ctla-4-deficient zebrafish exhibited reduced diversity and an altered composition of the intestinal microbiota. All these phenotypes closely resemble those found in mammalian IBD. Lastly, supplementation with Ctla-4-Ig successfully alleviated intestinal inflammation in these mutants. Altogether, our findings demonstrate the pivotal role of Ctla-4 in maintaining intestinal homeostasis. Additionally, they offer substantial evidence linking CTLA-4 to IBD and establish a novel zebrafish model for investigating both the pathogenesis and potential treatments.

Introduction

Inflammatory bowel disease (IBD), including Crohn's disease and ulcerative colitis, refers to a group of chronic relapsing inflammation disorders affecting the gastrointestinal tract, that have been increasing in prevalence worldwide [1]. The precise etiology of IBD has yet to be fully elucidated. Conventional epidemiological studies have indicated that IBD tends to run in families and is linked to genetic factors [2, 3]. However, research also suggested that susceptibility gene patterns differ significantly among various geographic populations. Current evidence points towards a complicated interaction involving host genetics, disrupted intestinal microbiota, environmental triggers, and abnormal immune responses [4-6]. Advances in genomic sequencing techniques have allowed for the identification of genetic variants associated with an increased risk of developing IBD. Among these, mutations in immune-related genes have received particular attention. Research on humans with Crohn's disease and mouse models of IBD has shown that genetically susceptible individuals exhibit defects in intracellular pattern-recognition receptors (PRRS), such as toll-like receptors (TLR) and nucleotide-binding oligomerization domain (NOD)-like receptors (NLRs), which are responsible for initiating innate immune responses to eliminate harmful bacteria [7, 8]. Genetic variations in the tumor necrosis factor ligand superfamily member 15 (TNFSF15) and interleukin 23 receptor (IL23R) genes, both involved in suppressing

inflammation, have been associated with an increased risk of developing Crohn's

disease [9, 10].

Cytotoxic T lymphocyte antigen-4 (CTLA-4), also known as CD152, is one of the

most well-established immune checkpoint molecules expressed predominantly on T

cells [11-13]. It primarily regulates the early stages of T-cell activation by attenuating

downstream signaling of the T cell receptor (TCR) [14-16]. Specifically, CTLA-4 has

a much higher affinity for CD80 and CD86 ligands compared to the co-stimulatory

receptor CD28 [17, 18]. By outcompeting CD28 for ligand binding, CTLA-4 provides

an inhibitory signal that impacts immunological synapse formation and inhibits T-cell

proliferation and activation [16, 19]. The immunomodulatory role of CTLA-4 in

maintaining immune homeostasis is highlighted through CTLA-4 knockout studies.

Germline CTLA-4-deficient is lethal for mice within 3 to 4 weeks due to massive T

lymphocyte proliferation and the release of inflammatory cytokines [20-22]. Compared

to wild-type T lymphocytes, CTLA-4-deficient T lymphocytes exhibit accelerated

development of Th2 cells, leading to significantly enhanced secretion of IL-4 and IL-5

[23, 24]. Additionally, conditional deletion of CTLA-4 in adult mice results in rapid

immune activation, multiorgan lymphocyte infiltration, and autoantibody production

[25]. Moreover, a selective deficiency of CTLA-4 in Treg cells is sufficient to induce

lymphoproliferation and autoimmune diseases in mice [26]. Similarly, associations

between polymorphic alleles of CTLA-4 and IBD in humans have been reported in

multiple studies [27-29]. Moreover, CTLA-4 is an intriguing target for novel immune

checkpoint blockade therapies in cancer treatment, while intestinal inflammation is a

common side effect in these clinical trials [30, 31]. Establishing a direct causal relationship between CTLA-4 and IBD has been challenging due to difficulties in finding appropriate models. The early lethality observed in CTLA-4-deficient mice added another layer of complexity to this issue. Zebrafish is a powerful model system for immunological and biomedical research, due to its versatility and high degree of conservation in innate and adaptive immunities [32, 33]. In our current study, we identified the Ctla-4 homology in zebrafish and successfully developed an adult vertebrate model with homozygous knockout of the *ctla-4* gene for the first time. These *ctla-4*-deficient (*ctla-4*$^{-/-}$) zebrafish survive but exhibit attenuated growth and weight loss. Notably, *ctla-4* deficiency leads to an IBD-like phenotype in zebrafish characterized by altered intestinal epithelial cells morphology, abnormal inflammatory response, defects in microbial stratification and composition. Mechanistically, Ctla-4 exerts its inhibitory function by competing with Cd28 for binding to Cd80/86. These findings establish the *ctla-4* knockout zebrafish as an innovative platform to elucidate CTLA-4 immunobiology, model human IBD, and develop novel therapeutic modalities.

Results

Identification of zebrafish Ctla-4

Through a homology search in the NCBI database, we identified the *ctla-4* gene (XM_005167519.4) on zebrafish chromosome 9, which exhibits an exon organization comparable to that of the human CTLA-4 gene (Fig. S1A-C). Zebrafish Ctla-4 is predicted to be a type I transmembrane protein with a molecular weight of

approximately 33 kDa, featuring structural characteristics of the immunoglobulin superfamily, which include an N-terminal signal peptide, a single IgV-like extracellular domain, a transmembrane region, and a cytoplasmic tail (Fig. 1A). Multiple amino acid sequence alignments revealed that Ctla-4 contains a ^{113}LFPPPY^{118} motif within the ectodomain and a tyrosine-based ^{206}YVKF^{209} motif in the distal C-terminal region (Fig. 1A). These motifs closely resemble the MYPPPY and YVKM motifs found in mammalian CTLA-4 homologs, which are essential for binding to CD80/CD86 ligands, as well as molecular internalization and signaling inhibition [14, 34, 35]. The IgV-like domain of Ctla-4 was characterized by two-layer β-sandwich and was conserved between zebrafish and humans (Fig. 1B). In contrast, zebrafish Cd28 features a SYPPPF motif in its extracellular region and a FYIQ motif in its intracellular tail, distinguishing it from Ctla-4. Additionally, zebrafish Ctla-4, similar to its counterparts in other species, carries a conserved extracellular ^{123}GNGT^{126} motif, which is absent in zebrafish Cd28 [36]. This structural distinction further differentiates Ctla-4 from Cd28 (Fig.1A, Fig. S1D). Consistent with this, phylogenetic analysis showed that Ctla-4 clusters with other known CTLA-4 homologs from different species with high bootstrap probability, whereas zebrafish Cd28 groups separately with other CD28s (Fig. S1C). Structurally, Ctla-4 exists as a dimer, and unlike the intracellular localization of CTLA-4 in mammals, Ctla-4 is found on the cell membrane (Fig. 1C and D). By analyzing the splenic scRNA-seq dataset we recently established [37], Ctla-4 was primarily expressed on the T cells, including the Cd4$^+$ T and Cd8$^+$ T cells (Fig. 1E). This result was verified by immunofluorescence assays on the splenic leukocytes (Fig. 1F).

Ctla-4 deficiency induces inflammatory bowel disease (IBD)-like phenotype

To further investigate the function of Ctla-4, we generated a *ctla-4⁻/⁻* zebrafish line with a 14-base deletion in the second exon of the *ctla-4* gene (Fig. 2A-C). The zebrafish appeared grossly normal in appearance; however, the body weight and size were significantly reduced compared with those of wild-type zebrafish (Fig. 2D and E). Anatomically, the *ctla-4⁻/⁻* zebrafish were featured by intestine shortening and splenomegaly, suggesting the occurrence of chronic inflammation in the intestines (Fig. 2F and G). For clarification, we first performed histological analysis on the anterior, mid, and posterior intestine segments using H&E staining. Compared to the wild-type zebrafish, the *ctla-4⁻/⁻* fish exhibited significant epithelial hyperplasia in the anterior intestine segment, accompanied with a small amount of mucosal inflammatory cell infiltration (Fig. 2H). Moreover, a noteworthy goblet cell loss, reduction of normal mucins and the accumulation of acidic mucins were also observed in *ctla-4⁻/⁻* anterior intestine, as detected through Alcian Blue and Periodic Acid-Schiff (AB-PAS) or PAS staining (Fig. 2I and J, Fig. S3A). A small amount of lymphocytic infiltration and mild epithelial damage occurred in the mid intestine segment of *ctla-4⁻/⁻* zebrafish (Fig. 2H). In posterior intestine of *ctla-4⁻/⁻* fish, the intestinal villi were markedly shortened, the epithelial barrier showed severely disrupted, and the intestinal wall became thinner, wherein the mucosal and transmural inflammatory cells were significantly infiltrated (Fig. 2H). Notably, the intestinal lumens in all three intestinal segments were enlarged in the ctla-4⁻/⁻ zebrafish, and the ratio between the length of the intestinal villi and the intestinal ring radius was higher in the ctla-4⁻/⁻ zebrafish intestines compared to those

in wild-type zebrafish (Figure S3B). In addition, the ultrastructure analysis revealed

that the epithelial cells of posterior intestine in *ctla-4$^{-/-}$* zebrafish exhibited alteration in

tight junction, the loss of adhesion junctions and desmosomes, and disruption of

microvilli (Fig. 2K). All these results strongly indicate that Ctla-4 plays a crucial role

in preserving intestinal homeostasis in zebrafish. The intestinal phenotype resulting

from Ctla-4 deficiency was similar to IBD in mammals.

Molecular mechanisms of Ctla-4 deficiency-induced IBD-like phenotype

To explore the potential molecular mechanisms of Ctla-4 deficiency-induced IBD-

like phenotype, we performed transcriptome profiling analysis of intestines from wild-

type and *ctla-4$^{-/-}$* zebrafish. We identified a total of 1,140 differentially expressed genes

(DEGs), among which 714 genes were up-regulated, and 426 genes were down-

regulated in *ctla-4$^{-/-}$* zebrafish (Fig. 3A and B). GO enrichment analysis showed that

DEGs or up-regulated genes in the top 10 enriched biological processes were associated

with the immune response and inflammatory response (Fig. 3C and D). Moreover, the

KEGG enrichment analyses indicated that the up-regulated DEGs are primarily

involved in the process of cytokine-cytokine receptor interaction and cell adhesion

molecules, which are also related to inflammation (Fig. S4A); however, the down-

regulated DEGs were significantly enriched in the process of metabolism (Fig. S4B).

The intestines of *ctla-4$^{-/-}$* zebrafish showed significant upregulation of *il1b*, *tnfa*,

myeloid-specific peroxidase (*mpx*), matrix metallopeptidase 9 (*mmp9*), chemokine (C-

X-C motif) ligand 8a (*cxcl8a*), and *il13*. In contrast, *il10*, a potent anti-inflammatory

cytokine, was markedly down-regulated in Ctla-4-deficient intestines (Fig. 3E). The

transcriptional change of these genes was confirmed by RT-qPCR (Fig. 3F). By

constructing the protein-protein interaction (PPI) network, we found that *il1b* was a

major cytokine that played a hub role in promoting the bowel inflammation of *ctla-4⁻/⁻*

zebrafish (Fig. 3G). Moreover, Gene set enrichment analysis (GSEA) showed that

genes involved in the lymphocyte chemotaxis, positive regulation of ERK1 and ERK2

cascade, Calcium and MAPK signaling pathways were positively enriched in *ctla-4⁻/⁻*

zebrafish intestines, implying a sensitized or activated state of lymphocytes due to the

absence of Ctla-4 (Fig. S4C and D). Notably, biological processes related to neutrophil

activation and chemotaxis were significantly enriched (Fig. 3C and D). Studies have

shown that neutrophils can induce histopathological effects through releasing matrix

metalloproteinases (MMPs), neutrophil elastase, and myeloperoxidase (MPO) [38]. To

confirm the association between neutrophils and Ctla-4-deficient intestinal

inflammation, the MPO level was examined. As a support, MPO activity was markedly

increased in the intestines and peripheral blood of *ctla-4⁻/⁻* zebrafish (Fig. 3H). Besides,

a number of biological markers or susceptibility genes of IBD observed in mammals,

including c-reactive protein 6 (*crp6*), *crp7*, MMPs, haptoglobin, *il23r*, insulin-like

growth factor binding protein 1 a (*igfbp1a*), cAMP responsive element modulator b

(*cremb*) and lymphocyte specific protein 1 b (*lsp1b*), were highly expressed in the *ctla-*

4⁻/⁻ zebrafish (Fig. 3I and J) [9, 39, 40], suggesting the presence of a conserved

molecular network underlying IBD pathogenesis between *ctla-4⁻/⁻* zebrafish and

mammalian models.

Cellular mechanisms of Ctla-4 deficiency-induced IBD-like phenotype

To investigate the cellular mechanisms underlying the IBD-like phenotype

induced by Ctla-4 deficiency, we performed scRNA-seq analysis on intestines from

wild-type and *ctla-4*[-/-] zebrafish using the 10× Genomics Chromium platform. We

obtained nine discrete clusters from 7,539 cells of wild-type and *ctla-4*[-/-] zebrafish (Fig.

4A). These clusters of cells were classified as enterocytes, enteroendocrine cells,

smooth muscle cells, neutrophils, macrophages, B cells, and a group of T/NK/ILC-like

cells based on their co-expression of lineage marker genes (Fig. 4B and C, Fig. S6A

and B). Due to severe epithelial disruption and inflammatory cell infiltration in *ctla-4*[-

/-] zebrafish intestines, we focused on the pathological process and immune reactions in

enterocytes and immune cell populations. KEGG analysis showed that apoptotic

pathway was highly enriched in enterocytes of *ctla4*[-/-] zebrafish, suggesting that

aberrant apoptosis contributes to the epithelial injury (Fig. S6C). Subsequently, we

conducted a TUNEL assay to detect apoptosis in the posterior intestines from both wild-

type and *ctla4*[-/-] zebrafish. The results showed a higher number of apoptotic cells in the

intestines of *ctla4*[-/-] zebrafish (Fig. S6D). Additionally, genes functionally involved in

the formation of tight and adhesion junctions, such as *oclna*, *cdh1*, *pcdh1b* and *cldn15a*,

were significantly down-regulated in enterocytes of *ctla-4*[-/-] zebrafish (Fig. 4D),

consistent with the pathological observation under electron microscope. Furthermore,

inflammation-related genes and pathways were significantly up-regulated and enriched

in neutrophils, B cells, and macrophages of *ctla-4*[-/-] zebrafish, suggesting active

inflammatory responses (Fig. 4E-G, Fig. S6E). By sub-clustering analysis, six

subpopulations were classified from T/NK/ILC-like cell groups based on their

expression with a set of marker genes. These subpopulations include Cd8$^+$ T cells,

ILC3-like cells, maturing Ccr7^{high} T cells, NKT-like cells, and two groups of Th2 cells

(Fig. 5A-C, Fig. S7A). The abundances of NKT-like and two subsets of Th2 cells were

significantly increased in the intestines of *ctla-4*$^{-/-}$ zebrafish (Fig. 5D-F). These findings

were further validated by RT-qPCR detection of their corresponding marker genes (Fig.

S7B and C). These cells exhibited high expression levels of *il13* (Fig. 5G and H).

Specifically, the second subset of Th2 cells was seldom observed in the intestine of

wild-type zebrafish, indicating their unique role in the pathogenesis of IBD-like

phenotype in *ctla-4*$^{-/-}$ zebrafish (Fig. 5D-F). KEGG analysis of up-regulated genes from

ctla-4$^{-/-}$ NKT-like and Th2 cells indicated that Ctla-4 deficiency is positively associated

with the inflammatory cytokine-cytokine receptor interaction, PPAR, calcium and

MAPK signaling pathways, cellular adhesion and mucosal immune responses (Fig. 5I

and J, Fig. S7D). Although the abundance of Cd8$^+$ T cells was not significantly changed

in Ctla-4-deficient intestines, the inflammatory genes and pathways were up-regulated

and enriched in the subset of T cells (Fig. S7E and F). Notably, the proportion of ILC3-

like cells was decreased, and they highly expressed *il17a/f1* and *il17a/f3* in the Ctla-4-

deficient intestines (Fig. 5D-F, and K). Investigations have consistently reported a

substantial decline in the population of ILC3s within the inflamed intestines and IL-

17A-secreting ILC3s play a significant role in the development of intestinal

inflammation [41-46]. Thus, the reduced ILC3-like cells and increased expression of

il17a/f1 and *il17a/f3* may be responsible for intestinal inflammation induced by Ctla-4

deficiency.

Decreased microbiota diversity in *ctla-4^{-/-}* zebrafish intestines

The intestinal microbiota plays a crucial role in host functions such as nutrient acquisition, metabolism, epithelial cell development and immunity. Notably, lower microbiota diversity has consistently been observed in patients with IBD phenotype [47, 48], making it a valuable indicator of host health. Therefore, we further analyzed whether microbes are involved in the Ctla-4-deficiency induced intestinal inflammation. The results revealed a significantly higher number of amplicon sequence variants (ASVs) in wild-type zebrafish intestines, with 730 ASVs unique to the wild-type group and 276 ASVs exclusively found in *ctla-4^{-/-}* group (Fig. 6A). Furthermore, the Shannon index and the Simpson index indicated a decreased microbial diversity in *ctla-4^{-/-}* zebrafish intestines (Fig. 6B) and the Principal Coordinate Analysis (PCoA) using Bray Curtis distance revealed a significant separation in the microbial composition between *ctla-4^{-/-}* group and the wild-type group (Fig. 6C). To gain insights into the microbial community composition, we analyzed the identified microbial populations at the class and family level. Alphaproteobacteria and Gammaproteobacteria were found to be the most prevalent classes. Relative to wild-type group, Ctla-4 deficiency resulted in a significant reduction in Alphaproteobacteria abundance. However, the Gammaproteobacteria, one of the main classes of Gamma-negative pathogenic bacteria expanded under inflammation conditions, was increased, although the change did not reach statistical significance (Fig. 6D and E) [49]. In addition, we observed a decreased relative abundance of short-chain fatty acids (SCFAs)-producing Bacilli and Verrucomicrobiae, the latter of which contributes to glucose homeostasis and intestinal

health (Fig. 6F and G) [50, 51]. Notably, the family-level analysis revealed a notable

enrichment of Enterobacteriaceae, overgrowing under host inflammatory conditions,

and the Shewanellaceae, serving as the most important secondary or opportunistic

pathogens, in *ctla-4*[-/-] zebrafish (Fig. 6H and I). To identify differentially abundant

bacterial taxa between the wild-type and *ctla-4*[-/-] zebrafish, we conducted Linear

discriminant analysis (LDA) effect size (LEfSe). The results showed that several

potentially opportunistic pathogens, including Gammaproteobacteria, Enterobacterales,

and Aeromonadales were found to be overrepresented in *ctla-4*[-/-] zebrafish (Fig. 6J). In

contrast, Actinobacteriota, Cetobacterium, and Planctomycetota (Planctomycetes) were

more abundant in the wild-type zebrafish. These findings strongly indicated an

association between Ctla-4 deficiency-induced gut inflammation and dysbiosis, as

characterized by decreased microbial diversity, loss of potentially beneficial bacteria,

and expansion of pathobionts.

Inhibitory role of Ctla-4 in T cell activation

Given that Ctla-4 is primarily expressed on T cells (Fig. 1E-F), its absence has

been shown to induce intestinal immune dysregulation, indicating a crucial role of this

molecule as a conserved immune checkpoint in T cell inhibition. Mechanistically, Ctla-

4 may inhibit T cell activation by obstructing the Cd80/86-Cd28 costimulatory pathway,

a mechanism conserved in mammalian species. To elucidate the regulatory role of Ctla-

4 in costimulatory signal-dependent T cell activation, we conducted a series of blockade

and activation assays using anti-Ctla-4 antibody, recombinant soluble Ctla-4-Ig (sCtla-

4-Ig), sCd28-Ig, and sCd80/86 proteins in a PHA-stimulating and mixed lymphocyte

reaction (MLR) model. In this system, sCtla-4-Ig and sCd28-Ig served as antagonists

to block membrane-bound Cd80/86, while sCd80/86 acted as an agonist for membrane-

bound Cd28 (Fig. S8A-C). As expected, the proliferation of lymphocytes from *ctla-4[-/-]*

zebrafish was more pronounced compared to wild-type controls, and the addition of

sCtla-4-Ig effectively suppressed this proliferation (Fig. 7A and B). These findings

indicate that the absence of Ctla-4 leads to enhanced lymphocyte activation, which can

be counteracted by sCtla-4 administration, underscoring the inhibitory function of Ctla-

4 in T cell regulation. Consistent with these results, blockade of Ctla-4 using anti-Ctla-

4 Ab significantly promoted the proliferation of lymphocytes from wild-type zebrafish

(Fig. 7C). Furthermore, sCd28-Ig administration inhibited the proliferation of

lymphocytes from *ctla-4[-/-]* zebrafish (Fig. 7D), whereas sCd80/86 promoted the

expansion of Ctla-4-deficient lymphocytes (Fig. 7E). Based on these results, we

concluded that the presence of Ctla-4 obstructs the Cd80/86-Cd28 mediated

costimulatory signaling, consequently impeding cell proliferation. To further

investigate the molecular interactions between Cd28, Ctla-4, and Cd80/86, we

employed AlphaFold2 to predict the structures of Cd80/86-Cd28 and Cd80/86-Ctla-4

complexes. A total of 25 models were generated for each complex and subsequently

aligned with Cd80/86. The predictions indicated that both Cd28 and Ctla-4 form a well-

defined interface with Cd80/86, utilizing the same binding site (Fig. S8D and E). This

well-defined interface was corroborated by lower predicted aligned error (PAE) scores

for each model, as marked by the red dashed square (Fig. 7F and G). Subsequently, co-

immunoprecipitation (Co-IP) assays were conducted to provide compelling evidence

for the molecular interactions between Cd80/86 and Cd28 or Ctla-4. Flow cytometry

analysis further revealed dose-dependent associations between Cd80/86 and Cd28 or

Ctla-4 in HEK293T cells (Fig. 7H, S8F). Additionally, microscale thermophoresis

assays demonstrated that Ctla-4 exhibits a higher binding affinity for Cd80/86 than

Cd28, as evidenced by a lower equilibrium association constant value ($K_D = 0.50 \pm 0.25$

μM vs. $K_D = 2.64 \pm 0.45\ \mu M$) (Fig. 7I). These findings suggest that Ctla-4 exerts its

inhibitory function by competing with Cd28 for binding Cd80/86.

sCtla-4-Ig alleviates IBD-like phenotype

As described above, engagement of Cd80/86 by sCtla-4-Ig effectively suppressed

T cell activation *in vitro* (Fig. 7B), indicating that sCtla-4-Ig holds promise as a

potential intervention for IBD-like phenotype. This is supported by the observation that

Ctla-4-deficient zebrafish treated with sCtla-4-Ig exhibited obvious body weight

restoration compared to those treated with the IgG isotype control (Fig. 8A). To provide

further evidence, histological analysis was performed on the posterior intestine, which

is known to experience severe tissue destruction under Ctla-4 deficient conditions. As

expected, Ctla-4-Ig treatment resulted in a significant decrease in lymphocyte

infiltration and an improvement in the epithelial barrier (Fig. 8B). Moreover, Ctla-4-Ig

treatment significantly reduced the expression of pro-inflammatory genes, including

il13, *tnfa*, *mpx*, *mmp9*, and *cxcl8a*, as well as *igfbp1a*, *cremb*, and *lsp1a*, which are

susceptibility genes for IBD observed in mammals (Fig. 8C and D). These findings

demonstrate that the supplementation of Ctla-4-Ig alleviates intestinal inflammation in

Ctla-4-deficient zebrafish, highlighting its potential as a therapeutic intervention for

330 CTLA-4 deficiency-induced IBD in mammals.

Discussion

As an essential negative regulator of T cell activation, dysfunction of CTLA-4 was

implicated in various diseases in both humans and murine models [20, 21, 52].

Numerous previous studies have established the connection between CTLA-4 and

autoimmune thyroiditis, Graves' disease, myocarditis, pancreatitis, multiple sclerosis,

rheumatoid arthritis, and type I diabetes [53-59]. However, the involvement of CTLA-

4 in IBD has been understudied. Several investigations have reported that

haploinsufficiency resulting from mutations in CTLA-4 in humans is associated with

IBD, and genome-wide association studies (GWAS) have implicated CTLA-4 as a

susceptibility gene for IBD [60-62]. Nevertheless, the exact contributions and

mechanisms of CTLA-4 deficiency in the occurrence and pathology of IBD remain

incompletely understood, primarily due to the lack of animal models attributable to the

lethality of CTLA-4 knockout in mice. In this study, we identified the Ctla-4 homolog

in zebrafish, and discovered that defect in Ctla-4 did not have a severe lethal effect, but

did show a clear IBD-like phenotype. This makes zebrafish an attractive animal model

for investigating the molecular and cellular mechanisms underlying Ctla-4 mediated

IBD.

Multiple lines of histopathological evidence demonstrated the IBD-like phenotype

in Ctla-4-deficienct zebrafish. Key features include epithelial hyperplasia, disruption of

epithelial integrity, loss of goblet cells, increased acidic mucus production, intestinal

lumen enlargement, inflammatory cell infiltration, and elevated expression of pro-inflammatory cytokines in the inflamed intestines. These characteristics, such as epithelial hyperplasia, goblet cell depletion, inflammatory cell infiltration, and upregulated pro-inflammatory cytokine expression, closely resemble those observed in the dextran sodium sulfate (DSS)-induced IBD model in mice [63]. Similarly, mononuclear cell infiltration and significant upregulation of the il1β cytokine have been reported in the trinitrobenzenesulfonic acid (TNBS)-induced IBD model in adult zebrafish [64]. In zebrafish larvae, the TNBS-induced IBD-like phenotype also exhibits an enlarged intestinal lumen, although goblet cell numbers were increased [65]. Additionally, neutrophilic inflammation and acidic mucin accumulation have been observed in the DSS-induced enterocolitis model in zebrafish larvae [66]. In contrast, the soybean meal-induced enteritis (SBMIE) phenotype in zebrafish larvae shows no significant structural changes in intestinal architecture, despite an increased number of neutrophils and lymphocytes [67]. In summary, Ctla-4 deficiency induces IBD-like phenotypes analogous to those typically elicited by drugs in mice and zebrafish, making this model a valuable tool for comprehending the pathophysiological mechanisms underlying IBD.

A transcriptomics study was conducted to investigate the mechanisms of Ctla-4-deficiency induced IBD. RNA-seq analysis demonstrated a significant upregulation of important inflammatory cytokines, such as *il1b* and *tnfa* in the Ctla-4-deficient intestines. This is consistent with studies showing that IL-1β and TNF-α act as crucial mediators in mammalian IBD models by disrupting epithelial junctions and inducing

apoptosis of epithelial cells [68, 69]. Conversely, the key anti-inflammatory cytokines,

such as *il10*, were downregulated. These findings highlight an imbalance between pro-

inflammatory and anti-inflammatory cytokines in the intestines of Ctla-4-deficient fish.

Consistently, the inflammatory signaling pathways associated with these upregulated

cytokines, such as the ERK1/2 and MAPK pathways, were positively enriched in

inflamed intestines. Single-cell RNA-seq analysis revealed the upregulation and

enrichment of these inflammatory cytokines and pathways in neutrophils, macrophages,

and B cells of inflamed intestines, indicating their active involvement in inflammatory

responses and as major sources of inflammatory signals. Additionally, there was a

marked increase in the abundance of Th2 subset cells in the inflamed intestines, these

cells exhibited high expression of *il13* and were significantly enriched in inflammatory

signaling pathways, indicating their activated state. These findings align with previous

studies indicating that T cells in CTLA-4-deficient mice exhibit a bias toward Th2

differentiation [23, 24]. Furthermore, IL-13, a key effector Th2 cytokine, has been

implicated in the pathogenesis of ulcerative colitis in mammals, where it directly

contributes to epithelial cell damage by disrupting tight junctions, inducing apoptosis,

and impairing cellular restitution [70]. Therefore, upregulated Il13 from Th2 cells may

be a significant contributor to the occurrence of intestine inflammation in Ctla-4-

deficient zebrafish. Notably, the proportion of ILC3-like cells was downregulated in the

inflamed intestines, consistent with recent studies reporting a substantial decline of

ILC3 in IBD patients [41-43]. ILC3 is the most abundant type of ILCs in the intestines

and plays a protective role in IBD in mammals by promoting epithelial cell proliferation

and survival, as well as enhancing intestinal barrier function through the production of IL-22 [44, 46]. Thus, the marked decrease in ILC3-like cells may exacerbate intestinal inflammation and damage.

IBD is frequently associated with alterations in gut microbiota composition, characterized by reduced microbial diversity and an imbalance between beneficial and pathogenic bacteria. The common feature of these changes is the expansion of Proteobacteria, particularly members of the Enterobacteriaceae family [71, 72]. Similarly, Ctla-4-deficient zebrafish exhibited significant enrichment of Enterobacteriaceae, alongside a decline in beneficial bacteria like Cetobacterium and an increase in opportunistic pathogens such as γ-Proteobacteria and Aeromonadales. These findings indicate shared patterns in microbial flora changes during intestinal inflammation. Previous studies suggest that reduced microbial diversity in IBD results from the loss of normal anaerobic bacteria, such as Bacteroides, Eubacterium, and Lactobacillus [73]. Concurrently, inflammation-driven increases in intestinal lumen oxygenation and the availability of nitrate and host-derived electron acceptors facilitate anaerobic respiration and Enterobacteriaceae proliferation [74]. These observations highlight the intricate interplay between IBD pathogenesis, gut microbial alterations, and host immune homeostasis. The zebrafish IBD-like model induced by Ctla-4 deficiency offers new insights into this research area. For instance, abnormal activation of Th2 cells may lead to dysfunction in downstream B cells and mucosa-associated immunity, which are crucial for maintaining symbiotic bacterial homeostasis in the intestines [50]. This suggests a potential link between Th2 cell changes and the

observed alterations in the intestinal microbial community in Ctla-4-deficient zebrafish. Moreover, Ctla-4 deficiency alters the proportion and activation of ILC3 cells and damages the intestinal epithelium, potentially shaping the inflammatory milieu and further disrupting gut microbial homeostasis. Ctla-4 also regulates T cell activation by inhibiting the Cd80/86 co-stimulatory pathway. These findings suggest a regulatory interplay between Ctla-4, ILC3 cells, Cd80/86-primed T cells, and gut microbiota in Ctla-4 deficiency-induced IBD. Recently, gut microbiota exposure has been found to induce local IL-23 production, which upregulates CTLA-4 on ILC3s. This supports immune regulation by reducing CD80/86 co-stimulatory signaling and increasing PD-L1 bioavailability on myeloid cells. Impairment of this pathway manifests in a substantial imbalance of effector and regulatory T cell responses, exacerbating intestinal inflammation [75]. These findings bolster our hypothesis and provide valuable insights into the complex interactions between gut microbiota, ILC3-mediated immune responses, and Cd80/86 signaling in Ctla-4 deficiency-induced IBD.

In conclusion, our study demonstrates that Ctla-4 serves as a potential genetic determinant of the IBD-like phenotype in zebrafish, although further research is necessary to conclusively identify the causative variant responsible for this association. The development of this zebrafish model offers a valuable tool for elucidating the mechanisms underlying the disease's pathophysiology. Nevertheless, a deeper understanding of the intricate interactions among immune cells, intestinal epithelial cells, and the microbiome in IBD remains an area warranting further investigation.

Materials and Methods

Experimental fish

The AB strain zebrafish (*Danio rerio*) of both sexes, 4-6 months of age with body weights ranging from 0.3 to 0.8 g and lengths of 3-4 cm, were reared in the laboratory in recirculating water at 26-28 °C under standard conditions as previously described [76]. All animal experiments were performed in compliance with legal regulations and approved by the Research Ethics Committee of Zhejiang University. For sampling, wild-type and Ctla-4-deficient zebrafish of varying ages were kept in separate tanks and labeled with their respective dates of birth. Wild-type zebrafish aged 4-6 months and Ctla-4-deficient zebrafish aged 4 months were used for the experiments.

Generation of Ctla-4-deficient zebrafish

CRISPR/Cas9 system was used to knock out the *ctla-4* gene. The targeting sequence 5'-CTCAGAGCCCTACTTCGCAA-3' was designed by optimized CRISPR Design (http://crispr.mit.edu/) and synthesized by T7 RNA polymerase and purified by MEGAclear Kit (AM1908; Invitrogen) *in vitro*. Cas9 protein (500 ng/μl, A45220P; Thermo Fisher Scientific) and purified RNA (90 ng/μl) were coinjected into one cell-stage wild-type embryos. For genotyping, DNA fragment was amplified with primers (F: 5'-TGTGACAGGAAAAGATGGAGAA-3' and R: 5'-GATCAGATCCACTCCTCCAAAG-3') at 94°C for 4 min followed by 35 cycles at 94°C for 30 s, 58°C for 30 s and 72°C for 30 s, culminating in a final extension at 72°C for 10 min. Subsequently, the PCR product was subjected to sequencing. The mutant alleles (-14 bp) were obtained. As with wild-type zebrafish, Ctla-4-deficient zebrafish

462 were reared in the laboratory in recirculating water at 26-28 °C under standard

463 conditions.

Preparation of recombinant proteins

For prokaryotic expression, the encoding sequences for the extracellular domains

of Ctla-4 and Cd80/86 (designated as soluble Cd80/86, sCd80/86) were amplified and

cloned into the pET-28a (+) and pCold-GST vectors. The primers used are shown in

Supplemental Table 1. The recombinant plasmid was transformed into *Escherichia coli*

BL21 (DE3) competent cells (TransGen Biotech) and induced with isopropyl-β-D-

thiogalactoside (IPTG, 0.5 mM) at 20°C for 12 h. After ultrasonication, the supernatants

were collected for purification. For eukaryotic Ctla-4-Ig and Cd28-Ig expression, the

extracellular domains of Ctla-4 and Cd28 were fused to the Fc region of human IgG1

[77], and cloned into the pAcGHLTc vector. The recombinant plasmids were transfected

into sf9 (*Spodoptera frugiperda*) cells with baculovirus vector DNA (AB Vector) under

the assistance of lipofectamine™ 3000 (Thermo Fisher Scientific). The cells were

cultured at 28°C for 3 days and subsequently dissolved in a lysing buffer (50 mM Tris-

HCl, pH 8.0, 150 mM NaCl, 1% Nonidet P-40, 1 mM PMSF). The recombinant Ctla-

4-Ig and Cd28-Ig proteins (designated as soluble Ctla-4-Ig, sCtla-4-Ig and sCd28-Ig)

were purified using Ni-NTA agarose affinity chromatography (QIAGEN) following the

manufacturer's protocol. Proteins were then separated on a 12% SDS-PAGE gel and

visualized through Coomassie Brilliant Blue R250 staining.

Preparation of polyclonal antibody

483   Antibody (Ab) against the Ctla-4 extracellular domain protein was produced

through a recombinant protein immunized approach as previously described [78]. Briefly, four-week-old male BALB/c mice (~15 g) were immunized with recombinant Ctla-4 protein with extracellular domain (25 μg) each time in CFA (Sigma-Aldrich) initially and then in IFA (Sigma-Aldrich) for four times thereafter at biweekly intervals. Seven days after the final immunization, serum samples were collected when anti-serum titers exceeded 1:10,000. The Ab was affinity purified by Protein-A Agarose Columns (Thermo Fisher Scientific), and its titer was examined by ELISA. The validity and specificity of the Ab was determined by Western blot analysis.

Subcellular localization

HEK293T cells were seeded into the 12-well plates (Corning) with cover glass and cultured in high-glucose DMEM (Gibco) supplemented with 10% FBS (Cell-Box) at 37°C in 5% CO_2 to allow growth until 40%-50% confluence. The cells were transfected with pEGFPN1-Ctla-4 plasmid DNA (0.8 μg) with the help of PEI reagent (3.2 μg per well) in accordance with the manufacturer's protocol. After transfection for 24 h, the cells were fixed with 4% paraformaldehyde (PFA; Sigma-Aldrich) and stained with CM-DiI (1 μM; Thermo Fisher Scientific) and DAPI (100 ng/ml; Sigma-Aldrich). The fluorescence images were obtained using a two-photon laser scanning confocal microscope (LSM710; Zeiss, Jena, Germany) with ×630 magnification.

Identification of monomer or dimer

HEK293T cells were transfected with pCDNA3.1-HA-Ctla-4 (0.8 μg) or pCDNA3.1-HA (0.8 μg) under the assistance of polyethylenimine (PEI; Sigma-Aldrich). After 48 h, the cells were lysed with precooling cell lysis buffer (Beyotime)

and the supernatants were mixed with non-reducing (without β-ME) or reducing (with

β-ME) loading buffer for Western blot analysis.

Immunofluorescence staining

Colocalization of Cd4-1/Cd8α and Ctla-4 was determined by immunofluorescence staining. Leukocytes were isolated from zebrafish's spleen, kidney, and peripheral blood by Ficoll-Hypaque (1.080 g/ml; Sangon Biotech) centrifugation at 2,500 rpm at 25°C for 25 min. After washing with D-Hank's solution, cells were fixed with 4% PFA at room temperature for 10 min, blocked with 2% BSA (Sigma-Aldrich), and incubated with primary Abs at 4°C for 2 h. The primary Abs included combinations of rabbit anti-CD4-1 and mouse anti-CTLA-4, or rabbit anti-CD8α and mouse anti-CTLA-4, which were produced in our laboratory as previously described [75]. Following another wash with D-Hank's solution, the cells were combined with secondary Abs, including FITC-conjugated goat anti-rabbit IgG and PE-conjugated goat anti-mouse IgG (Thermo Fisher Scientific), according to the manufacturer's instructions. After a final wash with D-Hank's solution, the cells were stained with DAPI (100 ng/ml) at room temperature for 10 min. Fluorescence images were captured using a two-photon laser confocal scanning microscope (LSM710; Zeiss, Jena, Germany) with ×630 magnification.

Myeloperoxidase activity measurement

The myeloperoxidase (MPO) activity in intestine and peripheral blood was quantified using a commercial colorimetric assay kit (Nanjing Jiancheng Bioengineering Institute, China) according to the manufacturer's instructions. Briefly, intestinal tissues were homogenized in extraction buffer to obtain a 5% homogenate,

while peripheral blood was mixed with extraction buffer at a 1:1 ratio. A 180 µl aliquot

of resultant mixture was incubated with 20 µl of reaction buffer for 15 min at 37℃.

Enzymatic activity was determined by measuring the changes in absorbance at 460 nm

using a 96-well plate reader. MPO activity was expressed as units per gram of wet

intestinal tissue or per milliliter of peripheral blood.

Co-immunoprecipitation and Western blot analysis

Co-immunoprecipitation (Co-IP) was performed to detect the interaction between

Cd28/Ctla-4 and Cd80/86. HEK293T cells were co-transfected with pLVX-mCherry-

C1-Cd28 (3 µg) and pEGFP-N1-Cd80/86 (3 µg) or pEGFP-N1-Ctla-4 (3 µg) and

pCDNA3.1-HA-Cd80/86 (3 µg) using PEI as a transfection reagent. At 48 h post-

transfection, the cells were lysed with pre-cooled cell lysis buffer (Beyotime). The

lysates were centrifuged at 12,000 × g for 8 minutes at 4°C, and the supernatants were

incubated with mouse anti-myc mAb (Abmart) or mouse anti-HA mAb (Abmart)

overnight at 4℃. Expression of the transfected plasmids was analyzed in the whole cell

lysates as an input control. The following day, the mixture was incubated with 50 µl of

protein A agarose beads (Thermo Fisher Scientific) for 4 h. The beads were washed 3

times with lysis buffer and mixed with loading buffer for SDS-PAGE separation. Target

proteins were transferred onto a 0.22-µm polyvinylidene difluoride (PVDF) membrane

(EMD Millipore) for Western blot analysis. The blots were blocked with 5% skimmed

milk, incubated with mouse or rabbit primary Abs overnight at 4℃, washed with TBST,

and then incubated with HRP-conjugated goat anti-mouse/rabbit IgG mAb (Abmart) at

room temperature for 1 h. Detection was performed using a gel imaging system (Tanon

550   4500).

Histopathological analysis

The anterior, mid, and posterior intestines (n = 3 replicates) were fixed in 4% PFA

overnight and embedded in paraffin. The tissues were cut into 4 μm sections and stained

with hematoxylin and eosin (H&E) for histopathological analysis. To evaluate the

severity of intestinal inflammation, histologic scores were determined based on

established criteria from a previous study [79]. Briefly, three independent parameters,

including inflammation severity, inflammation extent, and epithelial changes, were

assessed and scored as follows: inflammation severity (0 = none, 1 = minimal, 2 = mild,

3 = moderate, 4 = marked); inflammation extent (0 = none, 1 = mucosa, 2 = mucosa

and submucosa, 3 = transmural), epithelial changes (0 = none, 1 = minimal hyperplasia,

2 = mild hyperplasia, minimal goblet cell loss, 3 = moderate hyperplasia, mild goblet

cell loss, 4 = marked hyperplasia with moderate to marked goblet cell loss). Each

parameter was calculated and summed to obtain the overall score. Additionally, tissue

sections were stained with Periodic Acid-Schiff (PAS) or Alcian Blue and Periodic

Acid-Schiff (AB-PAS) reagent to evaluate the mucin components and goblet cell

numbers.

Transmission electron microscopy observation

568   The posterior intestines were cut into 0.2 cm segments and then split lengthwise

569   to expose the intestinal villi to the fixative fully. The samples were first fixed with 2.5%

570   glutaraldehyde in phosphate buffer (0.1 M, pH 7.0) overnight at 4°C, washed three

571   times in the phosphate buffer for 15 min at each step and post-fixed in 1% OsO_4 for 1

h. Following gradient acetone dehydration and Spurr resin infiltration (1:1 and 1:3 mixture of absolute acetone and the final Spurr resin mixture for 1 h and 3 h, and final Spurr resin mixture overnight), the specimens were placed in Eppendorf contained Spurr resin and heated at 70°C for overnight. The samples were sectioned using an ultratome (LEICA EM UC7). Then, the sections were stained with uranyl acetate and alkaline lead citrate for 10 min each and observed under a transmission electron microscopy (Hitachi Model H-7650).

Assessment of apoptosis by TUNEL

The posterior intestines were fixed by 4% paraformaldehyde and embedded in paraffin. Apoptosis was detected using TUNEL staining following the manufacturer's protocol (Beyotime). Briefly, deparaffinized sections were incubated with biotin-dUTP labeling solution (TdT Enzyme: Biotin-dUTP = 1: 9) for 1 h, followed by incubation with streptavidin-HRP for 30 min at room temperature. Positive signals were visualized using DAB chromogenic solution and counterstained with hematoxylin. Apoptotic cells and the area of the intestinal epithelium were quantified, and the apoptosis index was calculated as the number of apoptotic cells per 1×10^4 μm^2 observed using ImageJ software.

RNA-sequencing and bioinformatic analysis

Total RNAs were isolated from wild-type or *ctla-4$^{-/-}$* intestines (three biological replicates) using TRIzol reagent following the manufacturer's instructions (Takara). cDNA libraries were constructed using NEB Next Ultra Directional RNA Library Prep Kit (NEB), and sequencing was performed according to the Illumina Hiseq2500

standard protocol at LC Bio (Hangzhou, China). The differentially expressed genes

(DEGs) were identified with absolute Log_2 fold change > 1 and adjusted p-value < 0.05

by R package DESeq2. Gene Ontology (GO) enrichment and Kyoto encyclopedia of

genes and genomes (KEGG) enrichment analyses were performed by the OmicStudio

(http://www.omicstudio.cn/tool) tools. Gene-set enrichment analysis was performed

using software GSEA (v4.1.0, https://www.gsea-msigdb.org/gsea/index.jsp), and the

heatmap was generated using the R package ggplot2. For the protein-protein interaction

(PPI) networks, the DEGs were retrieved in STRING (version 11.5, https://string-

db.org/) database (combined score > 0.4), and the PPI network was visualized by

Cytoscape software (version 3.9.1) [80]. The betweenness centrality (BC) was

calculated using the CytoNCA plugin in Cytoscape software. The RNA-sequencing

(RNA-seq) data in this study were deposited in the Gene Expression Omnibus (GEO)

(http://www.ncbi.nlm.nih.gov/geo/) database.

Quantitative real-time PCR

The transcript abundance of target genes was quantified using quantitative real-

time PCR on a CFX Connect Real-Time PCR Detection System equipped with

Precision Melt Analysis$^{\text{TM}}$ Software (Bio-Rad, Cat. No. 1855200EM1). Total RNA

from intestines was extracted using TRIzol reagent (Takara Bio) and reverse transcribed

into cDNAs according to the manufacturer's protocol. PCR experiments were

performed in a total volume of 10 μl by using an SYBR Premix Ex Taq kit (Takara Bio).

The reaction mixtures were incubated for 2 min at 95℃, then subjected to 40 cycles of

15 s at 95℃, 15 s at 60℃, and 20 s at 72℃. Relative expression levels of the target

616 genes were calculated using the $2^{-\Delta\Delta ct}$ method with β-actin for normalization. Each PCR

617 trial was run in triplicate parallel reactions and repeated three times. The primers used

618 in the experiments are listed in Supplemental Table 1.

Single-cell RNA-sequencing analysis

The intestines from wild-type (30 fish) and *ctla-4*$^{-/-}$ zebrafish (30 fish) were

washed by D-Hank's and incubated with D-Hank's containing EDTA (0.37 mg/mL) and

DTT (0.14 mg/mL) for 20 min. The resulting supernatants were collected as fraction 1.

The remaining tissues were subsequently digested with type IV collagenase (0.15

mg/mL) for 1 h at room temperature and the resulting supernatants were collected as

fraction 2. Fractions 1 and 2 were combined and centrifuged at 350 g for 10 min. Cells

were then washed with D-Hank's and suspended in a 40% Percoll (Shanghai Yes

Service Biotech, China) solution (Percoll: FBS: L-15 medium = 4: 1: 5) and passed

through a 40-μm strainer (Bioland). The cell suspension was carefully layered onto 63%

Percoll (Percoll: FBS: L-15 medium = 6.3: 1: 2.7) and centrifuged at 400 g for 30 min

at room temperature. The cell layer at the interface was collected and washed with D-

Hank's at 400 g for 10 min. Cell quantity and viability were assessed using 0.4% trypan

blue staining, revealing that over 90% of the cells were viable. Single-cell samples

(8,047 cells in wild-type group, 8,321 cells in *ctla-4*$^{-/-}$ group) were submitted to the LC-

Bio Technology Co., Ltd (Hangzhou, China) for 10× Genomics library preparation and

data analysis assistance. Libraries were prepared using the Chromium™ Controller and

Chromium™ Single Cell 3' Library & Gel Bead Kit v2 (10× Genomics) according to

the manufacturer's protocol, and sequenced on an Illumina NovaSeq 6000 sequencing

system (paired-end multiplexing run, 150 bp) at a minimum depth of 20,000 reads per

cell. Sequencing results were demultiplexed and converted to FASTQ format using

Illumina bcl2fastq software and the data were aligned to Ensembl zebrafish genome

assembly GRCz11. Quality control was performed using the Seurat. DoubletFinder R

package was used to identify and filter the doublets (multiplets) [81]. The cells were

removed if they expressed fewer than 500 unique genes, or > 60% mitochondrial reads.

Cloud-based Cumulus v1.0 was used to perform the bath correction (using the Harmony

algorithm) on the aggregated gene-count matrices [82]. The number of cells after

filtration in the current study was 3,263 in wild-type and 4,276 in *ctla-4*[-/-] groups,

respectively. Cells were grouped into an optimal number of clusters for *de novo* cell

type discovery using Seurat's FindNeighbors() and FindClusters() functions, graph-

based clustering approach with visualization of cells being achieved through the use of

t-SNE or UMAP plots [83]. The cell types were determined using a combination of

marker genes identified from the literature and gene ontology. The marker genes were

visualized by dot plot and t-SNE plots to reveal specific individual cell types.

16S rRNA sequencing analysis

Intestinal contents were collected from both wild-type and *ctla-4*[-/-] zebrafish by

gently squeezing the intestines with fine-tipped tweezers, and the remaining intestines

were used for single-cell RNA-sequencing analysis. Contents from six fish were pooled

to form one replicate, with each experimental sample comprising four replicates. DNA

was extracted from the samples using the CTAB method, a protocol known for its

efficacy in isolating DNA from trace quantities. The quality of DNA was assessed

through agarose gel electrophoresis. Total DNA was amplified to construct sequencing libraries using primers (341F: 5'-CCTACGGGNGGCWGCAG-3'; 805R: 5'-GACTACHVGGGTATCTAATCC-3') targeting the V3-V4 regions of the 16S rRNA gene. The libraries were sequenced on an Illumina NovaSeq PE250 platform. Quality filtering was performed under specific conditions to obtain high-quality clean tags using fqtrim (v0.94). Chimeric sequences were removed using Vsearch software (v2.3.4). After dereplication using DADA2, a feature table and feature sequences were generated. Alpha diversity is applied in analyzing the complexity of species diversity for a sample through the Shannon and Simpson indices, with all calculations performed in QIIME2. Beta diversity analysis was also conducted in QIIME2, and the graphs were drawn by R package. Sequence alignment was performed using BLAST, and representative sequences were annotated using the SILVA database. Other diagrams were implemented using the R package (v3.5.2).

***In vitro* lymphocyte proliferation assay**

The leukocytes were prepared from the spleen, kidney, and peripheral blood of wild-type (10 fish) or *ctla-4*$^{-/-}$ (10 fish) zebrafish through Ficoll-Hypaque centrifugation. A total of 2×10^6 leukocytes from either wild-type or *ctla-4*$^{-/-}$ individuals were labeled with 1 μM carboxyfluorescein succinimidyl ester (CFSE; Thermo Fisher Scientific) at 25°C for 5 min. The reaction was terminated by adding a triploid volume of Leibovitz L-15 medium (Gibco) supplemented with 10% FBS, as previously described [84]. After washing with D-Hank's solution, the cells were cultured in L-15 medium containing 10% FBS in the presence or absence of PHA (5 μg/ml), recombinant Ctla-4-Ig (20 μg/ml),

Cd28-Ig (20 μg/ml), Cd80/86 (10 μg/ml) proteins or anti-Ctla-4 Ab (10 μg/ml) at 28℃

for 3 days. CFSE fluorescence intensity in the labeled co-cultures was analyzed using

a flow cytometer (FACSJazz; BD Biosciences) to assess cell division [85].

Prediction of protein interactions by AlphaFold2

AlphaFold2 (version 2.3.2; available at https://github.com/google-deepmind/

alphafold) was implemented on a high-performance computing cluster to predict the

structures of the Cd80/86 complexes with Cd28 and Ctla-4 [85]. The resulting models

were ranked based on their per-residue Local Distance Difference Test (pLDDT) scores,

which quantify the confidence level of each residue on a scale from 0 to 100. Residues

were color-coded according to their pLDDT scores, with higher values reflecting

greater confidence in the prediction. Furthermore, the Predicted Aligned Error (PAE)

scores were analyzed to identify well-defined interaction interfaces between Cd28 or

Ctla-4 and Cd80/86.

Microscale thermophoresis assay

The binding affinity between Cd80/86 and Cd28/Ctla-4 were measured through

microscale thermophoresis (MST) assays using a Monolith NT.115 instrument (Nano

Temper Technologies) as previously described [86]. In each assay, the labeled proteins

(Cd28/Ctla-4 with EGFP-tag) were incubated with varying concentrations of unlabeled

ligand protein (Cd80/86) for 10 min at room temperature. The initial protein

concentration of 3.2 μM was diluted into 16 different concentrations by doubling

dilution. The samples were then loaded into capillaries and analyzed at 25℃ by using

40% light emitting diode (LED) and medium MST power. The binding affinities of

Cd80/86 with Cd28 and Cd80/86 with Ctla-4 were examined using the same parameters.

Each assay was repeated three to five times, and dissociation constants (K_D) were

calculated using MO.Affinity Analysis software.

***In vivo* administration of sCtla-4-Ig**

An *in vivo* sCtla-4-Ig administration assay was conducted to evaluate the potential

therapeutic effect of sCTLA-4-Ig on intervention of a *ctla-4*-deficiency induced IBD-

like phenotype. For this procedure, the *ctla-4*$^{-/-}$ zebrafish were i.p administered with

recombinant soluble Ctla-4-Ig protein (sCtla-4-Ig, 2 μg/g body weight) every other day

until day 14. Fish that received an equal amount of human IgG isotype were used as

control. The dose of sCtla-4-Ig was chosen based on its ability to significantly inhibit

the mRNA expression of *il13* in Ctla-4-deficient zebrafish.

Statistical analysis

Statistical analysis and graphical presentation were performed with GraphPad

Prism 8.0. All data were presented as the mean ± SD of each group. Statistical

evaluation of differences was assessed using one-way ANOVA, followed by an

unpaired two-tailed *t*-teat. Statistical significance was defined as $^*p < 0.05$, $^{**}p < 0.01$,

$^{***}p < 0.001$, and $^{****}p < 0.0001$. All experiments were replicated at least three times.

Data availability

All data generated or analyzed during this study are included in this article and its

supplementary information files. The RNA-seq and scRNA-seq data for this study have

been deposited in NCBI Gene Expression Omnibus (GEO)

(https://www.ncbi.nlm.nih.gov/geo/) under accession numbers GSE255304 and GSE255303, respectively. The 16S rRNA sequencing data in this study have been deposited in the NCBI Sequence Read Archive (SRA) (https://www.ncbi.nlm.nih.gov/sra/) with an accession number of BioProject PRJNA1073727.

Acknowledgements

We are grateful to Bio-ultrastructure Analysis Laboratory at the Analysis Center of Agrobiology and Environmental Sciences, Zhejiang University, for their assistance in TEM sample preparation and observation. We also thank Hong Deng and Qiong Huang for their valuable advice and expertise in pathological analysis. Additionally, we acknowledge She-long Zhang for his support in two-photon laser confocal scanning microscope capture. This work was supported by grants from the National Natural Science Foundation of China (32173003) and the National Key Research and Development Program of China (2018YFD0900503, 2018YFD0900505).

Author Contributions

L. L. Q. and J. Z. S. conceived and designed the experiments. L. L. Q. and C. B. H. conducted the experiments. Data analysis was performed by L. L. Q., C. B. H., Q. Z., Y. W., A. F. L., L. X. X., Y. C., and J. Z. S. Reagents, materials, and analysis tools were contributed by L. X. X., D. D. F., and J. Z. S. The manuscript was written by L. L. Q., Y. C., L. X. X., and J. Z. S. All authors reviewed the manuscript and provided

feedback.

Conflict of Interest

The authors declare that they have no competing financial interests.

Ethical approval and consent to participate

All animal experiments were performed with the approval of the Ethics Committee for

Animal Experimentation of Zhejiang University.

REFERENCES

1. Hodson R (2016) Inflammatory bowel disease. Nature 540:S97‑S97.
https://10.1038/540S97a
2. Zhang Y Z, Li Y Y (2014) Inflammatory bowel disease: Pathogenesis. World J Gastroentero
20:91‑99. https://10.3748/wjg.v20.i1.91
3. Venema W T C U, Voskuil M D, Dijkstra G, Weersma R K, Festen E A M (2017) The genetic
background of inflammatory bowel disease: from correlation to causality. J Pathol
241:146‑158. https://10.1002/path.4817
4. Loftus E V (2004) Clinical epidemiology of inflammatory bowel disease: Incidence,
prevalence, and environmental influences. Gastroenterology 126:1504‑1517.
https://10.1053/j.gastro.2004.01.063
5. Khor B, Gardet A, Xavier R J (2011) Genetics and pathogenesis of inflammatory bowel
disease. Nature 474:307‑317. https://10.1038/nature10209
6. Neurath M F (2020) Host-microbiota interactions in inflammatory bowel disease. Nat Rev
Gastro Hepat 17:76‑77. https://10.1038/s41575‑019‑0248‑1
7. Kordjazy N, Haj-Mirzaian A, Haj-Mirzaian A, Rohani M M, Gelfand E W, Rezaei N *et al.*
(2018) Role of toll-like receptors in inflammatory bowel disease. Pharmacological
Research 129:204‑215. https://10.1016/j.phrs.2017.11.017
8. Horowitz J E, Warner N, Staples J, Crowley E, Gosalia N, Murchie R *et al.* (2021) Mutation
spectrum of NOD2 reveals recessive inheritance as a main driver of Early Onset Crohn's
Disease. Sci Rep-Uk 11:5595. https://10.1038/s41598‑021‑84938‑8
9. Duerr R H, Taylor K D, Brant S R, Rioux J D, Silverberg M S, Daly M J *et al.* (2006) A genome-
wide association study identifies IL23R as an inflammatory bowel disease gene. Science
314:1461‑1463. https://10.1126/science.1135245
10. Tremelling M, Berzuini C, Massey D, Bredin F, Price C, Dawson C *et al.* (2008) Contribution

of TNFSF15 gene variants to Crohn's disease susceptibility confirmed in UK population.
Inflamm Bowel Dis 14:733-737. https://10.1002/ibd.20399

11. Rudd C E, Schneider H (2003) Unifying concepts in CD28, ICOS and CTLA4 co-receptor
signalling. Nat Rev Immunol 3:544-556. https://10.1038/nri1131

12. Yang Y, Li X, Ma Z, Wang C, Yang Q, Byrne-Steele M *et al.* (2021) CTLA-4 expression by
B-1a B cells is essential for immune tolerance. Nat Commun 12:525.
https://10.1038/s41467-020-20874-x

13. Kim G R, Choi J M (2022) Current Understanding of Cytotoxic T Lymphocyte Antigen-4
(CTLA-4) Signaling in T-Cell Biology and Disease Therapy. Mol Cells 45:513-521.
https://10.14348/molcells.2022.2056

14. Marengere L E, Waterhouse P, Duncan G S, Mittrucker H W, Feng G S, Mak T W (1996)
Regulation of T cell receptor signaling by tyrosine phosphatase SYP association with
CTLA-4. Science 272:1170-1173. https://10.1126/science.272.5265.1170

15. Lee K M, Chuang E, Griffin M, Khattri R, Hong D K, Zhang W G *et al.* (1998) Molecular basis
of T cell inactivation by CTLA-4. Science 282:2263-2266.
https://10.1126/science.282.5397.2263

16. Yokosuka T, Kobayashi W, Takamatsu M, Sakata-Sogawa K, Zeng H, Hashimoto-Tane A
et al. (2010) Spatiotemporal Basis of CTLA-4 Costimulatory Molecule-Mediated Negative
Regulation of T Cell Activation. Immunity 33:326-339.
https://10.1016/j.immuni.2010.09.006

17. Linsley P S, Greene J L, Brady W, Bajorath J, Ledbetter J A, Peach R (1994) Human B7-1
(Cd80) and B7-2 (Cd86) Bind with Similar Avidities but Distinct Kinetics to Cd28 and Ctla-
4 Receptors. Immunity 1:793-801. https://10.1016/S1074-7613(94)80021-9

18. Collins A V, Brodie D W, Gilbert R J, Iaboni A, Manso-Sancho R, Walse B *et al.* (2002) The
interaction properties of costimulatory molecules revisited. Immunity 17:201-210.
https://10.1016/s1074-7613(02)00362-x

19. Saito T, Yokosuka T, Hashimoto-Tane A (2010) Dynamic regulation of T cell activation and
co-stimulation through TCR-microclusters. Febs Lett 584:4865-4871.
https://10.1016/j.febslet.2010.11.036

20. Tivol E A, Borriello F, Schweitzer A N, Lynch W P, Bluestone J A, Sharpe A H (1995) Loss of
Ctla-4 Leads to Massive Lymphoproliferation and Fatal Multiorgan Tissue Destruction,
Revealing a Critical Negative Regulatory Role of Ctla-4. Immunity 3:541-547.
https://10.1016/1074-7613(95)90125-6

21. Waterhouse P, Penninger J M, Timms E, Wakeham A, Shahinian A, Lee K P *et al.* (1995)
Lymphoproliferative Disorders with Early Lethality in Mice Deficient in Ctla-4. Science
270:985-988. https://10.1126/science.270.5238.985

22. Chambers C A, Sullivan T J, Allison J P (1997) Lymphoproliferation in CTLA-4-deficient
mice is mediated by costimulation-dependent activation of CD4(+) T cells. Immunity
7:885-895. https://10.1016/S1074-7613(00)80406-9

23. Khattri R, Auger J A, Griffin M D, Sharpe A H, Bluestone J A (1999) Lymphoproliferative
Disorder in CTLA-4 Knockout Mice Is Characterized by CD28-Regulated Activation of Th2
Responses. The Journal of Immunology 162:5784-5791.
https://10.4049/jimmunol.162.10.5784

24. Bour-Jordan H, Grogan J L, Tang Q Z, Auger J A, Locksley R M, Bluestone J A (2003) CTLA-

4 regulates the requirement for cytokine-induced signals in T$_H$2 lineage commitment.
Nature Immunology 4:182-188. https://10.1038/ni884

25. Klocke K, Sakaguchi S, Holmdahl R, Wing K (2016) Induction of autoimmune disease by
deletion of CTLA-4 in mice in adulthood. Proc Natl Acad Sci U S A 113:E2383-2392.
https://10.1073/pnas.1603892113

26. Wing K, Onishi Y, Prieto-Martin P, Yamaguchi T, Miyara M, Fehervari Z *et al.* (2008) CTLA-
4 control over Foxp3+ regulatory T cell function. Science 322:271-275.
https://10.1126/science.1160062

27. Repnik K, Potocnik U (2010) CT60 Single-Nucleotide Polymorphism Is Associated with
Slovenian Inflammatory Bowel Disease Patients and Regulates Expression of Isoforms.
DNA Cell Biol 29:603-610. https://10.1089/dna.2010.1021

28. Xia B, Crusius J B A, Wu J, Zwiers A, van Bodegraven A A, Peña A S (2002) CTLA-4 gene
polymorphisms in Dutch and Chinese patients with inflammatory bowel disease. Scand J
Gastroentero 37:1296-1300. https://10.1080/003655202761020579

29. Jiang T, Ge L Q, Chen Z T, Li C, Zhou F, Luo Y *et al.* (2010) Effect of cytotoxic T lymphocyte-
associated molecule 4 1661 gene polymorphism on its expression and transcription in
ulcerative colitis. J Digest Dis 11:369-375. https://10.1111/j.1751-2980.2010.00462.x

30. Bamias G, Delladetsima I, Perdiki M, Siakavellas S I, Goukos D, Papatheodoridis G V *et al.*
(2017) Immunological Characteristics of Colitis Associated with Anti-CTLA-4 Antibody
Therapy. Cancer Invest 35:443-455. https://10.1080/07357907.2017.1324032

31. Lo B C, Kryczek I, Yu J, Vatan L, Caruso R, Matsumoto M *et al.* (2024) Microbiota-
dependent activation of CD4+ T cells induces CTLA-4 blockade–associated colitis via Fcγ
receptors. Science 383:62-70. https://10.1126/science.adh8342

32. Cooper M D, Alder M N (2006) The evolution of adaptive immune systems. Cell 124:815-
822. https://10.1016/j.cell.2006.02.001

33. Lam S H, Chua H L, Gong Z, Lam T J, Sin Y M (2004) Development and maturation of the
immune system in zebrafish, Danio rerio: a gene expression profiling, in situ hybridization
and immunological study. Dev Comp Immunol 28:9-28. https://10.1016/s0145-
305x(03)00103-4

34. Peach R J, Bajorath J, Brady W, Leytze G, Greene J, Naemura J *et al.* (1994)
Complementarity-Determining Region-1 (Cdr1)-Analogous and Cdr3-Analogous
Regions in Ctla-4 and Cd28 Determine the Binding to B7-1. J Exp Med 180:2049-2058.
https://10.1084/jem.180.6.2049

35. Shiratori T, Miyatake S, Ohno H, Nakaseko C, Isono K, Bonifacino J S *et al.* (1997) Tyrosine
phosphorylation controls internalization of CTLA-4 by regulating its interaction with
clathrin-associated adaptor complex AP-2. Immunity 6:583-589. https://10.1016/s1074-
7613(00)80346-5

36. Bernard D, Riteau B, Hansen J D, Phillips R B, Michel F, Boudinot P *et al.* (2006)
Costimulatory receptors in a teleost fish: Typical CD28, elusive CTLA4. Journal of
Immunology 176:4191-4200. https://10.4049/jimmunol.176.7.4191

37. Hu C B, Wang J, Hong Y, Li H, Fan D D, Lin A F *et al.* (2023) Single-cell transcriptome
profiling reveals diverse immune cell populations and their responses to viral infection in
the spleen of zebrafish. Faseb J 37:e22951. https://10.1096/fj.202201505RRRR

38. Butin-Israeli V, Bui T M, Wiesolek H L, Mascarenhas L, Lee J J, Mehl L C *et al.* (2019)

Neutrophil-induced genomic instability impedes resolution of inflammation and wound
healing. J Clin Invest 129:712-726. https://10.1172/JCI122085

39.   Lees C W, Barrett J C, Parkes M, Satsangi J (2011) New IBD genetics: common pathways
with other diseases. Gut 60:1739-1753. https://10.1136/gut.2009.199679

40.   Lee J C, Biasci D, Roberts R, Gearry R B, Mansfield J C, Ahmad T et al. (2017) Genome-
wide association study identifies distinct genetic contributions to prognosis and
susceptibility in Crohn's disease. Nat Genet 49:262-268. https://10.1038/ng.3755

41.   Bernink J H, Peters C P, Munneke M, te Velde A A, Meijer S L, Weijer K et al. (2013) Human
type 1 innate lymphoid cells accumulate in inflamed mucosal tissues. Nat Immunol
14:221-229. https://10.1038/ni.2534

42.   Li J, Doty A L, Tang Y, Berrie D, Iqbal A, Tan S A et al. (2017) Enrichment of IL-17A(+) IFN-
gamma(+) and IL-22(+) IFN-gamma(+) T cell subsets is associated with reduction of
NKp44(+) ILC3s in the terminal ileum of Crohn's disease patients. Clin Exp Immunol
190:143-153. https://10.1111/cei.12996

43.   Martin J C, Chang C, Boschetti G, Ungaro R, Giri M, Grout J A et al. (2019) Single-Cell
Analysis of Crohn's Disease Lesions Identifies a Pathogenic Cellular Module Associated
with   Resistance   to   Anti-TNF   Therapy.   Cell   178:1493-1508   e1420.
https://10.1016/j.cell.2019.08.008

44.   Buonocore S, Ahern P P, Uhlig H H, Ivanov, II, Littman D R, Maloy K J et al. (2010) Innate
lymphoid cells drive interleukin-23-dependent innate intestinal pathology. Nature
464:1371-1375. https://10.1038/nature08949

45.   Ermann J, Staton T, Glickman J N, de Waal Malefyt R, Glimcher L H (2014) Nod/Ripk2
signaling in dendritic cells activates IL-17A-secreting innate lymphoid cells and drives
colitis in T-bet-/-.Rag2-/- (TRUC) mice. Proc Natl Acad Sci U S A 111:E2559-2566.
https://10.1073/pnas.1408540111

46.   Aparicio-Domingo P, Romera-Hernandez M, Karrich J J, Cornelissen F, Papazian N,
Lindenbergh-Kortleve D J et al. (2015) Type 3 innate lymphoid cells maintain intestinal
epithelial   stem   cells   after   tissue   damage.   J   Exp   Med   212:1783-1791.
https://10.1084/jem.20150318

47.   Ott S J, Musfeldt M, Wenderoth D F, Hampe J, Brant O, Folsch U R et al. (2004) Reduction
in diversity of the colonic mucosa associated bacterial microflora in patients with active
inflammatory bowel disease. Gut 53:685-693. https://10.1136/gut.2003.025403

48.   Manichanh C, Rigottier-Gois L, Bonnaud E, Gloux K, Pelletier E, Frangeul L et al. (2006)
Reduced diversity of faecal microbiota in Crohn's disease revealed by a metagenomic
approach. Gut 55:205-211. https://10.1136/gut.2005.073817

49.   Zhao Q, Chang H, Zheng J, Li P, Ye L, Pan R et al. (2023) A novel Trmt5-deficient zebrafish
model with spontaneous inflammatory bowel disease-like phenotype. Signal Transduct
Target Ther 8:86. https://10.1038/s41392-023-01318-6

50.   Xu Z, Takizawa F, Casadei E, Shibasaki Y, Ding Y, Sauters T J C et al. (2020) Specialization
of mucosal immunoglobulins in pathogen control and microbiota homeostasis occurred
early   in   vertebrate   evolution.   Sci   Immunol   5:3254-3271.
https://10.1126/sciimmunol.aay3254

51.   Belzer C, de Vos W M (2012) Microbes inside--from diversity to function: the case of
Akkermansia. ISME J 6:1449-1458. https://10.1038/ismej.2012.6

52. Hosseini A, Gharibi T, Marofi F, Babaloo Z, Baradaran B (2020) CTLA-4: From mechanism
to autoimmune therapy. Int Immunopharmacol 80:106221.
https://10.1016/j.intimp.2020.106221

53. Sun W H, Zhang X M, Wu J, Zhao W D, Zhao S X, Li M L (2019) Correlation of TSHR and
CTLA-4 Single Nucleotide Polymorphisms with Graves Disease. Int J Genomics
2019:6982623. https://10.1155/2019/6982623

54. Vergara A, De Felice M, Cesaro A, Gragnano F, Pariggiano I, Golia E *et al.* (2023) Immune-
Checkpoint Inhibitor-Related Myocarditis: Where We Are and Where We Will Go.
Angiology 12:33197231201929. https://10.1177/00033197231201929

55. Kheiralla K E K (2021) CTLA-4 (+49A/G) Polymorphism in Type 1 Diabetes Children of
Sudanese Population. Glob Med Genet 8:11-18. https://10.1055/s-0041-1723008

56. Lin T W, Hu Y C, Yang Y H, Chien Y H, Lee N C, Yu H H *et al.* (2022) CTLA-4 gene mutation
and multiple sclerosis: A case report and literature review. J Microbiol Immunol 55:545-
548. https://10.1016/j.jmii.2021.10.009

57. Cutolo M, Sulli A, Paolino S, Pizzorni C (2016) CTLA-4 blockade in the treatment of
rheumatoid arthritis: an update. Expert Rev Clin Immu 12:417-425.
https://10.1586/1744666x.2016.1133295

58. Chang M C, Chang Y T, Tien Y W, Liang P C, Jan I S, Wei S C *et al.* (2007) T-Cell regulatory
gene CTLA-4 Polymorphism/Haplotype association with autoimmune pancreatitis. Clin
Chem 53:1700-1705. https://10.1373/clinchem.2007.085951

59. Fathima N, Narne P, Ishaq M (2019) Association and gene-gene interaction analyses for
polymorphic variants in CTLA-4 and FOXP3 genes: role in susceptibility to autoimmune
thyroid disease. Endocrine 64:591-604. https://10.1007/s12020-019-01859-3

60. Zeissig S, Petersen B S, Tomczak M, Melum E, Huc-Claustre E, Dougan S K *et al.* (2015)
Early-onset Crohn's disease and autoimmunity associated with a variant in CTLA-4. Gut
64:1889-1897. https://10.1136/gutjnl-2014-308541

61. Angelino G, Cifaldi C, Zangari P, Di Cesare S, Di Matteo G, Chiriaco M *et al.* (2021) Gastric
cancer, inflammatory bowel disease and polyautoimmunity in a 17-year-old boy: CTLA-
4 deficiency successfully treated with Abatacept. Eur J Gastroen Hepat 33:E1051-E1056.
https://10.1097/Meg.0000000000002185

62. Liu J Z, van Sommeren S, Huang H L, Ng S C, Alberts R, Takahashi A *et al.* (2015)
Association analyses identify 38 susceptibility loci for inflammatory bowel disease and
highlight shared genetic risk across populations. Nat Genet 47:979-+.
https://10.1038/ng.3359

63. Kim J J, Shajib M S, Manocha M M, Khan W I (2012) Investigating Intestinal Inflammation
in DSS-induced Model of IBD. Jove-J Vis Exp 10.3791/3678 https://10.3791/3678

64. Geiger B, Zhen A W, Kokkotou E, Fraenkel P G (2013) Granulocytes Infiltrate The Intestine
In a Zebrafish Model Of Inflammatory Bowel Disease. Blood
122:https://10.1182/blood.V122.21.2276.2276

65. Fleming A, Jankowski J, Goldsmith P (2010) In Vivo Analysis of Gut Function and Disease
Changes in a Zebrafish Larvae Model of Inflammatory Bowel Disease: A Feasibility Study.
Inflamm Bowel Dis 16:1162-1172. https://10.1002/ibd.21200

66. Oehlers S H, Flores M V, Hall C J, Crosier K E, Crosier P S (2012) Retinoic acid suppresses
intestinal mucus production and exacerbates experimental enterocolitis. Dis Model Mech

5:457-467. https://10.1242/dmm.009365

67.   Coronado M, Solis C J, Hernandez P P, Feijóo C G (2019) Soybean Meal-Induced Intestinal
Inflammation in Zebrafish Is T Cell-Dependent and Has a Th17 Cytokine Profile. Frontiers
in Immunology 10:610-622. https://10.3389/fimmu.2019.00610

68.   Al-Sadi R, Guo S H, Ye D M, Dokladny K, Alhmoud T, Ereifej L *et al.* (2013) Mechanism of
IL-1b Modulation of Intestinal Epithelial Barrier Involves p38 Kinase and Activating
Transcription Factor-2 Activation. Journal of Immunology 190:6596-6606.
https://10.4049/jimmunol.1201876

69.   Al-Sadi R, Guo S H, Ye D M, Ma T Y (2013) TNF-α Modulation of Intestinal Epithelial Tight
Junction Barrier Is Regulated by ERK1/2 Activation of Elk-1. Am J Pathol 183:1871-1884.
https://10.1016/j.ajpath.2013.09.001

70.   Heller F, Florian P, Bojarski C, Richter J, Christ M, Hillenbrand B *et al.* (2005) Interleukin-13
is the key effector Th2 cytokine in ulcerative colitis that affects epithelial tight junctions,
apoptosis, and cell restitution. Gastroenterology 129:550-564.
https://10.1016/j.gastro.2005.05.002

71.   Shin N R, Whon T W, Bae J W (2015) Proteobacteria: microbial signature of dysbiosis in
gut microbiota. Trends Biotechnol 33:496-503. https://10.1016/j.tibtech.2015.06.011

72.   Winter S E, Lopez C A, Bäumler A J (2013) The dynamics of gut-associated microbial
communities during inflammation. Embo Rep 14:319-327.
https://10.1038/embor.2013.27

73.   Ott S J, Musfeldt M, Wenderoth D F, Hampe J, Brant O, Fölsch U R *et al.* (2004) Reduction
in diversity of the colonic mucosa associated bacterial microflora in patients with active
inflammatory bowel disease. Gut 53:685-693. https://10.1136/gut.2003.025403

74.   Hughes E R, Winter M G, Duerkop B A, Spiga L, de Carvalho T F, Zhu W H *et al.* (2017)
Microbial Respiration and Formate Oxidation as Metabolic Signatures of Inflammation-
Associated Dysbiosis. Cell Host Microbe 21:208-219. https://10.1016/j.chom.2017.01.005

75.   Ahmed A, Joseph A M, Zhou J, Horn V, Uddin J, Lyu M *et al.* (2024) CTLA-4-expressing
ILC3s restrain interleukin-23-mediated inflammation. Nature 630:976-983.
https://10.1038/s41586-024-07537-3

76.   Shi W, Shao T, Li J Y, Fan D D, Lin A F, Xiang L X *et al.* (2019) BTLA-HVEM Checkpoint Axis
Regulates Hepatic Homeostasis and Inflammation in a ConA-Induced Hepatitis Model in
Zebrafish. Journal of Immunology 203:2425-2442. https://10.4049/jimmunol.1900458

77.   Linsley P S, Brady W, Grosmaire L, Aruffo A, Damle N K, Ledbetter J A (1991) Binding of
the B cell activation antigen B7 to CD28 costimulates T cell proliferation and interleukin 2
mRNA accumulation. J Exp Med 173:721-730. https://10.1084/jem.173.3.721

78.   Shi W, Shao T, Li J Y, Fan D D, Lin A F, Xiang L X *et al.* (2019) BTLA-HVEM Checkpoint Axis
Regulates Hepatic Homeostasis and Inflammation in a ConA-Induced Hepatitis Model in
Zebrafish. J Immunol 203:2425-2442. https://10.4049/jimmunol.1900458

79.   Erben U, Loddenkemper C, Doerfel K, Spieckermann S, Haller D, Heimesaat M M *et al.*
(2014) A guide to histomorphological evaluation of intestinal inflammation in mouse
models. Int J Clin Exp Pathol 7:4557-4576.

80.   Kohl M, Wiese S, Warscheid B (2011) Cytoscape: software for visualization and analysis of
biological networks. Methods Mol Biol 696:291-303. https://10.1007/978-1-60761-987-
1_18

81.   McGinnis C S, Murrow L M, Gartner Z J (2019) DoubletFinder: Doublet Detection in Single-
1003   Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst 8:329-+.
1004   https://10.1016/j.cels.2019.03.003

1005   82.   Li B, Gould J, Yang Y, Sarkizova S, Tabaka M, Ashenberg O *et al.* (2020) Cumulus provides
1006   cloud-based data analysis for large-scale single-cell and single-nucleus RNA-seq. Nat
1007   Methods 17:793-798. https://10.1038/s41592-020-0905-x

1008   83.   Cronan M R, Hughes E J, Brewer W J, Viswanathan G, Hunt E G, Singh B *et al.* (2021) A
1009   non-canonical type 2 immune response coordinates tuberculous granuloma formation
1010   and epithelialization. Cell 184:1757-1774 e1714. https://10.1016/j.cell.2021.02.046

1011   84.   Quah B J C, Warren H S, Parish C R (2007) Monitoring lymphocyte proliferation and with
1012   the intracellular fluorescent dye carboxyfluorescein diacetate succinimidyl ester. Nature
1013   Protocols 2:2049-2056. https://10.1038/nprot.2007.296

1014   85.   Rieder S A, Wang J Y, White N, Qadri A, Menard C, Stephens G *et al.* (2021) B7-H7 (HHLA2)
1015   inhibits T-cell activation and proliferation in the presence of TCR and CD28 signaling. Cell
1016   Mol Immunol 18:1503-1511. https://10.1038/s41423-020-0361-7

1017   86.   Jerabek-Willemsen M, André T, Wanner R, Roth H M, Duhr S, Baaske P *et al.* (2014)
1018   MicroScale Thermophoresis: Interaction analysis and beyond. J Mol Struct 1077:101-113.
1019   https://10.1016/j.molstruc.2014.03.009

1020

1021

1022

Fig.1 Characterization of zebrafish Ctla-4. **A** Alignment of the Ctla-4 homologs from

different species generated with ClustalX and Jalview. The conserved and partially

conserved amino acid residues in each species are colored in hues graded from orange

to red, respectively. Key features, including conserved cysteine residues, functional

motifs, such as B7-binding motif, tyrosine phosphorylation site, and potential tyrosine

phosphorylation site, were indicated separately. The signal peptide, IgV-like domain,

transmembrane (TM) domain, and cytoplasmic domain were marked above the

sequence. **B** The tertiary structure of the zebrafish Ctla-4 ectodomain, as predicted by AlphaFold2, was compared with that of humans. The two pairs of disulfide bonds (Cys^{20}-Cys^{91}/Cys^{46}-Cys^{65} in zebrafish and Cys^{21}-Cys^{92}/Cys^{48}-Cys^{66} in humans) used to connect the two-layer β-sandwich, and the separate Cys residue (Cys^{119} in zebrafish and Cys^{120} in humans) involved in the dimerization of the proteins are indicated. Cysteine residues are represented in purple ball-and-stick models, and the identified or potential B7 binding sites are highlighted in blue. **C** Dimer of Ctla-4 was identified by Western blot under reducing (+β-ME) or non-reducing (-β-ME) conditions. The ctrl represents a control sample derived from cells transfected with an empty plasmid. The monomers and dimers were indicated by single and double arrows, respectively. **D** The subcellular localization of Ctla-4 protein was assessed in HEK293T cells transfected with pEGFPN1-Ctla-4 for 48 hours, imaged using a two-photon laser-scanning microscope (Original magnification, 630×). Nuclei were stained with DAPI (blue), and cell membranes were stained with DiI (red). **E** UMAP plots showing the relative distribution of common T cell markers (*cd4-1*, *cd8a* and *ctla-4*) based on a splenic single-cell RNA sequencing (scRNA-seq) dataset we recently established [37]. **F** Immunofluorescence staining of lymphocytes isolated from zebrafish blood, spleen, and kidney. Cells were stained with mouse anti-Ctla-4, together with rabbit anti-Cd4-1 or rabbit anti-Cd8α. DAPI stain shows the location of the nuclei. Images were obtained using a two-photon laser-scanning microscope (Original magnification, 630×).

1050

Fig.2 Examination on the IBD-like phenotype in *ctla-4⁻/⁻* zebrafish. **A** Generation of a

homozygous *ctla-4*-deficient (*ctla-4⁻/⁻*) zebrafish line through CRISPR/Cas9-based

knockout of *ctla-4* gene on chromosome 9. A 14-bp deletion mutation in exon 2 results

in a premature stop at codon 82, which is predicted to produce a defective Ctla-4 protein

containing 81 amino acids. **B** Genotyping of the deficiency of *ctla-4* gene by Sanger

sequencing. **C** Knockout efficiency of Ctla-4 selectively examined in spleen and gut

tissues of *ctla-4⁻/⁻* zebrafish by Western blot analysis. Gapdh serves as a loading control.

D Normal gross appearance of adult wild-type (WT) and *ctla-4⁻/⁻* zebrafish. **E** Body

weight statistics of WT and *ctla-4⁻/⁻* zebrafish (n = 30). **F** The change of intestine length

in WT and *ctla-4⁻/⁻* zebrafish. **G** The change of splenic size in WT and *ctla-4⁻/⁻* zebrafish.

H Representative H & E staining analysis of histopathological changes and quantitation

of histology scores in the anterior, mid and posterior intestines from WT and *ctla-4⁻/⁻*

zebrafish. Red arrows denote mucosal inflammatory cell infiltration, and black arrow

indicates transmural inflammatory cell infiltration. **I** AB-PAS staining was used to

analyze the mucin components and the number of goblet cells in anterior intestine from

WT and *ctla-4⁻/⁻* zebrafish (n = 5). **J** Quantitation analysis of goblet cells of each villus

in the foregut of WT and *ctla-4⁻/⁻* zebrafish (n = 8). **K** Observation of cell junctions

between intestinal epithelial cells in posterior intestines from WT and *ctla-4⁻/⁻* zebrafish

under TEM (Hitachi Model H-7650). White triangles indicate tight junctions, black

triangle indicates adhesion junctions, and red triangles indicate desmosomes. Data are

presented as mean ± standard deviation (SD). Statistical significance was assessed

through an unpaired Student's t test ($^*p < 0.05$; $^{**}p < 0.01$; $^{***}p < 0.001$; $^{****}p < 0.0001$).

1073

Fig.3 RNA-sequencing analysis of the molecular implications associated with the IBD-

like phenotype in *ctla-4^-/-* zebrafish. **A** Heatmap of different expressed genes between

the intestines from wild-type (WT) and *ctla-4^-/-* zebrafish. **B** Volcano plot showing the

up-/down-regulated genes in the intestines of *ctla-4^-/-* zebrafish compared with those of

WT zebrafish. Red represents up-regulated genes, while blue denotes down-regulated

genes. **C** GO analysis showing top 10 terms in biological processes of DEGs. **D** GO

analysis showing top 10 terms in biological processes of all up-regulated genes. **E**

Heatmap showing row-scaled expression of the representative differently expressed

inflammation and chemotaxis-related genes. **F** The mRNA expression levels of

important genes associated with inflammation and chemokines confirmed by real-time

qPCR. **G** Protein-protein interaction network was constructed using the DEGs. The

nodes represent the proteins (genes); the edges represent the interaction of proteins

(genes). **H** The MPO activity in the intestines (up) and peripheral blood (down). **I**

Heatmap showing row-scaled expression of the IBD biomarker genes and IBD-related

genes. **J** The mRNA expression levels of representative IBD biomarker genes and IBD-

related genes were analyzed by real-time qPCR. Data are presented as mean ± standard

deviation (SD). Statistical significance was assessed through an unpaired Student's t

test ($^{**}p < 0.01$; $^{***}p < 0.001$; $^{****}p < 0.0001$).

Fig.4 Single-cell RNA sequencing analysis of the major cell types associated with the

IBD-like phenotype in *ctla-4⁻/⁻* zebrafish. **A** Classification of cell types from zebrafish

intestines by tSNE embedding. **B** Dot plot showing the expression levels of lineage

marker genes and percentage of cells per cluster that express the gene of interest. **C**

Expression maps of T cell associated markers within the cell populations of the zebrafish intestines. **D** Heatmap showing the mean expression levels of genes associated with tight and adhesion junctions in enterocytes across samples from wild-type (WT) and *ctla-4*$^{-/-}$ zebrafish. **E** Heatmap showing the mean expression levels of inflammation-related genes involved in cytokine-cytokine receptor interactions in neutrophils from WT and *ctla-4*$^{-/-}$ zebrafish samples. **F** KEGG enrichment analysis showing the top 15 terms of up-regulated genes in neutrophils from the *ctla-4*$^{-/-}$ sample versus the WT sample. **G** KEGG enrichment analysis showing the top 15 terms of up-regulated genes in macrophages from the *ctla-4*$^{-/-}$ sample versus the WT sample.

1106

1107 **Fig.5** Single-cell RNA sequencing analysis of the subset immune-cells associated with

the IBD-like phenotype in *ctla-4*[-/-] zebrafish. **A** Classification of subset cells from the T/NK/ILC-like group by tSNE embedding. **B** Dot plot showing the mean expression levels of subset marker genes and percentage of cells per cluster that express the gene of interest. **C** Marker gene expression in individual cluster identifying this cluster as ILC3-like cells. **D** Changes in the composition of subset cells between samples from wild-type (WT) and *ctla-4*[-/-] zebrafish. A significantly increased Th2 subset (referred to as Th2 cells 2) in the *ctla-4*[-/-] sample was highlighted with a black dashed circle. **E** Histogram showing the different ratios of subset cells between the WT and *ctla-4*[-/-] samples. **F** Histogram presenting the different numbers of subset cells between the WT and *ctla-4*[-/-] samples. **G** Mean expression levels of the cytokine *il13* within different subset cells between the WT and *ctla-4*[-/-] samples. **H** Dot plot illustrating the mean expression of *il13* in T/NK/ILC-like cells from WT and *ctla-4*[-/-] zebrafish. **I** KEGG enrichment analysis showing the top 15 terms of the Th2 cells 2 genes from *ctla-4*[-/-] zebrafish. **J** KEGG enrichment analysis showing the top 15 terms of up-regulated genes in NKT-like cells. **K** Scatter plot showing the DEGs of ILC3-like cells in WT and *ctla-4*[-/-] zebrafish. The *il17a/f1* and *il17a/f3* was shown in the scatter plot.

1124

Fig.6 Alteration in microbial composition in the intestines of *ctla-4⁻/⁻* zebrafish. **A** Venn

diagram showing the number of ASVs in zebrafish intestinal microbiota. **B** Alpha-

diversity of microbes was calculated through Shannon index and Simpson index. **C**

Beta-diversity analyzed based on PCoA was shown by using Bray Curtis distance. **D**

The relative abundance of intestinal microbiota at the class level. **E-G** The relative

abundance of Alphaproteobacteria (**E**), Bacilli (**F**) and verrucomicrobiae (**G**) in the

intestines from the wild-type (WT) and *ctla-4⁻/⁻* zebrafish. *$p < 0.05$. **H** The relative

abundance of intestinal microbiota at the family level. **I** Heatmap showing row-scaled

expression of the differential abundances of bacterial communities at family level in

the WT and *ctla-4⁻/⁻* zebrafish ($p < 0.05$). **J** Cladogram representation of LEfSe analysis

showing the differentially abundant bacterial taxa between the intestines from WT (red)

 and *ctla-4⁻ᐟ⁻* (green) zebrafish ($p < 0.05$).

1137

1138

Fig.7 Examination on the inhibitory function of Ctla-4 in T cell activation. **A** Assessment of the proliferative activity of T cells from wild-type (WT) and *ctla-4⁻ᐟ⁻* zebrafish by a mixed lymphocyte reaction combined with PHA-stimulation. The CFSE dilution, which served as an indicator of lymphocyte proliferation, was measured through flow cytometry. **B** Assessment of the proliferative activity of lymphocytes from *ctla-4⁻ᐟ⁻* zebrafish by the administration of sCtla-4-Ig. **C** Assessment of the proliferative activity of lymphocytes from WT zebrafish by supplementing anti-Ctla-4 antibody. **D** Assessment of the proliferative activity of lymphocytes from *ctla-4⁻ᐟ⁻* zebrafish by the administration of sCd28-4-Ig. **E** Assessment of the proliferative activity of lymphocytes

from *ctla-4*[-/-] zebrafish by the administration of recombinant sCd80/86 protein. **F, G**

Interactions between Cd80/86 and Cd28 (**F**), and Cd80/86 and Ctla-4 (**G**) as predicted

by AlphaFold2. On the left are structural models depicting Cd80/86 in complex with

Cd28 or Ctla-4. The center panels display per-residue model confidence scores (pLDDT)

for each structure, using a color gradient from 0 to 100, where higher scores indicate

increased confidence. The right panels show the predicated aligned error (PAE) scores

for each model. The well-defined interfaces between Cd28 or Ctla-4 and Cd80/86 are

highlighted with red dashed squares. **H** The interaction between Cd80/86 and Cd28

(left), and Cd80/86 and Ctla-4 (right) were verified by Co-IP. **I** Binding affinities of the

Cd80/86 protein for the Cd28 and Ctla-4 proteins were measured by the microscale

thermophoresis (MST) assay. The K_D values are provided. Data are presented as mean

± standard deviation (SD), which were calculated from three independent experiments.

Statistical significance was assessed through an unpaired Student's t test ([**]$p < 0.01$;

[***]$p < 0.001$; ns denotes no statistical significance).

1162

Fig.8 In vivo inhibition of intestinal inflammation by sCtla-4-Ig. **A** Percent initial

weight of zebrafish after injection of the sCtla-4-Ig or the IgG isotype control. Each

group consisted of six zebrafish (n = 6). Data show means with SEM analyzed by two-

way ANOVA with Sidak's correction for multiple comparisons. **B** Representative H&E

staining analysis of histopathological changes and quantitation of histology scores in

the posterior intestine from *ctla-4*[-/-] zebrafish treated with sCtla-4-Ig or IgG isotype

control. Scale bar: 50 μm. **C** The mRNA expression levels of inflammation-related

genes in *ctla-4*[-/-] zebrafish treated with sCtla-4-Ig or IgG isotype control. **D** The mRNA

expression levels of IBD biomarker genes and IBD-related genes in *ctla-4*[-/-] zebrafish

treated with sCtla-4-Ig or IgG isotype control. The *p* value was generated by unpaired

two-tailed Student's *t-test*. **$^{**}p < 0.01$; $^{***}p < 0.001$; $^{****}p < 0.0001$.

1174

Supplemental Figures

Fig. S1 The organization, sequence and phylogenetic analysis of zebrafish *ctla-4* and

cd28 genes. **A** Comparison of the intron/exon organizations of *ctla-4* gene in zebrafish

and humans. Exons and introns are shown with black boxes and lines, and their size are

indicated by the numbers found above and below the sequences respectively. **B** The

nucleotide and amino acid sequences of *ctla-4* gene and Ctla-4 protein. The underline

indicates the signal peptide, the circles represent the conserved cysteine residues. **C**

Phylogenetic analysis of the relationship of Ctla-4 and Cd28 between zebrafish and

other species. An unrooted phylogenetic tree was constructed through the neighbor-

joining method, based on amino acid sequence alignments generated by ClustalX.

Bootstrap confidence values, derived from 500 replicates, are indicated as percentages

at each node. **D** Alignment of the Cd28 homologs from different species generated with

ClustalX and Jalview. The conserved and partially conserved amino acid residues in

each species are colored in hues graded from orange to red, respectively. The conserved

functional motifs, such as B7-binding motif, tyrosine phosphorylation site, and

potential tyrosine phosphorylation site, were indicated separately. The signal peptide,

IgV-like domain, transmembrane (TM) domain and cytoplasmic domain were marked

at the top of the sequence.

1194

1195

Fig. S2 Preparation of mouse anti-Ctla-4 antibody. **A** SDS-PAGE detection of the

recombinant Ctla-4 protein with extracellular domain (ECD). Lane 1, 2 and 3 represent

the protein markers, blank, and target protein, respectively. **B** Western blot analysis of

the mouse anti-EGFP and anti-Ctla-4 antibodies that bind to the recombinant Ctla-4-

EGFP fusion proteins expressed in HEK293T cells. **C** Western blot analysis of native

Ctla-4 protein in zebrafish intestinal tissues using mouse anti-Ctla-4 antibody.

1202

1203

Fig. S3 Histopathological analysis of intestines. **A** Periodic Acid-Schiff (PAS) staining

was used to analyze the mucin components in anterior intestine from wild-type (WT)

and *ctla-4⁻/⁻* zebrafish (n = 5). **B** The ratio of intestinal villi length to intestinal ring

radius was measured in the anterior, mid, and posterior intestines of WT and *ctla-4⁻/⁻*

zebrafish (n = 6). Statistical significance was assessed through an unpaired Student's t

test (*$p < 0.05$; **$p < 0.01$).

1210

1211

Fig. S4 Examination on the functional genes and pathways associated with the IBD-

like phenotype in *ctla-4*⁻/⁻ zebrafish. **A** Top 5 KEGG enrichment bar plot of up-regulated

genes in *ctla-4*⁻/⁻ zebrafish intestines versus wild-type (WT) zebrafish intestines. **B** Top

10 KEGG enrichment bar plot of down-regulated genes in *ctla-4*⁻/⁻ zebrafish intestines

versus WT zebrafish intestines. **C, D** Changes in the expression of genes associated

with lymphocyte chemotaxis, positive regulation of ERK1/ERK2 cascades, Calcium

and MAPK signaling pathways in the *ctla-4*⁻/⁻ zebrafish intestines analyzed by using a

collection of pre-defined gene sets retrieved from GO (**C**) and KEGG (**D**) database. The

p value, false discovery rates (FDR) and normalized enrichment score (NES) are shown

above each pathway graph.

1222

1223

Fig. S5 Quality control analysis of single-cell RNA sequencing data. **A** Identification

of the effective cell number of the sample. The blue line represents the effective cells

corresponding to barcodes, while the gray line denotes the background noise. **B** The

basic cellular metrics before and after filtering, including the total number of detected

genes (nFeature_RNA), the total number of UMIs (nCount_RNA), and the percentage

of reads mapped to mitochondrial genes (Percent.mito). **C** The scatter plot comparing

the cellular metrics before and after filtering, showing the relationship between

nCount_RNA and nFeature_RNA. The Pearson correlation coefficients are indicated

above the graph.

Fig. S6 Examination on the involvement of apoptotic process in epithelial cells and

expression of inflammation-related genes in neutrophils and B cells in the intestines of

ctla-4⁻/⁻ zebrafish. **A** Expression map of the epithelial markers within the cell

populations of the zebrafish intestines. **B** Expression map of the neutrophil markers

within the cell populations of the zebrafish intestines. **C** KEGG enrichment bar plot of

all differentially expressed genes (DEGs) from epithelial cells. **D** Quantification of

TUNEL-positive cells per 1×10^4 μm^2 in WT and *ctla-4⁻/⁻* posterior intestines (n = 5).

E Heatmap of inflammation-related genes in B cells from wild-type and *ctla-4⁻/⁻*

intestines. Statistical significance was assessed through an unpaired Student's t test (***p

< 0.001).

1244

1245

Fig. S7 Examination on the activation of T cell subsets in the intestines of *ctla-4*[-/-] zebrafish. **A** Marker gene expression in individual cluster identifies the cluster as NKT-like cells. **B** RT-qPCR confirms the mRNA expression levels of Th2 cell marker genes in the intestines of wild-type (WT) and *ctla-4*[-/-] zebrafish. **C** RT-qPCR validates the mRNA expression levels of NKT-like cell marker genes in the intestines of WT and *ctla-4*[-/-] zebrafish. **D** Heatmap illustrates up-regulated genes involved in cytokine-cytokine receptor interaction in NKT-like cells from WT and *ctla-4*[-/-] samples. **E** KEGG enrichment analysis reveals the top 15 terms of up-regulated genes in Cd8[+] T cells in *ctla-4*[-/-] samples versus WT samples. **F** Heatmap displays up-regulated genes involved in cytokine-cytokine receptor interaction in Cd8[+] T cells from WT and *ctla-4*[-/-] samples. Statistical significance was assessed through an unpaired Student's t test (*$p < 0.05$; **$p <0.01$; ***$p < 0.001$; ****$p < 0.0001$).

Fig. S8 Preparation of recombinant proteins and examination of their molecular

interactions. **A-C** SDS-PAGE detection of the purified recombinant soluble Ctla-4-Ig

(sCtla-4) (**A**) and sCd28-Ig (**B**) proteins and the Cd80/86 extracellular domain (ECD)

(**C**) with Coomassie brilliant blue staining. **D-E** The predicted molecular interactions

between Cd80/86 and Cd28 (D), as well as Cd80/86 and Ctla-4 (E), as modeled by

AlphaFold2. The structures are represented in a cartoon style, with Cd80/86, Cd28, and

Ctla-4 colored cyan, green, and magenta, respectively. A total of 25 models were

predicted for each complex and aligned with Cd80/86. **F** Flow cytometry analysis of

the interactions between Cd80/86 and Cd28 (top), and Cd80/86 and Ctla-4 (bottom).

Cd80/86 was expressed on HEK293T cells and incubated with varying concentrations

of fluorescently labeled sCd28-Ig or sCtla-4-Ig. Fluorescence intensity was detected by

flow cytometry to determine molecular interactions. Data are presented as mean $\pm$ SD,

derived from three independent experiments. Statistical significance was evaluated

using an unpaired Student's t test ($^{**}p < 0.01$; $^{***}p < 0.001$).