Unraveling the Role of Ctla-4 in Intestinal Immune Homeostasis: Insights from a novel Zebrafish Model of Inflammatory Bowel Disease

Lulu Qin[a], Chongbin Hu[a], Qiong Zhao[a], Yong Wang[a], Dongdong Fan[a], Aifu Lin[a], Lixin Xiang[a*], Ye Chen[a,c*], Jianzhong Shao[a,b*]

Affiliations:

[a]College of Life Sciences, Key Laboratory for Cell and Gene Engineering of Zhejiang Province, Zhejiang University, Hangzhou, China.

[b]Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, Qingdao, China

[c]Department of Genetic and Metabolic Disease, the Children's Hospital, Zhejiang University School of Medicine, National Clinical Research Center for Child Health, Hangzhou 310052, China.

[*]Corresponding authors: shaojz@zju.edu.cn, yechency@zju.edu.cn, and xianglx@zju.edu.cn

Address correspondence and reprint requests to Prof. Jianzhong Shao, Prof. Ye Chen, and Assoc. Prof. Lixin Xiang, College of Life Sciences, Zhejiang University, 866 YuHangTang Road, Hangzhou 310058, China. Tel.: +86 (571) 88206582; Fax: +86 (571) 88206582.

E-mail addresses: shaojz@zju.edu.cn (J.-z.S.), yechency@zju.edu.cn (Y.C), and xianglx@zju.edu.cn (L.-x.X.)

Abstract

Inflammatory bowel disease (IBD) is a chronic and relapsing immune-mediated disorder characterized by intestinal inflammation and epithelial injury. The underlying causes of IBD are not fully understood, but genetic factors have implicated in genome-wide association studies, including CTLA-4, an essential negative regulator of T cell activation. However, establishing a direct link between CTLA-4 and IBD has been challenging due to the early lethality of CTLA-4 knockout mice. In this study, we identified zebrafish Ctla-4 homolog and investigated its role in maintaining intestinal immune homeostasis by generating a Ctla-4-deficient ($ctla\text{-}4^{-/-}$) zebrafish line. These mutant zebrafish exhibit reduced weight, along with impaired epithelial barrier integrity and lymphocytic infiltration in their intestines. Transcriptomics analysis revealed upregulation of inflammation-related genes, disturbing immune system homeostasis. Moreover, single-cell RNA-sequencing analysis indicated increased Th2 cells and interleukin 13 expression, along with decreased innate lymphoid cells and upregulated proinflammatory cytokines. Additionally, Ctla-4-deficient zebrafish exhibited reduced diversity and an altered composition of the intestinal microbiota. All these phenotypes closely resemble those found in mammalian IBD. Lastly, supplementation with Ctla-4-Ig successfully alleviated intestinal inflammation in these mutants. Altogether, our findings demonstrate the pivotal role of Ctla-4 in maintaining intestinal homeostasis. Additionally, they offer substantial evidence linking CTLA-4 to IBD and establish a novel zebrafish model for investigating both the pathogenesis and potential treatments.

Key words: Ctla-4 deficiency; inflammatory bowel disease; molecular and cellular

mechanisms; zebrafish model

Introduction

Inflammatory bowel disease (IBD), including Crohn's disease and ulcerative

colitis, refers to a group of chronic relapsing inflammation disorders affecting the

gastrointestinal tract, that have been increasing in prevalence worldwide [1]. The

precise etiology of IBD has yet to be fully elucidated. Conventional epidemiological

studies have indicated that IBD tends to run in families and is linked to genetic factors

[2, 3]. However, research also suggested that susceptibility gene patterns differ

significantly among various geographic populations. Current evidence points towards

a complicated interaction involving host genetics, disrupted intestinal microbiota,

environmental triggers, and abnormal immune responses [4-6]. Advances in genomic

sequencing techniques have allowed for the identification of genetic variants associated

with an increased risk of developing IBD. Among these, mutations in immune-related

genes have received particular attention. Research on humans with Crohn's disease and

mouse models of IBD has shown that genetically susceptible individuals exhibit defects

in intracellular pattern-recognition receptors (PRRS), such as toll-like receptors (TLR)

and nucleotide-binding oligomerization domain (NOD)-like receptors (NLRs), which

are responsible for initiating innate immune responses to eliminate harmful bacteria [7,

8]. Genetic variations in the tumor necrosis factor ligand superfamily member 15

(TNFSF15) and interleukin 23 receptor (IL23R) genes, both involved in suppressing

inflammation, have been associated with an increased risk of developing Crohn's

disease [9, 10].

Cytotoxic T-cell associated antigen-4 (CTLA-4), also known as CD152, is one of

the most well-established immune checkpoint molecules expressed predominantly on

T cells. It primarily regulates the early stages of T-cell activation by attenuating

downstream signaling of the T cell receptor (TCR) [11-13]. Specifically, CTLA-4 has

a much higher affinity for CD80 and CD86 ligands compared to the co-stimulatory

receptor CD28 [14, 15]. By outcompeting CD28 for ligand binding, CTLA-4 provides

an inhibitory signal that impacts immunological synapse formation and inhibits T-cell

proliferation and activation [13, 16]. As expected, links between polymorphic alleles of

CTLA4 and IBD have been reported in multiple studies [17-19]. Moreover, CTLA-4 is

an intriguing target for novel immune checkpoint blockade therapies in cancer

treatment, while intestinal inflammation is a common side effect in these clinical trials

[20, 21]. Establishing a direct causal relationship between CTLA-4 and IBD has been

challenging due to difficulties in finding appropriate models. Early lethality was

observed in CTLA-4-deficient mice, adding to the complexity of this issue. Zebrafish

is a powerful model system for immunological and biomedical research, due to its

versatility and high degree of conservation in innate and adaptive immunities. In our

current study, we identified the Ctla-4 homology in zebrafish and successfully

developed an adult vertebrate model with homozygous knockout of the *ctla-4* gene for

the first time. These *ctla-4*-deficient (*ctla-4$^{-/-}$*) zebrafish survive but exhibit attenuated

growth and weight loss. Notably, *ctla-4* deficiency leads to an IBD-like phenotype in

zebrafish characterized by altered intestinal epithelial cells morphology, abnormal

inflammatory response, defects in microbial stratification and composition.

Mechanistically, Ctla-4 exerts its inhibitory function by competing with Cd28 for

binding to Cd80/86. These findings establish the *ctla-4* knockout zebrafish as an

innovative platform to elucidate CTLA-4 immunobiology, model human IBD, and

develop novel therapeutic modalities.

Results

Identification of zebrafish Ctla-4

Through a homology search in the NCBI database, we identified the *ctla-4* gene

on zebrafish chromosome 9 (Fig. S1A-C). Zebrafish Ctla-4 protein contains an N-

terminal signal peptide, a single IgV-like extracellular domain, a transmembrane region,

and a cytoplasmic tail (Fig. 1A). A ^{113}LFPPPY^{118} motif is present in the ectodomain of

Ctla-4, similar to the MYPPPY motif, a binding site for CD80/CD86 found in mammals

(Fig. 1A). Thus, we reasoned that the ^{13}LFPPPY^{118} motif may be a potential binding

site for Cd80/86 in zebrafish. The tyrosine-based YVKM motif in the cytoplasmic

domain in mammals, which is involved in internalization and signaling inhibition, is

absent in Ctla-4. By contrast, a unique tyrosine-based ^{206}YVKF^{209} motif was found in

the C-terminus of Ctla-4 molecule, which is relatively conserved in different fish

species (Fig. 1A). The IgV-like domain of Ctla-4 was characterized by two-layer β-

sandwich and was conserved between zebrafish and humans (Fig. 1B). Ctla-4 exists as

a dimer, and unlike the intracellular localization of CTLA-4 in mammals, Ctla-4 was

located on the cell membrane (Fig. 1C and D). Moreover, Ctla-4 was mainly expressed

on the T cells, including the Cd4$^+$ T cells and Cd8$^+$ T cells (Fig. 1E and F).

Ctla-4 deficiency induces inflammatory bowel disease (IBD)-like phenotype

To further investigate the function of Ctla-4, we generated a *ctla-4*$^{-/-}$ zebrafish line

with a 14-base deletion in the second exon of the *ctla-4* gene (Fig. 2A-C). The zebrafish

appeared grossly normal in appearance; however, the body weight and size were

significantly reduced compared with those of wild-type zebrafish (Fig. 2D and E).

Anatomically, the *ctla-4*$^{-/-}$ zebrafish were featured by intestine shortening and

splenomegaly, suggesting the occurrence of chronic inflammation in the intestines (Fig.

2F and G). For clarification, we first performed histological analysis on the anterior,

mid, and posterior intestine segments using H&E staining. Compared with the wild-

type zebrafish, the *ctla-4*$^{-/-}$ fish exhibited significant epithelial hyperplasia in the

anterior intestine segment, accompanied with a small amount of mucosal inflammatory

cell infiltration (Fig. 2H). This result indicates an enhanced regeneration of the intestine,

which is typically observed in cases of intestinal inflammation [22]. A noteworthy

goblet cell loss which marks the severity of intestinal inflammation was also observed

in *ctla-4*$^{-/-}$ anterior intestine, as quantified through PAS staining (Fig. 2I and J). Besides,

a small amount of lymphocytic infiltration and mild epithelial damage occurred in the

mid intestine segment of *ctla-4*$^{-/-}$ zebrafish (Fig. 2H). In posterior intestine of *ctla-4*$^{-/-}$

fish, the intestinal villi were markedly shortened, the epithelial barrier showed severely

disrupted, and the intestinal wall became thinner, wherein the mucosal and transmural

inflammatory cells were significantly infiltrated (Fig. 2H). Consistent with these

observations, ultrastructure analysis revealed that the epithelial cells of posterior intestine in *ctla-4*$^{-/-}$ zebrafish exhibited alteration in tight junction, the loss of adhesion junctions and desmosomes, and disruption of microvilli (Fig. 2K). All these results strongly indicate that Ctla-4 plays a crucial role in preserving intestinal homeostasis in zebrafish. The intestinal phenotype resulting from Ctla-4 deficiency was similar to IBD in mammals.

Moleculear mechanisms of Ctla-4 deficiency-induced IBD-like phenotype

To explore the potential molecular mechanisms of Ctla-4 deficiency-induced IBD-like phenotype, we performed transcriptome profiling analysis of intestines from wild-type and *ctla-4*$^{-/-}$ zebrafish. We identified a total of 1,140 differentially expressed genes (DEGs), among which 714 genes were up-regulated, and 426 genes were down-regulated in *ctla-4*$^{-/-}$ zebrafish (Fig. 3A and B). GO enrichment analysis showed that DEGs or up-regulated genes in the top 10 enriched biological processes were associated with the immune response and inflammatory response (Fig. 3C and D). Moreover, the KEGG enrichment analyses indicated that the up-regulated DEGs are primarily involved in the process of cytokine-cytokine receptor interaction and cell adhesion molecules, which are also related to inflammation (Fig. S3A); however, the down-regulated DEGs were significantly enriched in the process of metabolism (Fig. S3B). The intestines of *ctla-4*$^{-/-}$ zebrafish showed significant upregulation of *il1b*, *tnfa*, myeloid-specific peroxidase (*mpx*), matrix metallopeptidase 9 (*mmp9*), chemokine (C-X-C motif) ligand 8a (*cxcl8a*), and *il13*, a Th2-typic cytokine crucial for ulcerative colitis in mammals [23]. In contrast, *il10*, a potent anti-inflammatory cytokine, was

markedly down-regulated in Ctla-4-deficient intestines (Fig. 3E). The transcriptional change of these genes was confirmed by RT-qPCR (Fig. 3F). By constructing the PPI network, we found that *il1b* was a major cytokine that played a hub role in promoting the bowel inflammation of *ctla-4*$^{-/-}$ zebrafish (Fig. 3G). Moreover, Gene set enrichment analysis (GSEA) showed that genes involved in the lymphocyte chemotaxis, positive regulation of ERK1 and ERK2 cascade, Calcium and MAPK signaling pathways were positively enriched in *ctla-4*$^{-/-}$ zebrafish intestines, implying a sensitized or activated state of lymphocytes due to the absence of Ctla-4 (Fig. S3C and D). Notably, biological processes related to neutrophil activation and chemotaxis were significantly enriched (Fig. 3C and D). Studies have shown that neutrophils can induce histopathological effects through releasing matrix metalloproteinases (MMPs), neutrophil elastase, and myeloperoxidase (MPO) [24]. To confirm the association between neutrophils and Ctla-4-deficient intestinal inflammation, the MPO level was examined. As a support, MPO activity was markedly increased in the intestines and peripheral blood of *ctla-4*$^{-/-}$ zebrafish (Fig. 3H). Besides, a number of biological markers or susceptibility genes of IBD observed in mammals, including c-reactive protein 6 (*crp6*), *crp7*, MMPs, haptoglobin, *il23r*, insulin-like growth factor binding protein 1 a (*igfbp1a*), cAMP responsive element modulator b (*cremb*) and lymphocyte specific protein 1 b (*lsp1b*), were highly expressed in the *ctla-4*$^{-/-}$ zebrafish (Fig. 3I and J) [9, 25, 26], suggesting the presence of a conserved molecular network underlying IBD pathogenesis between *ctla-4*$^{-/-}$ zebrafish and mammalian models.

Cellular mechanisms of Ctla-4 deficiency-induced IBD-like phenotype

To investigate the cellular mechanisms underlying the IBD-like phenotype

induced by Ctla-4 deficiency, we performed scRNA-seq analysis on intestines from

wild-type and *ctla-4*$^{-/-}$ zebrafish using the 10× Genomics Chromium platform. We

obtained nine discrete clusters from 7,539 cells of wild-type and *ctla-4*$^{-/-}$ zebrafish (Fig.

4A). These clusters of cells were classified as enterocytes, enteroendocrine cells,

smooth muscle cells, neutrophils, macrophages, B cells, and a group of T/NK/ILC-like

cells based on their co-expression of lineage marker genes (Fig. 4B and C, Fig. S4A

and B). Due to severe epithelial disruption and inflammatory cell infiltration in *ctla-4*$^{-}$

$^{/-}$ zebrafish intestines, we focus on the pathological process and immune reactions in

enterocytes and immune cell populations. KEGG analysis showed that apoptotic

pathway was highly enriched in enterocytes of *ctla4*$^{-/-}$ zebrafish, suggesting that

aberrant apoptosis contributes to the epithelial injury (Fig. S4C). Additionally, genes

functionally involved in the formation of tight and adhesion junctions, such as *oclna*,

cdh1, *pcdh1b* and *cldn15a*, were significantly down-regulated in enterocytes of *ctla-4*$^{-}$

$^{/-}$ zebrafish (Fig. 4D), consistent with the pathological observation under electron

microscope. Furthermore, inflammation-related genes and pathways were significantly

up-regulated and enriched in neutrophils, B cells, and macrophages of *ctla-4*$^{-/-}$ zebrafish,

suggesting active inflammatory responses (Fig. 4E-G, Fig. S4D). By sub-clustering

analysis, six subpopulations were classified from T/NK/ILC-like cell groups based on

their expression with a set of marker genes. These subpopulations include Cd8^{+} T cells,

ILC3-like cells, maturing Ccr7^{high} T cells, NKT cells, and two groups of Th2 cells (Fig.

5A-C; Fig. S5A). The abundances of NKT and two subsets of Th2 cells were

significantly increased in the intestines of *ctla-4*$^{-/-}$ zebrafish (Fig. 5D-F). These cells exhibited high expression of *il13*, a key contributor to intestinal inflammation in mammals (Fig. 5G and H) [23]. Specifically, the second subset of Th2 cells was seldom observed in the intestine of wild-type zebrafish, indicating their unique role in the pathogenesis of IBD-like phenotype in *ctla-4*$^{-/-}$ zebrafish (Fig. 5D-F). KEGG analysis of up-regulated genes from *ctla-4*$^{-/-}$ NKT and Th2 cells indicated that Ctla-4 deficiency is positively associated with the inflammatory cytokine-cytokine receptor interaction, PPAR, calcium and MAPK signaling pathways, cellular adhesion and mucosal immune responses (Fig. 5I and J; Fig. S5B), suggesting high activity of NKT and Th2 cells in inflamed intestines [27, 28]. Although the abundance of Cd8$^+$ T cells was not significantly changed in Ctla-4-deficient intestines, the inflammatory genes and pathways were up-regulated and enriched in the subset of T cells (Fig. S5C and D). Notably, the proportion of ILC3-like cells was decreased, and they highly expressed *il17a/f1* and *il17a/f3* in the Ctla-4-deficient intestines (Fig. 5D-F, and K). Investigations have consistently reported a substantial decline in the population of ILC3s within the inflamed intestines and IL-17A-secreting ILC3s play a significant role in the development of intestinal inflammation [29-34]. Thus, the reduced ILC3-like cells and increased expression of *il17a/f1* and *il17a/f3* may be responsible for intestinal inflammation induced by Ctla-4 deficiency.

Decreased microbiota diversity in *ctla-4*$^{-/-}$ zebrafish intestines

The intestinal microbiota plays a crucial role in host functions such as nutrient acquisition, metabolism, epithelial cell development and immunity. Notably, lower

microbiota diversity has consistently been observed in patients with IBD phenotype [35, 36], making it a valuable indicator of host health. Therefore, we further analyzed whether microbes are involved in the Ctla-4-deficiency induced intestinal inflammation. The results revealed a significantly higher number of amplicon sequence variants (ASVs) in wild-type zebrafish intestines, with 730 ASVs unique to the wild-type group and 276 ASVs exclusively found in *ctla-4⁻ᐟ⁻* group (Fig. 6A). Furthermore, the Shannon index and the Simpson index indicated a decreased microbial diversity in *ctla-4⁻ᐟ⁻* zebrafish intestines (Fig. 6B) and the Principal Coordinate Analysis (PCoA) using Bray Curtis distance revealed a significant separation in the microbial composition between *ctla-4⁻ᐟ⁻* group and the wild-type group (Fig. 6C). To gain insights into the microbial community composition, we analyzed the identified microbial populations at the class and family level. Alphaproteobacteria and Gammaproteobacteria were found to be the most prevalent classes. Relative to wild-type group, Ctla-4 deficiency resulted in a significant reduction in Alphaproteobacteria abundance. However, the Gammaproteobacteria, one of the main classes of Gamma-negative pathogenic bacteria expanded under inflammation conditions, was increased, although the change did not reach statistical significance (Fig. 6D and E) [37]. In addition, we observed a decreased relative abundance of short-chain fatty acids (SCFAs)-producing Bacilli and Verrucomicrobiae, the latter of which contributes to glucose homeostasis and intestinal health (Fig. 6F and G) [38, 39]. Notably, the family-level analysis revealed a notable enrichment of Enterobacteriaceae, overgrowing under host inflammatory conditions, and the Shewanellaceae, serving as the most important secondary or opportunistic

pathogens, in *ctla-4*^{-/-} zebrafish (Fig. 6H and I). To identify differentially abundant

bacterial taxa between the wild-type and *ctla-4*^{-/-} zebrafish, we conducted Linear

discriminant analysis (LDA) effect size (LEfSe). The results showed that several

potentially opportunistic pathogens, including Gammaproteobacteria, Enterobacterales,

and Aeromonadales were found to be overrepresented in *ctla-4*^{-/-} zebrafish (Fig. 6J). In

contrast, Actinobacteriota, Cetobacterium, and Planctomycetota (Planctomycetes) were

more abundant in the wild-type zebrafish. These findings strongly indicated an

association between Ctla-4 deficiency-induced gut inflammation and dysbiosis, as

characterized by decreased microbial diversity, loss of potentially beneficial bacteria,

and expansion of pathobionts.

Inhibitory role of Ctla-4 in T cell activation

Given that Ctla-4 is primarily expressed on T cells (Fig. 1E-F), and its absence has

been shown to result in intestinal immune dysregulation, indicating a crucial role of this

molecule as a conserved immune checkpoint in T cell inhibition. For clarification, the

lymphocyte proliferation was examined by a series of blockade/inhibition assays using

anti-Ctla-4 Ab, recombinant soluble Ctla-4-Ig (sCtla-4-Ig), sCd28-Ig, and sCd80/86

proteins in a PHA-stimulating and mixed lymphocyte reaction (MLR) model (Fig. S6

A-C). Results showed that the lymphocyte proliferation was more pronounced in the

absence of Ctla-4, and the supplementation of sCtla-4-Ig inhibited cell proliferation,

suggesting the inhibitory function of Ctla-4 in T cells (Fig. 7A and B). Consistent with

these results, blockade of Ctla-4 by administering anti-Ctla-4 Ab significantly promoted

the proliferation of lymphocytes from wild-type zebrafish (Fig. 7C). Moreover,

administration of sCd28-Ig into the co-cultures also inhibited the proliferation of lymphocytes from *ctla-4⁻/⁻* zebrafish (Fig. 7D). Besides, the addition of Cd80/86 significantly promoted the expansion of Ctla-4-deficient lymphocytes (Fig. 7E). Based on these results, we concluded that the presence of Ctla-4 obstructs the Cd80/86-Cd28 mediated costimulatory signaling, consequently impeding cell proliferation. To further investigate the relationship between Cd28, Ctla-4 and Cd80/86, we assessed the interaction between these molecules *in vitro*. Our bioinformatics analysis by AlphaFold2, flow cytometry and Co-IP assays demonstrated the interaction between Cd28/Ctla-4 and Cd80/86 (Fig. 7F-H, Fig. S6D). Moreover, using a microscale thermophoresis assay, we found that Ctla-4 has higher affinity for Cd80/86 than Cd28, as indicated by a lower equilibrium association constant value ($K_D = 0.50 \pm 0.25$ μM vs. $K_D = 2.64 \pm 0.45$ μM) (Fig. 7I). These results suggested that Ctla-4 could exert its inhibitory function by competing with Cd28 for binding Cd80/86.

sCtla-4-Ig alleviates IBD-like phenotype

As described above, engagement of Cd80/86 by sCtla-4-Ig effectively suppressed T cell activation *in vitro* (Fig. 7B), indicating that sCtla-4-Ig holds promise as a potential intervention for IBD-like phenotype. This is supported by the observation that zebrafish treated with sCtla-4-Ig exhibited obvious body weight restoration compared to those treated with the IgG isotype control (Fig. 8A). To provide further evidence, histological analysis was performed on the posterior intestine, which is known to experience severe tissue destruction under Ctla-4 deficient conditions. As expected, Ctla-4-Ig treatment resulted in a significant decrease in lymphocyte infiltration and an

improvement in the epithelial barrier (Fig. 8B). Moreover, Ctla-4-Ig treatment

significantly reduced the expression of pro-inflammatory genes, including *il13*, *tnfa*,

mpx, *mmp9*, and *cxcl8a*, as well as *igfbp1a*, *cremb*, and *lsp1a*, which are susceptibility

genes for IBD observed in mammals (Fig. 8C and D). These findings demonstrate that

the supplementation of Ctla-4-Ig alleviates intestinal inflammation in Ctla-4-deficient

zebrafish, highlighting its potential as a therapeutic intervention for CTLA-4

deficiency-induced IBD in mammals.

294 Discussion

As an essential negative regulator of T cell activation, dysfunction of CTLA-4 was

implicated in various diseases in both humans and murine models [40-42]. Numerous

previous studies have established the connection between CTLA-4 and autoimmune

thyroiditis, Graves' disease, myocarditis, pancreatitis, multiple sclerosis, rheumatoid

arthritis, and type I diabetes [43-49]. However, the involvement of CTLA-4 in IBD has

been understudied. Several investigations have reported that haploinsufficiency

resulting from mutations in CTLA-4 in humans is associated with IBD, and genome-

wide association studies (GWAS) have implicated CTLA-4 as a susceptibility gene for

IBD [50-52]. Nevertheless, the exact contributions and mechanisms of CTLA-4

deficiency in the occurrence and pathology of IBD remain incompletely understood,

primarily due to the lack of animal models attributable to the lethality of CTLA-4

knockout in mice. In this study, we identified the Ctla-4 homolog in zebrafish, and

discovered that defect in Ctla-4 did not have a severe lethal effect, but did show a clear

IBD-like phenotype. This makes zebrafish an attractive animal model for investigating the molecular and cellular mechanisms underlying Ctla-4 mediated IBD. Multiple lines of experimental evidence demonstrated the IBD-like phenotype in Ctla-4-deficienct zebrafish. These include the observed destruction of epithelial integrity and infiltration of inflammatory cells in the inflamed intestines during histopathological analysis, the enrichment of inflammation-related pathways and the imbalance between pro-inflammatory and anti-inflammatory cytokines identified through transcriptome profiling, the abnormal composition of immune cell populations revealed by single-cell RNA sequencing, as well as the reduced diversity and altered composition of intestinal microbiota. These observations collectively suggest that Ctla-4 may serve as a potential genetic determinant of the IBD-like phenotype, although further research is required to definitively identify the causative variant responsible for this association. Moreover, the establishment of zebrafish models provides a valuable tool for comprehending the underlying mechanisms of the disease's pathophysiology.

A multi-omics study was conducted to investigate the mechanisms of Ctla-4-deficiency induced IBD. RNA-seq analysis demonstrated a significant upregulation of important inflammatory cytokines, such as *il1b* and *tnfa* in the Ctla-4-deficient intestines. This is consistent with studies showing that IL-1β and TNF-α act as crucial mediators in mammalian IBD models by disrupting epithelial junctions and inducing apoptosis of epithelial cells [53, 54]. Conversely, the key anti-inflammatory cytokines, such as *il10*, were downregulated. These findings highlight an imbalance between pro-inflammatory and anti-inflammatory cytokines in the intestines of Ctla-4-deficient fish.

Consistently, the inflammatory signaling pathways associated with these upregulated cytokines, such as the ERK1/2 and MAPK pathways, were positively enriched in inflamed intestines. Single-cell RNA-seq analysis revealed the upregulation and enrichment of these inflammatory cytokines and pathways in neutrophils, macrophages, and B cells of inflamed intestines, indicating their active involvement in inflammatory responses and as major sources of inflammatory signals. Additionally, there was a marked increase in the abundance of Th2 subset cells in the inflamed intestines. These Th2 cells exhibited high expression of *il13* and were significantly enriched in inflammatory signaling pathways, indicating their activated state. Previous research in mammals has identified IL-13 as the key effector Th2 cytokine in ulcerative colitis, directly causing damage to epithelial cells by affecting epithelial tight junctions, apoptosis, and cell restitution [23]. Therefore, upregulated Il13 from Th2 cells may be a significant contributor to epithelial injury in zebrafish model. Notably, the proportion of ILC3-like cells was downregulated in the inflamed intestines, consistent with recent studies reporting a substantial decline of ILC3 in IBD patients [29-31]. ILC3 is the most abundant type of ILCs in the intestines and plays a protective role in IBD in mammals by promoting epithelial cell proliferation and survival, as well as enhancing intestinal barrier function through the production of IL-22 [32, 34]. Thus, the marked decrease in ILC3-like cells may exacerbate intestinal inflammation and damage. Besides, abnormal activation of Th2 cells can lead to dysfunction in downstream B cells and mucosa-associated immunity, which is essential in maintaining the symbiotic bacterial homeostasis in intestines [38]. Thus, a potential correlation may exist between changes

in Th2 cells and the observed alterations in the intestinal microbial community in Ctla-4-deficient zebrafish. Overall, our findings provided insights into the occurrence of IBD. However, gaining a comprehensive understanding of the complex interactions among these immune cells, intestinal epithelial cells, and the microbiome in IBD requires further exploration.

Materials and Methods

Experimental fish

Wild-type AB zebrafish (*D rerio*) of both sexes, 4-6 months of age with body weights ranging from 0.3 to 0.8 g and lengths of 3-4 cm, were reared in the laboratory in recirculating water at 26-28 ℃ under standard conditions as previously described [55]. All animal experiments were performed in accordance with legal regulations and were approved by the Research Ethics Committee of Zhejiang University.

Generation of Ctla-4-deficient zebrafish

CRISPR/Cas9 system was used to knock out the *ctla-4* gene. The targeting sequence 5'-CTCAGAGCCCTACTTCGCAA-3' was designed by optimized CRISPR Design (http://crispr.mit.edu/) and synthesized by T7 RNA polymerase and purified by MEGAclear Kit (AM1908; Invitrogen) *in vitro*. Cas9 protein (500 ng/μl, A45220P; Thermo Fisher Scientific) and purified RNA (90 ng/μl) were coinjected into one cell-stage wild-type embryos. For genotyping, DNA fragment was amplified with primers (F: 5' -TGTGACAGGAAAAGATGGAGAA- 3' and R: 5'-GATCAGATCCACTCCTCCAAAG- 3') and subjected to sequencing. The mutant

alleles (-14 bp) were obtained and their offspring were used in experiments.

Histopathological analysis

The anterior, mid, and posterior intestines (n = 3 replicates) were fixed in 4% PFA

overnight and embedded in paraffin. The tissues were cut into 4 μm sections and stained

with Hematoxylin and eosin (H&E) for pathological analysis. To evaluate the severity

of intestinal inflammation, histologic scores were determined according to the criteria

supplied by previous publication [22]. Briefly, three independent parameters, including

inflammation, extent, and epithelial changes, were assessed and scored as follows: the

severity of inflammation (0 = none, 1 = minimal, 2 = mild, 3 = moderate, 4 = marked);

the level of inflammation extent (0 = none, 1 = mucosa, 2 = mucosa and submucosa, 3

= transmural), the degree of epithelial changes (0 = none, 1 = minimal hyperplasia, 2 =

mild hyperplasia, minimal goblet cell loss, 3 = moderate hyperplasia, mild goblet cell

loss, 4 = marked hyperplasia with moderate to marked goblet cell loss). Each parameter

was calculated and summed to obtain the overall score. The sections were stained with

Periodic Acid-Schiff (PAS) reagent to evaluate the number of goblet cells.

RNA-sequencing and bioinformatic analysis

Total RNAs were isolated from wild-type or *ctla-4$^{-/-}$* intestines (three biological

replicates) using TRIzol reagent following the manufacturer's instructions (Takara).

cDNA libraries were constructed using NEB Next Ultra Directional RNA Library Prep

Kit (NEB), and sequencing was performed according to the Illumina Hiseq2500

standard protocol at LC Bio (Hangzhou, China). The differentially expressed genes

(DEGs) were identified with absolute Log_2 fold change > 1 and adjusted *p*-value < 0.05

by R package DESeq2. Gene Ontology (GO) enrichment and Kyoto encyclopedia of genes and genomes (KEGG) enrichment analyses were performed by the OmicStudio (http://www.omicstudio.cn/tool) tools. Gene-set enrichment analysis was performed using software GSEA (v4.1.0, https://www.gsea-msigdb.org/gsea/index.jsp), and the heatmap was generated using the R package ggplot2. For the protein-protein interaction (PPI) networks, the DEGs were retrieved in STRING (version 11.5, https://string-db.org/) database (combined score > 0.4), and the PPI network was visualized by Cytoscape software (version 3.9.1) [56]. The betweenness centrality (BC) was calculated using the CytoNCA plugin in Cytoscape software. The RNA-sequencing (RNA-seq) data in this study were deposited in the Gene Expression Omnibus (GEO) (http://www.ncbi.nlm.nih.gov/geo/) database.

Quantitative real-time PCR

Total RNA from intestines was extracted using TRIzol reagent (Takara Bio) and reverse transcribed into cDNAs according to the manufacturer's protocol. PCR experiments were performed in a total volume of 10 μl by using an SYBR Premix Ex Taq kit (Takara Bio). The reaction mixtures were incubated for 2 min at 95℃, then subjected to 40 cycles of 15 s at 95℃, 15 s at 60℃, and 20 s at 72℃. Relative expression levels of the target genes were calculated using the $2^{-\Delta\Delta ct}$ method with β-actin for normalization. Each PCR trial was run in triplicate parallel reactions and repeated three times. The primers used in the experiments are listed in Supplemental Table 1.

Single-cell RNA-sequencing analysis

The intestines from wild-type and *ctla-4*[-/-] zebrafish were washed by D-Hank's and incubated with D-Hank's containing EDTA (0.37 mg/mL) and DTT (0.14 mg/mL) for 20 min. The resulting supernatants were collected as fraction 1. The remaining tissues were subsequently digested with type IV collagenase (0.15 mg/mL) for 1 h at room temperature and the resulting supernatants were collected as fraction 2. Fractions 1 and 2 were combined and centrifuged at 350 g for 10 min. Cells were then washed with D-Hank's and suspended in 40% percoll (percoll : FBS: L-15 medium = 4 : 1: 5) and passed through a 40-μm strainer. The cell suspension was carefully layered onto 63% percoll (percoll : FBS : L-15 medium = 6.3 : 1: 2.7) and centrifuged at 400 g for 30 min at room temperature. The cell layer of the interface was collected and washed with D-Hank's at 400 g for 10 min. Cell quantity and viability were assessed using 0.4% trypan blue staining, revealing that over 90% of the cells were viable. Single-cell samples were submitted to the LC-Bio Technology Co., Ltd (Hangzhou, China) for 10× Genomics library preparation and data analysis assistance. Libraries were prepared using the Chromium[TM] Controller and Chromium[TM] Single Cell 3' Library & Gel Bead Kit v2 (10× Genomics) according to the manufacturer's protocol, and sequenced on an Illumina NovaSeq 6000 sequencing system (paired-end multiplexing run, 150 bp) at a minimum depth of 20,000 reads per cell. Sequencing results were demultiplexed and converted to FASTQ format using Illumina bcl2fastq software and the data were aligned to Ensembl zebrafish genome assembly GRCz11. Quality control was performed using the Seurat. DoubletFinder R package was used to identify and filter the doublets (multiplets) [57]. The cells were removed if they expressed fewer than 500

unique genes, or > 60% mitochondrial reads. The number of cells after filtration in the

current study was 3,263 in wild-type and 4,276 in *ctla-4*$^{-/-}$ groups, respectively. Cells

were grouped into an optimal number of clusters for *de novo* cell type discovery using

Seurat's FindNeighbors() and FindClusters() functions, graph-based clustering

approach with visualization of cells being achieved through the use of t-SNE or UMAP

plots [58]. The cell types were determined using a combination of marker genes

identified from the literature and gene ontology. The marker genes were visualized by

dot plot and t-SNE plots to reveal specific individual cell types.

In vivo administration of sCtla-4-Ig

An *in vivo* sCtla-4-Ig administration assay was conducted to evaluate the potential

therapeutic effect of sCTLA-4-Ig on intervention of a *ctla-4*-deficiency induced IBD-

like phenotype. For this procedure, the *ctla-4*$^{-/-}$ zebrafish were i.p administered with

recombinant soluble Ctla-4-Ig protein (sCtla-4-Ig, 2 μg/g body weight) every other day

until day 14. Fish that received an equal amount of human IgG isotype were used as

control. The dose of sCtla-4-Ig was chosen based on its ability to significantly inhibit

the mRNA expression of *il13* in Ctla-4-deficient zebrafish.

Statistical analysis

Statistical analysis and graphical presentation were performed with GraphPad

Prism 8.0. All data were presented as the mean ± SD of each group. Statistical

evaluation of differences was assessed using one-way ANOVA, followed by an

unpaired two-tailed *t*-teat. Statistical significance was defined as $^{*}p < 0.05$, $^{**}p < 0.01$,

$^{***}p < 0.001$, and $^{****}p < 0.0001$. All experiments were replicated at least three times.

Data availability

All data generated or analyzed during this study are included in this article and its supplementary information files. The RNA-seq and scRNA-seq data for this study have been deposited in NCBI Gene Expression Omnibus (GEO) (https://www.ncbi.nlm.nih.gov/geo/) under accession numbers GSE255304 and GSE255303, respectively. The 16S rRNA sequencing data in this study have been deposited in the NCBI Sequence Read Archive (SRA) (https://www.ncbi.nlm.nih.gov/sra/) with an accession number of BioProject PRJNA1073727.

Acknowledgements

We are grateful to Bio-ultrastructure analysis Lab of Analysis center of Agrobiology and environmental sciences, Zhejiang Univ in the TEM sample preparation and observation. We thank Hong Deng and Qiong Huang for advice and expertise in pathological analysis. We also thank She-long Zhang for two-photon laser confocal scanning microscope capture. This work was supported by grants from the National Natural Science Foundation of China (32173003), the National Key Research and Development Program of China (2018YFD0900503, 2018YFD0900505).

Author Contributions

L. L. Q. and J. Z. S. conceived and designed the experiments. L. L. Q. and C. B.

H. performed the experiments. L. L. Q., C. B. H., Q. Z., Y. W., A. F. L., L. X. X., Y.

C., and J. Z. S. analyzed the data. L. X. X., D. D. F., and J. Z. S. contributed

reagents/materials/analysis tools. L. L. Q., Y. C., L. X. X. and J. Z. S. wrote the

manuscript. All authors reviewed the paper and provided comments.

Conflict of Interest

The authors declare that they have no competing financial interests.

Ethical approval and consent to participate

All animal experiments were performed with the approval of the Ethics Committee for

Animal Experimentation of Zhejiang University.

REFERENCES

1. Hodson R (2016) Inflammatory bowel disease. Nature 540:S97-S97. https://10.1038/540S97a
2. Zhang Y Z, Li Y Y (2014) Inflammatory bowel disease: Pathogenesis. World J Gastroentero 20:91-99. https://10.3748/wjg.v20.i1.91
3. Venema W T C U, Voskuil M D, Dijkstra G, Weersma R K, Festen E A M (2017) The genetic background of inflammatory bowel disease: from correlation to causality. J Pathol 241:146-158. https://10.1002/path.4817
4. Loftus E V (2004) Clinical epidemiology of inflammatory bowel disease: Incidence, prevalence, and environmental influences. Gastroenterology 126:1504-1517. https://10.1053/j.gastro.2004.01.063
5. Khor B, Gardet A, Xavier R J (2011) Genetics and pathogenesis of inflammatory bowel disease. Nature 474:307-317. https://10.1038/nature10209
6. Neurath M F (2020) Host-microbiota interactions in inflammatory bowel disease. Nat Rev Gastro Hepat 17:76-77. https://10.1038/s41575-019-0248-1
7. Kordjazy N, Haj-Mirzaian A, Haj-Mirzaian A, Rohani M M, Gelfand E W, Rezaei N *et al.* (2018) Role of toll-like receptors in inflammatory bowel disease. Pharmacological Research 129:204-215. https://10.1016/j.phrs.2017.11.017
8. Horowitz J E, Warner N, Staples J, Crowley E, Gosalia N, Murchie R *et al.* (2021) Mutation

spectrum of NOD2 reveals recessive inheritance as a main driver of Early Onset Crohn's
Disease. Sci Rep-Uk 11:5595. https://10.1038/s41598-021-84938-8

9.   Duerr R H, Taylor K D, Brant S R, Rioux J D, Silverberg M S, Daly M J *et al.* (2006) A genome-
wide association study identifies IL23R as an inflammatory bowel disease gene. Science
314:1461-1463. https://10.1126/science.1135245

10.   Tremelling M, Berzuini C, Massey D, Bredin F, Price C, Dawson C *et al.* (2008) Contribution
of TNFSF15 gene variants to Crohn's disease susceptibility confirmed in UK population.
Inflamm Bowel Dis 14:733-737. https://10.1002/ibd.20399

11.   Marengere L E, Waterhouse P, Duncan G S, Mittrucker H W, Feng G S, Mak T W (1996)
Regulation of T cell receptor signaling by tyrosine phosphatase SYP association with
CTLA-4. Science 272:1170-1173. https://10.1126/science.272.5265.1170

12.   Lee K M, Chuang E, Griffin M, Khattri R, Hong D K, Zhang W G *et al.* (1998) Molecular basis
of T cell inactivation by CTLA-4. Science 282:2263-2266.
https://10.1126/science.282.5397.2263

13.   Yokosuka T, Kobayashi W, Takamatsu M, Sakata-Sogawa K, Zeng H, Hashimoto-Tane A
et al. (2010) Spatiotemporal Basis of CTLA-4 Costimulatory Molecule-Mediated Negative
Regulation of T Cell Activation. Immunity 33:326-339.
https://10.1016/j.immuni.2010.09.006

14.   Linsley P S, Greene J L, Brady W, Bajorath J, Ledbetter J A, Peach R (1994) Human B7-1
(Cd80) and B7-2 (Cd86) Bind with Similar Avidities but Distinct Kinetics to Cd28 and Ctla-
4 Receptors. Immunity 1:793-801. https://10.1016/S1074-7613(94)80021-9

15.   Collins A V, Brodie D W, Gilbert R J, Iaboni A, Manso-Sancho R, Walse B *et al.* (2002) The
interaction properties of costimulatory molecules revisited. Immunity 17:201-210.
https://10.1016/s1074-7613(02)00362-x

16.   Saito T, Yokosuka T, Hashimoto-Tane A (2010) Dynamic regulation of T cell activation and
co-stimulation through TCR-microclusters. Febs Lett 584:4865-4871.
https://10.1016/j.febslet.2010.11.036

17.   Repnik K, Potocnik U (2010) CT60 Single-Nucleotide Polymorphism Is Associated with
Slovenian Inflammatory Bowel Disease Patients and Regulates Expression of Isoforms.
DNA Cell Biol 29:603-610. https://10.1089/dna.2010.1021

18.   Xia B, Crusius J B A, Wu J, Zwiers A, van Bodegraven A A, Peña A S (2002) CTLA-4 gene
polymorphisms in Dutch and Chinese patients with inflammatory bowel disease. Scand J
Gastroentero 37:1296-1300. https://10.1080/003655202761020579

19.   Jiang T, Ge L Q, Chen Z T, Li C, Zhou F, Luo Y *et al.* (2010) Effect of cytotoxic T lymphocyte-
associated molecule 4 1661 gene polymorphism on its expression and transcription in
ulcerative colitis. J Digest Dis 11:369-375. https://10.1111/j.1751-2980.2010.00462.x

20.   Bamias G, Delladetsima I, Perdiki M, Siakavellas S I, Goukos D, Papatheodoridis G V *et al.*
(2017) Immunological Characteristics of Colitis Associated with Anti-CTLA-4 Antibody
Therapy. Cancer Invest 35:443-455. https://10.1080/07357907.2017.1324032

21.   Lo B C, Kryczek I, Yu J, Vatan L, Caruso R, Matsumoto M *et al.* (2024) Microbiota-
dependent activation of CD4+ T cells induces CTLA-4 blockade–associated colitis via Fcγ
receptors. Science 383:62-70. https://10.1126/science.adh8342

22.   Erben U, Loddenkemper C, Doerfel K, Spieckermann S, Haller D, Heimesaat M M *et al.*
(2014) A guide to histomorphological evaluation of intestinal inflammation in mouse

559   models. Int J Clin Exp Pathol 7:4557-4576.

560   23.   Heller F, Florian P, Bojarski C, Richter J, Christ M, Hillenbrand B *et al.* (2005) Interleukin-13
561   is the key effector Th2 cytokine in ulcerative colitis that affects epithelial tight junctions,
562   apoptosis, and cell restitution. Gastroenterology 129:550-564.
563   https://10.1016/j.gastro.2005.05.002

564   24.   Butin-Israeli V, Bui T M, Wiesolek H L, Mascarenhas L, Lee J J, Mehl L C *et al.* (2019)
565   Neutrophil-induced genomic instability impedes resolution of inflammation and wound
566   healing. J Clin Invest 129:712-726. https://10.1172/JCI122085

567   25.   Lees C W, Barrett J C, Parkes M, Satsangi J (2011) New IBD genetics: common pathways
568   with other diseases. Gut 60:1739-1753. https://10.1136/gut.2009.199679

569   26.   Lee J C, Biasci D, Roberts R, Gearry R B, Mansfield J C, Ahmad T *et al.* (2017) Genome-
570   wide association study identifies distinct genetic contributions to prognosis and
571   susceptibility in Crohn's disease. Nat Genet 49:262-268. https://10.1038/ng.3755

572   27.   Sinkala M, Nkhoma P, Mulder N, Martin D P (2021) Integrated molecular characterisation
573   of the MAPK pathways in human cancers reveals pharmacologically vulnerable mutations
574   and gene dependencies. Communications Biology 4:https://10.1038/s42003-020-01552-
575   6

576   28.   Trebak M, Kinet J P (2019) Calcium signalling in T cells. Nature Reviews Immunology
577   19:154-169. https://10.1038/s41577-018-0110-7

578   29.   Bernink J H, Peters C P, Munneke M, te Velde A A, Meijer S L, Weijer K *et al.* (2013) Human
579   type 1 innate lymphoid cells accumulate in inflamed mucosal tissues. Nat Immunol
580   14:221-229. https://10.1038/ni.2534

581   30.   Li J, Doty A L, Tang Y, Berrie D, Iqbal A, Tan S A *et al.* (2017) Enrichment of IL-17A(+) IFN-
582   gamma(+) and IL-22(+) IFN-gamma(+) T cell subsets is associated with reduction of
583   NKp44(+) ILC3s in the terminal ileum of Crohn's disease patients. Clin Exp Immunol
584   190:143-153. https://10.1111/cei.12996

585   31.   Martin J C, Chang C, Boschetti G, Ungaro R, Giri M, Grout J A *et al.* (2019) Single-Cell
586   Analysis of Crohn's Disease Lesions Identifies a Pathogenic Cellular Module Associated
587   with Resistance to Anti-TNF Therapy. Cell 178:1493-1508 e1420.
588   https://10.1016/j.cell.2019.08.008

589   32.   Buonocore S, Ahern P P, Uhlig H H, Ivanov, II, Littman D R, Maloy K J *et al.* (2010) Innate
590   lymphoid cells drive interleukin-23-dependent innate intestinal pathology. Nature
591   464:1371-1375. https://10.1038/nature08949

592   33.   Ermann J, Staton T, Glickman J N, de Waal Malefyt R, Glimcher L H (2014) Nod/Ripk2
593   signaling in dendritic cells activates IL-17A-secreting innate lymphoid cells and drives
594   colitis in T-bet-/-.Rag2-/- (TRUC) mice. Proc Natl Acad Sci U S A 111:E2559-2566.
595   https://10.1073/pnas.1408540111

596   34.   Aparicio-Domingo P, Romera-Hernandez M, Karrich J J, Cornelissen F, Papazian N,
597   Lindenbergh-Kortleve D J *et al.* (2015) Type 3 innate lymphoid cells maintain intestinal
598   epithelial stem cells after tissue damage. J Exp Med 212:1783-1791.
599   https://10.1084/jem.20150318

600   35.   Ott S J, Musfeldt M, Wenderoth D F, Hampe J, Brant O, Folsch U R *et al.* (2004) Reduction
601   in diversity of the colonic mucosa associated bacterial microflora in patients with active
602   inflammatory bowel disease. Gut 53:685-693. https://10.1136/gut.2003.025403

603   36.   Manichanh C, Rigottier-Gois L, Bonnaud E, Gloux K, Pelletier E, Frangeul L *et al.* (2006)
604   Reduced diversity of faecal microbiota in Crohn's disease revealed by a metagenomic
605   approach. Gut 55:205-211. https://10.1136/gut.2005.073817

606   37.   Zhao Q, Chang H, Zheng J, Li P, Ye L, Pan R *et al.* (2023) A novel Trmt5-deficient zebrafish
607   model with spontaneous inflammatory bowel disease-like phenotype. Signal Transduct
608   Target Ther 8:86. https://10.1038/s41392-023-01318-6

609   38.   Xu Z, Takizawa F, Casadei E, Shibasaki Y, Ding Y, Sauters T J C *et al.* (2020) Specialization
610   of mucosal immunoglobulins in pathogen control and microbiota homeostasis occurred
611   early in vertebrate evolution. Sci Immunol 5:https://10.1126/sciimmunol.aay3254

612   39.   Belzer C, de Vos W M (2012) Microbes inside--from diversity to function: the case of
613   Akkermansia. ISME J 6:1449-1458. https://10.1038/ismej.2012.6

614   40.   Waterhouse P, Penninger J M, Timms E, Wakeham A, Shahinian A, Lee K P *et al.* (1995)
615   Lymphoproliferative Disorders with Early Lethality in Mice Deficient in Ctla-4. Science
616   270:985-988. https://10.1126/science.270.5238.985

617   41.   Tivol E A, Borriello F, Schweitzer A N, Lynch W P, Bluestone J A, Sharpe A H (1995) Loss of
618   Ctla-4 Leads to Massive Lymphoproliferation and Fatal Multiorgan Tissue Destruction,
619   Revealing a Critical Negative Regulatory Role of Ctla-4. Immunity 3:541-547.
620   https://10.1016/1074-7613(95)90125-6

621   42.   Hosseini A, Gharibi T, Marofi F, Babaloo Z, Baradaran B (2020) CTLA-4: From mechanism
622   to autoimmune therapy. Int Immunopharmacol 80:106221.
623   https://10.1016/j.intimp.2020.106221

624   43.   Sun W H, Zhang X M, Wu J, Zhao W D, Zhao S X, Li M L (2019) Correlation of TSHR and
625   CTLA-4 Single Nucleotide Polymorphisms with Graves Disease. Int J Genomics
626   2019:6982623. https://10.1155/2019/6982623

627   44.   Vergara A, De Felice M, Cesaro A, Gragnano F, Pariggiano I, Golia E *et al.* (2023) Immune-
628   Checkpoint Inhibitor-Related Myocarditis: Where We Are and Where We Will Go.
629   Angiology 12:33197231201929. https://10.1177/00033197231201929

630   45.   Kheiralla K E K (2021) CTLA-4 (+49A/G) Polymorphism in Type 1 Diabetes Children of
631   Sudanese Population. Glob Med Genet 8:11-18. https://10.1055/s-0041-1723008

632   46.   Lin T W, Hu Y C, Yang Y H, Chien Y H, Lee N C, Yu H H *et al.* (2022) CTLA-4 gene mutation
633   and multiple sclerosis: A case report and literature review. J Microbiol Immunol 55:545-
634   548. https://10.1016/j.jmii.2021.10.009

635   47.   Cutolo M, Sulli A, Paolino S, Pizzorni C (2016) CTLA-4 blockade in the treatment of
636   rheumatoid arthritis: an update. Expert Rev Clin Immu 12:417-425.
637   https://10.1586/1744666x.2016.1133295

638   48.   Chang M C, Chang Y T, Tien Y W, Liang P C, Jan I S, Wei S C *et al.* (2007) T-Cell regulatory
639   gene CTLA-4 Polymorphism/Haplotype association with autoimmune pancreatitis. Clin
640   Chem 53:1700-1705. https://10.1373/clinchem.2007.085951

641   49.   Fathima N, Narne P, Ishaq M (2019) Association and gene-gene interaction analyses for
642   polymorphic variants in CTLA-4 and FOXP3 genes: role in susceptibility to autoimmune
643   thyroid disease. Endocrine 64:591-604. https://10.1007/s12020-019-01859-3

644   50.   Zeissig S, Petersen B S, Tomczak M, Melum E, Huc-Claustre E, Dougan S K *et al.* (2015)
645   Early-onset Crohn's disease and autoimmunity associated with a variant in CTLA-4. Gut
646   64:1889-1897. https://10.1136/gutjnl-2014-308541

51.   Angelino G, Cifaldi C, Zangari P, Di Cesare S, Di Matteo G, Chiriaco M *et al.* (2021) Gastric
cancer, inflammatory bowel disease and polyautoimmunity in a 17-year-old boy: CTLA-
4 deficiency successfully treated with Abatacept. Eur J Gastroen Hepat 33:E1051-E1056.
https://10.1097/Meg.0000000000002185

52.   Liu J Z, van Sommeren S, Huang H L, Ng S C, Alberts R, Takahashi A *et al.* (2015)
Association analyses identify 38 susceptibility loci for inflammatory bowel disease and
highlight shared genetic risk across populations. Nat Genet 47:979-+.
https://10.1038/ng.3359

53.   Al-Sadi R, Guo S H, Ye D M, Dokladny K, Alhmoud T, Ereifej L *et al.* (2013) Mechanism of
IL-1b Modulation of Intestinal Epithelial Barrier Involves p38 Kinase and Activating
Transcription Factor-2 Activation. Journal of Immunology 190:6596-6606.
https://10.4049/jimmunol.1201876

54.   Al-Sadi R, Guo S H, Ye D M, Ma T Y (2013) TNF-α Modulation of Intestinal Epithelial Tight
Junction Barrier Is Regulated by ERK1/2 Activation of Elk-1. Am J Pathol 183:1871-1884.
https://10.1016/j.ajpath.2013.09.001

55.   Zhu L Y, Lin A F, Shao T, Nie L, Dong W R, Xiang L X *et al.* (2014) B Cells in Teleost Fish
Act as Pivotal Initiating APCs in Priming Adaptive Immunity: An Evolutionary Perspective
on the Origin of the B-1 Cell Subset and B7 Molecules. Journal of Immunology 192:2699-
2714. https://10.4049/jimmunol.1301312

56.   Kohl M, Wiese S, Warscheid B (2011) Cytoscape: software for visualization and analysis of
biological networks. Methods Mol Biol 696:291-303. https://10.1007/978-1-60761-987-
1_18

57.   McGinnis C S, Murrow L M, Gartner Z J (2019) DoubletFinder: Doublet Detection in Single-
Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst 8:329-+.
https://10.1016/j.cels.2019.03.003

58.   Cronan M R, Hughes E J, Brewer W J, Viswanathan G, Hunt E G, Singh B *et al.* (2021) A
non-canonical type 2 immune response coordinates tuberculous granuloma formation
and epithelialization. Cell 184:1757-1774 e1714. https://10.1016/j.cell.2021.02.046

59.   Hu C B, Wang J, Hong Y, Li H, Fan D D, Lin A F *et al.* (2023) Single-cell transcriptome
profiling reveals diverse immune cell populations and their responses to viral infection in
the spleen of zebrafish. Faseb J 37:e22951. https://10.1096/fj.202201505RRRR

Fig.1 Characterization of zebrafish Ctla-4. **A** Alignment of the Ctla-4 homologues from

different species generated with ClustalX and Jalview. The conserved and partially

conserved amino acid residues in each species are colored in hues graded from orange

to red, respectively. The conserved Cysteine residues, and conserved functional motifs,

such as B7-binding motif, Tyrosine phosphorylation site, and potential Tyrosine

phosphorylation site were indicated separately. The signal peptide, IgV-like domain,

transmembrane (TM) domain and cytoplasmic domain were marked at the top of the

sequence. **B** Tertiary protein structures of Ctla-4 ectodomains between humans and zebrafish were predicted by AlphaFold2. The two pairs of disulfide bonds (Cys^{20}-Cys^{91}/Cys^{46}-Cys^{65} in zebrafish and Cys^{21}-Cys^{92}/Cys^{48}-Cys^{66} in humans) used to connect the two-layer β-sandwich, and the separate Cys residue (Cys^{119} in zebrafish and Cys^{120} in humans) associated with the dimerization of the proteins are indicated. The Cys residues are represented in purple ball-and-stick and the identified or potential B7 binding sites are highlighted in blue. **C** Dimer of Ctla-4 was identified by Western blot under reducing (+β-ME) or non-reducing (-β-ME) conditions. The ctrl represents the empty control plasmid. The monomer is indicated by one arrow, the dimer by two arrows. **D** The subcellular localization of Ctla-4 protein was assessed in HEK293T cells transfected with pEGFPN1-Ctla-4 for 48 hours using a two-photon laser-scanning microscope (Original magnification, 630×). The nucleus was stained with DAPI (blue), the cell membrane was stained with DiI (red). **E** tSNE plots show the relative distribution of common T cell markers *cd4-1*, *cd8a* and *ctla-4*. The data are from splenic single cell RNA sequencing (scRNA-seq) dataset we recently established [59]. **F** Immunofluorescence staining of lymphocytes separated from blood, spleens, and kidneys of zebrafish. Cells were stained with mouse anti-Ctla-4, together with rabbit anti-Cd4-1 or rabbit anti-Cd8α. DAPI stain shows the location of the nuclei. The images were obtained by two-photon laser-scanning microscope (Original magnification, 630×).

Fig.2 Examination on the IBD-like phenotype in *ctla-4⁻/⁻* zebrafish. **A** Generation of a

homozygous *ctla-4*-deficient (*ctla-4⁻/⁻*) zebrafish line through CRISPR/Cas9-based

knockout of *ctla-4* gene on chromosome 9. A 14-bp deletion mutation in exon 2 results

in a premature stop at codon 82, which is predicted to produce a defective Ctla-4 protein

containing 81 amino acids. **B** Genotyping of the deficiency of *ctla-4* gene by Sanger

sequencing. **C** Knockout efficiency of Ctla-4 selectively examined in spleen and gut

tissues of *ctla-4*$^{-/-}$ zebrafish by Western blot analysis. Gapdh serves as a loading control.

D Normal gross appearance of adult wild-type (WT) and *ctla-4*$^{-/-}$ zebrafish. **E** Body

weight statistics of WT and *ctla-4*$^{-/-}$ zebrafish (n = 30). **F** The change of intestine length

in WT and *ctla-4*$^{-/-}$ zebrafish. **G** The change of splenic size in WT and *ctla-4*$^{-/-}$ zebrafish.

H Representative H＆E staining analysis of histopathological changes and quantitation

of histology score in anterior, mid and posterior intestines from WT and *ctla-4*$^{-/-}$

zebrafish. Red arrows indicate mucosal inflammatory cell infiltration, and black arrow

indicates transmural inflammatory cell infiltration. **I** PAS analysis indicates goblet cells

in anterior intestine from WT and *ctla-4*$^{-/-}$ zebrafish. **J** Quantitation analysis of goblet

cells of each villus in the foregut of WT and *ctla-4*$^{-/-}$ zebrafish (n = 8). **K** Observation

of cell junctions between intestinal epithelial cells in posterior intestines from WT and

ctla-4$^{-/-}$ zebrafish under TEM (Hitachi Model H-7650). White triangles indicate tight

junctions, black triangle indicates adhesion junctions, and red triangles indicate

desmosomes. Data are presented as mean ± standard deviation (SD). Statistical

significance was assessed through an unpaired Student's t test ($^*p < 0.05$; $^{**}p < 0.01$;

$^{***}p < 0.001$; $^{****}p < 0.0001$).

Fig.3 RNA-sequencing analysis of the molecular implications associated with the IBD-

like phenotype in *ctla-4⁻/⁻* zebrafish. **A** Heatmap of different expressed genes between

the intestines from wild-type (WT) and *ctla-4⁻/⁻* zebrafish. **B** Volcano plot showing the

up-/down-regulated genes in the intestines of *ctla-4⁻/⁻* zebrafish compared with those of

WT zebrafish. Red indicates up-regulated genes and blue indicates down-regulated

genes. **C** GO analysis showing top 10 terms in biological processes of DEGs. **D** GO

analysis showing top 10 terms in biological processes of all up-regulated genes. **E**

Heatmap of representative differently expressed inflammation and chemotaxis-related

genes. **F** The mRNA expression levels of important genes associated with inflammation

and chemokines confirmed by real-time qPCR. **G** Protein-protein interaction network

was constructed using the DEGs. The nodes represent the proteins (genes); the edges

represent the interaction of proteins (genes). **H** The MPO activity in the intestines (up)

and peripheral blood (down). **I** Heatmap of IBD biomarker genes and IBD-related genes.

J The mRNA expression levels of representative IBD biomarker genes and IBD-related

genes were analyzed by real-time qPCR. Data are presented as mean ± standard

deviation (SD). Statistical significance was assessed through an unpaired Student's t

test ($^{**}p < 0.01$; $^{***}p < 0.001$; $^{****}p < 0.0001$).

Fig.4 Single cell RNA-sequencing analysis of the major cell types associated with the

IBD-like phenotype in *ctla-4⁻/⁻* zebrafish. **A** Classification of cell types from zebrafish

intestines by tSNE nonlinear clustering. **B** Dot plot showing the expression levels of

lineage marker genes and percentage of cells per cluster that express the gene of interest.

C Expression maps of T cell associated markers within the cell populations of the

zebrafish intestines. **D** Heatmap of the tight/adhesion junction-related genes in

enterocytes among the samples from wild-type (WT) and *ctla-4*$^{-/-}$ zebrafish. **E** Heatmap

of inflammation-related genes involved in cytokine-cytokine receptor interaction in

neutrophils among the samples from WT and *ctla-4*$^{-/-}$ zebrafish. **F** KEGG enrichment

analysis showing the top 15 terms of up-regulated genes in neutrophils from the *ctla-4*$^{-}$

$^{/-}$ sample versus the WT sample. **G** KEGG enrichment analysis showing the top 15

terms of up-regulated genes in macrophages from the *ctla-4*$^{-/-}$ sample versus the WT

sample.

Fig.5 Single cell RNA-sequencing analysis of the subset immune-cells associated with

the IBD-like phenotype in *ctla-4⁻/⁻* zebrafish. **A** Classification of subset cells from the

T/NK/ILC-like group by tSNE nonlinear clustering. **B** Dot plot showing the expression

levels of subset marker genes and percentage of cells per cluster that express the gene

of interest. **C** Marker gene expression in individual cluster identifying this cluster as

ILC3-like cells. **D** Changes in the composition of subset cells between samples from

wild-type (WT) and *ctla-4⁻/⁻* zebrafish. A significantly increased Th2 subset (referred to

as Th2 cells 2) in the *ctla-4⁻/⁻* sample was highlighted with a black dashed circle. **E**

Histogram showing the different ratios of subset cells between the WT and *ctla-4⁻/⁻*

samples. **F** Histogram presenting the different numbers of subset cells between the WT

and *ctla-4⁻/⁻* samples. **G** Expression maps of the cytokine *il13* within different subset

cells between the WT and *ctla-4⁻/⁻* samples. **H** Dot plots illustrating the differential

expression of *il13* in T/NK/ILC-like cells from WT and *ctla-4⁻/⁻* zebrafish. **I** KEGG

enrichment analysis showing the top 15 terms of up-regulated genes in the subset of

Th2 cells 2. **J** KEGG enrichment analysis showing the top 15 terms of up-regulated

genes in NKT cells. **K** Scatter plot showing the DEGs of ILC3-like cells in WT and

ctla-4⁻/⁻ zebrafish. The *il17a/f1* and *il17a/f3* was shown in the scatter plot.

Fig.6 Alteration in microbial composition in the intestines of *ctla-4⁻ᐟ⁻* zebrafish. **A** Venn

diagram showing the number of ASVs in zebrafish intestinal microbiota. **B** Alpha-

diversity of microbes was calculated through Shannon index and Simpson index. **C**

Beta-diversity analyzed based on PCoA was shown by using Bray Curtis distance. **D**

The relative abundance of intestinal microbiota at the class level. **E-G** The relative

abundance of Alphaproteobacteria (**E**), Bacilli (**F**) and verrucomicrobiae (**G**) in the

intestines from the wild-type (WT) and *ctla-4⁻ᐟ⁻* zebrafish. *$p < 0.05$. **H** The relative

abundance of intestinal microbiota at the family level. **I** Heatmap analysis of the

differential abundances of bacterial communities at family level in the WT and *ctla-4⁻ᐟ⁻*

zebrafish ($p < 0.05$). **J** Cladogram representation of LEfSe analysis showing the

differentially abundant bacterial taxa between the intestines from WT (red) and *ctla-4⁻*

 ᐟ⁻ (green) zebrafish ($p < 0.05$).

Fig.7 Examination on the inhibitory function of Ctla-4 in T cell activation. **A** Assessment of the proliferative activity of T cells from wild-type and *ctla-4*⁻ᐟ⁻ zebrafish by a mixed lymphocyte reaction combined with PHA-stimulation. The CFSE dilution, which served as an indicator of lymphocyte proliferation, was measured through flow cytometry. **B** Assessment of the proliferative activity of lymphocytes from *ctla-4*⁻ᐟ⁻ zebrafish by the administration of sCtla-4-Ig. **C** Assessment of the proliferative activity of lymphocytes from wild-type zebrafish by supplementing anti-Ctla-4 antibody. **D** Assessment of the proliferative activity of lymphocytes from *ctla-4*⁻ᐟ⁻ zebrafish by the administration of sCd28-4-Ig. **E** Assessment of the proliferative activity of lymphocytes

from *ctla-4$^{-/-}$* zebrafish by the administration of recombinant Cd80/86 protein. **F, G**

Interactions between Cd80/86 and Cd28 (**F**), and Cd80/86 and Ctla-4 (**G**) as predicted

by AlphaFold2. On the left are structural models depicting Cd80/86 in complex with

Cd28 or Ctla-4. The center panels display per-residue model confidence scores (pLDDT)

for each structure, using a color gradient from 0 to 100, where higher scores indicate

increased confidence. The right panels show the predicated aligned error (PAE) scores

for each model. The well-defined interfaces between Cd28 or Ctla-4 and Cd80/86 are

highlighted with red dashed squares. **H** The interaction between Cd80/86 and Cd28

(left), and Cd80/86 and Ctla-4 (right) were verified by Co-IP. **I** Binding affinities of the

Cd80/86 protein for the Cd28 and Ctla-4 proteins were measured by the MST assay.

The K_D values are provided. Data are presented as mean ± standard deviation (SD),

which were calculated from three independent experiments. Statistical significance was

assessed through an unpaired Student's t test ($^{**}p < 0.01$; $^{***}p < 0.001$; ns denotes no

statistical significance).

Fig.8 In vivo inhibition of intestinal inflammation by Ctla-4-Ig. **A** Percent initial weight

of zebrafish after injection of the Ctla-4-Ig or the IgG isotype control. In each group, n

= 6. Data show means with SEM analyzed by two-way ANOVA with Sidak's correction

for multiple comparisons. **B** Representative H&E staining analysis of histopathological

changes and quantitation of histology score in the posterior intestine from Ctla-4-Ig or

IgG isotype control-supplemented *ctla-4*[-/-] zebrafish. Scale bar: 50 μm. **C** The mRNA

expression levels of inflammation-related genes in the Ctla-4-Ig or IgG isotype control

treatment groups of *ctla-4*[-/-] zebrafish. **D** The mRNA expression levels of IBD

biomarker genes and IBD-related genes in the Ctla-4-Ig or IgG isotype control

treatment groups of *ctla-4*[-/-] zebrafish. The *p* value was generated by unpaired two-

tailed Student's *t-test*. $^{**}p < 0.01$; $^{***}p < 0.001$; $^{****}p < 0.0001$.

Supplemental Figures

Fig. S1 The organization, sequence and phylogenetic analysis of zebrafish *ctla-4* gene.

A Comparison of the intron/exon organizations of *ctla-4* gene in zebrafish and humans.

Exons and introns are shown with black boxes and lines, and their size are indicated by

the numbers found above and below the sequences respectively. **B** The nucleotide and

amino acid sequences of *ctla-4* gene and Ctla-4 protein. The underline indicates the

signal peptide, the circles represent the conserved cysteine residues. **C** Phylogenetic

analysis of the relationship of Ctla-4 between zebrafish and other species. The unrooted

phylogenetic tree was constructed through the neighbor-joining method based on the

amino acid alignment (ClustalX) of amino acid sequences. The numbers represent the

percentage bootstrap confidence derived from 500 replicates.

Fig. S2 Preparation of mouse anti-Ctla-4 antibody. **A** SDS-PAGE detection of the

recombinant Ctla-4 protein with extracellular domain (ECD). Lane 1, 2 and 3 represent

the protein markers, blank, and target protein, respectively. **B** Western blot analysis of

the mouse anti-EGFP and anti-Ctla-4 antibodies that bind to the recombinant Ctla-4-

EGFP fusion proteins expressed in HEK293T cells. **C** Western blot analysis of native

Ctla-4 protein in zebrafish intestinal tissues using mouse anti-Ctla-4 antibody.

Fig. S3 Examination on the functional genes and pathways associated with the IBD-

like phenotype in *ctla-4*$^{-/-}$ zebrafish. **A** Top 5 KEGG enrichment bar plot of up-regulated

genes in *ctla-4⁻ᐟ⁻* zebrafish intestines versus wild-type zebrafish intestines. **B** Top 10

KEGG enrichment bar plot of down-regulated genes in *ctla-4⁻ᐟ⁻* zebrafish intestines

versus wild-type zebrafish intestines. **C, D** Changes in the expression of genes

associated with lymphocyte chemotaxis, positive regulation of ERK1/ERK2 cascades,

Calcium and MAPK signaling pathways in the *ctla-4⁻ᐟ⁻* zebrafish intestines analyzed by

using a collection of pre-defined gene sets retrieved from GO (**C**) and KEGG (**D**)

database. The *p* value, false discovery rates (FDR) and normalized enrichment score

(NES) are shown above each pathway graph.

Fig. S4 Examination on the involvement of apoptotic process in epithelial cells and

expression of inflammation-related genes in neutrophils and B cells in the intestines of

ctla-4⁻ᐟ⁻ zebrafish. **A** Expression map of the epithelial markers within the cell

populations of the zebrafish intestines. **B** Expression map of the neutrophil markers

within the cell populations of the zebrafish intestines. **C** KEGG enrichment bar plot of

all DEGs from epithelial cells. **D** Heatmap of inflammation-related genes in B cells

from wild-type and *ctla-4⁻ᐟ⁻* intestines.

Fig. S5 Examination on the activation of T cell subsets in the intestines of *ctla-4⁻ᐟ⁻*

zebrafish. **A** Marker gene expression in individual cluster identifying the cluster as

NKT cells. **B** Heatmap of up-regulated genes involved in cytokine-cytokine receptor

interaction in NKT cells among the wild-type and *ctla-4⁻ᐟ⁻* samples. **C** KEGG

enrichment analysis showing top 15 terms of up-regulated genes in Cd8⁺ T cells in *ctla-*

4⁻ᐟ⁻ samples versus wild-type samples. **D** Heatmap of up-regulated genes involved in

cytokine-cytokine receptor interaction in Cd8$^+$ T cells among the wild-type and *ctla-4$^-$*

$^{/-}$ samples.

Fig. S6 Preparation of recombinant proteins and examination on the molecular

interactions. **A-C** SDS-PAGE detection of the purified recombinant soluble Ctla-4-Ig

(**A**) and Cd28-Ig (**B**) proteins and the Cd80/86 extracellular domain (ECD) (**C**) with

Coomassie brilliant blue staining. **D** Flow cytometry analysis of the interaction between

Cd80/86 and Cd28 (up), as well as Cd80/86 and Ctla-4 (down). Data are presented as

mean ± SD, which were calculated from three independent experiments. Statistical

significance was evaluated using an unpaired Student's t test ($^{**}p < 0.01$; $^{***}p < 0.001$).