TITLE

CD2AP is Co-Expressed with Tropomyosin-Related Kinase A and Ras-Related Protein Rab-5A in

Cholinergic Neurons of the Murine Basal Forebrain

AUTHORS

Lindsey Avery Fitzsimons[1,2]   lfitzsimons@une.edu

Mohammad Atif-Sheikh[3]   matifsheikh@une.edu

Jayden Lovely[3]   jlovely@une.edu

Madison Mueth[2,4]   mmueth@une.edu

Makaela Rice[2]   mrice4@une.edu

Kevin Kotredes[5]   kevin.kotredes@jax.org

Gareth Howell[5]   gareth.howell@jax.org

Benjamin J Harrison[1,2]   bharrison2@une.edu

1.   Biomedical Sciences, College of Osteopathic Medicine, University of New England,

Biddeford. ME 04005

2.   Center for Excellence in the Neurosciences, University of New England. ME 04005

3.   College of Arts and Sciences, University of New England, Biddeford. ME 04005

4.   Graduate School of Biomedical Science and Engineering, University of Maine, Orono, ME

04469

5.   The Jackson Laboratory, Bar Harbor, ME 04609

KEYWORDS

CD2AP, cholinergic neurons, Alzheimer's Disease, Rab5, retrograde transport

AKNOWLEDGEMENTS

Funding for these experiments were provided by the National Institutes of Health (R01NS1215; PI: Benjamin J. Harrison) as well as from the University of New England, Center for Pain Research (NIGMS UNE COBRE #P30GM145497; PI: Ian Meng, PhD) and the Khan Foundation through the University of New England, College of Arts and Sciences (Biddeford, Maine, USA). Animal care services were generously provided by UNE COBRE Behavior Core; histology & imaging training and services were provided by the UNE COBRE Histology & Imaging Core Facilities. Both UNE Behavior Core and Imaging & Histology Core facilities are also supported by the UNE Center for Pain Research (NIGMS UNE COBRE #P30GM145497; PI: Ian Meng, PhD). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We wish to thank Christoph Straub for providing numerous tissues and experimental assistance as well as members of the Harrison lab who assisted with imaging, experiments and laboratory management, especially Peter Neufeld, BS. Graphical abstract was created with Biorender.com.

CONFLICT OF INTEREST STATEMENT

The authors declare no competing financial interests.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author

upon reasonable request

ABSTRACT

Basal forebrain cholinergic neurons project to the hippocampus and cortex, are critical for learning

and memory, and are central to the pathogenesis of Alzheimer's disease (AD). GWAS have

consistently shown that genomic variants at the *CD2AP* gene locus are associated with significant

increased risk of AD. GWAS studies have also shown that genetic variants in endocytosis genes,

including *RAB5A*, significantly increase susceptibility to AD. Previous work in our lab has shown

that CD2AP functions as a docking-scaffold/adaptor protein as a coordinator of nerve growth

factor (NGF) and trophic signaling in neurons. We have also demonstrated that CD2AP positively

regulates Rab5-mediated mechanisms of endocytosis in primary sensory neurons. The purpose

of this study was to perform an *in vivo* characterization of CD2AP expression in cholinergic

neurons of the brain regions most relevant to AD pathogenesis and to investigate the

colocalization of CD2AP and Rab5 in cholinergic neurons of the murine basal forebrain. Brain

tissue was perfused, harvested from ChAT^{BAC}-eGFP transgenic mice (N=4 male, N=4 female;

aged 10 mo), where cholinergic neurons (co-) express green fluorescence protein (GFP) in central

and peripheral neurons that express choline acetyltransferase (ChAT). Frozen tissue sections

were used to assess the specificity of the reporter in mouse brain along with localization of both

CD2AP and Rab5 (co-) expression using immunofluorescence (IF) analysis of ChAT-GFP+

neurons and primary antibodies against ChAT, CD2AP and Rab5. Image J software was used to

develop and optimize a colocalization assay for CD2AP and Rab5 puncta. Experiments were

repeated in a follow-up cohort of aged-adult mice (N=2 male, N=2 female; aged 18 mo). IF

expression of CD2AP was quantified in the basal forebrain, diagonal band of Broca (vDB), and

striatal regions and compared to results from the cortical regions of the adult mouse brain.

Colocalization of CD2AP was observed in the cell bodies of ChAT-GFP+ neurons of the striatum,

vDB and basal forebrain regions, where CD2AP expression intensity as well as the number of cell

bodies with positive signal increased incrementally. Colocalization analyses revealed near-

complete overlap of CD2AP and Rab5 expression in ChAT-GFP+ cholinergic neurons of the basal

forebrain region. We conclude that cholinergic neurons express CD2AP in healthy adult and

aged-adult mouse brains. These data provide the first evidence of quantifiable CD2AP protein

expression of cholinergic neurons specific to the diagonal band of Broca (vDB) and basal

forebrain. Together with previous research from our lab, these data support a role for CD2AP in

the pathogenesis of AD through orchestration of endocytosis and retrograde signaling. Ongoing

studies are underway to verify these findings in a novel AD mouse model that incorporates the

humanized variant of *CD2AP*, created by MODEL-AD, where we aim to further investigate how

CD2AP variants may affect mechanistic components of Rab5 endocytosis as well as subsequent

survival of cholinergic neurons in the context of known amyloid beta and Tau pathologies.

1 INTRODUCTION

CD2-associated protein (CD2AP) is a docking-scaffold/adaptor protein with multiple

binding interfaces for coordination of protein-protein interactions. It was first characterized as an

SH-3-containing protein that binds to CD-2 via the cytoplasmic domain (Dustin et al., 1998; Rouka

et al., 2015). CD2AP serves as a hub for signaling pathway components that converge on this

scaffold. When the required combination of interactions are formed, the corresponding

downstream signaling cascade is activated, and for example, the presence or absence of CD2AP-

binding interactions differentially modulates the amplitude of MAPK-ERK or PI3K signal

transduction (Kirsch et al., 2001). Known upstream binding-partners include Tropomyosin

receptor kinase-A (TrkA) and/or rearranged during transfection (RET); both of are activated by

glial cell line-derived neurotrophic factor (GDNF) family ligands (GFLs) including nerve growth

factor (NGF) (Harrison et al., 2016; Tsui & Pierchala, 2008). Known downstream effector binding-

partners include the PI3K regulatory subunit p85 (Harrison et al., 2016), p21ras (Qui & Green,

1992) and Akt (Sang et al., 2018).

Endocytosis is a cellular process through which cells absorb proteins on the external surface

of the cell by engulfing the molecular contents along with the segment of plasma membrane to

form an endosome. Endosomes may subsequently function as mobile compartmentalized

signaling platforms termed signalosomes (Hoeller et al., 2005). In neuronal cell populations,

retrograde trophic signaling requires endosomal signaling both to internalize the ligand-receptor

complex as well as for the propagation/transportation towards its downstream effectors located in

the cell body (Delcroix et al., n.d.; Hoeller et al., 2005; Ye et al., 2003).

Variants in genes associated with endocytosis may be neuro-protective, involved in

developing resilience against Alzheimer's Disease (AD) (Tesi et al., 2020). Conversely,

pathogenic variants have been consistently found in genome-wide association study (GWAS)

datasets, as factors associated with increased susceptibility to AD (Lambert et al., 2013; Tesi et

al., 2020). Specifically, the *Cd2ap* gene is an AD-predisposition locus, substantiated by multiple

GWAS with multiple populations (African American (Kunkle et al., 2021; Sherva et al., 2023),

Chinese (Xue et al., 2022), Northern and Western European ancestry (CEU)(Gockley et al.,

2021), UK Biobank (Khaire et al., 2022; Marioni et al., 2018)) that support the association of

Cd2ap variants with small but significant increases in incidence (Amlie-Wolf et al., 2019),

progression (Amlie-Wolf et al., 2019; van Bree et al., 2022), and severity (van Bree et al., 2022;

Xue et al., 2022) of Alzheimer's Disease (AD). CD2AP is expressed by multiple cell types in

multiple tissue types (Li et al., 2000), and may play a role in several mechanisms responsible for

the pathogenesis and/or progression of AD. For example, dysfunction of brain vasculature is a

pathophysiological mechanism of blood brain barrier deficits (Cochran et al., 2015) relevant to

multiple dementias, including AD, and is a well-established source of CD2AP expression (Asghari

et al., 2024; Koizumi et al., 2016; Liu et al., 2024). Outside of the vasculature, CD2AP is known to

be expressed by migratory dendritic cell populations (Li et al., 2000; Schaefer et al., 2017), renal

podocytes (Li et al., 2000; Schwarz et al., 2001), and sensory neurons of the peripheral nervous

system (Harrison et al., 2016). Thus, the translational relevance of GWAS studies of *Cd2ap* must

be interpreted carefully, and should serve as a confirmation of the relevance of endocytosis and

retrograde transport in AD. However, the mechanistic underpinning of *Cd2ap* variant-associated

risk, specific to the dynamic interplay between/among cell and tissue types remain poorly

characterized.

It is widely accepted that phosphorylated Tau protein, forming neurofibrillary tangles, impedes

conductance and function of neurons, resulting in the neurotoxicity and degradation seen in

healthy aging as well as in neurodegenerative disease like AD (Dickson et al., 1986; Kovacs et

al., 2017). A genetic association between CD2AP, tau-accumulation and subsequent neurotoxicity

(Nisbet et al., 2015) have been observed using drosophila homolog *cindr* in *a* loss of function

model of AD (Nisbet et al., 2015; Shulman et al., 2014). CD2AP scaffolding interactions regulate

Rab5-mediated endocytosis (Behl et al., 2022; Cormont et al., 2003a; Furusawa et al., 2019;

Gockley et al., 2021; Maninger et al., 2024; Rouka et al., 2015; Shen et al., 2020; Xu et al.,

2018a). Therefore, reduced CD2AP function may lead to impaired clearance and subsequent

accumulation of amyloid-beta (Aß) and tau aggregates (Behl et al., 2022).

Reduced kidney function and diagnosis with chronic kidney diseases (CKD) is an

established risk factor for cognitive decline and AD, although the mechanism has yet to be fully

elucidated (Nakagawa et al., 2017; Shi et al., 2018; Zhang et al., 2020). The biochemistry and

function of CD2AP was first characterized in T-lymphocytes of the immune system, where it was

described as an adhesion molecule responsible for mechanisms of protein segregation and

cytoskeletal polarity, and where it was first postulated to orchestrate dynamics of receptor

patterning via the cytoskeletal scaffold (Dustin et al., 1998). This initial characterization prompted

further exploration of CD2AP expression. In other cell and organ systems, and specifically in the

podocyte cells of the kidney, a highly-specialized, gate-keeping cell found in the glomerulus

(Dustin et al., 1998; Nakagawa et al., 2017; Shi et al., 2018; Shih et al., 1999; Zhang et al., 2020).

In podocytes, CD2AP is located to the actin-rich scaffolds of the slit-diaphragm, a charge-

selective opening through which vascular/metabolic waste products are filtered from circulating

blood and excreted from the body in urine. Loss of CD2AP is homozygous-lethal, due to the rapid-

onset of severe nephrotic syndrome and glomerular disease, resulting in kidney failure and death

in mice by 6 weeks of age due to failure of the slit-diaphragm (Li et al., 2000; Shih et al., 1999).

Thus, CD2AP functions as a mediator and organizer of F-Actin structures, where it colocalizes

with Rab4 in podocyte endosomes (Welsch et al., 2001, 2005). Genetic mutations and/or

impairment of CD2AP, therefore, not only impairs cellular cytoarchitectural scaffolding, but also

decreases endosomal-mediated degradation processing, influencing secretase and amyloid

precursor proteins (APP) transport and ultimately enabling the accumulation of APP segments

and tangles (Behl et al., 2022; Howe & Mobley, 2004; Iulita & Cuello, 2016).

CD2AP is also required for the maintenance of structure and permeability of systemic

blood vessels (e.g. epithelial and endothelial cells) (Hamano et al., 2002; Li et al., 2000; Vandal et

al., 2022). Interestingly, analysis of microvessels isolated from human patients with confirmed AD

observed *Cd2ap* expression to be linearly associated with cognitive decline and *Cd2ap* KO from

brain microvasculature (microvessels) in mice resulted in cognitive impairment (Vandal et al.,

2022). Interaction between the vasculature and the immune system has also been implicated in

the onset, progression and severity of AD (Bettcher et al., 2021; Heppner et al., 2015). In vascular

endothelial cells CD2AP regulates intercellular adhesion molecule-1 (ICAM-1) (Schaefer et al.,

2017). Loss of CD2AP initiates increased clustering of ICAM-1 on the endothelial cell surface,

leading to excessive leukocyte transmigration and subsequent pro-inflammatory signaling

pathways associated with kidney dysfunction, arthritis and atherosclerosis (Schaefer et al., 2017).

CD2AP has also been shown to be highly expressed in plasmacytoid dendritic cells (pDC), a

specialized subtype of dendritic cells known to initiate type-1 interferon signaling in response to

inflammation and in microbial pattern recognition (Srivatsan et al., 2013).

In neuronal cell populations, CD2AP has been shown to be expressed in embryonic dorsal

root ganglia (DRG) in E11 mouse embryos, and was identified as a key regulator molecule for

growth factor and hormone-mediated cell signaling mechanisms critical for mammalian

organogenesis and embryogenesis as a whole (Lehtonen et al., 2008). Previous studies from our

laboratory have demonstrated that CD2AP is expressed in mature primary sensory neurons of the

adult peripheral nervous system *in vivo (Harrison et al., 2016)*. Stimulation of these cells with

nerve growth factor (NGF), initiated CD2AP binding to Neurotrophic Receptor Tyrosine Kinase 1

(Gene name *Ntrk1*, Protein name: Tropomyosin Receptor Kinase A (TrkA)) and the PI3-kinase

effector (p85) , to upregulate the PI3K/AKT pathway, further demonstrating a functional role for

CD2AP as a positive coordinator of NGF-stimulated axon growth and branching (Harrison et al.,

2016). More recent experiments investigating the Ras and Rab Interactor 3 (RIN3), have

demonstrated that CD2AP is expressed in murine cholinergic neurons isolated from basal

forebrain, and that endosomal trafficking involving CD2AP *in vitro*, is reliant upon RIN3, a guanine

nucleotide exchange factor (GEF) in the Rab5 small GTPase family (Yoshida et al., 2021).

CD2AP has also been characterized in the nervous system of *Drosophila,* via its homologue, *cindr*

(Ojelade et al., 2019). *Cindr* expression can be found in adult *Drosophila* throughout the

brain/nervous system, and is predominantly expressed within neuropil and the surrounding

presynaptic terminals, where it is shown to function as a positive regulator and rate-limiting

component of synaptic transmission, calcium release, and proteostasis (Ojelade et al., 2019).

Silencing of *cindr* (CD2AP) robustly enhanced tau-toxicity in *Drosophila* (Shulman et al., 2014),

using a human mutant form of Tau shown to be associated with familial frontotemporal dementia

(Wittmann et al., 2001).

Previously, we observed that CD2AP is expressed in primary sensory neurons, and

demonstrated that this adaptor protein coordinates NGF-TrkA-PI3K trophic signaling and

retrograde transport via Rab5-decorated endosomes (Harrison et al., 2016). In the current study,

we observed that in central nervous system (CNS) neurons, CD2AP expression is predominantly

expressed in populations of cholinergic neurons in the basal forebrain and striatum. Cholinergic

projections from the basal forebrain to the cortex constitute the primary source of cholinergic input

to the forebrain (Engelhardt et al., 2007), and are also known to be the first to show signs of

degeneration in AD (Geula et al., 2021). Disruption of NGF-TrkA trophic signaling via interruption

of retrograde Rab5a endosomes is a key mechanism of AD pathogenesis (Iwano et al., 2023). We

therefore also examine the localization of CD2AP and TrkA with Rab5a endosomes in cholinergic

neurons and discuss a possible role for CD2AP in the trophic support of cholinergic neurons and

AD pathophysiology.

2 MATERIALS AND METHODS

2.1 Animals

Animal housing and procedures were approved by the Institutional Animal Care and Use

Committee of the University of New England consistent with federal regulations and guidelines. All

animals used for this study were ChAT[BAC]-eGFP mice (JAX Strain# 007902) purchased from The

Jackson Laboratory (Bar Harbor, Maine). Age and sex of cohorts are specified in appropriate

sections below.

2.2 Prediction of CD2AP mRNA Expression in Neuronal Populations

The DropViz online search tool was used to uncover putative populations of *Cd2ap*-expressing

adult mouse neurons (Saunders et al., 2018). The "Limit by Class" option was set to "Neuron" and

the top-10 results were retrieved using following individual search terms: "*Cd2ap*", "*Ntrk1*" and

"*Chat*". Results reflect the top-10 expressing populations for each of these individual search

terms.

2.3 Tissue Preparation & Frozen Tissue Sectioning

Six week-old, ChAT-GFP:Cre-+ mice (N=2 males, 2 females), were deeply anaesthetized with

0.1ml intraperitoneal Euthasol, before intracardial perfusion with 20ml phosphate buffered saline

(PBS) followed by 30ml cold 4% paraformaldehyde in PBS. Brains were immediately harvested,

and post fixed in cold 4% paraformaldehyde in PBS for 4 hours. Following fixation, tissues were

washed with PBS and incubated in 30% sucrose cryoprotectant for 48 hours. Tissues were then

embedded in OCT frozen tissue embedding medium and stored at -80 degrees Celsius prior to

sectioning. Coronal sectioning of mouse brains embedded in OCT was performed using a Leica

Cryostat 1900 (Leica Microsystems, USA). From each mouse, fifteen 12um thick, coronal brain

sections were sliced 1.77mm to 0.13mm rostral to the bregma onto a glass slide (VWR®,

Superfrost® Plus Micro Slide, Cat# 48311-703).

2.4 Tissue Processing and Immunofluorescent Staining

Sections were dried overnight, then briefly washed with PBS before incubation with blocking

solution (4% normal donkey serum, 0.3% Triton X-100 PBS) for 30 mins at room temperature.

Following blocking, slides were incubated with primary antibodies at the indicated dilutions over-

night (18 hours) at room temperature (Table 1). Slides were then washed 5 times for 5 minutes

with washing buffer (0.3% Triton X-100 in PBS) before incubation with fluorescent secondary

antibodies (Table 1) at room temperature for 1 hour. Finally, slides were washed 5 times for 5

minutes with washing buffer (0.3% Triton X-100 in PBS) before mounting with Fluoroshield

Mounting Medium with DAPI (Abcam, Cat# ab104139) using 1.5 thickness coverslips. The

following primary antibodies were used: Anti-Cd2ap (Proteintech, Cat# 51046-1-AP,

RRID:AB_2879436); anti-TrkA (R&D, Cat# AF1056, RRID:AB_2283049); anti-choline

acetyltransferase (ChAT; Sigma-Aldrich, Cat# AB144P, RRID:AB_2079751); anti-green

fluorescence protein (GFP; Aves Labs, Cat# GFP-1020, RRID:AB_10000240); and anti-Rab5

(Cell Signaling Technology, Cat# 46449, RRID:AB_2799303).

2.5 Thick tissue Immunofluorescence (iDISCO)

Preliminary characterization was performed using thick tissue sections (150μm) of ChAT-

GFP:Cre+ mouse brains embedded in 5% agarose in 1X PBS. Thick sections were cleared and

prepped for IF using iDISCO (Renier et al., 2014). Whole-section images of Chat-GFP+ brains

were imaged at 20X magnification using an epifluorescence microscopy slide scanner and images

were merged to generate a composite using the image stitching function (BZ-X800 Slide

Scanning Microscope, Keyence, USA). Working concentrations of primary and secondary

antibodies were increased for thick tissue section IF and are indicated in Table 1**.**

TABLE 1 Antibodies and reagents used for immunofluorescence analysis.

2.6 CD2AP Antibody Validation

The anti-CD2AP primary antibody (Proteintech, Cat# 51046-1-AP, RRID:AB_2879436) was

validated for use with qualitative and quantitative IF in two previous studies. Characterization of

this antibody was performed using fixed, frozen sections of mouse kidney tissue, demonstrating

signal specificity localized along the glomerular basement membrane, as expected (Yoshida et

al., 2021). Subsequent validation of anti-Cd2ap was performed using Lentiviral-mediated, *Cd2ap*-

knockdown tissue in human telomerase-immortalized retinal pigmented epithelial cells (hTERT-

RPE1) as well as murine NIH3T3 cells (Iwano et al., 2023). Investigators demonstrated that

expression of CD2AP was localized to the primary cilium in control cells, and was ablated in

knockdown tissues using both IF (See Figure 5, panels A-C, Iwano et al., 2023) and Western Blot

(See Figure 5, panel E Iwano et al., 2023). Additional antibody validation was performed in our

laboratory, using kidney tissue collected from ChAT-GFP+ mice, fixed and processed for frozen

tissue sectioning.

2.7 Quantification of CD2AP Fluorescence

2.7.1 Microscopy and Image Analysis

One slide/mouse (with all 15 sections) was imaged using an epifluorescence microscope (Leica

DMi8, Leica Microsystems) at 20X. Four color micrographs from alternate (odd numbered)

sections were then processed using FIJI software (ImageJ2; version 2.14.0/1.5f) (Rueden et al.,

2017). For fluorescence measurements in ChAT-GFP positive neurons, a binary mask was

created by thresholding the GFP channel. The mask was processed to remove noise by removing

foci<20μm in diameter. Masks were then layered over the ChAT antibody stain and then the

CD2AP antibody stain to acquire the respective fluorescent signals from GFP-positive neurons.

2.7.2 Estimation of Background Fluorescence

Populations of cell bodies of developing cholinergic neurons undergo a phenotypic switch into

adulthood, where multiple cholinergic cell subtypes are thought to be established based on

synaptic properties (Ahmed et al., 2019). In ChAT/GFP+ mice, GFP signals are sustained into

adulthood (Tallini et al., 2006), and represent a subpopulation of cortical interneurons which

synapse locally within the cortex, and are best characterized by their expression of vasoactive

intestinal peptide (Ahmed et al., 2019; Engelhardt et al., 2007). In the present study, we confirm

that this population of adult cortical neurons remain ChAT-GFP positive, but do not stain with

ChAT antibody. Therefore, we measured the background fluorescence of antibody stains using

signals from GFP-positive ChAT-negative neurons.

2.8 Immunofluorescence Colocalization Assays

Colocalization assays were performed on IF z-stack images collected using confocal microscopy

(Leica Stellaris) and Image J *EzColocalization* plugin (Stauffer et al., 2018). To restrict analysis to

Chat-GFP+ neurons, the polygon selection tool was used to trace the outline of the ChAT-GFP+

cell body from the GFP+ maximum projection, and the 'region of interest' (ROI) manager was

employed to demarcate the GFP+ border onto the Cd2ap (yellow) and Rab5 (magenta) split

channel maximal projections. Selection of images was restricted to those collected in the vDB

region of the basal forebrain, and an average of 10 images per section, 20 sections per animal, in

a total of 4 ChAT-GFP+ mice.

2.9 Statistical Analysis

All statistical analyses were conducted using Graphpad Prism 9. Data are presented as

individual animal regional mean values ± standard deviation (SD) and reflect quantification of

integrated density of fluorescence intensity, measured using FIJI (ImajeJ2). Collated data from

grouped animals was expressed as mean regional values ± standard error of the mean (SEM).

Data normality for individual animals as well as grouped data ware analyzed and confirmed using

the Kolmogorov-Smirnov test. Regional quantification of CD2AP expression was compared

between brain regions using a standard one-way analysis of variance (ANOVA) followed by

Tukey's post-hoc analysis. For all analyses, statistical significance was set at $p < 0.05$.

For immunofluorescence colocalization assays, degree of overlap for CD2AP and Rab5

punctuate colocalization was assessed using the R value calculated from Pearson correlation,

analysis. Additional analysis of colocalization degree of overlap was assessed using a Costas

array, and statistical significance of the Costas P value was $p < 0.05$.

3 RESULTS

3.1 Analysis of Publicly Available Single-Cell Sequencing Data Reveals that *Cd2ap* mRNA

is Expressed in TrkA-Expressing Cholinergic Neurons

Our previous work provided one of the first descriptions of the structural and mechanistic

dynamics of CD2AP in neurotrophin-mediated neuroplasticity, coordination of scaffolding

interactions between TrkA and AKT subunit p85 in neurons and the positive regulation of

retrograde signaling implemented through Rab5-decorated signalosomes (Harrison et al., 2016).

We further postulated that the retrograde, Rab5-mediated signaling mechanism, combined with

the dynamics of the CD2AP-TrkA complex in cholinergic neurons would be a reasonable,

potential mechanism(s), highly applicable to the pathogenesis of neurodegenerative disease like

AD (Harrison et al., 2016). Since this publication, germline variants in human *CD2AP* have

remained at the forefront of GWAS investigating risk(s) for development of AD in humans (Latina

et al., 2023; Rajabli et al., 2023; Xue et al., 2022).

The functional consequences of *Cd2ap* variants as well as the possibility of a potential role

of CD2AP in CNS neurons is currently unknown. Using human AD GWAS studies, we first

performed a review of the literature generated, with an emphasis on recent GWAS studies

published between 2021 to present (date of submission, June 2024). Since 2021, we report a total

of five GWAS studies for AD that identify *CD2AP* as a primary risk factor (and risk factor loci) for

the development of AD in humans (Table 2). Of the loci identified, specific variant rs9349407 was

correlated with increased risk for AD in Chinese populations (Gao et al., 2022) and specific variant

rs7738720 was correlated with increased risk of AD pathophysiology in individuals of African

ancestry (Sherva et al., 2023). Overall, we identified a total of 19 GWAS studies performed

worldwide (since the year 2010) where *CD2AP* loci have been positively associated with AD

pathophysiology (Table 2).

TABLE 2 Summary of GWAS studies from 2013 to present (June 2024) implicating *CD2AP* in the

pathogenesis and/or susceptibility to AD in humans.

Although there is a clear association between *Cd2ap* variants and an increased risk of AD,

it is currently unknown if *Cd2ap* (mRNA) or CD2AP (protein) is expressed in neurons of the CNS

in either mouse or human models of health and disease. In order to determine and characterize

CD2AP expression in CNS neurons, we performed an analysis of single-cell, neuronal, RNA-

sequencing data publicly-available and sourced from DropViz webtool (Saunders et al., 2018)

(Table 1). Our initial analysis of neuronal cell types ordered neurons by level of CD2AP

expression (Figure 1, magenta), expressed as transcripts per 100,000 in cluster. From these data,

we found that of the few neuronal populations expressing CD2AP, these neurons also expressed

the highest levels of *Ntrk1 (TrkA),* classifying them as cholinergic neuronal cell populations

(Figure 1**,** cyan). Highest expression levels of *Ntrk1* were found in cholinergic interneurons of the

striatum (Figure 1), along with neurons located in the globus pallidus (GP; Figure 1). Striatum

interneurons and GP neurons were found to also co-express the highest values of *Cd2ap* in the

top-10, highest Ntrk1-expressing (cholinergic) neurons of the adult mouse brain (Figure 1). In

other words, of all neuronal cell types present in the adult mouse brain, *Cd2ap* expression was

restricted to Trk-A expressing, cholinergic neurons (Figure 1).

3.2 CD2AP Protein is Expressed in Cholinergic Neurons of the Murine Basal Forebrain

To determine if CD2AP is expressed in ChAT neurons, (as predicted in Figure 1), we

performed an immunofluorescence analysis of frozen tissue sections from ChAT-GFP reporter

(ChAT-GFP+) mouse brains (Figure 2). Whole-section analysis of ChAT-GFP+ brains confirmed

the bilateral presence and expression of ChAT-GFP+ neurons throughout the superior thalamic

radiation (striatum), nucleus accumbens and substantia innominata (Figure 2, a). CD2AP

expression was highest in the basal forebrain region, where we observed the presence of ChAT-

GFP+ neurons (cyan) throughout the vDB. Co-localization of ChAT-GFP+ and anti-ChAT (yellow)

in presumed-cholinergic neurons of the vDB confirmed the specificity of the ChAT primary

antibody, as well as provided confirmation that these neurons are, in fact, acetylcholine-producing

(Figure 2). Of note, absence of ChAT+ staining (yellow) in Chat-GFP+ cell bodies (cyan) of the

cortical region also confirmed the specificity of the anti-ChAT antibody, as was originally

documented by Gamage and colleagues in their recent characterization of the ChAT-GFP

transgenic mouse brain (Gamage et al., 2023). The presence of ChAT-GFP+/anti-ChAT-negative

cell bodies in the cortex therefore serves as an effective, internal negative control tissue, and as

an additional verification for the specificity of the anti-ChAT antibody.

Next, we sought to quantify expression of CD2AP in cholinergic populations (Figure 3).

Confocal scans were collected from the following regions of adult mouse brains (N=4): Substantia

innominata (basal forebrain) region, vDB, the superior thalamic radiation (striatum), and the cortex

(Figure 3). Three-channel fluorescence signals were acquired from ChAT-GFP reporter (cyan),

CD2AP antibody (magenta) and ChAT antibody (yellow; Figure 3). Double-positive signal from

CD2AP and ChAT antibodies, were detected in the majority of ChAT-GFP+ cell bodies in the

basal forebrain, vDB and striatum (Figure 3). ChAT-GFP+ soma in the cortex region did not stain

with the anti-ChAT antibody (yellow). ChAT is expressed during a developmental window in

cortical neurons, and therefore, in the adult these neurons are ChAT-GFP+ and ChAT-antibody

negative (Gamage et al., 2023). Therefore, this served as a convenient negative control to

quantify background signals from ChAT and CD2AP antibody immunostaining. Note that,

although, CD2AP signal was not present in the cell bodies of cortical CHAT-GFP neurons, it was

expressed in cortical vascular structures and non-neuronal cells, as expected (Vandal et al.,

2022). Taken together, these results validate the specificity of CD2AP immunostaining, and

qualitatively show that CD2AP is co-expressed with ChAT in the substantia innominata (basal

forebrain), vDB, and striatum.

To quantify CD2AP expression, we used the imaging data documenting regional

distribution of CD2AP+ fluorescence in ChAT-GFP/ChAT-double-positive cells (Figure 3) and

performed a one-way ANOVA to compare and contrast CD2AP integrated density values in the

cortex (negative control), superior thalamic radiation, substantia innominata (basal forebrain) and

Broca's area (vDB; Figure 4, a-b). CD2AP fluorescence (integrated density) values of individual

animals were expressed as mean values ± SD (Figure 4, a), and collated group regional data

were expressed as mean values ± SEM (Figure 4, b). In the grouped regional mean data, the

lowest CD2AP fluorescence (mean integrated density ± SEM) was observed in the cortex (3449 ±

124), serving as a baseline/background measurement. Mean signal intensity from the superior

thalamic radiation was 4936± 27.6, the substantia innominata (basal forebrain) was 7428 ± 55.9

and Broca's area/vDB was 10929 ± 72.3 (Figure 4, b). A one-way ANOVA with Tukey post-hoc

analysis revealed a statistically significant difference among brain regions (p=0.0004).

Fluorescence expression of CD2AP was significantly higher in the Broca/vDB regions when

compared to background cortex (p=0.0085; Figure 4, b). Mean CD2AP expression in the vDB was

also statistically significantly greater when compared to the superior thalamic radiation

(p=0.0018), and the basal forebrain (p=0.04; Figure 4, b).

3.3 CD2AP is Colocalized to the Termini of Cortical Projections of ChAT-GFP+ Basal

Forebrain Neurons

During the quantification of CD2AP fluorescence expression, we observed an overlap in

signaling between CD2AP (magenta) and ChAT-GFP+ projections extending from the basal

forebrain up through the isocortical regions. Subsequent focused analyses sought to substantiate

these observations with the use of immunofluorescence to more thoroughly document CD2AP co-

expression with ChAT-GFP+ signal beginning with ChAT-GFP+ cell bodies in the basal forebrain

and continuing throughout the neuronal projections extending anteriorly and superiorly in

proximity to the hippocampus and within the isocortex (Figure 5). Preliminary analysis of thick

tissue sections from ChAT-GFP:Cre+ brains allowed investigators to characterize and visualize

both the spatial and dimensional layout of ChAT-GFP+ neurons, along with their respective

neuronal projections and termini from a three-dimensional perspective (Figure 5, top).

Confocal microscopy analysis of ChAT-GFP+ cell bodies was repeated to confirm the co-

labeling of ChAT-GFP+ (cyan) and CD2AP+ (magenta; magenta and cyan overlap is white) signal

in the cell bodies of cholinergic neurons of the basal forebrain (Figure 5, bottom left). Subsequent

analyses of ChAT-GFP+ neuronal projections from the basal forebrain confirmed the presence of

co-labeling with CD2AP in these cholinergic fibers (Figure 5, bottom right). Of note, vascular

expression of CD2AP is substantial and can easily be visualized throughout all regions of the

mouse brain. However, CD2AP co-labeling in ChAT-GFP+ projections are visualized in a

morphological pattern and smaller/finer distribution of specific staining that is morphologically

distinctive from CD2AP+ expression in blood vessels (Figure 5, bottom).

3.4 CD2AP is colocalized with Rab5-Positive, Early Endosomal Formations of ChAT-GFP+

Basal Forebrain Neurons.

To further examine CD2AP expression in cholinergic neurons, relative to endosomal

formation, we performed a series of immunofluorescence experiments to characterize and

quantify CD2AP colocalization with Rab5, a small GTPase, mediator and sorter of endosomal

formation (Figure 6). Immunofluorescence analysis of ChAT-GFP+ brain frozen tissue sections

revealed an abundance of CD2AP (yellow) and Rab5 (magenta) colocalization puncta, which

were interpreted as formed endosomes, and which were diffusely present in ChAT-GFP+ neurons

(cyan) throughout the basal forebrain tissue. CD2AP-Rab5-positive (CD2AP/Rab5+) endosomes

observed throughout the cell body of the ChAT-GFP+ neurons and are indicated by a white

arrowhead (Figure 6, a). We also observed clusters of CCD2AP/Rab5+ endosomes that did not

overlap with the ChAT-GFP+ signal of the ChAT-GFP+ cell bodies and were indicated by a red

arrowhead (Figure 6, b). In order to quantify colocalization of CD2AP (yellow) with Rab5+ vesicles

(magenta), we performed colocalization analyses of ChAT-GFP+ cell bodies (cyan; Figure 6).

Quantification of CD2AP/Rab5+ colocalization revealed a near-complete overlap between

CD2AP+ and Rab5+ punctuates (Figure 6). Mean Pearson correlation R values was 0.80, with a

Costas P value of 1.00, demonstrated in Figure 6.

4 DISCUSSION

By analysis of publicly available scRNA-seq data, we observed inherent expression of

Cd2ap specifically and exclusively in Trk-A expressing, cholinergic neurons of the central nervous

system. Using an *in vivo* reporter mouse for ChAT-GFP-positive cholinergic neurons, we

confirmed the presence and distribution of ChAT/GFP+ neurons in the superior thalamic radiation

(striatum), nucleus accumbens/substantia innominata, the basal forebrain, and the vDB.

Additionally, we identified and quantified the colocalization/expression of ChAT/GFP+/ChAT+ with

CD2AP in cholinergic neurons in each of these regions, most pronounced of which was quantified

in the vDB/basal forebrain. Notably, expression of CD2AP in ChAT/GFP+ axonal

branching/termini continues outside of the basal forebrain and up through the (iso-) cortex, posing

additional hypotheses surrounding a possible role/application for CD2AP in retrograde axonal

transport signaling in maintaining the viability and bidirectionality of basal forebrain

communication with the cortex in healthy aging and in AD pathogenesis. Co-staining of ChAT-

GFP+/CD2AP+ cell bodies with an antibody against Rab5 revealed near-complete colocalization

between CD2AP and Rab5 punctuate patterning, consistent with endosomal structural

configurations previously described in primary sensory neurons of the peripheral nervous system

and in cholinergic neurons *in vitro* (Harrison et al., 2016; Shen et al., 2020). In this study, we are

the first to characterize the expression of the Alzheimer associated gene, *Cd2ap,* in populations of

adult CNS neurons, *in vivo*.

CD2AP was first characterized as an adhesion molecule critical for the structural integrity,

filtration and overall function of the kidney, as determined using various genetic knockout/knock-

down studies *in vivo* and *in vitro* (Li et al., 2000; Schwarz et al., 2001; Shih et al., 1999)*.*

Concurrently, CD2AP was identified as a genetic risk factor for the development of AD in humans

(Hollingworth et al., 2011; Naj et al., 2011), a finding that has been supported by a substantial

number of studies across model organism species, including *Mus musculus* (Vandal et al., 2022),

Drosophila (Ojelade et al., 2019; Shulman et al., 2014), and *Caenorhabditis elegans* (Alvarez et

al., 2022; Kim et al., 2018). Previously established *Cd2ap* variants have even been incorporated

along with variants in *Trem2^{em1Adiuj}, App^{em1Adiuj}* and *Apoe$^{tm1.1(APOE*4)Adiuj}$* into a multi-genetic/causal,

humanized mouse model of AD (JAX Strain# 033873) (Jung et al., 2023; Pandey et al., 2023).

Although CD2AP is known to be expressed in tissues and cell types outside of the kidney (e.g.

primary sensory neurons (Harrison et al., 2016), endo/epithelial cells (Cochran et al., 2015; Li et

al., 2000) and migratory glial cells (Bao et al., 2012; Vandal et al., 2022), inherent expression of

CD2AP in cholinergic neurons of CNS (*in vivo*) has not previously been identified or quantified,

supporting the plausibility of a pathogenic mechanism of AD inherent to cholinergic neurons of the

CNS.

477   Early characterizations of CD2AP (previously also referred to as 'Cas ligand with multiple

Src homology (SH) 3 domains' (CMS)) identified that co-expression of active Rab4 and CD2AP *in*

vivo resulted in the enlargement of early endosome formations, was required (along with c-Cbl)

for proper endosomal formation, and finally, that CD2AP interactions with both Rab4 and Rab7

might also be involved in degradation of these endosomes (Cormont et al., 2003b). Interestingly,

CD2AP expression in conditionally immortalized mouse podocytes was shown to be directly co-

expressed with Rab4, but not Rab5, in late-endosomal processing (Welsch et al., 2005).

However, subsequent research from our laboratory and others, has indicated that CD2AP

interaction with specific Rab isoforms may, in fact, vary by cell/tissue type, where NGF signaling

via TrkA receptor in primary, DRG neurons, is dependent upon TrkA+/Rab5+ endosomes for

mediation of long-range NGF signaling (Furusawa et al., 2019; Harrison et al., 2016). In the

present study, we document the visual and quantitated colocalization of CD2AP and Rab5 *in vivo,*

where previous research has identified interactions with Rab3 (Iwano et al., 2023), Rab4

(Furusawa et al., 2019), and RIN (Rouka et al., 2015; Shen et al., 2020) or isolated with Rab5 *in*

vitro (Furusawa et al., 2019; Rouka et al., 2015). Evidence in support of a role for Rab5

specifically for endosomal formation and the propagation of trophic signaling in cholinergic

neurons (Behl et al., 2022; Harrison et al., 2016; Kawabata, 2022), in conjunction an established

association of impaired Rab5 signaling in brain tissue of deceased humans with confirmed AD (Xu

et al., 2018b), are consistent with the findings of the present study, and further support the

hypothesis that CD2AP-Rab5 endosomal signaling in cholinergic neurons of the basal forebrain is

a viable, mechanistic explanation for AD pathogenesis.

In the nervous system, CD2AP plays a regulatory role in neuronal cell survival

mechanisms, a process that is known to be trophic-factor dependent (Tsui & Pierchala, 2008).

Early investigations into the possible pathological relevance of trophic signaling in AD showed that

while retrograde transport was reduced, there is a specific reduction in NGF retrograde transport

observed from the hippocampus to the septum in basal forebrain cholinergic neurons in a Ts65Dn

mouse model of Down's Syndrome, also known to be applicable to mouse models of AD (Cooper

et al., 2001). In addition to what is known about CD2AP, Rab5-mediated endocytosis is also

hypothesized to be dysregulated in AD (Pensalfini et al., 2020; Xu et al., 2018b), a finding that is

also supported by human GWAS studies (Novikova et al., 2021; Tesi et al., 2020).

In the present manuscript, we reiterate the origin and pathogenic significance of

cholinergic neurons in AD, where current working hypotheses support the idea that disruption of

endocytosis may lead to accumulation of amyloid beta and/or the formation of neurofibrillary

tangles. We highlight and summarize the volume of GWAS studies supporting the idea that

variants in genes known to be critical for endocytosis and endocytic processes (like *Cd2ap* and

Rab5) and that are consistently associated with increased susceptibility to AD. We summarize

previous studies from our laboratory and others, supporting a role for CD2AP as a coordinator of

NGF signaling. Finally, we present new data demonstrating that CD2AP is expressed in both the

cell body as well as localization in the projections of cholinergic neurons in the CNS, reiterating

the working hypothesis that CD2AP may orchestrate endocytosis in cholinergic neurons and

formalizing the feasibility and relevance of this mechanism to AD pathogenesis.

5 CONCLUSION

CD2AP is expressed in multiple cell types critical for the function of various tissues/organs

like those of the kidney, vasculature, immune system, sensory neurons of the PNS, and now,

cholinergic neurons of the CNS. Taken together, we propose a working model through which the

targeting CD2AP-coordinated signals may serve as a viable target to prevent endocytosis-

mediated mechanisms of neurodegeneration and may also serve as a multifactorial, systems-

wide approach to improving AD outcomes.

REFERENCES

Ahmed, N. Y., Knowles, R., & Dehorter, N. (2019). New Insights Into Cholinergic Neuron Diversity. *Frontiers*

in Molecular Neuroscience, *12*, 204. https://doi.org/10.3389/fnmol.2019.00204

Alvarez, J., Alvarez-Illera, P., Santo-Domingo, J., Fonteriz, R. I., & Montero, M. (2022). Modeling

Alzheimer's Disease in Caenorhabditis elegans. *Biomedicines*, *10*(2), Article 2.

https://doi.org/10.3390/biomedicines10020288

Amlie-Wolf, A., Tang, M., Way, J., Dombroski, B., Jiang, M., Vrettos, N., Chou, Y.-F., Zhao, Y., Kuzma, A.,

Mlynarski, E. E., Leung, Y. Y., Brown, C. D., Wang, L.-S., & Schellenberg, G. D. (2019). Inferring the

Molecular Mechanisms of Noncoding Alzheimer's Disease-Associated Genetic Variants. *Journal of*

Alzheimer's Disease: JAD, *72*(1), 301–318. https://doi.org/10.3233/JAD-190568

Asghari, K., Niknam, Z., Mohammadpour-Asl, S., & Chodari, L. (2024). Cellular junction dynamics and

Alzheimer's disease: A comprehensive review. *Molecular Biology Reports*, *51*(1), 273.

https://doi.org/10.1007/s11033-024-09242-w

Bao, M., Hanabuchi, S., Facchinetti, V., Du, Q., Bover, L., Plumas, J., Chaperot, L., Cao, W., Qin, J., Sun, S.-C.,

& Liu, Y.-J. (2012). CD2AP/SHIP1 complex positively regulates plasmacytoid dendritic cell receptor

signaling by inhibiting the E3 ubiquitin ligase Cbl. *Journal of Immunology (Baltimore, Md.: 1950)*,

189(2), 786–792. https://doi.org/10.4049/jimmunol.1200887

Behl, T., Kaur, D., Sehgal, A., Singh, S., Makeen, H. A., Albratty, M., Abdellatif, A. A. H., Dachani, S. R., &

Bungau, S. (2022). Exploring the potential role of rab5 protein in endo-lysosomal impairment in

Alzheimer's disease. *Biomedicine & Pharmacotherapy = Biomedecine & Pharmacotherapie*, *148*,

112773. https://doi.org/10.1016/j.biopha.2022.112773

Bettcher, B. M., Tansey, M. G., Dorothée, G., & Heneka, M. T. (2021). Peripheral and central immune

system crosstalk in Alzheimer disease—A research prospectus. *Nature Reviews Neurology*, *17*(11),

Article 11. https://doi.org/10.1038/s41582-021-00549-x

Cochran, J. N., Rush, T., Buckingham, S. C., & Roberson, E. D. (2015). The Alzheimer's disease risk factor

CD2AP maintains blood–brain barrier integrity. *Human Molecular Genetics*, *24*(23), 6667–6674.

https://doi.org/10.1093/hmg/ddv371

Cooper, J. D., Salehi, A., Delcroix, J. D., Howe, C. L., Belichenko, P. V., Chua-Couzens, J., Kilbridge, J. F.,

Carlson, E. J., Epstein, C. J., & Mobley, W. C. (2001). Failed retrograde transport of NGF in a mouse

model of Down's syndrome: Reversal of cholinergic neurodegenerative phenotypes following NGF

infusion. *Proceedings of the National Academy of Sciences of the United States of America*, *98*(18),

10439–10444. https://doi.org/10.1073/pnas.181219298

Cormont, M., Metón, I., Mari, M., Monzo, P., Keslair, F., Gaskin, C., McGraw, T. E., & Le Marchand-Brustel,

Y. (2003a). CD2AP/CMS regulates endosome morphology and traffic to the degradative pathway

through its interaction with Rab4 and c-Cbl. *Traffic (Copenhagen, Denmark)*, *4*(2), 97–112.

https://doi.org/10.1034/j.1600-0854.2003.40205.x

Cormont, M., Metón, I., Mari, M., Monzo, P., Keslair, F., Gaskin, C., McGraw, T. E., & Le Marchand-Brustel,

Y. (2003b). CD2AP/CMS regulates endosome morphology and traffic to the degradative pathway

through its interaction with Rab4 and c-Cbl. *Traffic (Copenhagen, Denmark)*, *4*(2), 97–112.

https://doi.org/10.1034/j.1600-0854.2003.40205.x

Delcroix, J.-D., Valletta, J. S., Wu, C., Hunt, S. J., Kowal, A. S., & Mobley, W. C. (n.d.). *NGF Signaling in

Sensory Neurons: Evidence that Early Endosomes Carry NGF Retrograde Signals*.

Dickson, D. W., Yen, S.-H., Suzuki, K. I., Davies, P., Garcia, J. H., & Hirano, A. (1986). Ballooned neurons in

select neurodegenerative diseases contain phosphorylated neurofilament epitopes. *Acta*

Neuropathologica, *71*(3), 216–223. https://doi.org/10.1007/BF00688042

Dustin, M. L., Olszowy, M. W., Holdorf, A. D., Li, J., Bromley, S., Desai, N., Widder, P., Rosenberger, F., van

der Merwe, P. A., Allen, P. M., & Shaw, A. S. (1998). A novel adaptor protein orchestrates receptor

patterning and cytoskeletal polarity in T-cell contacts. *Cell*, *94*(5), 667–677.

https://doi.org/10.1016/s0092-8674(00)81608-6

Engelhardt, J. von, Eliava, M., Meyer, A. H., Rozov, A., & Monyer, H. (2007). Functional Characterization of

Intrinsic Cholinergic Interneurons in the Cortex. *Journal of Neuroscience*, *27*(21), 5633–5642.

https://doi.org/10.1523/JNEUROSCI.4647-06.2007

Furusawa, K., Takasugi, T., Chiu, Y.-W., Hori, Y., Tomita, T., Fukuda, M., & Hisanaga, S. (2019). CD2-

associated protein (CD2AP) overexpression accelerates amyloid precursor protein (APP) transfer

from early endosomes to the lysosomal degradation pathway. *Journal of Biological Chemistry*,

294(28), 10886–10899. https://doi.org/10.1074/jbc.RA118.005385

Gamage, R., Zaborszky, L., Münch, G., & Gyengesi, E. (2023). Evaluation of eGFP expression in the ChAT-

eGFP transgenic mouse brain. *BMC Neuroscience*, *24*(1), 4. https://doi.org/10.1186/s12868-023-

00773-9

Gao, S., Hao, J., Zhao, Y., Li, X., Wang, T., Han, Z., Sun, B., Sun, J., & Liu, G. (2022). An updated analysis of

the association between CD2-associated protein gene rs9349407 polymorphism and Alzheimer's

disease in Chinese population. *Frontiers in Neuroinformatics*, *16*, 1006164.

https://doi.org/10.3389/fninf.2022.1006164

Geula, C., Dunlop, S. R., Ayala, I., Kawles, A. S., Flanagan, M. E., Gefen, T., & Mesulam, M.-M. (2021). Basal

forebrain cholinergic system in the dementias: Vulnerability, resilience, and resistance. *Journal of*

Neurochemistry, *158*(6), 1394–1411. https://doi.org/10.1111/jnc.15471

Gockley, J., Montgomery, K. S., Poehlman, W. L., Wiley, J. C., Liu, Y., Gerasimov, E., Greenwood, A. K.,

Sieberts, S. K., Wingo, A. P., Wingo, T. S., Mangravite, L. M., & Logsdon, B. A. (2021). Multi-tissue

neocortical transcriptome-wide association study implicates 8 genes across 6 genomic loci in

Alzheimer's disease. *Genome Medicine*, *13*(1), 76. https://doi.org/10.1186/s13073-021-00890-2

Hamano, Y., Grunkemeyer, J. A., Sudhakar, A., Zeisberg, M., Cosgrove, D., Morello, R., Lee, B., Sugimoto,

H., & Kalluri, R. (2002). Determinants of Vascular Permeability in the Kidney Glomerulus *. *Journal*

of Biological Chemistry, *277*(34), 31154–31162. https://doi.org/10.1074/jbc.M204806200

Harrison, B. J., Venkat, G., Lamb, J. L., Hutson, T. H., Drury, C., Rau, K. K., Bunge, M. B., Mendell, L. M.,

Gage, F. H., Johnson, R. D., Hill, C. E., Rouchka, E. C., Moon, L. D. F., & Petruska, J. C. (2016). The

Adaptor Protein CD2AP Is a Coordinator of Neurotrophin Signaling-Mediated Axon Arbor

Plasticity. *Journal of Neuroscience*, *36*(15), 4259–4275. https://doi.org/10.1523/JNEUROSCI.2423-

15.2016

Heppner, F. L., Ransohoff, R. M., & Becher, B. (2015). Immune attack: The role of inflammation in

Alzheimer disease. *Nature Reviews Neuroscience*, *16*(6), Article 6.

https://doi.org/10.1038/nrn3880

Hoeller, D., Volarevic, S., & Dikic, I. (2005). Compartmentalization of growth factor receptor signalling.

Current Opinion in Cell Biology, *17*(2), 107–111. https://doi.org/10.1016/j.ceb.2005.01.001

Hollingworth, P., Harold, D., Sims, R., Gerrish, A., Lambert, J.-C., Carrasquillo, M. M., Abraham, R.,

Hamshere, M. L., Pahwa, J. S., Moskvina, V., Dowzell, K., Jones, N., Stretton, A., Thomas, C.,

Richards, A., Ivanov, D., Widdowson, C., Chapman, J., Lovestone, S., … Williams, J. (2011). Common

variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's

disease. *Nature Genetics*, *43*(5), 429–435. https://doi.org/10.1038/ng.803

Howe, C. L., & Mobley, W. C. (2004). Signaling endosome hypothesis: A cellular mechanism for long

distance communication. *Journal of Neurobiology*, *58*(2), 207–216.

https://doi.org/10.1002/neu.10323

Iulita, M. F., & Cuello, A. C. (2016). The NGF Metabolic Pathway in the CNS and its Dysregulation in Down

Syndrome and Alzheimer's Disease. *Current Alzheimer Research*, *13*(1), 53–67.

https://doi.org/10.2174/1567205012666150921100030

Iwano, T., Sobajima, T., Takeda, S., Harada, A., & Yoshimura, S. (2023). The Rab GTPase-binding protein

EHBP1L1 and its interactors CD2AP/CIN85 negatively regulate the length of primary cilia via actin

remodeling. *Journal of Biological Chemistry*, *299*(3). https://doi.org/10.1016/j.jbc.2023.102985

Jung, H., Lee, Y., Lee, S.-H., & Sohn, J.-H. (2023). Auditory or Audiovisual Stimulation Ameliorates Cognitive

Impairment and Neuropathology in ApoE4 Knock-In Mice. *International Journal of Molecular*

Sciences, *24*(2), Article 2. https://doi.org/10.3390/ijms24020938

Kawabata, S. (2022). Signaling abnormality leading to excessive/aberrant synaptic plasticity in Alzheimer's

disease. *Frontiers in Aging Neuroscience*, *14*, 1062519.

https://doi.org/10.3389/fnagi.2022.1062519

Khaire, A. S., Wimberly, C. E., Semmes, E. C., Hurst, J. H., & Walsh, K. M. (2022). An integrated genome and

phenome-wide association study approach to understanding Alzheimer's disease predisposition.

Neurobiology of Aging, *118*, 117–123. https://doi.org/10.1016/j.neurobiolaging.2022.05.011

Kim, W., Underwood, R. S., Greenwald, I., & Shaye, D. D. (2018). OrthoList 2: A New Comparative Genomic

Analysis of Human and Caenorhabditis elegans Genes. *Genetics*, *210*(2), 445–461.

https://doi.org/10.1534/genetics.118.301307

Kirsch, K. H., Georgescu, M.-M., Shishido, T., Langdon, W. Y., Birge, R. B., & Hanafusa, H. (2001). The

Adapter Type Protein CMS/CD2AP Binds to the Proto-oncogenic Protein c-Cbl through a Tyrosine

Phosphorylation-regulated Src Homology 3 Domain Interaction*. *Journal of Biological Chemistry*,

276(7), 4957–4963. https://doi.org/10.1074/jbc.M005784200

Koizumi, K., Wang, G., & Park, L. (2016). Endothelial Dysfunction and Amyloid-β-Induced Neurovascular

Alterations. *Cellular and Molecular Neurobiology*, *36*(2), 155–165.

https://doi.org/10.1007/s10571-015-0256-9

Kovacs, G. G., Robinson, J. L., Xie, S. X., Lee, E. B., Grossman, M., Wolk, D. A., Irwin, D. J., Weintraub, D.,

Kim, C. F., Schuck, T., Yousef, A., Wagner, S. T., Suh, E., Van Deerlin, V. M., Lee, V. M.-Y., &

Trojanowski, J. Q. (2017). Evaluating the Patterns of Aging-Related Tau Astrogliopathy Unravels

Novel Insights Into Brain Aging and Neurodegenerative Diseases. *Journal of Neuropathology &*

Experimental Neurology, *76*(4), 270–288. https://doi.org/10.1093/jnen/nlx007

Kunkle, B. W., Schmidt, M., Klein, H.-U., Naj, A. C., Hamilton-Nelson, K. L., Larson, E. B., Evans, D. A., De

Jager, P. L., Crane, P. K., Buxbaum, J. D., Ertekin-Taner, N., Barnes, L. L., Fallin, M. D., Manly, J. J.,

Go, R. C. P., Obisesan, T. O., Kamboh, M. I., Bennett, D. A., Hall, K. S., … Kukull, W. A. (2021). Novel

Alzheimer Disease Risk Loci and Pathways in African American Individuals Using the African

Genome Resources Panel: A Meta-analysis. *JAMA Neurology*, *78*(1), 102.

https://doi.org/10.1001/jamaneurol.2020.3536

Lambert, J. C., Ibrahim-Verbaas, C. A., Harold, D., Naj, A. C., Sims, R., Bellenguez, C., DeStafano, A. L., Bis, J.

C., Beecham, G. W., Grenier-Boley, B., Russo, G., Thorton-Wells, T. A., Jones, N., Smith, A. V.,

Chouraki, V., Thomas, C., Ikram, M. A., Zelenika, D., Vardarajan, B. N., … Amouyel, P. (2013). Meta-

analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nature*

Genetics, *45*(12), 1452–1458. https://doi.org/10.1038/ng.2802

Latina, V., Atlante, A., Malerba, F., La Regina, F., Balzamino, B. O., Micera, A., Pignataro, A., Stigliano, E.,

Cavallaro, S., Calissano, P., & Amadoro, G. (2023). The Cleavage-Specific Tau 12A12mAb Exerts an

Anti-Amyloidogenic Action by Modulating the Endocytic and Bioenergetic Pathways in Alzheimer's

Disease Mouse Model. *International Journal of Molecular Sciences*, *24*(11), Article 11.

https://doi.org/10.3390/ijms24119683

Lehtonen, S., Tienari, J., Londesborough, A., Pirvola, U., Ora, A., Reima, I., & Lehtonen, E. (2008). CD2-

associated protein is widely expressed and differentially regulated during embryonic

development. *Differentiation*, *76*(5), 506–517. https://doi.org/10.1111/j.1432-0436.2007.00255.x

Li, C., Ruotsalainen, V., Tryggvason, K., Shaw, A. S., & Miner, J. H. (2000). CD2AP is expressed with nephrin

in developing podocytes and is found widely in mature kidney and elsewhere. *American Journal of*

Physiology. Renal Physiology, *279*(4), F785-792. https://doi.org/10.1152/ajprenal.2000.279.4.F785

Liu, R., Collier, J. M., Abdul-Rahman, N.-H., Capuk, O., Zhang, Z., & Begum, G. (2024). Dysregulation of Ion

Channels and Transporters and Blood-Brain Barrier Dysfunction in Alzheimer's Disease and

Vascular Dementia. *Aging and Disease*. https://doi.org/10.14336/AD.2023.1201

Maninger, J.-K., Nowak, K., Goberdhan, S., O'Donoghue, R., & Connor-Robson, N. (2024). Cell type-specific

functions of Alzheimer's disease endocytic risk genes. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, *379*(1899), 20220378. https://doi.org/10.1098/rstb.2022.0378

Marioni, R. E., Harris, S. E., Zhang, Q., McRae, A. F., Hagenaars, S. P., Hill, W. D., Davies, G., Ritchie, C. W.,

Gale, C. R., Starr, J. M., Goate, A. M., Porteous, D. J., Yang, J., Evans, K. L., Deary, I. J., Wray, N. R.,

& Visscher, P. M. (2018). GWAS on family history of Alzheimer's disease. *Translational Psychiatry*,

8(1), Article 1. https://doi.org/10.1038/s41398-018-0150-6

Naj, A. C., Jun, G., Beecham, G. W., Wang, L.-S., Vardarajan, B. N., Buros, J., Gallins, P. J., Buxbaum, J. D.,

Jarvik, G. P., Crane, P. K., Larson, E. B., Bird, T. D., Boeve, B. F., Graff-Radford, N. R., De Jager, P. L.,

Evans, D., Schneider, J. A., Carrasquillo, M. M., Ertekin-Taner, N., … Schellenberg, G. D. (2011).

Common variants at MS4A4/MS4A6E, CD2AP, CD33 and EPHA1 are associated with late-onset

Alzheimer's disease. *Nature Genetics*, *43*(5), 436–441. https://doi.org/10.1038/ng.801

Nakagawa, T., Hasegawa, Y., Uekawa, K., & Kim-Mitsuyama, S. (2017). Chronic kidney disease accelerates

cognitive impairment in a mouse model of Alzheimer's disease, through angiotensin II.

Experimental Gerontology, *87*, 108–112. https://doi.org/10.1016/j.exger.2016.11.012

Nisbet, R. M., Polanco, J.-C., Ittner, L. M., & Götz, J. (2015). Tau aggregation and its interplay with amyloid-

β. *Acta Neuropathologica*, *129*(2), 207–220. https://doi.org/10.1007/s00401-014-1371-2

Novikova, G., Kapoor, M., Tcw, J., Abud, E. M., Efthymiou, A. G., Chen, S. X., Cheng, H., Fullard, J. F., Bendl,

J., Liu, Y., Roussos, P., Björkegren, J. L., Liu, Y., Poon, W. W., Hao, K., Marcora, E., & Goate, A. M.

(2021). Integration of Alzheimer's disease genetics and myeloid genomics identifies disease risk

regulatory elements and genes. *Nature Communications*, *12*(1), Article 1.

https://doi.org/10.1038/s41467-021-21823-y

Ojelade, S. A., Lee, T. V., Giagtzoglou, N., Yu, L., Ugur, B., Li, Y., Duraine, L., Zuo, Z., Petyuk, V., De Jager, P.

L., Bennett, D. A., Arenkiel, B. R., Bellen, H. J., & Shulman, J. M. (2019). Cindr, the Drosophila

Homolog of the CD2AP Alzheimer's Disease Risk Gene, Is Required for Synaptic Transmission and

Proteostasis. *Cell Reports*, *28*(7), 1799-1813.e5. https://doi.org/10.1016/j.celrep.2019.07.041

Pandey, R. S., Kotredes, K. P., Sasner, M., Howell, G. R., & Carter, G. W. (2023). Differential splicing of

neuronal genes in a Trem2*R47H mouse model mimics alterations associated with Alzheimer's

disease. *BMC Genomics*, *24*(1), 172. https://doi.org/10.1186/s12864-023-09280-x

Pensalfini, A., Kim, S., Subbanna, S., Bleiwas, C., Goulbourne, C. N., Stavrides, P. H., Jiang, Y., Lee, J.-H.,

Darji, S., Pawlik, M., Huo, C., Peddy, J., Berg, M. J., Smiley, J. F., Basavarajappa, B. S., & Nixon, R. A.

(2020). Endosomal Dysfunction Induced by Directly Overactivating Rab5 Recapitulates Prodromal

and Neurodegenerative Features of Alzheimer's Disease. *Cell Reports*, *33*(8), 108420.

https://doi.org/10.1016/j.celrep.2020.108420

Qui, M. S., & Green, S. H. (1992). PC12 cell neuronal differentiation is associated with prolonged p21ras

activity and consequent prolonged ERK activity. *Neuron*, *9*(4), 705–717.

https://doi.org/10.1016/0896-6273(92)90033-a

Rajabli, F., Benchek, P., Tosto, G., Kushch, N., Sha, J., Bazemore, K., Zhu, C., Lee, W.-P., Haut, J., Hamilton-

Nelson, K. L., Wheeler, N. R., Zhao, Y., Farrell, J. J., Grunin, M. A., Leung, Y. Y., Kuksa, P. P., Li, D.,

Lucio da Fonseca, E., Mez, J. B., … Naj, A. C. (2023). Multi-ancestry genome-wide meta-analysis of

56,241 individuals identifies LRRC4C, LHX5-AS1 and nominates ancestry-specific loci PTPRK ,

GRB14 , and KIAA0825 as novel risk loci for Alzheimer's disease: The Alzheimer's Disease Genetics

Consortium. *medRxiv: The Preprint Server for Health Sciences*, 2023.07.06.23292311.

https://doi.org/10.1101/2023.07.06.23292311

Renier, N., Wu, Z., Simon, D. J., Yang, J., Ariel, P., & Tessier-Lavigne, M. (2014). iDISCO: A Simple, Rapid

Method to Immunolabel Large Tissue Samples for Volume Imaging. *Cell*, *159*(4), 896–910.

https://doi.org/10.1016/j.cell.2014.10.010

Rouka, E., Simister, P. C., Janning, M., Kumbrink, J., Konstantinou, T., Muniz, J. R. C., Joshi, D., O'Reilly, N.,

Volkmer, R., Ritter, B., Knapp, S., von Delft, F., Kirsch, K. H., & Feller, S. M. (2015). Differential

Recognition Preferences of the Three Src Homology 3 (SH3) Domains from the Adaptor CD2-

associated Protein (CD2AP) and Direct Association with Ras and Rab Interactor 3 (RIN3). *The*

Journal of Biological Chemistry, *290*(42), 25275–25292. https://doi.org/10.1074/jbc.M115.637207

Rueden, C. T., Schindelin, J., Hiner, M. C., DeZonia, B. E., Walter, A. E., Arena, E. T., & Eliceiri, K. W. (2017).

ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics*, *18*(1), 529.

https://doi.org/10.1186/s12859-017-1934-z

Sang, Q., Sun, D., Chen, Z., & Zhao, W. (2018). NGF and PI3K/Akt signaling participate in the ventral motor

neuronal protection of curcumin in sciatic nerve injury rat models. *Biomedicine &*

Pharmacotherapy, *103*, 1146–1153. https://doi.org/10.1016/j.biopha.2018.04.116

Saunders, A., Macosko, E. Z., Wysoker, A., Goldman, M., Krienen, F. M., de Rivera, H., Bien, E., Baum, M.,

Bortolin, L., Wang, S., Goeva, A., Nemesh, J., Kamitaki, N., Brumbaugh, S., Kulp, D., & McCarroll, S.

A. (2018). Molecular Diversity and Specializations among the Cells of the Adult Mouse Brain. *Cell*,

174(4), 1015-1030.e16. https://doi.org/10.1016/j.cell.2018.07.028

Schaefer, A., van Duijn, T. J., Majolee, J., Burridge, K., & Hordijk, P. L. (2017). Endothelial CD2AP Binds the

Receptor ICAM-1 To Control Mechanosignaling, Leukocyte Adhesion, and the Route of Leukocyte

Diapedesis In Vitro. *The Journal of Immunology*, *198*(12), 4823–4836.

https://doi.org/10.4049/jimmunol.1601987

Schwarz, K., Simons, M., Reiser, J., Saleem, M. A., Faul, C., Kriz, W., Shaw, A. S., Holzman, L. B., & Mundel,

P. (2001, December 1). *Podocin, a raft-associated component of the glomerular slit diaphragm,*

interacts with CD2AP and nephrin. American Society for Clinical Investigation.

https://doi.org/10.1172/JCI12849

Shen, R., Zhao, X., He, L., Ding, Y., Xu, W., Lin, S., Fang, S., Yang, W., Sung, K., Spencer, B., Rissman, R. A.,

Lei, M., Ding, J., & Wu, C. (2020). Upregulation of RIN3 induces endosomal dysfunction in

Alzheimer's disease. *Translational Neurodegeneration*, *9*(1), 26. https://doi.org/10.1186/s40035-

020-00206-1

Sherva, R., Zhang, R., Sahelijo, N., Jun, G., Anglin, T., Chanfreau, C., Cho, K., Fonda, J. R., Gaziano, J. M.,

Harrington, K. M., Ho, Y.-L., Kremen, W. S., Litkowski, E., Lynch, J., Neale, Z., Roussos, P., Marra, D.,

Mez, J., Miller, M. W., … Logue, M. W. (2023). African ancestry GWAS of dementia in a large

military cohort identifies significant risk loci. *Molecular Psychiatry*, *28*(3), 1293–1302.

https://doi.org/10.1038/s41380-022-01890-3

Shi, Y., Liu, Z., Shen, Y., & Zhu, H. (2018). A Novel Perspective Linkage Between Kidney Function and

Alzheimer's Disease. *Frontiers in Cellular Neuroscience*, *12*.

https://www.frontiersin.org/articles/10.3389/fncel.2018.00384

Shih, N. Y., Li, J., Karpitskii, V., Nguyen, A., Dustin, M. L., Kanagawa, O., Miner, J. H., & Shaw, A. S. (1999).

Congenital nephrotic syndrome in mice lacking CD2-associated protein. *Science (New York, N.Y.)*,

286(5438), 312–315. https://doi.org/10.1126/science.286.5438.312

Shulman, J. M., Imboywa, S., Giagtzoglou, N., Powers, M. P., Hu, Y., Devenport, D., Chipendo, P., Chibnik, L.

B., Diamond, A., Perrimon, N., Brown, N. H., De Jager, P. L., & Feany, M. B. (2014). Functional

screening in Drosophila identifies Alzheimer's disease susceptibility genes and implicates Tau-

mediated mechanisms. *Human Molecular Genetics*, *23*(4), 870–877.

https://doi.org/10.1093/hmg/ddt478

Srivatsan, S., Swiecki, M., Otero, K., Cella, M., & Shaw, A. S. (2013). CD2-Associated Protein Regulates

Plasmacytoid Dendritic Cell Migration, but Is Dispensable for Their Development and Cytokine

Production. *The Journal of Immunology*, *191*(12), 5933–5940.

https://doi.org/10.4049/jimmunol.1300454

Stauffer, W., Sheng, H., & Lim, H. N. (2018). EzColocalization: An ImageJ plugin for visualizing and

measuring colocalization in cells and organisms. *Scientific Reports*, *8*(1), Article 1.

https://doi.org/10.1038/s41598-018-33592-8

Tallini, Y. N., Shui, B., Greene, K. S., Deng, K.-Y., Doran, R., Fisher, P. J., Zipfel, W., & Kotlikoff, M. I. (2006).

BAC transgenic mice express enhanced green fluorescent protein in central and peripheral

cholinergic neurons. *Physiological Genomics*, *27*(3), 391–397.

https://doi.org/10.1152/physiolgenomics.00092.2006

Tesi, N., van der Lee, S. J., Hulsman, M., Jansen, I. E., Stringa, N., van Schoor, N. M., Scheltens, P., van der

Flier, W. M., Huisman, M., Reinders, M. J. T., & Holstege, H. (2020). Immune response and

endocytosis pathways are associated with the resilience against Alzheimer's disease. *Translational*

Psychiatry, *10*(1), Article 1. https://doi.org/10.1038/s41398-020-01018-7

Tsui, C. C., & Pierchala, B. A. (2008). CD2AP and Cbl-3/Cbl-c Constitute a Critical Checkpoint in the

Regulation of Ret Signal Transduction. *Journal of Neuroscience*, *28*(35), 8789–8800.

https://doi.org/10.1523/JNEUROSCI.2738-08.2008

van Bree, E. J., Guimarães, R. L. F. P., Lundberg, M., Blujdea, E. R., Rosenkrantz, J. L., White, F. T. G.,

Poppinga, J., Ferrer-Raventós, P., Schneider, A.-F. E., Clayton, I., Haussler, D., Reinders, M. J. T.,

Holstege, H., Ewing, A. D., Moses, C., & Jacobs, F. M. J. (2022). A hidden layer of structural

variation in transposable elements reveals potential genetic modifiers in human disease-risk loci.

Genome Research, *32*(4), 656–670. https://doi.org/10.1101/gr.275515.121

Vandal, M., Institoris, A., Korin, B., Gunn, C., Lee, S., Lee, J., Bourassa, P., Mishra, R. C., Peringod, G., Jiang,

Y., Hirai, S., Belzil, C., Reveret, L., Tremblay, C., Hashem, M., Elias, E., Meilandt, B., Foreman, O.,

Rouse-Girma, M., … Nguyen, M. D. (2022). *The Alzheimer risk factor CD2AP causes dysfunction of*

the brain vascular network (p. 2020.12.10.419598). bioRxiv.

https://doi.org/10.1101/2020.12.10.419598

Welsch, T., Endlich, N., Gökce, G., Doroshenko, E., Simpson, J. C., Kriz, W., Shaw, A. S., & Endlich, K. (2005).

Association of CD2AP with dynamic actin on vesicles in podocytes. *American Journal of Physiology.*

Renal Physiology, *289*(5), F1134-1143. https://doi.org/10.1152/ajprenal.00178.2005

Welsch, T., Endlich, N., Kriz, W., & Endlich, K. (2001). CD2AP and p130Cas localize to different F-actin

structures in podocytes. *American Journal of Physiology-Renal Physiology*, *281*(4), F769–F777.

https://doi.org/10.1152/ajprenal.2001.281.4.F769

Wittmann, C. W., Wszolek, M. F., Shulman, J. M., Salvaterra, P. M., Lewis, J., Hutton, M., & Feany, M. B.

(2001). Tauopathy in Drosophila: Neurodegeneration Without Neurofibrillary Tangles. *Science*,

293(5530), 711–714. https://doi.org/10.1126/science.1062382

Xu, W., Fang, F., Ding, J., & Wu, C. (2018a). Dysregulation of Rab5-mediated endocytic pathways in

Alzheimer's disease. *Traffic*, *19*(4), 253–262. https://doi.org/10.1111/tra.12547

Xu, W., Fang, F., Ding, J., & Wu, C. (2018b). Dysregulation of Rab5-mediated endocytic pathways in

Alzheimer's disease. *Traffic (Copenhagen, Denmark)*, *19*(4), 253–262.

https://doi.org/10.1111/tra.12547

Xue, Y.-Y., Chen, Y.-H., Lin, R.-R., Huang, H.-F., Wu, Z.-Y., & Tao, Q.-Q. (2022). Alzheimer's disease

susceptibility locus in CD2AP is associated with increased cerebrospinal fluid tau levels in mild

cognitive impairment. *Neuroscience Letters*, *771*, 136419.

https://doi.org/10.1016/j.neulet.2021.136419

Ye, H., Kuruvilla, R., Zweifel, L. S., & Ginty, D. D. (2003). Evidence in Support of Signaling Endosome-Based

Retrograde Survival of Sympathetic Neurons. *Neuron*, *39*(1), 57–68.

https://doi.org/10.1016/S0896-6273(03)00266-6

Yoshida, S., Wei, X., Zhang, G., O'Connor, C. L., Torres, M., Zhou, Z., Lin, L., Menon, R., Xu, X., Zheng, W.,

Xiong, Y., Otto, E., Tang, C.-H. A., Hua, R., Verma, R., Mori, H., Zhang, Y., Hu, C.-C. A., Liu, M., … Qi,

L. (2021, April 1). *Endoplasmic reticulum–associated degradation is required for nephrin*

maturation and kidney glomerular filtration function. American Society for Clinical Investigation.

https://doi.org/10.1172/JCI143988

Zhang, C.-Y., He, F.-F., Su, H., Zhang, C., & Meng, X.-F. (2020). Association between chronic kidney disease

and Alzheimer's disease: An update. *Metabolic Brain Disease*, *35*(6), 883–894.

https://doi.org/10.1007/s11011-020-00561-y

FIGURE LEGENDS

FIGURE 1

CD2AP mRNA is expressed in Ntrk1 (TRKA)-expressing populations of cholinergic neurons of the

adult mouse brain. Summary of single-cell RNA-sequencing data from the adult mouse brain,

using the Dropviz webtool (Saunders et al., 2018). The top ten, highest *CD2AP*-expressing

populations are shown on the left (magenta) and corresponding expression levels of *Ntrk1* on the

right (cyan).

FIGURE 2

CD2AP protein is expressed in ChAT-positive neurons in vivo. (a) Representative confocal image

of ChAT-GFP mouse brain 20um section, stained with primary antibodies to ChAT (yellow) and

Cd2ap (magenta). (b) Inset indicated by yellow square; diagonal band of broca (vDB). CD2AP

signal is delectated in adult (ChAT-GFP+/ChAT-antibody+) cholinergic neurons.

FIGURE 3

Immunofluorescence analysis of ChAT-GFP neurons in regions of interest within the mouse brain.

Representative immunofluorescence images from 20um-thick frozen tissue sections. ChAT-GFP

signal (left column, cyan) was detected in ChAT-lineage neurons (ChAT-GFP+/αChAT-) in Cortex

(top row), Striatum, Brocca (vDB) and Substantia Innominata (bottom row). Fluorescence signal

from α (second column, yellow) was observed in neuron soma of adult (ChAT-GFP+/ αChAT+)

cholinergic neurons in Striatum (second row), Brocca (third row) and Substantia Innominata

(bottom row). Fluorescence signal of CD2AP (third column, magenta), was observed in adult

cholinergic neurons (ChAT-GFP+/ αChAT+), but not in cholinergic-lineage neurons (ChAT-

GFP+/αChAT-). Cyan arrows indicate ChAT-GFP+/αChAT- neuron cell bodies; magenta arrows

indicate (cortical) vascular αCD2AP; white arrows indicate ChAT-GFP+/ αChAT+ neuron cell

bodies that are also CD2AP+.

FIGURE 4

Quantification of CD2AP protein expression in cholinergic neurons. (a) Mean, regional expression

levels (expressed as integrated density mean ± SD) of Cd2ap in individual mice; violin clusters

reflect raw data point distribution and mean values ± SD. (b) Mean CD2AP immunofluorescence

signals in ChAT-GFP+ soma from indicated brain regions, from group data (N=3 animals). One-

way analysis of variance (ANOVA) with Tukey post-hoc analysis revealed Cd2ap expression in

the basal, striatal and optic regions of the mouse brain, with cortex serving as a background-

signal control. ****$p<0.0001$.

FIGURE 5

CD2AP expression in ChAT-GFP+ adult mouse brain projections. (a) Representative, whole-brain

imaging of ChAT-GFP+, sagittal sections (150um) embedded in agarose. (a; i) ChAT/GFP+

cholinergic neuron cell bodies originating in the basal forebrain; (a; ii) ChAT/GFP+ axonal

projections (white arrow) from basal forebrain on trajectory towards hippocampal/iso-cortical

regions. (b-c) Representative, regional examination of Cd2ap expression with

immunofluorescence analysis of ChAT-GFP+ neuron cell bodies in the basal forebrain (b; white

arrows) and staining patterning consistent with ChAT/GFP+ axonal projections (a; i, ii). Yellow box

indicates low magnification location of inset image at high magnification; cyan indicates

endogenous ChAT/GFP+ fluorescence signal; magenta indicates fluorescence signal from anti-

Cd2ap primary antibody; white arrows indicate areas of overlap between ChAT/GFP+ and

Cd2ap+.

FIGURE 6

CD2AP and Rab5 co-expression in ChAT/GFP-positive (ChAT/GFP+) neurons in adult mouse

brains (aged 10 months). (a) Representative immunofluorescence image from the vDB region

(Basal forebrain; 20um), demonstrating the colocalization and co-expression of CD2AP and Rab5

endosomal formations within and in proximity to cell bodies of ChAT/GFP+ cholinergic neurons.

White arrows indicate punctate endosomal formations localizing around/within the ChAT/GFP+

cell body. (b, top) Representative region of interest (ROI) outlined in cyan, indicating the outline of

a single, ChAT/GFP+ neuron cell body; area within ROI is the max projection of colocalized

Cd2ap/Rab5+ endosomes, a single example of which is indicated with a red arrow and circle. (b,

bottom) Representative, two-dimensional intensity histogram demonstrating the quantification

(correlation) of Cd2ap/Rab5+ endosomes.

TABLE 1. Antibodies used.

Primary Antibody	Source	Cat#	Lot#	Species/clonality	[working] IHC/IF (20um)	[working] IHC/IF (150um)
Anti-Cd2ap	Proteintech	51046-1-AP	23812	Rabbit polyclonal (IgG)	1:800	1:200
Anti-TrkA (Rat)	R&D	AF1056	VFA0118091	Goat polyclonal (IgG)	1:200	1:100
Anti-Choline Acetyltransferase (ChAT)	Millipore/Sigma	AB144P	3926290	Goat polyclonal (IgG)	1:200	1:100
Anti-Green Fluorescence Protein	Aves Labs	GFP-1020	GFP917979	Chicken (IgY)	1:1000	(1:500)10mg/mL
Anti-Rab5	Cell Signaling Technology	46449	3	Mouse monoclonal (IgG1; clone E6N8S)	1:200	N/A

Secondary Antibody/fluorophore	Source	Cat#	Lot#	Species	[working] IHC/IF (20um)	[working] IHC/IF (150um)
Alexa647		711-605-152	165592	Donkey-Anti-Rabbit	1:500	1:250
Cy3		705-165-147	130439	Donkey-Anti-Goat	1:500	1:250
Alexa488		703-545-155	139424	Donkey-Anti-Chicken	N/A	1:250
Cy3		705-165-147	130439	Donkey-Anti-Goat	N/A	1:250
Alexa647	Jackson ImmunoResearch	703-605-155	138591	Donkey-Anti-Chicken	N/A	1:250
Cy3		711-165-152	166393	Donkey-Anti-Rabbit	N/A	1:250
Alexa647		705-605-147	160039	Donkey-Anti-Goat	1:500	N/A
Cy3		715-165-151	130991	Donkey-Anti-Mouse	1:500	N/A
Alexa488		711-545-152	161527	Donkey-Anti-Rabbit	1:500	N/A

Other Reagents	Source	Cat#	Lot#	Species	[working] IHC/IF (20um)	[working] IHC/IF (150um)
Normal Donkey Serum	Jackson ImmunoResearch	167385	017-000-121	4%	8%	017-000-121
Mounting Medium with DAPI	Abcam	ab104139	1043154-3	n/a	n/a	ab104139

TABLE 2. Summary of GWAS studies from 2013 to present (June 2024) implicating Cd2ap in the pathogenesis and/or susceptibility to AD in humans.

YEAR	CITATION	OUTCOME	rsID
2023	Sherva R, Zhang R, Sahelijo N, et al. African ancestry GWAS of dementia in a large military cohort identifies significant risk loci. Mol Psychiatry. 2023;28(3):1293-1302.	Novel, significant SNPs near known AD risk genes TREM2, CD2AP, and ABCA7; peak variants in these genes differed from those previously reported in European and African ancestry cohorts.	rs7738720
2023	Charisis S, Lin H, Ray R, et al. Obesity impacts the expression of Alzheimer's disease-related genes: The Framingham Heart Study. Alzheimers Dement. 2023;19(8):3496-3505.	Gene expression was measured for 74 AD-related genes, derived by integrating genome-wide association study results with functional genomics data from the Framingham Heart Study. Obesity metrics were associated with the expression of 21 AD-related genes. The strongest associations were observed with CLU, CD2AP, KLC3, and FCER1G.	
2022	Gao S, Hao JW, Zhao YN, et al. An updated analysis of the association between CD2-associated protein gene rs9349407 polymorphism and Alzheimer's disease in Chinese population. Front Neuroinform. 2022;16:1006164.	Comparison of 11 independent studies from 8 articles exploring the correlation between rs9349407 variation and AD in Chinese. Additional meta-analysis based on fixed and random effect models were compiled to perform heterogeneity test. Use of additive model, dominant model, and recessive model for subgroup analysis demonstrated that the CD2AP rs9349407 polymorphism increases AD susceptibility in Chinese populations.	rs9349407
2022	Khaire AS, Wimberly CE, Semmes EC, Hurst JH, Walsh KM. An integrated genome and phenome-wide association study approach to understanding Alzheimer's disease predisposition. Neurobiol Aging. 2022;118:117-123.	"Phenome-wide association" of 23 known, LOAD-associated SNPs and 4:1 matched control SNPs using UK Biobank data; associations confirmed known LOAD risk loci near PICALM, CD2AP, SPI1, and NDUFAF6, and identified a novel risk locus in epidermal growth factor receptor.	
2022	Bellenguez, C., Küçükali, F., Jansen, I.E. et al. New insights into the genetic etiology of Alzheimer's disease and related dementias. Nat Genet 2022; 54:412–436	Two-stage GWAS of 111,326 clinically diagnosed/'proxy' AD cases and 677,663 controls; found 75 risk loci, including 42 novel sites at the time of analysis. Pathway enrichment analyses confirmed involvement of amyloid/tau pathways and highlighted microglia implication.	rs7767350 6:47517390:T:C
2021	Li Y, Laws SM, Miles LA, et al. Genomics of Alzheimer's disease implicates the innate and adaptive immune systems. Cell Mol Life Sci. 2021;78(23):7397-7426.	Comprehensive review of GWAS studies (with meta-analyses from 2013 up to 2021)	
2021	Kunkle BW, Schmidt M, Klein HU, et al. Novel Alzheimer Disease Risk Loci and Pathways in African American Individuals Using the African Genome Resources Panel: A Meta-analysis. JAMA Neurol. 2021;78(1):102-113.	Of 25 known AD loci in non-Hispanic, White individuals, only APOE, ABCA7, TREM2, BIN1, CD2AP, FERMT2, and WWOX were significantly associated with AD in African American individuals.	
2021	Schwartzentruber J, Cooper S, Liu JZ, Barrio-Hernandez I, Bello E, Kumasaka N et al. Genome-wide meta-analysis, fine-mapping and integrative prioritization implicate new Alzheimer's disease risk genes. Nat Genet 2021;53(3):392–402.	AD GWAS with focus on causal genes and variants; confirmed recently discovered AD genes (CD2AP) also the top gene within 500 kb by network score and AD Risk.	rs9381563 6:47432637:C:T
2019	Jansen IE, Savage JE, Watanabe K, Bryois J, Williams DM, Steinberg S et al. Genome-wide meta-analysis identifies new loci and functional pathways influencing Alzheimer's disease risk. Nat Genet 2019;51(3):404–413.	AD GWAS of clinically diagnosed AD and AD-by-proxy cases; meta-analysis results revealed 29 risk loci, implicating 215 potential causative genes (CD2AP); analyses support a role for lipid processing and degradation of amyloid precursor proteins in AD pathogenesis.	rs9381563 6:47432637:C:T
2019	Kunkle BW, Grenier-Boley B, Sims R, Bis JC, Damotte V, Naj AC et al. Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates Abeta, tau, immunity and lipid processing. Nat Genet 2019;51(3):414–430.	AD GWAS; confirmed eQTLs for genes in previously identified loci, including *CD2AP* and the *CD2AP* locus	rs9473117 6:47431284:A:C
2018	Marioni, R.E., Harris, S.E., Zhang, Q. et al. GWAS on family history of Alzheimer's disease. Transl Psychiatry 2018; 8, 99.	Self-report of parental history of Alzheimer's dementia for GWAS of 314,278 participants from UK Biobank; meta-analysis with published consortium data revealed three new loci, relevant for AD and neurodegeneration, including CD2AP.	rs9381563 6:47432637:C:T
2013	Lambert JC, Ibrahim-Verbaas CA, Harold D, Naj AC, Sims R, Bellenguez C et al. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. Nat Genet 2013; 45(12):1452–1458.	Meta-analysis of 17,008 cases and 37,154 controls, followed by replication in an independent cohort; confirmed previous GWAS findings (CD2AP) and identified 11 novel AD susceptibility loci: HLA-DRB5-DRB1, SORL1, PTK2B, SLC24A4, ZCWPW1, CELF1, NME8, FERMT2, CASS4, INPP5D, and MEF2C	rs10948363 6:47487762:A:G

(a)

Animal 1

Animal 2

Animal 3

(b)

(a)

ChAT/GFP+

Rab5

Cd2ap

ChAT/GFP+
Rab5
Cd2ap

10μm

Rab5
Cd2ap

10μm

63X/Zoom2.0

(b)

Region of Interest (ROI)

CD2AP/Rab5
Colocalization

Colocalization

Pearson R value: 0.80
Li's ICQ value: 0.241
Costes P-Value: 1.00