**<u>Satellite glial contact enhances differentiation and maturation of human induced sensory</u>**

**<u>neurons</u>**

Chelsey J. LeBlang[1,2], Maria F. Pazyra-Murphy[1], Elizabeth Silagi[1,2], Srestha Dasgupta[1,2],
Marianna Tsolias[1], Toussaint Miller[1], Veselina Petrova[2,3], Shannon Zhen[3], Kathryn Gerrish[4],
Vukasin Jovanovic[4], David Castellano[4], Pinar Ormanoglu[4], Carlos Tristan[4], Clifford J. Woolf[2,3],
Ozge Tasdemir-Yilmaz[1,2,5], Rosalind A. Segal*[1,2,5].

[1]Department of Cancer Biology, Dana Farber Cancer Institute, Boston, MA 02215

[2]Department of Neurobiology, Harvard Medical School, Boston, MA 02215

[3]F.M. Kirby Neurobiology Center, Boston Children's Hospital, Boston, MA 02215

[4]National Center for Advancing Translational Science, National Institute of Health, Bethesda,
MD, 20892

[5]co-senior author

*corresponding author: rosalind_segal@dfci.harvard.edu

**<u>Abstract:</u>** Induced sensory neurons (iSNs) generated from pluripotent stem cells are used to
model human peripheral neuropathies, however current differentiation protocols produce
sensory neurons with an embryonic phenotype. Peripheral glial cells contact sensory neurons
early in development and contribute to formation of the canonical pseudounipolar morphology,
but these signals are not encompassed in current iSN differentiation protocols. Here, we show
that terminal differentiation of iSNs in co-culture with rodent Dorsal Root Ganglion satellite glia
(rSG) advances their differentiation and maturation. Co-cultured iSNs develop a pseudounipolar
morphology through contact with rSGs. This transition depends on semaphorin-plexin guidance
cues and on glial gap junction signaling. In addition to morphological changes, iSNs terminally
differentiated in co-culture exhibit enhanced spontaneous action potential firing, more mature
gene expression, and increased susceptibility to paclitaxel induced axonal degeneration. Thus,
iSNs differentiated in coculture with rSGs provide a better model for investigating human
peripheral neuropathies.

**Keywords:** Satellite Glia, Pseudounipolar, iSN, DRG, Neuropathy, CIPN.

## Introduction:

Dorsal Root Ganglion (DRG) neurons are responsible for communicating somatosensory input from the outside world to the central nervous system[1]. They relay these signals through a single specialized T-shaped axon, which projects to both the periphery and the spinal cord[1–3]. DRG neurons include multiple phenotypic subtypes, which respond to specific stimuli and exhibit distinct patterns of gene expression[1,4,5]. DRG neurons are subject to a large number of disorders including hereditary neuropathies, diabetic neuropathy, and chemotherapy induced peripheral neuropathy[6], therefore developing appropriate human model systems is a major goal for researchers.

The use of differentiated human induced pluripotent stem cells (hiPSCs) has become a gold standard for studying disease pathology in relevant cell types and is increasingly accessible[7,8]. Over the past decade, several protocols have been generated to differentiate standard and patient derived hiPSCs into somatosensory neurons (iSNs)[9–15]. Researchers have developed these protocols and utilized iSNs to study neuropathic pain signaling[10,16–18], mechanisms underlying debilitating and painful neuropathies[19–24], and development of treatments for these disorders[17,23,25,26].

During iSN differentiation, iPSCs undergo changes mimicking embryonic development, altering their transcriptional program to progressively resemble neuroectoderm, neural crest progenitor cells, and ultimately sensory neurons, through treatment with combinations of small molecules over time and/or genetic manipulation[9–15]. Cells produced using these protocols express many hallmark proteins associated with sensory neuron identity, however iSNs exhibit an embryonic phenotype, different from that of postnatal neurons[4]. In the developing mammalian embryo, DRG neurons initially are bipolar in morphology and undergo a series of morphological changes leading to a pseudounipolar morphology before birth[27–29]. Current differentiation protocols generate iSNs that are predominantly bipolar in morphology, reflecting

the early embryonic state, or a multipolar morphology, similar to that observed following an

acute injury *in vivo*[30–32]. While developing sensory neurons alter their transcriptome over the

course of prenatal development and express transcription factors that contribute to neuronal

subtyping[1,33], iSNs remain functionally unspecialized. Therefore, modifications of iSN protocols

to achieve a more differentiated phenotype would greatly advance efforts to study postnatal

homeostasis and disease at a molecular level, in a way that is more translatable to human

health or disease conditions.

We have sought to leverage developmental cues to advance the maturation of differentiated

iSNs. While specific mechanisms underlying pseudounipolarization and subtyping are unknown,

*in vitro* studies show that primary sensory neurons co-cultured with mixed DRG glia, including

Schwann cells (SCs) and satellite glia (SGs), become pseudounipolar[34–36]. Yet, iSNs

differentiated in a co-culture with induced or primary SCs, undergo no morphological changes,

aside from myelination[16,37,38]. *In vivo,* intimate contact is initiated between sensory neurons and

satellite glia during the early developmental period[39,40], therefore we hypothesized that satellite

glia contribute to DRG sensory neuron maturation, and that terminal differentiation of iSNs in

coculture with satellite glial cells would advance this. Here, we show that physical contact

between satellite glia and iSNs during differentiation significantly advances morphological and

physiological development compared to those differentiated alone, and both semaphorin-plexin

signaling and gap junction signaling contribute to this maturation process. We also tested the

utility of this model and demonstrate that cocultured iSNs are more susceptible to toxic

neuropathy compared to iSNs alone. This work not only provides an advanced iSN model for

use by researchers in future studies, but also provides insight into mechanisms underlying

sensory neuron development.

**Results:**

**Human iSNs differentiated in coculture with embryonic rodent DRG glia exhibit advanced**
**morphology:**

Currently, differentiation protocols used to generate hiSNs from iPSCs produce sensory

neurons with predominantly immature morphologies[9–15]. While the molecular mechanisms

underlying embryonic transition to pseudounipolar morphology are unknown, our data (Fig.

S1A-B) and others[36,40] suggest that satellite glia may play a pivotal role in morphological

maturation, through contact with sensory neurons during this developmental period. To test this,

we used the Deng et al protocol[15] to terminally differentiate human induced neural crest

precursors (iNCPs) alone, or together with peripheral glia prepared from rat E14.5 DRGs (Fig.

1A). After one week, cells were fixed and stained to visualize neuronal morphology (TUJ1,

cytoskeletal marker, green) and contact with rDRG satellite glia (p75, NGF receptor, magenta)

(Fig 1B).

Morphologies of human neurons were identified and quantified using the rules specified in

the methods section. Morphology categories compared consisted of "immature" (including

unipolar and bipolar cells) "multipolar", "Pseudounipolar Transition (PUT)" (including PUT-Bulge,

PUT-Bell, and PU-Associated Branch), and "pseudounipolar (PU)" (including PU-Short and PU-

Long) (Fig. 1B). Frequencies observed for each morphology category (Table 1) indicate that

iSNs differentiated in coculture include significantly more transitioning and pseudounipolar cells

(PUT *adj. residual* = 7.215, PU *adj. residual* = 6.350) and fewer cells with an immature

morphology (immature *adj. residual* = -5.555, multipolar *adj. residual* = -3.653) compared to

iSNs differentiated alone (PUT *adj. residual* = -7.215, PU *adj. residual* = -6.350 ,immature *adj.*

*residual* = 5.555, multipolar *adj. residual* = 3.653) (Fig. 1C). No differences were observed

between iSNs derived from different iPSC cell lines. These results suggest that terminal

differentiation in co-culture with DRG glia enhances the maturation of sensory neuron

morphology.

To determine whether the ability of peripheral glia to promote a mature, pseudounipolar

morphology is restricted to only one of the protocols used to generate iSNs, we also tested our

co-culture protocol using RealDRG[TM] neurons from Anatomic. RealDRGs were terminally differentiated in coculture or alone, using Senso-MM maturation media by Anatomic as a control. RealDRGs cocultured with rDRG Glia in Deng et al., 2023[15] NCATS Maturation Media (NMM) undergo extensive morphologic changes, although the frequency of transition to pseudounipolar morphology was reduced compared to Deng et al. 2023[15] generated iSNs (Fig S1C-D). Treatment of RealDRGs cultures alone with NMM did not impact observed cell health or morphology but did result in the presence of neural precursor cells, which are not typically observed in RealDRG cultures maintained in Anatomic Senso-MM Media (Fig S1C-D).

| | Morphology Frequencies | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | Immature | | Multipolar | Pseudounipolar Transition (PUT) | | | Pseudounipolar (PU) | | |
| Condition | Unipolar | Bipolar | Multipolar | PUT-Bulge | PUT-Bell | PUT-AB | PU-Short | PU-Long | Totals |
| Coculture | 30 | 22 | 151 | 43 | 31 | 9 | 34 | 9 | 329 |
| iSN Alone | 0 | 101 | 174 | 11 | 1 | 0 | 0 | 0 | 287 |
| Totals | 153 | | 325 | 95 | | | 43 | | 616 |

Table 1. Raw frequency of morphology subtypes observed in co-cultured iSNs vs. iSNs terminally differentiated alone. $X^2$ frequency statistic calculated for major morphology groups: Immature, Multipolar PUT, and PU. ($X^2$= 111, df=3, p<0.001).

**Human iSNs differentiated in coculture with embryonic rodent DRG glia exhibit advanced physiology:**

We next assessed the impact of the coculture strategy on development of spontaneous electrophysiologic signaling in iSNs. DRG sensory neurons can transmit sensory input to the CNS via single action potential spikes, as well as synchronous action potential bursts[41,42]. While spontaneous activity is relatively low[41-45] increased spiking and bursting is observed in models of neuropathic pain both *in vivo*[41,42,45-48] and *in vitro*[42,43,45,49]. Therefore, it is crucial that iSNs develop physiologic signaling capability to study the pathogenesis of neuropathies and develop potential treatments.

Characterization of action potential firing using Multi-Electrode Array (MEA) recordings

shows that iSNs generated with the Deng et al. 2023[15] protocol exhibit nearly undetectable

baseline spontaneous activity after the typical two-week maturation period. Since peripheral

satellite glial cells play an important role in the regulation of somatosensory neuron firing[50–53],

we tested the hypothesis that iSNs differentiated in coculture would exhibit more mature

electrophysiologic properties by measuring spontaneous neural activity of iSNs differentiated in

coculture, or alone, using Axion MEA plates containing 16 electrodes/well. Activity was

measured on the Maestro Pro recording system for 5 minutes weekly over the span of 5-7

weeks, and neuronal spiking and bursting was analyzed using AxIS software. Overall,

cocultured iSNs became robustly active earlier than iSNs differentiated alone, as indicated by a

higher number of active electrodes beginning around week 3 of maturation (Fig. S1E-H). Data

was analyzed when an average of 7 or more electrodes were active per well in both iSN culture

conditions (Fig. 1E), this occurred between weeks 5-7 across 4 experimental replicates (Fig.

S1E-G). Glia alone were inactive and do not contribute to signals recorded in the coculture (Fig.

1E, Fig. S1E-H).

Mature iSNs differentiated in coculture were more active than iSNs differentiated alone (Fig.

1E), with a significantly higher number of spikes elicited over the recording period (Mann-

Whitney Test, p= 0.0162) (Fig. 1F), as well as a significantly higher Mean Firing Rate (MFR)

(Mann-Whitney Test, p= 0.013) (Fig. 1G). The average MFR observed in cocultured iSNs, 1.657

Hz, is comparable to MFRs previously reported in primary DRG neurons[54,55] and iSNs cultured

long-term[56]. No significant difference was observed in the total number of bursts recorded,

however the range was quite different, with a maximum burst number of 971 detected in

coculture and only 341 in iSNs alone (Fig. 1H). Interestingly, cocultured iSNs elicited

significantly longer burst trains (Mann-Whitney Test, p= <0.0001) (Fig. 1I), with a significantly

shorter period between bursts (Mann-Whitney Test, p= 0.0376) (Fig. 1J), thus spending more

time in a burst state. These results suggest that cocultured iSNs are more physiologically

mature than iSNs cultured alone.

**iNCPs terminally differentiated in co-culture are advanced in neuronal differentiation:**

To identify the transcriptional programs that promote iSN maturation in coculture, we carried

out RNA sequencing of iSNs terminally differentiated in coculture or alone. mRNA reads from

cocultures were aligned to both the human and rat genome to distinguish mRNAs derived from

hiSNs and rDRG glia respectively. PCA analysis confirmed that iSNs terminally differentiated in

co-culture cluster separately from iSNs differentiated alone, illustrating transcriptional

differences between these two groups (Fig. S2). Differential gene expression analysis yielded

476 significantly (p<0.05) upregulated genes in cocultured iSNs (Fig. 2A, Table S1). Genes

implicated in posterior thoracic neural crest development[57,58] (*HoxB9*, p adj.=6.922e$^{-15}$) and

proliferation[59] (*ID3,* p adj.=1.306e$^{-09}$ ) were among the top genes upregulated. Importantly, the

most highly upregulated gene, *NFIX* (p adj.=1.025e$^{-16}$), is associated with onset of neuronal

differentiation and quiescence of neural stem cells [60–62] signaling a shift towards a more mature

neuronal phenotype. Interestingly, *NFIX*, has also been shown to regulate neuroblast branching

in adult neurogenesis[63]. Additional changes in expression of genes underlying neuronal

morphogenesis include significant upregulation of *GPRIN3* (p adj.=6.437e$^{-12}$) and *RAI14* (p adj.=

4.910e$^{-09}$), proteins essential to axonal outgrowth[64,65] and dendritic development[66,67]

respectively, through regulation of actin cytoskeletal structure. These changes illustrate potential

transcriptional differences that underly the morphological maturation in cocultured iSNs.

Gene Ontology (GO) analysis provided further information on the upregulated gene

programs, with 185 significantly upregulated gene terms in cocultured iSNs compared to iSNs

alone (Fig. 2B, Table S2). The most significantly upregulated biological processes include

increased Bone Morphogenesis Pathway (BMP) signaling and anterior/posterior pattern

specification, which confirm commitment to the neural crest cell lineage[33,68] (Table S2). Several

biological processes related to neuronal differentiation and morphogenesis were also

significantly upregulated, including neural precursor proliferation, neuronal differentiation, and

neuronal fate commitment, (Fig 2B). Interestingly, we also observed a significant upregulation of

genes associated with axonogenesis (Fig 2B). These genes specifically influence axonal

specification, outgrowth, guidance, branching, and synaptogenesis (Table 2, Table S2), and

may potentially drive the pseudounipolarization process.

Functional maturation was also evident with the upregulated expression of biological

processes and cellular components related to synaptic structure and development, including

synaptic membrane proteins, receptors, and ion channels localized to synapses in DRG

neurons[4,69] (Fig 2B). Specifically, we observed increased expression of genes encoding sodium

ion channels (*SCN7a/SCN10a*) (Table S2)*. SCN10a* is highly expressed in DRG nociceptors,

encoding the sodium channel Nav1.8[4,70], critical for nociception[71–73]. We also observed

upregulation of muscarinic receptors (*CHRM2/CHRM3*) essential to cholinergic signaling in

DRG neurons[74–76], and increased expression of potassium ion channels (*KCNA1/KCNQ5)* that

limit action potential firing and maintain resting membrane potential in adult DRG sensory

neurons[77–79] (Table S2)*.* Lastly, we observed significant upregulation in AMPAR gene *GRIA2*

and GABA receptors (*GABRA1/GABRA5*) which are expressed in several adult DRG neuronal

subtypes[4,69] (Table S2)*.* Overall, these transcriptional changes may underly the observed

physiologic maturation of co-cultured iSN.

Conversely, 748 significantly (p<0.05) downregulated genes were identified in cocultured

iSNs compared to iSNs alone (Fig. 2A, Table S1). There was decreased expression of genes

implicated in early-stage neural crest development (*Lin28a*, p adj.=$1.249^{-21}$; *RTKN,* p adj.=$9.661^{-}$

$^{15})$[80,81] as well as a gene integral to differentiation of neural crest progenitors into mesenchymal

lineage cells (*ADAM12*, p adj.=$1.069^{-42})$[82] (Fig. 2A). We also observed a highly significant

decrease in *RGMB* expression (p adj.=$9.985^{-28}$), a BMP signaling partner which has been

shown to promote ventral neuronal differentiation from neural crest cells[83] (Fig. 2A).

Additionally, we observed a significant decrease in expression of genes associated with

peripheral gliogenesis, including myelinating Schwann cell markers (*Pou3F1*[84], p adj.=$4.872^{-12}$,

*SVIP*[85] p adj.=$5.928^{-15}$, *GPR17*[86] p adj.= $7.605^{-1}$, *MAL*[87] p adj.=$1.120^{-8}$, *PMP22*[88,89] p adj.=$1.325^{-5}$

, satellite glial markers (*TRDN*[89] p adj.=$4.190^{-20}$, *BCAN*[88,89]  p adj.=$7.605^{-11}$, *CHL1*[90] p

adj.=$1.863^{-7}$), and glial proteins integral to axonal development (*S100A6*[91] p adj.=$1.481^{-14}$ ,

*SEMA3B*[92] p adj.=$8.410^{-10}$,  *SEMA3G*[92] p adj.=$9.360^{-8}$)  (Fig. 2A) suggesting enhanced

commitment to a neuronal fate when iSNs are differentiated in co-culture.

Gene Ontology analysis yielded 134 significantly downregulated gene programs in iSNs

differentiated in coculture compared to iSNs differentiated alone (Fig. 2C, Table S2). Gene

programs associated with MHC I/II complex assembly and antigen presentation were among the

most significantly downregulated biological processes in cocultured iSNs (Table S2), consistent

with recent data that differentiated iNCPs express these proteins at low levels, if at all[93,94]. We

also observed significant downregulation in gene programs associated with gliogenesis, glial

differentiation, myelination, and axon ensheathment (Fig. 2C), consistent with results suggesting

iSNs differentiated in coculture are committed to a neuronal, rather than a glial, fate.

Interestingly, GO terms also included transcription factors implicated in the early specification of

DRG neuronal subtypes including *NTRK3* and *SHOX2* (expressed in proprioceptors), *RET*

(expressed in nonpeptidergic nociceptors and mechanoreceptors) *POU4f3* (expressed in a

subpopulation of peptidergic nociceptors) and *GFRA3* (expressed in a subpopulation of

peptidergic nociceptors)[1,4,95,96]  (Table S2). These results indicate that iSNs in co-culture are

committed to a sensory neuron fate and restrict their transcription factor repertoire, which may

advance development of functional DRG subtypes.

GO analysis also provides potential explanation for a change in neuronal physiology with

decreased expression of genes involved in the development, composition, and function of

synaptic membrane structures (Fig. 2C), including glutamate receptors, such as kainate

receptors (*GRIK1/GRIK3*), AMPAR *GRIA4,* and Delta receptor *GRID2*[97]*,* as well as purinergic

receptors (*P2RY1, P2RX6*) (Table S2). Interestingly, potassium ion channel expression (*KNCT2,*

*KNCJ3, KCNJ12, KCNS1*) was also decreased, which could contribute to the increased

excitability we observe in cocultured iSNs compared to iSNs differentiated alone[79] (Table S2).

Taken together transcriptomic analyses indicate that coculture of iNCPs with peripheral glia

enhances differentiation into more mature sensory neurons.

| Role in Axonogenesis | Gene Names |
|---|---|
| Specification | *CDKL5*[98]/*PICALM* [99] |
| Outgrowth | *CDKL5*[98,100]/*AUTS2*[101] /*LAMA2*[102] /*PRICKLE1*[103]/ *EIF4G2* [104]/ *TRPC5*[105] /*ADARB1*[106]/*ENAH*[107]/ *RAB10*[108]; *USP9X*[109]/ *SOCS3*[110] |
| Branching | *NTN1*[111] / *BCL11A*[112] |
| Guidance | *EXT1*[111] /*PAK3*[113] /*EFNB2*[114]/ *NTN1*[111,115]/ *UNC5C*[115], *GLI3*[116], *FOXD1*[117] |
| Synaptogenesis | *NTN1*[111]/*SLITRK1/3*[118] /*ADARB1*[119] / *BCL11A*[120] / *DLG5*[121] |

**Table 2. Genes upregulated in cocultured iSNs that are implicated in aspects of axonogenesis.**

**Rodent DRG satellite glia (rSGs) are responsible for advanced maturation of iSNs:**

Our studies demonstrate that iSNs differentiated in coculture with embryonic rat DRG cells

mature at an expedited rate compared to iSNs differentiated alone, however due to the protocol

used to prepare the rDRG cells, the glial subtype(s) driving this maturation was not initially clear.

The rodent DRG cell population includes satellite glia (SGs), myelinating and non-myelinating

Schwann cells (SCs), glial progenitor cells (GPCs), fibroblasts, endothelial cells, and immune

cells. When initially plated (Fig. 1A, Day 0-7), rDRG cells display a multitude of morphologies,

with small islands of semilunar cells and large flat cells, and this variety in cellular profiles is

maintained as long as the cells remain in Glia Base Media (Fig. S3A-B). However, the

morphology and cell number of rDRG glia changes upon addition of Deng et al. 2023[15] NCATS

Maturation Media (NMM), which is required for terminal differentiation of iSNs (Fig S3A-B). We

used RNA sequencing to determine the impact of both NMM treatment and iSN contact on

rodent glial identity and function (Fig. S3C), mRNA reads from co-cultured RNAs were aligned

to both the human and rat genome to differentiate mRNAs derived from hiSNs and rDRG glia

respectively. PCA analysis and hierarchal clustering (Fig. S3D-E) showed that rDRG glia treated

with NMM cluster together with cocultured rDRG glia treated with NMM. All glia treated with

NMM cluster separately from rDRG Glia Alone maintained in base media, suggesting that media

composition drives the transcriptional changes observed, while iSN contact leads to minimal

changes in the glial transcriptome.

DGE analysis yielded 1160 significantly (p<0.05) upregulated genes in NMM treated rDRG

glia compared to rDRG glia in Glia media  (Fig 3A, Table S3). Overall, top upregulated genes in

NMM treated glia are associated with mature satellite glia (SGs). The most upregulated genes,

*DDIT4* (p adj.=$1.196^{-39}$) and *INSIG1* (p adj.=$1.196^{-39}$), have previously been identified as DRG

SG specific markers in mouse sc-RNAseq studies[88,90]. *DDIT4* has been implicated in

suppression of myelination[122] and *INSIG1*[90] plays a role in cholesterol biosynthesis. Lipid

synthesis and metabolism are established functions of DRG SGs[123,124] and all top upregulated

genes with NMM treatment are involved in these processes including *PRR5*[125] (p adj.=$2.185^{-23}$),

*PCSK9*[126] (p adj.=$2.067^{-21}$), *HMGCS1*[89] (p adj.=$5.928^{-21}$), *TMEM97*[127] (p adj.=$9.604^{-19}$). and

*SCD*[89,90] (p adj.=$2.762^{-18}$). *HMGCS1* and *SCD*[89], and their related upregulated genes, *HMGCR*

(p adj.=$1.537^{-12}$) and *SCD2*[90] (p adj.=$4.821^{-15}$), have also previously been identified as DRG SG

gene makers in sc-RNA seq studies. GO analysis (Fig. 3B, Table S4) confirmed that lipid

biosynthesis and metabolism pathways were significantly upregulated in NMM cultures.

We also observed significant upregulation in biologic processes, cellular components, and

molecular functions related to extracellular matrix (ECM) production. These include increased

expression of SG genes related to collagen production, such as *COL16A1*, *COL1A1, COL3A1,*

and *COL5A1*[90], and other ECM related genes previously associated with SG identity, *FBLN5*[89,90]

and *ANXA5*[90] (Table S4). Significant upregulation in biological processes related to DNA

replication (Table S4) indicate that enrichment for SGs may be due to enhanced cellular

proliferation. Together these results support that treatment with NMM results in an increased

proportion of mature SGs in our cultures.

DGE analysis also yielded 933 significantly ($p<0.05$) downregulated genes in NMM treated

rDRG glia compared cultures maintained in base Glia Media (Fig. 3A, Table S3). Overall, these

results indicate a decrease in immune functions in NMM treated cultures. Top downregulated

genes were key markers of DRG macrophages including *STAB1*[128,129] (p adj.=$2.984^{-30}$), *LYZ2*[89]

(p adj.=$4.974^{-22}$), *CSF1R*[130] (p adj.=$9.916^{-20}$), *MSR1*[131] (p adj.=$1.090^{-19}$), and *AIF1*[89] (p

adj.=$4.859^{-16}$). GO analysis (Fig. 3C, Table S4) confirmed that the most downregulated

biological processes in NMM treated rDRG glia are related to various aspects of leukocyte

function including leukocyte differentiation, proliferation, homeostasis, activation, migration,

cytokine production, and chemotaxis. Significant decrease in expression of gene programs

related to both mononuclear cells (phagocytosis, macrophage activation, macrophage

differentiation) and lymphocytes (T-cell proliferation, T-cell activation, B-cell differentiation, B-

cell activation, adaptive immune response) are observed (Table S4). GO analysis also reveals

decreased expression of genes related to several aspects of actin cytoskeletal organization,

which may contribute to morphologic changes observed with NMM treatment. Together these

data indicate the composition of NMM results in an enrichment of SGs. When we compared

transcriptomic differences between co-cultured rDRG glia treated with NMM and rDRG glia

maintained in Glia Media, we found that differentially expressed genes were nearly identical to

those identified in NMM treated cultures without iSN contact (Fig. 3D-F, Table S5-6), indicating

that NMM treatment drives an enrichment of SGs in rDRG glial cultures.

To elucidate the impact of iSN contact on the transcriptome of SGs, we next compared gene

expression between co-cultured NMM treated glia and NMM treated glia alone. DGE analysis

yielded very few significant (p<0.05) differences, with only 20 significantly upregulated and 52

significantly downregulated genes in co-cultured NMM treated glia (Fig. 3G, Table S7). The

results indicate that iSN contact impacts the SG cell-cycle and morphology with upregulation of

genes related to DNA replication and repair (*CDC14A*[132], p adj.=0.0007), transcription

(*ZSCAN18,* p adj.= 0.0352; *TLE4* p adj.= 0.0393), translation (*QARS*[133], p adj.=$8.165^{-10}$),

cytoskeletal organization (*DNMBP*[134] p adj.= 0.047, *SRCIN1*[135] p adj.= 0.047)*, SLITRK2*[136] p

adj.=0.0007) and ion channels (*TRPM3* p adj.= 0.047*, GRIK2* p adj.= 0.008*, KCNK5* p adj.=

0.007). Several upregulated genes were previously linked to specialized SGs (*SLITRK2*[86]*,*

*TRPM3*[137]*, ITGB4*[90]*, GBP2*[90,137]), which may indicate changes in SG specific functional

phenotype when in contact with iSNs.  Conversely, downregulated genes (Fig. 3G, Fig. S3F,

Table S7-8) were associated with collagen production (*P4HA3*[138] p adj.= 0.013, *COL4A5*[139] p

adj.= 0.015) and cell-adhesion (*LTBP2*[140] p adj.= 0.008), which may indicate an alteration in cell

spreading and migration with neuronal contact.

Taken together, transcriptomic analyses illustrate that initial DRG cell preparations contain

multiple cell types, however maturation media enriches for satellite glia. SGs in coculture and

SGs cultured alone with NMM treatment are transcriptionally very similar, however minor

changes in morphological and functional aspects of SGs do occur with neuronal contact.

**Rodent SG (rSG) secreted factors alone do not induce a pseudounipolar morphology in**

**hiSNs:**

The transcriptomic analysis confirmed that rodent satellite glia (rSGs) contribute to the

maturation of iSNs, however the molecular mechanisms underlying this process are unknown.

We first asked whether this process is mediated by glial secreted factors, using a transwell co-

culture system (Fig. 4A). rSGs were cultured for one week on polycarbonate membranes

inserted into a well plate, and iSNs were then terminally differentiated in the same wells beneath

the membranes, sharing media but never physically touching rSGs, or in wells alone. iSNs were

then fixed and stained to visualize neuronal morphology (TUJ1, cytoskeletal marker, green;

BRN3a, iSN maturation marker, red) (Fig. 4B), while the rSGs on membranes were dyed and

visualized live after two weeks of culture to confirm that a healthy population was present.  iSNs

differentiated in transwell coculture and alone both presented with predominantly multipolar or

immature morphologies, with no significant transition to pseudounipolar morphology (Fig. 4C,

Table 3).

It is possible that glia only secrete factors critical for the maturation process after they have

made physical contact with iSNs. To test this possibility, iSNs plated alone were treated for six

days with spent media collected from cocultures (Fig. 4D-F, Table 4). While there are no

differences in pseudounipolar morphology, there is a higher frequency of immature neurons in

untreated iSNs (immature *adj. residual* = 3.544) compared to spent media treated iSNs

(immature *adj. residual* = -3.544), as well as an increased proportion of multipolar cells in spent

media treated iSNs (multipolar *adj. residual* = 2.954) compared to untreated iSNs (multipolar

*adj. residual* = -2.954). These results are consistent with the idea that rSG-iSN contact, rather

than secreted factors, are needed to induce pseudounipolar morphology.

| | **Morphology Frequencies** | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | **Immature** | | **Multipolar** | **Pseudounipolar Transition (PUT) + Pseudounipolar (PU)** | | | | | |
| **Condition** | **Unipolar** | **Bipolar** | **Multipolar** | **PUT-Bulge** | **PUT-Bell** | **PUT-AB** | **PU-Short** | **PU-Long** | **Totals** |
| Coculture | 0 | 34 | 167 | 18 | 3 | 0 | 1 | 0 | 223 |
| iSN Alone | 1 | 42 | 176 | 8 | 6 | 1 | 1 | 1 | 236 |
| **Totals** | 77 | | 343 | 39 | | | | | 459 |

**Table 3. Raw frequency of morphology subtypes observed in transwell co-cultured iSNs vs. iSNs terminally differentiated alone. $X^2$ frequency statistic calculated for major morphology groups: Immature, Multipolar, and PU(T). $X^2$= 1.562, df=2, p= 0.4579.**

| | Morphology Frequencies | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | **Immature** | | **Multipolar** | **Pseudounipolar Transition (PUT) + Pseudounipolar (PU)** | | | | | |
| **Condition** | **Unipolar** | **Bipolar** | **Multipolar** | **PUT-Bulge** | **PUT-Bell** | **PUT-AB** | **PU-Short** | **PU-Long** | **Totals** |
| Media Treated | 0 | 10 | 225 | 10 | 2 | 0 | 0 | 0 | 247 |
| Untreated | 0 | 26 | 159 | 1 | 3 | 2 | 4 | 0 | 195 |
| **Totals** | 36 | | 384 | 22 | | | | | 442 |

**Table 4. Raw frequency of morphology subtypes observed with and without coculture spent media treatment. $X^2$ frequency statistic calculated for major morphology groups: Immature, Multipolar, and PU(T). X2= 12.69, df=2, p= 0.0018.**

**Physical contact between rSGs and hiSNs promotes pseudounipolarization:**

To determine whether physical contact with satellite glia promotes pseudounipolarization, we designed an assay in which rDRG cells were cultured for two weeks in NMM, then fixed with methanol prior to coculture, preserving surface protein configurations. iSNs were subsequently cultured with the methanol fixed glial cells, in a normal coculture, or alone for one week (Fig. 5A). Cultures were then fixed and stained to visualize neuronal morphology (TUJ1, cytoskeletal marker, green) and contact with rSGs (p75, NGF receptor, magenta) (Fig. 5B). iSNs differentiated in methanol fixed coculture exhibit a phenotype intermediate between iSNs differentiated in normal coculture and those differentiated alone (Fig. 5C, Table 5). The Chi-square frequency test identified significant differences between observed and expected morphology frequencies across all three groups ($X^2$=131.3, df=6, p<0.001). Interestingly, these results are driven by significant differences between iSNs differentiated in normal coculture conditions (Immature *adj. residual* = -3.543, Multipolar *adj. residual*=-6.471, PUT *adj. residual*= 6.468, PU *adj. residual*=7.893) and iSNs differentiated alone (Immature *adj. residual* = 2.041, Multipolar *adj. residual* = 4.963, PUT *adj. residual*= -5.753, PU *adj. residual*= -3.893), while iSNs

differentiated in a methanol fixed coculture exhibited "expected" morphology frequencies, with

only significantly fewer pseudounipolar cells than expected (PU *adj. residual*= -4.007). Since

PU(T) cells make-up 14.02% of iSNs differentiated in methanol fixed coculture, compared to the

2.26% of iSNs differentiated alone, these results suggest that physical contact between rSGs

and iSNs contributes to the changes required for a transition to pseudounipolar morphology.

The degree to which such direct contact truly contributes to this process is unclear, as

methanol fixation prevents movement of rSGs in culture, and therefore physical contacts

between rSGs and iSNs may be reduced compared to normal cocultures. Overall, the average

number of rSG contacts per iSN was significantly reduced in the methanol fixed condition

compared to normal cocultures ($t$=23.04, df=470, p<0.0001), with an average of 2 contacts/iSN

in methanol fixed coculture vs. 10 contacts/iSN in normal cocultures (Fig. S4A). Significantly

more contacts were made to true pseudounipolar cells in methanol fixed coculture compared to

less advanced morphologies (Bonferroni post-hoc: PU vs. Multipolar p=0.0038; Bipolar

p=0.0053, PUT Bulge p=0.0097) (Fig. S4B). In fact, pseudounipolar iSNs in methanol fixed co-

culture were contacted to a similar degree to those in normal coculture, with an average of 9

rSG contacts/ iSN in methanol fixed coculture and 10 rSG contacts/iSN in normal coculture.

These results indicate that contact is essential to pseudounipolarization, but downstream

signaling and movements of live rSGs may also contribute to this process.

| | Morphology Frequencies | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | Immature | | Multipolar | Pseudounipolar Transition (PUT) | | | Pseudounipolar (PU) | | |
| Condition | Unipolar | Bipolar | Multipolar | PUT-Bulge | PUT-Bell | PUT-AB | PU-Short | PU-Long | Totals |
| Methanol Fixed Coculture | 2 | 37 | 189 | 35 | 2 | 1 | 4 | 0 | 270 |
| Coculture | 5 | 7 | 105 | 36 | 16 | 10 | 30 | 5 | 214 |
| iSN Alone | 1 | 28 | 144 | 4 | 0 | 0 | 0 | 0 | 177 |

| Totals | 80 | 438 | 104 | 39 | 661 |
|--------|----|----|----|----|----|

**Table 5: Raw frequencies of morphologies and morphology subtypes observed in methanol fixed coculture, normal coculture, and alone. $X^2$ frequency statistic calculated for major morphology groups: Immature, Multipolar, PUT and PU. X2=131.3, df=6, p<0.001.**

**Semaphorin-plexin and gap junction signaling play a role in pseudounipolarization of hiSNs:**

We identified two potential pathways whose constituents are appropriately expressed to enable contact mediated SG signaling to iSNs, namely, Sema6a-Plexin A4 and gap junctions. Sema6a, a transmembrane semaphorin protein that is expressed in satellite glia, repels or inhibits axonal growth in developing sensory neurons through engagement with the PlexinA4 receptor[141–144], which is highly expressed in hiSNs (Fig. S4C). Additionally, our sc-RNA seq data set from embryonic DRG satellite glia at the time of pseudounipolarization reveals high expression of *GJA1*, encoding the gap junction protein Connexin-43[88]. *GJA1* was also found to be highly expressed in all NMM treated rDRG glial cultures (Fig. S4D).

To test if these pathways promote a pseudounipolar morphology (Fig. 5D), we first knocked down expression of *SEMA6A* in the rSGs using LV-shRNA transduction and puromycin selection. Selected glia were allowed to repopulate and then hiSNs were plated for terminal differentiation in coculture with shRNA-Sema6a expressing glia (shSema6a-rSGs) or shRNA-GFP expressing glia (shGFP-rSGs, control). Cells were cocultured for one week and then fixed and stained to visualize neuronal morphology (TUJ1, cytoskeletal marker, red) and glial expression of Sema6a (Sema6a, magenta; GFP, green) (Fig. 5E). We found that the iSNs differentiated in coculture with shSema6a-rSGs transition to pseudounipolar morphology at a reduced rate compared to controls (Fig. 5F, Table 6), with significantly fewer PU(T) cells (PUT *adj. residual*=-4.772, PU *adj. residual*=-6.861) compared to controls (PUT *adj. residual*=4.772, PU *adj. residual*=6.861), and significantly more multipolar cells (*adj. residual*=9.39) compared to controls (Multipolar *adj. residual*=-9.39. These results indicate that Sema6a-PlexinA4 signaling contributes to pseudounipolarization in this coculture model and may also decrease the

411 incidence of multipolar morphology by inhibiting axonal outgrowth. However, the results also

412 suggest that plexin signaling is not the sole mechanism underlying iSN maturation, since PU(T)

413 cells make up only 10.28% of iSNs differentiated in shSema6a-rSG coculture.

| | Morphology Frequencies | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | Immature | | Multipolar | Pseudounipolar Transition (PUT) | | | Pseudounipolar (PU) | | |
| Condition | Unipolar | Bipolar | Multipolar | PUT-Bulge | PUT-Bell | PUT-AB | PU-Short | PU-Long | Totals |
| shRNA-Sema6a | 2 | 9 | 242 | 15 | 7 | 2 | 4 | 1 | 282 |
| ShRNA-GFP | 20 | 8 | 181 | 38 | 29 | 14 | 53 | 16 | 359 |
| Totals | 39 | | 423 | 105 | | | 74 | | 641 |

**Table 6. Raw frequency of morphologies observed in shRNA-Sema6a-SG coculture and shRNA-GFP-SG**
**Coculture. $X^2$ frequency statistic calculated for major morphology groups: Immature, Multipolar, PUT and PU.**
**$X^2$=94.62, df=3, p<0.001.**

To assess the role of satellite glial gap junctions in iSN maturation, we knocked down *GJA1*

in the rSGs and plated iSNs for terminal differentiation in coculture with shRNA-GJA1

expressing glia (shGJA1-rSGs) or shRNA-GFP expressing glia (shGFP-rSGs, control) for one

week (Fig 5D). Cocultures of hiSNs with shRNA-GJA1 expressing glia (shGJA1-rSGs) or

shRNA-GFP expressing glia (shGFP-rSGs, control) were stained to visualize neuronal

morphology (TUJ1, cytoskeletal marker, red) and glial expression of GJA1 (GJA1, magenta;

GFP, green) (Fig 5G). This experiment was carried out in parallel with the semaphorin

knockdown, and therefore we utilized a single group of sh-GFP RNA control samples. Strikingly,

iSNs differentiated in coculture with shGJA1-rSGs also transition to pseudounipolar morphology

at a reduced rate compared to controls (Fig. 5F, Table 7) with fewer PU(T) cells than expected

(PUT *adj. residual*=-5.537, PU *adj. residual*=-7.014), and significantly more multipolar cells than

expected (*adj. residual*=10.109). These results suggest that both gap junction and

Sema6a/PlexinA mediated signaling contribute to pseudounipolarization.

| | Morphology Frequencies | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | Immature | | Multipolar | Pseudounipolar Transition (PUT) | | | Pseudounipolar (PU) | | |
| Condition | Unipolar | Bipolar | Multipolar | PUT-Bulge | PUT-Bell | PUT-AB | PU-Short | PU-Long | Totals |
| shRNA-GJA1 | 3 | 9 | 277 | 12 | 7 | 4 | 6 | 1 | 319 |
| ShRNA-GFP | 20 | 8 | 181 | 38 | 29 | 14 | 53 | 16 | 359 |
| Totals | 40 | | 458 | 104 | | | 76 | | 678 |

**Table 7. Raw frequency of morphologies observed in shGJA1-rSG coculture and shGFP-rSG Coculture. $X^2$ frequency statistic calculated for major morphology groups: Immature, Multipolar, PUT and PU. $X^2$=107.5, df=3, p<0.001.**

**Cocultured iSNs provide a more physiologically relevant model of neuropathy:**

iPSC derived neurons often exhibit embryonic phenotypes, which reduces the ability to model complex molecular mechanisms and disease processes [9–15]. To determine whether cocultured iSNs provide a physiologically relevant model of peripheral neuropathy compared to iSNs differentiated alone, we examined the response of these cells to a common form of Chemotherapy Induced Peripheral Neuropathy (CIPN), paclitaxel induced peripheral neuropathy[145,146]. Patients with CIPN experience symptoms of allodynia, paresthesia, numbness, and pain in a stocking glove distribution, due to the degeneration of sensory nerve axon terminals[147]. We assessed the degeneration response of iSNs differentiated in coculture and alone when treated with low dose paclitaxel. At one week of terminal differentiation cultures were treated with paclitaxel chemotherapy or DMSO for 48 hours, then fixed and stained to visualize the neuronal cytoskeleton (TUJ1, Green) and gap junctions in both neurons and glia (GJA1, Magenta) (Fig. 6A,C). Neurodegeneration was estimated by quantifying TUJ1 signal.

We first tested the impact of low-dose paclitaxel on the degeneration of iSNs differentiated with the Deng et al. 2023[15] protocol, generated by NCATS. Analysis of TUJ1 expression showed that cocultured iSNs were significantly more susceptible to paclitaxel induced degeneration than iSNs differentiated alone (Bonferroni post-hoc: Coculture + Paclitaxel treatment vs. Coculture +

DMSO: p <0.0001; vs. iSN Alone + Paclitaxel: <0.0001; vs. iSN Alone + DMSO: p<0.0001),

which were not significantly impacted by treatment (Bonferroni post-hoc: iSN Alone + Paclitaxel

vs. iSN Alone + DMSO p >0.9999) (Fig. 6B). We next tested the impact of the chemotherapy

treatment on degeneration of RealDRG[TM] iSNs from Anatomic, which require a larger dose of

paclitaxel to elicit axonal degeneration[148]. Again, iSNs differentiated in coculture were more

susceptible to paclitaxel induced degeneration compared to iSNs differentiated alone

(Bonferroni post hoc: Coculture + Paclitaxel treatment vs. Coculture + DMSO: p <0.0001; vs.

iSN Alone + Paclitaxel: <0.0001; vs. iSN Alone + DMSO: p<0.0001), which were not significantly

impacted by treatment (iSN Alone + Paclitaxel vs. iSN Alone + DMSO p >0.9999) (Fig. 6D).

Together these data indicate that iSNs cocultured with satellite glia provide a more physiologic

model for toxin induced neuropathies.

**Discussion:**

The use of differentiated iPSCs to model human diseases has revolutionized our ability to

understand pathology and develop efficacious treatments. While researchers have begun to

utilize induced sensory neurons to understand peripheral neuropathies, current differentiation

methods produce iSNs with an embryonic phenotype that do not fully recapitulate postnatal

functions. In this study, we developed a protocol that enables maturation of iSNs in coculture

with rodent satellite glia. We found that contact between iSNs and rSGs, mediated by

semaphorin-plexin and glial gap junction signaling, promotes an adult morphology and

physiology in the sensory neurons. Ultimately, this system also enables an enhanced model of

neuropathy *in vitro*, by increasing iSN susceptibility to chemotherapy induced axonal

degeneration at physiologic doses.

These studies establish a role for satellite glia in morphologic development of DRG neurons

and demonstrate that we can leverage this relationship to reliably produce pseudounipolar iSNs.

Previous studies have implicated DRG supporting cells in the morphologic maturation of

sensory neurons, with mixed primary cultures of DRG neurons and glia facilitating the transition

of sensory neurons to a pseudounipolar morphology[34–36,40]. This result has been recapitulated in

induced neuro-mesodermal assembloids[149], however iSNs cocultured with induced[16] or primary

[21] Schwann cells alone do not exhibit advanced morphology.  Based on these studies, we tested

the possibility that satellite glia primarily contribute to sensory neuron maturation.  *In vivo,* DRG

sensory neurons increase physical contact with satellite glia during pseudounipolarization

through the development of perikaryal outgrowths that increase contact surface area[40] (Fig.

S1B),  In the current study, we show that physical contact between glial cells and iSNs is

essential to the development of a pseudounipolar morphology, and RNA sequencing and

immunohistochemistry indicate that specifically satellite glia are responsible for this effect.

Though glial contribution to sensory neuron maturation have been recognized previously,

the molecular mechanisms enabling development of pseudounipolar morphology have been

unclear. Since our results ruled out glial secreted factors as a direct mechanism of

pseudounipolarization, we explored potential contact mediated mechanisms driven by our RNA

sequencing data. In iSNs, we observed high expression levels of *Plexina4*, a receptor that

interacts with glial semaphorins Sema3a and Sema6a[141–144,150], which have both been

implicated in sensory neuron axon development[141–144,151–153]. Since class 3 semaphorins are

secreted, we assessed the role of Sema6a, a transmembrane semaphorin[154], in the

development of iSN axons. We found knockdown of glial *Sema6a* significantly reduced

pseudounipolar formation in cocultured iSNs and instead resulted in a high proportion of

multipolar cells. While there was a significant reduction in the proportion of pseudounipolar cells

following *Sema6a* knockdown in the satellite glia, this effect was not complete. Factors

contributing to observed low levels of pseudounipolarization may include incomplete knockdown

of *Sema6a*, compensation via secreted Sema3a, and the possibility that semaphorin-plexin

signaling is not the only mechanism underlying this process.

In the DRG, satellite glia communicate with adjacent satellite glia and sensory neurons

through gap junctions[155,156], contributing to the maintenance of neuronal homeostasis through

direct exchange of diverse signaling molecules[157,158]. While little is known regarding gap junction

signaling in sensory neuron axon development, communication through gap junctions correlates

with neural crest cell differentiation into neuronal cells[159]. Knockdown of glial *GJA1,* like

knockdown of Sema6a, significantly reduced the ability of glial cells to induce a transition of

iSNs to a pseudounipolar morphology. As was seen following knockdown of Sema6a,

knockdown of *GJA1* did not completely abolish pseudounipolarization. Again, this may be

explained by incomplete knockdown efficiency, compensation by an alternate connexin, or

another parallel signaling mechanism. Future studies need to be carried out to determine

whether glial-glial or glial neuronal gap junction signaling contributes to development of

pseudounipolar morphology.

RNA sequencing also revealed potential intracellular mediators of pseudounipolarization.

Multiple molecules implicated in axonal guidance were significantly upregulated in cocultured

iSNs, as illustrated in Table 2. These include *CDKL5,* a kinase linked to axonal polarization[98]

and *BCL11a,* a transcription factor that suppresses growth of supernumerary neurites and

regulates microtubule protein expression[112]. We also observed significant downregulation of

*RGMB* which promotes formation of multiple neurites[160]. The identification of cues and

molecular pathways that stimulate pseudounipolar morphology may enable further refinements

of iSN protocols.

Pseudounipolar morphology has multiple implications for sensory neuron physiology, as the

axon stem may act as an axon initial segment that potentiates spontaneous activity[36] or as a

filter, gating nociceptive action potentials[161,162]. Importantly, we demonstrate that terminal

differentiation in coculture yields more physiologically mature iSNs. Using MEA, we found that

iSNs in coculture become electrically active earlier than iSNs cultured alone, and fire at a rate

more consistent with rodent DRG neurons[54,55]. In addition to the development of an axon stem,

the observed differential expression of several ion channels and receptors may contribute to

mature firing rates. Upregulation of *SCN10a*, encoding Nav1.8, is of particular importance, as

this sodium channel maintains repetitive action potential firing in DRG nociceptors[163] and is

integral to pain signaling[72,73,164]. Increased expression of muscarinic receptors *CHRM2* and

*CHRM3* may also contribute to spontaneous activity, as acetylcholine has been implicated in the

development of bursting and spiking behavior in sympathetic neurons cocultured with satellite

glia[165]. Lastly, increased neuronal excitability may reflect a reduction in expression of several

potassium channels, namely *KNCT2*[166] and *KCNS1*[167], which have previously been linked to

hyperexcitability and neuropathic pain.

In this study our goal was to generate iSNs with a more mature adult-like phenotype that

would allow us to better model peripheral neuropathies, and we tested the efficacy of modeling

toxic neuropathy using chemotherapy treated cocultured iSNs. We found that cocultured iSNs

were significantly more susceptible to paclitaxel chemotherapy mediated axonal degeneration

compared to iSNs alone. Hyperexcitability of sensory neurons has been shown to contribute to

axonal degeneration across multiple neuropathies[6], including chemotherapy induced peripheral

neuropathy (CIPN)[145,146], and so the increased baseline excitability of cocultured iSNs may

enhance susceptibility to degeneration. Satellite glia may also directly contribute to neuronal

degeneration, as studies in rodent models of CIPN indicate that glial release of inflammatory

cytokines contribute to degeneration[168–170]. Overall, our results provide new avenues for

studying the potential involvement of satellite glial-sensory neuron interactions in the

pathogenesis of neuropathies, as this protocol can easily be used to model diabetic peripheral

neuropathy, inherited neuropathies, or axonal injury as well.

In summary, our work identifies molecular direct contact cues whereby satellite glia support

sensory neuron development and establishes an accessible protocol for the reliable

differentiation of mature iSNs which can be used to study physiological and pathological

changes in sensory neurons and develop novel treatments for peripheral neuropathies.

*Limitations of this Study:*

This study is limited by the use of rodent glial cells and human derived sensory neurons.

Currently, there are no publicly available protocols to differentiate human satellite glia, however

several protocols have been developed to induce Schwann Cells[38,171] which share a glial

progenitor cell lineage. Adjustments of these protocols such that satellite glia differentiation can

be optimized will allow for an advanced iSN differentiation protocol with exclusive use of human

cells, and aid in identification of molecular mechanisms underlying normal human sensory

neuron development and neuropathy.

**Methods:**

**Animal Care:** Sprague-Dawley rats were purchased from Charles River Laboratory. Animals

were cared for in accordance with NIH standards and guidelines. Experimentation was

approved by the Dana Farber Cancer Institute IACUC.

**rDRG-iSN Coculture:**

*Rodent DRG glial cell culture:* Sprauge-Dawley day E14.5 rat DRGs were dissected. DRGs

were trypsinized (1mg trypsin /mL HBSS) for 50 minutes at 37°C. 40k cells/well were plated on

uncoated Lab-Tek™ II Chambered Coverglass (Thermofisher cat#: 155409) with filtered Glia

Base Media (DMEM/F12, 10% FBS, 1% Pen. Strep., 0.08% glucose).  After a 30-minute

incubation at 37°C 7.5% $CO_{2.}$, chambers were vigorously shaken, and media was fully changed

to remove cells in suspension. Remaining glial cells were cultured in Glia Base Media at 37°C

7.5% $CO_2$ for one week with media changes 3x/ week.

*Induced Sensory Neuron (iSN) terminal differentiation and iSN- rDRG glia co-culture:* Frozen

Neural Precursor Cells (NPCs) were obtained from collaborators at NIH-NCATS. NPCs were

differentiated from NCRM5 (Research Grade, male, CD34+ cord blood)  and GR1.1 (Clinical

Grade, male CD34+ cord blood) iPSC lines at NIH-NCATS using the protocol outlined in Deng

et al., 2023[15]. Lab-Tek™ II Chambered Coverglass (Thermofisher cat#: 155409) were coated

with 0.005% poly-ornithine solution (Sigma-Alrich cat#: P4957) in dH2O 1 hr at 37°C, washed,

and then coated with 5 ug/mL mouse laminin (Thermofisher cat#: 23017015) in dH2O 1 hr at

37°C. NPCs were thawed in NCATS Maturation Media (DMEM/F12+Glutamax, 2% B27,1% N2,

25ng rh- BDNF/b-NGF/GDNF/NT3)    + CEPT rock inhibitor cocktail (50nM Chroman-1

(Medchem Express Cat#: HY-15392), 5uM Emricasan (Selleck cat#: S7775), 0.001% Polyamine

supplement (Sigma-Alrich cat#: P8483), 0.7uM Trans-ISRIB (Tocris cat#: 5284)). 15k cells/well

were plated in NCATS Maturation Media + CEPT on coated cultureware for terminal

differentiation alone (iSN-Only), or in coculture with rDRG glial cultures treated with NCATS

Maturation Media + CEPT. 24 hrs after plating, media was changed to NCATS Maturation Media

+ 1uM PD0332991 (Tocris cat#: 4786). Cells were cultured for one week at 37°C 5% $CO_{2.}$, with

½ media changes 3x/week. n=4 biological replicates, 2/ genetic iPSC line.

**Quantification of iSN morphology frequency:**

*Immunofluorescence:* Cells were fixed for downstream Immunofluorescence (IF) on Day 14.

First, cells were fixed with 2% paraformaldehyde (PFA) (EMS cat#: 15710; 1:1 4% PFA to spent

media) 10 minutes at room temperature (RT), and then in 4% PFA 20 minutes at RT. Cells were

washed 3x in PBS and stored at 4°C. Fixed cultures were permeabilized with 0.1% Triton-x in

PBS 10 minutes and blocked in blocking buffer (5% Normal Goat Serum (Cell Signaling Tech.

cat#: 5425S) in 1x PBS) for 1 hr at RT. Cultures were then incubated overnight at 4°C in primary

antibodies (Table 11) diluted in blocking buffer. The following day, cultures were washed and

incubated in Alexa-Fluor secondary antibodies (Table 12) and DAPI diluted 1:1000 in blocking

buffer for 1 hr at RT. Co-Cultures were stained with mouse anti-human nuclei-CY3 conjugate

(Sigma cat#: MAB1281C) for 2 hrs at RT to distinguish human neuronal nuclei from rodent.

Cells cultured in Lab-Tek™ II Chambered Coverglass were stored at 4°C in PBS prior to

imaging, and cells cultured on coverslips were mounted on slides with fluoromount

(Thermofisher cat#: OB10001).

*Imaging:* All photomicrographs of cells cultured in Lab-Tek™ II Chambered Coverglass were

collected on the Zeiss-980 airy-scan/confocal microscope. Images of all conditions were

obtained at 20x with 1.3x zoom and bidirectional scanning in SRY-4 Airy-Scan. Multiplex

imaging was utilized with 405, 488, 561, and 639 lasers at 800V gain, 1% laser power, and 0

offset. All images were obtained in a z-stack with 0.35um step size. Systematic random

sampling was utilized to image n=50-100 cells/biological replicate across 8-wells. Areas with

fused cell bodies and axonal fasciculations were omitted from sampling. All images were 3D

airy-scan post processed in Zen Blue software.

*Analysis:* All human cells in each photomicrograph were categorized into the following

morphologic groups : unipolar (one axonal projection off of the cell body, that does not

bifurcate), bipolar (two axonal projections off of opposite cell body poles, ~180° angle),

multipolar (more than two axonal projections), Pseudounipolar Transition (PUT)- Bulge (two

axonal projections off of the bottom of the cell body, axons form >90° angle), PUT-Bell (two

axonal projections off of the bottom of the cell body, axons form a 90° angle or less), PU-

Associated Branch (PU-AB,T-shaped pseudounipolar axon with a branch <10uM long), PU-

Short (T-shaped pseudounipolar axon, with an axon stem <60um long), PU-Long (T-shaped

pseudounipolar axon, with an axon stem >60um long). Human cells were distinguished from

rodent cells based on expression of human nuclei-Cy3 marker. Images were blinded and intra-

rater or inter-rater reliability was calculated to be >90%.  Morphology frequency was compared

across morphologic groups: Immature (unipolar and bipolar), Multipolar, Pseudounipolar

Transition (PUT-Bulge, PUT-Bell, PUT-AB), and Pseudounipolar (PU-Short and PU-Long) using

the Chi-square test in Graphpad Prism 10. Sub-morphologies were pooled into major

morphologic categories to reach chi-square test validity requirements. Adjusted residuals with

Bonferroni correction were calculated in excel using standard formulas.

**Multi-Electrode Array Recording:**

_Cell Culture:_ rDRG glia were dissected and plated in 48-well, 16 electrode Cytoview-MEA

plates (Axion) at 20k cells/well in Glia Media. After a 30-minute incubation at 37°C 7.5% $CO_2$,

plates were vigorously shaken, and media was fully changed to remove cells in suspension.

Remaining glial cells were cultured in Glia Media at 37°C 7.5% $CO_2$ for one week with media

changes 3x/ week. On the 7[th] day, iNCPs were thawed and plated in NCATS Maturation Media +

CEPT at 150k-200k cells/ well, in coculture or alone on coated wells (100ug/mL PDL 1 hr 37°C,

10ug/mL mouse laminin 1 hr 37°C). A fraction of the wells remained glial cells alone as a

control, treated with NCATS Maturation Media +CEPT. 24 hrs after plating, media was changed

to NCATS Maturation Media + 1uM PD0332991. Cells were cultured for 5-8 weeks at 37°C 5%

$CO_2$., with ½ media changes 3x/week. n=4 biological replicates; replicates 1-3 = iPSC Gr1.1,

replicate 4 = iPSC NCRM5.

_MEA Recording:_ MEA recordings were performed on the Maestro Pro recording system from

Axion at 37°C 5% $CO_2$, 1x/week for 5 minutes, after at least 5 minutes of acclimation. AxIS

navigator software was used to record standard Spontaneous Neural and Viability properties.

Burst Detection was enabled with the following specifications: Single Electrode Minimum #

Spikes=5, Network Minimum #Spikes=50, Maximum Interspike Interval= 100s, Mean Firing Rate

detection window= 100s. Recordings were carried out weekly, and data was analyzed when an

average of 7+ active electrodes was reached for both cocultured iSNs and iSNs alone.

Electrodes were considered active with a minimum of 5 spikes/minute.

_Analysis:_ Data was analyzed at the time point when an average of 7+ electrodes were active in

both conditions. Wells with 2 or less active electrodes were excluded from analysis. Glial cells

were not active, and therefore were excluded from all analyses aside from # of active

electrodes. Data did not follow a normal gaussian distribution, and therefore averages were

compared using the Mann-Whitney U test in Graphpad Prism 10. n=# wells, pooled from 4

biological replicates.

**RNA sequencing of iSN and rDRG Glia cocultures:**

_Cell Culture and RNA Collection:_ rDRG glia were dissected and plated in uncoated glass bottom

35mm Fluorodishes (WPI cat #: FD35-100) at 250k cells/dish in Glia Media. After 30 minutes at

37°C 7.5% $CO_2$, dishes were vigorously shaken, and media was fully changed to remove cells in

suspension. Remaining glial cells were cultured in Glia Media at 37°C 7.5% $CO_2$ for one week

with media changes 3x/ week. On Day 7, cells were either treated with NCATS Maturation

Media + 1uM PD0332991, maintained in Glia Media, or co-cultured with iSNs as described

above. iSNs were plated at 90k/dish in co-culture or alone, as described above. Cultures were

maintained at 37°C 5% $CO_2$, with ½ media changes 3x/week for an additional week. On Day

14, RNA was collected from Glia Media Treated rDRG Glia, NCATS Maturation Media Treated

rDRG Glia, iSN-rDRG Glia Co-Culture, and iSN-Only mass cultures using the RNeasy Plus Mini

Kit (Qiagen cat#: 74134). n=3-4 biological replicates; Coculture: n= 2 iPSC NCRM5, n=1 GR1.1.

iSN only: n=4 iPSC NCRM5. Glial cells derived from n=4 rats.

_RNA sequencing:_ mRNA bulk sequencing was carried out at the Dana Farber Cancer Institute

Molecular Biology Core Facility.  The genomics core team performed RNA QC, rRNA depletion,

library preparation (Roche Kapa mRNA Kit), and sequencing on the Illumina NovaSeq 6000.

Reads were aligned to both the rat (_Rattus norvegicus_) and human genome using STAR aligner

to separate rDRG glia and iSN reads in co-culture. GEO Accession #: GSE272868.

_Analysis:_ Normalized counts (with raw counts <3 set to NA) were analyzed for differential gene

expression using the DESeq2 R package version 1.42.0[172] with an adjusted p-value cutoff of

0.05. Gene Ontology Enrichment Analysis was performed using the clusterProfiler R package

version 4.10.0[173] with an adjusted p-value cutoff of 0.05.

**iSN-rSG co-culture in transwell compartmentalized system:**

*Cell Culture:* rDRG glia were dissected as described above and plated at 40k cells/well on

uncoated Millicell® Standing Cell Culture Inserts (MilliporeSigma cat#: PIHP01250). rDRG glia

were prepared as described above. On the 7th day, iNPCs were plated at 15k cells/ well on

sterile 12mm #1.5 coverslips, coated as described above, in the same plate for terminal

differentiation either in compartmentalized co-culture or alone. Cells were allowed to attach to

coverglass for 30 min at 37°C 5% $CO_2$, after which Glia Media was removed from rDRG glial

inserts and inserts were transferred to wells with NPCs dedicated for terminal differentiation in

compartmentalized co-culture. 24 hrs after iNPC plating, media was changed to NCATS

Maturation Media + 1uM PD0332991. Cells were cultured for one week at 37°C 5% $CO_2$., with ½

media changes 3x/week. On the 14th day, rDRG glial culture inserts were moved to new wells

and glia attached to insert were visualized with CellTracker Red CMTPX (Thermofisher cat#:

C34552). Cells were incubated with 12.5uM dye in DMEM/F12 for 30 min at 37deg 5% CO2,

after which media was changed to DMEM/F12 and live cells were imaged at 10x TRITC on the

Nikon-Ti E microscope. iSNs adherent to coverglass were fixed for downstream

Immunofluorescence (IF) as described above. n=3 biological replicates, iPSC Gr1.1.

*Imaging and Analysis:* Photomicrographs of cells cultured on coverslips were collected on the

Nikon Ni-E confocal microscope at 60x with oil immersion. Images were obtained with 405 and

488 lasers, at 30V gain 5% power and 561 30V gain 7% power. All images were obtained in a z-

stack with 0.35um step size. Systematic random sampling was utilized to image n=50-100

cells/biological replicate across 3 coverslips. Areas with fused cell bodies and axonal

fasciculations were omitted from sampling. Inter-rater reliability was calculated >90%.

Morphology frequencies and statistics were carried out as described above.

**699**   **iSN cultures treated with co-culture spent media:**

**700**   Spent media was collected from iSN-rDRG Glia Co-Cultures over the course of one week and

**701**   stored at 4°C. On Day 21, iNPCs were plated alone on chambered coverglass in NCATS

**702**   Maturation Media + CEPT as described above. After 24 hrs, media was either changed to ½

**703**   fresh NCATS Maturation Media + 1uM PD0332991 + ½ spent media or maintained in NCATS

**704**   maturation Media + 1uM PD0332991 as described above. Cells were cultured for one week with

**705**   3x full media changes maintaining specified ratios. On Day 28, cultures were fixed for IF and

**706**   images were acquired and analyzed as described. n=3 biological replicates; n= 2 iPSC NCRM5

**707**   n= 1 iPSC Gr1.1.

**708**   **iSN-rSG co-culture with glial methanol fixation:**

**709**   rDRG glia were dissected, plated, prepared, and cultured in chambered coverglass for one

**710**   week as described above. On Day 7, glia were treated with NCATS maturation media + CEPT,

**711**   and 24 hrs later media was changed to NCATS Maturation Media + 1uM PD0332991. Glia were

**712**   maintained in NCATS Maturation Media + 1uM PD0332991 an additional week with ½ media

**713**   changes 3x/week. On day 14, cells were fixed in 95% methanol 7 min at 4°C, washed, and

**714**   treated with NCATS maturation media + CEPT for downstream iSN co-culture. iNPCs were

**715**   thawed and plated at 15k cells/well for terminal differentiation in iSN-rDRG Glial Co-Culture with

**716**   methanol fixation, iSN-rDRG Glial Co-Culture without methanol fixation (prepared as described

**717**   above), or iSN-only culture (prepared as described above). 24 hrs after iNPC plating, media was

**718**   changed to NCATS Maturation Media + 1uM PD0332991. Cells were cultured for one week at

**719**   37°C 5% $CO_2$., with ½ media changes 3x/week. After one week of coculture, cells were fixed

**720**   with PFA for downstream IF and images and analyses were carried out as described above. n=4

**721**   biological replicates, 2/ iPSC line.

**722**   **iSN-rSG co-culture with glial *Sema6a* or *GJA1* sh-RNA knockdown:**

rDRG glia were dissected, plated, prepared, and cultured for five days as described above. On

Day 5, Glia were transduced with ShRNA-Sema6a-puro-GFPtag lentivirus (Millipore Sigma

Cat#: SHCLNV), ShRNA-GJA1-puro-GFPtag lentivirus (Millipore Sigma Cat#: SHCLNV), or

ShRNA-GFP-puro control lentivirus (Millipore Sigma cat#: SHC005V) in Glia Media for 24 hrs,

after which media was changed. Cells were maintained in Glia Media 72 hrs post-transduction

and evaluated for sh-RNA expression using live-imaging in brightfield and FITC on the Nikon-Ti

E microscope. Glia were then treated on Day 8 with 1ug/mL puromycin in Glia Media for 48hrs.

Media was changed, and selected cells were allowed to proliferate for four days. On Day 14,

iNPCs were plated at 15k cells/well in NCATS Maturation Media + CEPT for terminal

differentiation in iSN-rDRG Glia Knockdown Co-Culture, iSN rDRG Glia Knockdown Control Co-

culturel and iSN-rDRG Glia Co-Culture Control (prepared as described above). 24 hrs after

iNPC plating, media was changed to NCATS Maturation Media + 1uM PD0332991, and cells

were co-cultured for one week at 37°C 5% $CO_2$, with ½ media changes 3x/week. On Day 21,

co-cultures were fixed for IF as described. n=4 biological replicates, 2/ iPSC line.

**Modeling chemotherapy induced axonal degeneration:**

*Chemotherapy treatment of iSN co-culture with rDRG glia:* rDRG glia were prepared as

described above. iSNs generated with the Deng et al. 2023[15] protocol, were plated at 15k

cells/well in rDRG glial co-culture or alone and maintained in culture for 7 days as described

above. On day 14, cells were treated with 60nM Paclitaxel (Sigma Aldrich cat #: T7402) or 0.2%

DMSO (Sigma Aldrich cat#: D2650) in NCATS Maturation Media + 1uM PD0332991 for 48 hrs.

n=4, iPSC NCRM5. Anatomic RealDRG iSNs were plated at 5k cells/well in rDRG glial co-

culture or alone and maintained in culture for 7 days as described above. On day 14, cells were

treated with 600nM Paclitaxel (Sigma Aldrich cat #: T7402) or 0.2% DMSO (Sigma Aldrich cat#:

D2650) in NCATS Maturation Media + 1uM PD0332991 for 48 hrs. n=4, Anatomic RealDRGs

are derived from one genetic iPSC line. Cells were then fixed as described above for IF.

*Immunofluorescence (IF):* Fixed cultures were permeabilized with 0.1% Triton-x in PBS 10

minutes, and blocked in blocking buffer (5% Normal Donkey Serum (Sigma Aldrich cat#: D9663)

in PBS) for 1 hr at RT. Next, they were incubated overnight at 4°C in primary antibodies (Table

11) diluted in blocking buffer. The following day, cultures were washed and incubated in Alexa-

Fluor secondary antibodies (Table 12) and DAPI diluted 1:1000 in blocking buffer for 1 hr at RT.

Cells were washed and stored in PBS at 4°C prior to imaging.

*Imaging:* All photomicrographs of cells cultured in Lab-Tek™ II Chambered Coverglass were

collected on the Zeiss-980 airy-scan/confocal microscope. Images of all conditions were

obtained at 20x with 1.3x zoom and bidirectional scanning in SRY-4 Airy-Scan configuration.

Multiplex imaging was utilized with 405, 488, 561, lasers at 800V gain, 1% laser power, and 0

offset. All images were obtained in a z-stack with 0.35um step size. Systematic random

sampling was utilized to image n=16 fields/biological replicate across 6-wells. All images were

3D airy-scan post processed in Zen Blue software.

*Analysis:* Particle analysis was carried out on gray-scale TUJ1 channel image stacks. Threshold

was manually set at the brightest z-stack slice for each image, and percent area occupied by

TUJ1 pixels was quantified for each image stack. Average values were calculated for each field.

Field averages were compared between treatment groups and culture conditions using two-way

ANOVA analysis with the Bonferroni post-hoc test in Graphpad Prism. n=64 fields from 4

biological replicates.

| Primary Antibody | Supplier:Cat # | Concentration | Experiment(s) |
|---|---|---|---|
| rbt anti-BRN3a | Abcam: ab245230 | 1:250 | Co-culture in compartmentalized system, Spent media treatment |
| rbt anti-dsRed | Takara:632496 | 1:1000 | *in vivo* DRG morphology |
| gt anti-GJA1 | Abcam: ab87645 | 1:200 | Connexin-43/GJA1 knockdown co-culture, Chemotherapy treatment co-culture |

| | | | |
|---|---|---|---|
| rbt anti-Fabp7 | ThermoFisher: PA531864 | 1:200 | Identification of glial subtypes |
| Mse anti-human nuclei-CY3 | Millipore Sigma: MAB1281C | 1:200 | Co-culture,  RealDRG co-culture, Methanol fixation co-culture |
| rbt anti-MBP | Abcam: ab218011 | 1:250 | Identification of glial subtypes |
| rbt anti-P75/ngfr | Abcam: ab52987 | 1:100/ 1:200 | Co-culture,  RealDRG co-culture, Methanol fixation co-culture, Identification of glial subtypes |
| rbt anti-OCT6 | Abcam: ab259952 | 1:200 | Identification of glial subtypes |
| Mouse anti-S100b | Sigma: S2532 | 1:1000 | Identification of glial subtypes |
| rbt anti-Sema6a | Thermofisher: PA581009 | 1:100 | Sema6a knockdown co-culture |
| ms anti-TUJ1 | Biolegend: 801202 | 1:1000 | Co-culture, RealDRG co-culture,  Co-culture in compartmentalized system, Methanol fixation co-culture, Spent media treatment, Sema6a knockdown co-culture Connexin-43/GJA1 knockdown co-culture, Chemotherapy treatment co-culture |

767 **Table 11. Primary antibodies.**

| Secondary Antibody | Supplier:Cat # | Concentration | Experiment(s) |
|---|---|---|---|
| gt anti-Ms Alexa 488 | Thermofisher: A11001 | 1:1000 | Co-Culture, RealDRG co-culture, Methanol Fixation co-culture, Co-culture in compartmentalized system, Spent media treatment, Identification of Glial Subtypes |
| gt anti Rbt Alexa 647 | Thermofisher: A21245 | 1:1000 | Co-Culture, RealDRG co-culture, Methanol Fixation co-culture |
| gt anti-Rbt Alexa 546 | Thermofisher: A11035 | 1:1000 | Co-culture in compartmentalized system, Spent media treatment, Identification of Glial Subtypes |

| | | | Sema6a Knockdown coculture, Connexin-43/GJA1 knockdown coculture, CIPN chemotherapy treatment coculture |
|---|---|---|---|
| dnky anti-Ms Alexa 568 | Thermofisher: A10037 | 1:1000 | Sema6a Knockdown coculture, Connexin-43/GJA1 knockdown coculture, CIPN chemotherapy treatment coculture |
| dnky anti-Gt Alexa 647 | Thermofisher: A21447 | 1:1000 | Connexin-43/GJA1 knockdown coculture, Chemotherapy treatment coculture |
| dnky anti-Rbt Alexa 647 | Thermofisher: A31573 | 1:1000 | Sema6a knockdown coculture |
| dnky anti-Rbt Alexa 546 | Thermofisher: A10040 | 1:1000 | *in vivo* DRG morphology |

**Table 12. Secondary antibodies.**

**Supplemental Methods:**

**Animal Care:**

Sprague-Dawley rats were purchased from Charles River Laboratory. Ntrk2-CreER/C57BL/6
mice and lox-tdtomato/C57BL/6 mice were shared by Dr. David Ginty, Harvard Medical School.
PLP1-GFP/C57BL/6 mice were shared by Dr. Lisa Goodrich, Harvard Medical School. Animals
were cared for in accordance with NIH standards and guidelines. Experimentation was
approved by the Dana Farber Cancer Institute IACUC.

**Tamoxifen induced sparse labeling of Ntrk2+ sensory neurons with tdtomato *in vivo*:**

Ntrk-CrER mice were crossed with lox-tdtomato mice and subsequently crossed with PLP1-
GFP/C57BL/6 mice. Timed pregnant Ntrk-CreER/lox-tdtomato/PLP-1GFP mice were weighed
and tamoxifen (TMXF) dose was calculated at 40ug/g body weight.  2mg/mL Tamoxifen (TMXF)
was stored in methanol at -80C and the final TMXF dose was diluted in 100uL corn oil. Diluted
TMXF was vortexed 45 minutes and dried in a Speedvac to remove methanol from the mixture,
leaving TMXF in corn-oil alone. Injections were carried out intraperitoneally at E12.5-E14.5 to
sustain expression and sacrifice at E14.5, E16.5, E18.5, P5.  At time of sacrifice, embryonic
lumbar DRGs were dissected, and expression of tdtomato and PLP1-GFP was evaluated via
epifluorescence. DRGs were then drop fixed in 4% paraformaldehyde at 4°C overnight. The
next day, DRGs were washed 3x with 1XPBS and stored at 4°C in 1X PBS until staining. n=3+
embyros total per age group.

**Quantification of DRG sensory neuron morphology frequency *in vivo*:**

*Immunofluorescence*: Fixed DRGs were permeabilized 2 hrs in 0.5% tritonX in PBS, and then
blocked in 5% Normal Donkey Serum (NDS)/0.5% tritonX in PBS 4-5hrs at RT. They were then
incubated in rbt anti-dsRed (Takara Cat#: 632496) primary antibody diluted 1:1000 in blocking
buffer at 4°C overnight, and washed the next day with 1X PBS. Next, DRGs were incubated with
dnky anti-rbt 546 (Thermofisher Cat#: A10040) secondary antibody diluted 1:500 in blocking
buffer  for 3 hrs at RT. After staining, DRGs were stored at 4°C in 1XPBS, or gradually
transferred to methanol for tissue clearing.

*DRG Tissue Clearing:* Stained DRGs stored in PBS were gradually transferred to methanol with
10 minute incubations in the following concentrations: 25%, 50%, 75%, and 100%. After
methanol dehydration, each DRG was transferred to a slide and placed within a slide spacer
(VWR Cat#102096-614). BABB solution (1 Benzyl Alcohol:2 Benzyl Benzoate) was added to the
DRG and incubated 2-3min, after which it was replaced with fresh BABB, and steps were
repeated until the tissue was cleared. Then, a circular coverslip was mounted on top with tissue
embedded in BABB solution.

*Imaging:* Confocal images were acquired on the Nikon Ni-E confocal microscope, first at 20X to
visualize the entire DRG, and then at 40x to visualize individually labeled neurons. Images were
acquired at x resolution, 2.4 speed, 0.2-0.3um step, with 488 and 561 laser at 1.04 laser power,
36 gain, 0 offset. All cells labeled with tdtomato in the DRG were imaged from the top to bottom
of neuronal cell bodies.

*Analysis:* Images were blinded after acquisition, and NIS elements software was used to
visualize all cell morphology in 2D and/or 3D. All labeled cells were categorized into the
following morphologic groups: bipolar (elongated cell body, with an axon emanating from the
two opposite poles of the cell body with ~180° angle between), intermediate (round cell body
with two axons forming <180° angle between), and pseudounipolar (one axon stem emanating
from cell body, forming a T-Shape axon). Cells were considered apoptotic with condensed cell
bodies small, and no axons attached.

**Anatomic RealDRG iSN co-culture with rDRG glia:** Frozen RealDRG iSNs were purchased
from Anatomic (cat#: 1020). rDRG glia were dissected, plated, prepared, and cultured for one
week as described above. On Day 7, RealDRG iSNs were thawed in Chrono-Senso-MM
(Anatomic) and plated at 15k, 9k, or 5k cells/well either in rDRG glia co-culture in NCATS
Maturation Media. Lab-Tek™ II Chambered Coverglass were coated with 0.01% Poly-ornithine
solution for 1 hr at 37°C, washed with dPBS, and coated with iMatrix-511 (Reprocell
cat#:NP892-021) 1:50 in dPBS 3 hrs at 37°C for iSN-only culture control. After 24 hrs, media
was either changed to NCATS maturation Media + 1uM PD0332991. Cells were cultured for
one week with 3x ½ media changes. On Day 14, cells were fixed for downstream IF and
analysis as described in text. n=3 biological replicates at densities indicated.
825
**Treatment of rDRG glia with neuronal maturation media:** rDRG glia were dissected, plated,
prepared, and cultured for one week as described above. On Day 7, glia were either treated
with NCATS Maturation Media + CEPT and cultured at 37°C 5% $CO_2$ or maintained in Glia
Media at 37°C 7.5% $CO_2$. After 24 hrs, NCATS Maturation Media + CEPT was changed to
NCATS Maturation Media + 1uM PD0332991. Glia were matured for an additional week with ½
media changes 3x/week and fixed for downstream Immunofluorescence (IF) on Day 14. First,
cells were fixed with 2% paraformaldehyde (PFA) (EMS cat#: 15710; 1:1 4% PFA to spent
media) 10 minutes at room temperature (RT), and then in 4% PFA 20 minutes at RT. Cells were
washed 3x in PBS and stored at 4°C for downstream experimentation. n=4 biological replicates.
835
836  **Identification of Glial Subtypes in rDRG Glial cultures with IF:**
837  *Immunofluorescence:* Fixed rDRG glial cultures treated with and without NCATS maturation
838  medium were stained using the protocol described in text.
839
*Imaging:* All photomicrographs of cells cultured in Lab-Tek™ II Chambered Coverglass were
collected on the Zeiss-980 airy-scan/confocal microscope. Images of all conditions were
obtained at 20x with 1.3x zoom and bidirectional scanning in SRY-4 Airy-Scan configuration.
Multiplex imaging was utilized with 405, 488, 561, lasers. All images were obtained in a z-stack

with 0.35um step size. Systematic random sampling was utilized to image n=12-15
fields/biological replicate across 8-wells. All images were 2D airy-scan post processed in Zen
Blue software. n=4 biological replicates.
847
848 _Analysis:_ Particle analysis was carried out on glial marker channels. Threshold was manually
849 set at brightest z-stack slice for each image, and percent area occupied by particles, average
850 particle size, and average mean gray value were quantified for each image stack. Average
851 values were calculated for each field, and averages were compared between treatment groups
852 using the t-test function in excel where n=# of fields.
853

854 **Quantification of rSG-iSN contacts in coculture with methanol fixation:**

855 _Immunofluorescence and Imaging:_ IF and imaging were performed as described in text. n=4
856 biological replicates
857
_Analysis:_ Number of rSGs contacting individual iSNs was quantified using the CellCounter
plugin in FIJI. Contact was defined as apposition between P75+, human Nuc. Cy3- processes
with TUJ1+, Human Nuc. Cy3+ cell body and neurites. Number of rSGs were marked by
individual DAPI+, Human Nuc. Cy3- nuclei and rSGs contacting iSNs without a nucleus present
in the field were not counted. Contacts were counted per individual iSN, and each iSN was
categorized into defined morphologic categories. Average number of total glial contacts/cell was
compared using the t-test function and average number of total glial contacts/cell for each
morphologic category was compared using Two-way ANOVA with Bonferroni post-hoc test in
GraphPad Prism 10.
867

**Figure Legends:**

**Figure 1: Human iSNs differentiated in coculture with embryonic rodent DRG glia exhibit**
**more mature morphology and physiology.** A) Schematic illustrating protocol to terminally
differentiate iSNs in coculture with embryonic rat DRG glia (rDRG). Briefly, DRGs are dissected
from E14.5 rat embryos. After 30 minutes, cultureware is shaken and cells in suspension are
removed. Glia are cultured for 1 week, after which iNCPs are thawed and plated in coculture or
alone for 1 week of maturation. B)  Photomicrographs representing distinct iSN morphologies
observed. TUJ1= Green, p75= Magenta, Human Nuclei= Red, DAPI= Cyan. Scale=50um. PU=
pseudounipolar. PU-AB: white arrowhead indicates accessory branch. PU-Long: white
arrowhead indicates the PU-Long neuron in this field. C) Frequency of each morphology
observed in coculture and alone, displayed as percentages. 4 biological replicates, 2 iPSC cell
lines. D. Representative raster plots over 2.5 minutes of MEA recording. Black lines indicate
individual spikes and blue lines indicate bursts. E) Average number of active electrodes when
average was > 7 (5-7 weeks). Coculture n= 28 wells, iSN alone n=17 wells, Glia alone n=16
wells from 4 biological replicates. F-H) Number of spikes (F), mean firing rate (G), and number
of bursts (H) when average # active electrodes were >7. Wells with <2 active electrodes were
excluded. Coculture n= 26 wells, iSN alone n=15 wells from 4 biological replicates. I) Average
burst duration in seconds when average # active electrodes were >7. Wells with <2 active
electrodes and 0 bursts were excluded. Coculture n=24 wells, iSN alone n=14 wells from 4
biological replicates. J) Interburst Interval in seconds when average # active electrodes were
>7. Wells with <2 active electrodes and <2 bursts were excluded. Coculture n= 22 wells, iSN

alone n=13 wells from 4 biological replicates. F-J) Mann-Whitney U tests performed in
Graphpad Prism 10, p<0.05=significant, p<0.0001=****.

**Figure 2. iNCPs terminally differentiated in co-culture more advanced in neuronal**
**differentiation compared to iNCPs alone.** A) Volcano plot illustrating differential gene
expression in coculture vs. alone. B) GO analysis terms upregulated in coculture vs. iSNs alone,
related to neuronal development. C) GO analysis terms downregulated in coculture vs. iSNs
alone, related to neuronal development. Coculture n=3, iSN alone n=4. P adj.<0.05.

**Figure 3. Rodent DRG satellite glia (rSGs) are responsible for advanced maturation of**
**iSNs.** A) Volcano plot illustrating differential gene expression in NCATS Maturation Media
treated rDRG glia vs. rDRG glia maintained in base glia media B-C) GO analysis terms
upregulated (B) and downregulated (C) in NCATS Maturation Media treated rDRG glia vs. rDRG
glia maintained in base glia media. D) Volcano plot illustrating differential gene expression in
cocultured rDRG glia vs. rDRG glia maintained in base glia media E-F) GO analysis terms
upregulated (E) and downregulated (F) in cocultured rDRG glia vs. rDRG glia maintained in
base glia media. G) Volcano plot illustrating differential gene expression in cocultured rDRG glia
vs. NCATS Maturation Media treated rDRG glia. n=3-4/group. P adj.<0.05.

**Figure 4. rSG secreted factors do not induce pseudounipolar morphology in hiSNs.** A)
Schematic illustrating transwell coculture protocol. B) Representative photomicrographs of iSNs
cultured in transwell coculture or alone. TUJ1= Green, BRN3a=Red, DAPI= Blue. Scale= 50um.
C) Frequency of each morphology observed in coculture and alone, displayed as percentages
.n= 3 biological replicates D) Schematic illustrating protocol for iSN treatment with coculture
spent media. E) Representative photomicrographs of iSNs treated with coculture spent media or
untreated. TUJ1= Green, BRN3a=Red, DAPI= Blue. Scale=50um. F) Frequency of each
morphology observed in Coculture Media treated and untreated iSNs, displayed as
percentages. 3 biological replicates.

**Figure    5.    Physical    contact    between    rSGs    and    hiSNs    is    essential    to**
**pseudounipolarization.** A) Schematic illustrating protocol for iSN coculture with methanol fixed
glia. B) Representative photomicrographs of iSNs in coculture with methanol fixed glia, normal
glia, or alone. TUJ1= Green, p75=Magenta, Human Nuclei= Red, DAPI= Cyan. iSNs alone were
not stained for human nuclei. Scale= 50um. C) Frequency of each morphology observed in
cocultured iSNs with methanol fixed glia and normal glia, displayed as percentages. n=4
biological replicates. D) Schematic illustrating protocol for iSN coculture with Sema6a, GJA1,
and GFP knockdown in glia. All three conditions were run in parallel. E) Representative
photomicrographs of iSNs in coculture with SEMA6a knockdown glia or GFP knockdown glia.
Sema6a=Magenta, TUJ1= Red, GFP=Green, DAPI=Cyan. Scale=50um. F) Frequency of each

morphology observed in iSNs cocultured with SEMA6a knockdown, GFP knockdown, and GJA1
Knockdown glia, displayed as percentages. n=4 biological replicates. G) Representative
photomicrographs of iSNs in coculture with GJA1 knockdown glia or GFP knockdown glia.
GJA1=Magenta, TUJ1= Red, GFP=Green, DAPI=Cyan. Scale=50um.

**Figure 6. Cocultured iSNs are more susceptible to paclitaxel mediated**
**neurodegeneration.** A) Representative photomicrographs of Deng et al. 2023 generated iSNs
in coculture or alone, treated with either 0.2% DMSO or 60nM paclitaxel. TUJ1= Green,
GJA1=Magenta, Human Nuclei= Red, DAPI= Cyan. iSNs alone were not stained for human
nuclei. Scale= 50um. B) Percent area occupied by TUJ1 particles in cocultured iSNs and iSNs
differentiated alone + 0.2% DMSO or 60nM Paclitaxel. Two-Way ANOVA with Bonferroni post-
hoc calculated in Graphpad Prism 10, $p < 0.05$= significant, $p < 0.0001$=****. n=64 fields from 4
biological replicates. C) Representative photomicrographs of Anatomic RealDRG iSNs in
coculture or alone, treated with either 0.2% DMSO or 600nM paclitaxel. TUJ1= Green,
GJA1=Magenta, Human Nuclei= Red, DAPI= Cyan. iSNs alone were not stained for human
nuclei. Scale= 50um. D) Percent area occupied by TUJ1 particles in cocultured iSNs and iSNs
differentiated alone + 0.2% DMSO or 600nM Paclitaxel. Two-Way ANOVA with Bonferroni post-
hoc calculated in Graphpad Prism 10, $p < 0.05$= significant, $p < 0.0001$=****. n=64 fields from 4
biological replicates.

**Supplemental Figure Legends:**

**Figure S1. Human iSNs differentiated in coculture with embryonic rodent DRG glia**
**exhibit advanced morphology and physiology.** A) Timeline for pseudounipolar transition was
established *in vivo* in embryonic mice expressing tamoxifen induced NTRK2-tdtomato.
Morphology frequencies from whole DRGs were quantified and categorized as apoptotic (light
gray), bipolar (dark gray), intermediate (pink), or pseudounipolar (red). DRG NTRK2+ neurons
became pseudounipolar between days E14.5 and E16.5, indicated by black arrow. E14.5 3-5
DRGs were quantified. B) Photomicrographs illustrating physical contact between satellite glia
and DRG neurons during pseudounipolarization. Green= PLP1, Red= Ntrk-tdtomato. Scale=
20um. C-D) C. Representative photomicrographs of Anatomic RealDRGs terminally
differentiated in coculture or alone with NCATS maturation media and Anatomic's proprietary
media Senso-MM. D. Photomicrographs depicting transitional and pseudounipolar morphologies
observed in cocultured Anatomic RealDRGs. TUJ1=Green, p75=Magenta, human nuclei = Red,
DAPI=Cyan. RealDRGs alone were not stained for human nuclei. Scale=50um.E-H) Average
number of active electrodes during weekly MEA recordings for fourbiological replicates.
Cocultured iSNs=purple, iSNs alone= orange, Glia alone= green. Average >7 active electrodes
were observed in both cocultured iSNs and iSNs alone at 7 weeks in replicate 1, 5 weeks in
replicate 2, 6 weeks in replicate 3, and 5 weeks in replicate 4.

**Figure S2. iNCPs terminally differentiated in co-culture more advanced in neuronal**
**differentiation compared to iNCPs alone.** A) Principal component analysis and B) Hierarchal
Cluster analysis of RNA sequencing reads from cocultured iSN and iSNs alone.

**Figure S3. Figure 3. Rodent DRG satellite glia (rSGs) are responsible for advanced**
**maturation of iSNs.** A) Representative photomicrographs of glial cultures maintained in glial

base media for 1 week and treated with NCATS Maturation Media for 1 week (Maturation
Media) or maintained in glial base media for 2 weeks (Glia Media). S100b= Green,
FABP7/OCT6/P75= Red, DAPI= Cyan. Scale = 50 um. B) Representative photomicrographs of
glial cultures maintained in glial base media for 1 week and treated with NCATS Maturation
Media for 1 week (Maturation Media) or maintained in glial base media for 2 weeks (Glia Only),
iSNs in coculture and iSNs alone. Glial Cultures: S100b=Magenta, MBP=Green, DAPI=Cyan,
iSN cultures: TUJ1=Magenta, MBP=Green, DAPI= Cyan. No MBP signal was detected in any
culture condition. Scale=50um. C) Schematic of glial culture protocol for RNA sequencing. D-E)
Principal component analysis (D) and Hierarchal Cluster analysis (E) of RNA sequencing reads
from rDRG glia in coculture, treated with maturation media, and maintained in base glia media.
F) GO terms significantly downregulated in cocultured glia compared to glia treated with
maturation media. P adj.<0.05.

**Figure S4. Physical contact between rSGs and hiSNs is essential to**
**pseudounipolarization.** A) Total rSG-iSN contacts per individual iSN in coculture with methanol
fixed glia or in normal coculture. T-test performed in GraphPad Prism 10, ****=p<0.0001. B)
Average number of rSG-iSN contacts per individual iSN in coculture with methanol fixed glia or
in normal coculture, partitioned by morphology of iSN. Two-way ANOVA with Bonferroni post-
hoc test calculated in GraphPad Prism 10. P<0.05=*, P<0.01=**, P< 0.0001=****. C) PlexinA4
mRNA reads normalized to TUBB3 (TUJ1) mRNA reads in cocultured iSNs and iSNs
differentiated alone. D) GJA1 mRNA reads normalized to S100b mRNA reads in cocultured and
maturation media treated glia.

**Acknowledgements:** NCI R01 CA205255. Schematics created with Biorender.

**Declaration of Interests:** C.J.W. is a founder of Nocion, Quralis, and Blackbox Bio, and is a
member of the SAB of Lundbeck Pharma and Tafalgie Therapeutics. All other authors have no
interests to disclose.

**Author Contributions**: Conceptualization: CJL, OYT, RAS. Generation of iNCPs: KG, VJ, DC,
PO, TC. Performance of experiments: CJL, MFPM, ES, SD, VP, SZ, OYT. Analysis: CJL, MFPM,
MT, TM. Funding Acquisition: RAS. Project Administration: CJL, RAS. Supervision: RAS, CW.
Writing: CJL, RAS, with input from all authors.

**Supplemental Index:**

1. Document S1: Supplemental Methods and Figure Legends
2. Document S2: Supplemental Figures 1-4
3. Supplemental Tables Containing Large Datasets:
a. S1: Cocultured iSN vs iSN Alone DGE
b. S2: Cocultured iSN vs iSN Alone GO
c. S3: NMM treated rDRG Glia vs rDRG Glia alone in base DGE
d. S4: NMM treated rDRG Glia vs rDRG Glia alone in base GO
e. S5: Cocultured rDRG Glia vs rDRG Glia Alone in base DGE
f. S6: Cocultured rDRG Glia vs rDRG Glia alone in base GO
g. S7: Cocultured rDRG Glia vs NMM treated rDRG Glia DGE
h. S8: Cocultured rDRG Glia vs NMM Treated rDRG Glia GO

**References:**

1.    Meltzer, S., Santiago, C., Sharma, N., and Ginty, D.D. (2021). The cellular and molecular
basis of somatosensory neuron development. Neuron *109*, 3736–3757.
https://doi.org/10.1016/j.neuron.2021.09.004.

2.    Zenker, W., and Högl, E. (1976). The prebifurcation section of the axon of the rat spinal
ganglion cell. Cell Tissue Res. *165*, 345–363. https://doi.org/10.1007/BF00222438.

3.    Nascimento, A.I., Mar, F.M., and Sousa, M.M. (2018). The intriguing nature of dorsal root
ganglion neurons: Linking structure with polarity and function. Prog. Neurobiol. *168*, 86–
103. https://doi.org/10.1016/j.pneurobio.2018.05.002.

4.    Sharma, N., Flaherty, K., Lezgiyeva, K., Wagner, D.E., Klein, A.M., and Ginty, D.D.
(2020). The emergence of transcriptional identity in somatosensory neurons. Nature *577*,
392–398. https://doi.org/10.1038/s41586-019-1900-1.

5.    Körner, J., and Lampert, A. (2022). Functional subgroups of rat and human sensory
neurons: a systematic review of electrophysiological properties. Pflugers Arch. Eur. J.
Physiol. *474*, 367–385. https://doi.org/10.1007/s00424-021-02656-6.

6.    Saporta, M.A., Bacha, A., and Shy, M.E. (2022). Peripheral neuropathies. In
Neurobiology of Brain Disorders: Biological Basis of Neurological and Psychiatric
Disorders, Second Edition (Elsevier), pp. 165–184. https://doi.org/10.1016/B978-0-323-
85654-6.00006-X.

7.    Rowe, R.G., and Daley, G.Q. (2019). Induced pluripotent stem cells in disease modelling
and drug discovery. Nat. Rev. Genet. *20*, 377–388. https://doi.org/10.1038/s41576-019-
0100-z.

8.    Nicholson, M.W., Ting, C.Y., Chan, D.Z.H., Cheng, Y.C., Lee, Y.C., Hsu, C.C., Huang,
C.Y., and Hsieh, P.C.H. (2022). Utility of iPSC-Derived Cells for Disease Modeling, Drug
Development, and Cell Therapy. Cells *11*. https://doi.org/10.3390/cells11111853.

9.    Chambers, S.M., Qi, Y., Mica, Y., Lee, G., Zhang, X.J., Niu, L., Bilsland, J., Cao, L.,
Stevens, E., Whiting, P., et al. (2012). Combined small-molecule inhibition accelerates
developmental timing and converts human pluripotent stem cells into nociceptors. Nat.
Biotechnol. *30*, 715–720. https://doi.org/10.1038/nbt.2249.

10.   Guimarães, M.Z.P., De Vecchi, R., Vitória, G., Sochacki, J.K., Paulsen, B.S., Lima, I.,
Rodrigues da Silva, F., da Costa, R.F.M., Castro, N.G., Breton, L., et al. (2018).
Generation of iPSC-Derived Human Peripheral Sensory Neurons Releasing Substance P
Elicited by TRPV1 Agonists. Front. Mol. Neurosci. *11*, 1–17.
https://doi.org/10.3389/fnmol.2018.00277.

11.   Hulme, A.J., McArthur, J.R., Maksour, S., Miellet, S., Ooi, L., Adams, D.J., Finol-
Urdaneta, R.K., and Dottori, M. (2020). Molecular and Functional Characterization of
Neurogenin-2 Induced Human Sensory Neurons. Front. Cell. Neurosci. *14*, 1–18.
https://doi.org/10.3389/fncel.2020.600895.

12.   Umehara, Y., Toyama, S., Tominaga, M., Matsuda, H., Takahashi, N., Kamata, Y.,
Niyonsaba, F., Ogawa, H., and Takamori, K. (2020). Robust induction of neural crest
cells to derive peripheral sensory neurons from human induced pluripotent stem cells.
Sci. Rep. *10*, 1–10. https://doi.org/10.1038/s41598-020-60036-z.

13.    Nickolls, A.R., Lee, M.M., Espinoza, D.F., Szczot, M., Lam, R.M., Wang, Q., Beers, J.,
1053            Zou, J., Nguyen, M.Q., Solinski, H.J., et al. (2020). Transcriptional Programming of
1054            Human Mechanosensory Neuron Subtypes from Pluripotent Stem Cells. Cell Rep. *30*,
1055            932-946.e7. https://doi.org/10.1016/j.celrep.2019.12.062.

1056    14.    Kalia, A.K., Rösseler, C., Granja-Vazquez, R., Ahmad, A., Pancrazio, J.J., Neureiter, A.,
1057            Zhang, M., Sauter, D., Vetter, I., Andersson, A., et al. (2023). How to differentiate induced
1058            pluripotent stem cells into sensory neurons for disease modelling: a comparison of two
1059            protocols. Res. Sq., 1–47. https://doi.org/10.21203/rs.3.rs-3127017/v1.

1060    15.    Deng, T., Jovanovic, V.M., Tristan, C.A., Weber, C., Chu, P.H., Inman, J., Ryu, S.,
1061            Jethmalani, Y., Ferreira de Sousa, J., Ormanoglu, P., et al. (2023). Scalable generation of
1062            sensory neurons from human pluripotent stem cells. Stem Cell Reports *18*, 1030–1047.
1063            https://doi.org/10.1016/j.stemcr.2023.03.006.

1064    16.    Clark, A.J., Kaller, M.S., Galino, J., Willison, H.J., Rinaldi, S., and Bennett, D.L.H. (2017).
1065            Co-cultures with stem cell-derived human sensory neurons reveal regulators of peripheral
1066            myelination. Brain *140*, 898–913. https://doi.org/10.1093/brain/awx012.

1067    17.    McDermott, L.A., Weir, G.A., Themistocleous, A.C., Segerdahl, A.R., Blesneac, I.,
1068            Baskozos, G., Clark, A.J., Millar, V., Peck, L.J., Ebner, D., et al. (2019). Defining the
1069            Functional Role of Na V 1.7 in Human Nociception. Neuron *101*, 905-919.e8.
1070            https://doi.org/10.1016/j.neuron.2019.01.047.

1071    18.    Odawara, A., Shibata, M., Ishibashi, Y., Nagafuku, N., Matsuda, N., and Suzuki, I. (2022).
1072            In Vitro Pain Assay Using Human iPSC-Derived Sensory Neurons and Microelectrode
1073            Array. Toxicol. Sci. *188*, 131–141. https://doi.org/10.1093/toxsci/kfac045.

1074    19.    Wheeler, H.E., Wing, C., Delaney, S.M., Komatsu, M., and Dolan, M.E. (2015). Modeling
1075            chemotherapeutic neurotoxicity with human induced pluripotent stem cell-derived
1076            neuronal cells. PLoS One *10*, 1–16. https://doi.org/10.1371/journal.pone.0118020.

1077    20.    Vojnits, K., Mahammad, S., Collins, T.J., and Bhatia, M. (2019). Chemotherapy-Induced
1078            Neuropathy and Drug Discovery Platform Using Human Sensory Neurons Converted
1079            Directly from Adult Peripheral Blood. Stem Cells Transl. Med. *8*, 1180–1191.
1080            https://doi.org/10.1002/sctm.19-0054.

1081    21.    Xiong, C., Chua, K.C., Stage, T.B., Priotti, J., Kim, J., Altman-Merino, A., Chan, D., Saraf,
1082            K., Canato Ferracini, A., Fattahi, F., et al. (2020). Human Induced Pluripotent Stem Cell
1083            Derived Sensory Neurons are Sensitive to the Neurotoxic Effects of Paclitaxel. Clin.
1084            Transl. Sci., 1–14. https://doi.org/10.1111/cts.12912.

1085    22.    Chen, Y., Saski, Y., DiAntonio, A., and Milbrandt, J. (2021). SARM1 is required in human
1086            derived sensory neurons for injury- induced and neurotoxic axon degeneration. Exp.
1087            Neurol. *339*. https://doi.org/10.4049/jimmunol.1801473.The.

1088    23.    Schinke, C., Fernandez Vallone, V., Ivanov, A., Peng, Y., Körtvelyessy, P., Nolte, L.,
1089            Huehnchen, P., Beule, D., Stachelscheid, H., Boehmerle, W., et al. (2021). Modeling
1090            chemotherapy induced neurotoxicity with human induced pluripotent stem cell (iPSC) -
1091            derived sensory neurons. Neurobiol. Dis. *155*, 105391.
1092            https://doi.org/10.1016/j.nbd.2021.105391.

1093    24.    Snavely, A.R., Heo, K., Petrova, V., Ho, T.S.Y., Huang, X., Hermawan, C., Kagan, R.,
1094            Deng, T., Singeç, I., Chen, L., et al. (2022). Bortezomib-induced neurotoxicity in human
1095            neurons is the consequence of nicotinamide adenine dinucleotide depletion. DMM Dis.

Model. Mech. *15*. https://doi.org/10.1242/dmm.049358.

25.   Rana, P., Luerman, G., Hess, D., Rubitski, E., Adkins, K., and Somps, C. (2017).
Utilization of iPSC-derived human neurons for high-throughput drug-induced peripheral
neuropathy screening. Toxicol. Vitr. *45*, 111–118.
https://doi.org/10.1016/j.tiv.2017.08.014.

26.   Jayakar, S., Shim, J., Jo, S., Bean, B.P., Singeç, I., and Woolf, C.J. (2021). Developing
nociceptor-selective treatments for acute and chronic pain. Sci. Transl. Med. *13*.
https://doi.org/10.1126/scitranslmed.abj9837.

27.   Matsuda, S., and Uehara, Y. (1984). Prenatal development of the rat dorsal root ganglia -
A scanning electron-microscopic study. Cell Tissue Res. *235*, 13–18.
https://doi.org/10.1007/BF00213717.

28.   Barber, R.P., and Vaughn, J.E. (1986). Differentiation of dorsal root ganglion cells with
processes in their synaptic target zone of embryonic mouse spinal cord: a retrograde
tracer study. J. Neurocytol. *15*, 207–218. https://doi.org/10.1007/BF01611657.

29.   Matsuda, S., Baluk, P., Shimizu, D., and Fujiwara, T. (1996). Dorsal root ganglion neuron
development in chick and rat. Anat. Embryol. (Berl). *193*, 475–480.
https://doi.org/10.1007/BF00185878.

30.   Kiss, F. (1932). SYMPATHETIC ELEMENTS IN THE CRANIAL AND SPINAL GANGLIA.
J. Anat. *66*, 488–498.

31.   Fisher, B.Y.C., and Ranson, S.W. (1933). ON THE SO-CALLED SYMPATHETIC CELLS
IN TI. J. Anat. *68*, 1–10.

32.   Truex, R.C. (1943). DEGENERATE VERSUS MULTIPOLAR NEURONS INT SENSORY
GANGLIA. J. Nerv. Ment. Dis. *97*, 606.

33.   Bhatt, S., Diaz, R., and Trainor, P.A. (2013). Signals and switches in mammalian neural
crest cell differentiation. Cold Spring Harb. Perspect. Biol. *5*.
https://doi.org/10.1101/cshperspect.a008326.

34.   Mudge, A.W. (1984). Schwann cells induce morphological transformation of sensory
neurones in vitro. Nature *309*, 367–369. https://doi.org/10.1038/309367a0.

35.   Takahashi, K., and Ninomiya, T. (1987). Morphological changes of dorsal root ganglion
cells in the process-forming period. Prog. Neurobiol. *29*, 393–410.
https://doi.org/10.1016/0301-0082(87)90020-7.

36.   Nascimento, A.I., Da Silva, T.F., Fernandes, E.C., Luz, L.L., Mar, F.M., Safronov, B. V.,
and Sousa, M.M. (2022). Sensory neurons have an axon initial segment that initiates
spontaneous activity in neuropathic pain. Brain *145*, 1632–1640.
https://doi.org/10.1093/brain/awac078.

37.   Jones, I., Yelhekar, T.D., Wiberg, R., Kingham, P.J., Johansson, S., Wiberg, M., and
Carlsson, L. (2018). Development and validation of an in vitro model system to study
peripheral sensory neuron development and injury. Sci. Rep. *8*, 1–15.
https://doi.org/10.1038/s41598-018-34280-3.

38.   Kim, H.S., Kim, J.Y., Song, C.L., Jeong, J.E., and Cho, Y.S. (2020). Directly induced
human Schwann cell precursors as a valuable source of Schwann cells. Stem Cell Res.
Ther. *11*, 257. https://doi.org/10.1186/s13287-020-01772-x.

39.   Pannese, E. (1969). Electron microscopical study on the development of the satellite cell
1139           sheath in spinal ganglia. J. Comp. Neurol. *135*, 381–421.
1140           https://doi.org/10.1002/cne.901350403.

1141   40.   Matsuda, S., Kobayashi, N., Mominoki, K., Wakisaka, H., Mori, M., and Murakami, S.
1142           (2000). Morphological transformation of sensory ganglion neurons and satellite cells. Acta
1143           Anat. Nippon. *73*, 612–613.

1144   41.   Wall, P.D., and Devor, M. (1983). Sensory afferent impulses originate from dorsal root
1145           ganglia as well as from the periphery in normal and nerve injured rats. Pain *17*, 321–339.
1146           https://doi.org/10.1016/0304-3959(83)90164-1.

1147   42.   Amir, R., and Devor, M. (1997). Spike-evoked suppression and burst patterning in dorsal
1148           root ganglion neurons of the rat. J. Physiol. *501*, 183–196. https://doi.org/10.1111/j.1469-
1149           7793.1997.183bo.x.

1150   43.   Amir, R., Michaelis, M., and Devor, M. (2002). Burst discharge in primary sensory
1151           neurons: Triggered by subthreshold oscillations, maintained by depolarizing
1152           afterpotentials. J. Neurosci. *22*, 1187–1198. https://doi.org/10.1523/jneurosci.22-03-
1153           01187.2002.

1154   44.   Djouhri, L., Koutsikou, S., Fang, X., McMullan, S., and Lawson, S.N. (2006).
1155           Spontaneous pain, both neuropathic and inflammatory, is related to frequency of
1156           spontaneous firing in intact C-fiber nociceptors. J. Neurosci. *26*, 1281–1292.
1157           https://doi.org/10.1523/JNEUROSCI.3388-05.2006.

1158   45.   Ma, C., and LaMotte, R.H. (2007). Multiple sites for generation of ectopic spontaneous
1159           activity in neurons of the chronically compressed dorsal root ganglion. J. Neurosci. *27*,
1160           14059–14068. https://doi.org/10.1523/JNEUROSCI.3699-07.2007.

1161   46.   Liu, C.N., Wall, P.D., Ben-Dor, E., Michaelis, M., Amir, R., and Devor, M. (2000). Tactile
1162           allodynia in the absence of C-fiber activation: Altered firing properties of DRG neurons
1163           following spinal nerve injury. Pain *85*, 503–521. https://doi.org/10.1016/S0304-
1164           3959(00)00251-7.

1165   47.   Xie, W., Strong, J.A., Kim, D., Shahrestani, S., and Zhang, J.M. (2012). Bursting activity
1166           in myelinated sensory neurons plays a key role in pain behavior induced by localized
1167           inflammation of the rat sensory anglion. Neuroscience *206*, 212–223.
1168           https://doi.org/10.1016/j.neuroscience.2012.01.007.Bursting.

1169   48.   Zheng, Q., Xie, W., Lückemeyer, D.D., Lay, M., Wang, X.W., Dong, X., Limjunyawong,
1170           N., Ye, Y., Zhou, F.Q., Strong, J.A., et al. (2022). Synchronized cluster firing, a distinct
1171           form of sensory neuron activation, drives spontaneous pain. Neuron *110*, 209-220.e6.
1172           https://doi.org/10.1016/j.neuron.2021.10.019.

1173   49.   North, R.Y., Li, Y., Ray, P., Rhines, L.D., Tatsui, C.E., Rao, G., Johansson, C.A., Zhang,
1174           H., Kim, Y.H., Zhang, B., et al. (2019). Electrophysiological and transcriptomic correlates
1175           of neuropathic pain in human dorsal root ganglion neurons. Brain *142*, 1215–1226.
1176           https://doi.org/10.1093/brain/awz063.

1177   50.   Zhang, X., Chen, Y., Wang, C., and Huang, L.Y.M. (2007). Neuronal somatic ATP
1178           release triggers neuron-satellite glial cell communication in dorsal root ganglia. Proc. Natl.
1179           Acad. Sci. U. S. A. *104*, 9864–9869. https://doi.org/10.1073/pnas.0611048104.

51.   Rozanski, G.M., Li, Q., and Stanley, E.F. (2013). Transglial transmission at the dorsal

root ganglion sandwich synapse: Glial cell to postsynaptic neuron communication. Eur. J.
Neurosci. *37*, 1221–1228. https://doi.org/10.1111/ejn.12132.

52.  Huang, L.Y.M., Gu, Y., and Chen, Y. (2013). Communication between neuronal somata
and satellite glial cells in sensory ganglia. Glia *61*, 1571–1581.
https://doi.org/10.1002/glia.22541.

53.  Kim, Y.S., Anderson, M., Park, K., Zheng, Q., Agarwal, A., Gong, C., Saijilafu, Young,
L.A., He, S., LaVinka, P.C., et al. (2016). Coupled Activation of Primary Sensory Neurons
Contributes to Chronic Pain. Neuron *91*, 1085–1096.
https://doi.org/10.1016/j.neuron.2016.07.044.

54.  Newberry, K., Wang, S., Hoque, N., Kiss, L., Ahlijanian, M.K., Herrington, J., and Graef,
J.D. (2016). Development of a spontaneously active dorsal root ganglia assay using
multiwell multielectrode arrays. J. Neurophysiol. *115*, 3217–3228.
https://doi.org/10.1152/jn.01122.2015.

55.  Black, B.J., Atmaramani, R., Kumaraju, R., Plagens, S., Romero-Ortega, M., Dussor, G.,
Price, T.J., Campbell, Z.T., and Pancrazio, J.J. (2018). Adult mouse sensory neurons on
microelectrode arrays exhibit increased spontaneous and stimulus-evoked activity in the
presence of interleukin-6. J. Neurophysiol. *120*, 1374–1385.
https://doi.org/10.1152/jn.00158.2018.

56.  Hiranuma, M., Okuda, Y., Fujii, Y., Richard, J.P., and Watanabe, T. (2024).
Characterization of human iPSC-derived sensory neurons and their functional
assessment using multi electrode array. Sci. Rep. *14*, 1–8.
https://doi.org/10.1038/s41598-024-55602-8.

57.  Frith, T.J.R., Granata, I., Wind, M., Stout, E., Thompson, O., Neumann, K., Stavish, D.,
Heath, P.R., Ortmann, D., Hackland, J.O.S., et al. (2018). Human axial progenitors
generate trunk neural crest cells in vitro. Elife *7*, 1–27.
https://doi.org/10.7554/eLife.35786.

58.  Libby, A.R.G., Joy, D.A., Elder, N.H., Bulger, E.A., Krakora, M.Z., Gaylord, E.A.,
Mendoza-Camacho, F., Butts, J.C., and McDevitt, T.C. (2021). Axial elongation of
caudalized human organoids mimics aspects of neural tube development. Dev. *148*.
https://doi.org/10.1242/dev.198275.

59.  Kee, Y., and Bronner-Fraser, M. (2005). To proliferate or to die: Role of Id3 in cell cycle
progression and survival of neural crest progenitors. Genes Dev. *19*, 744–755.
https://doi.org/10.1101/gad.1257405.

60.  Heng, Y.H.E., McLeay, R.C., Harvey, T.J., Smith, A.G., Barry, G., Cato, K., Plachez, C.,
Little, E., Mason, S., Dixon, C., et al. (2014). NFIX regulates neural progenitor cell
differentiation during hippocampal morphogenesis. Cereb. Cortex *24*, 261–279.
https://doi.org/10.1093/cercor/bhs307.

61.  Martynoga, B., Mateo, J.L., Zhou, B., Andersen, J., Achimastou, A., Urbán, N., van den
Berg, D., Georgopoulou, D., Hadjur, S., Wittbrodt, J., et al. (2013). Epigenomic enhancer
annotation reveals a key role for NFIX in neural stem cell quiescence. Genes Dev. *27*,
1769–1786. https://doi.org/10.1101/gad.216804.113.

62.  Harris, L., Genovesi, L.A., Gronostajski, R.M., Wainwright, B.J., and Piper, M. (2015).
Nuclear factor one transcription factors: Divergent functions in developmental versus
adult stem cell populations. Dev. Dyn. *244*, 227–238. https://doi.org/10.1002/dvdy.24182.

63.    Zalucki, O., Harris, L., Harvey, T.J., Harkins, D., Widagdo, J., Oishi, S., Matuzelski, E.,
1226             Ling Hilary Yong, X., Schmidt, H., Anggono, V., et al. (2019). NFIX-mediated inhibition of
1227             neuroblast branching regulates migration within the adult mouse ventricular-
1228             subventricular zone. Cereb. Cortex *29*, 3590–3604.
1229             https://doi.org/10.1093/cercor/bhy233.

1230    64.    He, J.C., Neves, S.R., Jordan, J.D., and Iyengar, R. (2006). Role of the Go/i signaling
1231             network in the regulation of neurite outgrowth. Can. J. Physiol. Pharmacol. *84*, 687–694.
1232             https://doi.org/10.1139/Y06-025.

1233    65.    Bromberg, K., Iyengar, R., and He, J. (2011). Regulation of neurite outgrowth by Gi/o
1234             signaling pathways. Front Biosci *13*, 4544–4557.

1235    66.    Wolf, D., Hofbrucker-MacKenzie, S.A., Izadi, M., Seemann, E., Steiniger, F., Schwintzer,
1236             L., Koch, D., Kessels, M.M., and Qualmann, B. (2019). Ankyrin repeat-containing N-Ank
1237             proteins shape cellular membranes. Nat. Cell Biol. *21*, 1191–1205.
1238             https://doi.org/10.1038/s41556-019-0381-7.

1239    67.    Kim, S.J., Woo, Y., Kim, H.J., Goo, B.S., Nhung, T.T.M., Lee, S.A., Suh, B.K., Mun, D.J.,
1240             Kim, J.H., and Park, S.K. (2022). Retinoic acid-induced protein 14 controls dendritic spine
1241             dynamics associated with depressive-like behaviors. Elife *11*, 1–30.
1242             https://doi.org/10.7554/eLife.77755.

1243    68.    Christiansen, J.H., Coles, E.G., and Wilkinson, D.G. (2000). Molecular control of neural
1244             crest formation, migration and differentiation. Curr. Opin. Cell Biol. *12*, 719–724.
1245             https://doi.org/10.1016/S0955-0674(00)00158-7.

1246    69.    Lian, Y., Wu, C., Liu, L., and Li, X. (2024). Prediction of cell-cell communication patterns
1247             of dorsal root ganglion cells: single-cell RNA sequencing data analysis. Neural Regen.
1248             Res. *19*, 1367–1374. https://doi.org/10.4103/1673-5374.384067.

1249    70.    Bennett, D.L., Clark, X.A.J., Huang, J., Waxman, S.G., and Dib-Hajj, S.D. (2019). The
1250             role of voltage-gated sodium channels in pain signaling. Physiol. Rev. *99*, 1079–1151.
1251             https://doi.org/10.1152/physrev.00052.2017.

1252    71.    Jarvis, M.F., Honore, P., Shieh, C.C., Chapman, M., Joshi, S., Zhang, X.F., Kort, M.,
1253             Carroll, W., Marron, B., Atkinson, R., et al. (2007). A-803467, a potent and selective
1254             Nav1.8 sodium channel blocker, attenuates neuropathic and inflammatory pain in the rat.
1255             Proc. Natl. Acad. Sci. U. S. A. *104*, 8520–8525.
1256             https://doi.org/10.1073/pnas.0611364104.

1257    72.    Jones, J., Correll, D.J., Lechner, S.M., Jazic, I., Miao, X., Shaw, D., Simard, C., Osteen,
1258             J.D., Hare, B., Beaton, A., et al. (2023).  Selective Inhibition of Na V 1.8 with VX-548 for
1259             Acute Pain . N. Engl. J. Med. *389*, 393–405. https://doi.org/10.1056/nejmoa2209870.

1260    73.    Jo, S., Zhang, H.X.B., and Bean, B.P. (2023). Use-Dependent Relief of Inhibition of
1261             Nav1.8 Channels by A-887826. Mol. Pharmacol. *103*, 221–229.
1262             https://doi.org/10.1124/molpharm.122.000593.

1263    74.    Haberberger, R., Scholz, A., Kummer, W., and Kress, M. (2000). M2-receptor subtype
1264             does not mediate muscarine-induced increases in [Ca2+](i) in nociceptive neurons of rat
1265             dorsal root ganglia. J. Neurophysiol. *84*, 1934–1941.
1266             https://doi.org/10.1152/jn.2000.84.4.1934.

75.    Tata, A.M., Vilaró, M.T., and Mengod, G. (2000). Muscarinic receptor subtypes

expression in rat and chick dorsal root ganglia. Mol. Brain Res. *82*, 1–10.
https://doi.org/10.1016/S0169-328X(00)00165-0.

76.   De Angelis, F., Marinelli, S., Fioretti, B., Catacuzzeno, L., Franciolini, F., Pavone, F., and
Tata, A.M. (2014). M2 receptors exert analgesic action on DRG sensory neurons by
negatively modulating VR1 activity. J. Cell. Physiol. *229*, 783–790.
https://doi.org/10.1002/jcp.24499.

77.   Xian, X.C., and Nicol, G.D. (2007). Manipulation of the potassium channel Kv1.1 and its
effect on neuronal excitability in rat sensory neurons. J. Neurophysiol. *98*, 2683–2692.
https://doi.org/10.1152/jn.00437.2007.

78.   Wang, J.J., and Li, Y. (2016). KCNQ potassium channels in sensory system and neural
circuits. Acta Pharmacol. Sin. *37*, 25–33. https://doi.org/10.1038/aps.2015.131.

79.   Smith, P.A. (2020). K+ Channels in Primary Afferents and Their Role in Nerve Injury-
Induced Pain. Front. Cell. Neurosci. *14*. https://doi.org/10.3389/fncel.2020.566418.

80.   Bhattacharya, D., Rothstein, M., Azambuja, A.P., and Simoes-Costa, M. (2018). Control
of neural crest multipotency by wnt signaling and the Lin28/let-7 axis. Elife *7*, 1–24.
https://doi.org/10.7554/eLife.40556.

81.   Shoval, I., and Kalcheim, C. (2012). Antagonistic activities of Rho and Rac GTPases
underlie the transition from neural crest delamination to migration. Dev. Dyn. *241*, 1155–
1168. https://doi.org/10.1002/dvdy.23799.

82.   Kurisaki, T., Masuda, A., Osumi, N., Nabeshima, Y.I., and Fujisawa-Sehara, A. (1998).
Spatially- and temporally-restricted expression of meltrin α (ADAM12) and β (ADAM19) in
mouse embryo. Mech. Dev. *73*, 211–215. https://doi.org/10.1016/S0925-4773(98)00043-
4.

83.   Samad, T.A., Rebbapragada, A., Bell, E., Zhang, Y., Sidis, Y., Jeong, S.J., Campagna,
J.A., Perusini, S., Fabrizio, D.A., Schneyer, A.L., et al. (2005). DRAGON, a bone
morphogenetic protein co-receptor. J. Biol. Chem. *280*, 14122–14129.
https://doi.org/10.1074/jbc.M410034200.

84.   Zorick, T.S., Syroid, D.E., Arroyo, E., Scherer, S.S., and Lemke, G. (1996). The
transcription factors SCIP and Krox-20 mark distinct stages and cell fates in Schwann cell
differentiation. Mol. Cell. Neurosci. *8*, 129–145. https://doi.org/10.1006/mcne.1996.0052.

85.   Wu, J., Peng, D., Voehler, M., Sanders, C., and Li, J. (2013). Structure and Expression of
a Novel Compact Myelin Protein - Small VCP-Interacting Protein (SVIP). Biochem.
Biophys. Res. Commun. *440*. https://doi.org/10.1016/j.bbrc.2013.09.056.Structure.

86.   Kastriti, M.E., Faure, L., Von Ahsen, D., Bouderlique, T.G., Boström, J., Solovieva, T.,
Jackson, C., Bronner, M., Meijer, D., Hadjab, S., et al. (2022). Schwann cell precursors
represent a neural crest-like state with biased multipotency. EMBO J. *41*, 1–28.
https://doi.org/10.15252/embj.2021108780.

87.   Frank, M. (2000). MAL, a proteolipid in glycosphingolipid enriched domains: Functional
implications in myelin and beyond. Prog. Neurobiol. *60*, 531–544.
https://doi.org/10.1016/S0301-0082(99)00039-8.

88.   Tasdemir-Yilmaz, O.E., Druckenbrod, N.R., Olukoya, O.O., Dong, W., Yung, A.R.,
Bastille, I., Pazyra-Murphy, M.F., Sitko, A.A., Hale, E.B., Vigneau, S., et al. (2021).

Diversity of developing peripheral glia revealed by single-cell RNA sequencing. Dev. Cell
*56*, 2516-2535.e8. https://doi.org/10.1016/j.devcel.2021.08.005.

89. Avraham, O., Chamessian, A., Feng, R., Yang, L., Halevi, A.E., Moore, A.M., Gereau,
R.W., and Cavalli, V. (2022). Profiling the molecular signature of satellite glial cells at the
single cell level reveals high similarities between rodents and humans. Pain *163*, 2348–
2364. https://doi.org/10.1097/j.pain.0000000000002628.

90. Mapps, A.A., Thomsen, M.B., Boehm, E., Zhao, H., Hattar, S., and Kuruvilla, R. (2022).
Diversity of satellite glia in sympathetic and sensory ganglia. Cell Rep. *38*.
https://doi.org/10.1016/j.celrep.2022.110328.

91. Cinquina, V., Calvigioni, D., Piscitelli, F., Keimpema, E., Di Marzo, V., Verkhratsky, A.,
and Harkany, T. (2023). Astrocytes modulate neuronal development by S100A6
signaling. bioRxiv.

92. O'Malley, A.M., Shanley, D.K., Kelly, A.T., and Barry, D.S. (2014). Towards an
understanding of semaphorin signalling in the spinal cord. Gene *553*, 69–74.
https://doi.org/10.1016/j.gene.2014.10.005.

93. Fujii, S., Yoshida, S., Inagaki, E., Hatou, S., Tsubota, K., Takahashi, M., Shimmura, S.,
and Sugita, S. (2019). Immunological Properties of Neural Crest Cells Derived from
Human Induced Pluripotent Stem Cells. Stem Cells Dev. *28*, 28–43.
https://doi.org/10.1089/scd.2018.0058.

94. Mehler, V.J., Burns, C.J., Stauss, H., Francis, R.J., and Moore, M.L. (2020). Human
iPSC-Derived Neural Crest Stem Cells Exhibit Low Immunogenicity. Mol. Ther. Methods
Clin. Dev. *16*, 161–171. https://doi.org/10.1016/j.omtm.2019.12.015.

95. Li, C., Wang, S., Chen, Y., and Zhang, X. (2018). Somatosensory Neuron Typing with
High-Coverage Single-Cell RNA Sequencing and Functional Analysis. Neurosci. Bull. *34*,
200–207. https://doi.org/10.1007/s12264-017-0147-9.

96. Ray, P., Torck, A., Quigley, L., Wangzhou, A., Neiman, M., Rao, C., Lam, T., Kim, J.Y.,
Kim, T.H., Zhang, M.Q., et al. (2018). Comparative transcriptome profiling of the human
and mouse dorsal root ganglia: An RNA-seq-based resource for pain and sensory
neuroscience research https://doi.org/10.1097/j.pain.0000000000001217.

97. Fernández-Montoya, J., Avendaño, C., and Negredo, P. (2018). The glutamatergic
system in primary somatosensory neurons and its involvement in sensory input-
dependent plasticity. Int. J. Mol. Sci. *19*. https://doi.org/10.3390/ijms19010069.

98. Nawaz, M.S., Giarda, E., Bedogni, F., Montanara, P. La, Ricciardi, S., Ciceri, D., Alberio,
T., Landsberger, N., Rusconi, L., and Kilstrup-Nielsen, C. (2016). CDKL5 and shootin1
interact and concur in regulating neuronal polarization. PLoS One *11*.
https://doi.org/10.1371/journal.pone.0148634.

99. Bushlin, I., Petralia, R.S., Wu, F., Harel, A., Mughal, M.R., Mattson, M.P., and Yao, P.J.
(2008). Clathrin assembly protein AP180 and CALM differentially control axogenesis and
dendrite outgrowth in embryonic hippocampal neurons. J. Neurosci. *28*, 10257–10271.
https://doi.org/10.1523/JNEUROSCI.2471-08.2008.

100. Montanara, P. La, Hervera, A., Baltussen, L.L., Hutson, T.H., Palmisano, I., De Virgiliis,
F., Kong, G., Chadwick, J., Gao, Y., Bartus, K., et al. (2020). Cyclin-dependent-like
kinase 5 is required for pain signaling in human sensory neurons and mouse models. Sci.

1353            Transl. Med. *12*, 1–36. https://doi.org/10.1126/SCITRANSLMED.AAX4846.

1354   101.  Hori, K., and Hoshino, M. (2017). Neuronal migration and AUTS2 syndrome. Brain Sci. *7*.
1355            https://doi.org/10.3390/brainsci7050054.

1356   102.  Lentz, S.I., Miner, J.H., Sanes, J.R., and Snider, W.D. (1997). Distribution of the ten
1357            known laminin chains in the pathways and targets of developing sensory axons. J. Comp.
1358            Neurol. *378*, 547–561. https://doi.org/10.1002/(SICI)1096-
1359            9861(19970224)378:4<547::AID-CNE9>3.0.CO;2-2.

1360   103.  Liu, C., Lin, C., Whitaker, D.T., Bakeri, H., Bulgakov, O. V., Liu, P., Lei, J., Dong, L., Li,
1361            T., and Swaroop, A. (2013). Prickle1 is expressed in distinct cell populations of the
1362            central nervous system and contributes to neuronal morphogenesis. Hum. Mol. Genet.
1363            *22*, 2234–2246. https://doi.org/10.1093/hmg/ddt075.

1364   104.  Agrawal, M., and Welshhans, K. (2021). Local Translation Across Neural Development: A
1365            Focus on Radial Glial Cells, Axons, and Synaptogenesis. Front. Mol. Neurosci. *14*, 1–18.
1366            https://doi.org/10.3389/fnmol.2021.717170.

1367   105.  Kumar, S., Chakraborty, S., Barbosa, C., Brustovetsky, T., Brustovetsky, N., and
1368            Obukhov, A.G. (2012). Mechanisms controlling neurite outgrowth in a
1369            pheochromocytoma cell line: The role of TRPC channels. J. Cell. Physiol. *227*, 1408–
1370            1419. https://doi.org/10.1002/jcp.22855.

1371   106.  Joseph, D.J., Williams, D.J., and MacDermott, A.B. (2011). Modulation of neurite
1372            outgrowth by activation of calcium-permeable kainate receptors expressed by rat
1373            nociceptive-like dorsal root ganglion neurons. Dev. Neurobiol. *71*, 818–835.
1374            https://doi.org/10.1002/dneu.20906.

1375   107.  Bakos, J., Bacova, Z., Grant, S.G., Castejon, A.M., and Ostatnikova, D. (2015). Are
1376            Molecules Involved in Neuritogenesis and Axon Guidance Related to Autism
1377            Pathogenesis? NeuroMolecular Med. *17*, 297–304. https://doi.org/10.1007/s12017-015-
1378            8357-7.

1379   108.  Wang, T., Liu, Y., Xu, X.H., Deng, C.Y., Wu, K.Y., Zhu, J., Fu, X.Q., He, M., and Luo,
1380            Z.G. (2011). Lgl1 activation of Rab10 Promotes axonal membrane trafficking underlying
1381            neuronal polarization. Dev. Cell *21*, 431–444.
1382            https://doi.org/10.1016/j.devcel.2011.07.007.

1383   109.  Stegeman, S., Jolly, L.A., Premarathne, S., Gecz, J., Richards, L.J., Mackay-Sim, A., and
1384            Wood, S.A. (2013). Loss of Usp9x Disrupts Cortical Architecture, Hippocampal
1385            Development and TGFβ-Mediated Axonogenesis. PLoS One *8*, 1–12.
1386            https://doi.org/10.1371/journal.pone.0068287.

1387   110.  Luo, X., Salgueiro, Y., Beckerman, S., Lemmon, V., Tsoulfas, P., and Park, K. (2013).
1388            Three-dimensional evaluation of retinal ganglion cell axon regeneration and pathfinding in
1389            whole mouse tissue after injury. Exp. Neurol. *247*, 653–662.
1390            https://doi.org/10.1016/j.expneurol.2013.03.001.Three-dimensional.

1391   111.  Mutalik, S.P., and Gupton, S.L. (2021). Glycosylation in axonal guidance. Int. J. Mol. Sci.
1392            *22*. https://doi.org/10.3390/ijms22105143.

1393   112.  Kuo, T.Y., Hong, C.J., and Hsueh, Y.P. (2009). Bcl11A/CTIP1 regulates expression of
1394            DCC and MAP1b in control of axon branching and dendrite outgrowth. Mol. Cell.
1395            Neurosci. *42*, 195–207. https://doi.org/10.1016/j.mcn.2009.07.006.

113. Hall, A., and Lalli, G. (2010). Rho and Ras GTPases in Axon Growth, Guidance, and
Branching. Cold Spring Harb. Perspect. Biol., 1–18.

114. Luxey, M., Jungas, T., Laussu, J., Audouard, C., Garces, A., and Davy, A. (2013). Eph:
Ephrin-B1 Forward Signaling Controls Fasciculation Of Sensory And Motor Axons. Dev.
Biol. *383*, 264–274. https://doi.org/10.1016/j.ydbio.2013.09.010.

115. Laumonnerie, C., Da Silva, R. V., Kania, A., and Wilson, S.I. (2014). Netrin 1 and Dcc
signalling are required for confinement of central axons within the central nervous
system. Dev. *141*, 594–603. https://doi.org/10.1242/dev.099606.

116. Balmer, C.W., and Lamantia, A.S. (2004). Loss of Gli3 and Shh Function Disrupts
Olfactory Axon Trajectories. J. Comp. Neurol. *472*, 292–307.
https://doi.org/10.1002/cne.20053.

117. Butler, S.J., and Tear, G. (2007). Getting axons onto the right path: The role of
transcription factors in axon guidance. Development *134*, 439–448.
https://doi.org/10.1242/dev.02762.

118. Zhao, L.L., Zhang, T., Huang, W.X., Guo, T.T., and Gu, X.S. (2023). Transcriptional
regulatory network during axonal regeneration of dorsal root ganglion neurons: laser-
capture microdissection and deep sequencing. Neural Regen. Res. *18*, 2056–2066.
https://doi.org/10.4103/1673-5374.366494.

119. Shamay-Ramot, A., Khermesh, K., Porath, H.T., Barak, M., Pinto, Y., Wachtel, C.,
Zilberberg, A., Lerer-Goldshtein, T., Efroni, S., Levanon, E.Y., et al. (2015). Fmrp
Interacts with Adar and Regulates RNA Editing, Synaptic Density and Locomotor Activity
in Zebrafish. PLoS Genet. *11*, 1–28. https://doi.org/10.1371/journal.pgen.1005702.

120. John, A., Brylka, H., Wiegreffe, C., Simon, R., Liu, P., Jüttner, R., Bryan Crenshaw, I.,
Luyten, F.P., Jenkins, N.A., Copeland, N.G., et al. (2012). Bcl11a is required for neuronal
morphogenesis and sensory circuit formation in dorsal spinal cord development.
Development *139*, 1831–1841. https://doi.org/10.1242/dev.072850.

121. Wang, S.H.J., Celic, I., Choi, S.Y., Riccomagno, M., Wang, Q., Sun, L.O., Mitchell, S.P.,
Vasioukhin, V., Huganir, R.L., and Kolodkin, A.L. (2014). Dlg5 regulates dendritic spine
formation and synaptogenesis by controlling subcellular N-cadherin localization. J.
Neurosci. *34*, 12745–12761. https://doi.org/10.1523/JNEUROSCI.1280-14.2014.

122. Noseda, R., Belin, S., Piguet, F., Vaccari, I., Scarlino, S., Brambilla, P., Boneschi, F.M.,
Feltri, M.L., Wrabetz, L., Quattrini, A., et al. (2013). DDIT4/REDD1/RTP801 is a novel
negative regulator of schwann cell myelination. J. Neurosci. *33*, 15295–15305.
https://doi.org/10.1523/JNEUROSCI.2408-13.2013.

123. Gunsch, G., Paradie, E., and Townsend, K.L. (2023). Peripheral nervous system glia in
support of metabolic tissue functions. Trends Endocrinol. Metab. *34*, 622–639.
https://doi.org/10.1016/j.tem.2023.07.004.

124. Birren, S.J., Goodrich, L. V., and Segal, R.A. (2024). Satellite Glial Cells: No Longer the
Most Overlooked Glia. Cold Spring Harb. Perspect. Biol., a041367.
https://doi.org/10.1101/cshperspect.a041367.

125. Narayanan, S.P., Flores, A.I., Wang, F., and Macklin, W.B. (2009). Akt signals through
the mammalian target of rapamycin pathway to regulate CNS myelination. J. Neurosci.
*29*, 6860–6870. https://doi.org/10.1523/JNEUROSCI.0232-09.2009.

126. Lebeau, P., Al-Hashimi, A., Sood, S., Lhoták, Š., Yu, P., Gyulay, G., Paré, G., Wayne
Chen, S.R., Trigatti, B., Prat, A., et al. (2017). Endoplasmic reticulum stress and Ca2+
depletion differentially modulate the sterol regulatory protein PCSK9 to control lipid
metabolism. J. Biol. Chem. *292*, 1510–1523. https://doi.org/10.1074/jbc.M116.744235.

127. Bartz, F., Kern, L., Erz, D., Zhu, M., Gilbert, D., Meinhof, T., Wirkner, U., Erfle, H.,
Muckenthaler, M., Pepperkok, R., et al. (2009). Identification of Cholesterol-Regulating
Genes by Targeted RNAi Screening. Cell Metab. *10*, 63–75.
https://doi.org/10.1016/j.cmet.2009.05.009.

128. Elbaz, B., Yang, L., Vardy, M., Isaac, S., Rader, B.L., Kawaguchi, R., Traka, M., Woolf,
C.J., Renthal, W., and Popko, B. (2022). Sensory neurons display cell-type-specific
vulnerability to loss of neuron-glia interactions. Cell Rep. *40*, 111130.
https://doi.org/10.1016/j.celrep.2022.111130.

129. Meriau, P., Weill, L., Puccio, H., and Agulhon, C. (2024). Frataxin deficiency in
proprioceptive neurons is causal to inflammatory and glial responses in dorsal root
ganglia.

130. Krishnan, A., Bhavanam, S., and Zochodne, D. (2018). An intimate role for adult dorsal
root ganglia resident cycling cells in the generation of local macrophages and satellite
glial cells. J. Neuropathol. Exp. Neurol. *77*, 929–941. https://doi.org/10.1093/jnen/nly072.

131. Hall, B.E., Macdonald, E., Cassidy, M., Yun, S., Sapio, M.R., Ray, P., Doty, M., Nara, P.,
Burton, M.D., Shiers, S., et al. (2022). Transcriptomic analysis of human sensory neurons
in painful diabetic neuropathy reveals inflammation and neuronal loss. Sci. Rep. *12*, 1–
16. https://doi.org/10.1038/s41598-022-08100-8.

132. Mocciaro, A., Berdougo, E., Zeng, K., Black, E., Vagnarelli, P., Earnshaw, W., Gillespie,
D., Jallepalli, P., and Schiebel, E. (2010). Vertebrate cells genetically deficient for
Cdc14A or Cdc14B retain DNA damage checkpoint proficiency but are impaired in DNA
repair. J. Cell Biol. *189*, 631–639. https://doi.org/10.1083/jcb.200910057.

133. Park, B.S., Jo, H.W., and Jung, J. (2015). Expression profile of aminoacyl-tRNA
synthetases in dorsal root ganglion neurons after peripheral nerve injury. J. Mol. Histol.
*46*, 115–122. https://doi.org/10.1007/s10735-014-9601-4.

134. Ninomiya, K., Ohta, K., Yamashita, K., Mizuno, K., and Ohashi, K. (2021). PLEKHG4B
enables actin cytoskeletal remodeling during epithelial cell-cell junction formation. J. Cell
Sci. *134*. https://doi.org/10.1242/jcs.249078.

135. Repetto, D., Camera, P., Melani, R., Morello, N., Russo, I., Calcagno, E., Tomasoni, R.,
Bianchi, F., Berto, G., Giustetto, M., et al. (2014). p140Cap regulates memory and
synaptic plasticity through src-mediated and citron-N-mediated actin reorganization. J.
Neurosci. *34*, 1542–1553. https://doi.org/10.1523/JNEUROSCI.2341-13.2014.

136. Aruga, J., and Mikoshiba, K. (2003). Identification and characterization of Slitrk, a novel
neuronal transmembrane protein family controlling neurite outgrowth. Mol. Cell. Neurosci.
*24*, 117–129. https://doi.org/10.1016/S1044-7431(03)00129-5.

137. Chu, Y., Jia, S., Xu, K., Liu, Q., Mai, L., Liu, J., Fan, W., and Huang, F. (2023). Single-cell
transcriptomic profile of satellite glial cells in trigeminal ganglion. Front. Mol. Neurosci. *16*,
1–16. https://doi.org/10.3389/fnmol.2023.1117065.

138. Luo, Y., Xu, W., Chen, H., Warburton, D., Dong, R., Qian, B., Selman, M., Gualdie, J.,

Kolb, M., and Shi, W. (2015). A novel profibrotic mechanism mediated by TGF-β-stimulated collagen prolyl hydroxylase expression in fibrotic lung mesenchymal cells. J. Pathol. *236*, 13384–13394. https://doi.org/10.1002/path.4530.A.

139. Chen, P., Cescon, M., and Bonaldo, P. (2015). The Role of Collagens in Peripheral Nerve Myelination and Function. Mol. Neurobiol. *52*, 216–225. https://doi.org/10.1007/s12035-014-8862-y.

140. Vehviläinen, P., Hyytiäinen, M., and Keski-Oja, J. (2009). Matrix association of latent TGF-beta binding protein-2 (LTBP-2) is dependent on fibrillin-1. J. Cell. Physiol. *221*, 586–593. https://doi.org/10.1002/jcp.21888.

141. Suto, F., Ito, K., Uemura, M., Shimizu, M., Shinkawa, Y., Sanbo, M., Shinoda, T., Tsuboi, M., Takashima, S., Yagi, T., et al. (2005). Plexin-A4 mediates axon-repulsive activities of both secreted and transmembrane semaphorins and plays roles in nerve fiber guidance. J. Neurosci. *25*, 3628–3637. https://doi.org/10.1523/JNEUROSCI.4480-04.2005.

142. Mauti, O., Domanitskaya, E., Andermatt, I., Sadhu, R., and Stoeckli, E.T. (2007). Semaphorin6A acts as a gate keeper between the central and the peripheral nervous system. Neural Dev. *2*, 1–17. https://doi.org/10.1186/1749-8104-2-28.

143. Haklai-Topper, L., Mlechkovich, G., Savariego, D., Gokhman, I., and Yaron, A. (2010). Cis interaction between Semaphorin6A and Plexin-A4 modulates the repulsive response to Sema6A. EMBO J. *29*, 2635–2645. https://doi.org/10.1038/emboj.2010.147.

144. Lowry Curley, J., Catig, G.C., Horn-Ranney, E.L., and Moore, M.J. (2014). Sensory axon guidance with semaphorin 6A and nerve growth factor in a biomimetic choice point model. Biofabrication *6*. https://doi.org/10.1088/1758-5082/6/3/035026.

145. Pease-Raissi, S.E., Pazyra-Murphy, M.F., Li, Y., Wachter, F., Fukuda, Y., Fenstermacher, S.J., Barclay, L.A., Bird, G.H., Walensky, L.D., and Segal, R.A. (2017). Paclitaxel Reduces Axonal Bclw to Initiate IP3R1-Dependent Axon Degeneration. Neuron *96*, 373-386.e6. https://doi.org/10.1016/j.neuron.2017.09.034.

146. Li, Y., Pazyra-Murphy, M.F., Avizonis, D., Russo, M.T., Tang, S., Chen, C.Y., Hsueh, Y.P., Bergholz, J.S., Jiang, T., Zhao, J.J., et al. (2022). Sarm1 activation produces cADPR to increase intra-axonal Ca++ and promote axon degeneration in PIPN. J. Cell Biol. *221*. https://doi.org/10.1083/jcb.202106080.

147. Chen, X., Gan, Y., Au, N.P.B., and Ma, C.H.E. (2024). Current understanding of the molecular mechanisms of chemotherapy-induced peripheral neuropathy. Front. Mol. Neurosci. *17*. https://doi.org/10.3389/fnmol.2024.1345811.

148. Paulson, S., McCabe, G., Winkler, M., Findley, T., Pascua, A., Gordon, K., Walsh, P., and Truong, V. (2024). Scalable and Efficient Generation of iPSC-derived Human Sensory Neurons for CIPN and Pain Research. In SLAS.

149. Rockel, A.F., Wagner, N., Spenger, P., Ergün, S., and Wörsdörfer, P. (2023). Neuro-mesodermal assembloids recapitulate aspects of peripheral nervous system development in vitro. Stem Cell Reports *18*, 1155–1165. https://doi.org/10.1016/j.stemcr.2023.03.012.

150. Lu, D., Shang, G., He, X., Bai, X. chen, and Zhang, X. (2021). Architecture of the Sema3A/PlexinA4/Neuropilin tripartite complex. Nat. Commun. *12*, 1–13. https://doi.org/10.1038/s41467-021-23541-x.

151. Kitsukawa, T., Shimizu, M., Sanbo, M., Hirata, T., Taniguchi, M., Bekku, Y., Yagi, T., and

Fujisawa, H. (1997). Neuropilin-semaphorin III/D-mediated chemorepulsive signals play a
crucial role in peripheral nerve projection in mice. Neuron *19*, 995–1005.
https://doi.org/10.1016/S0896-6273(00)80392-X.

152.   Xu, X.M., Fisher, D.A., Zhou, L., White, F.A., Ng, S., Snidpper, W.D., and Luo, Y. (2000).
The transmembrane protein semaphorin 6A repels embryonic sympathetic axons. J.
Neurosci. *20*, 2638–2648. https://doi.org/10.1523/jneurosci.20-07-02638.2000.

153.   Lerman, O., Ben-Zvi, A., Yagil, Z., and Behar, O. (2007). Semaphorin3A accelerates
neuronal polarity in vitro and in its absence the orientation of DRG neuronal polarity in
vivo is distorted. Mol. Cell. Neurosci. *36*, 222–234.
https://doi.org/10.1016/j.mcn.2007.07.003.

154.   Nakamura, F., Kalb, R.G., and Strittmatter, S.M. (2000). Molecular basis of semaphorin-
mediated axon guidance. J. Neurobiol. *44*, 219–229. https://doi.org/10.1002/1097-
4695(200008)44:2<219::AID-NEU11>3.0.CO;2-W.

155.   Pannese, E. (2010). The structure of the perineuronal sheath of satellite glial cells
(SGCs) in sensory ganglia. Neuron Glia Biol. *6*, 3–10.
https://doi.org/10.1017/S1740925X10000037.

156.   Kolos, E.A., and Korzhevsky, D.E. (2024). Gap Junction Protein Connexin-43 in a Rat
Dorsal Root Ganglion. Cell tissue biol. *18*, 189–198.
https://doi.org/10.1134/S1990519X23700049.

157.   Retamal, M.A., Riquelme, M.A., Stehberg, J., and Alcayaga, J. (2017). Connexin43
hemichannels in satellite glial cells, can they influence sensory neuron activity? Front.
Mol. Neurosci. *10*, 1–9. https://doi.org/10.3389/fnmol.2017.00374.

158.   Hanani, M., and Spray, D.C. (2020). Emerging importance of satellite glia in nervous
system function and dysfunction. Nat. Rev. Neurosci. *21*, 485–498.
https://doi.org/10.1038/s41583-020-0333-z.

159.   Keane, R.W., Mehta, P.P., Rose, B., Honig, L.S., Loewenstein, W.R., and Rutishauser,
U. (1988). Neural differentiation, NCAM-mediated adhesion, and gap junctional
communication in neuroectoderm. A study in vitro. J. Cell Biol. *106*, 1307–1319.
https://doi.org/10.1083/jcb.106.4.1307.

160.   Ma, C.H.E., Brenner, G.J., Omura, T., Samad, O.A., Costigan, M., Inquimbert, P.,
Niederkofler, V., Salie, R., Sun, C.C., Lin, H.Y., et al. (2011). The BMP coreceptor RGMb
promotes while the endogenous BMP antagonist noggin reduces neurite outgrowth and
peripheral nerve regeneration by modulating BMP signaling. J. Neurosci. *31*, 18391–
18400. https://doi.org/10.1523/JNEUROSCI.4550-11.2011.

161.   Sundt, D., Gamper, N., and Jaffe, D.B. (2015). Spike propagation through the dorsal root
ganglia in an unmyelinated sensory neuron: A modeling study. J. Neurophysiol. *114*,
3140–3153. https://doi.org/10.1152/jn.00226.2015.

162.   Hao, H., Ramli, R., Wang, C., Liu, C., Shah, S., Mullen, P., Lall, V., Jones, F., Shao, J.,
Zhang, H., et al. (2023). Dorsal root ganglia control nociceptive input to the central
nervous system https://doi.org/10.1371/journal.pbio.3001958.

163.   Han, C., Estacion, M., Huang, J., Vasylyev, D., Zhao, P., Dib-Hajj, S.D., and Waxman,
S.G. (2015). Human Nav1.8: Enhanced persistent and ramp currents contribute to distinct
firing properties of human DRG neurons. J. Neurophysiol. *113*, 3172–3185.

1568   https://doi.org/10.1152/jn.00113.2015.

1569 164. Lai, J., Gold, M.S., Kim, C.S., Biana, D., Ossipov, M.H., Hunterc, J.C., and Porreca, F.
1570   (2002). Inhibition of neuropathic pain by decreased expression of the tetrodotoxin-
1571   resistant sodium channel, NaV1.8. Pain *95*, 143–152. https://doi.org/10.1016/S0304-
1572   3959(01)00391-8.

1573 165. Enes, J., Haburcák, M., Sona, S., Gerard, N., Mitchell, A.C., Fu, W., and Birren, S.J.
1574   (2020). Satellite glial cells modulate cholinergic transmission between sympathetic
1575   neurons. PLoS One *15*, 1–21. https://doi.org/10.1371/journal.pone.0218643.

1576 166. Tomasello, D.L., Hurley, E., Wrabetz, L., and Bhattacharjee, A. (2017). Slick (Kcnt2)
1577   sodium-activated potassium channels limit peptidergic nociceptor excitability and
1578   hyperalgesia. J. Exp. Neurosci. *11*. https://doi.org/10.1177/1179069517726996.

1579 167. Tsantoulas, C., Denk, F., Signore, M., Nassar, M.A., Futai, K., and McMahon, S.B.
1580   (2018). Mice lacking Kcns1 in peripheral neurons show increased basal and neuropathic
1581   pain sensitivity. Pain *159*, 1641–1651. https://doi.org/10.1097/j.pain.0000000000001255.

1582 168. Warwick, R.A., and Hanani, M. (2013). The contribution of satellite glial cells to
1583   chemotherapy-induced neuropathic pain. Eur. J. Pain (United Kingdom) *17*, 571–580.
1584   https://doi.org/10.1002/j.1532-2149.2012.00219.x.

1585 169. Leo, M., Schmitt, L.I., Kutritz, A., Kleinschnitz, C., and Hagenacker, T. (2021). Cisplatin-
1586   induced activation and functional modulation of satellite glial cells lead to cytokine-
1587   mediated modulation of sensory neuron excitability. Exp. Neurol. *341*.
1588   https://doi.org/10.1016/j.expneurol.2021.113695.

1589 170. Schmitt, L.I., Leo, M., Kutritz, A., Kleinschnitz, C., and Hagenacker, T. (2020). Activation
1590   and functional modulation of satellite glial cells by oxaliplatin lead to hyperexcitability of
1591   sensory neurons in vitro. Mol. Cell. Neurosci. *105*, 103499.
1592   https://doi.org/10.1016/j.mcn.2020.103499.

1593 171. Majd, H., Amin, S., Ghazizadeh, Z., Cesiulis, A., Arroyo, E., Lankford, K., Majd, A.,
1594   Farahvashi, S., Chemel, A.K., Okoye, M., et al. (2023). Deriving Schwann cells from
1595   hPSCs enables disease modeling and drug discovery for diabetic peripheral neuropathy.
1596   Cell Stem Cell *30*, 632-647.e10. https://doi.org/10.1016/j.stem.2023.04.006.

1597 172. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
1598   dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 1–21.
1599   https://doi.org/10.1186/s13059-014-0550-8.

1600 173. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L.,
1601   et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
1602   Innovation *2*, 100141. https://doi.org/10.1016/j.xinn.2021.100141.

1603

**A.**

**B.** Upregulated in Coculture vs. iSN Alone

**C.** Downregulated in Coculture vs. iSN Alone

**A.**

**Coculture**        **iSN Alone**

0.2% DMSO    60nM Paclitaxel      0.2% DMSO    60nM Paclitaxel

GJA1
Human
nuc
DAPI
TUJ1

GJA1
DAPI

**B.**

**% Area TUJ1**

%Area

40 — ****
30 —  ****  ****
20 —
10 —
0 —

Coculture    iSN Alone
Culture Condition

■ DMSO
■ Paclitaxel

**C.**

**Coculture**        **iSN Alone**

0.2% DMSO    600nM Paclitaxel      0.2% DMSO    600nM Paclitaxel

GJA1
Human
nuc
DAPI

GJA1
DAPI
TUJ1

**D.**

**% Area TUJ1**

%Area

20 — ****
15 —  ****  ****
10 —
5 —
0 —

Coculture    iSN Alone
Culture Condition

■ DMSO
■ Paclitaxel

S1.

**A.** Ntrk2+ neurons

% neuron morphology

apoptotic
bipolar
intermediate
pseudo-unipolar

| | E14.5 | E16.5 | E18.5 | P5 |
|---|---|---|---|---|
| #neurons: | 51 | 216 | 151 | 60 |

44%
50%
5%

14%
83%

1%
99%

100%

**B.** TUJ1  PLP1  Merge

20um

20um

**C.** Real DRG Coculture, NCATS Maturation   RealDRG, NCATS Maturation   RealDRG, SensoMM Maturation

TU1
P75
Human nuclei
DAPI

**D.** Bulge   Bell   PU-Short

**E.** Active Electrodes Replicate 1

Avg # Active Electrodes

Glia Alone
Coculture
iSN Alone

iSN weeks in culture

**F.** Active Electrodes Replicate 2

Avg # Active Electrodes

Glia Alone
Coculture
iSN Alone

iSN weeks in culture

**G.** Active Electrodes Replicate 3

Avg # Active Electrodes

Glia Alone
Coculture
iSN Alone

iSN weeks in culture

**H.** Active Electrodes Replicate 4

# Active Electrodes

Glia Alone
Coculture
iSN Alone

iSN weeks in culture

S2.

S3.

S4.