Satellite glial contact enhances differentiation and maturation of human iPSC-derived

sensory neurons

Chelsey J. LeBlang[1,2], Maria F. Pazyra-Murphy[1], Elizabeth S. Silagi[1,2], Srestha Dasgupta[1,2],

Marianna Tsolias[1], Toussaint Miller[1], Veselina Petrova[2,3], Shannon Zhen[3], Vukasin Jovanovic[4],

David Castellano[4], Kathryn Gerrish[4], Pinar Ormanoglu[4], Carlos Tristan[4], Ilyas Singeç[4], Clifford J.

Woolf[2,3], Ozge Tasdemir-Yilmaz[1,2,5], Rosalind A. Segal*[1,2,5].

[1]Department of Cancer Biology, Dana Farber Cancer Institute, Boston, MA 02215

[2]Department of Neurobiology, Harvard Medical School, Boston, MA 02215

[3]F.M. Kirby Neurobiology Center, Boston Children's Hospital, Boston, MA 02215

[4]Stem Cell Translation Laboratory, National Center for Advancing Translational Science,
National Institute of Health, Bethesda, MD, 20892

[5]co-senior author

*corresponding author: rosalind_segal@dfci.harvard.edu

Abstract: Sensory neurons generated from induced pluripotent stem cells (iSNs) are used to
model human peripheral neuropathies, however current differentiation protocols produce sensory
neurons with an embryonic phenotype. Peripheral glial cells contact sensory neurons early in
development and contribute to formation of the canonical pseudounipolar morphology, but these
signals are not encompassed in current iSN differentiation protocols. Here, we show that terminal
differentiation of iSNs in co-culture with rodent Dorsal Root Ganglion satellite glia (rSG) advances
their differentiation and maturation. Co-cultured iSNs develop a pseudounipolar morphology
through contact with rSGs. This transition depends on semaphorin-plexin guidance cues and on
glial gap junction signaling. In addition to morphological changes, iSNs terminally differentiated in
co-culture exhibit enhanced spontaneous action potential firing, more mature gene expression,
and increased susceptibility to paclitaxel induced axonal degeneration. Thus, iSNs differentiated
in coculture with rSGs provide a better model for investigating human peripheral neuropathies.

Keywords: Satellite Glia, Pseudounipolar, iSN, Sensory Neuron, DRG, Neuropathy, CIPN.

Introduction:

Dorsal Root Ganglion (DRG) neurons are responsible for communicating somatosensory input from the outside world to the central nervous system[1]. They relay these signals through a single specialized T-shaped axon, which projects to both the periphery and the spinal cord[1–3]. DRG neurons include multiple phenotypic subtypes, which respond to specific stimuli and exhibit distinct patterns of gene expression[1,4,5]. DRG neurons are subject to a large number of disorders including hereditary neuropathies, diabetic neuropathy, and chemotherapy induced peripheral neuropathy[6], therefore developing appropriate human model systems is a major goal for researchers.

The use of differentiated human induced pluripotent stem cells (hiPSCs) has become a gold standard for studying disease pathology in relevant cell types and is increasingly accessible[7,8]. Over the past decade, several protocols have been generated to differentiate standard and patient derived hiPSCs into somatosensory neurons (iSNs, hiPSC-derived sensory neurons)[9–15]. Researchers have developed these protocols and utilized iSNs to study neuropathic pain signaling[10,16–18], mechanisms underlying debilitating and painful neuropathies[19–24], and development of treatments for these disorders[17,23,25,26].

During sensory neuron differentiation, iPSCs undergo changes mimicking embryonic development, altering their transcriptional program to progressively resemble neuroectoderm, neural crest progenitor cells, and ultimately sensory neurons, through treatment with combinations of small molecules over time and/or genetic manipulation[9–15]. Cells produced using these protocols express many hallmark proteins associated with sensory neuron identity, however iSNs exhibit an embryonic phenotype, different from that of postnatal neurons[4]. In the developing mammalian embryo, DRG neurons initially are bipolar in morphology and undergo a series of morphological changes leading to a pseudounipolar morphology before birth[27–29]. Current differentiation protocols generate iSNs that are predominantly bipolar in morphology, reflecting

the early embryonic state, or a multipolar morphology, similar to that observed following an acute

injury *in vivo*[30–32]. While developing sensory neurons alter their transcriptome over the course of

prenatal development and express transcription factors that contribute to neuronal subtyping[1,33],

iSNs remain functionally unspecialized. Therefore, modifications of sensory neuron differentiation

protocols to achieve a further differentiated phenotype would greatly advance efforts to study

postnatal homeostasis and disease at a molecular level, in a way that is more translatable to

human health or disease conditions.

We have sought to leverage developmental cues to advance the maturation of differentiated

sensory neurons. While specific mechanisms underlying pseudounipolarization and subtyping are

unknown, *in vitro* studies show that primary sensory neurons co-cultured with mixed DRG glia,

including Schwann cells (SCs) and satellite glia (SGs), become pseudounipolar[34–36]. Yet, iSNs

differentiated in a co-culture with induced or primary SCs, undergo no morphological changes,

aside from myelination[16,37,38]. *In vivo,* intimate contact is initiated between sensory neurons and

satellite glia during the early developmental period[39,40]*,* therefore we hypothesized that satellite

glia contribute to DRG sensory neuron maturation, and that terminal differentiation of iSNs in

coculture with satellite glial cells would advance this. Here, we show that physical contact between

satellite glia and iSNs during differentiation significantly advances morphological and

physiological development compared to those differentiated alone, and both semaphorin-plexin

signaling and gap junction signaling contribute to this maturation process. We also tested the

utility of this model and demonstrate that cocultured iSNs are more susceptible to toxic

neuropathy compared to iSNs alone. This work not only provides an advanced iSN model for use

by researchers in future studies, but also provides insight into mechanisms underlying sensory

neuron development.

Results:

Human iSNs differentiated in coculture with embryonic rodent DRG glia exhibit advanced
morphology:

Currently, differentiation protocols used to generate iSNs from iPSCs produce sensory

neurons with predominantly immature morphologies[9–15]. While the molecular mechanisms

underlying embryonic transition to pseudounipolar morphology are unknown, our data (Fig. S1A-

B) and others[36,40] suggest that satellite glia may play a pivotal role in morphological maturation,

through contact with sensory neurons during this developmental period. To test this, we used our

previously published protocol[15] to terminally differentiate human iPSC-derived neural crest

precursors (iNCPs) alone, or together with peripheral glia prepared from rat E14.5 DRGs (Fig.

1A). After one week, cells were fixed and stained to visualize neuronal morphology (TUJ1,

cytoskeletal marker, green) and contact with rDRG satellite glia (p75, NGF receptor, magenta)

(Fig 1B).

Morphologies of human neurons were identified and quantified using the rules specified in the

methods section. Morphology categories compared consisted of "immature" (including unipolar

and bipolar cells) "multipolar", "Pseudounipolar Transition (PUT)" (including PUT-Bulge, PUT-Bell,

and PU-Associated Branch), and "pseudounipolar (PU)" (including PU-Short and PU-Long) (Fig.

1B). Frequencies observed for each morphology category (Table 1) indicate that iSNs

differentiated in coculture include significantly more transitioning and pseudounipolar cells (PUT

adj. residual = 7.215, PU *adj. residual* = 6.350) and fewer cells with an immature morphology

(immature *adj. residual* = -5.555, multipolar *adj. residual* = -3.653) compared to iSNs differentiated

alone (PUT *adj. residual* = -7.215, PU *adj. residual* = -6.350 ,immature *adj. residual* = 5.555,

multipolar *adj. residual* = 3.653) (Fig. 1C). No differences were observed between iSNs derived

from different iPSC cell lines. These results suggest that terminal differentiation in co-culture with

DRG glia enhances the maturation of sensory neuron morphology.

To determine whether the ability of peripheral glia to promote a mature, pseudounipolar

morphology is restricted to only one of the protocols used to generate iSNs, we also tested our

co-culture protocol using RealDRG[TM] neurons from Anatomic. RealDRGs were terminally

differentiated in coculture or alone, using Senso-MM maturation media by Anatomic as a control.

RealDRGs cocultured with rDRG Glia in NCATS Maturation Media (NMM)[15] undergo extensive

morphologic changes, although the frequency of transition to pseudounipolar morphology was

reduced compared to Deng et al. 2023[15] generated iSNs (Fig S1C-D). Treatment of RealDRGs

cultures alone with NMM did not impact observed cell health or morphology but did result in the

presence of neural precursor cells, which are not typically observed in RealDRG cultures

maintained in Anatomic Senso-MM Media (Fig S1C-D).

	Morphology Frequencies								
	Immature		**Multipolar**	**Pseudounipolar Transition (PUT)**			**Pseudounipolar (PU)**		
Condition	**Unipolar**	**Bipolar**	**Multipolar**	**PUT-Bulge**	**PUT-Bell**	**PUT-AB**	**PU-Short**	**PU-Long**	**Totals**
Coculture	30	22	151	43	31	9	34	9	329
iSN Alone	0	101	174	11	1	0	0	0	287
Totals	153		325	95			43		616

Table 1. Raw frequency of morphology subtypes observed in co-cultured iSNs vs. iSNs terminally differentiated
alone. X^2 frequency statistic calculated for major morphology groups: Immature, Multipolar PUT, and PU. (X^2=
111, df=3, p<0.001).

Human iSNs differentiated in coculture with embryonic rodent DRG glia exhibit advanced
physiology:

We next assessed the impact of the coculture strategy on development of spontaneous

electrophysiologic signaling in iSNs. DRG sensory neurons can transmit sensory input to the CNS

via single action potential spikes, as well as synchronous action potential bursts[41,42]. While

spontaneous activity is relatively low[41–45] increased spiking and bursting is observed in models of

neuropathic pain both *in vivo*[41,42,45–48] and *in vitro*[42,43,45,49]. Therefore, it is crucial that iSNs develop

physiologic signaling capability to study the pathogenesis of neuropathies and develop potential

treatments.

Characterization of action potential firing using Multi-Electrode Array (MEA) recordings shows

that iSNs generated with the Deng et al. 2023[15] protocol exhibit nearly undetectable baseline

spontaneous activity after the typical two-week maturation period. Since peripheral satellite glial

cells play an important role in the regulation of somatosensory neuron firing[50–53], we tested the

hypothesis that iSNs differentiated in coculture would exhibit more mature electrophysiologic

properties by measuring spontaneous neural activity of iSNs differentiated in coculture, or alone,

using Axion MEA plates containing 16 electrodes/well. Activity was measured on the Maestro Pro

recording system for 5 minutes weekly over the span of 5-7 weeks, and neuronal spiking and

bursting was analyzed using AxIS software. Overall, cocultured iSNs became robustly active

earlier than iSNs differentiated alone, as indicated by a higher number of active electrodes

beginning around week 3 of maturation (Fig. S1E-H). Data was analyzed when an average of 7

or more electrodes were active per well in both iSN culture conditions (Fig. 1E), this occurred

between weeks 5-7 across 4 experimental replicates (Fig. S1E-G). Glia alone were inactive and

do not contribute to signals recorded in the coculture (Fig. 1E, Fig. S1E-H).

Mature iSNs differentiated in coculture were more active than iSNs differentiated alone (Fig.

1E), with a significantly higher number of spikes elicited over the recording period (Mann-Whitney

Test, p= 0.0162) (Fig. 1F), as well as a significantly higher Mean Firing Rate (MFR) (Mann-

Whitney Test, p= 0.013) (Fig. 1G). The average MFR observed in cocultured iSNs, 1.657 Hz, is

comparable to MFRs previously reported in primary DRG neurons[54,55] and iSNs cultured long-

term[56]. No significant difference was observed in the total number of bursts recorded, however

the range was quite different, with a maximum burst number of 971 detected in coculture and only

341 in iSNs alone (Fig. 1H). Interestingly, cocultured iSNs elicited significantly longer burst trains

(Mann-Whitney Test, p= <0.0001) (Fig. 1I), with a significantly shorter period between bursts

(Mann-Whitney Test, p= 0.0376) (Fig. 1J), thus spending more time in a burst state. These results

suggest that cocultured iSNs are more physiologically mature than iSNs cultured alone.

iNCPs terminally differentiated in co-culture are advanced in neuronal differentiation:

To identify the transcriptional programs that promote iSN maturation in coculture, we carried

out RNA sequencing of iSNs terminally differentiated in coculture or alone. mRNA reads from

cocultures were aligned to both the human and rat genome to distinguish mRNAs derived from

hiSNs and rDRG glia respectively. PCA analysis confirmed that iSNs terminally differentiated in

co-culture cluster separately from iSNs differentiated alone, illustrating transcriptional differences

between these two groups (Fig. S2). Differential gene expression analysis yielded 476

significantly (p<0.05) upregulated genes in cocultured iSNs (Fig. 2A, Table S1). Genes implicated

in posterior thoracic neural crest development[57,58] (*HoxB9*, p adj.=$6.922e^{-15}$) and proliferation[59]

(*ID3,* p adj.=$1.306e^{-09}$) were among the top genes upregulated. Importantly, the most highly

upregulated gene, *NFIX* (p adj.=$1.025e^{-16}$), is associated with onset of neuronal differentiation and

quiescence of neural stem cells [60–62] signaling a shift towards a more mature neuronal phenotype.

Interestingly, *NFIX*, has also been shown to regulate neuroblast branching in adult

neurogenesis[63]. Additional changes in expression of genes underlying neuronal morphogenesis

include significant upregulation of *GPRIN3* (p adj.=$6.437e^{-12}$) and *RAI14* (p adj.= $4.910e^{-09}$),

proteins essential to axonal outgrowth[64,65] and dendritic development[66,67] respectively, through

regulation of actin cytoskeletal structure. These changes illustrate potential transcriptional

differences that underly the morphological maturation in cocultured iSNs.

Gene Ontology (GO) analysis provided further information on the upregulated gene

programs, with 185 significantly upregulated gene terms in cocultured iSNs compared to iSNs

alone (Fig. 2B, Table S2). The most significantly upregulated biological processes include

increased Bone Morphogenesis Pathway (BMP) signaling and anterior/posterior pattern

specification, which confirm commitment to the neural crest cell lineage[33,68] (Table S2). Several

biological processes related to neuronal differentiation and morphogenesis were also significantly

upregulated, including neural precursor proliferation, neuronal differentiation, and neuronal fate

commitment, (Fig 2B). Interestingly, we also observed a significant upregulation of genes

associated with axonogenesis (Fig 2B). These genes specifically influence axonal specification,

outgrowth, guidance, branching, and synaptogenesis (Table 2, Table S2), and may potentially

drive the pseudounipolarization process.

Functional maturation was also evident with the upregulated expression of biological

processes and cellular components related to synaptic structure and development, including

synaptic membrane proteins, receptors, and ion channels localized to synapses in DRG

neurons[4,69] (Fig 2B). Specifically, we observed increased expression of genes encoding sodium

ion channels (*SCN7a*/*SCN10a*) (Table S2). *SCN10a* is highly expressed in DRG nociceptors,

encoding the sodium channel Nav1.8[4,70], critical for nociception[71–73]. We also observed

upregulation of muscarinic receptors (*CHRM2/CHRM3*) essential to cholinergic signaling in DRG

neurons[74–76], and increased expression of potassium ion channels (*KCNA1/KCNQ5)* that limit

action potential firing and maintain resting membrane potential in adult DRG sensory neurons[77–

79] (Table S2). Lastly, we observed significant upregulation in AMPAR gene *GRIA2* and GABA

receptors (*GABRA1/GABRA5*) which are expressed in several adult DRG neuronal subtypes[4,69]

(Table S2). Overall, these transcriptional changes may underly the observed physiologic

maturation of co-cultured iSN.

Conversely, 748 significantly (p<0.05) downregulated genes were identified in cocultured iSNs

compared to iSNs alone (Fig. 2A, Table S1). There was decreased expression of genes

implicated in early-stage neural crest development (*Lin28a*, p adj.=1.249^{-21}; *RTKN,* p adj.=9.661-

15)[80,81] as well as a gene integral to differentiation of neural crest progenitors into mesenchymal

lineage cells (*ADAM12*, p adj.=1.069^{-42})[82] (Fig. 2A). We also observed a highly significant

decrease in *RGMB* expression (p adj.=9.985^{-28}), a BMP signaling partner which has been shown

to promote ventral neuronal differentiation from neural crest cells[83] (Fig. 2A). Additionally, we

observed a significant decrease in expression of genes associated with peripheral gliogenesis,

including myelinating Schwann cell markers (*Pou3F1*[84]*,* p adj.=4.872^{-12}, *SVIP*[85] p adj.=5.928^{-15},

GPR17[86] p adj.= 7.605^{-1}*, MAL*[87] p adj.=1.120^{-8}*, PMP22*[88,89] p adj.=1.325^{-5}, satellite glial markers

(*TRDN*[89] p adj.=4.190^{-20}, *BCAN*[88,89] p adj.=7.605^{-11}, *CHL1*[90] p adj.=1.863^{-7}), and glial proteins

integral to axonal development (*S100A6*[91] p adj.=1.481^{-14}*, SEMA3B*[92] p adj.=8.410^{-10}, *SEMA3G*[92]

p adj.=9.360^{-8}) (Fig. 2A) suggesting enhanced commitment to a neuronal fate when iSNs are

differentiated in co-culture.

Gene Ontology analysis yielded 134 significantly downregulated gene programs in iSNs

differentiated in coculture compared to iSNs differentiated alone (Fig. 2C, Table S2). Gene

programs associated with MHC I/II complex assembly and antigen presentation were among the

most significantly downregulated biological processes in cocultured iSNs (Table S2), consistent

with recent data that iNCPs express these proteins at low levels, if at all[93,94]. We also observed

significant downregulation in gene programs associated with gliogenesis, glial differentiation,

myelination, and axon ensheathment (Fig. 2C), consistent with results suggesting iSNs

differentiated in coculture are committed to a neuronal, rather than a glial, fate. Interestingly, GO

terms also included transcription factors implicated in the early specification of DRG neuronal

subtypes including *NTRK3* and *SHOX2* (expressed in proprioceptors), *RET* (expressed in

nonpeptidergic nociceptors and mechanoreceptors) *POU4f3* (expressed in a subpopulation of

peptidergic nociceptors) and *GFRA3* (expressed in a subpopulation of peptidergic

nociceptors)[1,4,95,96] (Table S2). These results indicate that iSNs in co-culture are committed to a

sensory neuron fate and restrict their transcription factor repertoire, which may advance

development of functional DRG subtypes.

GO analysis also provides potential explanation for a change in neuronal physiology with

decreased expression of genes involved in the development, composition, and function of

synaptic membrane structures (Fig. 2C), including glutamate receptors, such as kainate receptors

(*GRIK1/GRIK3*), AMPAR *GRIA4,* and Delta receptor *GRID2*[97]*,* as well as purinergic receptors

(*P2RY1, P2RX6*) (Table S2). Interestingly, potassium ion channel expression (*KNCT2, KNCJ3,*

KCNJ12, KCNS1) was also decreased, which could contribute to the increased excitability we

observe in cocultured iSNs compared to iSNs differentiated alone[79] (Table S2). Taken together

transcriptomic analyses indicate that coculture of iNCPs with peripheral glia enhances

differentiation into more mature sensory neurons.

Role in Axonogenesis	Gene Names
Specification	*CDKL5*[98]/*PICALM* [99]
Outgrowth	*CDKL5*[98,100]/*AUTS2*[101] /*LAMA2*[102] /*PRICKLE1*[103]/ *EIF4G2* [104]/ *TRPC5*[105] /*ADARB1*[106]/*ENAH*[107]/ *RAB10*[108]; *USP9X*[109]/ *SOCS3*[110]
Branching	*NTN1*[111] / *BCL11A*[112]
Guidance	*EXT1*[111] /*PAK3*[113] /*EFNB2*[114]/ *NTN1*[111,115]/ *UNC5C*[115], *GLI3*[116], *FOXD1*[117]
Synaptogenesis	*NTN1*[111]/*SLITRK1/3*[118] /*ADARB1*[119] / *BCL11A*[120] / *DLG5*[121]

Table 2. Genes upregulated in cocultured iSNs that are implicated in aspects of axonogenesis.

Rodent DRG satellite glia (rSGs) are responsible for advanced maturation of iSNs:

Our studies demonstrate that iSNs differentiated in coculture with embryonic rat DRG cells

mature at an expedited rate compared to iSNs differentiated alone, however due to the protocol

used to prepare the rDRG cells, the glial subtype(s) driving this maturation was not initially clear.

The rodent DRG cell population includes satellite glia (SGs), myelinating and non-myelinating

Schwann cells (SCs), glial progenitor cells (GPCs), fibroblasts, endothelial cells, and immune

cells. When initially plated (Fig. 1A, Day 0-7), rDRG cells display a multitude of morphologies,

with small islands of semilunar cells and large flat cells, and this variety in cellular profiles is

maintained as long as the cells remain in Glia Base Media (Fig. S3A-B). However, the morphology

and cell number of rDRG glia changes upon addition of Deng et al. 2023[15] NCATS Maturation

Media (NMM), which is required for terminal differentiation of iSNs (Fig S3A-B). We used RNA

sequencing to determine the impact of both NMM treatment and iSN contact on rodent glial

identity and function (Fig. S3C), mRNA reads from co-cultured RNAs were aligned to both the

human and rat genome to differentiate mRNAs derived from hiSNs and rDRG glia respectively.

PCA analysis and hierarchal clustering (Fig. S3D-E) showed that rDRG glia treated with NMM

cluster together with cocultured rDRG glia treated with NMM. All glia treated with NMM cluster

separately from rDRG Glia Alone maintained in base media, suggesting that media composition

drives the transcriptional changes observed, while iSN contact leads to minimal changes in the

glial transcriptome.

DGE analysis yielded 1160 significantly (p<0.05) upregulated genes in NMM treated rDRG

glia compared to rDRG glia in Glia media (Fig 3A, Table S3). Overall, top upregulated genes in

NMM treated glia are associated with mature satellite glia (SGs). The most upregulated genes,

DDIT4 (p adj.=1.196^{-39}) and *INSIG1* (p adj.=1.196^{-39}), have previously been identified as DRG SG

specific markers in mouse sc-RNAseq studies[88,90]. *DDIT4* has been implicated in suppression of

myelination[122] and *INSIG1*[90] plays a role in cholesterol biosynthesis. Lipid synthesis and

metabolism are established functions of DRG SGs[123,124] and all top upregulated genes with NMM

treatment are involved in these processes including *PRR5*[125] (p adj.=2.185^{-23}), *PCSK9*[126] (p

adj.=2.067^{-21}), *HMGCS1*[89] (p adj.=5.928^{-21}), *TMEM97*[127] (p adj.=9.604^{-19}). and *SCD*[89,90] (p

adj.=2.762^{-18}). *HMGCS1* and *SCD*[89], and their related upregulated genes, *HMGCR* (p adj.=1.537^{-}

^{12}) and *SCD2*[90] (p adj.=4.821^{-15}), have also previously been identified as DRG SG gene makers

in sc-RNA seq studies. GO analysis (Fig. 3B, Table S4) confirmed that lipid biosynthesis and

metabolism pathways were significantly upregulated in NMM cultures.

We also observed significant upregulation in biologic processes, cellular components, and

molecular functions related to extracellular matrix (ECM) production. These include increased

expression of SG genes related to collagen production, such as *COL16A1, COL1A1, COL3A1,*

and *COL5A1*[90], and other ECM related genes previously associated with SG identity, *FBLN5*[89,90]

and *ANXA5*[90] (Table S4). Significant upregulation in biological processes related to DNA

replication (Table S4) indicate that enrichment for SGs may be due to enhanced cellular

proliferation. Together these results support that treatment with NMM results in an increased

proportion of mature SGs in our cultures.

DGE analysis also yielded 933 significantly (p<0.05) downregulated genes in NMM treated

rDRG glia compared cultures maintained in base Glia Media (Fig. 3A, Table S3). Overall, these

results indicate a decrease in immune functions in NMM treated cultures. Top downregulated

genes were key markers of DRG macrophages including $STAB1$[128,129] (p adj.=2.984^{-30}), $LYZ2$[89]

(p adj.=4.974^{-22}), $CSF1R$[130] (p adj.=9.916^{-20}), $MSR1$[131] (p adj.=1.090^{-19}), and $AIF1$[89] (p adj.=4.859^{-}

^{16}). GO analysis (Fig. 3C, Table S4) confirmed that the most downregulated biological processes

in NMM treated rDRG glia are related to various aspects of leukocyte function including leukocyte

differentiation, proliferation, homeostasis, activation, migration, cytokine production, and

chemotaxis. Significant decrease in expression of gene programs related to both mononuclear

cells (phagocytosis, macrophage activation, macrophage differentiation) and lymphocytes (T-cell

proliferation, T-cell activation, B-cell differentiation, B-cell activation, adaptive immune response)

are observed (Table S4). GO analysis also reveals decreased expression of genes related to

several aspects of actin cytoskeletal organization, which may contribute to morphologic changes

observed with NMM treatment. Together these data indicate the composition of NMM results in

an enrichment of SGs. When we compared transcriptomic differences between co-cultured rDRG

glia treated with NMM and rDRG glia maintained in Glia Media, we found that differentially

expressed genes were nearly identical to those identified in NMM treated cultures without iSN

contact (Fig. 3D-F, Table S5-6), indicating that NMM treatment drives an enrichment of SGs in

rDRG glial cultures.

To elucidate the impact of iSN contact on the transcriptome of SGs, we next compared gene

expression between co-cultured NMM treated glia and NMM treated glia alone. DGE analysis

yielded very few significant (p<0.05) differences, with only 20 significantly upregulated and 52

significantly downregulated genes in co-cultured NMM treated glia (Fig. 3G, Table S7). The

results indicate that iSN contact impacts the SG cell-cycle and morphology with upregulation of

genes related to DNA replication and repair (*CDC14A*[132], p adj.=0.0007), transcription (*ZSCAN18,*

p adj.= 0.0352; *TLE4* p adj.= 0.0393), translation (*QARS*[133], p adj.=8.165^{-10}), cytoskeletal

organization (*DNMBP*[134] p adj.= 0.047, *SRCIN1*[135] p adj.= 0.047)*, SLITRK2*[136] p adj.=0.0007) and

ion channels (*TRPM3* p adj.= 0.047*, GRIK2* p adj.= 0.008*, KCNK5* p adj.= 0.007). Several

upregulated genes were previously linked to specialized SGs (*SLITRK2*[86]*, TRPM3*[137]*, ITGB4*[90],

GBP2[90,137]), which may indicate changes in SG specific functional phenotype when in contact with

iSNs. Conversely, downregulated genes (Fig. 3G, Fig. S3F, Table S7-8) were associated with

collagen production (*P4HA3*[138] p adj.= 0.013, *COL4A5*[139] p adj.= 0.015) and cell-adhesion

(*LTBP2*[140] p adj.= 0.008), which may indicate an alteration in cell spreading and migration with

neuronal contact.

Taken together, transcriptomic analyses illustrate that initial DRG cell preparations contain

multiple cell types, however maturation media enriches for satellite glia. SGs in coculture and SGs

cultured alone with NMM treatment are transcriptionally very similar, however minor changes in

morphological and functional aspects of SGs do occur with neuronal contact.

Rodent SG (rSG) secreted factors alone do not induce a pseudounipolar morphology in

iSNs:

The transcriptomic analysis confirmed that rodent satellite glia (rSGs) contribute to the

maturation of iSNs, however the molecular mechanisms underlying this process are unknown.

We first asked whether this process is mediated by glial secreted factors, using a transwell co-

culture system (Fig. 4A). rSGs were cultured for one week on polycarbonate membranes inserted

into a well plate, and iSNs were then terminally differentiated in the same wells beneath the

membranes, sharing media but never physically touching rSGs, or in wells alone. iSNs were then

fixed and stained to visualize neuronal morphology (TUJ1, cytoskeletal marker, green; BRN3a,

iSN maturation marker, red) (Fig. 4B), while the rSGs on membranes were dyed and visualized

live after two weeks of culture to confirm that a healthy population was present. iSNs differentiated

in transwell coculture and alone both presented with predominantly multipolar or immature

morphologies, with no significant transition to pseudounipolar morphology (Fig. 4C, Table 3).

It is possible that glia only secrete factors critical for the maturation process after they have

made physical contact with iSNs. To test this possibility, iSNs plated alone were treated for six

days with spent media collected from cocultures (Fig. 4D-F, Table 4). While there are no

differences in pseudounipolar morphology, there is a higher frequency of immature neurons in

untreated iSNs (immature *adj. residual* = 3.544) compared to spent media treated iSNs (immature

adj. residual = -3.544), as well as an increased proportion of multipolar cells in spent media treated

iSNs (multipolar *adj. residual* = 2.954) compared to untreated iSNs (multipolar *adj. residual* = -

2.954). These results are consistent with the idea that rSG-iSN contact, rather than secreted

factors, are needed to induce pseudounipolar morphology.

	Morphology Frequencies								
	Immature		**Multipolar**	**Pseudounipolar Transition (PUT) + Pseudounipolar (PU)**					
Condition	**Unipolar**	**Bipolar**	**Multipolar**	**PUT-Bulge**	**PUT-Bell**	**PUT-AB**	**PU-Short**	**PU-Long**	**Totals**
Coculture	0	34	167	18	3	0	1	0	223
iSN Alone	1	42	176	8	6	1	1	1	236
Totals	77		343	39					459

335 **Table 3. Raw frequency of morphology subtypes observed in transwell co-cultured iSNs vs. iSNs terminally**
336 **differentiated alone. X^2 frequency statistic calculated for major morphology groups: Immature, Multipolar, and**
337 **PU(T). X^2= 1.562, df=2, p= 0.4579.**

	Morphology Frequencies								
	Immature		**Multipolar**	**Pseudounipolar Transition (PUT) + Pseudounipolar (PU)**					
Condition	**Unipolar**	**Bipolar**	**Multipolar**	**PUT-Bulge**	**PUT-Bell**	**PUT-AB**	**PU-Short**	**PU-Long**	**Totals**

Media Treated	0	10	225	10	2	0	0	0	247
Untreated	0	26	159	1	3	2	4	0	195
Totals		36	384		22				442

Table 4. Raw frequency of morphology subtypes observed with and without coculture spent media treatment. X^2 frequency statistic calculated for major morphology groups: Immature, Multipolar, and PU(T). X2= 12.69, df=2, p= 0.0018.

Physical contact between rSGs and hiSNs promotes pseudounipolarization:

To determine whether physical contact with satellite glia promotes pseudounipolarization, we designed an assay in which rDRG cells were cultured for two weeks in NMM, then fixed with methanol prior to coculture, preserving surface protein configurations. iSNs were subsequently cultured with the methanol fixed glial cells, in a normal coculture, or alone for one week (Fig. 5A). Cultures were then fixed and stained to visualize neuronal morphology (TUJ1, cytoskeletal marker, green) and contact with rSGs (p75, NGF receptor, magenta) (Fig. 5B). iSNs differentiated in methanol fixed coculture exhibit a phenotype intermediate between iSNs differentiated in normal coculture and those differentiated alone (Fig. 5C, Table 5). The Chi-square frequency test identified significant differences between observed and expected morphology frequencies across all three groups (X^2=131.3, df=6, p<0.001). Interestingly, these results are driven by significant differences between iSNs differentiated in normal coculture conditions (Immature *adj. residual* = -3.543, Multipolar *adj. residual*=-6.471, PUT *adj. residual*= 6.468, PU *adj. residual*=7.893) and iSNs differentiated alone (Immature *adj. residual* = 2.041, Multipolar *adj. residual* = 4.963, PUT *adj. residual*= -5.753, PU *adj. residual*= -3.893), while iSNs differentiated in a methanol fixed coculture exhibited "expected" morphology frequencies, with only significantly fewer pseudounipolar cells than expected (PU *adj. residual*= -4.007). Since PU(T) cells make-up 14.02% of iSNs differentiated in methanol fixed coculture, compared to the 2.26% of iSNs differentiated alone, these results suggest that physical contact between rSGs and iSNs contributes to the changes required for a transition to pseudounipolar morphology.

The degree to which such direct contact truly contributes to this process is unclear, as methanol fixation prevents movement of rSGs in culture, and therefore physical contacts between

rSGs and iSNs may be reduced compared to normal cocultures. Overall, the average number of

rSG contacts per iSN was significantly reduced in the methanol fixed condition compared to

normal cocultures (t=23.04, df=470, p<0.0001), with an average of 2 contacts/iSN in methanol

fixed coculture vs. 10 contacts/iSN in normal cocultures (Fig. S4A). Significantly more contacts

were made to true pseudounipolar cells in methanol fixed coculture compared to less advanced

morphologies (Bonferroni post-hoc: PU vs. Multipolar p=0.0038; Bipolar p=0.0053, PUT Bulge

p=0.0097) (Fig. S4B). In fact, pseudounipolar iSNs in methanol fixed co-culture were contacted

to a similar degree to those in normal coculture, with an average of 9 rSG contacts/ iSN in

methanol fixed coculture and 10 rSG contacts/iSN in normal coculture. These results indicate that

contact is essential to pseudounipolarization, but downstream signaling and movements of live

rSGs may also contribute to this process.

	Morphology Frequencies								
	Immature		Multipolar	Pseudounipolar Transition (PUT)			Pseudounipolar (PU)		
Condition	Unipolar	Bipolar	Multipolar	PUT-Bulge	PUT-Bell	PUT-AB	PU-Short	PU-Long	Totals
Methanol Fixed Coculture	2	37	189	35	2	1	4	0	270
Coculture	5	7	105	36	16	10	30	5	214
iSN Alone	1	28	144	4	0	0	0	0	177
Totals	80		438	104			39		661

Table 5: Raw frequencies of morphologies and morphology subtypes observed in methanol fixed coculture,
normal coculture, and alone. X^2 frequency statistic calculated for major morphology groups: Immature,
Multipolar, PUT and PU. X2=131.3, df=6, p<0.001.

Semaphorin-plexin and gap junction signaling play a role in pseudounipolarization of
iSNs:

We identified two potential pathways whose constituents are appropriately expressed to enable

contact mediated SG signaling to iSNs, namely, Sema6a-Plexin A4 and gap junctions. Sema6a,

a transmembrane semaphorin protein that is expressed in satellite glia, repels or inhibits axonal

growth in developing sensory neurons through engagement with the PlexinA4 receptor[141–144],

which is highly expressed in hiSNs (Fig. S4C). Additionally, our sc-RNA seq data set from

embryonic DRG satellite glia at the time of pseudounipolarization reveals high expression of

GJA1, encoding the gap junction protein Connexin-43[88]. *GJA1* was also found to be highly

expressed in all NMM treated rDRG glial cultures (Fig. S4D).

To test if these pathways promote a pseudounipolar morphology (Fig. 5D), we first knocked

down expression of *SEMA6A* in the rSGs using LV-shRNA transduction and puromycin selection.

Selected glia were allowed to repopulate and then hiSNs were plated for terminal differentiation

in coculture with shRNA-Sema6a expressing glia (shSema6a-rSGs) or shRNA-GFP expressing

glia (shGFP-rSGs, control). Cells were cocultured for one week and then fixed and stained to

visualize neuronal morphology (TUJ1, cytoskeletal marker, red) and glial expression of Sema6a

(Sema6a, magenta; GFP, green) (Fig. 5E). We found that the iSNs differentiated in coculture with

shSema6a-rSGs transition to pseudounipolar morphology at a reduced rate compared to controls

(Fig. 5F, Table 6), with significantly fewer PU(T) cells (PUT *adj. residual*=-4.772, PU *adj. residual*=-

6.861) compared to controls (PUT *adj. residual*=4.772, PU *adj. residual*=6.861), and significantly

more multipolar cells (*adj. residual*=9.39) compared to controls (Multipolar *adj. residual*=-9.39.

These results indicate that Sema6a-PlexinA4 signaling contributes to pseudounipolarization in

this coculture model and may also decrease the incidence of multipolar morphology by inhibiting

axonal outgrowth. However, the results also suggest that plexin signaling is not the sole

mechanism underlying iSN maturation, since PU(T) cells make up only 10.28% of iSNs

differentiated in shSema6a-rSG coculture.

	Morphology Frequencies								
	Immature		Multipolar	Pseudounipolar Transition (PUT)			Pseudounipolar (PU)		
Condition	Unipolar	Bipolar	Multipolar	PUT-Bulge	PUT-Bell	PUT-AB	PU-Short	PU-Long	Totals

	Unipolar	Bipolar	Multipolar	PUT-Bulge	PUT-Bell	PUT-AB	PU-Short	PU-Long	Totals
shRNA-Sema6a	2	9	242	15	7	2	4	1	282
ShRNA-GFP	20	8	181	38	29	14	53	16	359
Totals	39		423	105			74		641

Table 6. Raw frequency of morphologies observed in shRNA-Sema6a-SG coculture and shRNA-GFP-SG Coculture. X^2 frequency statistic calculated for major morphology groups: Immature, Multipolar, PUT and PU. $X^2=94.62$, df=3, p<0.001.

To assess the role of satellite glial gap junctions in iSN maturation, we knocked down *GJA1* in the rSGs and plated iSNs for terminal differentiation in coculture with shRNA-GJA1 expressing glia (shGJA1-rSGs) or shRNA-GFP expressing glia (shGFP-rSGs, control) for one week (Fig 5D). Cocultures of hiSNs with shRNA-GJA1 expressing glia (shGJA1-rSGs) or shRNA-GFP expressing glia (shGFP-rSGs, control) were stained to visualize neuronal morphology (TUJ1, cytoskeletal marker, red) and glial expression of GJA1 (GJA1, magenta; GFP, green) (Fig 5G). This experiment was carried out in parallel with the semaphorin knockdown, and therefore we utilized a single group of sh-GFP RNA control samples. Strikingly, iSNs differentiated in coculture with shGJA1-rSGs also transition to pseudounipolar morphology at a reduced rate compared to controls (Fig. 5F, Table 7) with fewer PU(T) cells than expected (PUT *adj. residual*=-5.537, PU *adj. residual*=-7.014), and significantly more multipolar cells than expected (*adj. residual*=10.109). These results suggest that both gap junction and Sema6a/PlexinA mediated signaling contribute to pseudounipolarization.

	Morphology Frequencies								
	Immature		Multipolar	Pseudounipolar Transition (PUT)			Pseudounipolar (PU)		
Condition	Unipolar	Bipolar	Multipolar	PUT-Bulge	PUT-Bell	PUT-AB	PU-Short	PU-Long	Totals
shRNA-GJA1	3	9	277	12	7	4	6	1	319
ShRNA-GFP	20	8	181	38	29	14	53	16	359
Totals	40		458	104			76		678

Table 7. Raw frequency of morphologies observed in shGJA1-rSG coculture and shGFP-rSG Coculture. X^2 frequency statistic calculated for major morphology groups: Immature, Multipolar, PUT and PU. $X^2=107.5$, df=3, p<0.001.

Cocultured iSNs provide a more physiologically relevant model of neuropathy:

iPSC derived neurons often exhibit embryonic phenotypes, which reduces the ability to model

complex molecular mechanisms and disease processes [9–15]. To determine whether cocultured

iSNs provide a physiologically relevant model of peripheral neuropathy compared to iSNs

differentiated alone, we examined the response of these cells to a common form of Chemotherapy

Induced Peripheral Neuropathy (CIPN), paclitaxel induced peripheral neuropathy[145,146]. Patients

with CIPN experience symptoms of allodynia, paresthesia, numbness, and pain in a stocking

glove distribution, due to the degeneration of sensory nerve axon terminals[147]. We assessed the

degeneration response of iSNs differentiated in coculture and alone when treated with low dose

paclitaxel.   At one week of terminal differentiation cultures were treated with paclitaxel

chemotherapy or DMSO for 48 hours, then fixed and stained to visualize the neuronal

cytoskeleton (TUJ1, Green) and gap junctions in both neurons and glia (GJA1, Magenta) (Fig.

6A,C). Neurodegeneration was estimated by quantifying TUJ1 signal.

We first tested the impact of low-dose paclitaxel on the degeneration of iSNs differentiated

with the Deng et al. 2023[15] protocol, generated by NCATS. Analysis of TUJ1 expression showed

that cocultured iSNs were significantly more susceptible to paclitaxel induced degeneration than

iSNs differentiated alone (Bonferroni post-hoc: Coculture + Paclitaxel treatment vs. Coculture +

DMSO: $p < 0.0001$; vs. iSN Alone + Paclitaxel: <0.0001; vs. iSN Alone + DMSO: $p<0.0001$), which

were not significantly impacted by treatment (Bonferroni post-hoc: iSN Alone + Paclitaxel vs. iSN

Alone + DMSO $p > 0.9999$) (Fig. 6B). We next tested the impact of the chemotherapy treatment

on degeneration of RealDRG[TM] iSNs from Anatomic, which require a larger dose of paclitaxel to

elicit axonal degeneration[148]. Again, iSNs differentiated in coculture were more susceptible to

paclitaxel induced degeneration compared to iSNs differentiated alone (Bonferroni post hoc:

Coculture + Paclitaxel treatment vs. Coculture + DMSO: $p < 0.0001$; vs. iSN Alone + Paclitaxel:

<0.0001; vs. iSN Alone + DMSO: $p<0.0001$), which were not significantly impacted by treatment

(iSN Alone + Paclitaxel vs. iSN Alone + DMSO $p > 0.9999$) (Fig. 6D). Together these data indicate

that iSNs cocultured with satellite glia provide a more physiologic model for toxin induced

neuropathies.

Discussion:

The use of differentiated iPSCs to model human diseases has revolutionized our ability to

understand pathology and develop efficacious treatments. While researchers have begun to

utilize induced sensory neurons to understand peripheral neuropathies, current differentiation

methods produce iSNs with an embryonic phenotype that do not fully recapitulate postnatal

functions. In this study, we developed a protocol that enables maturation of iSNs in coculture with

rodent satellite glia. We found that contact between iSNs and rSGs, mediated by semaphorin-

plexin and glial gap junction signaling, promotes an adult morphology and physiology in the

sensory neurons. Ultimately, this system also enables an enhanced model of neuropathy *in vitro*,

by increasing iSN susceptibility to chemotherapy induced axonal degeneration at physiologic

doses.

These studies establish a role for satellite glia in morphologic development of DRG neurons

and demonstrate that we can leverage this relationship to reliably produce pseudounipolar iSNs.

Previous studies have implicated DRG supporting cells in the morphologic maturation of sensory

neurons, with mixed primary cultures of DRG neurons and glia facilitating the transition of sensory

neurons to a pseudounipolar morphology[34–36,40]. This result has been recapitulated in induced

neuro-mesodermal assembloids[149], however iSNs cocultured with iPSC-derived[16] or primary [21]

Schwann cells alone do not exhibit advanced morphology. Based on these studies, we tested the

possibility that satellite glia primarily contribute to sensory neuron maturation. *In vivo,* DRG

sensory neurons increase physical contact with satellite glia during pseudounipolarization through

the development of perikaryal outgrowths that increase contact surface area[40] (Fig. S1B), In the

current study, we show that physical contact between glial cells and iSNs is essential to the

development of a pseudounipolar morphology, and RNA sequencing and immunohistochemistry

indicate that specifically satellite glia are responsible for this effect.

Though glial contribution to sensory neuron maturation have been recognized previously,

the molecular mechanisms enabling development of pseudounipolar morphology have been

unclear. Since our results ruled out glial secreted factors as a direct mechanism of

pseudounipolarization, we explored potential contact mediated mechanisms driven by our RNA

sequencing data. In iSNs, we observed high expression levels of *Plexina4*, a receptor that

interacts with glial semaphorins Sema3a and Sema6a[141–144,150], which have both been implicated

in sensory neuron axon development[141–144,151–153]. Since class 3 semaphorins are secreted, we

assessed the role of Sema6a, a transmembrane semaphorin[154], in the development of iSN axons.

We found knockdown of glial *Sema6a* significantly reduced pseudounipolar formation in

cocultured iSNs and instead resulted in a high proportion of multipolar cells. While there was a

significant reduction in the proportion of pseudounipolar cells following *Sema6a* knockdown in the

satellite glia, this effect was not complete. Factors contributing to observed low levels of

pseudounipolarization may include incomplete knockdown of *Sema6a*, compensation via

secreted Sema3a, and the possibility that semaphorin-plexin signaling is not the only mechanism

underlying this process.

In the DRG, satellite glia communicate with adjacent satellite glia and sensory neurons

through gap junctions[155,156], contributing to the maintenance of neuronal homeostasis through

direct exchange of diverse signaling molecules[157,158]. While little is known regarding gap junction

signaling in sensory neuron axon development, communication through gap junctions correlates

with neural crest cell differentiation into neuronal cells[159]. Knockdown of glial *GJA1,* like

knockdown of Sema6a, significantly reduced the ability of glial cells to induce a transition of iSNs

to a pseudounipolar morphology. As was seen following knockdown of Sema6a, knockdown of

GJA1 did not completely abolish pseudounipolarization. Again, this may be explained by

incomplete knockdown efficiency, compensation by an alternate connexin, or another parallel

signaling mechanism. Future studies need to be carried out to determine whether glial-glial or

glial neuronal gap junction signaling contributes to development of pseudounipolar morphology.

RNA sequencing also revealed potential intracellular mediators of pseudounipolarization.

Multiple molecules implicated in axonal guidance were significantly upregulated in cocultured

iSNs, as illustrated in Table 2. These include *CDKL5,* a kinase linked to axonal polarization[98] and

BCL11a, a transcription factor that suppresses growth of supernumerary neurites and regulates

microtubule protein expression[112]. We also observed significant downregulation of *RGMB* which

promotes formation of multiple neurites[160]. The identification of cues and molecular pathways that

stimulate pseudounipolar morphology may enable further refinements of iSN protocols.

Pseudounipolar morphology has multiple implications for sensory neuron physiology, as the

axon stem may act as an axon initial segment that potentiates spontaneous activity[36] or as a filter,

gating nociceptive action potentials[161,162]. Importantly, we demonstrate that terminal differentiation

in coculture yields more physiologically mature iSNs. Using MEA, we found that iSNs in coculture

become electrically active earlier than iSNs cultured alone, and fire at a rate more consistent with

rodent DRG neurons[54,55]. In addition to the development of an axon stem, the observed differential

expression of several ion channels and receptors may contribute to mature firing rates.

Upregulation of *SCN10a*, encoding Nav1.8, is of particular importance, as this sodium channel

maintains repetitive action potential firing in DRG nociceptors[163] and is integral to pain

signaling[72,73,164]. Increased expression of muscarinic receptors *CHRM2* and *CHRM3* may also

contribute to spontaneous activity, as acetylcholine has been implicated in the development of

bursting and spiking behavior in sympathetic neurons cocultured with satellite glia[165]. Lastly,

increased neuronal excitability may reflect a reduction in expression of several potassium

channels, namely *KNCT2*[166]   and *KCNS1*[167], which have previously been linked to

hyperexcitability and neuropathic pain.

In this study our goal was to generate iSNs with a more mature adult-like phenotype that
would allow us to better model peripheral neuropathies, and we tested the efficacy of modeling
toxic neuropathy using chemotherapy treated cocultured iSNs. We found that cocultured iSNs
were significantly more susceptible to paclitaxel chemotherapy mediated axonal degeneration
compared to iSNs alone. Hyperexcitability of sensory neurons has been shown to contribute to
axonal degeneration across multiple neuropathies[6], including chemotherapy induced peripheral
neuropathy (CIPN)[145,146], and so the increased baseline excitability of cocultured iSNs may
enhance susceptibility to degeneration. Satellite glia may also directly contribute to neuronal
degeneration, as studies in rodent models of CIPN indicate that glial release of inflammatory
cytokines contribute to degeneration[168–170]. Overall, our results provide new avenues for studying
the potential involvement of satellite glial-sensory neuron interactions in the pathogenesis of
neuropathies, as this protocol can easily be used to model diabetic peripheral neuropathy,
inherited neuropathies, or axonal injury as well.

In summary, our work identifies molecular direct contact cues whereby satellite glia support
sensory neuron development and establishes an accessible protocol for the reliable differentiation
of mature iSNs which can be used to study physiological and pathological changes in sensory
neurons and develop novel treatments for peripheral neuropathies.

Limitations of this Study:

This study is limited by the use of rodent glial cells and human derived sensory neurons. Currently,
there are no publicly available protocols to differentiate human satellite glia, however several
protocols have been developed to induce Schwann Cells[38,171] which share a glial progenitor cell
lineage. Adjustments of these protocols such that satellite glia differentiation can be optimized will
allow for an advanced iSN differentiation protocol with exclusive use of human cells, and aid in
identification of molecular mechanisms underlying normal human sensory neuron development
and neuropathy.

Methods:

Animal Care: Sprague-Dawley rats were purchased from Charles River Laboratory. Animals were cared for in accordance with NIH standards and guidelines. Experimentation was approved by the Dana Farber Cancer Institute IACUC.

iPSC Culture: Induced pluripotent stem cells (iPSCs) from LiPSC-GR1.1 (RRID:CVCL_RL65, Male, Hispanic, normal, <1D, umbilical cord blood, Lonza Inc. via NIH RMP, MCB p28) and NCRM5 cell lines (RRID: CVCL_1E75, Male, normal, <1D, umbilical cord blood, Lonza Inc. via NIH RMP, MCB p28) were cultured in feeder-free conditions in Essential-8 Medium (Thermo Fisher; A1517001) using cell culture plates or flasks coated with vitronectin (Thermo Fisher; A14700). iPSCs were passaged with EDTA (Thermo Fisher; 15575020) every 3 days at a 1:6 ratio. Cells were treated with the CEPT cocktail for 24 hours after passaging (Chroman 1, MedChem; HY-15392; Emricasan, SelleckChem S7775, Polyamine supplement (1000X), Sigma-Aldrich P8483, Trans-ISRIB, R&D Systems 5284)[172]. Basic characterization including, authentication using G-banded karyotyping, and Mycoplasma detection using Mycoalert kit (Lonza; LT07-318) were carried out at p28. Pluripotency was assessed at p30 using the TaqMan hPSC Scorecard assay (Thermo Fisher; A15876) and analysis software, and significance was determined using the *Student's t-test*. iPSCs (p30) were differentiated into neural crest precursors (NCPs) and immature sensory neurons (SNs) using our published protocol[15].

rDRG-iSN Coculture:

Rodent DRG glial cell culture: Sprauge-Dawley day E14.5 rat DRGs were dissected. DRGs were trypsinized (1mg trypsin /mL HBSS) for 50 minutes at 37°C. 40k cells/well were plated on uncoated Lab-Tek™ II Chambered Coverglass (Thermofisher cat#: 155409) with filtered Glia Base Media (DMEM/F12, 10% FBS, 1% Pen. Strep., 0.08% glucose). After a 30-minute incubation at 37°C 7.5% CO_2., chambers were vigorously shaken, and media was fully changed

to remove cells in suspension. Remaining glial cells were cultured in Glia Base Media at 37°C

7.5% CO_2 for one week with media changes 3x/ week.

Terminal differentiation and iSN- rDRG glia co-culture: Frozen Neural Precursor Cells (NPCs)

were differentiated by NIH-NCATS Stem Cell Laboratory using the protocol outlined in Deng et

al., 2023[15]. Lab-Tek™ II Chambered Coverglass (Thermofisher cat#: 155409) were coated with

0.005% poly-ornithine solution (Sigma-Alrich cat#: P4957) in dH2O 1 hr at 37°C, washed, and

then coated with 5 ug/mL mouse laminin (Thermofisher cat#: 23017015) in dH2O 1 hr at 37°C.

NPCs were thawed in NCATS Maturation Media (DMEM/F12+Glutamax, 2% B27,1% N2, 25ng

rh- BDNF/b-NGF/GDNF/NT3) + CEPT rock inhibitor cocktail (50nM Chroman-1 (Medchem

Express Cat#: HY-15392), 5uM Emricasan (Selleck cat#: S7775), 0.001% Polyamine supplement

(Sigma-Alrich cat#: P8483), 0.7uM Trans-ISRIB (Tocris cat#: 5284)). 15k cells/well were plated in

NCATS Maturation Media + CEPT on coated cultureware for terminal differentiation alone (iSN-

Only), or in coculture with rDRG glial cultures treated with NCATS Maturation Media + CEPT. 24

hrs after plating, media was changed to NCATS Maturation Media + 1uM PD0332991 (Tocris cat#:

4786). Cells were cultured for one week at 37°C 5% CO_2., with ½ media changes 3x/week.

For each experiment throughout this study, we indicate the number of biological replicates used,

with each biological sample representing an experiment done with one set of iSNs that were

plated and tested individually. The experiment was subsequently replicated on additional sets of

cells plated and tested on a different day (the biological replicates). n=4 biological replicates for

experiments performed in Figure 1C, n= 2 NCRM5 and n=2 GR1.1.

Quantification of iSN morphology frequency:

Immunofluorescence: Cells were fixed for downstream Immunofluorescence (IF) on Day 14. First,

cells were fixed with 2% paraformaldehyde (PFA) (EMS cat#: 15710; 1:1 4% PFA to spent media)

10 minutes at room temperature (RT), and then in 4% PFA 20 minutes at RT. Cells were washed

3x in PBS and stored at 4°C. Fixed cultures were permeabilized with 0.1% Triton-x in PBS 10

minutes and blocked in blocking buffer (5% Normal Goat Serum (Cell Signaling Tech. cat#:

5425S) in 1x PBS) for 1 hr at RT. Cultures were then incubated overnight at 4°C in primary

antibodies (Table 11) diluted in blocking buffer. The following day, cultures were washed and

incubated in Alexa-Fluor secondary antibodies (Table 12) and DAPI diluted 1:1000 in blocking

buffer for 1 hr at RT. Co-Cultures were stained with mouse anti-human nuclei-CY3 conjugate

(Sigma cat#: MAB1281C) for 2 hrs at RT to distinguish human neuronal nuclei from rodent. Cells

cultured in Lab-Tek™ II Chambered Coverglass were stored at 4°C in PBS prior to imaging, and

cells cultured on coverslips were mounted on slides with fluoromount (Thermofisher cat#:

OB10001).

Imaging: All photomicrographs of cells cultured in Lab-Tek™ II Chambered Coverglass were

collected on the Zeiss-980 airy-scan/confocal microscope. Images of all conditions were obtained

at 20x with 1.3x zoom and bidirectional scanning in SRY-4 Airy-Scan. Multiplex imaging was

utilized with 405, 488, 561, and 639 lasers at 800V gain, 1% laser power, and 0 offset. All images

were obtained in a z-stack with 0.35um step size. Systematic random sampling was utilized to

image n=50-100 cells/biological replicate across 8-wells. Areas with fused cell bodies and axonal

fasciculations were omitted from sampling. All images were 3D airy-scan post processed in Zen

Blue software.

Analysis: All human cells in each photomicrograph were categorized into the following

morphologic groups : unipolar (one axonal projection off of the cell body, that does not bifurcate),

bipolar (two axonal projections off of opposite cell body poles, ~180° angle), multipolar (more than

two axonal projections), Pseudounipolar Transition (PUT)- Bulge (two axonal projections off of the

bottom of the cell body, axons form >90° angle), PUT-Bell (two axonal projections off of the bottom

of the cell body, axons form a 90° angle or less), PU-Associated Branch (PU-AB,T-shaped

pseudounipolar axon with a branch <10uM long), PU-Short (T-shaped pseudounipolar axon, with

an axon stem <60um long), PU-Long (T-shaped pseudounipolar axon, with an axon stem >60um

long). Human cells were distinguished from rodent cells based on expression of human nuclei-

Cy3 marker. Images were blinded and intra-rater or inter-rater reliability was calculated to be

>90%. Morphology frequency was compared across morphologic groups: Immature (unipolar

and bipolar), Multipolar, Pseudounipolar Transition (PUT-Bulge, PUT-Bell, PUT-AB), and

Pseudounipolar (PU-Short and PU-Long) using the Chi-square test in Graphpad Prism 10. Sub-

morphologies were pooled into major morphologic categories to reach chi-square test validity

requirements. Adjusted residuals with Bonferroni correction were calculated in excel using

standard formulas.

Multi-Electrode Array Recording:

Cell Culture: rDRG glia were dissected and plated in 48-well, 16 electrode Cytoview-MEA plates

(Axion) at 20k cells/well in Glia Media. After a 30-minute incubation at 37°C 7.5% CO_2, plates

were vigorously shaken, and media was fully changed to remove cells in suspension. Remaining

glial cells were cultured in Glia Media at 37°C 7.5% CO_2 for one week with media changes 3x/

week. On the 7th day, iNCPs were thawed and plated in NCATS Maturation Media + CEPT at

150k-200k cells/ well, in coculture or alone on coated wells (100ug/mL PDL 1 hr 37°C, 10ug/mL

mouse laminin 1 hr 37°C). A fraction of the wells remained glial cells alone as a control, treated

with NCATS Maturation Media +CEPT. 24 hrs after plating, media was changed to NCATS

Maturation Media + 1uM PD0332991. Cells were cultured for 5-8 weeks at 37°C 5% CO_2, with ½

media changes 3x/week. n=4 biological replicates; replicates 1-3 = iPSC Gr1.1, replicate 4 = iPSC

NCRM5.

MEA Recording: MEA recordings were performed on the Maestro Pro recording system from

Axion at 37°C 5% CO_2, 1x/week for 5 minutes, after at least 5 minutes of acclimation. AxIS

navigator software was used to record standard Spontaneous Neural and Viability properties.

Burst Detection was enabled with the following specifications: Single Electrode Minimum #

Spikes=5, Network Minimum #Spikes=50, Maximum Interspike Interval= 100s, Mean Firing Rate

detection window= 100s. Recordings were carried out weekly, and data was analyzed when an

average of 7+ active electrodes was reached for both cocultured iSNs and iSNs alone. Electrodes

were considered active with a minimum of 5 spikes/minute.

Analysis: Data was analyzed at the time point when an average of 7+ electrodes were active in

both conditions. Wells with 2 or less active electrodes were excluded from analysis. Glial cells

were not active, and therefore were excluded from all analyses aside from # of active electrodes.

Data did not follow a normal gaussian distribution, and therefore averages were compared using

the Mann-Whitney U test in Graphpad Prism 10. n=# wells, pooled from 4 biological replicates.

RNA sequencing of iSN and rDRG Glia cocultures:

Cell Culture and RNA Collection: rDRG glia were dissected and plated in uncoated glass bottom

35mm Fluorodishes (WPI cat #: FD35-100) at 250k cells/dish in Glia Media. After 30 minutes at

37°C 7.5% CO_2, dishes were vigorously shaken, and media was fully changed to remove cells in

suspension. Remaining glial cells were cultured in Glia Media at 37°C 7.5% CO_2 for one week

with media changes 3x/ week. On Day 7, cells were either treated with NCATS Maturation Media

+ 1uM PD0332991, maintained in Glia Media, or co-cultured with iSNs as described above. iSNs

were plated at 90k/dish in co-culture or alone, as described above. Cultures were maintained at

37°C 5% $CO_{2.}$, with ½ media changes 3x/week for an additional week. On Day 14, RNA was

collected from Glia Media Treated rDRG Glia, NCATS Maturation Media Treated rDRG Glia, iSN-

rDRG Glia Co-Culture, and iSN-Only mass cultures using the RNeasy Plus Mini Kit (Qiagen cat#:

74134). n=3-4 biological replicates; Coculture: n= 2 iPSC NCRM5, n=1 GR1.1. iSN only: n=4

iPSC NCRM5. Glial cells derived from n=4 rats.

RNA sequencing: mRNA bulk sequencing was carried out at the Dana Farber Cancer Institute

Molecular Biology Core Facility. The genomics core team performed RNA QC, rRNA depletion,

library preparation (Roche Kapa mRNA Kit), and sequencing on the Illumina NovaSeq 6000.

Reads were aligned to both the rat (*Rattus norvegicus*) and human genome using STAR aligner

to separate rDRG glia and iSN reads in co-culture. GEO Accession #: GSE272868.

Analysis: Normalized counts (with raw counts <3 set to NA) were analyzed for differential gene

expression using the DESeq2 R package version 1.42.0[173] with an adjusted p-value cutoff of 0.05.

Gene Ontology Enrichment Analysis was performed using the clusterProfiler R package version

4.10.0[174] with an adjusted p-value cutoff of 0.05.

iSN-rSG co-culture in transwell compartmentalized system:

Cell Culture: rDRG glia were dissected as described above and plated at 40k cells/well on

uncoated Millicell® Standing Cell Culture Inserts (MilliporeSigma cat#: PIHP01250). rDRG glia

were prepared as described above. On the 7th day, iNPCs were plated at 15k cells/ well on sterile

12mm #1.5 coverslips, coated as described above, in the same plate for terminal differentiation

either in compartmentalized co-culture or alone. Cells were allowed to attach to coverglass for 30

min at 37°C 5% CO_2, after which Glia Media was removed from rDRG glial inserts and inserts

were transferred to wells with NPCs dedicated for terminal differentiation in compartmentalized

co-culture. 24 hrs after iNPC plating, media was changed to NCATS Maturation Media + 1uM

PD0332991. Cells were cultured for one week at 37°C 5% CO_2., with ½ media changes 3x/week.

On the 14th day, rDRG glial culture inserts were moved to new wells and glia attached to insert

were visualized with CellTracker Red CMTPX (Thermofisher cat#: C34552). Cells were incubated

with 12.5uM dye in DMEM/F12 for 30 min at 37deg 5% CO2, after which media was changed to

DMEM/F12 and live cells were imaged at 10x TRITC on the Nikon-Ti E microscope. iSNs adherent

to coverglass were fixed for downstream Immunofluorescence (IF) as described above. n=3

biological replicates, iPSC Gr1.1.

Imaging and Analysis: Photomicrographs of cells cultured on coverslips were collected on the

Nikon Ni-E confocal microscope at 60x with oil immersion. Images were obtained with 405 and

488 lasers, at 30V gain 5% power and 561 30V gain 7% power. All images were obtained in a z-

stack with 0.35um step size. Systematic random sampling was utilized to image n=50-100

cells/biological replicate across 3 coverslips. Areas with fused cell bodies and axonal

fasciculations were omitted from sampling. Inter-rater reliability was calculated >90%. Morphology

frequencies and statistics were carried out as described above.

iSN cultures treated with co-culture spent media:

Spent media was collected from iSN-rDRG Glia Co-Cultures over the course of one week and

stored at 4°C. On Day 21, iNPCs were plated alone on chambered coverglass in NCATS

Maturation Media + CEPT as described above. After 24 hrs, media was either changed to ½ fresh

NCATS Maturation Media + 1uM PD0332991 + ½ spent media or maintained in NCATS

maturation Media + 1uM PD0332991 as described above. Cells were cultured for one week with

3x full media changes maintaining specified ratios. On Day 28, cultures were fixed for IF and

images were acquired and analyzed as described. n=3 biological replicates; n= 2 iPSC NCRM5

n= 1 iPSC Gr1.1.

iSN-rSG co-culture with glial methanol fixation:

rDRG glia were dissected, plated, prepared, and cultured in chambered coverglass for one week

as described above. On Day 7, glia were treated with NCATS maturation media + CEPT, and 24

hrs later media was changed to NCATS Maturation Media + 1uM PD0332991. Glia were

maintained in NCATS Maturation Media + 1uM PD0332991 an additional week with ½ media

changes 3x/week. On day 14, cells were fixed in 95% methanol 7 min at 4°C, washed, and treated

with NCATS maturation media + CEPT for downstream iSN co-culture. iNPCs were thawed and

plated at 15k cells/well for terminal differentiation in iSN-rDRG Glial Co-Culture with methanol

fixation, iSN-rDRG Glial Co-Culture without methanol fixation (prepared as described above), or

iSN-only culture (prepared as described above). 24 hrs after iNPC plating, media was changed

to NCATS Maturation Media + 1uM PD0332991. Cells were cultured for one week at 37°C 5%

$CO_{2.}$, with ½ media changes 3x/week. After one week of coculture, cells were fixed with PFA for

downstream IF and images and analyses were carried out as described above. n=4 biological

replicates, n=2 NCRM5, n=2 GR1.1.

iSN-rSG co-culture with glial *Sema6a* or *GJA1* sh-RNA knockdown:

rDRG glia were dissected, plated, prepared, and cultured for five days as described above. On

Day 5, Glia were transduced with ShRNA-Sema6a-puro-GFPtag lentivirus (Millipore Sigma Cat#:

SHCLNV), ShRNA-GJA1-puro-GFPtag lentivirus (Millipore Sigma Cat#: SHCLNV), or ShRNA-

GFP-puro control lentivirus (Millipore Sigma cat#: SHC005V) in Glia Media for 24 hrs, after which

media was changed. Cells were maintained in Glia Media 72 hrs post-transduction and evaluated

for sh-RNA expression using live-imaging in brightfield and FITC on the Nikon-Ti E microscope.

Glia were then treated on Day 8 with 1ug/mL puromycin in Glia Media for 48hrs. Media was

changed, and selected cells were allowed to proliferate for four days. On Day 14, iNPCs were

plated at 15k cells/well in NCATS Maturation Media + CEPT for terminal differentiation in iSN-

rDRG Glia Knockdown Co-Culture, iSN rDRG Glia Knockdown Control Co-culturel and iSN-rDRG

Glia Co-Culture Control (prepared as described above). 24 hrs after iNPC plating, media was

changed to NCATS Maturation Media + 1uM PD0332991, and cells were co-cultured for one week

at 37°C 5% $CO_{2.}$, with ½ media changes 3x/week. On Day 21, co-cultures were fixed for IF as

described. n=4 biological replicates, 2/ iPSC line.

Modeling chemotherapy induced axonal degeneration:

Chemotherapy treatment of iSN co-culture with rDRG glia: rDRG glia were prepared as described

above. iSNs generated with the Deng et al. 2023[15] protocol, were plated at 15k cells/well in rDRG

glial co-culture or alone and maintained in culture for 7 days as described above. On day 14, cells

were treated with 60nM Paclitaxel (Sigma Aldrich cat #: T7402) or 0.2% DMSO (Sigma Aldrich

cat#: D2650) in NCATS Maturation Media + 1uM PD0332991 for 48 hrs. n=4, iPSC NCRM5.

Anatomic RealDRG iSNs were plated at 5k cells/well in rDRG glial co-culture or alone and

maintained in culture for 7 days as described above. On day 14, cells were treated with 600nM

Paclitaxel (Sigma Aldrich cat #: T7402) or 0.2% DMSO (Sigma Aldrich cat#: D2650) in NCATS

Maturation Media + 1uM PD0332991 for 48 hrs. n=4, Anatomic RealDRGs are derived from one

genetic iPSC line. Cells were then fixed as described above for IF.

Immunofluorescence (IF): Fixed cultures were permeabilized with 0.1% Triton-x in PBS 10

minutes, and blocked in blocking buffer (5% Normal Donkey Serum (Sigma Aldrich cat#: D9663)

in PBS) for 1 hr at RT. Next, they were incubated overnight at 4°C in primary antibodies (Table

11) diluted in blocking buffer. The following day, cultures were washed and incubated in Alexa-

Fluor secondary antibodies (Table 12) and DAPI diluted 1:1000 in blocking buffer for 1 hr at RT.

Cells were washed and stored in PBS at 4°C prior to imaging.

Imaging: All photomicrographs of cells cultured in Lab-Tek™ II Chambered Coverglass were

collected on the Zeiss-980 airy-scan/confocal microscope. Images of all conditions were obtained

at 20x with 1.3x zoom and bidirectional scanning in SRY-4 Airy-Scan configuration. Multiplex

imaging was utilized with 405, 488, 561, lasers at 800V gain, 1% laser power, and 0 offset. All

images were obtained in a z-stack with 0.35um step size. Systematic random sampling was

utilized to image n=16 fields/biological replicate across 6-wells. All images were 3D airy-scan post

processed in Zen Blue software.

Analysis: Particle analysis was carried out on gray-scale TUJ1 channel image stacks. Threshold

was manually set at the brightest z-stack slice for each image, and percent area occupied by

TUJ1 pixels was quantified for each image stack. Average values were calculated for each field.

Field averages were compared between treatment groups and culture conditions using two-way

ANOVA analysis with the Bonferroni post-hoc test in Graphpad Prism. n=64 fields from 4 biological

replicates.

Primary Antibody	Supplier:Cat #	Concentration	Experiment(s)
rbt anti-BRN3a	Abcam: ab245230	1:250	Co-culture in compartmentalized system, Spent media treatment
rbt anti-dsRed	Takara:632496	1:1000	*in vivo* DRG morphology
gt anti-GJA1	Abcam: ab87645	1:200	Connexin-43/GJA1 knockdown co-culture, Chemotherapy treatment co-culture
rbt anti-Fabp7	ThermoFisher: PA531864	1:200	Identification of glial subtypes
Mse anti-human nuclei-CY3	Millipore Sigma: MAB1281C	1:200	Co-culture, RealDRG co-culture, Methanol fixation co-culture
rbt anti-MBP	Abcam: ab218011	1:250	Identification of glial subtypes
rbt anti-P75/ngfr	Abcam: ab52987	1:100/ 1:200	Co-culture, RealDRG co-culture, Methanol fixation co-culture, Identification of glial subtypes
rbt anti-OCT6	Abcam: ab259952	1:200	Identification of glial subtypes
Mouse anti-S100b	Sigma: S2532	1:1000	Identification of glial subtypes
rbt anti-Sema6a	Thermofisher: PA581009	1:100	Sema6a knockdown co-culture
ms anti-TUJ1	Biolegend: 801202	1:1000	Co-culture, RealDRG co-culture, Co-culture in compartmentalized system, Methanol fixation co-culture, Spent media treatment, Sema6a knockdown co-culture Connexin-43/GJA1 knockdown co-culture, Chemotherapy treatment co-culture

769   **Table 11. Primary antibodies.**

Secondary	Supplier:Cat #	Concentration	Experiment(s)

Antibody			
gt anti-Ms Alexa 488	Thermofisher: A11001	1:1000	Co-Culture, RealDRG co-culture, Methanol Fixation co-culture, Co-culture in compartmentalized system, Spent media treatment, Identification of Glial Subtypes
gt anti Rbt Alexa 647	Thermofisher: A21245	1:1000	Co-Culture, RealDRG co-culture, Methanol Fixation co-culture
gt anti-Rbt Alexa 546	Thermofisher: A11035	1:1000	Co-culture in compartmentalized system, Spent media treatment, Identification of Glial Subtypes
dnky anti-Ms Alexa 568	Thermofisher: A10037	1:1000	Sema6a Knockdown coculture, Connexin-43/GJA1 knockdown coculture, CIPN chemotherapy treatment coculture
dnky anti-Gt Alexa 647	Thermofisher: A21447	1:1000	Connexin-43/GJA1 knockdown coculture, Chemotherapy treatment coculture
dnky anti-Rbt Alexa 647	Thermofisher: A31573	1:1000	Sema6a knockdown coculture
dnky anti-Rbt Alexa 546	Thermofisher: A10040	1:1000	*in vivo* DRG morphology

Table 12. Secondary antibodies.

<u>Figure Legends:</u>

Figure 1: Human iSNs differentiated in coculture with embryonic rodent DRG glia exhibit
more mature morphology and physiology. A) Schematic illustrating protocol to terminally
differentiate iSNs in coculture with embryonic rat DRG glia (rDRG). Briefly, DRGs are dissected
from E14.5 rat embryos. After 30 minutes, cultureware is shaken and cells in suspension are
removed. Glia are cultured for 1 week, after which iNCPs are thawed and plated in coculture or
alone for 1 week of maturation. B) Photomicrographs representing distinct iSN morphologies
observed. TUJ1= Green, p75= Magenta, Human Nuclei= Red, DAPI= Cyan. Scale=50um. PU=
pseudounipolar. PU-AB: white arrowhead indicates accessory branch. PU-Long: white arrowhead
indicates the PU-Long neuron in this field. C) Frequency of each morphology observed in
coculture and alone, displayed as percentages. 4 biological replicates, 2 iPSC cell lines. D.
Representative raster plots over 2.5 minutes of MEA recording. Black lines indicate individual

spikes and blue lines indicate bursts. E) Average number of active electrodes when average was
> 7 (5-7 weeks). Coculture n= 28 wells, iSN alone n=17 wells, Glia alone n=16 wells from 4
biological replicates. F-H) Number of spikes (F), mean firing rate (G), and number of bursts (H)
when average # active electrodes were >7. Wells with <2 active electrodes were excluded.
Coculture n= 26 wells, iSN alone n=15 wells from 4 biological replicates. I) Average burst duration
in seconds when average # active electrodes were >7. Wells with <2 active electrodes and 0
bursts were excluded. Coculture n=24 wells, iSN alone n=14 wells from 4 biological replicates. J)
Interburst Interval in seconds when average # active electrodes were >7. Wells with <2 active
electrodes and <2 bursts were excluded. Coculture n= 22 wells, iSN alone n=13 wells from 4
biological replicates. F-J) Mann-Whitney U tests performed in Graphpad Prism 10,
$p < 0.05$=significant, $p < 0.0001$=****.

Figure 2. iNCPs terminally differentiated in co-culture more advanced in neuronal
differentiation compared to iNCPs alone. A) Volcano plot illustrating differential gene
expression in coculture vs. alone. B) GO analysis terms upregulated in coculture vs. iSNs alone,
related to neuronal development. C) GO analysis terms downregulated in coculture vs. iSNs
alone, related to neuronal development. Coculture n=3, iSN alone n=4. P adj.<0.05.

Figure 3. Rodent DRG satellite glia (rSGs) are responsible for advanced maturation of iSNs.
A) Volcano plot illustrating differential gene expression in NCATS Maturation Media treated rDRG
glia vs. rDRG glia maintained in base glia media B-C) GO analysis terms upregulated (B) and
downregulated (C) in NCATS Maturation Media treated rDRG glia vs. rDRG glia maintained in
base glia media. D) Volcano plot illustrating differential gene expression in cocultured rDRG glia
vs. rDRG glia maintained in base glia media E-F) GO analysis terms upregulated (E) and
downregulated (F) in cocultured rDRG glia vs. rDRG glia maintained in base glia media. G)
Volcano plot illustrating differential gene expression in cocultured rDRG glia vs. NCATS
Maturation Media treated rDRG glia. n=3-4/group. P adj.<0.05.

Figure 4. rSG secreted factors do not induce pseudounipolar morphology in iSNs. A)
Schematic illustrating transwell coculture protocol. B) Representative photomicrographs of iSNs
cultured in transwell coculture or alone. TUJ1= Green, BRN3a=Red, DAPI= Blue. Scale= 50um.
C) Frequency of each morphology observed in coculture and alone, displayed as percentages .n=
3 biological replicates D) Schematic illustrating protocol for iSN treatment with coculture spent
media. E) Representative photomicrographs of iSNs treated with coculture spent media or
untreated. TUJ1= Green, BRN3a=Red, DAPI= Blue. Scale=50um. F) Frequency of each
morphology observed in Coculture Media treated and untreated iSNs, displayed as percentages.
3 biological replicates.

Figure 5. Physical contact between rSGs and iSNs is essential to pseudounipolarization.
A) Schematic illustrating protocol for iSN coculture with methanol fixed glia. B) Representative
photomicrographs of iSNs in coculture with methanol fixed glia, normal glia, or alone. TUJ1=
Green, p75=Magenta, Human Nuclei= Red, DAPI= Cyan. iSNs alone were not stained for human
nuclei. Scale= 50um. C) Frequency of each morphology observed in cocultured iSNs with
methanol fixed glia and normal glia, displayed as percentages. n=4 biological replicates. D)
Schematic illustrating protocol for iSN coculture with Sema6a, GJA1, and GFP knockdown in glia.
All three conditions were run in parallel. E) Representative photomicrographs of iSNs in coculture
with SEMA6a knockdown glia or GFP knockdown glia. Sema6a=Magenta, TUJ1= Red,
GFP=Green, DAPI=Cyan. Scale=50um. F) Frequency of each morphology observed in iSNs
cocultured with SEMA6a knockdown, GFP knockdown, and GJA1 Knockdown glia, displayed as
percentages. n=4 biological replicates. G) Representative photomicrographs of iSNs in coculture
with GJA1 knockdown glia or GFP knockdown glia. GJA1=Magenta, TUJ1= Red, GFP=Green,
DAPI=Cyan. Scale=50um.

Figure 6. Cocultured iSNs are more susceptible to paclitaxel mediated neurodegeneration.
A) Representative photomicrographs of Deng et al. 2023 generated iSNs in coculture or alone,
treated with either 0.2% DMSO or 60nM paclitaxel. TUJ1= Green, GJA1=Magenta, Human
Nuclei= Red, DAPI= Cyan. iSNs alone were not stained for human nuclei. Scale= 50um. B)
Percent area occupied by TUJ1 particles in cocultured iSNs and iSNs differentiated alone + 0.2%
DMSO or 60nM Paclitaxel. Two-Way ANOVA with Bonferroni post-hoc calculated in Graphpad
Prism 10, p<0.05= significant, p<0.0001=****. n=64 fields from 4 biological replicates. C)
Representative photomicrographs of Anatomic RealDRG iSNs in coculture or alone, treated with
either 0.2% DMSO or 600nM paclitaxel. TUJ1= Green, GJA1=Magenta, Human Nuclei= Red,
DAPI= Cyan. iSNs alone were not stained for human nuclei. Scale= 50um. D) Percent area
occupied by TUJ1 particles in cocultured iSNs and iSNs differentiated alone + 0.2% DMSO or
600nM Paclitaxel. Two-Way ANOVA with Bonferroni post-hoc calculated in Graphpad Prism 10,
p<0.05= significant, p<0.0001=****. n=64 fields from 4 biological replicates.

Acknowledgements: NCI R01 CA205255. Schematics created with Biorender.

Declaration of Interests: C.J.W. is a founder of Nocion, Quralis, and Blackbox Bio, and is a
member of the SAB of Lundbeck Pharma and Tafalgie Therapeutics. I.S. is currently CSO at
FUJIFILM Cellular Dynamics. All other authors have no interests to disclose.

Author Contributions: Conceptualization: CJL, OYT, RAS. Generation of NCPs: KG, VJ, DC,
PO, CT, IS. Performance of experiments: CJL, MFPM, ES, SD, VP, SZ, OYT. Analysis: CJL,
MFPM, MT, TM. Funding Acquisition: RAS. Project Administration: CJL, RAS. Supervision: RAS,
CW. Writing: CJL, RAS, with input from all authors.

Supplemental Methods:

Animal Care:

Sprague-Dawley rats were purchased from Charles River Laboratory. Ntrk2-CreER/C57BL/6 mice
and lox-tdtomato/C57BL/6 mice were shared by Dr. David Ginty, Harvard Medical School. PLP1-
GFP/C57BL/6 mice were shared by Dr. Lisa Goodrich, Harvard Medical School. Animals were

cared for in accordance with NIH standards and guidelines. Experimentation was approved by
the Dana Farber Cancer Institute IACUC.

Tamoxifen induced sparse labeling of Ntrk2+ sensory neurons with tdtomato *in vivo*:

Ntrk-CrER mice were crossed with lox-tdtomato mice and subsequently crossed with PLP1-
GFP/C57BL/6 mice. Timed pregnant Ntrk-CreER/lox-tdtomato/PLP-1GFP mice were weighed
and tamoxifen (TMXF) dose was calculated at 40ug/g body weight. 2mg/mL Tamoxifen (TMXF)
was stored in methanol at -80C and the final TMXF dose was diluted in 100uL corn oil. Diluted
TMXF was vortexed 45 minutes and dried in a Speedvac to remove methanol from the mixture,
leaving TMXF in corn-oil alone. Injections were carried out intraperitoneally at E12.5-E14.5 to
sustain expression and sacrifice at E14.5, E16.5, E18.5, P5. At time of sacrifice, embryonic
lumbar DRGs were dissected, and expression of tdtomato and PLP1-GFP was evaluated via
epifluorescence. DRGs were then drop fixed in 4% paraformaldehyde at 4°C overnight. The next
day, DRGs were washed 3x with 1XPBS and stored at 4°C in 1X PBS until staining. n=3+ embryos
total per age group.

Quantification of DRG sensory neuron morphology frequency *in vivo*:

Immunofluorescence: Fixed DRGs were permeabilized 2 hrs in 0.5% tritonX in PBS, and then
blocked in 5% Normal Donkey Serum (NDS)/0.5% tritonX in PBS 4-5hrs at RT. They were then
incubated in rbt anti-dsRed (Takara Cat#: 632496) primary antibody diluted 1:1000 in blocking
buffer at 4°C overnight, and washed the next day with 1X PBS. Next, DRGs were incubated with
dnky anti-rbt 546 (Thermofisher Cat#: A10040) secondary antibody diluted 1:500 in blocking
buffer for 3 hrs at RT. After staining, DRGs were stored at 4°C in 1XPBS, or gradually transferred
to methanol for tissue clearing.

DRG Tissue Clearing: Stained DRGs stored in PBS were gradually transferred to methanol with
10 minute incubations in the following concentrations: 25%, 50%, 75%, and 100%. After methanol
dehydration, each DRG was transferred to a slide and placed within a slide spacer (VWR
Cat#102096-614). BABB solution (1 Benzyl Alcohol:2 Benzyl Benzoate) was added to the DRG
and incubated 2-3min, after which it was replaced with fresh BABB, and steps were repeated until
the tissue was cleared. Then, a circular coverslip was mounted on top with tissue embedded in
BABB solution.

Imaging: Confocal images were acquired on the Nikon Ni-E confocal microscope, first at 20X to
visualize the entire DRG, and then at 40x to visualize individually labeled neurons. Images were
acquired at x resolution, 2.4 speed, 0.2-0.3um step, with 488 and 561 laser at 1.04 laser power,
36 gain, 0 offset. All cells labeled with tdtomato in the DRG were imaged from the top to bottom
of neuronal cell bodies.

Analysis: Images were blinded after acquisition, and NIS elements software was used to visualize
all cell morphology in 2D and/or 3D. All labeled cells were categorized into the following
morphologic groups: bipolar (elongated cell body, with an axon emanating from the two opposite
poles of the cell body with ~180° angle between), intermediate (round cell body with two axons
forming <180° angle between), and pseudounipolar (one axon stem emanating from cell body,
forming a T-Shape axon). Cells were considered apoptotic with condensed cell bodies small, and
no axons attached.

Anatomic RealDRG iSN co-culture with rDRG glia: Frozen RealDRG iSNs were purchased
from Anatomic (cat#: 1020). rDRG glia were dissected, plated, prepared, and cultured for one
week as described above. On Day 7, RealDRG iSNs were thawed in Chrono-Senso-MM
(Anatomic) and plated at 15k, 9k, or 5k cells/well either in rDRG glia co-culture in NCATS
Maturation Media. Lab-Tek™ II Chambered Coverglass were coated with 0.01% Poly-ornithine

solution for 1 hr at 37°C, washed with dPBS, and coated with iMatrix-511 (Reprocell cat#:NP892-
021) 1:50 in dPBS 3 hrs at 37°C for iSN-only culture control. After 24 hrs, media was either
changed to NCATS maturation Media + 1uM PD0332991. Cells were cultured for one week with
3x ½ media changes. On Day 14, cells were fixed for downstream IF and analysis as described
in text. n=3 biological replicates at densities indicated.

Treatment of rDRG glia with neuronal maturation media: rDRG glia were dissected, plated,
prepared, and cultured for one week as described above. On Day 7, glia were either treated with
NCATS Maturation Media + CEPT and cultured at 37°C 5% CO_2 or maintained in Glia Media at
37°C 7.5% CO_2. After 24 hrs, NCATS Maturation Media + CEPT was changed to NCATS
Maturation Media + 1uM PD0332991. Glia were matured for an additional week with ½ media
changes 3x/week and fixed for downstream Immunofluorescence (IF) on Day 14. First, cells were
fixed with 2% paraformaldehyde (PFA) (EMS cat#: 15710; 1:1 4% PFA to spent media) 10
minutes at room temperature (RT), and then in 4% PFA 20 minutes at RT. Cells were washed 3x
in PBS and stored at 4°C for downstream experimentation. n=4 biological replicates.

Identification of Glial Subtypes in rDRG Glial cultures with IF:
Immunofluorescence: Fixed rDRG glial cultures treated with and without NCATS maturation
medium were stained using the protocol described in text.

Imaging: All photomicrographs of cells cultured in Lab-Tek™ II Chambered Coverglass were
collected on the Zeiss-980 airy-scan/confocal microscope. Images of all conditions were obtained
at 20x with 1.3x zoom and bidirectional scanning in SRY-4 Airy-Scan configuration. Multiplex
imaging was utilized with 405, 488, 561, lasers. All images were obtained in a z-stack with 0.35um
step size. Systematic random sampling was utilized to image n=12-15 fields/biological replicate
across 8-wells. All images were 2D airy-scan post processed in Zen Blue software. n=4 biological
replicates.

Analysis: Particle analysis was carried out on glial marker channels. Threshold was manually set
at brightest z-stack slice for each image, and percent area occupied by particles, average particle
size, and average mean gray value were quantified for each image stack. Average values were
calculated for each field, and averages were compared between treatment groups using the t-test
function in excel where n=# of fields.

Quantification of rSG-iSN contacts in coculture with methanol fixation:

Immunofluorescence and Imaging: IF and imaging were performed as described in text. n=4
biological replicates

Analysis: Number of rSGs contacting individual iSNs was quantified using the CellCounter plugin
in FIJI. Contact was defined as apposition between P75+, human Nuc. Cy3- processes with
TUJ1+, Human Nuc. Cy3+ cell body and neurites. Number of rSGs were marked by individual
DAPI+, Human Nuc. Cy3- nuclei and rSGs contacting iSNs without a nucleus present in the field
were not counted. Contacts were counted per individual iSN, and each iSN was categorized into
defined morphologic categories. Average number of total glial contacts/cell was compared using
the t-test function and average number of total glial contacts/cell for each morphologic category
was compared using Two-way ANOVA with Bonferroni post-hoc test in GraphPad Prism 10.

Supplemental Figure Legends:

Figure S1. Human iSNs differentiated in coculture with embryonic rodent DRG glia exhibit advanced morphology and physiology. A) Timeline for pseudounipolar transition was established *in vivo* in embryonic mice expressing tamoxifen induced NTRK2-tdtomato. Morphology frequencies from whole DRGs were quantified and categorized as apoptotic (light gray), bipolar (dark gray), intermediate (pink), or pseudounipolar (red). DRG NTRK2+ neurons became pseudounipolar between days E14.5 and E16.5, indicated by black arrow. E14.5 3-5 DRGs were quantified. B) Photomicrographs illustrating physical contact between satellite glia and DRG neurons during pseudounipolarization. Green= PLP1, Red= Ntrk-tdtomato. Scale= 20um. C-D) C. Representative photomicrographs of Anatomic RealDRGs terminally differentiated in coculture or alone with NCATS maturation media and Anatomic's proprietary media Senso-MM. D. Photomicrographs depicting transitional and pseudounipolar morphologies observed in cocultured Anatomic RealDRGs. TUJ1=Green, p75=Magenta, human nuclei = Red, DAPI=Cyan. RealDRGs alone were not stained for human nuclei. Scale=50um.E-H) Average number of active electrodes during weekly MEA recordings for fourbiological replicates. Cocultured iSNs=purple, iSNs alone= orange, Glia alone= green. Average >7 active electrodes were observed in both cocultured iSNs and iSNs alone at 7 weeks in replicate 1, 5 weeks in replicate 2, 6 weeks in replicate 3, and 5 weeks in replicate 4.

Figure S2. iNCPs terminally differentiated in co-culture more advanced in neuronal differentiation compared to iNCPs alone. A) Principal component analysis and B) Hierarchal Cluster analysis of RNA sequencing reads from cocultured iSN and iSNs alone.

Figure S3. Figure 3. Rodent DRG satellite glia (rSGs) are responsible for advanced maturation of iSNs. A) Representative photomicrographs of glial cultures maintained in glial base media for 1 week and treated with NCATS Maturation Media for 1 week (Maturation Media) or maintained in glial base media for 2 weeks (Glia Media). S100b= Green, FABP7/OCT6/P75= Red, DAPI= Cyan. Scale = 50 um. B) Representative photomicrographs of glial cultures maintained in glial base media for 1 week and treated with NCATS Maturation Media for 1 week (Maturation Media) or maintained in glial base media for 2 weeks (Glia Only), iSNs in coculture and iSNs alone. Glial Cultures: S100b=Magenta, MBP=Green, DAPI=Cyan, iSN cultures: TUJ1=Magenta, MBP=Green, DAPI= Cyan. No MBP signal was detected in any culture condition. Scale=50um. C) Schematic of glial culture protocol for RNA sequencing. D-E) Principal component analysis (D) and Hierarchal Cluster analysis (E) of RNA sequencing reads from rDRG glia in coculture, treated with maturation media, and maintained in base glia media. F) GO terms significantly downregulated in cocultured glia compared to glia treated with maturation media. P adj.<0.05.

Figure S4. Physical contact between rSGs and hiSNs is essential to pseudounipolarization. A) Total rSG-iSN contacts per individual iSN in coculture with methanol fixed glia or in normal coculture. T-test performed in GraphPad Prism 10, ****=p<0.0001. B) Average number of rSG-iSN contacts per individual iSN in coculture with methanol fixed glia or in normal coculture, partitioned by morphology of iSN. Two-way ANOVA with Bonferroni post-hoc test calculated in GraphPad Prism 10. P<0.05=*, P<0.01=**, P< 0.0001=****. C) PlexinA4 mRNA reads normalized to TUBB3 (TUJ1) mRNA reads in cocultured iSNs and iSNs

999 differentiated alone. D) GJA1 mRNA reads normalized to S100b mRNA reads in cocultured and
1000 maturation media treated glia.

Supplemental Index:

References:

1. Meltzer, S., Santiago, C., Sharma, N., and Ginty, D.D. (2021). The cellular and molecular basis of somatosensory neuron development. Neuron *109*, 3736–3757. https://doi.org/10.1016/j.neuron.2021.09.004.

2. Zenker, W., and Högl, E. (1976). The prebifurcation section of the axon of the rat spinal ganglion cell. Cell Tissue Res. *165*, 345–363. https://doi.org/10.1007/BF00222438.

3. Nascimento, A.I., Mar, F.M., and Sousa, M.M. (2018). The intriguing nature of dorsal root ganglion neurons: Linking structure with polarity and function. Prog. Neurobiol. *168*, 86–103. https://doi.org/10.1016/j.pneurobio.2018.05.002.

4. Sharma, N., Flaherty, K., Lezgiyeva, K., Wagner, D.E., Klein, A.M., and Ginty, D.D. (2020). The emergence of transcriptional identity in somatosensory neurons. Nature *577*, 392–398. https://doi.org/10.1038/s41586-019-1900-1.

5. Körner, J., and Lampert, A. (2022). Functional subgroups of rat and human sensory neurons: a systematic review of electrophysiological properties. Pflugers Arch. Eur. J. Physiol. *474*, 367–385. https://doi.org/10.1007/s00424-021-02656-6.

6. Saporta, M.A., Bacha, A., and Shy, M.E. (2022). Peripheral neuropathies. In Neurobiology of Brain Disorders: Biological Basis of Neurological and Psychiatric Disorders, Second Edition (Elsevier), pp. 165–184. https://doi.org/10.1016/B978-0-323-85654-6.00006-X.

7. Rowe, R.G., and Daley, G.Q. (2019). Induced pluripotent stem cells in disease modelling and drug discovery. Nat. Rev. Genet. *20*, 377–388. https://doi.org/10.1038/s41576-019-0100-z.

8. Nicholson, M.W., Ting, C.Y., Chan, D.Z.H., Cheng, Y.C., Lee, Y.C., Hsu, C.C., Huang, C.Y., and Hsieh, P.C.H. (2022). Utility of iPSC-Derived Cells for Disease Modeling, Drug Development, and Cell Therapy. Cells *11*. https://doi.org/10.3390/cells11111853.

9. Chambers, S.M., Qi, Y., Mica, Y., Lee, G., Zhang, X.J., Niu, L., Bilsland, J., Cao, L., Stevens, E., Whiting, P., et al. (2012). Combined small-molecule inhibition accelerates developmental timing and converts human pluripotent stem cells into nociceptors. Nat.

Biotechnol. *30*, 715–720. https://doi.org/10.1038/nbt.2249.

10. Guimarães, M.Z.P., De Vecchi, R., Vitória, G., Sochacki, J.K., Paulsen, B.S., Lima, I.,
Rodrigues da Silva, F., da Costa, R.F.M., Castro, N.G., Breton, L., et al. (2018).
Generation of iPSC-Derived Human Peripheral Sensory Neurons Releasing Substance P
Elicited by TRPV1 Agonists. Front. Mol. Neurosci. *11*, 1–17.
https://doi.org/10.3389/fnmol.2018.00277.

11. Hulme, A.J., McArthur, J.R., Maksour, S., Miellet, S., Ooi, L., Adams, D.J., Finol-
Urdaneta, R.K., and Dottori, M. (2020). Molecular and Functional Characterization of
Neurogenin-2 Induced Human Sensory Neurons. Front. Cell. Neurosci. *14*, 1–18.
https://doi.org/10.3389/fncel.2020.600895.

12. Umehara, Y., Toyama, S., Tominaga, M., Matsuda, H., Takahashi, N., Kamata, Y.,
Niyonsaba, F., Ogawa, H., and Takamori, K. (2020). Robust induction of neural crest
cells to derive peripheral sensory neurons from human induced pluripotent stem cells.
Sci. Rep. *10*, 1–10. https://doi.org/10.1038/s41598-020-60036-z.

13. Nickolls, A.R., Lee, M.M., Espinoza, D.F., Szczot, M., Lam, R.M., Wang, Q., Beers, J.,
Zou, J., Nguyen, M.Q., Solinski, H.J., et al. (2020). Transcriptional Programming of
Human Mechanosensory Neuron Subtypes from Pluripotent Stem Cells. Cell Rep. *30*,
932-946.e7. https://doi.org/10.1016/j.celrep.2019.12.062.

14. Kalia, A.K., Rösseler, C., Granja-Vazquez, R., Ahmad, A., Pancrazio, J.J., Neureiter, A.,
Zhang, M., Sauter, D., Vetter, I., Andersson, A., et al. (2023). How to differentiate induced
pluripotent stem cells into sensory neurons for disease modelling: a comparison of two
protocols. Res. Sq., 1–47. https://doi.org/10.21203/rs.3.rs-3127017/v1.

15. Deng, T., Jovanovic, V.M., Tristan, C.A., Weber, C., Chu, P.H., Inman, J., Ryu, S.,
Jethmalani, Y., Ferreira de Sousa, J., Ormanoglu, P., et al. (2023). Scalable generation of
sensory neurons from human pluripotent stem cells. Stem Cell Reports *18*, 1030–1047.
https://doi.org/10.1016/j.stemcr.2023.03.006.

16. Clark, A.J., Kaller, M.S., Galino, J., Willison, H.J., Rinaldi, S., and Bennett, D.L.H. (2017).
Co-cultures with stem cell-derived human sensory neurons reveal regulators of peripheral
myelination. Brain *140*, 898–913. https://doi.org/10.1093/brain/awx012.

17. McDermott, L.A., Weir, G.A., Themistocleous, A.C., Segerdahl, A.R., Blesneac, I.,
Baskozos, G., Clark, A.J., Millar, V., Peck, L.J., Ebner, D., et al. (2019). Defining the
Functional Role of Na V 1.7 in Human Nociception. Neuron *101*, 905-919.e8.
https://doi.org/10.1016/j.neuron.2019.01.047.

18. Odawara, A., Shibata, M., Ishibashi, Y., Nagafuku, N., Matsuda, N., and Suzuki, I. (2022).
In Vitro Pain Assay Using Human iPSC-Derived Sensory Neurons and Microelectrode
Array. Toxicol. Sci. *188*, 131–141. https://doi.org/10.1093/toxsci/kfac045.

19. Wheeler, H.E., Wing, C., Delaney, S.M., Komatsu, M., and Dolan, M.E. (2015). Modeling
chemotherapeutic neurotoxicity with human induced pluripotent stem cell-derived
neuronal cells. PLoS One *10*, 1–16. https://doi.org/10.1371/journal.pone.0118020.

20. Vojnits, K., Mahammad, S., Collins, T.J., and Bhatia, M. (2019). Chemotherapy-Induced
Neuropathy and Drug Discovery Platform Using Human Sensory Neurons Converted
Directly from Adult Peripheral Blood. Stem Cells Transl. Med. *8*, 1180–1191.
https://doi.org/10.1002/sctm.19-0054.

21.   Xiong, C., Chua, K.C., Stage, T.B., Priotti, J., Kim, J., Altman-Merino, A., Chan, D., Saraf,
1085   K., Canato Ferracini, A., Fattahi, F., et al. (2020). Human Induced Pluripotent Stem Cell
1086   Derived Sensory Neurons are Sensitive to the Neurotoxic Effects of Paclitaxel. Clin.
1087   Transl. Sci., 1–14. https://doi.org/10.1111/cts.12912.

1088   22.   Chen, Y., Saski, Y., DiAntonio, A., and Milbrandt, J. (2021). SARM1 is required in human
1089   derived sensory neurons for injury- induced and neurotoxic axon degeneration. Exp.
1090   Neurol. *339*. https://doi.org/10.4049/jimmunol.1801473.The.

1091   23.   Schinke, C., Fernandez Vallone, V., Ivanov, A., Peng, Y., Körtvelyessy, P., Nolte, L.,
1092   Huehnchen, P., Beule, D., Stachelscheid, H., Boehmerle, W., et al. (2021). Modeling
1093   chemotherapy induced neurotoxicity with human induced pluripotent stem cell (iPSC) -
1094   derived sensory neurons. Neurobiol. Dis. *155*, 105391.
1095   https://doi.org/10.1016/j.nbd.2021.105391.

1096   24.   Snavely, A.R., Heo, K., Petrova, V., Ho, T.S.Y., Huang, X., Hermawan, C., Kagan, R.,
1097   Deng, T., Singeç, I., Chen, L., et al. (2022). Bortezomib-induced neurotoxicity in human
1098   neurons is the consequence of nicotinamide adenine dinucleotide depletion. DMM Dis.
1099   Model. Mech. *15*. https://doi.org/10.1242/dmm.049358.

1100   25.   Rana, P., Luerman, G., Hess, D., Rubitski, E., Adkins, K., and Somps, C. (2017).
1101   Utilization of iPSC-derived human neurons for high-throughput drug-induced peripheral
1102   neuropathy screening. Toxicol. Vitr. *45*, 111–118.
1103   https://doi.org/10.1016/j.tiv.2017.08.014.

1104   26.   Jayakar, S., Shim, J., Jo, S., Bean, B.P., Singeç, I., and Woolf, C.J. (2021). Developing
1105   nociceptor-selective treatments for acute and chronic pain. Sci. Transl. Med. *13*.
1106   https://doi.org/10.1126/scitranslmed.abj9837.

1107   27.   Matsuda, S., and Uehara, Y. (1984). Prenatal development of the rat dorsal root ganglia -
1108   A scanning electron-microscopic study. Cell Tissue Res. *235*, 13–18.
1109   https://doi.org/10.1007/BF00213717.

1110   28.   Barber, R.P., and Vaughn, J.E. (1986). Differentiation of dorsal root ganglion cells with
1111   processes in their synaptic target zone of embryonic mouse spinal cord: a retrograde
1112   tracer study. J. Neurocytol. *15*, 207–218. https://doi.org/10.1007/BF01611657.

1113   29.   Matsuda, S., Baluk, P., Shimizu, D., and Fujiwara, T. (1996). Dorsal root ganglion neuron
1114   development in chick and rat. Anat. Embryol. (Berl). *193*, 475–480.
1115   https://doi.org/10.1007/BF00185878.

1116   30.   Kiss, F. (1932). SYMPATHETIC ELEMENTS IN THE CRANIAL AND SPINAL GANGLIA.
1117   J. Anat. *66*, 488–498.

1118   31.   Fisher, B.Y.C., and Ranson, S.W. (1933). ON THE SO-CALLED SYMPATHETIC CELLS
1119   IN TI. J. Anat. *68*, 1–10.

1120   32.   Truex, R.C. (1943). DEGENERATE VERSUS MULTIPOLAR NEURONS INT SENSORY
1121   GANGLIA. J. Nerv. Ment. Dis. *97*, 606.

1122   33.   Bhatt, S., Diaz, R., and Trainor, P.A. (2013). Signals and switches in mammalian neural
1123   crest cell differentiation. Cold Spring Harb. Perspect. Biol. *5*.
1124   https://doi.org/10.1101/cshperspect.a008326.

1125   34.   Mudge, A.W. (1984). Schwann cells induce morphological transformation of sensory
1126   neurones in vitro. Nature *309*, 367–369. https://doi.org/10.1038/309367a0.

35. Takahashi, K., and Ninomiya, T. (1987). Morphological changes of dorsal root ganglion
cells in the process-forming period. Prog. Neurobiol. *29*, 393–410.
https://doi.org/10.1016/0301-0082(87)90020-7.

36. Nascimento, A.I., Da Silva, T.F., Fernandes, E.C., Luz, L.L., Mar, F.M., Safronov, B. V.,
and Sousa, M.M. (2022). Sensory neurons have an axon initial segment that initiates
spontaneous activity in neuropathic pain. Brain *145*, 1632–1640.
https://doi.org/10.1093/brain/awac078.

37. Jones, I., Yelhekar, T.D., Wiberg, R., Kingham, P.J., Johansson, S., Wiberg, M., and
Carlsson, L. (2018). Development and validation of an in vitro model system to study
peripheral sensory neuron development and injury. Sci. Rep. *8*, 1–15.
https://doi.org/10.1038/s41598-018-34280-3.

38. Kim, H.S., Kim, J.Y., Song, C.L., Jeong, J.E., and Cho, Y.S. (2020). Directly induced
human Schwann cell precursors as a valuable source of Schwann cells. Stem Cell Res.
Ther. *11*, 257. https://doi.org/10.1186/s13287-020-01772-x.

39. Pannese, E. (1969). Electron microscopical study on the development of the satellite cell
sheath in spinal ganglia. J. Comp. Neurol. *135*, 381–421.
https://doi.org/10.1002/cne.901350403.

40. Matsuda, S., Kobayashi, N., Mominoki, K., Wakisaka, H., Mori, M., and Murakami, S.
(2000). Morphological transformation of sensory ganglion neurons and satellite cells. Acta
Anat. Nippon. *73*, 612–613.

41. Wall, P.D., and Devor, M. (1983). Sensory afferent impulses originate from dorsal root
ganglia as well as from the periphery in normal and nerve injured rats. Pain *17*, 321–339.
https://doi.org/10.1016/0304-3959(83)90164-1.

42. Amir, R., and Devor, M. (1997). Spike-evoked suppression and burst patterning in dorsal
root ganglion neurons of the rat. J. Physiol. *501*, 183–196. https://doi.org/10.1111/j.1469-
7793.1997.183bo.x.

43. Amir, R., Michaelis, M., and Devor, M. (2002). Burst discharge in primary sensory
neurons: Triggered by subthreshold oscillations, maintained by depolarizing
afterpotentials. J. Neurosci. *22*, 1187–1198. https://doi.org/10.1523/jneurosci.22-03-
01187.2002.

44. Djouhri, L., Koutsikou, S., Fang, X., McMullan, S., and Lawson, S.N. (2006).
Spontaneous pain, both neuropathic and inflammatory, is related to frequency of
spontaneous firing in intact C-fiber nociceptors. J. Neurosci. *26*, 1281–1292.
https://doi.org/10.1523/JNEUROSCI.3388-05.2006.

45. Ma, C., and LaMotte, R.H. (2007). Multiple sites for generation of ectopic spontaneous
activity in neurons of the chronically compressed dorsal root ganglion. J. Neurosci. *27*,
14059–14068. https://doi.org/10.1523/JNEUROSCI.3699-07.2007.

46. Liu, C.N., Wall, P.D., Ben-Dor, E., Michaelis, M., Amir, R., and Devor, M. (2000). Tactile
allodynia in the absence of C-fiber activation: Altered firing properties of DRG neurons
following spinal nerve injury. Pain *85*, 503–521. https://doi.org/10.1016/S0304-
3959(00)00251-7.

47. Xie, W., Strong, J.A., Kim, D., Shahrestani, S., and Zhang, J.M. (2012). Bursting activity
in myelinated sensory neurons plays a key role in pain behavior induced by localized

inflammation of the rat sensory anglion. Neuroscience *206*, 212–223.
https://doi.org/10.1016/j.neuroscience.2012.01.007.Bursting.

48.   Zheng, Q., Xie, W., Lückemeyer, D.D., Lay, M., Wang, X.W., Dong, X., Limjunyawong,
N., Ye, Y., Zhou, F.Q., Strong, J.A., et al. (2022). Synchronized cluster firing, a distinct
form of sensory neuron activation, drives spontaneous pain. Neuron *110*, 209-220.e6.
https://doi.org/10.1016/j.neuron.2021.10.019.

49.   North, R.Y., Li, Y., Ray, P., Rhines, L.D., Tatsui, C.E., Rao, G., Johansson, C.A., Zhang,
H., Kim, Y.H., Zhang, B., et al. (2019). Electrophysiological and transcriptomic correlates
of neuropathic pain in human dorsal root ganglion neurons. Brain *142*, 1215–1226.
https://doi.org/10.1093/brain/awz063.

50.   Zhang, X., Chen, Y., Wang, C., and Huang, L.Y.M. (2007). Neuronal somatic ATP
release triggers neuron-satellite glial cell communication in dorsal root ganglia. Proc. Natl.
Acad. Sci. U. S. A. *104*, 9864–9869. https://doi.org/10.1073/pnas.0611048104.

51.   Rozanski, G.M., Li, Q., and Stanley, E.F. (2013). Transglial transmission at the dorsal
root ganglion sandwich synapse: Glial cell to postsynaptic neuron communication. Eur. J.
Neurosci. *37*, 1221–1228. https://doi.org/10.1111/ejn.12132.

52.   Huang, L.Y.M., Gu, Y., and Chen, Y. (2013). Communication between neuronal somata
and satellite glial cells in sensory ganglia. Glia *61*, 1571–1581.
https://doi.org/10.1002/glia.22541.

53.   Kim, Y.S., Anderson, M., Park, K., Zheng, Q., Agarwal, A., Gong, C., Saijilafu, Young,
L.A., He, S., LaVinka, P.C., et al. (2016). Coupled Activation of Primary Sensory Neurons
Contributes to Chronic Pain. Neuron *91*, 1085–1096.
https://doi.org/10.1016/j.neuron.2016.07.044.

54.   Newberry, K., Wang, S., Hoque, N., Kiss, L., Ahlijanian, M.K., Herrington, J., and Graef,
J.D. (2016). Development of a spontaneously active dorsal root ganglia assay using
multiwell multielectrode arrays. J. Neurophysiol. *115*, 3217–3228.
https://doi.org/10.1152/jn.01122.2015.

55.   Black, B.J., Atmaramani, R., Kumaraju, R., Plagens, S., Romero-Ortega, M., Dussor, G.,
Price, T.J., Campbell, Z.T., and Pancrazio, J.J. (2018). Adult mouse sensory neurons on
microelectrode arrays exhibit increased spontaneous and stimulus-evoked activity in the
presence of interleukin-6. J. Neurophysiol. *120*, 1374–1385.
https://doi.org/10.1152/jn.00158.2018.

56.   Hiranuma, M., Okuda, Y., Fujii, Y., Richard, J.P., and Watanabe, T. (2024).
Characterization of human iPSC-derived sensory neurons and their functional
assessment using multi electrode array. Sci. Rep. *14*, 1–8.
https://doi.org/10.1038/s41598-024-55602-8.

57.   Frith, T.J.R., Granata, I., Wind, M., Stout, E., Thompson, O., Neumann, K., Stavish, D.,
Heath, P.R., Ortmann, D., Hackland, J.O.S., et al. (2018). Human axial progenitors
generate trunk neural crest cells in vitro. Elife *7*, 1–27.
https://doi.org/10.7554/eLife.35786.

58.   Libby, A.R.G., Joy, D.A., Elder, N.H., Bulger, E.A., Krakora, M.Z., Gaylord, E.A.,
Mendoza-Camacho, F., Butts, J.C., and McDevitt, T.C. (2021). Axial elongation of
caudalized human organoids mimics aspects of neural tube development. Dev. *148*.
https://doi.org/10.1242/dev.198275.

59. Kee, Y., and Bronner-Fraser, M. (2005). To proliferate or to die: Role of Id3 in cell cycle
progression and survival of neural crest progenitors. Genes Dev. *19*, 744–755.
https://doi.org/10.1101/gad.1257405.

60. Heng, Y.H.E., McLeay, R.C., Harvey, T.J., Smith, A.G., Barry, G., Cato, K., Plachez, C.,
Little, E., Mason, S., Dixon, C., et al. (2014). NFIX regulates neural progenitor cell
differentiation during hippocampal morphogenesis. Cereb. Cortex *24*, 261–279.
https://doi.org/10.1093/cercor/bhs307.

61. Martynoga, B., Mateo, J.L., Zhou, B., Andersen, J., Achimastou, A., Urbán, N., van den
Berg, D., Georgopoulou, D., Hadjur, S., Wittbrodt, J., et al. (2013). Epigenomic enhancer
annotation reveals a key role for NFIX in neural stem cell quiescence. Genes Dev. *27*,
1769–1786. https://doi.org/10.1101/gad.216804.113.

62. Harris, L., Genovesi, L.A., Gronostajski, R.M., Wainwright, B.J., and Piper, M. (2015).
Nuclear factor one transcription factors: Divergent functions in developmental versus
adult stem cell populations. Dev. Dyn. *244*, 227–238. https://doi.org/10.1002/dvdy.24182.

63. Zalucki, O., Harris, L., Harvey, T.J., Harkins, D., Widagdo, J., Oishi, S., Matuzelski, E.,
Ling Hilary Yong, X., Schmidt, H., Anggono, V., et al. (2019). NFIX-mediated inhibition of
neuroblast branching regulates migration within the adult mouse ventricular-
subventricular zone. Cereb. Cortex *29*, 3590–3604.
https://doi.org/10.1093/cercor/bhy233.

64. He, J.C., Neves, S.R., Jordan, J.D., and Iyengar, R. (2006). Role of the Go/i signaling
network in the regulation of neurite outgrowth. Can. J. Physiol. Pharmacol. *84*, 687–694.
https://doi.org/10.1139/Y06-025.

65. Bromberg, K., Iyengar, R., and He, J. (2011). Regulation of neurite outgrowth by Gi/o
signaling pathways. Front Biosci *13*, 4544–4557.

66. Wolf, D., Hofbrucker-MacKenzie, S.A., Izadi, M., Seemann, E., Steiniger, F., Schwintzer,
L., Koch, D., Kessels, M.M., and Qualmann, B. (2019). Ankyrin repeat-containing N-Ank
proteins shape cellular membranes. Nat. Cell Biol. *21*, 1191–1205.
https://doi.org/10.1038/s41556-019-0381-7.

67. Kim, S.J., Woo, Y., Kim, H.J., Goo, B.S., Nhung, T.T.M., Lee, S.A., Suh, B.K., Mun, D.J.,
Kim, J.H., and Park, S.K. (2022). Retinoic acid-induced protein 14 controls dendritic spine
dynamics associated with depressive-like behaviors. Elife *11*, 1–30.
https://doi.org/10.7554/eLife.77755.

68. Christiansen, J.H., Coles, E.G., and Wilkinson, D.G. (2000). Molecular control of neural
crest formation, migration and differentiation. Curr. Opin. Cell Biol. *12*, 719–724.
https://doi.org/10.1016/S0955-0674(00)00158-7.

69. Lian, Y., Wu, C., Liu, L., and Li, X. (2024). Prediction of cell-cell communication patterns
of dorsal root ganglion cells: single-cell RNA sequencing data analysis. Neural Regen.
Res. *19*, 1367–1374. https://doi.org/10.4103/1673-5374.384067.

70. Bennett, D.L., Clark, X.A.J., Huang, J., Waxman, S.G., and Dib-Hajj, S.D. (2019). The
role of voltage-gated sodium channels in pain signaling. Physiol. Rev. *99*, 1079–1151.
https://doi.org/10.1152/physrev.00052.2017.

71. Jarvis, M.F., Honore, P., Shieh, C.C., Chapman, M., Joshi, S., Zhang, X.F., Kort, M.,
Carroll, W., Marron, B., Atkinson, R., et al. (2007). A-803467, a potent and selective

Nav1.8 sodium channel blocker, attenuates neuropathic and inflammatory pain in the rat.
Proc. Natl. Acad. Sci. U. S. A. *104*, 8520–8525.
https://doi.org/10.1073/pnas.0611364104.

72. Jones, J., Correll, D.J., Lechner, S.M., Jazic, I., Miao, X., Shaw, D., Simard, C., Osteen,
J.D., Hare, B., Beaton, A., et al. (2023). Selective Inhibition of Na V 1.8 with VX-548 for
Acute Pain . N. Engl. J. Med. *389*, 393–405. https://doi.org/10.1056/nejmoa2209870.

73. Jo, S., Zhang, H.X.B., and Bean, B.P. (2023). Use-Dependent Relief of Inhibition of
Nav1.8 Channels by A-887826. Mol. Pharmacol. *103*, 221–229.
https://doi.org/10.1124/molpharm.122.000593.

74. Haberberger, R., Scholz, A., Kummer, W., and Kress, M. (2000). M2-receptor subtype
does not mediate muscarine-induced increases in [Ca2+](i) in nociceptive neurons of rat
dorsal root ganglia. J. Neurophysiol. *84*, 1934–1941.
https://doi.org/10.1152/jn.2000.84.4.1934.

75. Tata, A.M., Vilaró, M.T., and Mengod, G. (2000). Muscarinic receptor subtypes
expression in rat and chick dorsal root ganglia. Mol. Brain Res. *82*, 1–10.
https://doi.org/10.1016/S0169-328X(00)00165-0.

76. De Angelis, F., Marinelli, S., Fioretti, B., Catacuzzeno, L., Franciolini, F., Pavone, F., and
Tata, A.M. (2014). M2 receptors exert analgesic action on DRG sensory neurons by
negatively modulating VR1 activity. J. Cell. Physiol. *229*, 783–790.
https://doi.org/10.1002/jcp.24499.

77. Xian, X.C., and Nicol, G.D. (2007). Manipulation of the potassium channel Kv1.1 and its
effect on neuronal excitability in rat sensory neurons. J. Neurophysiol. *98*, 2683–2692.
https://doi.org/10.1152/jn.00437.2007.

78. Wang, J.J., and Li, Y. (2016). KCNQ potassium channels in sensory system and neural
circuits. Acta Pharmacol. Sin. *37*, 25–33. https://doi.org/10.1038/aps.2015.131.

79. Smith, P.A. (2020). K+ Channels in Primary Afferents and Their Role in Nerve Injury-
Induced Pain. Front. Cell. Neurosci. *14*. https://doi.org/10.3389/fncel.2020.566418.

80. Bhattacharya, D., Rothstein, M., Azambuja, A.P., and Simoes-Costa, M. (2018). Control
of neural crest multipotency by wnt signaling and the Lin28/let-7 axis. Elife *7*, 1–24.
https://doi.org/10.7554/eLife.40556.

81. Shoval, I., and Kalcheim, C. (2012). Antagonistic activities of Rho and Rac GTPases
underlie the transition from neural crest delamination to migration. Dev. Dyn. *241*, 1155–
1168. https://doi.org/10.1002/dvdy.23799.

82. Kurisaki, T., Masuda, A., Osumi, N., Nabeshima, Y.I., and Fujisawa-Sehara, A. (1998).
Spatially- and temporally-restricted expression of meltrin α (ADAM12) and β (ADAM19) in
mouse embryo. Mech. Dev. *73*, 211–215. https://doi.org/10.1016/S0925-4773(98)00043-
4.

83. Samad, T.A., Rebbapragada, A., Bell, E., Zhang, Y., Sidis, Y., Jeong, S.J., Campagna,
J.A., Perusini, S., Fabrizio, D.A., Schneyer, A.L., et al. (2005). DRAGON, a bone
morphogenetic protein co-receptor. J. Biol. Chem. *280*, 14122–14129.
https://doi.org/10.1074/jbc.M410034200.

84. Zorick, T.S., Syroid, D.E., Arroyo, E., Scherer, S.S., and Lemke, G. (1996). The
transcription factors SCIP and Krox-20 mark distinct stages and cell fates in Schwann cell

1300 differentiation. Mol. Cell. Neurosci. *8*, 129–145. https://doi.org/10.1006/mcne.1996.0052.

1301 85. Wu, J., Peng, D., Voehler, M., Sanders, C., and Li, J. (2013). Structure and Expression of
1302  a Novel Compact Myelin Protein - Small VCP-Interacting Protein (SVIP). Biochem.
1303  Biophys. Res. Commun. *440*. https://doi.org/10.1016/j.bbrc.2013.09.056.Structure.

1304 86. Kastriti, M.E., Faure, L., Von Ahsen, D., Bouderlique, T.G., Boström, J., Solovieva, T.,
1305  Jackson, C., Bronner, M., Meijer, D., Hadjab, S., et al. (2022). Schwann cell precursors
1306  represent a neural crest-like state with biased multipotency. EMBO J. *41*, 1–28.
1307  https://doi.org/10.15252/embj.2021108780.

1308 87. Frank, M. (2000). MAL, a proteolipid in glycosphingolipid enriched domains: Functional
1309  implications in myelin and beyond. Prog. Neurobiol. *60*, 531–544.
1310  https://doi.org/10.1016/S0301-0082(99)00039-8.

1311 88. Tasdemir-Yilmaz, O.E., Druckenbrod, N.R., Olukoya, O.O., Dong, W., Yung, A.R.,
1312  Bastille, I., Pazyra-Murphy, M.F., Sitko, A.A., Hale, E.B., Vigneau, S., et al. (2021).
1313  Diversity of developing peripheral glia revealed by single-cell RNA sequencing. Dev. Cell
1314  *56*, 2516-2535.e8. https://doi.org/10.1016/j.devcel.2021.08.005.

1315 89. Avraham, O., Chamessian, A., Feng, R., Yang, L., Halevi, A.E., Moore, A.M., Gereau,
1316  R.W., and Cavalli, V. (2022). Profiling the molecular signature of satellite glial cells at the
1317  single cell level reveals high similarities between rodents and humans. Pain *163*, 2348–
1318  2364. https://doi.org/10.1097/j.pain.0000000000002628.

1319 90. Mapps, A.A., Thomsen, M.B., Boehm, E., Zhao, H., Hattar, S., and Kuruvilla, R. (2022).
1320  Diversity of satellite glia in sympathetic and sensory ganglia. Cell Rep. *38*.
1321  https://doi.org/10.1016/j.celrep.2022.110328.

1322 91. Cinquina, V., Calvigioni, D., Piscitelli, F., Keimpema, E., Di Marzo, V., Verkhratsky, A.,
1323  and Harkany, T. (2023). Astrocytes modulate neuronal development by S100A6
1324  signaling. bioRxiv.

1325 92. O'Malley, A.M., Shanley, D.K., Kelly, A.T., and Barry, D.S. (2014). Towards an
1326  understanding of semaphorin signalling in the spinal cord. Gene *553*, 69–74.
1327  https://doi.org/10.1016/j.gene.2014.10.005.

1328 93. Fujii, S., Yoshida, S., Inagaki, E., Hatou, S., Tsubota, K., Takahashi, M., Shimmura, S.,
1329  and Sugita, S. (2019). Immunological Properties of Neural Crest Cells Derived from
1330  Human Induced Pluripotent Stem Cells. Stem Cells Dev. *28*, 28–43.
1331  https://doi.org/10.1089/scd.2018.0058.

1332 94. Mehler, V.J., Burns, C.J., Stauss, H., Francis, R.J., and Moore, M.L. (2020). Human
1333  iPSC-Derived Neural Crest Stem Cells Exhibit Low Immunogenicity. Mol. Ther. Methods
1334  Clin. Dev. *16*, 161–171. https://doi.org/10.1016/j.omtm.2019.12.015.

1335 95. Li, C., Wang, S., Chen, Y., and Zhang, X. (2018). Somatosensory Neuron Typing with
1336  High-Coverage Single-Cell RNA Sequencing and Functional Analysis. Neurosci. Bull. *34*,
1337  200–207. https://doi.org/10.1007/s12264-017-0147-9.

1338 96. Ray, P., Torck, A., Quigley, L., Wangzhou, A., Neiman, M., Rao, C., Lam, T., Kim, J.Y.,
1339  Kim, T.H., Zhang, M.Q., et al. (2018). Comparative transcriptome profiling of the human
1340  and mouse dorsal root ganglia: An RNA-seq-based resource for pain and sensory
1341  neuroscience research https://doi.org/10.1097/j.pain.0000000000001217.

97. Fernández-Montoya, J., Avendaño, C., and Negredo, P. (2018). The glutamatergic
system in primary somatosensory neurons and its involvement in sensory input-
dependent plasticity. Int. J. Mol. Sci. *19*. https://doi.org/10.3390/ijms19010069.

98. Nawaz, M.S., Giarda, E., Bedogni, F., Montanara, P. La, Ricciardi, S., Ciceri, D., Alberio,
T., Landsberger, N., Rusconi, L., and Kilstrup-Nielsen, C. (2016). CDKL5 and shootin1
interact and concur in regulating neuronal polarization. PLoS One *11*.
https://doi.org/10.1371/journal.pone.0148634.

99. Bushlin, I., Petralia, R.S., Wu, F., Harel, A., Mughal, M.R., Mattson, M.P., and Yao, P.J.
(2008). Clathrin assembly protein AP180 and CALM differentially control axogenesis and
dendrite outgrowth in embryonic hippocampal neurons. J. Neurosci. *28*, 10257–10271.
https://doi.org/10.1523/JNEUROSCI.2471-08.2008.

100. Montanara, P. La, Hervera, A., Baltussen, L.L., Hutson, T.H., Palmisano, I., De Virgiliis,
F., Kong, G., Chadwick, J., Gao, Y., Bartus, K., et al. (2020). Cyclin-dependent-like
kinase 5 is required for pain signaling in human sensory neurons and mouse models. Sci.
Transl. Med. *12*, 1–36. https://doi.org/10.1126/SCITRANSLMED.AAX4846.

101. Hori, K., and Hoshino, M. (2017). Neuronal migration and AUTS2 syndrome. Brain Sci. *7*.
https://doi.org/10.3390/brainsci7050054.

102. Lentz, S.I., Miner, J.H., Sanes, J.R., and Snider, W.D. (1997). Distribution of the ten
known laminin chains in the pathways and targets of developing sensory axons. J. Comp.
Neurol. *378*, 547–561. https://doi.org/10.1002/(SICI)1096-
9861(19970224)378:4<547::AID-CNE9>3.0.CO;2-2.

103. Liu, C., Lin, C., Whitaker, D.T., Bakeri, H., Bulgakov, O. V., Liu, P., Lei, J., Dong, L., Li,
T., and Swaroop, A. (2013). Prickle1 is expressed in distinct cell populations of the
central nervous system and contributes to neuronal morphogenesis. Hum. Mol. Genet.
22, 2234–2246. https://doi.org/10.1093/hmg/ddt075.

104. Agrawal, M., and Welshhans, K. (2021). Local Translation Across Neural Development: A
Focus on Radial Glial Cells, Axons, and Synaptogenesis. Front. Mol. Neurosci. *14*, 1–18.
https://doi.org/10.3389/fnmol.2021.717170.

105. Kumar, S., Chakraborty, S., Barbosa, C., Brustovetsky, T., Brustovetsky, N., and
Obukhov, A.G. (2012). Mechanisms controlling neurite outgrowth in a
pheochromocytoma cell line: The role of TRPC channels. J. Cell. Physiol. *227*, 1408–
1419. https://doi.org/10.1002/jcp.22855.

106. Joseph, D.J., Williams, D.J., and MacDermott, A.B. (2011). Modulation of neurite
outgrowth by activation of calcium-permeable kainate receptors expressed by rat
nociceptive-like dorsal root ganglion neurons. Dev. Neurobiol. *71*, 818–835.
https://doi.org/10.1002/dneu.20906.

107. Bakos, J., Bacova, Z., Grant, S.G., Castejon, A.M., and Ostatnikova, D. (2015). Are
Molecules Involved in Neuritogenesis and Axon Guidance Related to Autism
Pathogenesis? NeuroMolecular Med. *17*, 297–304. https://doi.org/10.1007/s12017-015-
8357-7.

108. Wang, T., Liu, Y., Xu, X.H., Deng, C.Y., Wu, K.Y., Zhu, J., Fu, X.Q., He, M., and Luo,
Z.G. (2011). Lgl1 activation of Rab10 Promotes axonal membrane trafficking underlying
neuronal polarization. Dev. Cell *21*, 431–444.
https://doi.org/10.1016/j.devcel.2011.07.007.

109.   Stegeman, S., Jolly, L.A., Premarathne, S., Gecz, J., Richards, L.J., Mackay-Sim, A., and
1387   Wood, S.A. (2013). Loss of Usp9x Disrupts Cortical Architecture, Hippocampal
1388   Development and TGFβ-Mediated Axonogenesis. PLoS One *8*, 1–12.
1389   https://doi.org/10.1371/journal.pone.0068287.

1390   110.   Luo, X., Salgueiro, Y., Beckerman, S., Lemmon, V., Tsoulfas, P., and Park, K. (2013).
1391   Three-dimensional evaluation of retinal ganglion cell axon regeneration and pathfinding in
1392   whole mouse tissue after injury. Exp. Neurol. *247*, 653–662.
1393   https://doi.org/10.1016/j.expneurol.2013.03.001.Three-dimensional.

1394   111.   Mutalik, S.P., and Gupton, S.L. (2021). Glycosylation in axonal guidance. Int. J. Mol. Sci.
1395   *22*. https://doi.org/10.3390/ijms22105143.

1396   112.   Kuo, T.Y., Hong, C.J., and Hsueh, Y.P. (2009). Bcl11A/CTIP1 regulates expression of
1397   DCC and MAP1b in control of axon branching and dendrite outgrowth. Mol. Cell.
1398   Neurosci. *42*, 195–207. https://doi.org/10.1016/j.mcn.2009.07.006.

1399   113.   Hall, A., and Lalli, G. (2010). Rho and Ras GTPases in Axon Growth, Guidance, and
1400   Branching. Cold Spring Harb. Perspect. Biol., 1–18.

1401   114.   Luxey, M., Jungas, T., Laussu, J., Audouard, C., Garces, A., and Davy, A. (2013). Eph:
1402   Ephrin-B1 Forward Signaling Controls Fasciculation Of Sensory And Motor Axons. Dev.
1403   Biol. *383*, 264–274. https://doi.org/10.1016/j.ydbio.2013.09.010.

1404   115.   Laumonnerie, C., Da Silva, R. V., Kania, A., and Wilson, S.I. (2014). Netrin 1 and Dcc
1405   signalling are required for confinement of central axons within the central nervous
1406   system. Dev. *141*, 594–603. https://doi.org/10.1242/dev.099606.

1407   116.   Balmer, C.W., and Lamantia, A.S. (2004). Loss of Gli3 and Shh Function Disrupts
1408   Olfactory Axon Trajectories. J. Comp. Neurol. *472*, 292–307.
1409   https://doi.org/10.1002/cne.20053.

1410   117.   Butler, S.J., and Tear, G. (2007). Getting axons onto the right path: The role of
1411   transcription factors in axon guidance. Development *134*, 439–448.
1412   https://doi.org/10.1242/dev.02762.

1413   118.   Zhao, L.L., Zhang, T., Huang, W.X., Guo, T.T., and Gu, X.S. (2023). Transcriptional
1414   regulatory network during axonal regeneration of dorsal root ganglion neurons: laser-
1415   capture microdissection and deep sequencing. Neural Regen. Res. *18*, 2056–2066.
1416   https://doi.org/10.4103/1673-5374.366494.

1417   119.   Shamay-Ramot, A., Khermesh, K., Porath, H.T., Barak, M., Pinto, Y., Wachtel, C.,
1418   Zilberberg, A., Lerer-Goldshtein, T., Efroni, S., Levanon, E.Y., et al. (2015). Fmrp
1419   Interacts with Adar and Regulates RNA Editing, Synaptic Density and Locomotor Activity
1420   in Zebrafish. PLoS Genet. *11*, 1–28. https://doi.org/10.1371/journal.pgen.1005702.

1421   120.   John, A., Brylka, H., Wiegreffe, C., Simon, R., Liu, P., Jüttner, R., Bryan Crenshaw, I.,
1422   Luyten, F.P., Jenkins, N.A., Copeland, N.G., et al. (2012). Bcl11a is required for neuronal
1423   morphogenesis and sensory circuit formation in dorsal spinal cord development.
1424   Development *139*, 1831–1841. https://doi.org/10.1242/dev.072850.

1425   121.   Wang, S.H.J., Celic, I., Choi, S.Y., Riccomagno, M., Wang, Q., Sun, L.O., Mitchell, S.P.,
1426   Vasioukhin, V., Huganir, R.L., and Kolodkin, A.L. (2014). Dlg5 regulates dendritic spine
1427   formation and synaptogenesis by controlling subcellular N-cadherin localization. J.
1428   Neurosci. *34*, 12745–12761. https://doi.org/10.1523/JNEUROSCI.1280-14.2014.

122. Noseda, R., Belin, S., Piguet, F., Vaccari, I., Scarlino, S., Brambilla, P., Boneschi, F.M.,
Feltri, M.L., Wrabetz, L., Quattrini, A., et al. (2013). DDIT4/REDD1/RTP801 is a novel
negative regulator of schwann cell myelination. J. Neurosci. *33*, 15295–15305.
https://doi.org/10.1523/JNEUROSCI.2408-13.2013.

123. Gunsch, G., Paradie, E., and Townsend, K.L. (2023). Peripheral nervous system glia in
support of metabolic tissue functions. Trends Endocrinol. Metab. *34*, 622–639.
https://doi.org/10.1016/j.tem.2023.07.004.

124. Birren, S.J., Goodrich, L. V., and Segal, R.A. (2024). Satellite Glial Cells: No Longer the
Most Overlooked Glia. Cold Spring Harb. Perspect. Biol., a041367.
https://doi.org/10.1101/cshperspect.a041367.

125. Narayanan, S.P., Flores, A.I., Wang, F., and Macklin, W.B. (2009). Akt signals through
the mammalian target of rapamycin pathway to regulate CNS myelination. J. Neurosci.
29, 6860–6870. https://doi.org/10.1523/JNEUROSCI.0232-09.2009.

126. Lebeau, P., Al-Hashimi, A., Sood, S., Lhoták, Š., Yu, P., Gyulay, G., Paré, G., Wayne
Chen, S.R., Trigatti, B., Prat, A., et al. (2017). Endoplasmic reticulum stress and Ca2+
depletion differentially modulate the sterol regulatory protein PCSK9 to control lipid
metabolism. J. Biol. Chem. *292*, 1510–1523. https://doi.org/10.1074/jbc.M116.744235.

127. Bartz, F., Kern, L., Erz, D., Zhu, M., Gilbert, D., Meinhof, T., Wirkner, U., Erfle, H.,
Muckenthaler, M., Pepperkok, R., et al. (2009). Identification of Cholesterol-Regulating
Genes by Targeted RNAi Screening. Cell Metab. *10*, 63–75.
https://doi.org/10.1016/j.cmet.2009.05.009.

128. Elbaz, B., Yang, L., Vardy, M., Isaac, S., Rader, B.L., Kawaguchi, R., Traka, M., Woolf,
C.J., Renthal, W., and Popko, B. (2022). Sensory neurons display cell-type-specific
vulnerability to loss of neuron-glia interactions. Cell Rep. *40*, 111130.
https://doi.org/10.1016/j.celrep.2022.111130.

129. Meriau, P., Weill, L., Puccio, H., and Agulhon, C. (2024). Frataxin deficiency in
proprioceptive neurons is causal to inflammatory and glial responses in dorsal root
ganglia.

130. Krishnan, A., Bhavanam, S., and Zochodne, D. (2018). An intimate role for adult dorsal
root ganglia resident cycling cells in the generation of local macrophages and satellite
glial cells. J. Neuropathol. Exp. Neurol. *77*, 929–941. https://doi.org/10.1093/jnen/nly072.

131. Hall, B.E., Macdonald, E., Cassidy, M., Yun, S., Sapio, M.R., Ray, P., Doty, M., Nara, P.,
Burton, M.D., Shiers, S., et al. (2022). Transcriptomic analysis of human sensory neurons
in painful diabetic neuropathy reveals inflammation and neuronal loss. Sci. Rep. *12*, 1–
16. https://doi.org/10.1038/s41598-022-08100-8.

132. Mocciaro, A., Berdougo, E., Zeng, K., Black, E., Vagnarelli, P., Earnshaw, W., Gillespie,
D., Jallepalli, P., and Schiebel, E. (2010). Vertebrate cells genetically deficient for
Cdc14A or Cdc14B retain DNA damage checkpoint proficiency but are impaired in DNA
repair. J. Cell Biol. *189*, 631–639. https://doi.org/10.1083/jcb.200910057.

133. Park, B.S., Jo, H.W., and Jung, J. (2015). Expression profile of aminoacyl-tRNA
synthetases in dorsal root ganglion neurons after peripheral nerve injury. J. Mol. Histol.
46, 115–122. https://doi.org/10.1007/s10735-014-9601-4.

134. Ninomiya, K., Ohta, K., Yamashita, K., Mizuno, K., and Ohashi, K. (2021). PLEKHG4B

enables actin cytoskeletal remodeling during epithelial cell-cell junction formation. J. Cell
Sci. *134*. https://doi.org/10.1242/jcs.249078.

135. Repetto, D., Camera, P., Melani, R., Morello, N., Russo, I., Calcagno, E., Tomasoni, R.,
Bianchi, F., Berto, G., Giustetto, M., et al. (2014). p140Cap regulates memory and
synaptic plasticity through src-mediated and citron-N-mediated actin reorganization. J.
Neurosci. *34*, 1542–1553. https://doi.org/10.1523/JNEUROSCI.2341-13.2014.

136. Aruga, J., and Mikoshiba, K. (2003). Identification and characterization of Slitrk, a novel
neuronal transmembrane protein family controlling neurite outgrowth. Mol. Cell. Neurosci.
24, 117–129. https://doi.org/10.1016/S1044-7431(03)00129-5.

137. Chu, Y., Jia, S., Xu, K., Liu, Q., Mai, L., Liu, J., Fan, W., and Huang, F. (2023). Single-cell
transcriptomic profile of satellite glial cells in trigeminal ganglion. Front. Mol. Neurosci. *16*,
1–16. https://doi.org/10.3389/fnmol.2023.1117065.

138. Luo, Y., Xu, W., Chen, H., Warburton, D., Dong, R., Qian, B., Selman, M., Gualdie, J.,
Kolb, M., and Shi, W. (2015). A novel profibrotic mechanism mediated by TGF-β-
stimulated collagen prolyl hydroxylase expression in fibrotic lung mesenchymal cells. J.
Pathol. *236*, 13384–13394. https://doi.org/10.1002/path.4530.A.

139. Chen, P., Cescon, M., and Bonaldo, P. (2015). The Role of Collagens in Peripheral Nerve
Myelination and Function. Mol. Neurobiol. *52*, 216–225. https://doi.org/10.1007/s12035-
014-8862-y.

140. Vehviläinen, P., Hyytiäinen, M., and Keski-Oja, J. (2009). Matrix association of latent
TGF-beta binding protein-2 (LTBP-2) is dependent on fibrillin-1. J. Cell. Physiol. *221*,
586–593. https://doi.org/10.1002/jcp.21888.

141. Suto, F., Ito, K., Uemura, M., Shimizu, M., Shinkawa, Y., Sanbo, M., Shinoda, T., Tsuboi,
M., Takashima, S., Yagi, T., et al. (2005). Plexin-A4 mediates axon-repulsive activities of
both secreted and transmembrane semaphorins and plays roles in nerve fiber guidance.
J. Neurosci. *25*, 3628–3637. https://doi.org/10.1523/JNEUROSCI.4480-04.2005.

142. Mauti, O., Domanitskaya, E., Andermatt, I., Sadhu, R., and Stoeckli, E.T. (2007).
Semaphorin6A acts as a gate keeper between the central and the peripheral nervous
system. Neural Dev. *2*, 1–17. https://doi.org/10.1186/1749-8104-2-28.

143. Haklai-Topper, L., Mlechkovich, G., Savariego, D., Gokhman, I., and Yaron, A. (2010).
Cis interaction between Semaphorin6A and Plexin-A4 modulates the repulsive response
to Sema6A. EMBO J. *29*, 2635–2645. https://doi.org/10.1038/emboj.2010.147.

144. Lowry Curley, J., Catig, G.C., Horn-Ranney, E.L., and Moore, M.J. (2014). Sensory axon
guidance with semaphorin 6A and nerve growth factor in a biomimetic choice point
model. Biofabrication *6*. https://doi.org/10.1088/1758-5082/6/3/035026.

145. Pease-Raissi, S.E., Pazyra-Murphy, M.F., Li, Y., Wachter, F., Fukuda, Y.,
Fenstermacher, S.J., Barclay, L.A., Bird, G.H., Walensky, L.D., and Segal, R.A. (2017).
Paclitaxel Reduces Axonal Bclw to Initiate IP3R1-Dependent Axon Degeneration. Neuron
96, 373-386.e6. https://doi.org/10.1016/j.neuron.2017.09.034.

146. Li, Y., Pazyra-Murphy, M.F., Avizonis, D., Russo, M.T., Tang, S., Chen, C.Y., Hsueh,
Y.P., Bergholz, J.S., Jiang, T., Zhao, J.J., et al. (2022). Sarm1 activation produces
cADPR to increase intra-axonal Ca++ and promote axon degeneration in PIPN. J. Cell
Biol. *221*. https://doi.org/10.1083/jcb.202106080.

147.   Chen, X., Gan, Y., Au, N.P.B., and Ma, C.H.E. (2024). Current understanding of the
molecular mechanisms of chemotherapy-induced peripheral neuropathy. Front. Mol.
Neurosci. *17*. https://doi.org/10.3389/fnmol.2024.1345811.

148.   Paulson, S., McCabe, G., Winkler, M., Findley, T., Pascua, A., Gordon, K., Walsh, P., and
Truong, V. (2024). Scalable and Efficient Generation of iPSC-derived Human Sensory
Neurons for CIPN and Pain Research. In SLAS.

149.   Rockel, A.F., Wagner, N., Spenger, P., Ergün, S., and Wörsdörfer, P. (2023). Neuro-
mesodermal assembloids recapitulate aspects of peripheral nervous system development
in vitro. Stem Cell Reports *18*, 1155–1165. https://doi.org/10.1016/j.stemcr.2023.03.012.

150.   Lu, D., Shang, G., He, X., Bai, X. chen, and Zhang, X. (2021). Architecture of the
Sema3A/PlexinA4/Neuropilin tripartite complex. Nat. Commun. *12*, 1–13.
https://doi.org/10.1038/s41467-021-23541-x.

151.   Kitsukawa, T., Shimizu, M., Sanbo, M., Hirata, T., Taniguchi, M., Bekku, Y., Yagi, T., and
Fujisawa, H. (1997). Neuropilin-semaphorin III/D-mediated chemorepulsive signals play a
crucial role in peripheral nerve projection in mice. Neuron *19*, 995–1005.
https://doi.org/10.1016/S0896-6273(00)80392-X.

152.   Xu, X.M., Fisher, D.A., Zhou, L., White, F.A., Ng, S., Snidpper, W.D., and Luo, Y. (2000).
The transmembrane protein semaphorin 6A repels embryonic sympathetic axons. J.
Neurosci. *20*, 2638–2648. https://doi.org/10.1523/jneurosci.20-07-02638.2000.

153.   Lerman, O., Ben-Zvi, A., Yagil, Z., and Behar, O. (2007). Semaphorin3A accelerates
neuronal polarity in vitro and in its absence the orientation of DRG neuronal polarity in
vivo is distorted. Mol. Cell. Neurosci. *36*, 222–234.
https://doi.org/10.1016/j.mcn.2007.07.003.

154.   Nakamura, F., Kalb, R.G., and Strittmatter, S.M. (2000). Molecular basis of semaphorin-
mediated axon guidance. J. Neurobiol. *44*, 219–229. https://doi.org/10.1002/1097-
4695(200008)44:2<219::AID-NEU11>3.0.CO;2-W.

155.   Pannese, E. (2010). The structure of the perineuronal sheath of satellite glial cells
(SGCs) in sensory ganglia. Neuron Glia Biol. *6*, 3–10.
https://doi.org/10.1017/S1740925X10000037.

156.   Kolos, E.A., and Korzhevsky, D.E. (2024). Gap Junction Protein Connexin-43 in a Rat
Dorsal Root Ganglion. Cell tissue biol. *18*, 189–198.
https://doi.org/10.1134/S1990519X23700049.

157.   Retamal, M.A., Riquelme, M.A., Stehberg, J., and Alcayaga, J. (2017). Connexin43
hemichannels in satellite glial cells, can they influence sensory neuron activity? Front.
Mol. Neurosci. *10*, 1–9. https://doi.org/10.3389/fnmol.2017.00374.

158.   Hanani, M., and Spray, D.C. (2020). Emerging importance of satellite glia in nervous
system function and dysfunction. Nat. Rev. Neurosci. *21*, 485–498.
https://doi.org/10.1038/s41583-020-0333-z.

159.   Keane, R.W., Mehta, P.P., Rose, B., Honig, L.S., Loewenstein, W.R., and Rutishauser,
U. (1988). Neural differentiation, NCAM-mediated adhesion, and gap junctional
communication in neuroectoderm. A study in vitro. J. Cell Biol. *106*, 1307–1319.
https://doi.org/10.1083/jcb.106.4.1307.

160.   Ma, C.H.E., Brenner, G.J., Omura, T., Samad, O.A., Costigan, M., Inquimbert, P.,

Niederkofler, V., Salie, R., Sun, C.C., Lin, H.Y., et al. (2011). The BMP coreceptor RGMb promotes while the endogenous BMP antagonist noggin reduces neurite outgrowth and peripheral nerve regeneration by modulating BMP signaling. J. Neurosci. *31*, 18391–18400. https://doi.org/10.1523/JNEUROSCI.4550-11.2011.

161. Sundt, D., Gamper, N., and Jaffe, D.B. (2015). Spike propagation through the dorsal root ganglia in an unmyelinated sensory neuron: A modeling study. J. Neurophysiol. *114*, 3140–3153. https://doi.org/10.1152/jn.00226.2015.

162. Hao, H., Ramli, R., Wang, C., Liu, C., Shah, S., Mullen, P., Lall, V., Jones, F., Shao, J., Zhang, H., et al. (2023). Dorsal root ganglia control nociceptive input to the central nervous system https://doi.org/10.1371/journal.pbio.3001958.

163. Han, C., Estacion, M., Huang, J., Vasylyev, D., Zhao, P., Dib-Hajj, S.D., and Waxman, S.G. (2015). Human Nav1.8: Enhanced persistent and ramp currents contribute to distinct firing properties of human DRG neurons. J. Neurophysiol. *113*, 3172–3185. https://doi.org/10.1152/jn.00113.2015.

164. Lai, J., Gold, M.S., Kim, C.S., Biana, D., Ossipov, M.H., Hunterc, J.C., and Porreca, F. (2002). Inhibition of neuropathic pain by decreased expression of the tetrodotoxin-resistant sodium channel, NaV1.8. Pain *95*, 143–152. https://doi.org/10.1016/S0304-3959(01)00391-8.

165. Enes, J., Haburcák, M., Sona, S., Gerard, N., Mitchell, A.C., Fu, W., and Birren, S.J. (2020). Satellite glial cells modulate cholinergic transmission between sympathetic neurons. PLoS One *15*, 1–21. https://doi.org/10.1371/journal.pone.0218643.

166. Tomasello, D.L., Hurley, E., Wrabetz, L., and Bhattacharjee, A. (2017). Slick (Kcnt2) sodium-activated potassium channels limit peptidergic nociceptor excitability and hyperalgesia. J. Exp. Neurosci. *11*. https://doi.org/10.1177/1179069517726996.

167. Tsantoulas, C., Denk, F., Signore, M., Nassar, M.A., Futai, K., and McMahon, S.B. (2018). Mice lacking Kcns1 in peripheral neurons show increased basal and neuropathic pain sensitivity. Pain *159*, 1641–1651. https://doi.org/10.1097/j.pain.0000000000001255.

168. Warwick, R.A., and Hanani, M. (2013). The contribution of satellite glial cells to chemotherapy-induced neuropathic pain. Eur. J. Pain (United Kingdom) *17*, 571–580. https://doi.org/10.1002/j.1532-2149.2012.00219.x.

169. Leo, M., Schmitt, L.I., Kutritz, A., Kleinschnitz, C., and Hagenacker, T. (2021). Cisplatin-induced activation and functional modulation of satellite glial cells lead to cytokine-mediated modulation of sensory neuron excitability. Exp. Neurol. *341*. https://doi.org/10.1016/j.expneurol.2021.113695.

170. Schmitt, L.I., Leo, M., Kutritz, A., Kleinschnitz, C., and Hagenacker, T. (2020). Activation and functional modulation of satellite glial cells by oxaliplatin lead to hyperexcitability of sensory neurons in vitro. Mol. Cell. Neurosci. *105*, 103499. https://doi.org/10.1016/j.mcn.2020.103499.

171. Majd, H., Amin, S., Ghazizadeh, Z., Cesiulis, A., Arroyo, E., Lankford, K., Majd, A., Farahvashi, S., Chemel, A.K., Okoye, M., et al. (2023). Deriving Schwann cells from hPSCs enables disease modeling and drug discovery for diabetic peripheral neuropathy. Cell Stem Cell *30*, 632-647.e10. https://doi.org/10.1016/j.stem.2023.04.006.

172. Chen, Y., Tristan, C.A., Chen, L., Jovanovic, V.M., Malley, C., Chu, P.H., Ryu, S., Deng,

T., Ormanoglu, P., Tao, D., et al. (2021). A versatile polypharmacology platform promotes
cytoprotection and viability of human pluripotent and differentiated cells (Springer US)
https://doi.org/10.1038/s41592-021-01126-2.

173.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 1–21.
https://doi.org/10.1186/s13059-014-0550-8.

174.   Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L.,
et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
Innovation *2*, 100141. https://doi.org/10.1016/j.xinn.2021.100141.

1610

A.

Coculture | **iSN Alone**

0.2% DMSO | 60nM Paclitaxel | 0.2% DMSO | 60nM Paclitaxel

GJA1
Human
nuc.
DAPI
TUJ1

B.

% Area TUJ1

C.

Coculture | **iSN Alone**

0.2% DMSO | 600nM Paclitaxel | 0.2% DMSO | 600nM Paclitaxel

D.

% Area TUJ1

S1.

A.

Ntrk2+ neurons

% neuron morphology

apoptotic
bipolar
intermediate
pseudo-unipolar

44%
50%
5%
83%
14%
99%
1%
100%

E14.5 E16.5 E18.5 P5
#neurons: 51 216 151 60

B.

TUJ1   PLP1   Merge

20um

20um

C.

Real DRG Coculture, NCATS Maturation

RealDRG, NCATS Maturation

RealDRG, SensoMM Maturation

TUJ1
P75
Human nuclei
DAPI

D.

Bulge   Bell   PU-Short

E.

Active Electrodes Replicate 1

Avg # Active Electrodes

Glia Alone
Coculture
iSN Alone

1 week 2 weeks 3 weeks 4 weeks 5 weeks 6 weeks 7 weeks

iSN weeks in culture

F.

Active Electrodes Replicate 2

Avg # Active Electrodes

Glia Alone
Coculture
iSN Alone

1 week 2 weeks 3 weeks 4 weeks 5 weeks

iSN weeks in culture

G.

Active Electrodes Replicate 3

Avg # Active Electrodes

Glia Alone
Coculture
iSN Alone

1 week 2 weeks 3 weeks 4 weeks 5 weeks 6 weeks 7 weeks

iSN weeks in culture

H.

Active Electrodes Replicate 4

Active Electrodes

Glia Alone
Coculture
iSN Alone

1 week 2 weeks 3 weeks 4 weeks 5 weeks

iSN weeks in culture

S2.

S3.

S4.