*Title*:   Structural diversity and distribution of NMCP-class nuclear lamina proteins in

streptophytic algae

*Authors and Affiliations*:

Brendan S. Kosztyo[1], and Eric J. Richards[2*]

[1] Cornell University, College of Agriculture and Life Sciences, Ithaca, NY 14853

[2] Boyce Thompson Institute, Ithaca, NY 14853

*Author for Correspondence*:

Eric J. Richards

Boyce Thompson Institute

533 Tower Road

Ithaca, NY 14853

USA

Ph: 607-254-1234

**ABSTRACT**

Nuclear Matrix Constituent Proteins (NMCPs) in plants function like animal lamins, providing

the structural foundation of the nuclear lamina and regulating nuclear organization and

morphology. Although they are well-characterized in angiosperms, the presence and structure of

NMCPs in more distantly related species, such as streptophytic algae, are relatively unknown.

The rapid evolution of NMCPs throughout the plant lineage has caused a divergence in protein

sequence that makes similarity-based searches less effective. Structural features are more likely

to be conserved compared to primary amino acid sequence; therefore, we developed a filtration

protocol to search for diverged NMCPs based on four physical characteristics: intrinsically

disordered content, isoelectric point, number of amino acids, and the presence of a central coiled-

coil domain. By setting parameters to recognize the properties of bona fide NMCP proteins in

angiosperms, we filtered seven complete proteomes from streptophytic algae species and

identified strong NMCP candidates in five taxa in the Classes Zygnematophyceae,

Charophyceae, and Klebsormidophyceae. Through analysis of these proteins, we observed

structural variance in domain size between NMCPs in algae and land plants, as well as a single

block of amino acid conservation. Our analysis indicates that NMCPs are absent in the

Mesostigmatophyceae. The presence versus absence of NMCP proteins in different streptophytic

algae allows us to draw inferences about the evolution of NMCP proteins and their relationship

to different forms of mitosis throughout the plant lineage.

**SIGNIFICANCE STATEMENT**

All eukaryotic organisms contain a membrane-bound nucleus, which holds a cell's DNA, but

plants and animals have distinct sets of proteins that make up the scaffolding essential for

nuclear integrity and organization – a structure called the nuclear lamina. The major nuclear

lamina proteins in plants and animals share structural features, and we exploited these

characteristics to identify a key class of nuclear lamina proteins in streptophytic algae, allowing

us to chart the distribution of these proteins across photosynthetic organisms and to gain insight

into the evolution of nuclear organization and dynamics during cell division. Our results indicate

that the major class of nuclear lamina protein in plants evolved independently from structurally

similar nuclear lamina proteins in animals.

**KEYWORDS**

• Streptophyte

• Open mitosis

• Nuclear organization

• Proteome filtration

• Coiled-coil domain

• Intrinsically disordered domain

## INTRODUCTION

In eukaryotes, the nucleus is central to the integration and execution of cell functions, including storing the genetic material and regulating its expression. A complex of proteins called the nuclear lamina (NL), which resides directly below the inner nuclear membrane, is responsible for the organization of the nucleus and its indirect attachment to the cytoskeleton, as well as for genome and transcription factor regulation (Gruenbaum and Foisner, 2015). In metazoans, lamins are the major structural protein of the NL and the founding member of the intermediate filament superfamily (Dechat *et al.*, 2010).

Despite being ancestral and well conserved, lamins are not found constitutively throughout the eukaryotic lineage. The absence of lamin homologs could be explained by an extreme divergence in certain eukaryotic lineages (Dittmer and Misteli, 2011). Alternatively, the NL could have independent origins, being composed of distinct proteins recruited to form the NL through convergent evolution (Koreny and Field, 2016). Distinguishing between these two hypotheses requires assembly of a comprehensive catalog of lamin homologs and potential functional analogs across divergent eukaryotic lineages.

In the non-metazoan *Dictyostelium discoideum*, NE-81 represents an evolutionary precursor to the A- and B-type lamins found in vertebrates (Batsios *et al.*, 2012; Odell and Lammerding, 2023). This *Dictyostelium* protein has the structural features of metazoan lamins, with which it shares primary amino acid sequence similarity (Krüger *et al.*, 2012). Further, transgenic expression of NE-81 in a lamin A/C deficient cell line partially complements the mechanical defects exhibited by nuclei in those cells (Odell, Gräf and Lammerding, 2023). In other non-

metazoan groups, distinct NL proteins exist that are difficult to reconcile as lamin homologs. For

example, in Trypanosomes, NUP-1 and NUP-2 are NL proteins that appear to function

analogously to lamins, but these proteins lack sequence similarity with lamins and share only

some structural features (e.g., long coiled-coil domains in NUP-1; see Figure 1) (DuBois *et al.*,

2012; Maishman *et al.*, 2016; Padilla-Mejia *et al.*, 2021).

The situation in plants is more complex. Plants lack recognizable lamins yet nuclei in plant cells

have a NL structure (Masuda *et al.*, 1993; Fiserova, Kiseleva and Goldberg, 2009). The major

component of this structure in plant cells are Nuclear Matrix Constituent Proteins (NMCPs), first

discovered in the angiosperm *Daucus carota* (carrots) (Masuda *et al.*, 1997; Kimura, Kuroda and

Masuda, 2010; Ciska and Moreno Díaz de la Espina, 2013). Functional studies in *Arabidopsis*

*thaliana*, demonstrate that NMCPs (also called CRWNs, for CROWED NUCLEI, in this

species) are involved in the regulation of nuclear size and shape as well as heterochromatin

organization (Dittmer *et al.*, 2007; Sakamoto and Takagi, 2013; Wang, Dittmer and Richards,

2013; Sakamoto *et al.*, 2022). NMCPs in angiosperms are essential for survival (Wang, Dittmer

and Richards, 2013; Blunt *et al.*, 2023), but the sole *NMCP* gene can be deleted without affecting

viability in the liverwort *Marchantia polymorpha*, despite the fact that disruption of three-

dimensional chromatin organization results (Wang *et al.*, 2021). These functional studies indicate

that NMCPs play similar roles to lamins, but these major NL proteins in plants do not share

significant amino acid sequence similarity with lamins. Nor do they contain diagnostic features

of lamins, such as a C-terminal CAAX box or an immunoglobulin-fold domain (Ciska *et al.*,

2019). Nonetheless, NMCPs and lamins share structural characteristics, as illustrated in Figure 1.

The most obvious similarity is that NMCPs and lamins have a tripartite structure defined by a

long central coiled-coil domain. Another feature shared by NMCPs and lamins is the high

percentage of intrinsically disordered regions (IDRs) in the N- and C-terminal domains (Huang

*et al.*, 2021; Reimann *et al.*, 2023; Qin *et al.*, 2011). These structural similarities suggest that

NMCPs could be highly diverged lamin homologs, but these similarities are also consistent with

the convergent evolution hypothesis.

To determine if NMCPs are distant lamin homologs or evolutionarily distinct, we broadened our

search for NMCPs in the 'green lineage,' encompassing land plants and photosynthetic algae

(Merchant *et al.*, 2007). All land plants examined to date contain NMCPs, including

angiosperms, gymnosperms, ferns, liverworts and mosses (Poulet *et al.*, 2016; Ciska *et al.*,

2019). Recent similarity-based studies have shown that the presence of NMCPs in the green

lineage extends as far as streptophytic algae, a sister group to land plants, but obvious NMCPs

are absent in more distantly related green algae (Chlorophyta) (Koreny and Field, 2016; Ciska *et*

*al.*, 2019).

Recognizable NMCPs are extremely divergent, suggesting that the primary amino acid sequence

similarity might not be an effective approach to find ancestral orthologs (Ciska et al., 2019).

Moreover, unlike mammalian lamins, NMCPs lack conserved domains, such as diagnostic post-

translational modification sites, a C-terminal CaaX prenylation motif, and an immunoglobulin

domain, making sequenced-based identification even more challenging (Ciska and Moreno Díaz

de la Espina, 2013). We reasoned that overall structural organization would more likely be

conserved and therefore more useful in searching for distantly related proteins. Accordingly, we

designed a method of proteomic filtration using protein properties rather than amino acid

sequence similarity to search for highly diverged NMCP orthologs in the green lineage. One

criterion involved filtering proteins based on polypeptide size (> 800 amino acids) and acidic

isoelectric point (IEP, pH 4.0 to 6.0). A second criterion was the presence of an extended coiled-

coil domain. A third critical criterion of our filtration protocol was a high incidence of

intrinsically disordered regions (IDRs), diagnostic of all characterized NMCPs and also found in

animal lamins (see Materials and Methods). Intrinsically disordered regions (IDRs) are domains

within proteins that lack a defined three-dimensional shape (Piovesan et al., 2022). One

possibility is that having a malleable shape allows for plasticity in binding, not attainable by

ordered proteins, and dynamic structural ordering (Nishiyama *et al.*, 2018). Phase separation is

another emerging theory for IDR function, in which cellular components transition between fluid

and more rigid states of matter, often occupying a medial state (Molliex et al., 2015). IDRs

promote these phase transitions by acting as linkers and stimulating sticky interactions with other

proteins, mediated by weak hydrophobic interactions (Banjade et al., 2015; Boeynaems et al.,

2018). Phase separation and condensate formation from such interactions among proteins could

contribute to these proteins' organization of the nuclear lamina.

We focused our search on well-characterized genomes and deduced proteomes of streptophytic

algae, which comprise diverse photosynthetic organisms in sister lineages to land plants, with the

aim of understanding the evolutionary history of the main component of the plant NL. Our

results indicate that NMCP-class NL proteins are restricted to a subset of streptophytic algae, and

we found no evidence for transitional states with features more closely related to animal lamins.

These findings support the view that organisms in the plant kingdom contain a convergently

evolved NL composed of diverse NMCP proteins, rather than specialized plant-specific lamin

orthologs.

**Figure 1.** Structural comparison of NL proteins in different eukaryotes

**RESULTS**

We began our study by selecting sequenced genomes from seven different taxa in four different

Classes of streptophytic algae (see Figure 2). In addition, we examined one green algal

(chlorophytic) species (*Chlamydomonas reinhardtii*), one moss species (*Physcomitrium patens*),

and one angiosperm species (*Arabidopsis thaliana*) as reference taxa bracketing the streptophytic

algae.

**Figure 2.** Phylogenetic relationships among species that were used in this study.

**Figure 3.** Summary of proteome filtration pathway

Deduced proteomes from the sequenced genomes were processed following the filtration steps

diagrammed in Figure 3. The first characteristic used to filter each proteome was the presence of

intrinsically disordered regions (IDRs), which are diagnostic of NMCP and lamin proteins (see

Materials and Methods). We use the program Metapredict to locate regions predicted to have

high disordered content and to filter for protein size. NMCP proteins are typically very large; in

all known land plants NMCPs are ≥1000 amino acids, therefore, we selected a cutoff of 800

amino acids for filtration. Independently, we used the program Emboss to filter for protein

isoelectric point (IEP). Known NMCPs have an acidic IEP, around 5.3. We chose a threshold of

IEP between 4-6 to filter the deduced proteomes. The results from these three steps were

collected, and the tool Venny: Venn Diagrams (https://bioinfogp.cnb.csic.es/tools/venny/) was

used to select proteins that satisfied all three filters.

With the condensed list, we used the program DeepCoil to generate the probability of finding a

coiled-coil domain at each amino acid. These results were expressed graphically and analyzed

visually for similarity to the NMCP coiled-coil region. The proteins selected had a central coiled-

coil region that spanned at least 100 amino acids. The coiled-coil and IDR scores were overlayed

for each protein, and the composite graphical representations were visually inspected to generate

a list of finalists.

We applied this filtration protocol to the *A. thaliana* proteome and recovered seven protein

"finalists," four of which were bona fide NMCPs (i.e., CRWN1-4). The remaining three proteins

are predicted to be filament-like proteins, including TCS1 (TRICHOME CELL SHAPE 1),

which is a cortical microtubule binding protein (Chen *et al.*, 2016). Two additional proteins of

unknown function were found (At1g47900 and the SMAD/FHA domain protein encoded by

*At2g45460*). In the moss, *Physcomitrium patens*, three finalists were identified by the filtration

protocol, two of which are proteins that show a high degree of amino acid similarity with *A.*

*thaliana* NMCPs. These two proteins were previously identified as NMCP homologs in *P.*

*patens* (Wang, Dittmer and Richards, 2013; Ciska *et al.*, 2019). The third protein

(XP_024361755) shows a high degree of similarity with Arabidopsis TCS1 and is a likely moss

homolog of this protein. These findings suggest that the filtration protocol is robust and identifies

known NMCPs in land plant proteomes, but also retrieves a small number of additional proteins

that are likely false positives not involved in nuclear lamina formation.

Consistent with this expectation, the filtration protocol identified six proteins from

*Chlamydomonas reinhardtii,* a member of the green algae family, that lacks known NMCPs and

was used as a negative control. Four of these proteins are characterized to have roles outside of

the nucleus, including a kinesin motor domain-containing protein (A0A2K3CRT1), a predicted

tropomyosin (A0A2K3DS11), a centrosomal protein (A0A2K3D6E5), and a protein implicated

in Golgi vesicle trafficking (A0A2K3D1Z8). The remaining two Chlamydomonas proteins

identified by our filter did not have a predicted or characterized function.

**Table 1.** Stepwise protein count through filtration

| Species | Proteome | Disorder | Size | IEP | Size Disorder IEP | Coiled Coil (SDI) | Filtered Finalists | NMCP Candidates |
|---|---|---|---|---|---|---|---|---|
| *Arabidopsis thaliana* | 48266 | 21874 | 3539 | 12777 | 596 | 21 | 7 | 4 |
| *Physcomitrium patens* | 48022 | 28682 | 8900 | 12134 | 1985 | 19 | 3 | 2 |
| *Mesotaenium endlicherianum* | 11080 | 7022 | 1476 | 3047 | 296 | 26 | 5 | 1 |
| *Mesostigma viride* | 16232 | 8231 | 3045 | 4090 | 566 | 49 | 11 | 0 |
| *Spirogloea muscicola* | 27137 | 15841 | 3549 | 7006 | 640 | 42 | 13 | 0 |
| *Chara braunii* | 35672 | 21605 | 8157 | 11176 | 2223 | 19 | 5 | 1 |
| *Penium margaritaceum* | 52333 | 36805 | 2094 | 13075 | 612 | 14 | 1 | 1 |
| *Chlorokybus atmophyticus* | 8693 | 4339 | 851 | 2168 | 125 | 9 | 7 | 0 |
| *Klebsormidium nitens* | 16283 | 7846 | 1305 | 4957 | 544 | 128 | 10 | 1 |
| *Chlamydomonas reinhardtii* | 19526 | 12939 | 5051 | 4440 | 1033 | 46 | 6 | 0 |

After confirming the filtration protocol can identify NMCPs from whole proteomes with a modest number of false positives, seven streptophytic algal species were analyzed using the same approach. Once all four physical characteristics (size, disorder, IEP, and coiled-coil region similarity) were filtered, finalists were analyzed to examine their structural organization. Proteins with confirmed coiled-coil domains were put in a graphical overlay that combined coiled-coil and intrinsically disorder region prediction on a per-residue basis. Graphical analysis allowed for the protein to be visualized holistically and compared to angiosperm NMCP, to assess organizational similarity.

Each species filtered returned between 1 and 13 finalist proteins after this graphical analysis. We

know from the filtration of the control proteomes that a few false positives can be expected.

Therefore, another step was needed to differentiate NMCP candidate proteins from the false

positives. To supplement our sequence-agnostic approach, we applied multiple sequence

alignments to identify amino acid motifs that distinguished among the finalists generated by the

filtration protocol. As shown in Figure 4, a single protein from four of the seven streptophytic

algae species examined contained a 50-amino acid region of similarity in the coiled-coil region

(see the rightmost column of Table 1). The region corresponds to a short, degenerate but

conserved region in the N-terminus of the coiled-coil region previously identified in angiosperm

NMCPs (Masuda *et al.*, 1997).

```
Arabidopsis CRWN1   DPRILPEKISELEKELFEYQHSMGLLLIEKKEWSSQYEALQQAFEEVNEC
Physcomitrium       DRAALHARIAALEAELYDYQYNMGLLLLQRKTWSSQVDELKAAVADAQGT
Mesotaenium         ERGALQARVEELEDELYEYQYNMGVLLMERAEWQPKFDDARVAHAEAEEA
Penium              ERDELMLKVADLGAQLHDTQYNMGLLLMEQAKWNTQYEDARAALAEAEED
Chara               ERNFMVDHIKQLEDELYKYQYHMGLLLMERKSLLADNAKLDSMKKEYEEI
Klebsormidium       EKEELQRQLDDVERELADYRYNLGVLILEHKKCKPQIDELEKALQRTREE
                    :     :   ::  :    :* . :: :*:*::::         .              .
```

**Figure 4.** Multiple sequence alignment of a weakly conserved block within algal NMCPs

After identifying this region of sequence similarity in the coiled-coil domain, we wanted to

determine if the filtration protocol had missed any other NMCP candidates containing this

sequence. The 50-amino acid region of the *P. patens* NMCP protein shown in Figure 4 was used

as an independent query in a series of local BLAST searches against the proteome of each algal

species. The candidate proteins observed in *M. endlicherianum, C. braunii, P. margaritaceum,*

and *K. nitens* returned as significant matches (see Supplementary Table 1). In *P. margaritaceum,*

a second protein was identified (pm015243g0010), which was annotated as a putative NMCP in

the existing genome annotation (Jiao *et al.*, 2020). This protein has a very high sequence

similarity with the NMCP candidate identified in *P. margaritaceum,* however, both the N- and

C-terminal domains are truncated relative to the bona fide candidate (pm001739g0030).

Consequently, the shorter protein was discarded in our filtration protocol because it lacked

significant intrinsically disordered regions at the N- and C-termini. Other top "hits" returned

from the local BLAST searches had insignificant scores. Existing annotation and additional

protein family analysis of these less significant "hits" revealed relationships to characterized

proteins outside the nuclear lamina. These findings suggest that there is only one NMCP protein

in *M. endlicherianum, C. braunii,* and *K. nitens*, and possibly two in *P. margaritaceum*, but we

found no evidence for NMCPs in *S. muscicola, C. atmosphyticus,* nor *M. viride*.

We next investigated whether these absences might result from genome misannotation. Using the

50-amino acid sequence of *P. patens,* we searched, using tBLASTn, both the transcriptomes and

the genomes of three streptophytic algae taxa apparently lacking NMCPs. The transcriptome and

genome scans of *C. atmosphyticus* and *M. viride* returned a few proteins in the search with

marginally significant e-values; these proteins were the same as those returned in the BLAST

search against the algal proteomes (see Supplementary Table 1). These results strengthen our

initial conclusion that NMCP proteins are absent in *C. atmosphyticus,* and *M. viride*.

For *S. muscicola*, scans against the annotated transcriptome failed to return NMCP candidate

proteins; however, tBLASTn searches against the genome identified three independent sequence

scaffolds with putative exonic sequences capable of encoding a peptide with similarity to the

conserved 50-amino acid region (see Supplementary Figure 1A). By analyzing RNA-seq reads

aligned to genome scaffolds, we reconstructed the gene structure of three nearly complete NMCP

genes in *S. muscicola* (see Supplementary Figure 1B and Supplementary File 1). The three

deduced Spirogloea NMCP proteins share high identity (>80%) and contain a characteristic

extended coiled-coil region and a C-terminal domain with a high intrinsic disorder score

(Supplementary Figure 1C). Gaps in genome sequence obscure the true nature of the N-terminal

domain, but each gene is expressed based on paralog-specific polymorphisms in the RNA-seq

reads. A whole genome triplication was observed in *S. muscicola* (Cheng *et al.*, 2019), consistent

with our finding of three highly similar NMCP paralogs in this species.

**Figure 5.** Conserved NMCP motifs across algal candidates

We next examined the structural similarities among the four streptophytic algal NMCPs recovered by our filtration protocol. Although these proteins have a similar domain organization, they share only low amino acid sequence similarity, with the highest degree of sequence identity in the coiled-coil domain (21-29%) (Supplementary Figure 2). The four streptophytic algal proteins contain some of the amino acid motifs outside of the coiled-coil region that were originally identified by Ciska et al. (2019) as conserved among NMCP proteins in land plants and algae (Figure 5 and Supplementary Figure 3).

The length of NMCP proteins and their constituent domains varies between angiosperms and

streptophytic algae. Figure 5 illustrates a trend of increased size of NMCPs, particularly the

intrinsically disordered region, moving from angiosperms to moss to streptophytic algae

(Supplementary Figure 4). When the sizes of the three domains of the tripartite NMCPs are

compared within and between land plants and streptophytic algae, we noted a larger variance

among the algal proteins (see Figure 6). The size of the N-terminal domain averages over 150

amino acids for the four algal proteins identified here, roughly twice the size of N-terminal

domains of typical land plant NMCPs (Figure 5). A similar trend is observed for both the coiled-

coil and the C-terminal region in streptophytic algal NMCPs compared to land plant NMCPs,

although the differences in this sample comparison are not statistically significant (Figure 6).

These comparisons demonstrate that considerable variation in NMCP domain size is tolerated in

streptophytic algae while preserving the overall tripartite organization of the protein.

**Figure 6.** Comparison of the size of each tripartite domain in NMCPs between land plants and

algae

**Figure 7.** Distribution of NMCP proteins in the green lineage

**DISCUSSION**

NMCPs in land plants resemble lamins in structural organization, as illustrated in Figure 1. We reasoned that we might be able to identify transitional stages in the evolution of NMCPs by looking more closely for structural homologs in streptophytic algae in sister lineages of land plants. Would it be possible to start with validated NMCP-class proteins in land plants and bridge out through algae toward the Last Eukaryotic Common Ancestor to identify a connection to lamins? Or are NMCPs truly plant-specific (that is, limited to the 'green lineage')? If the latter, can we identify NMCPs in all lineages of photosynthetic organisms?

We developed a filtering protocol that considers the physical characteristics of proteins as criteria to search a proteome for potential homologs without dependence on sequence similarity. NMCP candidates could be identified in streptophytic algae through the analysis of four parameters: intrinsically disordered region content, peptide size, isoelectric point, and the presence of an extended coiled-coil domain. Candidate proteins were assessed by visualizing the organization of the tripartite structure with reference to known angiosperm NMCPs, such as *Arabidopsis* CRWN proteins. The filtration protocol (Figure 3) reduced the number of candidates to a small pool (Table 1), and we introduced a final step based on the presence of a weakly conserved region of 50 amino acids (Figure 4).

Through this process, we identified NMCP proteins in five streptophytic algal taxa: *Klebsormidium nitens*; *Chara braunii*; *Mesotaenium endlicherianum*; *Penium margaritaceum*; and *Spirogloea muscicola* (Supplementary Table 2). The *Spirogleoa muscicola* genome, which underwent a recent triplication (Cheng *et al.*, 2019), encodes three closely related NMCPs

(Supplementary Figure 1 and Supplementary File 1), but the remaining four taxa contain only a

single NMCP protein. Although we found a second NMCP paralog candidate in *Penium*, this

protein contains only the coiled-coil region. It is possible that this second predicted *Penium*

protein was derived from a full-length progenitor NMCP or that the unusual protein structure

stems from misannotation of the gene.

Despite retention of a characteristic tripartite structure, the size of the domains varies

considerably among the five algal NMCPs (Figure 5), and the overall size of the algal proteins

tends to be larger than their counterparts in land plants (Figure 6). Further, the algal NMCPs

share very limited amino acid similarity (Supplementary Figure 2), suggesting that primary

sequence can diverge significantly if the proteins maintain the structural characteristics that

define the NMCP family. Nonetheless, a few short sequence motifs shared among NMCP

proteins can be identified among the algal proteins (see Figures 4 and 5) (Ciska *et al.*, 2019).

Absent, however, is a well-conserved C-terminal motif, which is present in many NMCP

proteins in land plants and implicated in ABA regulation (Supplementary Figure 3) (Zhao *et al.*,

2016). The function of the few short amino acid motifs shared by algal NMCPs is unknown; it is

possible that these regions represent sites for interaction with core nucleoskeletal or nuclear

membrane proteins.

A previous study by Koreny and Field (2016) uncovered NMCP candidates in streptophytic

algae in the genera *Coleochaete, Klebsormidium, Nitella,* and possibly *Spirogyra*, and Ciska et

al. (2019) added additional analysis. These NMCP proteins were identified using a similarity-

based search from transcriptomic datasets, but the limitations of such an approach were

acknowledged by Koreny and Field (2016), citing the extreme divergence of these proteins. In

addition, similarity among even unrelated coiled-coil regions could lead to a high background of

false positives (Koreny and Field, 2016). We examined the three novel algal NMCP candidates

identified by Koreny and Field (2016) using the structural characteristics described in this study

(note that the previous study also found the protein from *Klebsormidium* uncovered in our

filtration protocol). Two of the three NMCP candidates conform to our structural criteria

(JV767595 from *Nitella mirabilis*; and GB-SL01031242 from *Coleochaete orbicularis*);

however, the *Spirogyra* protein (GB-SM01021289 from *Spirogyra pratensis*) is a relatively short

protein (634 amino acids) that lacks significant intrinsically disordered regions.

While NMCPs are clearly present in different streptophytic algae, these nuclear lamina proteins

are not found in all streptophytes. We conclude that NMCPs are absent from the proteomes of

*Chlorokybus atmophyticus* and *Mesostigma viride* based on the following results. First, we

analyzed the proteins in our finalist candidate list shown in Table 1 for the possibility that they

might represent previously unrecognized NMCP variants. However, these candidates were

related to proteins with roles outside the nuclear lamina and therefore are unlikely to be highly

diverged NMCP homologs. Second, to identify homologs that might not have the diagnostic

tripartite domain organization, we searched the seven complete proteomes using a query

corresponding to a 50-amino acid block in the coiled-coil region from a *Physcomitrium patens*

NMCP (Figure 4). No new NMCP candidates were recovered by this approach from proteomes

of *Chlorokybus atmophyticus* or *Mesostimga viride*. We did, however, identify an additional

NMCP-like protein from *Penium margaritaceum* that eluded the filtering protocol because it

lacked disordered domains as noted above. Third, to account for NMCP proteins potentially

overlooked due to errors in genome annotation, we performed tBLASTn searches with the same

query against algal transcriptomes, as well as genome sequences, and found no new candidates.

These findings argue that *Chlorokybus atmophyticus* and *Mesostimga viride* lack NMCP

proteins.

Mapping the presence and absence of NMCP proteins relative to the phylogeny of streptophytic

algae demonstrates that NMCPs do not appear at the boundary between green algae

(Chlorophyta) and streptophytic algae. The streptophytes most closely related to green algae,

*Mesostigma viride* and *Chlorokybus atmophyticus,* do not have a recognizable NMCP. Moving

toward lineages more closely related to land plants, NMCPs first appear in Klebsormidophyceae,

represented by *Klebsormidium nitens*. Taking a further step toward land plants, NMCPs are

present in the Charophyceae. Within the Zygnematophyceae, we found NMCPs in all three

different taxa examined. These data indicate that NMCP proteins were first acquired in the green

lineage at the branch point distinguishing the extant Classes Mesostigmatophyceae and

Klebsormidophyceae (Figure 7). Our results, which demonstrate that NMCP-class nuclear

lamina proteins are present in some but not all taxa in the green lineage, are consistent with the

convergent evolution model for the nuclear lamina in streptophytes. We found no evidence for

intermediate NMCP variants, resembling ancestral lamins, corresponding to more basal nodes of

the green lineage. Thus, we conclude that NMCPs are plant-specific proteins with an

independent origin and are not highly diverged lamins.

The distribution of NMCP proteins in streptophytic algae provides the opportunity to test the

hypothesis that NMCPs play a role in the control of nuclear dynamics disassembly and

reassembly during mitosis, analogous to lamins (Goldman *et al.*, 2002). According to this

hypothesis, NMCPs would be required to carry out an open mitosis, such as that which occurs in

land plants, where the nuclear envelope breaks down before chromosomal segregation and

reassembles after division (Rose, 2007). In closed mitosis, the nuclear envelope remains intact

throughout division, and the nucleus partitions into two daughter nuclei during cytokinesis

(Heath, 1986). Consistent with the hypothesis, green algae (Chlorophyta), which undergo closed

mitosis, lack NMCPs (Heath, 1986; Koreny and Field, 2016). Bracketed by green algae and land

plants, the streptophytic algae display a range of different mitotic types, including open, semi-

closed, and closed. In some cases, streptophytes that contain NMCPs, including in the genus

*Klebsormidium, Nitella*, and *Chara*, exhibit an open mitosis (Heath, 1986). The

Zygnematophyceae have a semi-closed mitosis, characterized by partial disassembly of the

nuclear envelope (Van Den Hoek, Mann and Jahns, 1995). This consideration suggests that all

taxa in Zygnematophyceae should contain an NMCP to mediate a partial nuclear breakdown, and

our results correspond to this prediction (Figure 7). The state of the nucleus during mitotic

division, however, is not necessarily predicted by presence or absence of an NMCP homolog. In

certain streptophytes more distantly related to land plants, open mitosis occurs without NMCPs.

*Mesostigma* is believed to be the final genus included in the Streptophyta at the boundary

between the Chlorophyta (Rodríguez-Ezpeleta *et al.*, 2007). Its sister group, represented in our

study by *Chlorokybus atmophyticus*, has an open mitosis but does not have an NMCP (Lokhorst,

Sluiman and Star, 1988).

Therefore, no simple correlation exists between the presence/absence of NMCPs and the form of

mitosis in streptophytic algae. The specific roles played by NMCPs in nuclear dynamics remain

to be defined, but we anticipate that investigations exploiting the diversity of nuclear types and

nuclear lamina components in streptophytic algae will help elucidate these roles.

**MATERIALS AND METHODS**

Each proteomic filtration consisted of four general steps, compounding the previous results

before advancing. Four physical characteristics were assessed for filtration: Amino acid chain

length, intrinsically disordered regions, isoelectric point, and coiled-coil patterns. In Arabidopsis,

CRWN proteins are well studied and provide a convenient positive control and a point of

reference for the filtration methods. For example, the CRWN1 protein is 1132 amino acids long,

and a threshold of 800 was chosen to account for potential candidates with fewer amino acids

than CRWN1. The coiled-coil region of CRWN1 is about 700 amino acids long and has a short

N-terminus of about 50 amino acids.

Metapredict, a machine learning-based intrinsic disorder prediction software, generates IDR

predictions for each protein from the amino acid sequence (Emenecker, Griffith and Holehouse,

2021). For all proteomic datasets, IDRs were predicted for each individual protein, and values

for angiosperms and gymnosperms were compared to the existing literature to confirm their

accuracy. Metapredict returned disorder probability scores describing the confidence of a

disordered region on a per residue basis across each protein for a complete proteome. These

scores were consolidated to a per protein basis by converting per residue scores to a binary

system. The developers of Metapredict recommend that a score of 0.3 constitutes significant

disorder; subsequently, the threshold of the binary systems was set at 0.3. Taking the binary

results per residue, each protein was assigned a percentage of residues satisfying the threshold.

The output reported the number of total residues, as well as the number of disordered amino

acids and total number of amino acids, allowing for amino acid chain length to be filtered

simultaneously with disorder.

The proteins' isoelectric points were calculated in parallel with filtration for overall size and

disorder content. The isoelectric point (IEP) of a protein, or pKi, describes the average pKa of

each amino acid residue over an entire protein. The IEP allows for the visualization of a protein's

optimal conditions to function. Lamin isoelectric points are typically acidic, ranging from about

4-6 pKi (Krohne and Benavente, 1986).  The bioinformatics package EMBOSS was used for

local isoelectric point prediction in UNIX (Rice, Longden and Bleasby, 2000). Each protein

returned a single score which was then filtered to satisfy the acidic range of lamin pKi.

Filtered lists of proteins for size, IDR, and IEP were compiled and consolidated using Venny's

biological Venn diagram tool to select for proteins that satisfied all three criteria

(https://bioinfogp.cnb.csic.es/tools/venny/). A FASTA file of only the selected proteins was then

generated and filtered for the coiled-coil domain.

The presence of a long coiled-coil domain is strongly characteristic of lamin proteins. To model

these regions computationally, the program DeepCoil, a coiled-coil domain prediction program,

was used to assign each residue a probability of being part of a coiled-coil in the tertiary

structure (Ludwiczak *et al.*, 2019). The scores of each protein were visualized graphically and

analyzed for similar patterns to the coiled-coil domain of CRWN1 in *A. thaliana.* A long central

coiled-coil, with a disordered N-terminus of about 100 amino acids and a disordered C-terminus

of varying length is indicative of an NMCP-like protein. Each coiled-coil output was assessed

visually for these characteristics and was assigned a similarity grouping from 1 (no coiled-coil)

to 5 (very similar coiled-coil pattern).

**ACKNOWLEDGEMENTS**

We are grateful for the work of Rachel Christopherson, who worked on an early version of this project. We thank the Adrian Powell, Ryan Preble, and other members of the Boyce Thompson Institute Computational Biology Center for their technical support and advice. This work was supported by funding from the National Science Foundation [grant URoL-2022048 to EJR].

line

*Data Availability*

Genome and Proteome Sources

The *Arabidopsis thaliana* proteome was obtained via TAIR (www.arabidopsis.org). The proteome for *Penium margariticeum* was obtained from the Zhangjun Fei lab (Jiao *et al.*, 2020). The proteome for *Klebsormidium nitens* (Hori et al 2014), *Chlamydomonas reinhardtii* (Merchant *et al.*, 2007)*,* and *Physcomitrium patens* (Rensing *et al.*, 2008) were obtained via the NCBI's GenBank genome portal. The proteomic information for the remaining algae: *Mesotaenium endlicherianum* (Cheng *et al.*, 2019)*, Spirogloea musicola* (Cheng *et al.*, 2019)*, Mesostigma viride* (Wang *et al.*, 2020)*, Chlorokybus atmophyticus* (Wang *et al.*, 2020)*,* and *Chara braunii* (Nishiyama *et al.*, 2018) were downloaded from the JGI genome portal (https://genome.jgi.doe.gov/portal/). Links to each dataset are provided in Supplementary Table 3.

488   *Code Availability*

489   The code used for the filtration pipeline in this manuscript can be accessed at

490   https://github.com/brendankz/Proteome-filtration.

*Figure 1:* Structural comparison of NL proteins in different eukaryotes. Scores for intrinsically disordered (blue) and coiled-coil (red) regions were mapped per residue for human lamin A, human lamin B1, Trypanosome NUP-1 and -2, NE81 in *Dictyostelium*, and CRWN1 in *Arabidopsis*. Human lamins, CRWN1, and NE81 share an overall tripartite organization, with disordered regions flanking a long, central coiled-coil region. NE81 differs from human lamins and CRWN proteins in having an elongated N-terminus. The Trypanosome proteins lack this tripartite organization. Note that the proteins lengths are normalized and not drawn to scale.

*Figure 2:* Phylogenetic relationships among species that were used in this study. Branch lengths are proportional to evolutionary distance, based on Figure 1 from Cheng *et al.* (2019). The modified position of Spirogloea between land plants and  Zygnematophyceae is supported by the recent work of Hess *et al.* (2022). Four Classes of streptophytic algae were examined: Zygnematophyceae, Charophyceae, Klebsormidophyceae, and Mesostigmatophyceae. Charophyceae, Klebsormidophyceae, and Mesostigmatophyceae are closely related and grouped together (in orange) because of similarity in their morphological characteristics. Green algae (Chlorophyta) are more diverged from land plants (represented here by Angiosperms and Mosses) than streptophytic algae.

*Figure 3:* Summary of proteome filtration pathway. Each filtration step is represented by a separate box. The method used to execute each filtration step is shown in the upper half of each box, and the characteristic being evaluated is shown in the lower portion. Intrinsic disorder is

represented in blue, size is represented in green, isoelectric point is shown in purple, and coiled-

coil is shown in red.

*Figure 4:* Multiple sequence alignment of a weakly conserved block within algal NMCPs. The

50 amino acids of the N-terminus of the coiled-coil regions of NMCP-like candidates and two

reference NMCPs are shown. The core of the motif is a well-conserved MGLL sequence, which

lies close to the N-terminal border of the extended coiled-coil domain of the proteins. Amino

acids in green are polar, red are non-polar, blue are negatively charged, and pink are positively

charged. The *Physcomitrium* protein, NMCP1, used in the multiple sequence alignment is

A0A2K1L4A2 encoded by the *PHYPA_003649* gene.

*Figure 5:* Conserved NMCP motifs across algal candidates. Algal candidate NMCPs and *P.*

*patens* NMCP1 (A0A2K1L4A2_PHYPA) aligned showing the differences in domain size and

presence of conserved regions identified in Ciska et al. (2019). Spirogloea is represented by

NMCP A in this figure. Regions in blue represent intrinsic disordered domains, while regions in

red represent the coiled-coil domain. The total length of each protein is indicated to the right, #

of amino acids; domains are drawn to scale. The yellow box represents the 50-amino acid motif

at the beginning of the coiled-coil domain described in this study (Figure 4). The head domain

characterized as motif 11 in Ciska et al. (2019) is represented in green. The tail motif 5 is

represented in aqua, motif 14 is represented in orange, and motif 4 is represented in purple (see

Supplementary Figure 3).

*Figure 6:* Comparison of the size of each tripartite domain in NMCPs between land plants and

algae. Length of different domains (# of amino acids) of NMCP proteins from land plants

compared to streptophytic algae. Representative proteins from land plants (n=8) are shown in

gray, including *A. thaliana* CRWN1 and 4; *S. lycopersicum* NMCP1B and 2; *Salvinia cucullata*

NMCP1 and 2; and both *P. patens* NMCPs. NMCP proteins from streptophytic algae (n=5) are

shown in green and are composed of the identified NMCP candidates from the filtering protocol

in *S. muscicola*, *M. endlicherianum, C. braunii*, and *K. nitens*, and *P. margaritaceum*. Error bars

indicate standard deviation. The asterisk over the algae N-terminus and coiled-coil domain

indicates a significant difference in domain sizes assessed using a Tukey HSD test (p<0.01).

*Figure 7:* Distribution of NMCP proteins in the green lineage. Presence or absence of NMCP or

NMCP-like protein by species and number of proteins identified. Color scheme of groups

corresponds to that shown in Figure 2. Branch lengths are proportional to the tree shown in

Figure 2. The hypothesized acquisition of an NMCP protein is indicated by a triangle.

*Table 1:* Genomic and proteomic file sources. Links to proteome sources for all species

considered in this study.

*Table 2:* Stepwise protein count through filtration. This table summarizes the protein number

present after each filtration step was executed. *Proteome* describes the total number of proteins

before filtration. *Disorder* corresponds to the proteins satisfying the Metapredict threshold of 0.3.

*Size* shows the number of proteins over 800 amino acids in length. IEP represents the proteins

that had an isoelectric point in the range of 4-6 pKi. *Size, Disorder, IEP* describes the number of

proteins that satisfy all three requirements. *Coiled-Coil* shows the number of proteins with a

structurally similar coiled-coil region via DeepCoil (EMBL) per residue predictions, of the

proteins that were included in SDI. *Filtered Finalists* describes the number of proteins, of those

that had satisfied the previous requirements, with a similar graphical distribution of disorder per

residue, which also have protein duplicates removed to describe the number of discrete proteins

considered. *Candidates* combines the data from the coiled-coil analysis, mapping the per residue

disorder scores with the per residue coiled-coil pattern to look for graphical similarities to a

typical NMCP map and ensure both motifs are accounted for and is compared with a multiple

sequence alignment to known NMCPs.

**REFERENCES**

Batsios, P. *et al.* (2012) 'A lamin in lower eukaryotes?', *Nucleus (Austin, Tex.)*, 3(3), pp. 237–

243. Available at: https://doi.org/10.4161/nucl.20149.

Blunt, E.L. *et al.* (2023) 'The nuclear lamina is required for proper development and nuclear

shape distortion in tomato', *Journal of Experimental Botany*, 74(18), pp. 5500–5513. Available

at: https://doi.org/10.1093/jxb/erad294.

Chen, L. *et al.* (2016) 'TCS1, a Microtubule-Binding Protein, Interacts with KCBP/ZWICHEL

to Regulate Trichome Cell Shape in Arabidopsis thaliana', *PLoS genetics*, 12(10), p. e1006266.

Available at: https://doi.org/10.1371/journal.pgen.1006266.

Cheng, S. *et al.* (2019) 'Genomes of Subaerial Zygnematophyceae Provide Insights into Land

Plant Evolution', *Cell*, 179(5), pp. 1057-1067.e14. Available at:

https://doi.org/10.1016/j.cell.2019.10.019.

Ciska, M. *et al.* (2019) 'Evolutionary history and structure of nuclear matrix constituent proteins,

the plant analogues of lamins', *Journal of Experimental Botany*, 70(10), pp. 2651–2664.

Available at: https://doi.org/10.1093/jxb/erz102.

Ciska, M. and Moreno Díaz de la Espina, S. (2013) 'NMCP/LINC proteins', *Plant Signaling &*

*Behavior*, 8(12), p. e26669. Available at: https://doi.org/10.4161/psb.26669.

Dechat, T. *et al.* (2010) 'Nuclear Lamins', *Cold Spring Harbor Perspectives in Biology*, 2(11), p.

a000547. Available at: https://doi.org/10.1101/cshperspect.a000547.

Dittmer, T.A. *et al.* (2007) 'LITTLE NUCLEI Genes Affecting Nuclear Morphology in

Arabidopsis thaliana', *The Plant Cell*, 19(9), pp. 2793–2803. Available at:

https://doi.org/10.1105/tpc.107.053231.

Dittmer, T.A. and Misteli, T. (2011) 'The lamin protein family', *Genome Biology*, 12(5), p. 222.

Available at: https://doi.org/10.1186/gb-2011-12-5-222.

DuBois, K.N. *et al.* (2012) 'NUP-1 Is a Large Coiled-Coil Nucleoskeletal Protein in

Trypanosomes with Lamin-Like Functions', *PLOS Biology*, 10(3), p. e1001287. Available at:

https://doi.org/10.1371/journal.pbio.1001287.

Emenecker, R.J., Griffith, D. and Holehouse, A.S. (2021) 'Metapredict: a fast, accurate, and

easy-to-use predictor of consensus disorder and structure', *Biophysical Journal*, 120(20), pp.

4312–4319. Available at: https://doi.org/10.1016/j.bpj.2021.08.039.

Fatafta, H. *et al.* (2021) 'Chapter Five - Molecular simulations of IDPs: From ensemble

generation to IDP interactions leading to disorder-to-order transitions', in V.N. Uversky (ed.)

*Progress in Molecular Biology and Translational Science*. Academic Press (Dancing Protein

Clouds: Intrinsically Disordered Proteins in the Norm and Pathology, Part C), pp. 135–185.

Available at: https://doi.org/10.1016/bs.pmbts.2021.06.003.

Fiserova, J., Kiseleva, E. and Goldberg, M.W. (2009) 'Nuclear envelope and nuclear pore

complex structure and organization in tobacco BY-2 cells', *The Plant Journal*, 59(2), pp. 243–

255. Available at: https://doi.org/10.1111/j.1365-313X.2009.03865.x.

Goldman, R.D. *et al.* (2002) 'Nuclear lamins: building blocks of nuclear architecture', *Genes &*

*Development*, 16(5), pp. 533–547. Available at: https://doi.org/10.1101/gad.960502.

Hawkins, J., Davis, L. and Bodén, M. (2007) 'Predicting Nuclear Localization', *Journal of*

*Proteome Research*, 6(4), pp. 1402–1409. Available at: https://doi.org/10.1021/pr060564n.

Heath, I.B. (1986) 'Nuclear Division: A Marker for Protist Phylogeny?', *Progress in*

*Protistology*, 1, pp. 115–162.

Hess, S. *et al.* (2022) 'A phylogenomically informed five-order system for the closest relatives of

land plants', *Current Biology*, 32(20), pp. 4473-4482.e7. Available at:

https://doi.org/10.1016/j.cub.2022.08.022.

Hori, K. *et al.* (2014) 'Klebsormidium flaccidum genome reveals primary factors for plant

terrestrial adaptation', *Nature Communications*, 5, p. 3978. Available at:

https://doi.org/10.1038/ncomms4978.

Huang, S. *et al.* (2021) 'A phase-separated nuclear GBPL circuit controls immunity in plants',

*Nature*, 594(7863), pp. 424–429. Available at: https://doi.org/10.1038/s41586-021-03572-6.

Jiao, C. *et al.* (2020) 'The Penium margaritaceum Genome: Hallmarks of the Origins of Land

Plants', *Cell*, 181(5), pp. 1097-1111.e12. Available at: https://doi.org/10.1016/j.cell.2020.04.019.

Kimura, Y., Kuroda, C. and Masuda, K. (2010) 'Differential nuclear envelope assembly at the

end of mitosis in suspension-cultured Apium graveolens cells', *Chromosoma*, 119(2), pp. 195–

204. Available at: https://doi.org/10.1007/s00412-009-0248-y.

Koreny, L. and Field, M.C. (2016) 'Ancient Eukaryotic Origin and Evolutionary Plasticity of

Nuclear Lamina', *Genome Biology and Evolution*, 8(9), pp. 2663–2671. Available at:

https://doi.org/10.1093/gbe/evw087.

Krohne, G. and Benavente, R. (1986) 'The nuclear lamins. A multigene family of proteins in

evolution and differentiation', *Experimental Cell Research*, 162(1), pp. 1–10. Available at:

https://doi.org/10.1016/0014-4827(86)90421-0.

Krüger, A. *et al.* (2012) 'Characterization of NE81, the first lamin-like nucleoskeleton protein in

a unicellular organism', *Molecular Biology of the Cell*, 23(2), pp. 360–370. Available at:

https://doi.org/10.1091/mbc.E11-07-0595.

Lemieux, C., Otis, C. and Turmel, M. (2007) 'A clade uniting the green algae Mesostigma viride

and Chlorokybus atmophyticus represents the deepest branch of the Streptophyta in chloroplast

genome-based phylogenies', *BMC Biology*, 5(1), p. 2. Available at: https://doi.org/10.1186/1741-

7007-5-2.

Liang, Z. *et al.* (2020) 'Mesostigma viride Genome and Transcriptome Provide Insights into the

Origin and Evolution of Streptophyta', *Advanced Science (Weinheim, Baden-Wurttemberg,*

*Germany)*, 7(1), p. 1901850. Available at: https://doi.org/10.1002/advs.201901850.

Lokhorst, G.M., Sluiman, H.J. and Star, W. (1988) 'The Ultrastructure of Mitosis and

Cytokinesis in the Sarcinoid Chlorokybus Atmosphyticus (Chlorophyta, Charophyceae)

Revealed by Rapid Freeze Fixation and Freeze Substitution', *Journal of Phycology*, 24, pp. 237–

248.

Ludwiczak, J. *et al.* (2019) 'DeepCoil-a fast and accurate prediction of coiled-coil domains in

protein sequences', *Bioinformatics (Oxford, England)*, 35(16), pp. 2790–2795. Available at:

https://doi.org/10.1093/bioinformatics/bty1062.

Maishman, L. *et al.* (2016) 'Co-dependence between trypanosome nuclear lamina components in

nuclear stability and control of gene expression', *Nucleic Acids Research*, 44(22), pp. 10554–

10570. Available at: https://doi.org/10.1093/nar/gkw751.

Masuda, K. *et al.* (1993) 'Residual structure and constituent proteins of the peripheral framework

of the cell nucleus in somatic embryos from Daucus carota L.', *Planta*, 191(4), pp. 532–540.

Available at: https://doi.org/10.1007/BF00195755.

Masuda, K. *et al.* (1997) 'Peripheral Framework of Carrot Cell Nucleus Contains a Novel

Protein Predicted to Exhibit a Long α-Helical Domain', *Experimental Cell Research*, 232(1), pp.

173–181. Available at: https://doi.org/10.1006/excr.1997.3531.

Merchant, S.S. *et al.* (2007) 'The Chlamydomonas genome reveals the evolution of key animal

and plant functions', *Science (New York, N.Y.)*, 318(5848), pp. 245–250. Available at:

https://doi.org/10.1126/science.1143609.

Nishiyama, T. *et al.* (2018) 'The Chara Genome: Secondary Complexity and Implications for

Plant Terrestrialization', *Cell*, 174(2), pp. 448-464.e24. Available at:

https://doi.org/10.1016/j.cell.2018.06.033.

Odell, J., Gräf, R. and Lammerding, J. (2023) 'Heterologous expression of Dictyostelium

discoideum NE81 in mouse embryo fibroblasts reveals conserved mechanoprotective roles of

lamins', *Molecular Biology of the Cell*, p. mbc.E23-05-0193. Available at:

https://doi.org/10.1091/mbc.E23-05-0193.

Odell, J. and Lammerding, J. (2023) 'Lamins as structural nuclear elements through evolution',

*Current Opinion in Cell Biology*, 85, p. 102267. Available at:

https://doi.org/10.1016/j.ceb.2023.102267.

Padilla-Mejia, N.E. *et al.* (2021) 'A hub-and-spoke nuclear lamina architecture in trypanosomes',

*Journal of Cell Science*, 134(12), p. jcs251264. Available at: https://doi.org/10.1242/jcs.251264.

Peter, A. and Stick, R. (2015) 'Evolutionary aspects in intermediate filament proteins', *Current*

*Opinion in Cell Biology*, 32, pp. 48–55. Available at: https://doi.org/10.1016/j.ceb.2014.12.009.

Poulet, A. *et al.* (2016) 'Exploring the evolution of the proteins of the plant nuclear envelope',

*Nucleus*, 8(1), pp. 46–59. Available at: https://doi.org/10.1080/19491034.2016.1236166.

Qin, Z. *et al.* (2011) 'Structure and stability of the lamin A tail domain and HGPS mutant',

*Journal of structural biology*, 175(3), pp. 425–433. Available at:

https://doi.org/10.1016/j.jsb.2011.05.015.

Reimann, T.M. *et al.* (2023) 'The large GTPase AtGBPL3 links nuclear envelope formation and

morphogenesis to transcriptional repression', *Nature Plants*, 9(5), pp. 766–784. Available at:

https://doi.org/10.1038/s41477-023-01400-5.

Rensing, S.A. *et al.* (2008) 'The Physcomitrella genome reveals evolutionary insights into the

conquest of land by plants', *Science (New York, N.Y.)*, 319(5859), pp. 64–69. Available at:

https://doi.org/10.1126/science.1150646.

Rice, P., Longden, I. and Bleasby, A. (2000) 'EMBOSS: the European Molecular Biology Open

Software Suite', *Trends in genetics: TIG*, 16(6), pp. 276–277. Available at:

https://doi.org/10.1016/s0168-9525(00)02024-2.

Rodríguez-Ezpeleta, N. *et al.* (2007) 'Phylogenetic Analyses of Nuclear, Mitochondrial, and

Plastid Multigene Data Sets Support the Placement of Mesostigma in the Streptophyta',

*Molecular Biology and Evolution*, 24(3), pp. 723–731. Available at:

https://doi.org/10.1093/molbev/msl200.

Rose, A.-K. (2007) 'Open mitosis: nuclear envelope dynamics.', in *In Cell Division Control in*

*Plants, ed.* Heidelberg, Germany: Springer-Verlag, pp. 207–30.

Sakamoto, Y. and Takagi, S. (2013) 'LITTLE NUCLEI 1 and 4 regulate nuclear morphology in

Arabidopsis thaliana', *Plant & Cell Physiology*, 54(4), pp. 622–633. Available at:

https://doi.org/10.1093/pcp/pct031.

Sakamoto, T. *et al.* (2022) 'Two-step regulation of centromere distribution by condensin II and

the nuclear envelope proteins', *Nature Plants*, 8(8), pp. 940–953. Available at:

https://doi.org/10.1038/s41477-022-01200-3.

Uversky, V.N. (2013) 'Intrinsically Disordered Proteins', in S. Maloy and K. Hughes (eds)

*Brenner's Encyclopedia of Genetics (Second Edition)*. San Diego: Academic Press, pp. 124–126.

Available at: https://doi.org/10.1016/B978-0-12-374984-0.01698-3.

Van Den Hoek, Mann, D.G. and Jahns, H.M. (1995) 'Chlorophyta: Class 10.

Zygnematophyceae', in *Algae: An Introduction to Phycology*. Cambridge University Press, pp.

461-473.

Wang, N. *et al.* (2021) 'Characterization of a Plant Nuclear Matrix Constituent Protein in

Liverwort', *Frontiers in Plant Science*, 12. Available at:

https://www.frontiersin.org/articles/10.3389/fpls.2021.670306

Wang, S. *et al.* (2020) 'Genomes of early-diverging streptophyte algae shed light on plant

terrestrialization', *Nature Plants*, 6(2), pp. 95–106. Available at: https://doi.org/10.1038/s41477-

019-0560-3.

Zhao, W. *et al.* (2016) 'The Arabidopsis CROWDED NUCLEI genes regulate seed germination

by modulating degradation of ABI5 protein', *Journal of Integrative Plant Biology*, 58(7), pp.

669–678. Available at: https://doi.org/10.1111/jipb.12448.

769
770
771