1

Symmetry facilitated the evolution of

heterospecificity and high-order stoichiometry in vertebrate hemoglobin

Carlos R. Cortez-Romero (1), Jixing Lyu (4), Arvind S. Pillai (2,5), Arthur Laganowsky (4), Joseph W. Thornton (2,3)

Department of (1) Cell and Molecular Biology, (2) Ecology and Evolution, and (3) Human Genetics, University of Chicago, Chicago, IL, 60637. (4) Department of Chemistry, Texas A&M University, College Station, TX, 77843. (5) Institute of Protein Design, University of Washington, Seattle, WA, 98195

Correspondence: joet1@uchicago.edu

Competing Interest Statement: The authors declare no competing interests.

Classification: Major classification - Biological Sciences **Minor classifications –** Biochemistry/Evolution

Keywords: protein evolution, molecular complexes, multimeric proteins, heterodimers, evolution of specificity, gene duplication, isology

This PDF file includes:
Main Text
Figures 1 to 6
Supplementary figures 1 to 13
Material and methods
Citations

ABSTRACT

Many proteins form paralogous multimers – molecular complexes in which evolutionarily related
proteins are arranged into specific quaternary structures. Little is known about the mechanisms
by which they acquired their stoichiometry (the number of total subunits in the complex) and
heterospecificity (the preference of subunits for their paralogs rather than other copies of the
same protein). Here we use ancestral protein reconstruction and biochemical experiments to
study historical increases in stoichiometry and specificity during the evolution of vertebrate
hemoglobin (Hb), a $\alpha_2\beta_2$ heterotetramer that evolved from a homodimeric ancestor after a gene
duplication. We show that the mechanisms for this evolutionary transition were simple. One
hydrophobic substitution in subunit β after the gene duplication was sufficient to cause the
ancestral dimer to homotetramerize with high affinity across a new interface. During this same
interval, a single-residue deletion in subunit α at the older interface conferred specificity for the
heterotetrameric form and the *trans*-orientation of subunits within it. These sudden transitions in
stoichiometry and specificity were possible because the interfaces in Hb are isologous –
involving the same surface patch on interacting subunits, rotated 180° relative to each other.
This architecture amplifies the impacts of individual mutations on stoichiometry and specificity,
especially in higher-order complexes, and allows single substitutions to differentially affect
heteromeric vs homomeric interactions. Our findings suggest that elaborate and specific
symmetrical molecular complexes may often evolve via simple genetic and physical
mechanisms.

Significance statement

Many molecular complexes are made up of proteins related by gene duplication, but how these
assemblies evolve is poorly understood. Using ancestral protein reconstruction and biochemical
experiments, we dissected how vertebrate hemoglobin, which comprises two copies each of two
related proteins, acquired this architecture from a homodimeric ancestor. Each aspect of this
transition – from dimer to tetramer and homomer to heteromer – had a simple genetic basis: a
single-site amino acid change in each protein drove these changes in size and specificity.
These transitions were possible because hemoglobin's architecture is symmetric, which
amplified the effect of small biochemical changes on the assembly of the complex. Many protein
complexes are symmetrical, suggesting that they too may have evolved via simple genetic
mechanisms.

INTRODUCTION

Protein multimers – associations of multiple protein subunits arranged in specific quaternary architectures – carry out most biochemical functions in living cells (1, 2). The mechanisms by which these complexes evolved their stoichiometry and specificity present some puzzling questions (2-10). Multimers assemble via interfaces that typically contain dozens of sterically and electrostatically complementary residues, and higher-than-dimeric stoichiometries (tetramers, octamers, etc.) use several such interfaces on each subunit (11). This seems to imply that many sequence substitutions would be required for a new multimeric assembly to originate during evolution.

A second complication is that many multimers are composed of paralogs -- proteins related to each other by gene duplication (12). Paralogs are genetically and structurally indistinguishable when generated by duplication, so initially they assemble indiscriminately into homomers and heteromers. Most complexes, however, have evolved specificity for either the homomeric or heteromeric form, with the latter being the most common outcome (12, 13). How specificity evolves is unclear, because mutations that affect multimerization are expected to cause correlated effects on the affinities of homomerization and heteromerization (6,12,14). The structural similarity of paralogs seems to imply that substitutions in both paralogs are required to confer any specificity at all. This complication is magnified for higher-order paralogous multimers, in which one might expect that every interface must evolve specificity to mediate assembly into the complex's particular architecture.

A critical factor in the evolution of specificity and high-order stoichiometry may be whether a multimer assembles through symmetrical interfaces. In many complexes, identical or paralogous subunits bind each other using an isologous interface – a form of symmetry in which a surface patch on one subunit binds to the same patch on its partner but rotated 180 degrees relative to each other (1). Isologous complexes might, in principle, have the potential to evolve changes in stoichiometry and specificity through simpler mechanisms than nonisologous head-to-tail interfaces. A single substitution appears twice across the interface(s) of an isologous homodimer or heterotetramer, four times in a homotetramer, etc. (Fig. 1A). Mutations that weakly affect affinity on their own can therefore confer large effects on the assembly of isologous multimers (1,5,9,15-17). Isology also changes the way that mutations can affect specificity. In a nonisologous interface, specificity requires mutations on both surfaces so that the tails are recognizably different from each other and each head prefers one tail over the other. In an isologous interface, however, a substitution on the surface of just one subunit has the potential to differentially affect the affinity of each kind of complex, because it will appear twice in the interface of a homomer, once in the heteromer, and not at all in the other homomer (Fig. 1A).

Little is known about the historical evolution of heterospecific complexes or the role of symmetry in this process, especially in high-order complexes. Biochemical and protein engineering studies have addressed the determinants of binding affinity in both homomeric and heteromeric interfaces of extant proteins (20-26). But the genetic and structural mechanisms by which those

interactions were acquired long ago are often different from their derived forms in the present
(27). Ancestral sequence reconstruction (ASR) can address this limitation by experimentally
characterizing the effects of historical sequence changes when introduced into ancestral
proteins. ASR has been used to understand the evolution of specificity after duplication in head-
to-tail paralogous heteromers (18,19) and in multimers composed of unrelated proteins, which
are by definition asymmetrical (26). But we know of no studies that have addressed how
isologous heteromers historically evolved their specificity or how specificity in high-order
complexes was acquired. A recent in silico analysis predicted that it should be possible for
specificity in heterodimers to evolve rapidly after gene duplication through small perturbations in
binding energy (28), but the underlying mechanisms and historical relevance of this
phenomenon are unknown.

Here we use ASR to study the evolution of higher-order stoichiometry and specificity in
vertebrate hemoglobin (Hb), the major carrier of oxygen in the blood of jawed vertebrates. Hb is
a paralogous $\alpha_2\beta_2$ heterotetramer ((17), Fig. 1B), assembly of which is mediated by two distinct
and isologous interface patches (IF1 and IF2). Each subunit of the tetramer uses its IF1 to bind
IF1 of a paralogous subunit; two of these heterodimers bind to each other using the IF2 on each
subunit to make the tetramer ((29), Fig. 1B). Hbα and Hbβ descend from a gene duplication
deep in the vertebrate lineage (Fig. 1C), and their sequences retain sufficient phylogenetic
signal to allow high-confidence reconstruction of ancestral Hb protein sequences. Using ASR,
we recently showed experimentally that extant Hb evolved its heterotetrameric architecture in
two phases from a monomeric precursor via a homodimeric intermediate (17). In the first phase,
prior to the gene duplication that yielded paralogous α and β lineages, a monomeric ancestor
evolved the capacity to homodimerize with moderate affinity across IF1. In the second phase –
after the gene duplication but before the last common ancestor of all vertebrates – binding
across IF2 was acquired, yielding the tetrameric stoichiometry, and specificity for the
heteromeric form $\alpha_2\beta_2$ also evolved (Fig. 1C).

Here we characterize the genetic and physical mechanisms that mediated the evolutionary
transition from homodimer to heterotetramer in this second phase. By experimentally
characterizing reconstructed ancestral hemoglobin subunits and the effects of historical
sequence changes on them, we address the following questions: 1) How many substitutions
were required to confer tetrameterization across IF2, and what thermodynamic and structural
mechanisms mediated their effects? 2) Did the evolution of specificity for the heterotetrameric
form require sequence changes at one or both interfaces, in one or both subunits, and what
physical mechanisms drove the acquisition of this specificity? 3) How did the symmetry of Hb's
two interfaces affect this evolutionary transition to a high-order, heterospecific architecture? 4)
Does a mutational propensity favor increased molecular complexity during the evolution of
isologous complexes?

RESULTS

Evolution of tetrameric stoichiometry. We first sought to identify the historical sequence
changes that conferred tetramerization after duplication of the ancestral homodimer Anc$\alpha\beta$. We

focused on the branch leading from the duplication of Ancαβ to Ancβ (the Hbβ subunit in the
last common ancestor of jawed vertebrates), because Ancβ heterotetramerizes with Ancα (the
Hbα subunit in the jawed vertebrate ancestor) and, like extant Hbβs, also homotetramerizes
with itself. We previously found two amino acid replacements that occurred on the branch
which, if introduced together into Ancαβ, are sufficient to confer high-affinity assembly into
homotetramers (17). One of these (q40W) is buried in the IF2 interface, whereas the other
(t37V) makes contacts across both IF1 and IF2 (Fig. 1D. 4, using lower and upper case to
denote ancestral and derived amino acids, respectively). W40 is strictly conserved in
Hbβ subunits throughout the jawed vertebrates, and V37 is conserved in Hbβ of most taxa (**Fig.**
S1).

Here we isolated the individual contributions of each amino acid changer by introducing them
singly into Ancαβ and characterizing their effect on assembly into tetramers using size-exclusion
chromatography (SEC) and native mass spectrometry (nMS) (30,31). We found that q40W
alone is sufficient to recapitulate the evolution of Hb's tetrameric stoichiometry. Ancαβ forms
only dimers in SEC at 100 μM of total protein subunits; by contrast, the mutant Ancαβ$_{q40W}$ is
tetrameric, with occupancy of the tetramer similar to that observed in the derived Ancα + Ancβ
complex and human Hb (Fig. 1E). We used nMS across a titration series to measure the affinity
with which dimers associate into tetramers and found that the tetramerization affinity of
Ancαβ$_{q40W}$ (Kd 10 μM) is stronger than that of Ancα + Ancβ (29 μM) and human Hb (34 μM)
(Fig. 1F). The conclusion that q40W is sufficient to confer tetramerization is robust to statistical
uncertainty about the ancestral sequence: similar experiments using a different reconstruction of
Ancαβ that incorporates alternative residues at all ambiguously reconstructed sites yields
almost identical results (Fig. S2).

The other historical replacement, t37V, is not sufficient to confer tetramerization. Mutant
Ancαβ$_{t37V}$ confers no detectable tetramer occupancy by SEC, even at 1 mM (Fig. 1G), and it
displays no measurable affinity to form tetramers using nMS (Fig. 1H). When combined with
q40W, however, t37V does increase affinity of the dimer-tetramer transition by a factor of 6
compared to the effect of q40W alone (Fig. 1D; Fig. S3).

In principle, a sequence change could also facilitate tetramerization by increasing affinity of the
monomer-to-dimer transition; by increasing the effective concentration of dimers, more
tetramers would be produced at a given protein concentration, even if affinity of the dimer-
tetramer transition were unchanged. Using nMS, we found that t37V improves the monomer-
dimer affinity of Ancαβ by >100-fold (Fig. 1H; Fig. S3). Substitution q40W, in contrast, has no
effect on monomer-dimer affinity. These findings are consistent with the structural location of
these residues -- t37V contributes to both IF1 and IF2 and q40W to IF2 only -- and they explain
why t37V does not confer tetramerization on its own but enhances the impact of q40W.

A likely physical mechanism for the effect of q40W is that tryptophan's bulky hydrophobic side
chain nestles into a hydrophobic divot on the IF2 surface of the facing subunit, and is further
strengthened by a hydrogen bond to 102D (32). To test this hypothesis, we identified alternative
amino acid replacements with similar biochemical properties and measured whether they also

could have caused Ancαβ to evolve into a tetramer. Like tryptophan, the bulky hydrophobic
residues phenylalanine or tyrosine at this position also confer tetramerization, albeit at affinity
slightly weaker than q40W but similar to that of Ancα+Ancβ and human Hb. The greater affinity
of tryptophan may be due to its longer side chain, which buries more hydrophobic surface area
across the interface; the hydrogen bond with 102D could make a small contribution but is not
necessary, because phenylalanine confers tetramerization but provides no hydrogen bond
donor. Leucine, in contrast, which has a smaller volume and no hydrogen bonding capacity,
confers no measurable tetramerization (Fig. 1I). High-affinity homotetramerization could
therefore have evolved via any of three different aromatic replacements at site 40.

Taken together, these data indicate that replacing the amino acid at a single residue position
was sufficient to confer tetramerization during historical Hb evolution, and several alternative
replacements at the same site could also have caused the acquisition of this higher-order
stoichiometry.

Isology facilitated IF2 evolution. How could a single amino acid replacement cause such a
dramatic change in stoichiometry? The Hb tetramer can be viewed as two heterodimers, each of
which is mediated by isologous assembly across IF1 (the larger interface); these heterodimers
then bind to each other isologously across IF2. We hypothesized that this doubly symmetrical
architecture allowed substitution q40W to confer the dimer-tetramer evolutionary transition,
because isology causes the derived amino acid to appear four times in the homotetramer and
twice in the heterotetramer.

If this hypothesis is correct, then assembly across IF2 by the derived Hb protein should require
assembly across IF1 to multiply the intrinsic affinity of IF2 (Fig. 1A). We tested this prediction by
introducing q40W into Ancαβ but doing so under conditions that prevent assembly across IF1.
We first compromised dimerization across IF1 genetically by reverting the IF1 surface to the
ancestral states of the monomeric ancestor AncMH; these mutations abolish dimer occupancy,
leaving a monomers-only population at 20 µM (Fig. 2A). We then introduced q40W into these
IF1-ablated mutants and assessed stoichiometry using nMS. As predicted, these proteins do not
form any observable dimers or tetramers (detection limit ~1 µM) (Fig. S4). Similar results are
found when we used the combination of t37V/q40W to confer association across IF2 or the
mutation P127R – which introduces unsatisfied positive charges into IF1 -- to compromise IF1.
(Fig. 2B). The IF2 mutations do not compromise heme binding or solubility, because the mutant
proteins are purifiable and heme-bound in nMS.

We also tested whether assembly across IF2 could have been historically acquired before
dimerization across IF1 evolved. We introduced t37V/q40W into the ancestral monomer AncMH
– which existed before the evolution of dimerization -- and tested whether dimer assembly
across IF2 can be conferred in this background. As predicted, only monomers were observed,
with no dimers or higher stoichiometries detected (Fig. 2C). Acquisition of multimerization
across IF2 by q40W and by the pair t37V/q40W therefore depends on the prior evolution of
dimerization via IF1.

These observations can be explained by a simple model in which the two symmetrical interfaces
contribute independently to the energy of binding. A single iteration of IF2 is too weak to confer
measurable binding of two monomers into a dimer; however, IF1 is stronger and mediates
assembly of dimers. Each IF1-mediated dimer presents two iterations of the IF2 surface patch,
doubling the total energy of IF2-mediated assembly of dimers into tetramers. Because of the
exponential relationship between energy and occupancy, a weak IF2 can therefore confer high-
affinity binding but only if IF1 is already present. The affinities that we measured are consistent
with this simple model. If the energy of dimer-to-tetramer assembly is twice that of monomer-
dimer binding using the same interface, then the Kd of IF2-mediated tetramerization should be
the square of the Kd of IF2-mediated dimerization (Fig. 2D). The Kd of the dimer-tetramer
transition by Ancαβ $_{t37V/q40W}$ across IF2 is 1 mM, which predicts that the affinity of IF2-mediated
monomer-dimer transition when IF1 is compromised should be ~ 1mM. Consistent with this
prediction, we detected no dimer occupancy by Ancαβ $_{t37V/q40W;\ IF1reverted}$ using an assay that can
quantify Kd up to 400 μM (see Methods). This simple additive model therefore explains most –
and possibly all -- of the difference in affinity conferred when IF2 is doubled in the symmetrical
tetramer. The dependence of assembly across IF2 upon the presence of IF1 does not imply any
direct physical interaction between the interfaces or any conformational change in one interface
caused by binding at the other. We cannot rule out the possibility that IF1 binding may also
allosterically modify IF2 and increase its affinity beyond the additive effect conferred by
isologous repetition alone; however, any such effect must be relatively small,.

Taken together, these data indicate that the isologous architecture of IF1 and IF2 facilitated the
evolution of the Hb tetramer via substitution q40W. Without this doubly symmetrical architecture,
IF2 would have been too weak to mediate multimerization. The dependence of q40W's effect on
the presence of IF1 also creates contingency and order-dependence in the evolution of the Hb
complex. We previously showed that IF1 evolved before the duplication of the dimeric ancestor
Ancαβ (17). Our present results show that if that IF1-mediated dimer had never evolved,
substitution q40W at IF2 would not have been sufficient to drive the acquisition of the tetrameric
stoichiometry, and the ancestral Hb protein would have remained a monomer. If events had
occurred in the opposite order – with the affinity-enhancing substitution at IF2 occurring first –
this intermediate ancestor would have been a monomer; when the substitutions that confer
binding across IF1 did occur, they would have triggered an immediate evolutionary transition
from monomer to tetramer.

Heteromeric specificity evolved at a single interface. We next focused on understanding the
evolution of Hb's specificity for the heterotetrameric form, which was acquired during the same
phylogenetic interval after the duplication of Ancαβ. Our first question was whether specificity for
heteromeric interactions was conferred by sequence changes at IF1, IF2, or both. Our
previously published experiments suggest that evolutionary changes at IF2 confer no specificity:
when all historical substitutions that occurred at the IF2 surface during the post-duplication
interval are introduced into Ancαβ and this protein is coexpressed with Ancα, an indiscriminate
mixture of homotetramers, $\alpha_1\beta_3$ heterotetramers, and $\alpha_2\beta_2$ heterotetramers is produced (17). We
therefore hypothesized that heterospecificity of the Hb tetramer is encoded entirely by IF1, such

that Ancα and Ancβ specifically heterodimerize across IF1, and these heterodimers then bind to
each other via a nonspecific IF2, yielding $\alpha_2\beta_2$ heterotetramers.

This hypothesis makes two predictions: 1) IF1 mediates specific assembly of α and β subunits
into heterodimers, and 2) this specificity is sufficient to account for the heterospecificity of $\alpha_2\beta_2$
heterotetramer. To test the first hypothesis, we characterized the specificity of hetero- vs
homodimer assembly by IF1 under two different conditions in which no binding across IF2
occurs. First, we diluted a coexpressed mixture of Ancα and Ancβ to concentrations at which
dimers rather than tetramers assemble: at 50 µM, only heterodimers and heterotetramers form;
at 5 µM, only heterodimers are observed (Fig. 3A). IF2 does not mediate assembly of
monomers into dimers in the absence of IF1 (Fig. 2A, 2B), so these heterodimers must be IF1-
mediated, indicating that IF1 is heterospecific (Fig. 3A). Second, we expressed Ancα and Ancβ
separately and mixed them at equal and moderate concentration; because tetramerization
requires co-folding, only IF1 dimers form (33), and these are predominantly heterodimers (Fig.
3B, Fig. S5). Finally, we engineered protein Ancβ' – a variant of Ancβ in which all IF2 residues
that were substituted between Ancαβ and Ancβ are reverted to the ancestral state, thus
abolishing binding across IF2– and found that it also forms predominantly heterodimers when
mixed with Ancα (Fig. 3C, Fig. S5). Together, these data indicate that the derived IF1 is specific,
preferentially mediating assembly into heterodimers.

To test the second prediction – that heterospecificity mediated by IF1 is sufficient to drive
specific assembly of $\alpha_2\beta_2$ heterotetramers even if IF2 is nonspecific – we measured the affinities
of homomerization and heteromerization across IF1 and used these measurements to predict
their effects on tetramer specificity in the absence of any specificity at IF2. Using nMS and
Ancβ', we found that IF1's heterodimerization affinity (Kd=0.6 µM) is slightly worse than its
homodimerization affinity (0.2 µM), but both are far better than the Ancα homodimer (24 µM)
(Fig. 3D, S6, S7, S8, S9). We then predicted occupancy of each stoichiometry as the
concentration of Hb subunits changes, given these affinities at IF1 and assuming that IF2 has a
dimer-to-tetramer affinity of 30 µM, as measured in Ancα + Ancβ, with no preference for
homomeric or heteromeric binding (Fig. 1D). At low concentrations, the system produces almost
exclusively IF1-mediated heterodimers. The predominance of heterodimers is attributable to
Ancα's weak homodimerization affinity; the excess of unbound Ancα subunits causes Ancβ
subunits to preferentially heterodimerize rather than homodimerize at equilibrium, even though
Ancβ's homodimerization affinity is slightly stronger than its heterodimerization affinity (Fig. 3D).
As protein concentration increases, these dimers begin to assemble with each other across IF2
into tetramers. The excess of heterodimers over homodimers means that the vast majority of
the tetramers are heterotetramers, even though IF2 itself does not distinguish between subunit
types. At physiologically relevant concentrations of 3mM total Hb subunits (34), the population is
dominated by $\alpha_2\beta_2$ heterotetramers, with a small fraction of heterodimers and virtually no
homotetramers (Fig. 3d; right panel).

Taken together, these data establish that the measured specificity of IF1 alone mediates highly
specific assembly of Ancα+ Ancβ into heterotetramers, even when IF2 is entirely nonspecific --

which our previous experiments suggest is the case – because IF1 is a much stronger interface
than IF2. The historical acquisition of heterospecificity across IF1 after the Ancαβ gene
duplication is therefore sufficient to account for the evolution of Hb's heterotetrameric
architecture.

Heteromeric specificity evolved primarily by reducing homodimerization affinity of Ancα.
Given our finding that heterospecificity evolved at the IF1 interface, we next sought to
characterize whether the acquisition of specificity was driven by evolutionary changes in the α
subunit, the β subunit, or both.

The heterospecificity of a pair of dimerizing proteins can be quantified in energetic terms as the
difference in the ΔG of binding between the heterodimer and the mean of the two homodimers
($\Delta\Delta G_{spec}$; see methods for calculation). If $\Delta\Delta G_{spec} = 0$, then the fractional occupancy of the
heterodimer at saturating and equal concentrations of subunits will be 50%, as will the sum of
the homodimers; this is true even if the homodimer ΔGs are very different from each other, as
long as the heterodimer ΔG is halfway between them. By contrast, if $\Delta\Delta G_{spec} < 0$, then
heterodimers will account for the majority of dimers; conversely, if $\Delta\Delta G_{spec} > 0$, homodimers
together will predominate (Fig. 4A-C). Hetero- or homospecificity thus arises when two paralogs
contribute nonadditively to dimerization. Whether or not the system is hetero- or homospecific,
the two homodimers will have equal occupancies to each other at saturating conditions:
irrespective of the fraction of subunits assembled into heterodimers, the remaining subunits will
be at equal concentrations and by definition will be well above the homodimerization Kds (28).

We quantified the heterospecificity of Ancα and Ancβ at IF1 by estimating $\Delta\Delta G_{spec}$. We used
nMS to measure the homodimer and heterodimer affinities of Ancα and Ancβ', which contains
all substitutions that occurred along the Ancβ branch except those that mediate tetramerization
across IF2, allowing us to prevent tetramerization and thus isolate the specificity effects at IF1.
From these affinities, we calculated the ΔG of binding and the expected fractional occupancy of
each dimer at high and equal concentration of subunits. For Ancα+Ancβ', we found that
$\Delta\Delta G_{spec} = -1.3$ (in units of kT) and heterodimer occupancy of 82% (Fig. 4D). This represents the
total specificity acquired by the two diverging paralogs after the duplication of of Ancαβ, which
by definition had no specificity. This specificity was acquired because of evolutionary changes in
all three relevant affinities. Relative to the ancestral dimerization affinity of Ancαβ, Ancα's
energy of homodimerization became worse ($\Delta\Delta G = 0.9$) while homodimerization by Ancβ'
improved substantially ($\Delta\Delta G = -3.7$). The heterodimer affinity improved by $\Delta\Delta G = -2.7$,
substantially more than the average of the two homodimers, yielding the observed strong
preference for the heterodimer.

We next sought to isolate the contribution of the evolutionary changes that occurred along each
of the two branches. To measure the specificity acquired along the branch leading to Ancα, we
measured affinities and calculated $\Delta\Delta G_{spec}$ when Ancα is mixed with the deeper ancestor
Ancαβ. This pair of proteins is heterospecific, with $\Delta\Delta G_{spec} = -1.2$ (expected heterodimer
occupancy 76%). Changes in the α subunit alone therefore account for >90% of the total
$\Delta\Delta G_{spec}$ that was acquired by the entire Ancα+Ancβ' system. This specificity is acquired via a

2.6-fold reduction in Ancα's homodimerization affinity compared to the Ancαβ ancestor and a
1.8-fold improvement in heterodimer affinity (Fig. 4E; Fig. S7B & D).

To isolate the contribution to IF1 specificity of evolutionary changes that occurred along the
branch to Ancβ, we measured affinities when Ancβ' is mixed with Ancαβ. This pair of proteins is
weakly heterospecific, with $\Delta\Delta G_{spec}$= –0.3 and expected heterodimer occupancy of just 58%.
The specificity is weak because both the homodimer and heterodimer affinity improved, but the
deviation of the heterodimer from the average of the homodimers is very small (Fig. 4F; Fig.
S7A &C).

Finally, we assessed whether the evolutionary changes in the Hbα subunit and those in the Hbβ
subunit interacted with each other nonindependently. If the changes affect specificity entirely
independently, $\Delta\Delta G_{spec}$ should equal the sum of the $\Delta\Delta G_{spec}$ acquired on each of the two
branches (–1.2 + –0.3 = –1.5). The observed $\Delta\Delta G_{spec}$ = –1.3, suggesting a very weak negative
interaction between changes in the two subunits, which makes the complex slightly less
heterospecific than expected if the substitutions were independent (Fig. 4G).

Taken together, these data indicate that the IF1 specificity acquired by the derived complex
Ancα + Ancβ is primarily attributable to substitutions in the α subunit, with substitutions in the β
subunit making a much smaller contribution; nonadditive interactions between the two sets of
changes did not contribute to the evolution of heterospecificity. The most important factor was
that Ancα became much worse at binding itself than at binding Ancβ. Ancβ, by contrast, became
slightly worse at binding Ancα than binding itself (Fig. 4G).

A one-residue deletion was the primary evolutionary cause of heterospecificity. We next
sought to identify the particular historical substitutions in Ancα that conferred this heteromeric
specificity on IF1. Only three sequence changes occurred on the branch from Ancαβ to Ancα: a
single-residue deletion of a histidine at site 3 (ΔH3), a five-residue deletion in helix D (ΔD), and
an amino acid replacement (v140A). ΔH3 is on the protein's N-terminal loop near IF1, and ΔD
directly contributes to the interface. Substitution v140A is biochemically conservative and far
away from the interface. The deletions are conserved to the present in Hbα subunits throughout
the jawed vertebrates, including humans, whereas the amino acid at site 140 varies (Fig. S1).
We therefore focused first on the effects of the deletions.

To isolate the contribution of each deletion to the evolution of specificity, we introduced each
one singly into Ancαβ and measured its effect on affinity and specificity when the mutant protein
is mixed with Ancαβ. We found that introducing ΔH3 alone confers substantial specificity,
recapitulating >80% of the Ancα's acquired heterospecificity for Ancαβ ($\Delta\Delta G_{spec}$ = –1.0 out of a
total $\Delta\Delta G_{spec}$ = –1.2 acquired along this branch) and >75% of the specificity acquired along both
branches by the entire Ancα+Ancβ' complex ($\Delta\Delta G_{spec}$ = –1.3, Figs. 5A, C). ΔH3 enhances
specificity by improving heterodimer affinity and reducing homodimer affinity, with both Kds very
close to those of Ancα (Fig. 5A; Fig. S10A & C).

The other deletion, ΔD, removes several residues that directly interact with the other subunit
across IF1, but introducing this change into Ancαβ had a much weaker effect on specificity
($\Delta\Delta G_{spec}$ = –0.4, Fig. 5B; Fig. S10B & D). When the contributions of ΔH3 and ΔD to specificity
are added together, they slightly exceed the specificity of Ancα (–1.4 rather than –1.3),
suggesting the possibility of a very weak negative epistatic interaction between them or a small
countervailing effect of the third change v104A. Taken together, these results indicate that ΔH3
was a large-effect historical sequence change that accounted for most of the specificity
historically acquired by the derived Hb complex.

Structural mechanisms for the gain in specificity. We next considered the structural
mechanisms by which ΔH3 conferred specificity by increasing heterodimer affinity and reducing
homodimer affinity. For a mutation to have these opposite effects, it must yield favorable
interactions when introduced into one side of the interface (in the heterodimer) but have
deleterious effects when introduced twice (in the homodimer). In principle, two kinds of
mechanisms could cause these opposite effects. Either 1) the mutated residue could interact
directly with the same residue on the other subunit favorably when one is in the derived state
but unfavorably when both are, or 2) the symmetry of the interface could be imperfect, such that
introducing the mutation on one side of the interface is favorable but introducing it again onto
the other side is net-unfavorable. The first scenario does not pertain in this case. Residue H3 is
part of the N-terminal loop, which does not participate directly in IF1 but instead packs against
helix H, which does contribute to IF1. But neither helix H nor the N-terminal loop contact the
same elements in the other subunit across the interface (Fig. 5D). Asymmetry in the interface is
therefore the likely of cause ΔH3's differential effects on heterodimer vs. homodimer affinity.

To gain insight into the possible nature of this asymmetry and potential mechanisms by which
ΔH3 affects specificity, we modeled the structures of the Ancαβ homodimer, the Ancαβ$_{\Delta H2}$
homodimer, and the heterodimer of these two proteins. The modeled Ancαβ homodimer
contains a subtle asymmetry: on one end of IF1, residue 130H on helix H sits close to 33R on
the opposite subunit, which allows a cross-interface hydrogen bond to form; on the other end of
the interface, the two residues are slightly further away from each other, leaving their hydrogen-
bonding potential unsatisfied when bound (Fig. 5E & F)**.** In the heterodimer, deleting His2 from
one subunit repairs this unfavorable interaction. Specifically, the deletion shortens the N-
terminal loop and changes its packing interaction against helix H, which causes helix H to slide
along the interface by ~1 Å compared to its position in the unmutated Ancαβ homodimer (Figs.
5D, 5G). 130H moves closer to 37T on the other subunit, allowing it to form a new hydrogen
bond across the interface, and several other interactions across the interface are also
enhanced. On the other end of the isologous interactions, the favorable interactions found in the
homodimer remain intact. This provides a potential structural explanation for how ΔH3 improves
heterodimer affinity (Figs. 5D, G).

The modeled Ancαβ$_{\Delta H3}$ homodimer structure is notably asymmetric and suggests a possible
mechanism by which introducing ΔH3 into both subunits reduces affinity (Fig. 5H). One side
displays the favorable new cross-interface interactions caused by ΔH3 in the heterodimer,

including the 130H-37T hydrogen bond. On the other side, however, the effect of the deletion is
very different: ΔH3 again causes helix H to slide along the interface, but on this side the
movement of 130H breaks the ancestral 130H-33R hydrogen bond, and 37T is also too far
away to interact favorably. This leaves the side chains of both 130H and 33R unsatisfied,
reducing homodimer affinity. In total, the homodimer of Ancαβ$_{ΔH3}$ contains three unsatisfied
hydrogen-bond donors/acceptors at these sites, whereas only one and two are unsatisfied in the
heterodimer and the ancestral homodimer, respectively.

These hypothesized mechanisms appear to have persisted over time. The same pattern of
interactions are found in the modeled structures of the hetero- and homodimers of Ancα + Ancβ
(Fig. S11). High-resolution crystal structures of extant hemoglobin also show notable
asymmetries in the multimerization interfaces, which exceed the deviation expected given the
resolution of the structures (35). These structures include some of the particular asymmetrical
interactions observed in our ancestral models: in the human Hb heterotetramer, 33R hydrogen
bonds across IF1 to residue 130, but this interaction is again lacking in the homodimer of human
Hbα, leaving 33R unsatisfied, potentially explaining the weak homomeric affinity of Hbα (Fig.
S11). At least some of the mechanisms of heterodimer specificity suggested by the structural
models of the ancestral proteins are therefore present in the known structures of its present-day
descendants. Structural models are prone to error, and the asymmetries we observed are
subtle; further research will be required to definitively characterize potential asymmetries in the
ancestral multimers.

Multiple historical sets of substitutions could have conferred heterospecificity. If
specificity in an isologous interface can evolve simply by causing nonadditive impacts on the
binding energies of heterodimer and homodimers, then there should be many mutations that
have the potential to make the interface specific in one direction or another. Indeed, if the
interface's symmetry is imperfect, then most mutations that affect affinity should impart
specificity to some degree.

To test this hypothesis, we measured the effect on specificity of subsets of changes that
occurred along the Ancβ lineage, which the results above show had strong effects on affinity
when introduced all together. First, we tested the five substitutions that that occurred at the IF1
surface (Fig. 5E & 5F). We introduced these changes into Ancαβ (creating protein Ancαβ$_{IF1}$) and
measured affinity and specificity when this protein is mixed with Ancαβ. These substitutions
yield a highly heterospecific complex (ΔΔG$_{spec}$ = –2.2, heterodimer occupancy 90%, Fig. 6A; Fig.
S12A, C, & E). Unlike the Ancα substitutions, the Ancβ$_{IF1}$ substitutions confer heterospecificity
by improving both homodimer and heterodimer affinity, but they improve the latter by more than
the former.

Because Ancαβ$_{IF1}$ is specific in complex with Ancαβ, we wondered whether it would also be
specific with Ancα. We found that this complex is barely heterospecific (ΔΔG$_{spec}$ = –0.1, Fig. 6B),
implying that other substitutions on the branch leading to Ancβ but not on the interface must
have contributed to the evolution of specificity between Ancαβ$_{IF1}$ and Ancα. We therefore
introduced an additional set of five historical substitutions that occurred in Ancβ but one

structural layer away from IF1 (see ref. 17). This protein (Ancαβ$_{IF1+Adjacent}$) has strong
heterospecificity when mixed with Ancα ($\Delta\Delta G_{spec}$ = –2.0, heterodimer occupancy >85%, Fig. 6D;
Fig. S12B, D, & F), because these mutations together improve both heterodimer and
homodimer affinity, but with a larger improvement in the heterodimer. It is also moderately
heterospecific when mixed with Ancαβ ($\Delta\Delta G_{spec}$ = –0.9).

Finally, we tested the effect of the adjacent substitutions on their own and found that they confer
specificity when mixed with Ancαβ ($\Delta\Delta G_{spec}$ = –0.9). These mutations impart specificity by
causing almost identical changes in homo- and heterodimer affinity. They also confer some
heterospecificity when Ancαβ$_{Adjacent}$ is mixed with Ancα ($\Delta\Delta G_{spec}$ = –0.6, Fig. S13A-D).

There are therefore several distinct sets of substitutions that occurred during history, and which
can be sufficient to confer heterospecificity on their own (and in various combinations), and they
do so via distinct patterns of effects on affinity. This degeneracy of mechanisms for evolving
specificity arises because there are many ways in which the energy of binding can change
nonadditively between heterodimer and homodimer. In every case, heteromeric specificity
rather than preference for the homomer was the result.

DISCUSSION

This work provides a mechanistic history of the evolutionary transition from the ancestral Ancαβ
homodimer to the derived Hb heterotetramer (summarized in Fig. S1). Each transition was
driven by a very simple genetic mechanism: a single substitution at IF2 conferred high affinity
tetramerization, and a single amino acid deletion at IF1 conferred heteromeric specificity. These
two key sequence changes have remained conserved in the descendant Hbα and Hbβ subunits
of all extant jawed vertebrates (Fig. S1). These transitions were both facilitated by the isologous
architecture of Hb's two interfaces, which creates a propensity for mutation to produce high-
order complexes and heterospecificity.

Symmetry facilitated evolution of the tetrameric stoichiometry. We found that
tetramerization across IF2 was driven primarily by a single replacement to a bulky hydrophobic
amino acid (q40W). In biochemical studies of extant protein interfaces, much of the free energy
change in protein-protein binding is attributable to interactions of bulky hydrophobic residues
with hydrophobic surface indentations (36), and mutations to bulky hydrophobic amino acids
can drive assembly into high-order multimers (9,37-40). Similar substitutions during history may
have been driving mechanisms during the evolution not only of Hb but of other molecular
complexes, as well.
487
The majority of complexes assemble through isologous interfaces (41). It has been suggested
that this must imply that isology confers some selective benefit by improving protein function (1).
Our results suggest an alternative explanation. If mutations are much more likely to produce
isologous complexes than nonisologous ones, then isologous complexes will predominate in
nature, even if there is no systematic fitness difference between the two types of multimer. We
found that although IF2 is intrinsically weak and mutation q40W cannot confer dimerization on

its own, it can drive tetramerization if its effects are multiplied in an isologous higher-order
complex. By contrast, If the interfaces were non-isologous -- with q40W interacting with a
hydrophobic divot on some other surface of the facing subunit – then this favorable interaction
would appear only once, and it would be insufficient to substantially improve binding energy and
confer meaningful tetramer occupancy. By this explanation, isologous complexes are abundant
because they are easier to produce by mutation than head-to-tail multimers, not more likely to
be fixed by selection.
501
It has been observed that in high-order multimers, the interface with higher affinity usually
evolves before the lower-affinity interface(s) (42-45). Hb evolution displays this pattern, with the
stronger interface IF1 evolving before IF2 (17). It has been suggested that this pattern is
attributable to selection: by this hypothesis, selection favors evolutionary intermediates that
contain the high-affinity interface, because those containing only the low-affinity interface
assemble slowly and/or misassemble into anomalous complexes (42,43). Our work here
suggests a different explanation: it is easier for mutations to generate a new interface that
confers a higher stoichiometry if a strong interface is already present, because the affinity of the
new interface is multiplied by iteration in an isologous complex. By contrast, low-affinity
interfaces do not confer multimerization on their own, so if the low-affinity interface were to
evolve first, then the effects of mutations on the second interface would not be multiplied. In
complexes with multiple interfaces, the stronger interface tends to be older not because such
trajectories improve fitness but because mutation is more likely to build elaborate complexes in
this historical order.
516
One interface confers specificity on a higher-order multimer. Our experiments show that
evolutionary change at just one of Hb's interfaces was sufficient to confer specific assembly into
heterotetramers. Specificity at IF1 alone was sufficient to mediate the heterospecificity of the
tetramer because this interface is so much stronger than IF2: IF1 mediates the specific
assembly of heterodimers, which assemble into heterotetramers across IF2, even though IF2
itself confers little or no specificity.
523
The specificity of IF1 and the isology of the complex also explains the *trans* conformation of
Hb's quaternary structure, in which each Hbα subunit binds one Hbβ subunit across IF1 and a
different Hbβ across IF2. The alternative *cis* conformation -- in which Hbα is paired with an Hbα
(and Hbβ with Hbβ) across one of the interfaces – is never observed. Although IF2 imposes little
or no specificity, its isologous orientation necessarily means that the two IF1-mediated
heterodimers must be rotated 180° relative to each other, placing each Hbα across IF2 from the
Hbβ of the other heterodimer. In the *cis* conformation, the heterodimers would not be rotated
180° relative to each other, and all the favorable interactions that IF2 comprises would not form;
residue 40W, for example, would not face the hydrophobic divot on IF2 across the interface.
Given the heterospecificity of IF1, isology constrains the Hb tetramer to its *trans* $\alpha_2\beta_2$
architecture.
535

536   These observations suggest a simple and potentially general mechanism for the evolution of
537   specificity in the quaternary structures of high-order multimers. Specificity need not evolve at
538   every interface in the complex, especially if the interfaces are isologous. Rather, mutations need
539   only make the stronger interface specific to confer assembly into particular high-order
540   architectures.
541
542   **Symmetry allowed specificity to evolve in one subunit.** We found that a single genetic
543   change in one paralog – a one residue deletion in Ancα -- was sufficient to confer IF1's
544   heterospecificity. This result contrasts with prior studies of nonisologous complexes, in which
545   heterospecificity evolved because of genetic changes in both interacting subunits
546   (7,13,18,19,21,24-26).
547
This difference in historical genetic mechanism reflects the opportunities presented by the two
different types of multimeric architecture. In nonisologous complexes, a mutation in the "head"
of one duplicate gene will not be sufficient to distinguish between its own tail and that of its
paralog (unless it somehow changes the conformation of both distinct surfaces). In an isologous
complex like Hb, however, a change in one subunit can confer specificity, because it makes the
interface different between the heterodimer, the mutated homodimer, and the unmutated
homodimer.
555
Acquiring specificity in an isologous interface requires the mutation to nonadditively change the
affinity of the heterodimer relative to the homodimers. If the symmetry of such interfaces were
perfect, a mutation in one subunit would affect interactions across the interface identically on
each side of the interface, resulting in additive effects on affinity. Nonadditivity would arise only
if mutations affect sites that interact with each other across the rotated interface. This would
require either a mutation at the precise axis of rotational symmetry or multiple mutations at
several sites.
563
If the symmetry is imperfect, however, a single mutation can affect interactions differently when
it appears twice in the homodimer versus when it occurs once in the heterodimer. Imperfect
asymmetry could facilitate the evolution of specificity in many complexes. Virtually all isologous
interfaces contain subtle asymmetries (46). This imperfection arises for two reasons: perfect
symmetry is entropically unfavorable, and amino acids near the axis seldom face each other
with perfect symmetry, because each amino acid itself is asymmetrical, and this asymmetry
propagates elsewhere in the interface (46,47). Extant human hemoglobin is one of many
examples of isologous interfaces in which asymmetry is imperfect (35,48). Isologous interfaces
therefore provide a starting point for homo- or heterospecificity to be acquired by substitutions in
a single subunit.
574
Specificity evolved through a single mutation. We found that a single mutation – deletion of
residue His2 in the alpha subunit – conferred most of the heterospecificity of Ancα + Ancβ. This
simple mechanism was possible because only a small change in relative binding energy is

required to yield substantial changes in specificity. The IF1 of Ancα+Ancβ' mediates 80%

heterodimer occupancy at equal and saturating concentrations, but its specificity is quite

moderate $\Delta\Delta G_{spec}$ = –1.3). This difference in binding energy is less than that associated with a

typical hydrogen bond or burial of a large hydrophobic residue. The structural differences in

physical interactions across the homodimer vs. heterodimer interfaces in our modeled structures

could easily yield energetic differences of this magnitude, although the particular form of

asymmetry in ancestral hemoglobin complexes remains uncertain. Recent in silico work also

found that small differences in ΔG can cause large differences in occupancy between

homodimers and heterodimers (28).

Why do such subtle differences in energy have such large impacts on specificity? Mutations that

cause a modest deviation in binding energies can cause large changes in occupancy because

of the nonlinear Boltzmann relationship between these quantities (Fig. 4C). Moreover, specificity

is determined by the deviation from additivity in the heterodimer relative to the homodimers, so

small differences in the free energy of binding will propagate into even larger changes in

specificity. Because of this intrinsic sensitivity, we predict that the evolution of specificity in

paralogous complexes with symmetrical interfaces will often be attributable to one or a few

genetic changes with relatively small energetic effects and subtle structural mechanisms. That

specificity can evolve so easily also implies that paralog interference after gene duplication

(49,50) may often be easily resolved through one or a few mutations.

If specificity can be acquired by small deviations from energetic additivity in either direction, one

might expect that homomeric and heteromeric specificity would be equally likely to evolve. But

empirical observations suggest that heteromers evolve much more frequently after gene

duplication (12,13). Our findings suggest a plausible explanation for this pattern. The modeled

structures suggest that the critical mutation for conferring specificity on Hb does so because

imperfect asymmetry in the interface creates a kind of antagonistic pleiotropy: a favorable

interaction occurs when the mutation is introduced once in the heteromer, but it fails to produce

the same favorable contact and even disrupts a different favorable contact when introduced

again on the other side of the interface in the homomer. Heterospecificity will result whenever

asymmetry causes antagonistic pleiotropy like this, such that a favorable interaction can be

optimized when it is iterated once but not twice. In contrast, homomeric specificity requires a

mutation to be even more favorable the second time it is introduced on the other side of an

interface. For this to occur, imperfect symmetry must synergistically enhance the interactions

caused by the two iterations of the mutation in the homodimer. This scenario seems far less

likely than an antagonistic effect, because favorable interactions are constrained in many ways,

requiring fairly precise compatibility of polarity, size, angle, etc. The imperfect symmetry of

isologous interfaces may therefore create a mutational propensity that favors the evolution of

heteromeric over homomeric specificity.

Taken together, our observations contribute to a growing body of evidence that complex

multimeric complexes can evolve through simple genetic mechanisms (5, 15, 37, 39,51-55). In

Hb evolution, a single substitution in one of the duplicated genes was sufficient to cause a

doubling in stoichiometry from dimer to tetramer, and a single-residue deletion at one interface
in the other subunit was sufficient to confer strong preference for the $\alpha_2\beta_2$ heterotetrameric form.
Although other substitutions enhanced these effects, and others may have permitted or
entrenched them (5,56), our data indicate that discrete evolutionary increases in complexity can
occur by very short mutational paths from simpler ancestral forms. The major effects of these
small sequence changes was possible because they took place in the context of an isologous
complex, and it is likely that its symmetry was slightly imperfect. Many multimers share these
structural properties, so we predict that, when other multimeric complexes are studied in detail,
simple mechanisms will be found to have driven their historical elaboration.

MATERIALS AND METHODS

Sequence data, alignment, phylogeny, and ancestral sequence reconstruction. The reconstructed ancestral sequences used here are the same as those reported previously (17). Briefly, 177 amino acid sequences of hemoglobin and related paralogs were collected and aligned. The maximum likelihood (ML) phylogenetic tree was inferred using the AIC best-fit model, LG+G+F (57,58). The phylogeny was rooted using as outgroups neuroglobin and globin X, which are found in both deuterostomes and protostomes and diverged prior to the gene duplications that produced vertebrate myoglobin and the hemoglobin subunits. Ancestral sequence reconstruction was performed using the empirical Bayes method (62), given the alignment, ML phylogeny, ML branch lengths, and ML model parameters. Reconstructed ancestors that were used in this study have been deposited previously in GenBank (IDs MT079112, MT079113, MT079114, MT079115).

The historical mutations that we introduced into those ancestral proteins are the following. For the set *IF1-reverted*, all sites in IF1 that were substituted on the branch leading to Ancb are reverted to the ancestral state found in Ancab; the mutations introduced are V36t, Y38h, V115a, V119e, H130r, D134e. For the set *IF2-reverted*, all sites that were substituted in IF2 on the branch leading to Ancb are reverted to the ancestral state found in Ancab; the mutations introduced are T37v, W40q, R43t, H100r, E104h. For the set *IF1*, all sites at IF1 that were substituted between Ancab and Ancb are changed to the derived state found in Ancb; the mutations introduced are t37V, k58M, r107K, h130Q, d134Q4. For the set *Adjacent*, five sites adjacent to IF1 that were substituted between Ancab and Ancb are changed to the derived state found in Ancb; the mutations introduced are h47S, s60N, q62K, a96S, h97E. The set *IF1+Adjacent* is the union of the sets *IF1* and *Adjacent*. Deletion DD deletes residues a54, e55, a56, i57, and k58 from Ancab.

Recombinant protein expression. Coding sequences for reconstructed ancestral proteins were optimized for expression in *Escherichia coli* using IDT Codon Optimization and synthesized *de novo* as gBlocks (IDT). Coding sequences were cloned by Gibson assembly into vector pLIC (3) under control of a T7 polymerase promoter. For co-expression of Ancα+Ancβ, a polycistronic operon was constructed under control of a T7 promoter and separated by a spacer containing a stop codon and ribosome binding site, as described in (63).

BL21 (DE3) *Escherichia coli* cells (New England Biolabs) were heat-shock transformed and plated onto Luria broth (LB) containing 50 ug/mL carbenicillin. For the starter culture, a single colony was inoculated into 50 mL of LB with 1:1000 dilution of working-stock carbenicillin and grown overnight. 5 mL of the starter culture were inoculated into a larger 500-mL terrific broth (TB) mixture containing the appropriate antibiotic concentration. Cells were grown at 37° C and shaken at 225 rpm in an incubator until they reached an optical density at 600 nm of 0.6-0.8.

For expression of single globin proteins, 100 uM of isopropyl-β-D-1-thiogalactopyranoside (IPTG) and 25 mg/500 mL of hemin were added to each culture. Expression of single proteins in culture were done overnight at 22° C. Cells were collected by centrifugation at 4,000g and

671   stored at -80° C until protein purification. Coexpressed proteins were induced using 500 mM
672   IPTG expression with 25 mg/500 mL hemin for 4 hours at 37°C. Cells were collected by
673   centrifugation at 4,000g, immediately followed by purification.
674
675   Human hemoglobin was bought commercially (Sigma-Aldridge) and resuspended in PBS.
676
677   We attempted to co-express and purify Anc$\alpha\beta_{\Delta H3}$ in complex with Anc$\alpha\beta_{40W}$, but we were not
678   able to identify conditions at which the two species could be expressed and purified to near-
679   equal concentrations.
680
Protein purification by ion exchange. All singly expressed proteins (all ancestral globins
except Ancα+Ancβ) were purified using ion exchange chromatography. All buffers were vacuum
filtered through a 0.2 µM PFTE membrane (Omnipore). After expression, cells were
resuspended in 30 mL of 50 mM Tris-Base (pH 6.88). The resuspended cells were placed in a
10 mL falcon tube and lysed using a FB505 sonicator (1s on/off for three cycles, each 1 minute).
The lysate was saturated with CO, transferred to a 30 mL round bottom tube, and centrifuged at
20,000g for 60 minutes to separate supernatant from non-soluble cell debris. The supernatant
was collected and syringe-filtered using HPX Millex Durapore filters (Millipore) to further remove
debris. A HiTrap SP cation exchange (GE) column was attached to an FPLC system (Biorad)
and equilibrated in 50 mM Tris-Base (pH 6.88). The lysate was passed over the column. 50 mL
of 50 mM Trise-Base (pH 6.88) was run through the SP column to remove weakly bound non-
target soluble products. Elution of bound ancestral Hbs was performed with 100-mL gradient of
50mM Tris-Base 1 M NaCl (pH 6.88) buffer which was run through the column from 0% to
100%. 1.5 mL fractions were captured during the gradient process, all fractions containing red
eluant were put into an Amicon ultra-15 tube and concentrated by centrifugation at 4,000g to a
final volume of 1 mL. For additional purification, concentrated sample was injected into a HiPrep
16/60 Sephacryl S-100 HR size exclusion chromatography (SEC) column. The column was
equilibrated in phosphate buffered saline (PBS) at pH 7.4. Purified ancestral globins elute at
different volumes depending on the protein's complex stoichiometry: 48-52 for tetramers, 56-60
for dimers, and 65-67 for monomers. The purified proteins were concentrated as mentioned
above and then flash frozen with liquid nitrogen.
702
703   For experiments in which two proteins were singly expressed and purified and then mixed
704   together, expression and purification of each protein were performed as described above. The
705   concentration of each protein was then quantified using the Hemoglobin Assay Kit (Sigma).
706   Proteins were then mixed together at 50 µM each for nMS. This procedure was performed in
707   triplicate to assess technical error introduced by the quantification and mixing process.
708
Protein purification by zinc affinity chromatography. Coexpressed proteins Ancα + Ancβ
were purified using zinc-affinity chromatography, which was performed using a HisTrap metal
affinity column (GE) on a Biorad NGC Quest. Nickle ions were stripped from the column (buffer
100 mM EDTA, 100 mM NaCl, 20 mM TRIS, pH 8.0), followed by five column volumes of water.
To attach zinc to the column, 0.1 M ZnSO$_4$ was passed over until conductance was stable,
approximately 5 column volumes, followed by five column volumes of water. After expression,

cells were resuspended in a 50 mL lysis buffer (20 mM Tris, 150 mM Nacl, 10% glycerol (v/v),
1mM BME, 0.05% Tween-20, and 1 Roche Protease EDTA-free inhibitor tablet, pH 7.40),
sonicated as described above, and the lysate passed through the prepared column. To remove
non-specifically bound protein, the column was washed with 50 mL of lysis buffer. Bound protein
was then eluted across a gradient of imidazole concentrations (0 to 500 mM) in a total of 100
mL elution buffer (20 mM Tris, 150 mM NaCl, 500 mM imidazole, 10% glycerol, and 1 mM BME,
pH 7.4). 1 mL fractions were collected. The fraction corresponding to the second peak of UV
absorbance at 280 nm has a visible red color and was collected and concentrated as described
above. The concentrated solution was injected into a Biorad ENrich 650 10 x 300 columns for
additional purification and eluted in PBS buffer.

Size exclusion chromatography assay. For protein concentrations from 0 to 500 µM, size
exclusion chromatography was performed using a Superdex 75 increase 10/300 GL column
(GE) equilibrated in PBS, then injected with 250 µL of sample using a 2 mL injection loop on an
Biorad NGC Quest FPLC and monitored by absorbance at 280 nm. For proteins at
concentration 1 mM, a HiPrep 16/60 Sephacryl S-100 HR was equilibrated in PBS using an
AKTAprime FPLC, then injected with 1mL sample and monitored by absorbance at 280 nm.

Native Mass Spectrometry. Protein samples were buffer exchanged into 200mM ammonium
acetate using either a centrifugal buffer exchange device (Micro Bio-Spin P-6 Gel, Bio-Rad) or a
dialysis device (Slide-A-Lyzer MINI Dialysis Unit, 10000 MWCO, Thermo) prior to native MS
experiments. Samples were loaded into gold-coated glass capillaries made in-house and
introduced to Synapt G1 HDMS instrument (Waters corporation) equipped with a 32k RF
generator (30). The instrument was set to a source pressure of 5.47 mbar, capillary voltage of
1.75 kV, sampling cone voltage of 20 V, extractor cone voltage of 5.0 V, trap collision voltage of
10 V, collision gas (Argon) flow rate of 2 mL/min (2.65 x 10 -2 mbar), and T-wave settings
(velocity/height) for trap, IMS and transfer of 100 ms -1 /0.2 V, 300 ms -1 /16.0 V, and 100 ms -1
/10.0 V, respectively. The source temperature (70 °C) and trap bias (30 V) were optimized. Part
of the native MS experiments were conducted by Thermo Scientific Exactive Plus Orbitrap with
Extended Mass Range (EMR) with tuning as follow: source DC offset of 15 V, injection flatapole
DC to 13 V, inter flatapole lens to 5, bent flatapole DC to 4, transfer multipole DC to 3 and C
trap entrance lens to 0, trapping gas pressure to 5.0 with the CE to 10, spray voltage to 1.50 kV,
capillary temperature to 100 °C, maximum inject time to 100 ms. Mass spectra were acquired
with a setting of 8750 resolution, microscans set to 1 and averaging set to 100. Mass spectra
were deconvoluted using Unidec (65).

Calculating multimerization affinity of homdimers. To estimate Kd of the monomer-to-
homodimer transition of singly expressed proteins, we performed nMS at variable protein
concentrations (P_{tot}). The occupancy of each oligomeric state at each concentration was
calculated as the proportion of all globin subunits in that state, based on the summed areas
under the corresponding peaks in the native MS spectrum. The fraction of subunits assembled
into dimers (*Fd*) includes dimers and tetramers and is defined as

758
$$Fd = \frac{2x_d + 4x_t}{(x_m + 2x_d + 4x_t)},$$

760   where x_m, x_d, and x_t are the total signal intensities of all peaks corresponding to the

761   monomeric, dimeric and tetrameric stoichiometries, respectively. Nonlinear regression was used

762   to find the best-fit value of Kd of dimerization using the equation:

764
$$F_d = \frac{1}{P_{tot}} * \frac{(4P_{tot} + K_d) - \sqrt{(4P_{tot} + K_d)^2 - 16P_{tot}^2}}{4}$$

766   As an alternative model of homodimerization, we also used a version of the Hill-Langmuir

767   equation:

769
$$F_d = \frac{P_{tot}}{P_{tot} + K_d}$$

The Hill-Langmuir model, which is typically used for ligand-receptor binding, does not account

for depletion of free monomeric subunits as homodimerization takes place and is therefore not a

valid model in this case; we used it solely to determine the robustness of the estimated Kd to

the specific binding equation used.

Calculating multimerization affinity of homtetramers. To estimate the Kd of the homodimer–

homotetramer transition, the fraction of subunits assembled into tetramers is defined as

779
$$F_t = \frac{4x_t}{(2x_d + 4x_t)}.$$

781   The concentration of all dimers is defined as

783
$$P_d = F_d \times P_{tot}.$$

785   Nonlinear regression was then used to find the Kd of tetramerization using the equation:

787
$$F_t = \frac{1}{P_d} * \frac{(4P_d + K_d) - \sqrt{(4P_d + K_d)^2 - 16P_d^2}}{4}$$

Parameters were estimated using the curvefit script in the Scipy package (66). The 95%

confidence interval on the K_d was estimated as 1.96 times the estimated standard error.

Calculating multimerization affinity of heteromers. To determine the Kd of

heterodimerization, we used nMS to measure stoichiometries across a titration series in which

one protein's concentration was held constant at 50 μM and the other was added at variable

concentration (1 to 50 μM). From the nMS spectrum, we estimated the proportion of the

heterodimer and the two homodimers as

$$F_{\alpha\alpha} = \frac{2x_{\alpha\alpha}}{\left(2x_{\alpha\alpha} + 2x_{\alpha\beta} + 2x_{\beta\beta} + x_\alpha + x_\beta\right)}$$

799
$$F_{\alpha\beta} = \frac{2x_{\alpha\beta}}{\left(2x_{\alpha\alpha} + 2x_{\alpha\beta} + 2x_{\beta\beta} + x_\alpha + x_\beta\right)}$$

800
$$F_{\beta\beta} = \frac{2x_{\beta\beta}}{\left(2x_{\alpha\alpha} + 2x_{\alpha\beta} + 2x_{\beta\beta} + x_\alpha + x_\beta\right)}$$

802   where each *x* represents the signal intensity of all peaks corresponding to the species denoted

803   in the subscript. The dissociation constant for each dimer is defined as $Kd_1 = \frac{x_\alpha^2}{x_{\alpha\alpha}}$, $Kd_2 = \frac{x_\beta^2}{x_{\beta\beta}}$,

804   and $Kd_3 = \frac{x_\alpha x_\beta}{x_{\alpha\beta}}$. By substitution, $F_{\alpha\beta}$ can be expressed as

806
$$F_{\alpha\beta} = \frac{\sqrt{Kd_1 * Kd_2 * F_{\alpha\alpha} * F_{\beta\beta}}}{Kd_3}$$

Kd_3 was estimated using this equation by nonlinear regression, where $F_{\alpha\alpha}, F_{\alpha\beta}$ and $F_{\beta\beta}$ were

measured using the titration series, and the affinities Kd_1 and Kd_2 were assigned the values

estimated in the homodimerization experiments described above.

Prediction of homodimer and heterodimer occupancy at high concentrations. The

occupancy of each dimer at physiologically relevant concentrations (1 mM total globin subunits)

was predicted as follows, because nMS is limited to concentrations <100μM. In a mixture of two

types of globins *A* and *B*, the total concentration of each subunit can be expressed in terms of

the concentration of monomers [A] and [B] in the mixture:

818
$$[A]_{tot} = [A] + [AB] + 2[AA] = [A] + \frac{[A][B]}{Kd_3} + \frac{2[A]^2}{Kd_1}$$

819
$$[B]_{tot} = [B] + [AB] + 2[BB] = [B] + \frac{[A][B]}{Kd_3} + \frac{2[B]^2}{Kd_2}$$

We used these equations to predict [A] and [B] at any value of C_A and C_B given the

experimentally estimated Kds. The concentration of each dimer was then estimated using the

equations $[AA] = \frac{[A]^2}{Kd_1}$, $[BB] = \frac{2[A][B]}{Kd_3}$, and $[BB] = \frac{[B]^2}{Kd_2}$.

Establishing the upper limit of IF2 Kd. We estimated the minimum Kd of assembly across IF2

by Ancαβ 37V+40W; IF1 removed, because no homotetramer was observed using nMS at a protein

concentration of 20 μM. The minimum detection limit for dimers in the nMS assay is 1 μM. Kd is

defined as $Kd = \frac{[M]^2}{[D]}$, where $[M]$ and $[D]$ are the equilibrium concentrations of monomer and
dimer, respectively. Therefore
830

$$Kd_{min} = \frac{(20 * 10^{-6})^2 M}{1 * 10^{-6} M} = 400 \, uM$$

832
833 **Determining ΔΔG of specificity.** Specificity for heterodimer assembly between two paralogs
834 can be defined as the difference between the additive affinity of the heterodimer and the
835 measured affinity of the heterodimer, using ΔGs derived measured dimerization affinity for two
836 homodimers and their respective heterodimer. The additive affinity of the heterodimer is defined
837 as the averaged ΔG of both homodimers:
838

$$\Delta G_{heterodimer}^{additive} = \frac{\Delta G_{homodimer\ 1} + \Delta G_{homodimer\ 2}}{2}$$

840
841 Specificity is then the difference between the additive and measured heterodimer ΔG.
842

$$\Delta\Delta G_{spec} = \Delta G_{heterodimer}^{measured} - \Delta G_{heterodimer}^{additive}$$

844
845 This metric is analogous to the coupling energy, which expresses the deviation of the measured
846 DG for a double mutant from that expected given the DGs of two single mutants assuming
847 additivity (67-69).
848
849 **Quantifying non-additive effect on specificity between Ancα and Ancβ.** The non-additive
850 effect on specificity can be defined as the difference between the predict and measured ΔΔG of
851 the derived complex Ancα + Ancβ.
852

$$\Delta\Delta\Delta G = \Delta\Delta G_{\alpha + \beta} - \left(\Delta\Delta G_\alpha + \Delta\Delta G_\beta\right).$$

854
855 **Prediction of monomer, dimer, and tetramer occupancies with no IF2 specificity.** The
856 occupancy of monomers, dimers, and tetramers between 1 mM and 4 mM predicted was
857 calculated as follows. The concentration of subunit in each stoichiometric species can be
858 expressed in terms of the concentration of monomers [A] and [B]:
859
860 $$[A]_{tot} = [A] + [AB] + 2[AA] + [ABBB] + 2[AABB]$$
861

$$= [A] + \frac{[A][B]}{Kd_3} + \frac{2[A]^2}{Kd_1} + \frac{\frac{[A][B]^3}{Kd_2 * Kd_3}}{Kd_4} + \frac{\frac{2[A]^2[B]^2}{Kd_2^2}}{Kd_4}$$

863
864
865 $$[B]_{tot} = [B] + [AB] + 2[BB] + 2[AABB] + 3[ABBB] + 4[BBBB]$$
866

867
$$= [B] + \frac{[A][B]}{Kd_3} + \frac{2[B]^2}{Kd_2} + \frac{\frac{2[A]^2[B]^2}{Kd_2^2}}{Kd_4} + \frac{\frac{3[A][B]^3}{Kd_2 * Kd_3}}{Kd_4} + \frac{\frac{4[B]^4}{Kd_3^2}}{Kd_4}$$

869 We used these equations to predict [A] and [B] across a range of $[A]_{tot}$ and $[B]_{tot}$ values given
870 previously measured equilibrium constants. Predicted [A] and [B] concentrations were used to
871 calculate the concentration of homodimers and heterodimers as described above, and the
872 concentration of tetramers were calculated using the following equations:

873
$$[BBBB] = \frac{\frac{[B]^4}{Kd_2^2}}{Kd_4}$$

875
$$[ABBB] = \frac{\frac{[A][B]^3}{Kd_2 * Kd_3}}{Kd_4}$$

877
$$[AABB] = \frac{\frac{[A]^2[B]^2}{Kd_3^2}}{Kd_4}$$

where $[BBBB]$ corresponds to the concentration of homotetramer, $[ABBB]$ is concentration of
$\alpha_1\beta_3$ tetramers, and $[AABB]$ is the concentration of $\alpha_2\beta_2$ heterotetramers.

Homology models. SWISS-Model was used to generate a structural model of the Ancαβ_{q40W}
homotetramer using the crystal structure of the human Hbβ homotetramer (PDB 1CMB) as
template, which was then refined using Rosetta's Fast Relax protocol, which energetically
minimizes the initial structure via small adjustments to the backbone and side chain torsion
angles (70). PyMOL V2.1 was used to visualize the proteins and capture images.

IF1-mediated homodimers were generated by the same procedure, except for homodimers of
Ancα or Ancαβ$_{\Delta D}$, for which the homodimer of human Hbα (PDB 3S48) was used as template.
IF1-mediated heterodimers were generated by the same procedure but using the
heterotetramer of human Hb (PDB 4HHB). For PyMol PSE file containing these models, see
Supplementary file.

Acknowledgements and funding sources
We thank members of the Thornton and Laganowsky labs for helpful advice and comments.
Supported by NIGMS R01GM131128, R35 GM14533601, 1T32GM13978201.

Author Contributions: CRCR: project conception and design, protein expression and
purification, chromatography experiments, structural modeling, data analysis, and writing (initial
draft and revisions). JL: design, execution, and analysis of mass spectrometry experiments, and
writing (revisions). AP: project conception and design, protein expression and purification,
chromatography experiments, data analysis, writing (initial draft and revisions). AL: design and

analysis of mass spectrometry experiments, writing (revisions). JWT: project conception and

design, data analysis, project management, funding acquisition, writing (initial draft and

revisions).

Citations

1. D. S. Goodsell, A. J. Olson, Structural symmetry and protein function. *Annual Review of Biophysics and Biomolecular Structure* **29**, 105-153 (2000).
2. J. A. Marsh, S. A. Teichmann, Structure, dynamics, assembly, and evolution of protein complexes. *Annu Rev Biochem* **84**, 551-575 (2015).
3. M. Lynch, The Evolution of multimeric protein assemblages. *Molecular Biology and Evolution* **29**, 1353-13 (2012).
4. M. Lynch, Evolutionary diversification of the multimeric states of proteins. *Proceedings of the National Academy of Sciences* **110**, E2821-E2828 (2013).
5. A. S. Pillai, G. K. A. Hochberg, J. W. Thornton, Simple mechanisms for the evolution of protein complexity. *Protein Sci* **31**, e4449 (2022).
6. G. K. A. Hochberg *et al.*, Structural principles that enable oligomeric small heat-shock protein paralogs to evolve distinct functions. *Science* **359**, 930-935 (2018).
7. E. J. Capra, B. S. Perchuk, J. M. Skerker, M. T. Laub, Adaptive mutations that prevent crosstalk enable the expansion of paralogous signaling protein families. *Cell* **150**, 222-232 (2012).
8. S. E. Ahnert, J. A. Marsh, H. Hernández, C. V. Robinson, S. A. Teichmann, Principles of assembly reveal a periodic table of protein complexes. *Science* **350**, 245 (2015).
9. H. Garcia-Seisdedos, C. Empereur-Mot, N. Elad, E. D. Levy, Proteins evolve on the edge of supramolecular self-assembly. *Nature* **548**, 244-247 (2017).
10. G. Diss *et al.*, Gene duplication can impart fragility, not robustness, in the yeast protein interaction network. *Science* **355**, 630-634 (2017).
11. S. Jones, J. M. Thornton, Principles of protein-protein interactions. *Proc Natl Acad Sci U S A* **93**, 13-20 (1996).
12. S. Mallik, D. S. Tawfik, Determining the interaction status and evolutionary fate of duplicated homomeric proteins. *PLOS Computational Biology* **16**, e1008145 (2020).
13. A. Marchant *et al.*, The role of structural pleiotropy and regulatory evolution in the retention of heteromers of paralogs. *Elife* **8** (2019).
14. J. B. Pereira-Leal, E. D. Levy, C. Kamp, S. A. Teichmann, Evolution of protein complexes by duplication of homomeric interactions. *Genome Biol* **8**, R51 (2007).
15. D. Grueninger *et al.*, Designed protein-protein association. *Science* **319**, 206-209 (2008).
16. J. Monod, J. Wyman, J.-P. Changeux, On the nature of allosteric transitions: a plausible model. *J Mol Biol* **12**, 88-118 (1965).
17. A. S. Pillai *et al.*, Origin of complexity in haemoglobin evolution. *Nature* **581**, 480-485 (2020).
18. G. C. Finnigan, V. Hanson-Smith, T. H. Stevens, J. W. Thornton, Evolution of increased complexity in a molecular machine. *Nature* **481**, 360-364 (2012).
19. J. R. EmLaw *et al.*, A single historical substitution drives an increase in acetylcholine receptor complexity. *Proc Natl Acad Sci U S A* **118** (2021).
20. O. Ashenberg, K. Rozen-Gagnon, M. T. Laub, A. E. Keating, Determinants of homodimerization specificity in histidine kinases. *J Mol Biol* **413**, 222-235 (2011).
21. D. Ding *et al.*, Co-evolution of interacting proteins through non-contacting and non-specific mutations. *Nat Ecol Evol* **6**, 590-603 (2022).

22. D. A. Ghose, K. E. Przydzial, E. M. Mahoney, A. E. Keating, M. T. Laub, Marginal
specificity in protein interactions constrains evolution of a paralogous family. *Proc Natl*
Acad Sci U S A **120**, e2221163120 (2023).

23. C. J. McClune, A. Alvarez-Buylla, C. A. Voigt, M. T. Laub, Engineering orthogonal
signalling pathways

24. C. D. Aakre *et al.*, Evolving new protein-protein interaction specificity through
promiscuous intermediates. *Cell* **163**, 594-606 (2015).

25. T. V. Lite *et al.*, Uncovering the basis of protein-protein interaction specificity with a
combinatorially complete library. *Elife* **9** (2020).

26. I. Nocedal, M. T. Laub, Ancestral reconstruction of duplicated signaling proteins reveals
the evolution of signaling specificity. *eLife* **11**, e77346 (2022).

27. G. K. A. Hochberg, J. W. Thornton, Reconstructing ancient proteins to understand the
causes of structure and function. *Annual Review of Biophysics* **46**, 247-269 (2017).

28. A. F. Cisneros, L. Nielly-Thibault, S. Mallik, E. D. Levy, C. R. Landry, Mutational biases
favor complexity increases in protein interaction networks after gene duplication. *Mol*
Syst Biol **20**, 549-572 (2024).

29. M. F. Perutz, H. Muirhead, J. M. Cox, L. C. G. Goaman, Three-dimensional fourier
synthesis of horse oxyhaemoglobin at 2.8 Å resolution: the atomic model. *Nature* **219**,
131-139 (1968).

30. A. C. Leney, A. J. R. Heck, Native mass spectrometry: what is in the name? *Journal of*
the American Society for Mass Spectrometry **28**, 5-13 (2016).

31. E. Stellwagen, "Chapter 23 gel filtration" in *Methods in Enzymology,* R. R. Burgess, M.
P. Deutscher, Eds. (Academic Press, 2009), **vol. 463**, pp. 373-385.

32. L. Kiger *et al.*, Thermodynamic studies on the equilibrium properties of a series of
recombinant betaW37 hemoglobin mutants. *Biochemistry* **37**, 4336-4345 (1998).

33. B. L. Boys, L. Konermann, Folding and Assembly of Hemoglobin Monitored by
Electrospray Mass Spectrometry Using an On-line Dialysis System. *Journal of the*
American Society for Mass Spectrometry **18**, 8-16 (2007).

34. G. Snyder, B. Sheafor, Red blood ells: centerpiece in the evolution of the vertebrate
circulatory system. *Am. Zool.* **39** (1999).

35. G. Fermi, M. F. Perutz, B. Shaanan, R. Fourme, The crystal structure of human
deoxyhaemoglobin at 1.74 Å resolution. *Journal of Molecular Biology* **175**, 159-174
(1984).

36. A. A. Bogan, K. S. Thorn, Anatomy of hot spots in protein interfaces. *J Mol Biol* **280**, 1-9
(1998).

37. J. Jee, I.-J. L. Byeon, J. M. Louis, A. M. Gronenborn, The point mutation A34F causes
dimerization of GB1. *Proteins: Structure, Function, and Bioinformatics* **71**, 1420-1431
(2008).

38. H. Garcia Seisdedos, T. Levin, G. Shapira, S. Freud, E. D. Levy, Mutant libraries reveal
negative design shielding proteins from supramolecular self-assembly and relocalization
in cells. *Proceedings of the National Academy of Sciences* **119**, e2101117119 (2022).

39. C.-S. Chen *et al.*, How to change the oligomeric state of a circular protein assembly:
switch from 11-subunit to 12-subunit TRAP suggests a general mechanism. *PLOS ONE*
6, e25296 (2011).
40. B. Kuhlman, J. W. O'Neill, D. E. Kim, K. Y. Zhang, D. Baker, Conversion of monomeric
protein L to an obligate dimer by computational protein design. *Proc Natl Acad Sci U S A*
98, 10687-10691 (2001).
41. H. Schweke *et al.*, An atlas of protein homo-oligomerization across domains of life. *Cell*
187, 999-1010.e1015 (2024).
42. J. A. Marsh *et al.*, Protein complexes are under evolutionary selection to assemble via
ordered pathways. *Cell* **153**, 461-470 (2013).
43. E. D. Levy, E. B. Erba, C. V. Robinson, S. A. Teichmann, Assembly reflects evolution of
protein complexes. *Nature* **453**, 1262-1265 (2008).
44. R. P. Bahadur, F. Rodier, J. Janin, A dissection of the protein–protein interfaces in
icosahedral virus capsids. *Journal of Molecular Biology* **367**, 574-590 (2007).
45. E. T. Powers, D. L. Powers, A perspective on mechanisms of protein tetramer formation.
Biophysical Journal **85**, 3587-3599 (2003).
46. J. T. Brennecke, B. L. de Groot, Quantifying asymmetry of multimeric proteins. *The*
Journal of Physical Chemistry A **122**, 7924-7930 (2018).
47. M. Bonjack-Shterengartz, D. Avnir, The near-symmetry of proteins. *Proteins: Structure,*
Function, and Bioinformatics **83**, 722-734 (2015).
48. M. F. Perutz, G. Fermi, B. Luisi, B. Shaanan, R. C. Liddington, Stereochemistry of
cooperative mechanisms in hemoglobin. *Accounts of Chemical Research* **20**, 309-321
(1987).
49. C. R. Baker, V. Hanson-Smith, A. D. Johnson, Following gene duplication, paralog
interference constrains transcriptional circuit evolution. *Science* **342**, 104-108 (2013).
50. J. T. Brigham, J. E. Brown, A. Rodríguez-Marí, J. M. Catchen, J. W. Thornton, Evolution
of a new Function by degenerative mutation in cephalochordate steroid receptors. *PLOS*
Genetics **4**, e1000191 (2008).
51. S. Liu et al., Nonnatural protein–protein interaction-pair design by key residues grafting.
Proceedings of the National Academy of Sciences **104**, 5330-5335 (2007).
52. G. Guntas, C. Purbeck, B. Kuhlman, Engineering a protein–protein interface using a
computationally designed library. *Proceedings of the National Academy of Sciences* **107**,
19296-19301 (2010).
53. D. P. Anderson et al., Evolution of an ancient protein function involved in organized
multicellularity in animals. *Elife* **5**, e10147 (2016).
54. L. Schulz et al., Evolution of increased complexity and specificity at the dawn of form I
Rubiscos. *Science* **378**, 155-160 (2022).
55. F. L. Sendker et al., Emergence of fractal geometries in the evolution of a metabolic
enzyme. *Nature* **628**, 894-900 (2024).
56. G. K. A. Hochberg *et al.*, A hydrophobic ratchet entrenches molecular complexes.
Nature **588**, 503-508 (2020).
57. D. Posada, T. R. Buckley, Model Selection and Model Averaging in Phylogenetics:
Advantages of Akaike Information Criterion and Bayesian Approaches Over Likelihood
Ratio Tests. *Systematic Biology* **53**, 793-808 (2004).

58. S. Q. Le, O. Gascuel, An improved general amino acid replacement matrix. *Mol Biol Evol*
25, 1307-1320 (2008).
59. C. Natarajan *et al.*, Expression and purification of recombinant hemoglobin in
Escherichia coli. *PLoS One* **6**, e20176 (2011).
60. M. T. Marty *et al.*, Bayesian deconvolution of mass and ion mobility spectra: from binary
interactions to polydisperse ensembles. *Analytical Chemistry* **87**, 4370-4376 (2015).
61. L. G. Nivón, R. Moretti, D. Baker, A Pareto-Optimal Refinement Method for Protein
Design Scaffolds. *PLOS ONE* **8**, e59004 (2013).
62.   Z. Yang, PAML: a program package for phylogenetic analysis by maximum likelihood.
Comput Appl Biosci **13**, 555-556 (1997).
63. J. Yang, Z. Zhang, X. A. Zhang, Q. Luo, A ligation-independent cloning method using
nicking DNA endonuclease. *Biotechniques* **49**, 817-821 (2010).
64. C. Natarajan *et al.*, Expression and purification of recombinant hemoglobin in
Escherichia coli. *PLoS One* **6**, e20176 (2011).
65. M. T. Marty *et al.*, Bayesian deconvolution of mass and ion mobility spectra: from binary
interactions to polydisperse ensembles. *Analytical Chemistry* **87**, 4370-4376 (2015).
66. P. Virtanen *et al.*, SciPy 1.0: fundamental algorithms for scientific computing in Python.
Nature Methods **17**, 261-272 (2020).
67. P. J. Carter, G. Winter, A. J. Wilkinson, A. R. Fersht, The use of double mutants to
detect structural changes in the active site of the tyrosyl-tRNA synthetase (Bacillus
stearothermophilus). *Cell* **38**, 835-840 (1984).
68. A. Horovitz, Double-mutant cycles: a powerful tool for analyzing protein structure and
function. *Folding and Design* **1**, R121-R126 (1996).
69. J. A. Wells, Additivity of mutational effects in proteins. *Biochemistry* **29**, 8509-8517
(1990).
70. L. G. Nivón, R. Moretti, D. Baker, A Pareto-Optimal Refinement Method for Protein
Design Scaffolds. *PLOS ONE* **8**, e59004 (2013).
1061

FIGURES

Figure 1. A single substitution confers tetramerization on an ancestral dimer. (A) A
substitution in one subunit can potentially affect specificity and stoichiometry in an isologous
interface. *Top*: After duplication of an isologous homodimer (gray), a substitution that occurs in
one paralog (red box) appears twice in the interface of a homodimer (red circles), once in a
heterodimer, and not at all in the other homodimer (blue). *Bottom*: One substitution (blue circle)
in an isologous interface appears twice in a homodimer (*left*), twice in a heterotetramer (*middle*),
and four times in a homotetramer (*right*), multiplying its effects on affinity. Dark and light gray,
paralogous subunits. **(B)** *Top*: Interfaces in the human Hb heterotetramer (PDB 4HHB). Pink,
Hbα; blue, Hbβ; α_1 and β_1 are in lighter hues than α_2 and β_2. IF1 surfaces (orange) mediate α_1-
β_1 and α_2-β_2 interactions; yellow surfaces (IF2) mediate α_1-β_2 and α_2-β_1 interactions. Only
interfaces involving α_1 are shown. Inset, α_1 subunit rotated away from the rest of the tetramer to
show IF1 and IF2. *Bottom*: Isology of IF1 and IF2. Helices contributing to each interface are
shown and labeled. Balls and sticks: on each helix, one residue's side chain is shown to
visualize symmetry. **(C)** Evolution of tetrameric stoichiometry on the phylogeny of Hb and
related globins. Icons, oligomeric states determined by experimental characterization of
reconstructed ancestral proteins (16). Acquisition of interfaces of IF1 and IF2 is shown (16). **(D)**
Key residues V37 and W40 that were substituted in Ancb. Cyan cartoon helix, b_1 subunit. Pink
and violet surfaces, a subunits that interact with b_1 via IF1 and IF2, respectively. Dotted lines to
red or blue spheres, hydrogen bonds to oxygen or nitrogen atoms, respectively (PDB 4HHB).
(E,G) Effect of historical substitutions on stoichiometry, as measured by size exclusion

chromatography. The ancestral dimer Ancαβ and the tetramers Ancα+Ancβ and human
hemoglobin (HsHb) are shown for comparison. Protein concentration at 100 mM (E) or 1 mM
(G). **(F)** Dimer-to-tetramer affinity of reconstructed ancestral Hb subunits containing historical
substitutions q40W and t37V, measured by native mass spectrometry across a titration series.
Points, fraction of dimers that are incorporated into tetramers. Lines, best-fit binding curves.
Estimated Kd and 95% confidence interval are shown. **(H)** Effect of historical substitutions on
monomer-dimer affinity measured by native MS. **(I)** Effect on dimer-tetramer affinity of
nonhistorical hydrophobic mutations in at residue 40, measured by native MS.

1092

1093

Figure 2. Multimerization across IF2 requires IF1. (A) IF1-mediated dimerization can be
compromised by mutations. Relative occupancy of each stoichiometry as measured by native
MS at at 20 mM total protein is shown for the ancestral dimer Ancαβ (top), Ancαβ$_{IF1 reverted}$
(middle, a variant of Ancαβ in which all IF1 residues are reverted to the ancestral state found in
AncMH), and Ancαβ-P127R (bottom, in which a mutation known to compromise IF1-mediated
dimerization has been introduced). (B) Compromising IF1 prevents assembly across IF2.
Relative occupancy of Ancαβ$_{40W + 37V}$ with and without mutations that compromise IF1-mediated
dimerization. (C) AncMH, which does not dimerize across IF1, cannot multimerize across IF2,
even when mutations sufficient to confer IF2-mediated mutimerization in Ancαβ are introduced.
(D) Observed (black) and expected (red) affinities of Ancαβ +q40W interfaces. Expected Kd of a
single iteration of IF2 (top) equals the square root of the measured apparent Kd when two
iterations are present (bottom). Expected apparent Kd of two iterations of IF1 (right) equals the
square of the measured Kd of a single IF1 (left).

1107

1108

Figure 3. Heterotetramer specificity is conferred by specificity at IF1. (A) Occupancy (as
fraction of all Hb subunits) when Ancα +Ancβ are coexpressed, measured by native MS. At 50
uM total protein, heterotetramers and heterodimers predominate (left). At 5 uM (right) – at which
assembly occurs only across the high-affinity interface (IF1) -- all dimers are heterodimers. **(B)**
Occupancy of subunits in stoichiometries as measured by nMS when Ancα and Ancβ are
separately expressed and then mixed at 50 mM each; IF2-mediated tetramer assembly does
not occur under these conditions, and dimers are predominantly heterodimers. Error bars
represent SEM for three replicates. **(C)** Percent occupancy of stoichiometries when Ancα and
Ancβ' (Ancβ with all derived IF2 surface residues reverted to the state in Ancαβ) are expressed
separately and then mixed at 50 uM. **(D)** Predicted occupancy of multimeric stoichiometries if
IF1 is specific and IF2 is nonspecific. Left: binding scheme with experimentally estimated Kds
(in mM) for IF1 and IF2-mediated multimerization by Ancα + Ancβ, assuming that all IF2 Kds are
equal (for Kds, see Fig. 4D and 1D). Right: expected occupancies of each monomer, dimer, and
tetramer, given the binding scheme at left. Occupancies are expressed as the fraction of all
subunits in each species.

1124

Figure 4. Contribution of historical changes in each subunit to the acquisition of heterospecificity. (A) Theoretical example of affinities and occupancy in a system of dimers with no specificity. *Top*: ΔG of dimerization for homodimers (XX and YY) and heterodimers (XY), in units of kT. In the absence of specificity, ΔG of the heterodimer equals the average of the homodimers (dotted line). *Bottom*: expected fractional occupancies of dimers at 1 mM per subunit and dissociation constants (Kd), given the ΔGs in the top panel. In the absence of specificity, heterodimer occupancy = 50%. **(B)** Example of a system with preference for the heterodimer. ΔΔG (the deviation of the heterodimer ΔG from the average of the homodimers) is shown. *Bottom*: Kd and predicted occupancy of each dimer at 1 mM. **(C)** Relationship between ΔΔG and heteromeric occupancy at 1 mM per subunit, assuming the ΔGs of homodimerization for as shown in panel A. **(D)** Specificity of IF1 dimerization in system of Ancα+Ancβ'. *Top*: expected fractional occupancies at 1 mM, given Kds assessed by nMS (shown above each bar, with 95% confidence interval). *Bottom*: ΔGs and ΔΔG given measured Kds. Dotted line, expected occupancies in the absence of specificity. **(E)** Specificity of IF1 acquired on the branch leading from Ancαβ to Ancα, shown as occupancy and ΔGs of the Ancαβ + Ancα system. The number of subunits that contain historical changes in each dimer is shown relative to the Ancαβ homodimer. **(F)** Specificity of IF1 acquired on the branch leading from Ancαβ to Ancβ, shown as occupancy and ΔGs of Ancαβ + Ancβ'. **(G)** Interaction effect on specificity when evolutionary changes leading from Ancαβ to Ancα (pink) and Ancβ' (blue) are combined. Homodimer of Ancαβ (gray) and each heterodimer are plotted by their ΔG. The observed ΔΔG of each heterodimer in combination Ancαβ is shown (see panels D-F). If the specificity acquired in the two subunits affects heterodimerization independently, then ΔΔG of Ancα+Ancβ will equal the sum of the ΔΔGs, yielding a parallelogram. The deviation from this expectation is shown.

1148

Figure 5. Effect of historical sequence changes on specificity. (A) Specificity of Ancaβ$_{ΔH3}$ with Ancaβ, as in Fig. 4. **(B)** Specificity of Ancaβ$_{ΔD}$ with Ancaβ. **(C)** Gain in specificity caused by various sets of historical mutations, relative to Ancaβ. Anca+Ancβ, all changes on both post-duplication branches. Anca, all changes on the branch leading to Anca. ΔH2 and ΔD, deletions that occurred on the Anca branches. **(D)** Models of Ancaβ homodimer and Ancaβ$_{ΔH3}$ + Ancaβ heterodimer. The N-terminal helix and the portion of IF1 involving helix H is shown. Grey surface, Ancaβ subunit common to both models. Grey cartoon, other Ancaβ subunit in the homodimer; pink cartoon, Ancaβ$_{ΔH3}$ subunit in the heterodimer. Yellow, 3H residue deleted in Ancaβ$_{ΔH3}$. Helix H side chains in the interface are shown as sticks. The hydrogen bond in the heterodimer from 130H to 37T (red surface) is shown (dotted line). **(E)** A portion of IF1 in the Ancaβ homodimer model, showing the isologous interactions with imperfect symmetry between 130H and 33R. Orange dashed-line, hydrogen bond. The two subunits are colored different shades of gray. The surface of the light-gray subunit is shown. **(F ,G, H)** Key residues in IF1 with hydrogen bonds that are affected by ΔH3 in the homodimers and heterodimer of Ancaβ and Ancaβ$_{ΔH3}$. Top, cartoon of key contacts. The two iterations of these interactions across the isologous interface are shown, one each in light or dark hue. Blue and red, nitrogen and oxygen atoms, respectively. Dotted lines, hydrogen bonds. The change in position of the H-helix caused by ΔH2 is shown. Bottom, structural alignment of the two iterations of the isologous interface in each dimer. Each dimer structure was duplicated exactly and then aligned to the original by targeting one subunit of the copy to align to the other subunit of the original. Hues correspond to the isologous iterations in the cartoon above.

1170

Figure 6. Other subsets of historical substitutions confer heterospecificity on IF1.
Affinities measured by nMS, predicted occupancy based on those Kds at 1 mM each subunit,
and $\Delta\Delta G_{spec}$ are shown for A) Ancaβ + Ancaβ_{IF1}, which contains the five substitutions at the IF1
surface that occurred in the Ancβ lineage; B) Ancaβ + Anca$\beta_{IF1\ +\ adjacent}$, which also includes 4
additional substitutions in Ancβ near but not on the interface; C) Ancα + Ancaβ_{IF1} , and D) Ancα
+ Anca$\beta_{IF1\ +\ adjacent}$.

1177

1178 **SUPPLEMENTARY FIGURES**

1179
1180
Fig. S1. Key sequence changes are conserved in extant hemoglobin subunits. A) Multimerization
state and residues at key sequence sites in extant Hb subunits from major jawed vertebrate taxa.
Myoglobin is the closest paralogous protein in the globin family. For complete alignment, see
https://tinyurl.com/yc6kvdb8. **B**) Alignment of extant human and reconstructed ancestral Hb subunits is
shown. Historical sequence changes that confer heterospecificity (ΔH3 in Hbα) and tetramerization
(q40W in Hbβ) are labeled and shown with arrows. Additional substitutions that occurred on the branch
leading to Ancβ and assayed in this paper are also labeled, including changes on the surface of IF1 (1),
IF2 (2), and at sites adjacent to IF1 (A). **C**) Alignment of N-terminal portion of Hb subunits from species
representative of major vertebrate taxa. The deletion of residue 3 in alpha subunits is marked. **D**)
Summary of key sequence changes that occurred after the duplication of Anc$\alpha\beta$. Multimerization states
of reconstructed ancestral proteins is shown.
1192

1193

1194
1195
1196 **Fig. S2. Effect of q40W tetramerization is robust to statistical uncertainty**. (A) Relative
1197 occupancy of monomer, dimer, and tetramer of Ancαβ$_{Alt. all}$, an alternative reconstruction of
1198 Ancαβ that contains the second most likely state at all ambiguously reconstructed sites,
1199 measured at 20 μM total protein using native MS. (B) Relative occupancy Ancαβ$_{Alt. all}$ with
1200 substitution q40W.
1201

1202

1203
1204 **Fig. S3. Native mass spectrometry spectra.** nMS spectra across a concentration series is
1205 shown for A) human Hb, B) Ancα + Ancβ, and (C) Ancαβ. Peaks corresponding to monomers,
1206 dimers, and tetramer are labeled.

Fig. S4. The effect of q40W on tetramerization depends on IF1. (A) Relative occupancy of Ancαβ$_{q40W}$, measured by native MS at 20 µM total protein. (B) Relative occupancy of Ancαβ$_{q40W\_IF1\text{-}reverted}$, which contains mutation q40W, as well as reversions to the ancestral state found in AncMH of all residues that were substituted between AncMH and Ancαβ.

Fig. S5. Heterodimer occupancy of Ancα and Ancβ is near equilibrium after mixing. (A)
The percent of all dimers that are heterodimers, measured by nMS when proteins are mixed at
50 μM each and allowed to incubate for 0, 1, 3, 6, or 24 hours. Black line and points, Ancα +
Ancβ (which only dimerize when expressed separately and then mixed). Grey line and points,
Ancα + Ancβ' (Ancβ in which IF2 surface substitutions are reverted to their ancestral state in
Ancαβ, thus preventing tetramerization). Each dot shows the mean of three replicates; error
bars, SEM. (B) Affinity of monomer-to-heterodimer assembly measured by nMS immediately
upon mixing of Ancα and Ancβ. Ancα was kept constant at 50 μM, while the concentration of
Ancβ varied. Points, fraction of all subunits in the mixture that are incorporated into
heterodimers. Line, best-fit binding curve. Estimated Kd and 95% confidence interval are
shown. (C) Estimated heterodimerization affinity measured as in panel B, but 1 hour after
mixing.
1231

1232

1233
1234
1235 **Fig. S6. Heterodimerization by Ancα+Ancβ'.** Monomer-to-heterodimer assembly measured
1236 by nMS. Ancα was kept constant at 50 μM while Ancβ' was at variable concentration. Points,
1237 fraction of all subunits in the mixture that are incorporated into heterodimers at each
1238 concentration. Line, best-fit binding curve. Estimated Kd and 95% confidence interval are
1239 shown.
1240

1241
Fig. S7. Dimerization by Ancα and Ancβ' (A-B) Homodimerization by Ancβ' (panel A) and by
Ancα (B). measured by nMS across a titration series. Each point shows the fraction of subunits
incorporated into dimers as the concentration of protein varied. Best-fit binding curve, Kd, and
95% confidence interval are shown. (C-D) Heterodimerization by mixtures of Ancαβ+ Ancβ (C)
and Ancαβ+Ancα and Ancα+Ancαβ (D). Each point shows the fraction of all subunits
incorporated into heterodimers. In each case, one protein was held constant at 50 μM while the
other was varied.
1249

1250

1251
1252
Fig. S8. Estimated affinity of the monomer-dimer transition by the Ancα homodimer is
robust to the binding model used. We fit two different binding equations (curves) to the
stoichiometries of **Ancα** measured using nMS. The Langmuir-Hill model (black line) assumes
that the concentration of free monomeric subunits is not depleted by dimerization. The
dimerization model used throughout the rest of this paper accounts for this depletion (described
in Materials and Methods). The estimated Kds (shown with their 95% confidence intervals) are
statistically indistinguishable.

1260

Fig. S9. Dimerization affinities and occupancy of Ancαβ. (A) Expected fractional

occupancies of homodimer and heterodimers when Ancαβ is mixed at equal concentrations with

Ancαβ$_{his}$ (500 μM each), given the measured dimerization affinities (shown above each column,

with 95% confidence interval). Ancαβ$_{his}$ is Ancαβ with an N-terminal polyhistidine tag, which

allows the masses of the three kinds of dimer to be distinguished. (B-C) Homodimerization by

Ancαβ$_{his}$ and heterodimerization by affinity of Ancαβ+ Ancαβ$_{his}$, measured and represented as in

Fig. S5.

1268

1269
1270 **Fig. S10. Effect of historical deletions on dimerization.** (A-B) Homodimerization and (C-D)
1271 Heterodimerization by mixtures, measured and represented as in Fig. S5.
1272
1273

1274
Fig. S11. Nonadditive interactions that contribute to specificity are conserved in derived Hb
complexes. In the modeled homodimers and heterodimers of Ancα+Ancβ (panels A, B) and X-ray crystal
structure of human hemoglobin (PDB 4HHB and 3S48), the figure shows the key IF1 residues with
nonadditive interactions in Ancαβ+Ancαβ$_{\Delta H3}$ (see Fig. 5G for comparison). Top, cartoon of key contacts.
The two iterations of these interactions across the isologous interface are shown, one each in light or dark
hue. Blue and red, nitrogen and oxygen atoms, respectively. Dotted lines, hydrogen bonds. Bottom,
structural alignment of the two iterations of the isologous interface in each dimer. Each dimer structure
was duplicated exactly and then aligned to the original by targeting one subunit of the copy to align to the
other subunit of the original. Hues correspond to the isologous iterations in the cartoon above

1284
1285
1286 **Fig. S12. Homodimerization by** Ancαβ$_{IF1}$ and Ancαβ$_{IF1 + Adjacent}$ (A,B) and heterodimerization by
1287 those proteins when mixed with Ancαβ (C,D) or Ancα (E,F). Measurements and representation
1288 as in Fig. S5.
1289
1290

1291
Fig. S13. Dimerization affinity and occupancies for Ancαβ$_{Adjacent}$. Expected fractional
occupancies of homodimer and heterodimers when Ancαβ$_{Adjacent}$ Is mixed with Ancαβ (A) or
Ancα (B), each at (500 μM), given the measured dimerization affinities (shown above each
column, with 95% confidence interval). Inset, ΔG of each dimerization (measured in units of kT),
with ΔG_{spec} of the heterodimer shown. (C,D,E) Measurement of binding affinities, measured and
represented as in Fig. S5.
1298