A novel method for integrating genomic and Tn-Seq data to identify common *in vivo* fitness mechanisms across multiple bacterial species

Derrick E. Fouts[1*], Thomas H. Clarke[1], Aric N. Brown[2], Elizabeth N. Ottosen[4], Caitlyn L. Holmes[2,3], Bridget S. Moricz[3], Sophia Mason[3], Ritam Sinha[4], Geoffrey B. Severin[2], Mark T. Anderson[2], Victor DiRita[4], Michael A. Bachman[3] and Harry L. T. Mobley[2*]

[1]The J. Craig Venter Institute, Rockville, MD, USA.

[2]Department of Microbiology and Immunology, University of Michigan Medical School

[3]Department of Pathology, University of Michigan Medical School

[4]Department of Microbiology and Molecular Genetics, Michigan State University College of Natural Sciences

*To whom correspondence should be addressed:

dfouts@jcvi.org

hmobley@umich.edu

Abstract

Sepsis is life-threatening organ dysfunction due to an unregulated immune response to infection. Bacteremia is a leading cause of sepsis, and members of the *Enterobacterales* cause nearly half of bacteremia cases annually. While previous Tn-Seq studies to identify novel bacteremia-fitness genes have provided valuable insight into virulence mechanisms, evidence for common pathways across species was lacking. To identify common fitness pathways in five bacteremia-caused *Enterobacterales* species, we utilized the JCVI pan-genome pipeline to integrate Tn-Seq fitness data with multiple available functional data types. Core genes from species pan-genomes were used to construct a multi-species core pan-genome, producing 2,850 core gene clusters found in four out of the five species. Integration of Tn-Seq fitness data enabled identification of 373 protein clusters that were conserved in all five species. A scoring rubric was applied to these clusters, which incorporated Tn-Seq fitness defects, operon localization, and antibiotic susceptibility data to identify seven common bacteremia-fitness pathways. Mutations in *tatC* showed reduced fitness *in vivo* and increased susceptibility to beta-lactams that were restored following *tatC* complementation *in trans*. By integrating known operon structures and antibiotic susceptibility with Tn-Seq fitness data, common genes within the core pan-genome emerged and revealed mechanisms that are essential for colonization of, or survival in, the mammalian bloodstream. Our prediction and validation of *tatC* as a common bacteremia fitness factor and contributor of antibiotic resistance supports the utility of this bioinformatic approach. This study represents a major step forward to identify novel targets of therapy against these deadly widespread sepsis infections.

Author Summary

Bacteremia is a leading cause of sepsis, a life-threatening condition where an unregulated immune response to infection causes systemic organ failure. Nearly half of bacteremia cases are caused by members of the Gram-negative bacterial taxonomic order *Enterobacterales*. Given the public health impact of bacteremia and the reduction of existing antibiotic treatment options, novel strategies are needed to combat these infections. Pan-genome software was used to predict seven shared fitness pathways in these bacteria that may serve as novel targets for treatment of bacteremia. Briefly, a scoring rubric was applied to shared pan-genome clusters, which incorporated multiple data types, including Tn-Seq fitness defects, operon localization, and antibiotic susceptibility data to rank and prioritize fitness genes. To provide an example of our predictions, mutations were constructed in one prioritized gene, *tatC*, which showed both reduced fitness in mice and increased susceptibility to *beta*-lactam antibiotics; complementation restored fitness and antibiotic susceptibility to wild type levels. This study takes a novel bioinformatics approach to build a core pan-genome across multiple distantly related bacteria to integrate computational and experimental data to predict important shared fitness genes and represents a major step forward toward identifying novel targets of therapy against these deadly widespread life-threatening infections.

Introduction

Sepsis is the life-threatening organ dysfunction resulting from a dysregulated and overwhelming immune response to infection. There are an estimated 49 million cases of sepsis that occur annually worldwide [1] and in the United States sepsis accounts for one in every 2-3 hospital deaths [2]. Bacteremia, or the presence of bacteria in the bloodstream, is the leading cause of sepsis with Gram-negative species belonging to the *Enterobacterales* order among the most common causes of bacteremia [3–5]. Drug resistance among this cohort of pathogens is a major limitation to effective treatment [4,6] and contributes to the large burden of antibiotic resistant bacterial threats recognized by the Centers for Disease Control and Prevention [7]. Given the public health impact of *Enterobacterales* bacteremia and the waning efficacy of existing treatments, novel strategies are needed to combat these infections. To develop such approaches, it is critical to first understand the genetic and molecular basis by which these bacteria cause bloodstream infections (BSI).

Bacteremia can be divided into three phases of pathogenesis: initial primary site infection, dissemination to the bloodstream, and growth and survival in blood and filtering organs [8]. The primary site can also serve as a reservoir for the pathogen that intermittently re-seeds the bloodstream and prolongs systemic infection. We have previously determined that *Enterobacterales* species replicate rapidly in the liver and spleen during bacteremia [9]. For some tissue and species combinations, bacterial burdens decrease over time reflecting an ability of the host immune system to overcome this rapid proliferation, while in other instances bacterial growth outpaces clearance and bacterial numbers rapidly expand. This balance between a bacterium's ability to proliferate and withstand immune-mediated clearance can be collectively described as infection fitness.

To define the genetic requirements for "bacteremia-fitness," we previously performed a series of

transposon sequencing (Tn-Seq) studies in which bacterial genes disrupted by transposon insertion

mutations are assessed for their contribution to bacteremia-fitness [10–13]. These studies have

resulted in the identification and characterization of fitness genes ranked by impact for four

Enterobacterales BSI pathogens; *Escherichia coli*, *Klebsiella pneumoniae*, *Citrobacter freundii*,

and *Serratia marcescens*. Fitness gene identification has also allowed us to build models of

metabolism and characterize bacterial pathways required for replication and survival in the host,

enabling opportunities for the development of novel therapeutic strategies against these antibiotic

resistant pathogens. While the Tn-Seq studies have provided valuable insight on an organismal

level, comprehensive information on shared and unique fitness strategies across Gram-negative

species is lacking, due in part to the unique nature and depth of each transposon insertion library.

By integrating the biological infection fitness data with comprehensive pan-genomic information

on gene content, operon structure, and gene function across species we can gain a better

understanding of infection processes for this important group of pathogens.

Our pan-genome pipeline [14] uses PanOCT [15] to cluster orthologous proteins to identify core,

variable and singleton gene or protein clusters and provides the data structure to easily link gene

functions such as GO terms and antimicrobial resistance (AMR) genes as well as linkage of genes

into genomic islands. The core genome of a bacterial pathogen is the collection of genes shared by

all or nearly all strains of a species. This core genome is generally enriched for critical functions

including energy production, amino acid metabolism, metabolite transport, nucleotide metabolism,

and translational machinery [16]. On the other hand, the accessory genome is the set of genes that

vary across strains of a species. These genes are often involved in protein secretion and defense

against innate host immunity, as well as many niche-specific functions that include traits such as

fimbriae, toxins, or iron acquisition systems, which reside in so-called "flexible genomic islands"

or fGIs [17].

In this report, we first constructed individual species pan-genomes to identify genes shared within

E. coli, K. pneumoniae, C. freundii, S. marcescens, and *Enterobacter hormaechei* genomes. Each

species pan-genome represents the sum of the core and accessory genes for all strains of a species

under study. To add biological context, known operon structures, virulence gene predictions, AMR

genes and Tn-Seq bacteremia-fitness data were overlaid onto each pan-genome, visually depicted

by using the program *PanACEA* [18]. Then we took each species' core genes to build a multi–

species pan-genome, which identified the core genes shared across these species (*i.e*., the core of

cores). By integrating our multi-species pan-genome and genome-wide Tn-Seq fitness data, we

were able to infer fitness genes in *E. hormaechei,* which lacked Tn-Seq data. We identified 373

protein clusters that were conserved in all five species and predicted to contribute to bacteremia in

at least one of them. Applying a scoring rubric to these bacteremia protein clusters, which

incorporated the magnitude of a cluster's fitness defect as predicted by Tn-Seq, its operon

localization, and published antibiotic susceptibility data, seven common bacteremia-fitness

pathways were identified. Finally, to validate our findings within the prioritized list of bacteremia-

fitness genes, the twin-arginine translocation (Tat) system was selected for validation of

independent mutations for reduced fitness *in vivo* and for increased susceptibility to beta-lactam

antibiotics *in vitro*. The TatC transmembrane protein strongly contributes to *C. freundii* bacteremia

fitness [11] and was likewise predicted to be a bacteremia-fitness factor in *K. pneumoniae* [13],

but not in the *S. marcescens [10]*, *E. coli* [12] or *E. hormaechei*, which lacked Tn-Seq data.

Independent insertional inactivation of *tatC* showed reduced fitness *in vivo*, confirming one of our

bioinformatics fitness predictions, even in the three species lacking Tn-Seq evidence. The *tatC*

mutants also showed increased susceptibility to beta-lactam and beta-lactam plus inhibitor that

was restored following *tatC* complementation *in trans*. This raises the exciting possibility that

inhibitors of these gene products could simultaneously decrease *in vivo* fitness and sensitize the

bacteria to FDA-approved standard-of-care antibiotics. Combined, this study provides an atlas of

gene conservation across the *Enterobacterales* and the fitness contribution of these genes to

bacteremia, a powerful resource that can be leveraged to advance our understanding of

pathogenesis and explore novel therapeutic targets.

Results

Species Pan-genomes. Protein-level pan-genomes of five common sepsis-causing

Enterobacterales species were constructed for two key reasons. First, to integrate multiple data

types and Tn-Seq datasets to decipher common bacteremia-fitness factors across species. Second,

to generate core pan-genomes that can be used downstream to build a "Multi-Species Core" (MSC)

pan-genome that represents highly conserved genes that may be targets of interest for future

therapies that would treat infections from multiple species (**Fig 1**). Proteins from representative

genomes of *E. coli*, *K. pneumoniae*, *S. marcescens, C. freundii,* and *E. hormaechei* were used to

first build species pan-genomes using PanOCT separately within the JCVI pan-genome pipeline

(**S1 Table, S1-5 Datasets**). Centroids (*i.e*., orthologous protein cluster representatives) of core and

accessory pan-genome clusters were searched against curated databases of bacteremia-fitness

genes as defined by transposon insertion-site sequencing (**S2 Table**), classic *E. coli* "essential"

genes (**S3 Table**), virulence genes [19] (**S4 Table**), and AMR [20] genes (**Table 1**).

Table 1. Pan-genome Statistics

Pan-genome	#Genomes	#Clusters	#Paralogs (#clusters; fitness genes@)	#Core proteins	#Accessory proteins	#Singleton proteins	# core pan-chrom asmbls (cycle; chain)‡	#fGIs (cycle; chain)	#Bacteremia-fitness@ (core; accessory)	#Essential^ (core; accessory)	#VFDB^ (core; accessory)	#AMR (core; accessory)
*C. freundii**	28	13,825	698 (2,743; 3)	3,997	5,211	4,617	4 (2; 2)	2,336 (7; 2,329)	172 (90; 82)	659 (630; 29)	213 (140; 73)	142 (66; 76)
*E. hormaechei**	58	13,713	770 (2,730; NA)	3,821	5,601	4,291	1 (1; 0)	2,971 (13; 2,958)	NA	605 (583; 22)	239 (130; 109)	140 (62; 78)
*E.coli/Shigella spp.**	281	37,889	3,279 (19,551; 57)	3,690	14,899	19,300	3 (1; 2)	13,438 (21; 13,417)	233 (117; 116)	755 (677; 78)	1,575 (158; 1475)	160 (67; 93)
*K. pneumoniae**	205	24,203	1,788 (8,006; 2)	4,453	4,970	14,780	1 (1; 0)	7,232 (20; 7,212)	56 (33; 23)	600 (573; 27)	492 (159; 358)	171 (68; 103)
*S. marcescens**	48	12,874	529 (1,705; 6)	4,077	4,984	3,813	3 (0; 3)	2,922 (6; 2,916)	201 (117; 84)	466 (459; 7)	192 (102; 90)	96 (56; 40)
Multi-species Core (70% cutoff)*	5	7,332	155 (35; 20)	2,850	1,369	3,113	49 (0; 49)	1,649 (0; 1,649)	500 (424; 76)	571 (502; 69)	272 (122; 150)	87 (38; 49)
Multi-species Core (100% cutoff)†	5	7,332	ND	2,232	5,100	3,113	ND	ND	373	455	55	27

*70% core definition; †100% core definition; ‡cycle = a circular pan-chromosome, chain = a linear pan-chromosome; @from Tn-Seq studies; ^Using

BLASTP matches with a percent id of >70% and e-value < 1x10^{-5}; NA = Not Applicable; ND = Not Determined.

The core genomes of each species varied from 3,690 protein clusters in *E. coli* to 4,453 in *K.*

pneumoniae. Based on 281 genomes, *E. coli* had the largest pan-genome, encoding 37,889 protein

clusters, which is 8 times the number of proteins encoded by the average *E. coli* genome (*i.e.*,

4,661 proteins) (**Table 1, S1 Table**). *K. pneumoniae* had the second largest pan-genome with

24,203 protein clusters across 205 genomes, but the number of proteins of this pan-genome was

4.5 times that of the average *K. pneumoniae* genome (*i.e.*, 5,322 proteins). The remaining species'

pan-genome number of proteins were between 2.7-3.0 times larger than their respective genome

averages. The number of strain-specific protein clusters from the species pan-genomes ranged

from a minimum of 0 to a maximum of 488 (*Escherichia*: 0 to 488, *Klebsiella*: 1 to 296,

Enterobacter: 0 to 280, *Citrobacter*: 4 to 449, and *Serratia*: 0 to 449) (**S1 Table**), indicating that

each species has a large and variable accessory genome.

Since our pan-genome software, PanOCT, does not collapse or discard paralogs, we were able to

determine the extent of paralogy within each species pan-genome. Of the 37,889 total *E. coli* pan-

genome protein clusters, 52% were paralogous, reducing the size of the pan-genome to 18,338

protein clusters, which is still considerably larger than the other species pan-genomes (**Table 1**).

Klebsiella had the second highest number of paralogs (33%) followed by *Enterobacter*/*Citrobacter*

(20% each) and *Serratia* (13%).

A total of 569 CARD/RGI-predicted AMR protein clusters were identified across all 4 species

with Tn-Seq data (*i.e.*, 160 in *E. coli*, 171 in *K. pneumoniae*, 142 in *C. freundii*, and 96 in *S.*

marcescens) (**Table 1**). The distribution of AMR genes across the core and accessory species pan-

genomes was roughly even, but with slightly more in the accessory pan-genome (45% core, 55%

accessory) except for *S. marcescens* (58% core, 42% accessory). In stark contrast, genes essential

for growth under laboratory conditions were skewed towards core (95% core, 5% accessory).

Predicted virulence protein clusters varied across species with *E. coli* and *K. pneumoniae* having

more clusters in the accessory pan-genome.

Fitness genes across species.

A total of 695 published bacteremia-fitness genes derived from four Tn-Seq studies [10–13] of *E.*

coli (247 genes), *K. pneumoniae* (74 genes), *C. freundii* (172 genes), *S. marcescens* (202 genes)

(**S2 Table)** were mapped to 662 species pan-genome clusters (**Table 1**). The distribution of

species-specific fitness gene clusters across the core and accessory pan-genomes showed slightly

more in the core pan-genome (54% core, 46% accessory), with *K. pneumoniae* (59% core, 41%

accessory) and *S. marcescens* (58% core, 42% accessory) each having more bias toward core

genes. This indicates that many genes important for causing bacteremia are highly conserved

within each species and may be conserved across species as well.

Multi-species Pan-genome.

To identify genes conserved across species and determine if they are multi-species bacteremia-

fitness factors, core protein sequences from each of the five species-level pan-genomes were

combined and analyzed in a second pan-genome run - the MSC pan-genome. For this MSC pan-

genome, clusters were labeled as core if they contained proteins from at least four out of five

species core pan-genomes. There were 2,232 clusters shared across all five species and 2,850

shared in four of five species (**Table 1,** 100% and 70% cutoff respectively**,** and **Fig 2A**). *Serratia*

had the largest number of unique core protein clusters within its species core genome at 1,124 with

Klebsiella having the second largest number of unique core clusters at 892 (**Fig 2A**). Despite

Escherichia having the most accessory genes (**Table 1**) in the species pan-genome, it only has 385

unique core protein clusters within the MSC pan-genome. All genera but *Serratia* shared 344

clusters, supporting the newly proposed family reclassification of *Serratia* out of the

Enterobacteriaceae family *[21].*

Identification of common bacteremia-fitness and virulence genes

Next, we analyzed the MSC pan-genome for bacteremia-fitness genes (**S5 Table, S6 Dataset**).

Using a criterion of identification by Tn-Seq in at least one species of the 7,332 proteins used in

the MSC pan-genome run, 500 were identified as core bacteremia-fitness factors conserved in 4

of 5 species and 373 were conserved across all species (**Table 1, Fig 2B**). These bacteremia-fitness

clusters included proteins from multiple species with published Tn-Seq results implicating them

as fitness factors. For example, 11 clusters contained Tn-Seq fitness factors across 3 species: 7

from *C. freundii*, *K. pneumoniae*, and *S. marcescens*, and 4 from *C. freundii*, *E. coli*, and *S.*

marcescens (**S6 Table**). Combining these pan-genome clusters with Tn-Seq data can be used to

predict potential fitness genes in species and strains that lack experimentally derived data.

MSC pan-genome clusters with homology to known virulence factors were identified by searching

against the virulence factor database (VFDB) [19]. Out of 272 total clusters assigned to virulence

functions (**Table 1**), 55 (20%) were conserved across all five genera (**Fig 2C**). The maximum

number of shared virulence factors between four, three and two species was 49

(*Citrobacter*:*Enterobacter*:*Escherichia*:*Serratia*), 12 (*Citrobacter*:*Enterobacter*:*Escherichia*) and

7 (*Citrobacter*:*Escherichia*), respectively. *K. pneumoniae* had the highest number of species-

specific virulence factors with 45. We then determined how many MSC pan-genome clusters

predicted to be bacteremia-fitness factors were also homologous to known virulence factors (**S7**

Table). Of the 500 bacteremia-fitness clusters and 272 virulence clusters conserved in 4 or more

species, 34 were predicted bacteremia-fitness factors associated with known virulence mechanisms

(*e.g.*, LPS and capsular production for immune modulation, flagella and fimbriae for motility and

adherence, and genes for the acquisition of iron and magnesium).

Using an orthologous approach to identifying multi-species fitness factors, we searched for core

operons that contained distinct predicted fitness genes across species. We identified 369 operons

that contained at least one bacteremia-fitness factor in the MSC pan-genome (**S8 Table**). Of the

67 operons that contained more than one bacteremia-fitness factor in the MSC pan-genome, 14

operons contained bacteremia-fitness genes from 3 species, 29 from 2 species, and 24 from one

species (**S8 Table**). One prominent example was the operon encoding the Enterobacterial common

antigen (ECA), a 12-gene locus with 9 bacteremia-fitness genes identified in either *S. marcescens*,

C. freundii, or *K. pneumoniae* (**Fig 3**). The WzxE ECA flippase within this operon is a predicted

fitness   factor   in   all   3   species   while   WecA   (UDP-*N*-acetylglucosamine-UP   *N*-

acetylglucosaminephosphotransferase)   and   RffH   (glucose-1-phosphate   thymidylyltransferase)

proteins were fitness factors in two species.

To prioritize genes for future investigation and to identify conserved biological processes

contributing to bacteremia we devised a scoring rubric using four genotypic and phenotypic

characteristics (see Materials and Methods for details). The MSC pan-genome bacteremia-fitness

genes were individually scored based on the magnitude of their fitness defect predicted in any

bacteremia Tn-Seq screen, whether the same gene was predicted to be a bacteremia-fitness factor

in multiple Tn-Seq screens (*i.e.*, multiple species), if multiple bacteremia-fitness factors were

encoded in the same operon, and whether mutation of that gene was previously found to increase

antibiotic susceptibility of *E. coli* BW25113 [22,23]. Scores were summed in two ways: first as

individual genes to distill the most meaningful bacteremia-fitness factors from the 500 total MSC

pan-genome bacteremia-fitness clusters identified, and then as the total score of bacteremia-fitness

factors encoded in the same operon to identify biological pathways of interest (**S9 Table**).

Application of this scoring rubric led to the identification of 32 and 73 individual bacteremia-

fitness factors having scores two and one standard deviations above the mean cluster score,

respectively (**S1 Fig**). Additionally, summation of the individual scores for bacteremia-fitness

factors encoded within the same operon led to the identification of 11 operons with total scores 0.5

standard deviations above the mean operon score. Investigation of the gene functions encoded by

these highly-scored individual bacteremia-fitness factors and operons revealed seven biological

processes that are predicted to contribute to the full virulence capacity of *Enterobacterales* during

bacteremia (**Table 2**). These processes include maintenance of the proton motive force, resistance

to antimicrobial peptides and complement, protein and small molecule transport, DNA repair and

homologous recombination, shikimate biosynthesis, global gene regulation (pre- and post-

transcription), and resistance to oxidative stress. While not an exhaustive accounting of all

biological processes that contribute to bacteremia or of the functions encoded within the top

scoring genes and operons, these seven biological functions (**Table 2**) include, at a minimum, 19

of the 32 (59%) fitness genes with scores two standard deviations above the mean and 40 of the

73 (55%) fitness genes with scores at least one standard deviation above the mean (**S1 Fig**).

Additionally, these seven biological functions encompass genetic elements encoded by 10 of the

11 highest scoring bacteremia-fitness factor rich operons identified.

Table 2. Core Biological Processes Predicted to be Required for Full Virulence

Capacity of *Enterobacterales* during Bacteremia using a Multifactorial

Prioritization Scoring Rubric*

Function	Gene(s)
1. Maintenance of Proton-Motive Force across the inner membrane	
ATP Synthase	*atpA‡B†D†EG‡H†*
Electron Transport: Ubiquinone synthesis	*ubiC†GH*
Iron-Sulfur Cluster biosynthesis for Cytochrome maturation	*iscS‡U†A†, hscB‡A†*
2. Resistance to Antimicrobial Peptides and Complement	
Modification of Lipid A of LPS	*arnAD‡E†F†T*
LPS inner core synthesis	*gmhB†*
Synthesis of Enterobacterial Common Antigen	*wecA(rfe)†, wzzE, wecB†, rffH(rfbA)‡, wecE(rffA)†, wzxE‡, wecF, wzyE, rffM*
Sensitivity to antimicrobial peptides	*sapA‡CDF*
Periplasmic protease inactivates complement complex	*prc†*
3. Transport	
Phosphate (ABC) transport	*pstSC‡A‡, phoU‡*
Twin arginine transporter	*tatC‡*
4. DNA repair and homologous recombination	
DNA recombination	*ruvA‡BC‡*
Tyrosine recombinase	*xerC‡D†*
5. Shikimate Biosynthesis	

Production of siderophores, aromatic amino acids, folates, and quinones (*e.g.*, *ubiGH*)	*aroACB†EK†*
6. Global gene regulation	
<u>A</u>erobic <u>r</u>espiration <u>c</u>ontrol protein	*arcAB†*
Nucleoid-binding regulator Fis	*fis†*
Nitrogen gene regulation sigma factor	*rpoN‡*
Glutamine synthetase and glutamine sensing	*glnA†D†*
mRNA regulation by binding sRNAs to target mRNAs	*proQ‡*
7. Oxidative stress	
Oxidative stress response	*apaGH, pdxA‡, sspA‡*

Genes listed here were predicted as a bacteremia-fitness factor in at least one Tn-Seq screen for four species (*E. coli,*

K. pneumoniae, C. freundii and *S. marcescens*) in a murine model of bacteremia and are predicted to contribute to

these seven biological processes. Genes were scored among all 500 MSC pan-genome bacteremia-fitness clusters in

the four species. Genes followed by (‡) or (†) received scores >2 or >1 standard deviation above the mean gene score,

respectively.

Validation of a representative core fitness gene function for virulence phenotype in a

bacteremia mouse model. In our Tn-Seq screens, numerous genes have been predicted to

represent fitness factors during experimental bacteremia. This is a first step in identifying relevant

genes that are critical for survival in the bloodstream. To validate our findings within the prioritized

MSC genes, the twin-arginine translocation (Tat) system was selected for further study. The Tat

system, under Transport functions (**Table 2**), facilitates translocation of folded proteins across the

cytoplasmic membrane in bacteria and while the system itself (TatABC) is widely conserved, the

translocated proteins that serve as substrates for the system often vary between species [24] and

may therefore alter the requirement for the translocation system in different organisms. The *tatC*

allele was one of the highest-scoring predicted bacteremia-fitness factors (**Table 2** and **S9 Table**).

The TatC transmembrane protein strongly contributes to *C. freundii* bacteremia fitness [11] and

was likewise predicted to be a bacteremia-fitness factor in *K. pneumoniae* [13]　(**S2 Table**).

However, neither the *S. marcescens [10]* nor *E. coli* [12] Tn-Seq findings identified TatC as a

significant fitness factor (**S2 Table**), perhaps due to lack of sufficient representation of *tatC*

mutants in the inoculum or the initial transposon library, or genuine species-specific differences

in Tat requirements.

To directly test whether TatC was a conserved bacteremia-fitness factor among the remaining four

species of interest, *tatC* mutants were constructed in *S. marcescens*, *K. pneumoniae, E. hormaechei*

and *E. coli* followed by competition infections whereby the wild-type strain and the differentially

marked *tatC* mutant strain were co-inoculated in a murine infection model. For all four tested

species, the *tatC* mutant was significantly outcompeted by the wild-type strain when bacteria were

inoculated into the bloodstream via tail vein injection (TVI), as indicated by competitive indices

that were significantly below zero (**Fig 4A-D**). Furthermore, *tatC* was required in every tested

organ via this model, demonstrating that folded protein translocation was universally important

not only across species but within different infection niches.

Since the TVI route represents direct bacteremia, we additionally sought to determine how TatC

contributes to disseminated infection via a bacteremic pneumonia model. Following

retropharyngeal inoculation (RI), *S. marcescens* and *K. pneumoniae tatC* mutants were

outcompeted in the lungs of infected mice (**Fig 4E-F**). For *S. marcescens*, an even greater

competitive disadvantage was observed for bacteria that had escaped the primary infection site and

disseminated to the spleen, liver, and kidneys (**Fig 4A and E**). This result was further supported

by a trend toward lower recovery of the *tatC* mutant from the bloodstream and suggests that Tat

translocation is critically important for escape from the lungs or survival in distal sites for this

organism. Similarly, *K. pneumoniae tatC* mutants that disseminated from the primary infection

site were also significantly outcompeted by wild-type bacteria (**Fig 4B and F**). Together, these

results establish the importance of the Tat translocation system, and very likely its substrate

proteins, in Gram-negative bacteremia and demonstrate the value of comparing Tn-Seq fitness hits

between species.

Contribution of conserved fitness genes to intrinsic antibiotic resistance. Previous work in *E.*

coli revealed that many core genes had contributions to antibiotic resistance that were not

predictable based on their known functions [22,23]. Genes that contribute to both bacteremia

fitness and antibiotic resistance could be attractive therapeutic targets as disruption of their

function in combination with antibiotic therapy would be expected to have synergistic efficacy.

Therefore, we mapped published data on the antibiotic susceptibility of a comprehensive panel of

E. coli mutants to multiple classes of antibiotics to our MSC pan-genome. *E. coli* mutants in *tatC*

were previously shown to have increased susceptibility to beta-lactam antibiotics. Unlike *E. coli*,

the species *K. pneumoniae*, *S. marcescens*, *E. hormaechei* and *C. freundii* have intrinsic resistance

to ampicillin due to chromosomally-encoded beta-lactamases. To test whether disruption of *tatC*

by either a transposon insertion (*tatC::tn*) or deletion (Δ*tatC::nptII*) could counteract this intrinsic

resistance, we used Epsilometer tests to determine the minimum inhibitory concentration (MIC)

of ampicillin or piperacillin-tazobactam for the wild-type and *tatC* mutants of these three species.

The most striking finding was in *K. pneumoniae*, where disruption of *tatC* by either approach

lowered the MIC to ≤ 8, which is a level that would be considered susceptible to ampicillin in

clinical care of an infection, and is similar to disruption of the beta-lactamase itself (*bla::tn*) (**Table**

3, Fig 5). This increased susceptibility to ampicillin in *K. pneumoniae* was counteracted by

provision of either *tatC* (pACYC*tatC*) or the *tatABCD* operon (pACYC*tatABCD*) on a multicopy

plasmid, but not the empty vector control (pACYCev). When *tatC* was interrupted in a

carbapenem-resistant isolate of *E. hormaechei*, the resistance profile shifted from resistant to

sensitive dose dependent (SDD) to piperacillin-tazobactam (**Table 3**). The resistance phenotype

of the *tatC* mutant was returned to parental levels when *tatABC* was provided on a plasmid. In the

case of *S. marcescens*, which has an extremely high resistance to ampicillin (> 256 µg/ml),

mutation of *tatC* caused a substantial reduction in MIC, which could be rescued by provision of

the *tatC* in trans from a multicopy plasmid (pBB26) and not the vector control (pBBR1MCS-5)

(**Table 3**). In *C. freundii*, we tested piperacillin-tazobactam as a beta-lactam and beta-lactamase

inhibitor combination without intrinsic resistance and found that a *tatC* mutation also lowered the

MIC and this phenotype was likewise rescued by provision of *tatC* (pBB4 (*tatC*+)) in trans and

not the vector control ((pBBR1MCS-5; **Table 3**). Combined with animal models of infection, these

data demonstrate that *tatC* represents a conserved bacteremia-fitness factor that also contributes to

intrinsic resistance to beta-lactam antibiotics in *Enterobacterales* and has the potential to restore

the use of antibiotics to which these species have intrinsic or acquired resistance.

Table 3. Susceptibility of *tatC* mutants to Beta-lactam antibiotics

Species (strain)	Genotype	Antibiotic	Mean MIC (µg/ml)	CLSI Interpretation
K. pneumoniae (KPPR1)	wild-type	ampicillin	24.0	I
	wild-type + pACYCev		18.7	I
	tatC::tn		8.0	S
	tatC::*nptII*		6.7	S
	ftn-e		3.0	S
	tatC::tn + pACYCev		6.7	S
	tatC::tn + pACYC*tatC*		14.7	I
	tatC::tn + pACYC*tatABCD*		18.7	I
S. marcescens (UMH9)	wild-type	ampicillin	>256	R
	tatC::*nptII* + pBBR1MCS-5		85.3	R
	tatC::*nptII* + pBB26 (*tatC+*)		>256	R
C. freundii (UMH14)	wild-type	piperacillin-tazobactam	6	S
	tatC::*nptII* + pBBR1MCS-5		0.75	S
	tatC::*nptII* + pBB4 (*tatC+*)		5.3	S
E. hormaechei (CRE14)	wild-type	piperacillin-tazobactam	96	R
	tatC::*acc(3)IV* + pBBR1Hyg		16	SDD
	tatC::*acc(3)IV* + pBBR1Hyg_*tatABC*		85.3	R

S=sensitive, SDD=sensitive dose dependent, I=intermediate, R=resistant

Discussion

In this report, we leveraged the JCVI pan-genome pipeline as a tool to integrate multiple disparate data types to identify key pathways and mechanisms used by *Enterobacterales* to survive in the bloodstream. Within the core genes (2,850/2,232, 70%/100% core, respectively) shared by *E. coli/Shigella spp., K. pneumoniae, S. marcescens*, *C. freundii,* and *E. hormaechei,* statistically significant bacteremia-fitness factors were identified in core clusters (424/373, 70%/100% core, respectively) containing at least one bacteremia-fitness factor identified in any of the four Tn-Seq studies performed in our TVI model of bacteremia. Using a multifactorial scoring rubric, we propose prioritization of 73 of the 500 total conserved bacteremia-fitness factors found in the MSC pan-genome, that have scores at least one standard deviation above the mean gene score, for future investigation. Of these 73 prioritized fitness factor genes, 36 were predicted by Tn-Seq to contribute to bacteremia in at least two species and 41 were found to increase antibiotic susceptibility in *E. coli* BW25113 when mutated [22,23]. Summation of individual gene scores encoded in the same operon revealed 11 operons enriched for bacteremia-fitness factors with scores 0.5 standard deviations above the mean. Investigation of the biological functions encoded by these operons and high scoring bacteremia fitness factors revealed a minimum of seven common biological functions that significantly contribute to *Enterobacterales* fitness during bacteremia. These data support a model in which full virulence of *Enterobacterales* during bacteremia in-part requires the seven biological functions outlined in **Table 2** and discussed below.

To aid in the discovery of potential novel antibiotic targets to clear bacteremia, we conducted species-level and family-level pan-genomic analyses that identified the core pathogenic genome

of these *Enterobacterales* species. The identification of unique and common virulence

determinants among these Gram-negative bacterial pathogens also provided substantial advances

toward determining mechanisms of pathogenesis for these pathogens that disseminate to and

survive in the bloodstream. By integrating the pan-*Enterobacterales* genome, genome-wide Tn-

Seq fitness data, and published lists of essential genes [25–29], antibiotic mutational susceptibility

[22], virulence [19] and operon data [30,31], we have identified 373 conserved fitness genes across

all five species that were shown to be important in at least one species during experimental

bacteremia. Inhibition of these bacteremia-fitness factors *in vivo* could allow the immune system

to clear infecting bacteria more rapidly or even kill them directly. Furthermore, 95 mutants in these

bacteremia-fitness genes also confer increased susceptibility to common antibiotics in a laboratory

strain of *E. coli [22,23]*. This raises the exciting possibility that inhibitors of these gene products

can simultaneously decrease *in vivo* fitness and sensitize the bacteria to FDA-approved standard-

of-care antibiotics.

The most serious consequence of bacteremia, sepsis, is characterized by an overwhelming innate

immune response to pathogen-associated molecular patterns (PAMPs) such as LPS,

peptidoglycan, flagella, and other macromolecules produced by pathogenic bacteria that

translocate into the bloodstream from the urinary tract, lung, wounds, intraperitoneal cavity, and

the gut. Interaction of PAMPs with Toll-like receptors triggers a well-understood cascade that can

lead to catastrophic events including coagulation, vascular permeability, and death [32]. In

contrast, bacterial factors that allow survival of bacteria in the bloodstream are poorly understood.

The literature has primarily focused on *E. coli* for Gram-negative pathogens and includes surveys

of isolates that identify their phylogenetic group and their "traditional" virulence factors assayed

by PCR of relevant genes. These studies implicate ExPEC strains of the B2 and D phylogroups

(although other groups are also found in lower proportions) [33,34]. Virulence factors associated

with bacteremic isolates include *papC, afa, sfa* (adhesins), *fyuA, iutA, iroN* (iron acquisition),

microcins (small bacteriocins), *cnf1* (cytotoxic necrotizing factor), and *hlyA* (hemolysin) [34–39].

Isolates are more often serum-resistant than control strains and represented by certain O-antigen

types [40,41].

Mechanistic studies for evaluation of specific genes in animal models of bacteremia have been

conducted only sparingly [42]. Investigations of fitness genes for other Gram-negative species in

bacteremia are rare [43,44]. In practice, characteristics of Gram-negative bacterial species causing

bacteremia have been hard to identify. However, Tn-Seq has become a powerful tool to

simultaneously assess the contribution of all non-essential genes in animal models of infection.

The technique, employed prior to conducting this project, provided a standard approach that we

have exploited. While there have been more than 200 published Tn-Seq studies dealing with

bacterial pathogens *in vitro* or *in vivo* (*e.g.*, [12,39,45–81]), only a fraction of these have addressed

Gram-negative   pathogens   in   animal   models   of   infection;   for   example:

[12,39,45,52,56,58,61,68,70,71,75,79,81–94]. Indeed, only a few studies [68,95,96], beyond our

work [10,12,13,83,97–99], have sought fitness and virulence factors in models of bacteremia.

Several conserved and shared *Enterobacterales* genes were identified, following integration of Tn-

Seq with operon structure and AMR, that fall into common bacterial cell functions of which we

discuss seven (**Table 2**). These included 1) maintenance of proton-motive force across the inner

membrane, 2) resistance to antimicrobial peptides and complement, 3) substrate transport, 4)

genome maintenance, 5) shikimate biosynthesis, 6) global gene regulation, and 7) oxidative stress.

While it is beyond the scope of this study to discuss all these findings, it is instructive to summarize

notable predicted bacteremia-fitness factors below in the context of the seven highlighted common

bacterial cell processes that support survival of *Enterobacterales* species in the mammalian

bloodstream.

Maintenance of proton-motive force. Tn-Seq studies of *Enterobacterales* species in the murine

model of bacteremia clearly implicate the importance of genes that support aerobic respiration and

maintenance of a proton-motive force. First, multiple hits in the operon encoding ATP Synthase

underscore the importance of this enzyme, central to the chemiosmotic theory, as critical for

infection. This function is clearly supported by electron transport which depends on synthesis of

electron transport chain cofactors such as ubiquinone that are embedded in the cytochromes. In

addition, this system is dependent on the biosynthesis of Fe-S clusters that carry electrons within

cytochromes as critical to maintenance of proton motive force. The ISC system is encoded by the

iscRSUA-hscBA-fdx-iscX operon and predominantly carries out the Fe-S cluster biosynthesis under

normal conditions used for cytochrome maturation and proton motive force. Indeed, iron-sulfur

(Fe-S) proteins contribute to several cellular processes that include respiration, nitrogen fixation,

and gene regulation. *E. coli* and closely related enterobacteria possess two pathways [100]. While

oxygen is tightly bound by hemoglobin within erythrocytes circulating in the blood, we

nevertheless presume that this milieu represents an aerobic environment, and that the bacterial

species undergo aerobic respiration and thus would require the activities of ATP synthase coupled

with the electron transport chain.

Resistance to antimicrobial peptides, complement, and phagocytosis. Enterobacterial

Common Antigen is a critical outer membrane component in *Enterobacterales* that may affect

resistance to antimicrobial peptides and complement [101]. Interestingly, while identified by Tn-

Seq in *K. pneumoniae*, *C. freundii*, and *S. marcescens* (**Fig 3**), none of the ECA genes were

identified by Tn-Seq as fitness factors in *E. coli* suggesting that either the biological role of ECA

during infection differs in this species or that ECA genes were not adequately targeted by

transposon mutagenesis in *E. coli*. While this discrepancy is currently under investigation, the clear

benefit of such a comparison is in forming testable hypotheses regarding the conserved function

of ECA or other shared operons across Gram-negative bacteremia species. Combined, our findings

provide strong evidence that production of ECA is an important fitness determinant since mutation

of even a single gene in the processive ECA biosynthesis pathway may result in a lack of ECA

production or its proper localization in the bacterium. Clearly, surface-exposed macromolecules

on the Gram-negative outer membrane, when present or absent or biochemically modified, may

modulate surface charge. These changes in charge can confer, in some cases, resistance to binding

by complement components and host-derived antimicrobial peptides, thus conferring protection

from these host innate immune responses.

Likewise, GmhB (YaeD) is a multi-species fitness factor involved in biosynthesis of the LPS inner

core. Specifically, GmhB is a D,D-heptose 1,7-bisphosphate phosphatase that is a member of the

ADP-heptose enzymatic cascade [102]. While ADP-heptose is a part of the LPS inner core, it is

also a soluble inflammatory mediator recognized by host cytosolic receptors [103,104]. GmhB has

been validated as a bacteremia fitness factor in *K. pneumoniae, E. coli,* and *C. freundii [13,105]*.

Interestingly, GmhB is dispensable for fitness in the lung but is required for spleen and liver fitness

during bacteremia. Thus, GmhB is likely linked to bacteremia fitness across species in a manner

unique to bloodstream infection. Indeed, exponential replication in the spleen at 24 h during

murine bacteremia [9] may require substantial LPS export. In the absence of GmhB, lower

abundance of normal LPS may be produced, leaving Gram-negative species more susceptible to

killing by phagocytosis by immune cells such as splenic macrophages.

Transport. Twin-arginine translocation system [106] represents perhaps the most critical

transport system and fitness factor overall required for bacteremia. Specifically, *tatC* allele was

demonstrated to make a significant contribution to the fitness of all four species tested in this study,

as well as *C. freundii* [11], in a murine model of bacteremia (**Fig 4**). While it is possible that

disruption of the TatC transmembrane component may have unintended consequences, such as

altering bacterial envelope integrity, the most likely explanation for this shared phenotype is a

requirement for appropriate localization of Tat-secreted proteins for infection fitness. Indeed, we

have previously attributed at least some of the fitness defect associated with *C. freundii tatC*

mutation to SufI loss of function [11]. Whether SufI is required for infection of these additional

species remains to be determined. However, it may also be informative in future work to compare

the repertoire of Tat-secreted proteins in each species using the conserved N-terminal twin-

arginine motif, since it is additionally possible that the requirement for TatC is due to species-

specific translocation of different proteins.

In contrast, the Pst phosphate transport system imports one critical nutrient and is also predicted

to contribute to bacterial survival in the bloodstream. The *pst* operon encodes a high-affinity inner

membrane ABC transport system of inorganic phosphate [107]. The system includes the permeases

PstA (C) and PstC (C, K), the regulator PhoU (C), the periplasmic phosphate binding protein PstS

(C), and the ATP-binding protein PstB (not a predicted bacteremia-fitness factor). Phosphate

concentrations are sensed by PhoBR, with PhoB as the histidine kinase and PhoR as the response

regulator. The Pst system is required for signal transduction in response to inorganic phosphate,

and a mutant in any of the genes causes constitutive expression of the Pho operon encoding alkaline

phosphatase. This leads to pleiotropic effects in the cell that attenuate virulence and lower the MIC

to polymyxin [108]. This includes reduction in the amount of capsule, decreased resistance to

serum, antimicrobial peptides, acid and oxidative stress, and decreased production of type 1

fimbriae. Under phosphate starvation, bacteria modify their phospholipids, including lipid A. Pst

mutants also lead to imbalance of cyclopropane and unsaturated fatty acids and an overall increase

in outer membrane permeability. Some components of the Pst system (*e.g.*, PhoA, PhoE) have

detectable antiserum in previously infected patients with hemolytic uremic syndrome. The Pst

system, clearly implicated as a requisite mechanism for phosphate import and predicted as a fitness

operon in two of the four species, could also represent an attractive target for novel therapeutics,

since blocking of any of the proteins could cause pleiotropic attenuation of virulence and fitness.

DNA repair and recombination. DNA recombinases XerC and RuvABC were predicted to be

critical fitness factors for *Enterobacterales* species. RuvABC enzyme complexes are involved in

DNA recombination and repair and facilitate Holliday junction branch migration and resolution

[109,110]. RuvABC repairs damage by catalyzing homologous exchanges between damaged and

undamaged DNA [111]. XerC responds to DNA damage mediated by reactive oxygen species

produced by neutrophils during infection and thus defends against the innate immune system in

the bloodstream. The recombinase is also required for induction of the SOS response, with the

result that a mutant defective in *xerC* is more susceptible to a range of DNA-damaging antibiotics

including ciprofloxacin and immune-mediated killing [112].

Shikimate biosynthesis. Iron acquisition is required by Gram-negative bacilli and all bacterial

species to survive in a host where free iron is found in negligible concentrations. While some iron

salt import systems contribute (*e.g.*, ferric citrate uptake [113]) to iron acquisition, the highest

affinity and effective iron acquisition systems include siderophores that bind iron with

extraordinarily high affinity. Genes that contribute to the synthesis of specific siderophores

including enterobactin and salmochelin provide the ability to survive in the bloodstream.

Shikimate biosynthesis [114], a seven-step enzymatic process that terminates with the production

of chorismate, is one such pathway that contributes to the biosynthesis of some siderophores, like

enterobactin [115]. Chorismate originating from the shikimate biosynthesis pathway is also a

fundamental precursor for the biosynthesis of folates, aromatic amino acids, and the quinones;

ubiquinone (mentioned above) and menaquinone, which contribute to aerobic- and anaerobic

respiration, respectively. Alleles encoding five of the seven enzymatic steps of the shikimate

biosynthesis pathway were predicted to contribute to bacterial fitness during bacteremia in at least

one Tn-Seq study, with two alleles, *aroB* and *aroK*, being scored in the top 73 prioritized

bacteremia-fitness factors (**S2 Table, S9 Table**). Strains carrying mutations within the shikimate

biosynthesis pathway likely experience a fitness defect in the host resulting from the cumulative

pleiotropic effects of folate deprivation, aromatic amino acid auxotrophy, loss of energy

production through respiration, and iron starvation. In agreement with its prediction as a

bacteremia-fitness pathway in our studies, the shikimate biosynthesis pathway has been shown to

play a fundamental role during infection for many bacterial pathogens in diverse disease models

[83,116–118].

Global Regulation. ArcA is a global gene regulator of aerobic respiration that contributes to the

maintenance of a proton motive force when oxygen is available. In a recent study [119], we

constructed deletion mutants in the aerobic respiration control sensor *arcA* across four species and

confirmed that ArcA is a bacteremia-fitness factor in *K. pneumoniae*, *C. freundii*, and *S.*

marcescens but not *E. coli*. ArcA is conserved across the *Enterobacterales* and clearly contributes

to fitness in a murine bacteremia model [119]. Because this regulator, along with ArcB, mediates

transition from aerobic to anaerobic growth, mutants in *arcA* do not grow well under anaerobic

conditions. In addition, *arcA* mutants grew more slowly than isogenic wild-type strains when

cultured under iron-limited conditions because of their inability to easily shift to anaerobic growth.

Clearly ArcA, in conjunction with ArcB, promotes aerobic growth during bloodstream infections

but also provides flexibility to shift to microaerobic or anaerobic growth when oxygen is more

limited in the portions of blood-filtering organs distal from the artery [120].

Oxidative stress. The *pdxA* operon contributes to the alleviation of oxidative stress. This multi-

functional operon contains six genes, *lptD-surA-pdxA-rsmA-apaGH*, five of which are predicted

bacteremia-fitness genes. The operon begins with *lptD*, an essential gene encoding a protein for

lipopolysaccharide assembly [121]. *surA* encodes a periplasmic ATP-independent chaperone

protein involved in the proper folding of outer membrane proteins, including LptD [122]. *pdxA*

encodes 4-Hydroxythreonine-4-phosphate dehydrogenase (UniProt P19624) as part of the

pyridoxal -5'-phosphate (Vitamin B6) synthesis pathway that is required for many steps in

biosynthesis and transformations of amino acids [121]. Whereas most prokaryotes have a simple

pathway encoded by *pxtST*, gammaproteobacteria have a complex pathway spread across multiple

operons. In contrast, Vitamin B6 is an essential nutrient for humans. The *rsmA* (a.k.a. *ksgA*) locus

encodes a (adenine-N6, N6-)-dimethyltransferase that methylates two adenines in the 16S

ribosomal subunit as part of ribosomal biogenesis [123]. While the *apaG* gene product has an

unknown function, the downstream *apaH* locus encodes a diadenosine tetraphosphatase (UniProt

P05637) that degrades diadenosine tetraphosphate (Ap4A). Under disulfide stress, Ap4A

accumulates as a by-product of amino-acyl tRNA synthetases and ApaH is deactivated. This leads

to capping of RNA transcripts with Ap4A and other dinucleotides with adenosine (Np4A) that

ApaH removes and targets the RNA for degradation. The overall function of this operon and the

effect of these genes on bacteremia fitness is not entirely clear. It may be that this operon is required

for fitness under stresses encountered *in vivo*, including oxidative/disulfide stress that would affect

periplasmic protein folding (*lptD, surA*) and induce the *apaH* system to regulate RNA stability and

amino acid usage (*pdxA*) in protein synthesis (*rsmA*). Whereas 5 of 6 genes of this operon are

bacteremia-fitness factors, only 3 of 7 genes involved in Vitamin B6 synthesis are fitness factors,

one of which (*serC* in *C. freundii*), is also needed for serine biosynthesis.

Conclusions. The use of our pan-genome pipeline as a powerful tool for integrating multiple data

types and sources to infer biological functions was demonstrated in this study. Comparison of the

four *Enterobacterales* Tn-Seq datasets revealed, at first inspection, limited common mutations and

functional insight. It wasn't until those results were mapped to orthologous protein clusters and

overlaid upon operon structure that common pathways emerged (**Table 2**). This enabled the

prediction of common genes within this core pan-genome and brought to light mechanisms that

are essential for colonization of, or survival in, the mammalian bloodstream. This represents a step

forward in the quest to identify novel targets of therapy against these deadly widespread infections

often accompanied by development of life-threatening sepsis. Our prediction and subsequent

validation of *tatC* as a common bacteremia fitness factor (**Fig 4**) and further demonstration of

increased antibiotic susceptibility of strains carrying mutations in this allele (**Fig 5**, **Table 3**)

supports the utility of our bioinformatic analyses. Having predicted the importance of many more

genes (**S9 Table**) and pathways, highlighted in **Table 2**, it is now critical to follow up this study

by investigating the importance of these bacteremia-fitness factor genes across representative

species of the *Enterobacterales* and validate their attenuation in the murine model of bacteremia.

Materials and Methods

Species-level pan-genome analysis. The pan-genome for the five *Enterobacteriaceae* species

were constructed using all the proteins from a curated list of publicly available genomes for each

species using the JCVI pan-genome pipeline [14]. Specifically, we used the "user_core_cutoff"

branch of the JCVI pan-genome pipeline found on GitHub [124]. These included the reference

genomes used for Tn-Seq as well as genome quality, UMH and common laboratory strains, and

epidemiological importance. The number of genomes in each species-level pan-genome run

ranged from a minimum of 28 (*Citrobacter*) to a maximum of 281 (*Escherichia*) and consisted

mainly of a single species per genus (*e.g.*, *C. freundii, E. hormaechei, K. pneumoniae,* and *Serratia*

marcescens) where only the *Escherichia* pan-genome also contained *Shigella* spp. The default

parameters for the pan-genome pipeline were used, except for a core cutoff (i.e.,

run_pangenome.pl --core_cutoff) of 70% of genomes as opposed to the 95% default.

Enterobacteriaceae **Multi-Species Core (MSC) Pan-genome analysis.** The pan-genome pipeline

results were used to create a core "pseudo-genome" for each species from the sequences of the

core centroids and their location on the species-level consensus pan-chromosome as shown in the

Core.att attribute output file [125]. These core pseudo-genomes from the five species were

integrated in a MSC pan-genome using the same parameters including the 70% core cutoff. A

consensus MSC pan-chromosome was constructed as described for the species-level pan-

chromosomes. For all species pan-genomes and the MSC pan-genome, the TIGR roles and AMR

were assigned to centroids using the best match for the HMMer3 [126] alignment of the

TIGRFAMs [127] and the CARD [20], respectively as part of the pan-genome pipeline [14]. Other

high-level gene features were integrated into the pan-genomes as described below.

Integration of Tn-Seq bacteremia-fitness genes with pan-genomes. As the original Tn-Seq

fitness experiments were performed on single bacterial strains [10–13] whose genomes were

included in the individual species pan-genomes (*e.g.*, *C. freundii* UMH14 biosample

SAMN07729546, *E. coli* CFT073 biosample SAMN02604094, *K. pneumoniae* ATCC 43816

KPPR1 biosample SAMN02982872, and *S. marcescens* UMH9 biosample SAMN06164063), we

were able to annotate the fitness gene clusters using their unique gene identifiers (**S2 Table**). If

the species pan-genome bacteremia-fitness clusters were core (*i.e.*, orthologous sequences present

in greater than 70% of the genomes), they were used in the MSC pan-genome and the MSC pan-

genome clusters were labeled a fitness centroid.

Integration of operon structures, essential and virulence factors, and KEGG ontology.

Experimentally-determined operon structures for *E. coli* and *Klebsiella[30,31,128]* were obtained

and mapped to the MSC pan-genome centroids with BLASTP (reciprocal best matches with e-

value $< 1 \times 10^{-5}$). For the *E. coli* operons, additional information, such as the Gene Symbol and the

Blattner identifier, were also included (**S5 Table**). To include as many Blattner identifiers as

possible for the MSC pan-genome clusters, in instances where the MSC clusters were not in

operons, we used the Blattner identifier and the gene symbol of any gene from the genome of *E.*

coli str. K-12 substr. MG1655 (BioSample SAMN02604091) with a good BLASTP match (same

cutoff as above) to the MSC pan-genome clusters.

Genes that were found to be essential for growth of *E. coli* K-12 and derivatives under laboratory

growth conditions were obtained from five different manuscripts/databases[25–29]. A unique set

of "essential" genes was created (**S3 Table**) from these combined data sets and assigned the unique

E. coli K-12 composite (ECK*) identifier. BLASTP was used to map the translated essential

proteins to the *Escherichia* pan-genome centroids.

Similarly, we used the full set of protein sequences from the Virulence Factor Database (VFDB)

(**S4 Table**) [19] downloaded on August 31, 2021, to do a similar BLASTP mapping to species-

level pan-genome centroids. If the reciprocal best matches had an e-value $< 1 \times 10^{-5}$ and an

additional requirement that the BLASTP match had to have more than 70% identity, then the

centroid was annotated as "Essential" or a "Virulence Factor." The cluster annotation for the

species-level pan-genome was passed to the MSC pan-genomes as described for bacteremia-fitness

proteins above.

Additionally, KEGG annotations were mapped to the MSC centroids using hmmsearch v 3.3.2

[129] against the KEGG ontology dataset collected in KOfam [130] with an e-value $< 1\times10^{-5}$ (**S5**

Table).

Venn Diagrams. Venn Diagrams were created with R with the Venn Package [131], using data

from **S5 Table.** For each of the Venn diagrams comprising **Fig 2,** a different list of MSC cluster

IDs was used as input for each species in the MSC pan-genome to count the numbers of shared

orthologs in: A) all MSC clusters; B) all MSC clusters with at least one Tn-Seq bacteremia-fitness

protein; and C) all MSC clusters with at least one virulence factor protein.

Pan-chromosome visualizations. Browsable pan-chromosome visualizations, including the

fitness, antimicrobial resistance, and TIGRFam annotations were constructed for each species pan-

genome and the MSC pan-genome with PanACEA using default parameters (**S1-5 Dataset, S5**

Table) [18]. The species pan-genome comparison was constructed with data from **S5 Table**, using

a polar graph from *ggplot2* [132]. Orthologs for each MSC cluster were identified in each species

as well as the relative position in each species pan-genome relative to *dnaA*. MSC clusters that

were not present in a species were noted with a gap. The radial position of orthologs on the figure

was the relative location of the *Escherichia* ortholog, while the color was based on the location of

ortholog in each species. The fitness clusters with an *Escherichia* ortholog are shown at the

Escherichia ortholog position.

Prioritization of fitness genes and operons. We ranked MSC pan-genome bacteremia-fitness

clusters and operons identified among the four species Tn-Seq data using a scoring rubric. The

rubric was based on four additive criteria: 1) The fold-change of the fitness defect associated with

transposon interruption of a gene in any one species, 2) a single gene being identified as a fitness

factor in multiple species, 3) whether multiple fitness factors were encoded in the same operon,

and 4) a fitness factor single-gene-knockout was previously found to confer increased antibiotic

susceptibility in *E. coli* BW25113 [22,23].

The top 60 bacteremia-fitness factors in any one species Tn-Seq data set (*i.e.*, the top 60 least fit

transposon mutants) were awarded points based on the magnitude of the fitness defect determined

in any one species. The top 1 - 20 least fit bacteremia-fitness factors were awarded 3 points, 21 -

40 were awarded 2 points, and 41 - 60 were awarded 1 point. All scores for each MSC bacteremia-

fitness factor were then summed across species. (*e.g.*, *wzxE* received 6 total points in this category

by accumulating 3 points each for providing the 6th and 15th greatest fitness defects when mutated

in *S. marcescen*s UMH9 and *K. pneumoniae* KPPR1, respectively.) One item of note; *K.*

pneumoniae KPPR1 bacteremia Tn-Seq [13] only identified 55 fitness factors, therefore this

species was slightly underrepresented in this scoring category.

The MSC pan-genome bacteremia-fitness factor clusters were also awarded points based on their

identification in multiple species Tn-Seq datasets, regardless of the magnitude of their defect. If a

cluster member was identified as a bacteremia-fitness factor in three species or two species it was

awarded 3 points or 2 points, respectively. (*e.g.*, *wzxE* received 3 points for being identified as a

bacteremia-fitness factor in Tn-Seq datasets from *C. freundii* UMH14, S. *marcescen*s UMH9, and

K. pneumoniae KPPR1.)

The MSC pan-genome bacteremia-fitness factor clusters were awarded points based on the

identification of at least one other bacteremia-fitness factor encoded in the same operon, regardless

of which species any one fitness factor was identified in. Fitness factors found encoded in an

operon with 5 to 9 other fitness factors were awarded 3 points, those encoded with 3 to 4 other

fitness factors were awarded 2 points, and those encoded in an operon with 1 to 2 other fitness

factors were awarded 1 point. An additional point was awarded to fitness factors found in an

operon where at least 50% of the genes in that operon were found to be fitness factors. (*e.g.*, *wzxE*

is encoded in a 12 gene operon with 9 other fitness factors being identified in at least one species'

Tn-Seq dataset. Therefore, *wzxE* received a total of 4 points; 3 points for being encoded in an

operon with 5 to 9 other fitness factors and 1 additional point for greater than 50% of the genes in

that operon being identified as fitness factors.)

Finally, MSC pan-genome bacteremia-fitness factor clusters were given additional points if

mutation of that gene in *E. coli* BW25113 conferred increased susceptibility to an antibiotic

[22,23]. Fitness factor clusters received another 2 points if mutation of the orthologous gene in the

Keio collection was found to have increased sensitivity to any antibiotics, regardless of the mode

of action.

Generation of *tatC* mutant derivatives and complementation constructs. Mutations in *tatC*

were generated by lambda red recombineering using established protocols [10,11,83,133]. For *C.*

freundii, *E. coli*, *K. pneumoniae,* and *S. marcescens*, the *nptII* kanamycin resistance cassette was

PCR-amplified from pKD4[133,134] and targeted to an in-frame deletion of the *tatC* ORF via 5'-

end homologous sequences (**S10 Table**). For *E. hormaechei,* the *acc(3)IV* apramycin resistance

cassette was amplified from pUC18-miniTn7T-Apr [135]. Recombination was facilitated by Red

functions encoded on pKD46, pSIM19, or pSIM18 depending on the species mutated[133,134].

All mutations were confirmed by sequencing of PCR-amplified alleles or by PCR amplicon size

and recombineering plasmids were cured prior to phenotypic analysis.

Genetic complementation of the *S. marcescens tatC* mutation was achieved by insertion of the

UMH9 *tatC* ORF into plasmid vector pBBR1MCS-5 [136] via isothermal assembly, followed by

sequencing verification. The resulting plasmid, pBB26, was transformed into *S. marcescens* via

electroporation. Genetic complementation of the *K. pneumoniae tatC* mutation with the

pAYCYC$_{tatC}$ plasmid was achieved by using SOE PCR [137] to connect the promoter of the *tat*

operon to the *tatC* ORF. The pACYC$_{tatABCD}$ plasmid insert was made by amplifying *tatABCD*

including its native promoter. The amplicons were inserted into pACYC184 at the *Hin*dIII and

*Sal*I sites and the resulting plasmids were sequenced for verification. Correct plasmids were

transformed into *K. pneumoniae* via electroporation. The *E. hormaechei tatC* mutant was

complemented by insertion of wild-type *tatABC* into pBBR1Hyg, a hygromycin resistant

derivative of pBBR1MCS-5 (this study), by isothermal assembly. The resulting plasmid,

pBBR1Hyg_*tatABC* was transformed into *E. hormaechei* by electroporation.

Murine bacteremia models. All experiments involving the use of laboratory animals were

conducted with protocols approved by the University of Michigan or Michigan State University

Institutional Animal Care and Use Committee and were performed in accordance with the Office

of Laboratory Animal Welfare guidelines. Wild-type and *tatC* mutant derivatives were cultured

separately in LB medium, then washed with PBS and normalized to the final density prior to

mixing 1:1. Tail vein injections were performed on 7-8 week old C57BL/6J mice as previously

described for all tested species [9,42] and organs were collected for homogenization and CFU

determination at 24 h post-injection. For the bacteremic pneumonia model, 1×10^7 CFU *S.*

marcescens or $1x10^6$ CFU *K. pneumoniae* was administered retropharyngeally into lightly

anesthetized mice, as previously described [83]. Mice were sacrificed at 24 h post-inoculation for

CFU determinations. Competitive indices for both infection models were determined as the ratio

of mutant/wild-type bacteria after 24 h compared to that same ratio in the inoculum.

Determination of minimum inhibitory concentration. The minimum inhibitory concentration

(MIC) of ampicillin and piperacillin-tazobactam was determined using Epsilometer (E-test) strips.

This experiment was carried out following the guidelines of the Clinical Microbiology Procedure

Handbook [138]. Bacteria were cultured on Lysogeny Broth (LB) agar, then inoculated in 3 mL

of LB broth the following day. After reaching OD_{600} of 0.1, cultures were spread onto Mueller-

Hinton agar using a sterile cotton-swab and allowed to dry. Then E-test strips were placed on the

dried plates and incubated at 37°C overnight. The MIC was determined by observing the point at

which the zone of clearing touches the test strip. The Clinical and Laboratory Standards Institute

(CLSI) interpretations were obtained from *Performance Standards for Antimicrobial*

Susceptibility Testing [139].

Acknowledgements

744   We would like to thank all members of the Gram-negative bacilli bacteremia team for their

745   contribution to this manuscript.

References

1. Rudd KE, Johnson SC, Agesa KM, Shackelford KA, Tsoi D, Kievlan DR, et al. Global,

regional, and national sepsis incidence and mortality, 1990-2017: analysis for the Global

Burden of Disease Study. Lancet. 2020;395: 200–211.

2. Liu V, Escobar GJ, Greene JD, Soule J, Whippy A, Angus DC, et al. Hospital deaths in

patients with sepsis from 2 independent cohorts. JAMA. 2014;312: 90–92.

3. Wisplinghoff H, Bischoff T, Tallent SM, Seifert H, Wenzel RP, Edmond MB. Nosocomial

bloodstream infections in US hospitals: analysis of 24,179 cases from a prospective

nationwide surveillance study. Clin Infect Dis. 2004;39: 309–317.

4. Thaden JT, Li Y, Ruffin F, Maskarinec SA, Hill-Rorie JM, Wanda LC, et al. Increased

Costs Associated with Bloodstream Infections Caused by Multidrug-Resistant Gram-

Negative Bacteria Are Due Primarily to Patients with Hospital-Acquired Infections.

Antimicrob Agents Chemother. 2017;61. doi:10.1128/AAC.01709-16

5. Anderson DJ, Moehring RW, Sloane R, Schmader KE, Weber DJ, Fowler VG Jr, et al.

Bloodstream infections in community hospitals in the 21st century: a multicenter cohort

study. PLoS One. 2014;9: e91713.

6. Diekema DJ, Hsueh P-R, Mendes RE, Pfaller MA, Rolston KV, Sader HS, et al. The

Microbiology of Bloodstream Infection: 20-Year Trends from the SENTRY Antimicrobial

Surveillance Program. Antimicrob Agents Chemother. 2019;63. doi:10.1128/AAC.00355-

19

7.   Centers For Disease Control And Prevention, Centers for Disease Control and Prevention

(U.S.). Antibiotic resistance threats in the United States, 2019. 2019.

doi:10.15620/cdc:82532

8.   Holmes CL, Anderson MT, Mobley HLT, Bachman MA. Pathogenesis of Gram-Negative

Bacteremia. Clin Microbiol Rev. 2021;34. doi:10.1128/CMR.00234-20

9.   Anderson MT, Brown AN, Pirani A, Smith SN, Photenhauer AL, Sun Y, et al. Replication

Dynamics for Six Gram-Negative Bacterial Species during Bloodstream Infection. MBio.

2021;12: e0111421.

10. Anderson MT, Mitchell LA, Zhao L, Mobley HLT. Capsule Production and Glucose

Metabolism Dictate Fitness during Bacteremia. MBio. 2017;8. doi:10.1128/mBio.00740-17

11. Anderson MT, Mitchell LA, Zhao L, Mobley HLT. Citrobacter freundii fitness during

bloodstream infection. Sci Rep. 2018;8: 11792.

12. Subashchandrabose S, Smith SN, Spurbeck RR, Kole MM, Mobley HLT. Genome-wide

detection of fitness genes in uropathogenic *Escherichia coli* during systemic infection.

PLoS Pathog. 2013;9: e1003788.

13. Holmes CL, Wilcox AE, Forsyth V, Smith SN, Moricz BS, Unverdorben LV, et al.

Klebsiella pneumoniae causes bacteremia using factors that mediate tissue-specific fitness

and resistance to oxidative stress. PLoS Pathog. 2023;19: e1011233.

14. Inman JM, Sutton GG, Beck E, Brinkac LM, Clarke TH, Fouts DE. Large-scale

comparative analysis of microbial pan-genomes using PanOCT. Bioinformatics. 2019;35:

1049–1050.

15. Fouts DE, Brinkac L, Beck E, Inman J, Sutton G. PanOCT: Automated Clustering of

Orthologs Using Conserved Gene Neighborhood for Pan-Genomic Analysis of Bacterial

Strains and Closely Related Species. Nucleic Acids Res. 2012;40: e172.

16. Tettelin H, Masignani V, Cieslewicz MJ, Donati C, Medini D, Ward NL, et al. Genome

analysis of multiple pathogenic isolates of *Streptococcus agalactiae*: implications for the

microbial "pan-genome." Proc Natl Acad Sci U S A. 2005;102: 13950–13955.

17. Rodriguez-Valera F, Ussery DW. Is the pan-genome also a pan-selectome? F1000Res.

2012;1: 16.

18. Clarke TH, Brinkac LM, Inman JM, Sutton G, Fouts DE. PanACEA: a bioinformatics tool

for the exploration and visualization of bacterial pan-chromosomes. BMC Bioinformatics.

2018;19: 246.

19. Chen L, Yang J, Yu J, Yao Z, Sun L, Shen Y, et al. VFDB: a reference database for

bacterial virulence factors. Nucleic Acids Res. 2005;33: D325–8.

20. McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, et al. The

comprehensive antibiotic resistance database. Antimicrob Agents Chemother. 2013;57:

3348–3357.

21. Janda JM, Abbott SL. The Changing Face of the Family (Order:): New Members,

Taxonomic Issues, Geographic Expansion, and New Diseases and Disease Syndromes. Clin

Microbiol Rev. 2021;34. doi:10.1128/CMR.00174-20

22. Tamae C, Liu A, Kim K, Sitz D, Hong J, Becket E, et al. Determination of antibiotic

hypersensitivity among 4,000 single-gene-knockout mutants of *Escherichia coli*. J

Bacteriol. 2008;190: 5981–5988.

23. Liu A, Tran L, Becket E, Lee K, Chinn L, Park E, et al. Antibiotic sensitivity profiles

determined with an *Escherichia coli* gene knockout collection: generating an antibiotic bar

code. Antimicrob Agents Chemother. 2010;54: 1393–1403.

24. Palmer T, Stansfeld PJ. Targeting of proteins to the twin-arginine translocation pathway.

Mol Microbiol. 2020;113: 861–871.

25. Gerdes SY, Scholle MD, Campbell JW, Balázsi G, Ravasz E, Daugherty MD, et al.

Experimental determination and system level analysis of essential genes in Escherichia coli

MG1655. J Bacteriol. 2003;185: 5673–5684.

26. Hashimoto M, Ichimura T, Mizoguchi H, Tanaka K, Fujimitsu K, Keyamura K, et al. Cell

size and nucleoid organization of engineered *Escherichia coli* cells with a reduced genome.

Mol Microbiol. 2005;55: 137–149.

27. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of

Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol

Syst Biol. 2006;2: 2006.0008.

28. Goodall ECA, Robinson A, Johnston IG, Jabbari S, Turner KA, Cunningham AF, et al. The

Essential Genome of K-12. MBio. 2018;9. doi:10.1128/mBio.02096-17

29. PEC (Profiling of *E. coli* Chromosome). [cited 25 Jan 2023]. Available:

https://shigen.nig.ac.jp/ecoli/pec/

30. Salgado H, Santos-Zavaleta A, Gama-Castro S, Millán-Zárate D, Díaz-Peredo E, Sánchez-

Solano F, et al. RegulonDB (version 3.2): transcriptional regulation and operon organization

in Escherichia coli K-12. Nucleic Acids Res. 2001;29: 72–74.

31. Seo J-H, Hong JS-J, Kim D, Cho B-K, Huang T-W, Tsai S-F, et al. Multiple-omic data

analysis of *Klebsiella pneumoniae* MGH 78578 reveals its transcriptional architecture and

regulatory features. BMC Genomics. 2012;13: 679.

32. Cohen J. The immunopathogenesis of sepsis. Nature. 2002;420: 885–891.

33. Chakraborty A, Adhikari P, Shenoy S, Saralaya V. Genomic analysis and clinical

importance of *Escherichia coli* isolate from patients with sepsis. Indian J Pathol Microbiol.

2015;58: 22–26.

34. Rodríguez-Baño J, Mingorance J, Fernández-Romero N, Serrano L, López-Cerero L,

Pascual A, et al. Virulence profiles of bacteremic extended-spectrum β-lactamase-producing

Escherichia coli: association with epidemiological and clinical features. PLoS One. 2012;7:

e44238.

35. Budič M, Rijavec M, Petkovšek Z, Zgur-Bertok D. *Escherichia coli* bacteriocins:

antimicrobial efficacy and prevalence among isolates from patients with bacteraemia. PLoS

One. 2011;6: e28769.

36. Ito R, Shindo Y, Kobayashi D, Ando M, Jin W, Wachino J-I, et al. Molecular

epidemiological characteristics of *Klebsiella pneumoniae* associated with bacteremia among

patients with pneumonia. J Clin Microbiol. 2015;53: 879–886.

37. Krawczyk B, Śledzińska A, Szemiako K, Samet A, Nowicki B, Kur J. Characterisation of

Escherichia coli isolates from the blood of haematological adult patients with bacteraemia:

translocation from gut to blood requires the cooperation of multiple virulence factors. Eur J

Clin Microbiol Infect Dis. 2015;34: 1135–1143.

38. Mora-Rillo M, Fernández-Romero N, Navarro-San Francisco C, Díez-Sebastián J, Romero-

Gómez MP, Fernández FA, et al. Impact of virulence genes on sepsis severity and survival

in *Escherichia coli* bacteremia. Virulence. 2015;6: 93–100.

39. Wang N, Ozer EA, Mandel MJ, Hauser AR. Genome-wide identification of *Acinetobacter*

baumannii genes necessary for persistence in the lung. MBio. 2014;5: e01163–14.

40. Cha MK, Kang C-I, Kim SH, Cho SY, Ha YE, Wi YM, et al. Comparison of the

microbiological characteristics and virulence factors of ST131 and non-ST131 clones

among extended-spectrum β-lactamase-producing Escherichia coli causing bacteremia.

Diagn Microbiol Infect Dis. 2016;84: 102–104.

41. Ciesielczuk H, Doumith M, Hope R, Woodford N, Wareham DW. Characterization of the

extra-intestinal pathogenic *Escherichia coli* ST131 clone among isolates recovered from

urinary and bloodstream infections in the United Kingdom. J Med Microbiol. 2015;64:

1496–1503.

42. Smith SN, Hagan EC, Lane MC, Mobley HLT. Dissemination and systemic colonization of

uropathogenic *Escherichia coli* in a murine model of bacteremia. MBio. 2010;1.

doi:10.1128/mBio.00262-10

43.   Bruhn KW, Pantapalangkoor P, Nielsen T, Tan B, Junus J, Hujer KM, et al. Host fate is

869   rapidly determined by innate effector-microbial interactions during *Acinetobacter*

870   *baumannii* bacteremia. J Infect Dis. 2015;211: 1296–1305.

871   44.   Koenigs A, Zipfel PF, Kraiczy P. Translation Elongation Factor Tuf of *Acinetobacter*

872   *baumannii* Is a Plasminogen-Binding Protein. PLoS One. 2015;10: e0134418.

873   45.   Skurnik D, Roux D, Aschard H, Cattoir V, Yoder-Himes D, Lory S, et al. A comprehensive

874   analysis of in vitro and in vivo genetic fitness of *Pseudomonas aeruginosa* using high-

875   throughput sequencing of transposon libraries. PLoS Pathog. 2013;9: e1003582.

876   46.   van Opijnen T, Camilli A. A fine scale phenotype-genotype virulence map of a bacterial

877   pathogen. Genome Res. 2012;22: 2541–2551.

878   47.   Baugh L, Gallagher LA, Patrapuvich R, Clifton MC, Gardberg AS, Edwards TE, et al.

879   Combining functional and structural genomics to sample the essential *Burkholderia*

880   structome. PLoS One. 2013;8: e53851.

881   48.   Brutinel ED, Gralnick JA. Anomalies of the anaerobic tricarboxylic acid cycle in

882   *Shewanella oneidensis* revealed by Tn-seq. Mol Microbiol. 2012;86: 273–283.

883   49.   Burghout P, Zomer A, van der Gaast-de Jongh CE, Janssen-Megens EM, Françoijs K-J,

884   Stunnenberg HG, et al. *Streptococcus pneumoniae* folate biosynthesis responds to

885   environmental CO2 levels. J Bacteriol. 2013;195: 1573–1582.

886   50.   Byrne RT, Chen SH, Wood EA, Cabot EL, Cox MM. *Escherichia coli* genes and pathways

887   involved in surviving extreme exposure to ionizing radiation. J Bacteriol. 2014;196: 3534–

3545.

51. Carter R, Wolf J, van Opijnen T, Muller M, Obert C, Burnham C, et al. Genomic analyses

of pneumococci from children with sickle cell disease expose host-specific bacterial

adaptations and deficits in current interventions. Cell Host Microbe. 2014;15: 587–599.

52. Chaudhuri RR, Morgan E, Peters SE, Pleasance SJ, Hudson DL, Davies HM, et al.

Comprehensive assignment of roles for *Salmonella typhimurium* genes in intestinal

colonization of food-producing animals. PLoS Genet. 2013;9: e1003456.

53. de Vries SPW, Eleveld MJ, Hermans PWM, Bootsma HJ. Characterization of the molecular

interplay between *Moraxella catarrhalis* and human respiratory tract epithelial cells. PLoS

One. 2013;8: e72193.

54. Dembek M, Barquist L, Boinett CJ, Cain AK, Mayho M, Lawley TD, et al. High-

throughput analysis of gene essentiality and sporulation in Clostridium difficile. MBio.

2015;6: e02383.

55. Dong TG, Ho BT, Yoder-Himes DR, Mekalanos JJ. Identification of T6SS-dependent

effector and immunity proteins by Tn-seq in *Vibrio cholerae*. Proc Natl Acad Sci U S A.

2013;110: 2623–2628.

56. Fu Y, Waldor MK, Mekalanos JJ. Tn-Seq analysis of *Vibrio cholerae* intestinal colonization

reveals a role for T6SS-mediated antibacterial activity in the host. Cell Host Microbe.

2013;14: 652–663.

57. Gallagher LA, Shendure J, Manoil C. Genome-scale identification of resistance functions in

Pseudomonas aeruginosa using Tn-seq. MBio. 2011;2: e00315–10.

58. Gawronski JD, Wong SMS, Giannoukos G, Ward DV, Akerley BJ. Tracking insertion

mutants within libraries by deep sequencing and a genome-wide screen for *Haemophilus*

genes required in the lung. Proc Natl Acad Sci U S A. 2009;106: 16422–16427.

59. Goodman AL, McNulty NP, Zhao Y, Leip D, Mitra RD, Lozupone CA, et al. Identifying

genetic determinants needed to establish a human gut symbiont in its habitat. Cell Host

Microbe. 2009;6: 279–289.

60. Johnson JG, Livny J, Dirita VJ. High-throughput sequencing of *Campylobacter jejuni*

insertion mutant libraries reveals mapA as a fitness factor for chicken colonization. J

Bacteriol. 2014;196: 1958–1967.

61. Kamp HD, Patimalla-Dipali B, Lazinski DW, Wallace-Gadsden F, Camilli A. Gene fitness

landscapes of *Vibrio cholerae* at important stages of its life cycle. PLoS Pathog. 2013;9:

e1003800.

62. Khatiwara A, Jiang T, Sung S-S, Dawoud T, Kim JN, Bhattacharya D, et al. Genome

scanning for conditionally essential genes in *Salmonella enterica* Serotype Typhimurium.

Appl Environ Microbiol. 2012;78: 3098–3107.

63. Langereis JD, Zomer A, Stunnenberg HG, Burghout P, Hermans PWM. Nontypeable

Haemophilus influenzae carbonic anhydrase is important for environmental and intracellular

survival. J Bacteriol. 2013;195: 2737–2746.

64. Luan S-L, Chaudhuri RR, Peters SE, Mayho M, Weinert LA, Crowther SA, et al.

Generation of a Tn5 transposon library in *Haemophilus parasuis* and analysis by

transposon-directed insertion-site sequencing (TraDIS). Vet Microbiol. 2013;166: 558–566.

65.   Santa Maria JP Jr, Sadaka A, Moussa SH, Brown S, Zhang YJ, Rubin EJ, et al. Compound-

gene interaction mapping reveals distinct roles for *Staphylococcus aureus* teichoic acids.

Proc Natl Acad Sci U S A. 2014;111: 12510–12515.

66.   McDonough E, Lazinski DW, Camilli A. Identification of in vivo regulators of the *Vibrio*

cholerae xds gene using a high-throughput genetic selection. Mol Microbiol. 2014;92: 302–

315.

67.   Moule MG, Hemsley CM, Seet Q, Guerra-Assunção JA, Lim J, Sarkar-Tyson M, et al.

Genome-wide saturation mutagenesis of *Burkholderia pseudomallei* K96243 predicts

essential genes and novel targets for antimicrobial development. MBio. 2014;5: e00926–13.

68.   Palace SG, Proulx MK, Lu S, Baker RE, Goguen JD. Genome-wide mutant fitness profiling

identifies nutritional requirements for optimal growth of *Yersinia pestis* in deep tissue.

MBio. 2014;5. doi:10.1128/mBio.01385-14

69.   Remmele CW, Xian Y, Albrecht M, Faulstich M, Fraunholz M, Heinrichs E, et al.

Transcriptional landscape and essential genes of *Neisseria gonorrhoeae*. Nucleic Acids Res.

2014;42: 10579–10595.

70.   Troy EB, Lin T, Gao L, Lazinski DW, Camilli A, Norris SJ, et al. Understanding barriers to

Borrelia burgdorferi dissemination during infection using massively parallel sequencing.

Infect Immun. 2013;81: 2347–2357.

71. Turner KH, Everett J, Trivedi U, Rumbaugh KP, Whiteley M. Requirements for

Pseudomonas aeruginosa acute burn and chronic surgical wound infection. PLoS Genet.

2014;10: e1004518.

72. Valentino MD, Foulston L, Sadaka A, Kos VN, Villet RA, Santa Maria J Jr, et al. Genes

contributing to *Staphylococcus aureus* fitness in abscess- and infection-related ecologies.

MBio. 2014;5: e01729–14.

73. Verhagen LM, de Jonge MI, Burghout P, Schraa K, Spagnuolo L, Mennens S, et al.

Genome-wide identification of genes essential for the survival of *Streptococcus pneumoniae*

in human saliva. PLoS One. 2014;9: e89541.

74. Weerdenburg EM, Abdallah AM, Rangkuti F, Abd El Ghany M, Otto TD, Adroub SA, et al.

Genome-wide transposon mutagenesis indicates that *Mycobacterium marinum* customizes

its virulence mechanisms for survival and replication in different hosts. Infect Immun.

2015;83: 1778–1788.

75. Wiles TJ, Norton JP, Russell CW, Dalley BK, Fischer KF, Mulvey MA. Combining

quantitative genetic footprinting and trait enrichment analysis to identify fitness

determinants of a bacterial pathogen. PLoS Genet. 2013;9: e1003716.

76. Xiao J, Sun S, Liu Z, Fan C, Zhu B, Zhang D. Analysis of key genes for the survival of

Pantoea agglomerans under nutritional stress. Int J Biol Macromol. 2023;253: 127059.

77. Kado T, Akbary Z, Motooka D, Sparks IL, Melzer ES, Nakamura S, et al. A cell wall

synthase accelerates plasma membrane partitioning in mycobacteria. Elife. 2023;12.

doi:10.7554/eLife.81924

78. Ma J, Ahmed MAH, Shao S, Zhang Y, Wang Q, Yin K. The QseE-QseF two-component

system: A key mediator of epinephrine-regulated virulence in the marine pathogen

Edwardsiella piscicida. Microbiol Res. 2024;279: 127561.

79. Lo H-Y, Long DR, Holmes EA, Penewit K, Hodgson T, Lewis JD, et al. Transposon

sequencing identifies genes impacting invasion in a human macrophage model. Infect

Immun. 2023;91: e0022823.

80. Kim G-L, Hooven TA, Norambuena J, Li B, Boyd JM, Yang JH, et al. Growth and Stress

Tolerance Comprise Independent Metabolic Strategies Critical for *Staphylococcus aureus*

Infection. MBio. 2021;12: e0081421.

81. Shull LM, Camilli A. Transposon Sequencing of *Vibrio cholerae* in the Infant Rabbit Model

of Cholera. Methods Mol Biol. 2018;1839: 103–116.

82. Turner KH, Wessel AK, Palmer GC, Murray JL, Whiteley M. Essential genome of

Pseudomonas aeruginosa in cystic fibrosis sputum. Proc Natl Acad Sci U S A. 2015;112:

4110–4115.

83. Bachman MA, Breen P, Deornellas V, Mu Q, Zhao L, Wu W, et al. Genome-Wide

Identification of *Klebsiella pneumoniae* Fitness Genes during Lung Infection. MBio.

2015;6: e00775.

84. Gutierrez MG, Yoder-Himes DR, Warawa JM. Comprehensive identification of virulence

factors required for respiratory melioidosis using Tn-seq mutagenesis. Front Cell Infect

Microbiol. 2015;5: 78.

85.   Lourdault K, Matsunaga J, Haake DA. High-Throughput Parallel Sequencing to Measure

990   Fitness of *Leptospira interrogans* Transposon Insertion Mutants during Acute Infection.

991   PLoS Negl Trop Dis. 2016;10: e0005117.

992   86.   Russell CW, Mulvey MA. The Extraintestinal Pathogenic *Escherichia coli* Factor RqlI

993   Constrains the Genotoxic Effects of the RecQ-Like Helicase RqlH. PLoS Pathog. 2015;11:

994   e1005317.

995   87.   Stacy A, Fleming D, Lamont RJ, Rumbaugh KP, Whiteley M. A Commensal Bacterium

996   Promotes Virulence of an Opportunistic Pathogen via Cross-Respiration. MBio. 2016;7.

997   doi:10.1128/mBio.00782-16

998   88.   Troy EB, Lin T, Gao L, Lazinski DW, Lundt M, Camilli A, et al. Global Tn-seq analysis of

999   carbohydrate utilization and vertebrate infectivity of *Borrelia burgdorferi*. Mol Microbiol.

2016;101: 1003–1023.

89.   Lourdault K, Matsunaga J, Evangelista KV, Haake DA. High-throughput Parallel

Sequencing to Measure Fitness of *Leptospira interrogans* Transposon Insertion Mutants

During Golden Syrian Hamster Infection. J Vis Exp. 2017. doi:10.3791/56442

90.   Forsyth VS, Mobley HLT, Armbruster CE. Transposon Insertion Site Sequencing in a

Urinary Tract Model. Methods Mol Biol. 2019;2021: 297–337.

91.   Eichelberger KR, Sepúlveda VE, Ford J, Selitsky SR, Mieczkowski PA, Parker JS, et al.

Tn-Seq Analysis Identifies Genes Important for *Yersinia pestis* Adherence during Primary

Pneumonic Plague. mSphere. 2020;5. doi:10.1128/mSphere.00715-20

92. Perpich JD, Yakoumatos L, Stocke KS, Lewin GR, Ramos A, Yoder-Himes DR, et al.

Porphyromonas gingivalis Tyrosine Kinase Is a Fitness Determinant in Polymicrobial

Infections. Infect Immun. 2022;90: e0017022.

93. Belanger CR, Dostert M, Blimkie TM, Lee AH-Y, Dhillon BK, Wu BC, et al. Surviving the

host: Microbial metabolic genes required for growth of in physiologically-relevant

conditions. Front Microbiol. 2022;13: 1055512.

94. Price SL, Thibault D, Garrison TM, Brady A, Guo H, Kehl-Fie TE, et al. Droplet Tn-Seq

identifies the primary secretion mechanism for yersiniabactin in *Yersinia pestis*. EMBO

Rep. 2023;24: e57369.

95. Capel E, Barnier J-P, Zomer AL, Bole-Feysot C, Nussbaumer T, Jamet A, et al. Peripheral

blood vessels are a niche for blood-borne meningococci. Virulence. 2017;8: 1808–1819.

96. Brauer AL, White AN, Learman BS, Johnson AO, Armbruster CE. d-Serine Degradation by

Proteus mirabilis Contributes to Fitness during Single-Species and Polymicrobial Catheter-

Associated Urinary Tract Infection. mSphere. 2019;4. doi:10.1128/mSphere.00020-19

97. Subashchandrabose S, Smith S, DeOrnellas V, Crepin S, Kole M, Zahdeh C, et al.

Acinetobacter baumannii Genes Required for Bacterial Survival during Bloodstream

Infection. mSphere. 2016;1. doi:10.1128/mSphere.00013-15

98. Anderson MT, Mitchell LA, Mobley HLT. Cysteine Biosynthesis Controls *Serratia*

marcescens Phospholipase Activity. J Bacteriol. 2017;199. doi:10.1128/JB.00159-17

99. Armbruster CE, Forsyth VS, Johnson AO, Smith SN, White AN, Brauer AL, et al. Twin

arginine translocation, ammonia incorporation, and polyamine biosynthesis are crucial for

Proteus mirabilis fitness during bloodstream infection. PLoS Pathog. 2019;15: e1007653.

100.   Yuda E, Tanaka N, Fujishiro T, Yokoyama N, Hirabayashi K, Fukuyama K, et al.

Mapping the key residues of SufB and SufD essential for biosynthesis of iron-sulfur

clusters. Sci Rep. 2017;7: 9387.

101.   Rai AK, Mitchell AM. Enterobacterial Common Antigen: Synthesis and Function of an

Enigmatic Molecule. MBio. 2020;11. doi:10.1128/mBio.01914-20

102.   Kneidinger B, Marolda C, Graninger M, Zamyatina A, McArthur F, Kosma P, et al.

Biosynthesis pathway of ADP-L-glycero-beta-D-manno-heptose in *Escherichia coli*. J

Bacteriol. 2002;184: 363–369.

103.   Milivojevic M, Dangeard A-S, Kasper CA, Tschon T, Emmenlauer M, Pique C, et al.

ALPK1 controls TIFA/TRAF6-dependent innate immunity against heptose-1,7-

bisphosphate of gram-negative bacteria. PLoS Pathog. 2017;13: e1006224.

104.   Zhou P, She Y, Dong N, Li P, He H, Borio A, et al. Alpha-kinase 1 is a cytosolic innate

immune receptor for bacterial ADP-heptose. Nature. 2018;561: 122–126.

105.   Holmes CL, Smith SN, Gurczynski SJ, Severin GB, Unverdorben LV, Vornhagen J, et al.

The ADP-Heptose Biosynthesis Enzyme GmhB is a Conserved Gram-Negative Bacteremia

Fitness Factor. Infect Immun. 2022;90: e0022422.

106.   Berks BC, Sargent F, Palmer T. The Tat protein export pathway. Mol Microbiol.

2000;35: 260–274.

107.   Willsky GR, Malamy MH. Characterization of two genetically separable inorganic

phosphate transport systems in *Escherichia coli*. J Bacteriol. 1980;144: 356–365.

108.   Lamarche MG, Dozois CM, Daigle F, Caza M, Curtiss R 3rd, Dubreuil JD, et al.

Inactivation of the pst system reduces the virulence of an avian pathogenic *Escherichia coli*

O78 strain. Infect Immun. 2005;73: 4138–4145.

109.   Dillingham MS, Kowalczykowski SC. RecBCD enzyme and the repair of double-

stranded DNA breaks. Microbiol Mol Biol Rev. 2008;72: 642–71.

110.   Wyatt HDM, West SC. Holliday junction resolvases. Cold Spring Harb Perspect Biol.

2014;6: a023192.

111.   Rice DW, Rafferty JB, Artymiuk PJ, Lloyd RG. Insights into the mechanisms of

homologous recombination from the structure of RuvA. Curr Opin Struct Biol. 1997;7:

798–803.

112.   Ledger EVK, Lau K, Tate EW, Edwards AM. XerC Is Required for the Repair of

Antibiotic- and Immune-Mediated DNA Damage in *Staphylococcus aureus*. Antimicrob

Agents Chemother. 2023;67: e0120622.

113.   Frick-Cheng AE, Sintsova A, Smith SN, Pirani A, Snitkin ES, Mobley HLT. Ferric

Citrate Uptake Is a Virulence Factor in Uropathogenic *Escherichia coli*. MBio. 2022;13:

e0103522.

114.   Herrmann KM, Weaver LM. THE SHIKIMATE PATHWAY. Annu Rev Plant Physiol

Plant Mol Biol. 1999;50: 473–503.

115.   Dosselaere F, Vanderleyden J. A metabolic node in action: chorismate-utilizing enzymes

1070   in microorganisms. Crit Rev Microbiol. 2001;27: 75–131.

1071   116.   Karki HS, Ham JH. The roles of the shikimate pathway genes, *aroA* and *aroB*, in

1072   virulence, growth and UV tolerance of *Burkholderia glumae* strain 411gr-6. Mol Plant

1073   Pathol. 2014;15: 940–947.

1074   117.   Hernanz Moral C, Flaño del Castillo E, López Fierro P, Villena Cortés A, Anguita

1075   Castillo J, Cascón Soriano A, et al. Molecular characterization of the *Aeromonas hydrophila*

1076   *aroA* gene and potential use of an auxotrophic *aroA* mutant as a live attenuated vaccine.

1077   Infect Immun. 1998;66: 1813–1821.

1078   118.   Felgner S, Frahm M, Kocijancic D, Rohde M, Eckweiler D, Bielecka A, et al. aroA-

1079   Deficient *Salmonella enterica* Serovar Typhimurium Is More Than a Metabolically

1080   Attenuated Mutant. MBio. 2016;7. doi:10.1128/mBio.01220-16

1081   119.   Brown AN, Anderson MT, Smith SN, Bachman MA, Mobley HLT. Conserved Metabolic

1082   Regulator ArcA Responds to Oxygen Availability, Iron Limitation, and Cell Envelope

1083   Perturbations during Bacteremia. bioRxiv. 2023. p. 2023.04.18.537286.

1084   doi:10.1101/2023.04.18.537286

1085   120.   Holland SK, Kennan RP, Schaub MM, D'Angelo MJ, Gore JC. Imaging oxygen tension

1086   in liver and spleen by 19F NMR. Magn Reson Med. 1993;29: 446–458.

1087   121.   Tramonti A, Nardella C, di Salvo ML, Barile A, D'Alessio F, de Crécy-Lagard V, et al.

1088   Knowns and Unknowns of Vitamin B Metabolism in. EcoSal Plus. 2021;9.

1089   doi:10.1128/ecosalplus.ESP-0004-2021

1090   122.   Stull F, Betton J-M, Bardwell JCA. Periplasmic Chaperones and Prolyl Isomerases.

1091   EcoSal Plus. 2018;8. doi:10.1128/ecosalplus.ESP-0005-2018

1092   123.   Maksimova E, Kravchenko O, Korepanov A, Stolboushkina E. Protein Assistants of

1093   Small Ribosomal Subunit Biogenesis in Bacteria. Microorganisms. 2022;10.

1094   doi:10.3390/microorganisms10040747

1095   124.   GitHub - JCVenterInstitute/PanGenomePipeline: PanGenomePipeline. In: GitHub

1096   [Internet]. [cited 19 Jan 2024]. Available:

1097   https://github.com/JCVenterInstitute/PanGenomePipeline

1098   125.   Chan AP, Sutton G, DePew J, Krishnakumar R, Choi Y, Huang X-Z, et al. A novel

1099   method of consensus pan-chromosome assembly and large-scale comparative analysis

1100   reveal the highly flexible pan-genome of *Acinetobacter baumannii*. Genome Biol. 2015;16:

1101   143.

1102   126.   Mistry J, Finn RD, Eddy SR, Bateman A, Punta M. Challenges in homology search:

1103   HMMER3 and convergent evolution of coiled-coil regions. Nucleic Acids Res. 2013;41:

1104   e121.

1105   127.   Haft DH, Loftus BJ, Richardson DL, Yang F, Eisen JA, Paulsen IT, et al. TIGRFAMs: a

1106   protein family resource for the functional identification of proteins. Nucleic Acids Res.

1107   2001;29: 41–43.

1108   128.   Santos-Zavaleta A, Salgado H, Gama-Castro S, Sánchez-Pérez M, Gómez-Romero L,

1109   Ledezma-Tejeida D, et al. RegulonDB v 10.5: tackling challenges to unify classic and high

1110   throughput knowledge of gene regulation in *E. coli* K-12. Nucleic Acids Res. 2019;47:

D212–D220.

129. Eddy SR. Accelerated Profile HMM Searches. PLoS Comput Biol. 2011;7: e1002195.

130. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al.

KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score

threshold. Bioinformatics. 2020;36: 2251–2252.

131. Dusa A. Draw Venn Diagrams [R package venn version 1.12]. 2024 [cited 12 Jan 2024].

Available: https://CRAN.R-project.org/package=venn

132. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer Science & Business

Media; 2009.

133. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia*

coli K-12 using PCR products. Proc Natl Acad Sci U S A. 2000;97: 6640–6645.

134. Datta S, Costantino N, Court DL. A set of recombineering plasmids for gram-negative

bacteria. Gene. 2006;379: 109–115.

135. Ducas-Mowchun K, De Silva PM, Crisostomo L, Fernando DM, Chao T-C, Pelka P, et

al. Next Generation of Tn-Based Single-Copy Insertion Elements for Use in Multi- and Pan-

Drug-Resistant Strains of *Acinetobacter baumannii*. Appl Environ Microbiol. 2019;85.

doi:10.1128/AEM.00066-19

136. Kovach ME, Elzer PH, Hill DS, Robertson GT, Farris MA, Roop RM 2nd, et al. Four

new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying different

antibiotic-resistance cassettes. Gene. 1995;166: 175–176.

137.   You C, Zhang Y-HP. Simple cloning and DNA assembly in *Escherichia coli* by

prolonged overlap extension PCR. Methods Mol Biol. 2014;1116: 183–192.

138.   Dwived HP, Brocco F, Matuschek E. Section 7.5 Gradient Diffusion Tests. In: Leber AL,

Burnham C-AD, editors. Clinical Microbiology Procedures Handbook. ASM Press; 2023.

139.   M100 Performance Standards for Antimicrobial Susceptibility Testing. Clinical and

Laboratory Standards Institute; 2023.

1137

Figure Captions

Fig 1. Funnel Chart Detailing the Flow of Bioinformatics Processes used in this Study. The chart is read from top to bottom beginning with the numbers of genomes from sepsis-causing *Enterobacterales* and ending with the number of predicted conserved fitness pathways. Pan-genomes were constructed using the JCVI pan-genome pipeline. Tn-Seq data was from previously published studies. Orthologous protein clusters were considered to be core if an ortholog was present in at least 70% of the input genomes.

Fig 2. Proteins Shared Across Five bacteremia-causing *Enterobacterales species* in the MSC Pan-genome. All orthologous protein clusters (**A**), the clusters labeled as "bacteremia-fitness" from Tn-Seq data (**B**), and the clusters labeled as "virulence factor" with matches to the VFDB (**C**) from the MSC pan-genome were binned, counted, and placed into a Venn diagram by whether clusters contained proteins from *C. freundii*, *E. hormaechei*, *E. coli/Shigella* spp., *K. pneumoniae*, and *S. marcescens*. Singleton clusters, representing species-specific core genes, are noted in the outermost ellipses of the Venn diagram. The Venn diagram is not to scale.

Fig 3. The ECA operon is a shared bacteremia-fitness locus between species. Genes within the conserved ECA operon are represented by arrows and colored lines indicate genes that were identified as significant bacteremia-fitness genes in previously published Tn-Seq studies.

Fig 4. Validation of *tatC* contributions to bacteremia fitness. Mice were inoculated via tail vein injection (TVI) or retropharyngeal inoculation (RI) with a mixture of wild-type and *tatC* mutant

derivatives of *S. marcescens* (A and E), *K. pneumoniae* (B and F), *E. coli* (C), or *E. hormaechei*

(D). Viable bacteria were enumerated in the spleen (S), liver, (L), kidneys (K), lungs (Lg) or blood

(B) at 24 h post-inoculation and the relative fitness of each strain was determined as the log

competitive index. Only *S. marcescens* was assayed in the kidneys based on previously determined

infection profiles [8]. Mean indices (solid line) that deviated significantly from the hypothetical

value of zero (dotted line), which represents neutral fitness, were determined by one-sample t-test:

* ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), **** ($P<0.0001$). Red symbols designate samples

from which *tatC* mutant bacteria were at or below the limit of detection.

1169

1170   **Fig 5. Disruption of *K. pneumoniae tatC* confers susceptibility to ampicillin.** Epsilometer (E-

1171   test) results for ampicillin are shown with minimum inhibitory concentration (MIC) at the point

1172   the zone of clearing touches the test strip for A) WT KPPR1 + pACYCev (pACYC184), B) KPPR1

1173   *bla* transposon mutant VK055_0891_bla, C) KPPR1 Δ*tatC* + pACYCev and D) KPPR1 Δ*tatC* +

1174   pACYC*tatC*.

1175

1176

Supporting information

S1 Table. Genomes Used in this Study.

S1 Dataset. *E. coli* **PanACEA Output.**

S2 Dataset. *K. pneumoniae* **PanACEA Output.**

S3 Dataset. *E. hormaechei* **PanACEA Output.**

S4 Dataset. *C. freundii* **PanACEA Output.**

S5 Dataset. *S. marcescens* **PanACEA Output.**

S2 Table. Tn-Seq Fitness Genes used in this Study.

S3 Table. *E. coli* **Essential Genes used in this Study.**

S4 Table. Virulence Genes used in this Study.

S5 Table. Multi-species Core Pan-genome Clusters.

S6 Dataset. MSC Pan-genome PanACEA Output.

S6 Table. Tn-Seq Fitness Factors Shared Across 3 Species.

S7 Table. Bacteremia-fitness and Virulence MSC clusters.

S8 Table. Fitness Genes Mapped to Known Operons.

S9 Table. Ranking of Fitness Genes.

S1 Fig. Distribution of Bacteremia-fitness factor centroid scores. Summary of total centroid scores for all 500 bacteremia-fitness factors scored in S9 Table. The vertical dotted line depicts the mean total centroid score (2.52 points). Light gray box indicates the 73 bacteremia-fitness factors with a total centroid score 1 standard deviation (SD) above the mean (i.e., genes receiving at least 6 total points, and includes both † and ‡ genes highlighted in Table 2). The dark gray box indicates the 32 bacteremia-fitness factors with a total centroid score 2 SDs above the mean (i.e., genes receiving at least 8 total points, including only ‡ genes highlighted in Table 2).

1200   **S10 Table. Oligonucleotide Primers Used in this Study.**

1201

Fig 3

wecA *wzzE* *wecB* *wecC* *rffG* *rffH* *wecD* *wecE* *wzxE* *wecF* *wzyE* *wecG*

Tn-seq fitness gene

S. marcescens *C. freundii* *K. pneumoniae*

Fig 4