Amidase and Lysozyme Dual Functions in TseP Reveal a New Family of Chimeric Effectors in the Type VI Secretion System

Running title: Chimeric effector with dual functions

Zeng-Hang Wang[1,2], Ying An[2], Ting Zhao[3,4], Tong-Tong Pei[2], Dora Yuping Wang[2,5], Xiaoye Liang[2], Wenming Qin[3*], Tao Dong[2*]

Affiliations:

[1] State Key Laboratory of Microbial Metabolism, Joint International Research Laboratory of Metabolic & Developmental Sciences, School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai, 200240, China

[2] Department of Immunology and Microbiology, School of Life Sciences, Southern University of Science and Technology, 518055, Shenzhen, China

[3] National Facility for Protein Science in Shanghai, Shanghai Advanced Research Institute, Chinese Academy of Sciences, Shanghai, 201210, People's Republic of China

[4] School of Pharmaceutical Sciences, Wuhan University, Wuhan, China

[5] Department of Physiology, University of Toronto, Toronto, Canada

* Correspondence to dongt@sustech.edu.cn；qinwenming@sari.ac.cn

Keywords: protein secretion, effector, peptidoglycan, amidase, synthetic biology

Abstract

Peptidoglycan (PG) serves as an essential target for antimicrobial development. An overlooked reservoir of antimicrobials lies in the form of PG-hydrolyzing enzymes naturally produced for polymicrobial competition, particularly those associated with the type VI secretion system (T6SS). Here we report that a T6SS effector TseP, from *Aeromonas dhakensis*, represents a family of effectors with dual amidase-lysozyme activities. *In vitro* PG-digestion coupled with LC-MS analysis revealed the N-domain's amidase activity, which is neutralized by either catalytic mutations or the presence of the immunity protein TsiP. The N-domain, but not the C-domain, of TseP is sufficient to restore T6SS secretion in T6SS-defective mutants, underscoring its critical structural role. Using pull-down and secretion assays, we showed that these two domains interact directly with a carrier protein VgrG2 and can be secreted separately. Homologs in *Aeromonas hydrophila* and *Pseudomonas syringae* exhibited analogous dual functions. Additionally, N- and C-domains display distinctive GC contents, suggesting an evolutionary fusion event. By altering the surface charge through structural-guided design, we engineered the TseP^{C4+} effector that successfully lyses otherwise resistant *Bacillus subtilis* cells, enabling the T6SS to inhibit *B. subtilis* in a contact-independent manner. This research uncovers TseP as a new family of bifunctional chimeric effectors targeting PG, offering a potential strategy to harness these proteins in the fight against antimicrobial resistance.

Significance Statement

Antimicrobial resistance urgently demands global interventions, and the bacteria cell wall remains a promising target. Our research introduces a novel family of bifunctional, cell-wall-damaging T6SS effectors. More importantly, we demonstrate an effective strategy to enable an otherwise ineffective enzyme to target both Gram-negative and Gram-positive bacteria. Our findings highlight a promising path forward using cell-wall-damaging effectors, a largely untapped resource, in the fight against antimicrobial resistance.

Introduction

Antimicrobial resistance (AMR) is a critical global health crisis, with bacterial AMR implicated in approximately 4.95 million deaths in 2019, and predictions estimating up to 10 million deaths per year by 2050 (1, 2). Developing novel antimicrobials is thus of paramount importance but also a highly challenging task. Bacterial cell walls, mainly composed of peptidoglycan (PG) and crucial for survival, serve as a primary target for antimicrobials. However, the discovery of conventional PG-inhibiting small molecules is impeded by fewer choices and increasing resistance (3, 4). The naturally occurring PG-lysing enzymes, secreted by bacteria for competition in polymicrobial environments (5–8), remain underexplored for AMR treatment.

As a widespread molecular weapon in Gram-negative bacteria, the type VI secretion system (T6SS) can directly inject PG-lysing enzymes and other antibacterial toxins into neighboring microbes and anti-eukaryotic toxins to yeast and mammalian cells(7, 9–15). The conserved structure of T6SS consists of a double tubular structure, a baseplate TssEFGK complex, and a transmembrane TssJLM complex (16–19). A spear-like inner tube, made of stacks of Hcp hexamers, is ejected outward upon contraction of an outer sheath made of stacks of TssB/C hexamers, and a VgrG-PAAR spike complex sitting atop the Hcp tube is delivered concomitantly (12, 20, 21). Although the ejected Hcp spear can penetrate neighboring bacterial cells, T6SS-elicited damages are largely dependent on the secreted effectors rather than the penetration (22–24). Numerous T6SS effectors have been predicted in Gram-negative bacteria, and known effectors display diverse functions including damaging the cell wall and membrane, DNase, and metal scavenging (6, 25–30). Despite the functional diversity of T6SS effectors, PG-targeting effectors seem to be most commonly associated with the T6SSs.

The conserved structure of PG comprises repeating disaccharide units of N-acetylglucosamine (NAG) and N-acetylmuramic acid (NAM), with a short peptide (most commonly L-Ala-D-Glu-*m*-DAP-D-Ala-D-Ala) attached to each NAM (31). Adjacent peptide chains of PG are cross-linked, forming a continuous network that envelops bacterial cells (31). T6SS effectors known to target PG fall into two

categories, T6SS amidase/endopeptidase (Tae) families cleaving the peptides and T6SS glycoside hydrolase (Tge) families cleaving the glycan backbone. Tae effectors have been found to cleave bonds between $_D$-Glu and mDAP, between mDAP and $_D$-Ala, and between NAM and $_L$-Ala (7, 27, 32, 33). Tge effectors possess β-(1-4)-N-acetyl-muramidase activities and N-acetylglucosaminidase activities (5, 6, 27, 34). The recent discovery of Tse4 in *Acinetobacter baumannii*, cleaving both the glycan strand and the crosslinked peptide cross-bridge, suggests dual enzymatic activities could be found in a single effector (10). However, such bifunctionality among effectors is rare.

As a ubiquitous waterborne pathogen, *Aeromonas dhakensis* can cause soft tissue infection and bacteremia in fish and humans (35, 36). The T6SS of *A. dhakensis* type strain SSU is active under laboratory conditions and can secrete three known antimicrobial effectors, a membrane-targeting TseC, a lysozyme-like TseP, and a nuclease TseI (25, 37–40). Our previous findings have established that TseP serves a structural role because its expression can restore the T6SS secretion in the Δ*3eff* mutant lacking all three effector genes, and the C-terminal domain, TseP^C, possesses lysozyme activities (38). However, the function of the N-terminal domain, TseP^N, which lacks any recognizable conserved motif, has remained elusive. This study unveils TseP^N is capable of cleaving the peptide link between NAM and $_L$-Ala as an amidase. Both TseP^N and TseP^C can be secreted separately by the T6SS, but only TseP^N can compensate for T6SS defects, indicating its critical role as a structural component. Additionally, TseP^N exhibits approximately 10% higher GC content than TseP^C, and homologs of TseP^N and TseP^C exist independently in divergent species. This disparity in GC content and the presence of domain-specific homologs suggest an evolutionary fusion event. We further show that TseP^C can be engineered to hydrolyze resistant Gram-positive *Bacillus subtilis* cells, conferring the T6SS with the capability to kill Gram-positive bacteria in a contact-independent manner. This research identifies a novel class of amidase-lysozyme bifunctional effectors, provides insights into effector evolution, and highlights the potential of engineering these effectors as antimicrobials.

Results

TseP^N and TseP^C can be independently secreted by the T6SS

To delineate the regions of TseP crucial for T6SS-mediated secretion, we constructed two TseP truncations, TseP^N and TseP^C, and then expressed them in the Δ*3eff* mutant using the pBAD arabinose-inducible vectors. Consistent with our previous results, protein secretion assays show that the Δ*3eff* mutant, akin to the T6SS null Δ*vasK*, failed to secrete Hcp (Figure 1A). Complementation with TseP and TseP^N, but not TseP^C, restored Hcp secretion, despite at a reduced level relative to the wild type (Figure 1A). We next tested whether the T6SS assembly is restored using a chromosomal sheath-labeled construct TssB-sfGFP. Fluorescence microscopy analysis showed that complementation with both TseP and TseP^N, but not the TseP^C, restored sheath assembly (Figure 1, B and C). Because both Hcp secretion and sheath assembly are hallmarks of T6SS assembly, these findings highlight the important structural role of the TseP N-terminus.

Interestingly, we noticed that TseP^N, but not TseP^C, was secreted (Figure 1A). The lack of TseP^C secretion might be due to a defective T6SS. To test this, we expressed different plasmid-borne TseP constructs in a panel of SSU mutant strains. Western blot analysis not only confirmed the secretion of TseP and TseP^N but also detected the secretion of TseP^C, albeit to a lesser extent, in both wild-type and the Δ*tseP* cells (Figure 1D). Considering that VgrG2 is the carrier protein for TseP delivery (38), we next tested whether TseP^N and TseP^C could interact with VgrG2. Pull-down analysis showed direct interaction of VgrG2 with TseP^N and TseP^C, in contrast to the sfGFP and MBP negative controls (Figure 1E). These results indicate TseP^N and TseP^C can be secreted independently by the T6SS.

TseP^N is a Zn^{2+}-dependent amidase

When analyzing TseP-mediated PG hydrolysis products using ultraperformance liquid chromatography coupled with mass spectrometry (UPLC-MS), we detected two distinctive peaks and determined their corresponding products to be NAM-NAG disaccharides and Tri/Tetra peptides (Figure 2, A and B), the latter suggesting an amidase activity. The released peptides were also detected in samples

treated with TseP^N and the catalytically inactive C-terminal lysozyme mutant TseP^{E663A}, but not with

TseP^C or its inactive TseP$^{C-E663A}$ mutant, indicating that the amidase activity is attributed to TseP^N.

Sequence comparison with known amidases suggests that the TseP^N contains a predicted Zn^{2+}

binding domain with several conserved residues H19, H23, H109, H339, H359, and E539 (Figure 2C).

To test the effect of these residues on the TseP^N amidase, we replaced these sites with alanines using site-

directed mutagenesis and then purified a set of TseP^N mutant proteins. Subsequent PG-digestion assays,

coupled with UPLC-MS analysis, showed significantly reduced amidase activity in these mutants

compared to TseP^N (Figure 2D).

To test the role of metal ions for TseP's amidase activity, we added the chelating agent EDTA to

the PG-digestion mixture, which abolished the amidase activity (Figure 2E). To further identify the

specific metal ion involved, we also purified TseP in the presence of EDTA and then supplemented it

with Zn^{2+}, Ca^{2+} and a combination of both. PG-digestion assays revealed that adding Zn^{2+}, but not Ca^{2+},

restored the amidase function (Figure 2E).

Given the dual functionality of TseP, we sought to determine if TsiP, known as the immunity

protein neutralizing the lysozyme activity, could inhibit the amidase, by mixing purified TsiP with TseP,

TseP^N, and TseP^C in varying molar ratios. Enzymatic assays using purified PG showed that TsiP could

effectively inhibit both amidase activity and peptidoglycan hydrolysis activity of TseP, TseP^N and TseP^C

(Figure 2F, Supplemental Figure S1). Additionally, pull-down assays showed that TsiP interacts with not

only TseP^C but also TseP^N (Figure 2G). Collectively, these results indicate that TseP^N is a Zn^{2+}-dependent

amidase.

Amidase activity is not required for T6SS assembly or lysozyme function

To determine whether the amidase activity is important for the structural role of TseP^N, we

expressed these amidase-inactive mutants in the Δ*3eff* TssB-sfGFP mutant and tested T6SS-related

functions. Secretion analysis revealed that expression of TseP^{H339A}, but not the other mutants, restored

Hcp secretion similar to wild-type TseP (Figure 3A). As control, we examined the expression of these

TseP variants and found comparable signals in the cell lysates and secretion samples. Observations of

TssB-sfGFP sheath assembly in the Δ*3eff* TssB-sfGFP mutant further supported these findings, with the

TseP^{H339A} strain showing dynamic assembly of T6SS similar to the strain expressing the wild-type

protein (Figure 3B, Supplemental Figure S2A). Further analysis through bacterial competition assays,

with *E. coli* MG1655 strain as prey, mirrored these results (Figure 3C). The expression of TseP^{H339A} and

TseP showed comparable killing abilities while the expression of other mutants demonstrated severely

impaired killing activities as the T6SS defective Δ*3eff* (Figure 3C). Although it is unclear why the other

mutations failed to complement, the H339A amidase-inactive mutation exhibiting a similar

complementary role to wild type TseP suggests that the amidase activity is not required for T6SS

assembly.

Additionally, we tested whether there is any synergistic effect between amidase and the TseP^{C}-

mediated lysozyme activities by comparing PG degradation treated with TseP, TseP^{C}, and TseP^{H339A}.

Results showed that TseP exhibited higher activities than the TseP^{C} alone, but there was no difference

between TseP and the TseP^{H339A} mutant (Figure 3D, Supplemental Figure S2B). This indicates that

amidase and lysozyme activities can operate as independent modules within the protein.

TseP homologs exhibit similar dual functions

Homolog searching analysis reveals a broad distribution of TseP homologs across Gram-negative

bacteria (Figure 4A, Supplemental Figure S3). To examine the dual functionality of these homologs, we

selected two representative proteins that have not been previously characterized: AHA_1849 from

Aeromonas hydrophila which is a closely related homolog from the same genus, and PSPTO_5204 from

Pseudomonas syringae, a more distantly related homolog. To determine whether AHA_1849 and

PSPTO_5204 share amidase and lysozyme activities as TseP, we expressed and purified their respective

N- and C-terminal truncated mutants, AHA_1849^{N} (1-640aa), AHA_1849^{C} (641-851aa), PSPTO_5204^{N}

(1-343aa), and PSPTO_5204^{C} (344-588aa) (Supplemental Figure S3). We also constructed inactivating

mutations in their predicted catalytic residues for each protein to serve as controls (Supplemental Figure

S3). Subsequent PG-hydrolysis and UPLC-MS analysis confirmed that both AHA_1849 and

PSPTO_5204 exhibit dual amidase and lysozyme activities, capable of cleaving both peptide and glycan

chains (Figure 4, B and C).

Additionally, by analyzing the GC-contents of the two *Aeromonas* homologs, we found that the

GC-content of TseP^N is about 10% higher than that of TseP^C and 10% lower than the upstream VgrG

(Figure 4D). This suggests that these two domains might have been horizontally transferred and

subsequently fused during evolution. To explore this hypothesis further, we searched for homolog

proteins with similar structures to either TseP^N or TseP^C in the Foldseek Search AFDB50 database (41),

yielding 1000 proteins with comparable predicted structures. We identified TseP^N-only homologs in

various *Aeromonas* species, including a specific homolog in *A. schubertii* possessing only the TseP^N

domain, whereas TseP^C-homologs were discovered in several other species, such as *Fulvimonas soli* and

Vibrio campbellii (Figure 4E). The comparison of their predicted structures reveals high similarity

(Figure 4F, Supplemental Table S2 and S3), further supporting the notion that TseP^N and TseP^C domains

may have been fused through a recombination event, resulting in a chimeric, dual-functional TseP.

Crystallographic analysis of TseP^C reveals a wide active groove

While testing TseP-mediated hydrolysis of purified PG from Gram-negative bacteria, we noticed

that both TseP and TseP^C displayed more efficient digestion than the commonly used hen-egg lysozyme

(Figure 5A). This enhanced activity suggests that TseP may utilize a unique mechanism for cell wall

degradation. To further investigate this, we determined the X-ray crystal structure of TseP^C to a resolution

of 2.27Å (R$_{work}$=0.189 and R$_{free}$=0.227), by molecular replacement with one molecule in the asymmetric

unit (Table 1).

The crystal structure revealed that TseP^C comprises a smaller lobe of seven helices and a larger

lobe of ten helices, separated by a concave groove (Figures 5B, Supplemental Figure S4A). The surface

electrostatic potential of TseP^C, calculated using the Adaptive Poisson-Boltzmann Solver (APBS),

consists of both electronegative and electropositive regions, unlike the predominantly electronegative

groove of lysozyme (Figure 5C, Supplemental Figure S4B). The electropositive surface of TseP^C consists

of residues R723 and K796, with an adjacent residue Y720 corresponding to W62 in lysozyme, altering

which is known to affect substrate binding (42). Additionally, the previously predicted catalytic dyad

(E655 and E663) was located near the groove. Although this structural arrangement resembles that of

other lysozyme family members, TseP^C exhibits a less compact conformation with a notably wider

groove compared to the typical lysozyme (Figure 5D). Previous studies have shown that the lysozyme

possesses catalytic sites at E53 and D70 in the substrate-binding groove (42), while the active residues

in the TseP^C structure are E654 and E663, respectively (Figure 5D). To test the effect of E663 for TseP^C

activity, we purified a mutant E663D (TseP$^{C-E663D}$) protein and perform PG digestion assays. The results

show that TseP$^{C-E663D}$ lost the activity, suggesting the greater distance within the groove requires this

glutamate residue (Figure 5E). These results collectively highlight the key structural features that may

account for the increased activity of TseP^C relative to the hen-egg lysozyme.

188 Engineered TseP^C can lyse Gram-positive cells

To test whether TseP can degrade various cell walls, we purified PG from *B. subtilis*, a model

Gram-positive bacterium, for use as substrate. However, we failed to detect any PG hydrolysis treated

with TseP or TseP^C, in contrast to the lysozyme (Figure 6A). Further purification of PG with NaOH

treatment, removing PG-associated peptides, led to effective hydrolysis of PG by TseP and TseP^C,

suggesting a difference in substrate accessibility between the lysozyme and TseP. By comparing protein

surface electrostatic potentials between the lysozyme and TseP^C, we noticed a more electronegative

surface patch in TseP^C, suggesting a potential electrostatic repulsion from negatively charged PG (Figure

6B, Supplemental Figure S4B). To increase electropositive regions while minimizing impact on activity,

we mutated four negative charge residues distant from the active site and obtained a quadruple mutant

TseP^{C4+} (Figure 6B). Activity assays confirmed that TseP^{C4+} retained PG-hydrolyzing activities (Figure

6C). To test whether the engineered TseP^{C4+} was able to lyse *B. subtilis*, we treated *B. subtilis* cells with

purified TseP^C and TseP^{C4+} and found that cells were lysed effectively by TseP^{C4+} but not TseP^C (Figure

6D, Supplemental Figure S4C). These results indicate that modulating the surface charge is an effective

strategy to expand the target range of PG-lysing effectors.

Next, we tested whether the engineered TseP^{C4+} could be delivered by the T6SS for inhibiting *B.*

subtilis. Protein secretion analysis showed a significant TseP^{C4+} signal in secretion samples, despite at a

lower level than the wild type TseP^{C} (Figure 6E). When *B. subtilis* was co-incubated with the Δ*tseP*

mutant expressing the engineered TseP^{C4+} in liquid culture, survival of *B. subtilis* was significantly

reduced in a T6SS-dependent manner (Figure 6F, Supplemental Figure S4D). These data collectively

suggest that the T6SS can be armed with engineered TseP^{C4+} to gain Gram-positive killing capabilities

in a contact-independent manner.

Discussion

The T6SS plays a pivotal role in bacterial competition, using a variety of antibacterial effectors

to kill competing cells. The exploration of effector functions holds the promise of unlocking a largely

untapped source of novel antimicrobials, presenting innovative approaches to address the escalating

antimicrobial resistance crisis. Here we report that TseP represents a novel family of dual-functional

chimeric effectors, exhibiting both amidase and lysozyme activities. Our study reveals that the N-

terminus of TseP not only serves a critical structural role in T6SS assembly but also acts as a Zn^{2+}-

dependent amidase. The N- and C-domains are both secreted and functionally independent, and may be

subject to an evolutionary fusion event to form the chimeric full-length TseP. Such multifunctionality

underscores a complex evolutionary strategy aimed at promoting T6SS-mediated competitiveness.

Among the known T6SS effectors, those targeting the bacterial PG layer represent a prevalent

category. Certain T6SSs are capable of translocating multiple PG-targeting effectors. For instance, the

H1-T6SS in *Pseudomonas aeruginosa* deploys Tse1 and Tse3, while *Vibrio cholerae* secretes VgrG3

and TseH (5–7, 9, 43, 44). Tse1 and TseH exhibit endopeptidase activities, whereas Tse3 and VgrG3

have potent lysozyme activities (5–7, 9, 43, 44). Yet, it remains relatively uncommon to identify effectors

that possess dual functionality, capable of cleaving both the glycan chains and the cross-linked peptides

within the bacterial cell wall. To date, the sole established example is Tse4 from *A. baumannii*, which

acts both as a transglycosylase to cleave the glycan strand and as an endopeptidase to cut the peptide

cross-bridge (10, 45, 46). Our enzymatic assays reveal that TseP distinguishes itself from Tse4 with dual

roles as a lysozyme and an amidase, as well as a structural role for T6SS assembly. Further homology

searches and biochemical validations of homolog activities substantiate that TseP constitutes a new

family of dual-functional effectors.

Our data demonstrate that the N-terminus of TseP can modulate T6SS assembly, which provides

further evidence that effectors play a crucial structural role in T6SS assembly (23, 37, 47, 48). Although

this notion has now been shown in several species, the molecular details remain elusive. The assembly

of a non-contractile mutant T6SS in *V. cholerae* in the absence of effectors suggests that effectors might

be needed to stabilize a polymerizing T6SS (49). In comparison to TseP^C, the N-terminus not only

contributes to T6SS assembly but also exhibits 10% higher GC content. Additionally, both N and C-

terminus can be independently secreted by the T6SS. Using structure-assisted homolog search, we have

identified TseP^N and TseP^C-only proteins in different species. These data suggest that these two domains

may result from independent effector proteins and have undergone an evolutionary fusion event.

Previous studies have shown that expressing TseP in the periplasm is highly toxic, but this

toxicity is mitigated when catalytic mutations are introduced into the C-terminal lysozyme domain (38).

This indicates that the N-terminal amidase is either less toxic or is tightly regulated by the concentration

of Zn^{2+} in the growth medium. Although the total intracellular Zn^{2+} concentration during growth in LB

has been reported to be around 0.1mM—approximately 100 times higher than the Zn^{2+} level in the LB

medium—the bioavailable (free) Zn^{2+} concentration within *E. coli* cells remains exceedingly low (50).

A similar dependency on metal ions was observed with another cell-wall-targeting endopeptidase

effector, TseH, whose activity is dependent on the environmental concentrations of Mg^{2+} and Ca^{2+} (51).

Such metal ion dependencies might lead to misbelief about these effectors being cryptic or inactive,

complicating the discovery and functional analysis of these important proteins.

TseP is predicted to contain an EF-hand calcium-binding domain with a "helix-loop-helix" motif,

a feature previously unreported in T6SS effectors (Supplemental Figure S5A). However, mutations

introduced to this domain, aimed at exploring its role in T6SS assembly and TseP toxicity, did not reveal

any critical effect on TseP functions under the conditions tested (Supplemental Figure S5, B and C). It is

possible that this EF-hand motif is active under more Ca^{2+} stringent conditions or during infections.

Further study is required to decipher its physiological function. Additionally, while constructing TseP

variants with different residue changes, we noticed that even single residue changes could reduce the

expression, secretion or the structural role of TseP, highlighting the technical challenge of heterologous

effector engineering (Figure 3A, Supplemental Figure S6, A and B).

Because the outer membrane of Gram-negative bacteria generally prohibits the entry of cell-wall

lysing proteins, the relatively exposed Gram-positive bacteria are likely more amenable to such

treatments. However, the T6SS, exclusively found in Gram-negative bacteria, primarily targets Gram-

negative species. To date, only two T6SSs, identified in *Acidovorax citrulli* and *A. baumannii* (10, 13),

have shown the capability to kill Gram-positive bacteria. Additionally, no Gram-positive cell wall lysing

effectors have been reported; for instance, Tse4 from *A. baumannii* cannot directly lyse *B. subtilis*

cultures but requires T6SS delivery. Our biochemical characterization and engineering of TseP[C] not only

establishes its exceptional enzymatic activity but also demonstrates an effective strategy to acquire Gram-

positive-cell lysing activities, as well as to equip the T6SS of *A. dhakensis* with a contact-independent

killing ability. These results highlight the potential of antimicrobial effector engineering in addressing

antimicrobial resistance.

In conclusion, our study unveils a new family of dual functional chimeric T6SS effectors and

demonstrates the potential for bioengineering these proteins to broaden their antimicrobial spectrum. The

engineered TseP[C] represents the first effector variant capable of directly lysing *B. subtilis* cultures, paying

the way for engineering the diverse set of T6SS cell-wall targeting effectors. The urgent need for new

antimicrobials necessitates the exploration of innovative strategies. The insights from our research into

T6SS and its diverse effectors signal promising directions for future work in microbial biology and

antimicrobial therapy. When enhanced through protein engineering, these enzymes have the potential to

address bacterial infections not only in medical settings but also in food safety and animal health.

Materials and Methods

Bacterial strains and growth conditions

All strains and plasmids used in this study are displayed in Supplemental Table S1. Strains were routinely

cultivated in Lysogeny Broth (LB) ([w/v] 1% tryptone, 0.5% yeast extract, 0.5% NaCl) under aerobic

conditions at 37 ℃. Antibiotics were supplemented at the following concentrations: streptomycin (100

µg/ml), kanamycin (50 µg/ml), gentamicin (20 µg/ml), chloramphenicol (25 µg/ml for *E. coli*, 2.5 µg/ml

for SSU). All plasmids were constructed using standard molecular techniques, and their sequences were

verified through Sanger sequencing. All strains and plasmids are available upon request.

Protein secretion assay

Secretion assays were performed as previously described (38). Briefly, cells were grown in LB to

$OD_{600}=1$ and collected by centrifugation at $2,500 \times g$. The pellets were resuspended in fresh LB and

incubated at 28 ℃ for 2 h with 0.01% (w/v) L-arabinose. After incubation, cells were pelleted again by

centrifugation at $2,500 \times g$ for 8 min. Cell pellets were then resuspended in SDS-loading dye and used

as cell lysate samples. The supernatants were centrifuged again at $12,000 \times g$ for 30 s and then

precipitated in 10% (v/v) TCA (trichloroacetic acid) at -20 ℃ for 30 min. Precipitated proteins were

collected by centrifugation at $15,000 \times g$ for 30 min at 4 ℃, and then washed twice with acetone. The

air-dried pellets were resuspended in SDS-loading dye and used as the secreted protein samples. All the

samples were boiled for 10 min before SDS-PAGE analysis.

Western blotting analysis

Proteins were separated using SDS-PAGE and subsequently transferred onto a PVDF membrane (Bio-Rad). Prior to antibody incubation, the membrane was blocked with a solution of 5% (w/v) non-fat milk dissolved in TBST buffer (consisting of 50 mM Tris, 150 mM NaCl, 0.1% [v/v] Tween-20, pH 7.6) for 1 h at room temperature. The membrane was then incubated sequentially with primary and secondary antibodies. The Clarity ECL solution (Bio-Rad) was utilized for signal detection. Monoclonal antibodies specific to RpoB (RNA polymerase beta subunit) were sourced from Biolegend (Product # 663905). Additionally, polyclonal antibodies to Hcp were customized by Shanghai Youlong Biotech (39). HRP-linked secondary antibodies were purchased from Beyotime Biotechnology (Product # A0208 and # A0216, respectively).

Protein expression and purification

Proteins were expressed using pET28a vector with the His tag in *E. coli* BL21 (DE3). Strains containing the recombinant plasmid were grown in LB at 37 °C. When the optical density of the culture reached OD_{600} of 0.6, isopropyl-β-D-thiogalactopyranoside (IPTG) was added to a final concentration of 1 mM, and the culture was incubated at 20 °C for 16 h. The cells were collected by centrifugation at $2,500 \times g$ for 10 minutes at 4°C, resuspended in lysis buffer (20 mM Tris-HCl, 300 mM NaCl, 10 mM imidazole, pH 8.0), and lysed by sonication. After sonication, the lysate was centrifuged at $12,000 \times g$ for 60 min at 4 °C, and the supernatant was collected. Ni-NTA affinity chromatography was used to purify the target protein, followed by gel filtration using Superdex 200 pg 10/300 column (GE Healthcare). The protein was finally concentrated to 15 mg/ml. All samples were visualized by SDS-PAGE and stained with Coomassie brilliant blue dye. Gel images for all purified proteins used in this study are shown in Supplemental Figure S7.

Crystallization, X-ray data collection and structure determination

TseP[C] was crystallized at 18 °C using the hanging drop method. Equal volumes (1:1) of the protein solution and reservoir solution, containing 0.1 M sodium formate (pH 7.0) and 12% (w/v) PEG 3350,

were mixed. The crystals were cryoprotected in a mother liquor containing 20% v/v glycol and were

flash-cooled at 100 K.

An X-ray diffraction data set of TseP[C] was collected at beamline BL19U1 (wavelength, 0.97852 Å) at

the Shanghai Synchrotron Radiation Facility. Diffraction data was auto-processed by an aquarium

pipeline. The crystals were diffracted to 2.27 Å and belonged to the space group C 2 2 21 with unit cell

dimensions of a = 44.48 Å, b = 138.8 Å and c = 95.83 Å. This structure was solved by molecular

replacement with PHASER using the previously reported SPN1S endolysin structure (PDB ID: 4OK7)

as the searching model and further polished by COOT and PHENIX. refine (52, 53). All structure figures

were generated using PyMOL (https://pymol.org/2/) software (54).

Peptidoglycan purification

Peptidoglycan was extracted as previously described (55). Bacteria were cultured at 37 °C for 8 h. Cells

were then centrifuged at 4 °C for 30 min at 3,000 × g and the cell pellets were washed with ultrapure

water. After washing, the bacteria were resuspended in ultrapure water to an OD_{600} of 70-100 and then

added dropwise to an equal volume of boiling 8% SDS solution. The mixture was slowly stirred while

boiling for 3 h and then cooled naturally to room temperature. The precipitate was collected at 20 °C for

1 h at 100,000 × g and washed with ultrapure water five times. To remove the residual proteins and DNA,

the precipitate was resuspended in 10 ml of 10 mM Tris-HCl (pH 7.0) and treated with 5 mg DNase I

and 16 mg trypsin at 37 °C for 16 h. After treatment, the mixture was centrifuged at room temperature

for 1 h at 100,000 × g, and the pellets were dissolved and treated with an equal volume of boiling 8%

SDS solution for another 3 h. The purified peptidoglycan was obtained after washing with ultrapure water,

freeze-dried, and stored at -20°C.

In vitro **enzymatic assays and enzyme kinetics**

To characterize the *in vitro* enzymatic activity of TseP and its homologs, 500 μg of purified *E. coli* PG

was co-incubated with an equal amount of proteins in a 0.1 ml reaction system at 37 °C for 12 h. After

centrifugation, the supernatant was collected and subjected to analysis by ultraperformance liquid

chromatography-quadrupole time of flight mass spectrometry (UPLC-QTOF-MS). The Acquity UPLC HSS T3 column was chosen as the solid phase, with solvent A (water with 0.1% [v/v] formic acid) and solvent B (methanol) as the mobile phase. Analyte separation was achieved by a linear gradient of solvent B, which was gradually increased from 1% to 20% over 65 min. Thereafter, a rapid gradient adjustment was followed to increase the concentration of solvent B from 20% to 100% in 1 min, with a retention period of 5 min to ensure complete separation. Finally, the system was flushed with 99% solvent A for 4 min to complete the process.

To measure the enzyme kinetics of TseP^C, a dye-release method was used to detect the PG hydrolyzing. Prior to use, the purified PG was labelled by Remazol Brilliant Blue (RBB) following a previously described method (56). Briefly, purified PG was incubated with 20 mM RBB in 0.25 M NaOH for 12 h at 37°C. The RBB-labelled PG was collected via centrifugation for 10 min at $21,400 \times g$ and washed with ultrapure water several times until no RBB dye could be detected in the supernatant. This RBB-labelled PG was then applied for the enzyme kinetics analysis. In each reaction, 10 nM enzyme was mixed with different concentrations of RBB-labelled PG at 37°C for 10 min. Undigested PG was pelleted by centrifugation at $21,400 \times g$ for 10 min and the dye release in the supernatant was quantified by measuring its absorbance at OD$_{595}$.

Bacterial competition assays

Killer and prey strains were grown aerobically in liquid LB medium until they reached OD$_{600}$ values of 1 and 2, respectively. Later, the cells were harvested via centrifugation and gently resuspended in fresh LB broth. Killer and prey cells mixed at a ratio of 5:1 were deposited onto LB-agar plates supplemented with 0.01% (w/v) L-arabinose and co-incubated for 6 h at 28°C. Afterward, the mixed cells were recovered in fresh LB broth and vigorously vortexed to dislodge the agar. For the Gram-positive bacterial killing, the SSU and *B. subtilis* cells were mixed at a ratio of 20:1 with 0.01% L-arabinose in liquid LB medium for 24 h at 28°C. The mixtures were then serially diluted and plated onto specific antibiotic-containing plates.

373 **Bioinformatics analysis**

374 The gene sequences of *A. dhakensis* SSU were retrieved from the draft genome assembly (GenBank

375 accession NZ_JH815591.1) and confirmed by Sanger sequencing. The resulting sequences were then

376 aligned using Clustal Omega (57), and the alignment was visualized using ESPript with its default

377 settings (https://espript.ibcp.fr)(58).

378 **Data availability**

379 The data that support the findings of this study are available within the article or from the corresponding

380 author upon reasonable request.

381 **Protein Data Bank and accession code**

382 The atomic coordinate and structure factor of TseP^C have been deposited in the Protein Data Bank with

383 accession code 8XCL.

384 **Acknowledgements**

This work was supported by funding from National Natural Science Foundation of China (32030001)

and National Key R&D Program of China (2020YFA0907200). The funders had no role in study design,

data collection and interpretation, or the decision to submit the work for publication.

385 **Author contributions**

T.D. conceived the project. Z-H.W., T-T.P., D.Y.W., and X.L. performed research and data analysis. Z-

-H.W., Y.A., T.Z., and W.Q. performed the crystallography and structural analysis. T.D. and Z-H.W..

wrote the manuscript with assistance from Y.A.

Competing interests
The authors declare no competing interests.

References

1. C. J. L. Murray, *et al.*, Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet* **399**, 629–655 (2022).
2. J. O'Neill, Tackling drug-resistant infections globally: final report and recommendations. *Wellcome Collect.* (2016). Available at: https://wellcomecollection.org/works/thvwsuba/items.

3. R. A. Bonomo, β-lactamases: a focus on current challenges. *Cold Spring Harb. Perspect. Med.* **7**, a025239 (2017).

4. S. M. Drawz, R. A. Bonomo, Three decades of β-lactamase inhibitors. *Clin. Microbiol. Rev.* **23**, 160–201 (2010).

5. T. M. Brooks, D. Unterweger, V. Bachmann, B. Kostiuk, S. Pukatzki, Lytic activity of the *Vibrio cholerae* type VI secretion toxin VgrG-3 is inhibited by the antitoxin TsaB. *J. Biol. Chem.* **288**, 7618–7625 (2013).

6. T. G. Dong, B. T. Ho, D. R. Yoder-Himes, J. J. Mekalanos, Identification of T6SS-dependent effector and immunity proteins by Tn-seq in *Vibrio cholerae*. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 2623–2628 (2013).

7. A. B. Russell, *et al.*, Type VI secretion delivers bacteriolytic effectors to target cells. *Nature* **475**, 343–347 (2011).

8. J. C. Whitney, *et al.*, A broadly distributed toxin family mediates contact-dependent antagonism between gram-positive bacteria. *eLife* **6**, e26938 (2017).

9. R. D. Hood, *et al.*, A type VI secretion system of *Pseudomonas aeruginosa* targets a toxin to bacteria. *Cell Host Microbe* **7**, 25–37 (2010).

10. N.-H. Le, V. Pinedo, J. Lopez, F. Cava, M. F. Feldman, Killing of Gram-negative and Gram-positive bacteria by a bifunctional cell wall-targeting T6SS effector. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2106555118 (2021).

11. A. T. Ma, S. McAuley, S. Pukatzki, J. J. Mekalanos, Translocation of a *Vibrio cholerae* type VI secretion effector requires bacterial endocytosis by host cells. *Cell Host Microbe* **5**, 234–243 (2009).

12. J. D. Mougous, *et al.*, A virulence locus of *Pseudomonas aeruginosa* encodes a protein secretion apparatus. *Science* **312**, 1526–1530 (2006).

13. T.-T. Pei, *et al.*, Delivery of an Rhs-family nuclease effector reveals direct penetration of the gram-positive cell envelope by a type VI secretion system in *Acidovorax citrulli*. *mLife* **1**, 66–78 (2022).

14. S. Pukatzki, *et al.*, Identification of a conserved bacterial protein secretion system in *Vibrio cholerae* using the *Dictyostelium* host model system. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 1528–1533 (2006).

15. K. Trunk, *et al.*, The type VI secretion system deploys antifungal effectors against microbial competitors. *Nat. Microbiol.* **3**, 920–931 (2018).

16. M. Basler, M. Pilhofer, P. G. Henderson, J. G. Jensen, J. Mekalanos, Type VI secretion requires a dynamic contractile phage tail-like structure. *Nature* **483**, 182–186 (2012).

17. Y. R. Brunet, A. Zoued, F. Boyer, B. Douzi, E. Cascales, The type VI secretion TssEFGK-VgrG phage-like baseplate is recruited to the TssJLM membrane complex via multiple contacts and serves as assembly platform for tail tube/sheath polymerization. *PLoS Genet.* **11**, e1005545 (2015).

18. B. T. Ho, T. G. Dong, J. J. Mekalanos, A view to a kill: the bacterial type 6 secretion system. *Cell Host Microbe* **15**, 9–21 (2014).

19. J. Wang, M. Brodmann, M. Basler, Assembly and subcellular localization of bacterial type VI secretion systems. *Annu. Rev. Microbiol.* **73**, 621–638 (2019).

20. S. Pukatzki, *et al.*, Identification of a conserved bacterial protein secretion system in *Vibrio cholerae* using the *Dictyostelium* host model system. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 1528–1533 (2006).

21. M. M. Shneider, *et al.*, PAAR-repeat proteins sharpen and diversify the type VI secretion system spike. *Nature* **500**, 350–353 (2013).

22. F. Kamal, *et al.*, Differential cellular response to translocated toxic effectors and physical penetration by the type VI secretion system. *Cell Rep.* **31**, 107766 (2020).

23. X. Liang, *et al.*, An onboard checking mechanism ensures effector delivery of the type VI secretion system in *Vibrio cholerae*. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 23292–23298 (2019).

24. A. Vettiger, M. Basler, Type VI secretion system substrates are transferred and reused among sister cells. *Cell* **167**, 99-110.e12 (2016).

25. X. Liang, *et al.*, Identification of divergent type VI secretion effectors using a conserved chaperone domain. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 9106–9111 (2015).

26. S. Pukatzki, A. T. Ma, A. T. Revel, D. Sturtevant, J. J. Mekalanos, Type VI secretion system translocates a phage tail spike-like protein into target cells where it cross-links actin. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 15508–15513 (2007).

27. A. B. Russell, *et al.*, A widespread bacterial type VI secretion effector superfamily identified using a heuristic approach. *Cell Host Microbe* **11**, 538–549 (2012).

28. D. Salomon, *et al.*, Marker for type VI secretion system effectors. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 9271–9276 (2014).

29. M. Si, *et al.*, Manganese scavenging and oxidative stress response mediated by type VI secretion system in *Burkholderia thailandensis*. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E2233–E2242 (2017).

30. J. C. Whitney, *et al.*, An interbacterial NAD(P)$^+$ glycohydrolase toxin requires elongation factor Tu for delivery to target cells. *Cell* **163**, 607–619 (2015).

31. W. Vollmer, D. Blanot, M. A. De Pedro, Peptidoglycan structure and architecture. *FEMS Microbiol. Rev.* **32**, 149–167 (2008).

32. J. Ma, M. Sun, Z. Pan, C. Lu, H. Yao, Diverse toxic effectors are harbored by *vgrG* islands for interbacterial antagonism in type VI secretion system. *Biochim. Biophys. Acta BBA - Gen. Subj.* **1862**, 1635–1643 (2018).

33. R. E. Hernandez, R. Gallegos-Monterrosa, S. J. Coulthurst, Type VI secretion system effector proteins: Effective weapons for bacterial competitiveness. *Cell. Microbiol.* **22**, e13241 (2020).

34. J. C. Whitney, *et al.*, Identification, structure, and function of a novel type VI secretion septidoglycan glycoside hydrolase effector-immunity pair. *J. Biol. Chem.* **288**, 26616–26624 (2013).

35. P.-L. Chen, B. Lamy, W.-C. Ko, *Aeromonas dhakensis*, an increasingly recognized human pathogen. *Front. Microbiol.* **7**, 793 (2016).

36. H. Kitagawa, *et al.*, *Aeromonas dhakensis* is not a rare cause of *Aeromonas* bacteremia in Hiroshima, Japan. *J. Infect. Chemother.* **26**, 316–320 (2020).

37. X. Liang, *et al.*, VgrG spike dictates PAAR requirement for the assembly of the type VI secretion system. *J. Bacteriol.* e0035622 (2023). https://doi.org/10.1128/jb.00356-22.

38. X. Liang, *et al.*, Characterization of lysozyme-like effector TseP reveals the dependence of type VI secretion system (T6SS) secretion on effectors in *Aeromonas dhakensis* strain SSU. *Appl. Environ. Microbiol.* **87**, e0043521 (2021).

39. T.-T. Pei, *et al.*, Intramolecular chaperone-mediated secretion of an Rhs effector toxin by a type VI secretion system. *Nat. Commun.* **11**, 1865 (2020).

40. G. Suarez, *et al.*, Molecular characterization of a functional type VI secretion system from a clinical isolate of *Aeromonas hydrophila*. *Microb. Pathog.* **44**, 344–361 (2008).

41. M. van Kempen, *et al.*, Fast and accurate protein structure search with Foldseek. *Nat. Biotechnol.* **42**, 243–246 (2024).

42. K. Maenaka, *et al.*, Dissection of protein-carbohydrate interactions in mutant hen egg-white lysozyme complexes and their hydrolytic activity. *J. Mol. Biol.* **247**, 281–293 (1995).

43. E. Altindis, T. Dong, C. Catalano, J. Mekalanos, Secretome analysis of *Vibrio cholerae* type VI secretion system reveals a new effector-immunity pair. *mBio* **6**, e00075 (2015).

44. S. J. Hersch, *et al.*, Envelope stress responses defend against type six secretion system attacks independently of immunity proteins. *Nat. Microbiol.* **5**, 706–714 (2020).

45. S. Chou, *et al.*, Structure of a peptidoglycan amidase effector targeted to Gram-negative bacteria by the type VI secretion system. *Cell Rep.* **1**, 656–664 (2012).

46. A. Radkov, *et al.*, Antibacterial potency of type VI amidase effector toxins is dependent on substrate topology and cellular context. *eLife* **11**, e79796 (2022).

47. X. Liang, *et al.*, VgrG-dependent effectors and chaperones modulate the assembly of the type VI secretion system. *PLOS Pathog.* **17**, e1010116 (2021).

48. C.-F. Wu, *et al.*, Effector loading onto the VgrG carrier activates type VI secretion system assembly. *EMBO Rep.* **21**, e47961 (2019).

49. M. S. Stietz, X. Liang, H. Li, X. Zhang, T. G. Dong, TssA–TssM–TagA interaction modulates type VI secretion system sheath-tube assembly in *Vibrio cholerae*. *Nat. Commun.* **11** (2020).

50. C. E. Outten, and T. V. O'Halloran, Femtomolar sensitivity of metalloregulatory proteins controlling zinc homeostasis. *Science* **292**, 2488–2492 (2001).

51. M.-X. Tang, *et al.*, Abiotic factors modulate interspecies competition mediated by the type VI secretion system effectors in *Vibrio cholerae*. *ISME J.* **16**, 1765–1775 (2022).

52. P. D. Adams, *et al.*, PHENIX: building new software for automated crystallographic structure determination. *Acta Crystallogr. D Biol. Crystallogr.* **58**, 1948–1954 (2002).

53. P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta Crystallogr. D Biol. Crystallogr.* **60**, 2126–2132 (2004).

54. Schrödinger, LLC, The PyMOL molecular graphics system, Version 1.8. (2015).

55. B. Glauner, Separation and quantification of muropeptides with high-performance liquid chromatography. *Anal. Biochem.* **172**, 451–464 (1988).

56. L. Yang, Y. Gan, L. Yang, B. Jiang, J. Tang, Peptidoglycan hydrolysis mediated by the amidase AmiC and its LytM activator NlpD is critical for cell separation and virulence in the phytopathogen *Xanthomonas campestris*. *Mol. Plant Pathol.* **19**, 1705–1718 (2018).

57. F. Sievers, *et al.*, Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.* **7**, 539 (2011).

58. X. Robert, P. Gouet, Deciphering key features in protein structures with the new ENDscript server. *Nucleic Acids Res.* **42**, W320–W324 (2014).

59. K. R. Park, *et al.*, Structural mechanism underlying regulation of human EFhd2/Swiprosin-1 actin-bundling activity by Ser183 phosphorylation. *Biochem. Biophys. Res. Commun.* **483**, 442–448 (2017).

Correspondence and request for materials should be addressed to T. Dong.

Figure legends

Figure 1. Structural roles and independent secretion of TseP domains. A, Secretion analysis of TseP, TseP^N, and TseP^C in the SSU triple effector deletion mutant (Δ*3eff*). A schematic of the TseP N-terminus (TseP^N, 1-603 aa) and C-terminus (TseP^C, 604-845 aa) is depicted at the top. Hcp serves as a positive control for T6SS secretion. Hcp, RpoB, and 3V5-tagged TseP proteins were detected using specific antibodies. **B,** Time-lapse imaging of VipA-sfGFP signals in the Δ*3eff* mutant complemented with different TseP variants. Each sample was captured every 10 s for 5 min and temporally color-coded. Color scale used to temporally color code the VipA-sfGFP signals is shown at the bottom. A 30- × 30-μm representative field of cells is shown. Scale bars, 5 μm. **C,** Statistical analysis of T6SS sheath assemblies in the Δ*3eff* mutant complemented with different TseP variants. Error bars indicate the mean ± standard deviation of three biological replicates, and statistical significance was calculated using a two-tailed Student's *t*-test. ns, not significant; **, $P < 0.01$. **D,** Secretion analysis of TseP, TseP^N, and TseP^C in SSU wild type, Δ*vasK*, Δ*tseP*, and Δ*3eff* mutants. For **B** and **D**, TseP, TseP^N, and TseP^C were tagged with a 3V5 C-terminal tag and expressed on pBAD vectors. RpoB serves as an equal loading and autolysis control. **E,** Pull-down analysis of VgrG2 with TseP, TseP^N, and TseP^C. His-tagged VgrG2 and 3V5-tagged TseP, TseP^N, or TseP^C were used. His-tagged sfGFP and 3V5-tagged MBP were used as controls.

Figure 2. Functional analysis of TseP reveals an amidase activity. A, *In vitro* amidase activity of the TseP, TseP^N, TseP^C, and the lysozyme inactivated mutants TseP^{E663A} and TseP$^{C-E663A}$. Cell-wall digestion products after incubation with the TseP or its mutants were analyzed by the UPLC/MS. **B,** MS analysis of cell-wall digestion products (NAM-NAC, tetrapeptides, and tripeptides) following treatment with TseP. **C,** Protein sequence alignment of the TseP amidase domain with other amidase homologs (top), and the structural superimposition of the TseP amidase domain and 3SLU (bottom). Cartoon representations of TseP and 3SLU are shown in green and cyan, respectively. The key residues involved in Zn^{2+} binding are shown in a stick model, and the zinc ion is indicated by the green sphere. **D,** *In vitro* amidase activity of TseP^N and its amidase site mutated variants. **E,** *In vitro* amidase activity of TseP under cationic conditions of 2 mM EDTA, Zn^{2+}, Ca^{2+}, or a combination of both cations. TseP represents protein purified without EDTA treatment while TseP^{EDTA} refers to protein purified in presence of EDTA. **F,** PG digestion analysis of TseP^N with or without TsiP. The immunity protein TsiP was incubated with TseP^N on ice for 12 h before being mixed with PG. Products in **D**, **E**, and **F** were analyzed by

UPLC-QTOF MASS. **G,** Pull-down analysis of TsiP with TseP, TseP^N, and TseP^C. His-tagged TsiP and 3V5-tagged TseP, TseP^N, or TseP^C were used. His-tagged sfGFP and 3V5-tagged MBP were used as controls.

Figure 3. Amidase activity of TseP is not essential for T6SS assembly or lysozyme function. A, Secretion analysis of Hcp in the Δ*3eff* mutant complemented with different TseP variants. RpoB serves as an equal loading and autolysis control. Hcp, RpoB, and 3V5-tagged TseP proteins were detected using specific antibodies. **B,** Time-lapse imaging of VipA-sfGFP signals in the Δ*3eff* mutant complemented with TseP or its amidase-inactive mutant TseP^{H339A}. Each sample was captured every 10 s for 5 min and temporally color-coded. Color scale used to temporally color code the VipA-sfGFP signals is shown at the right. A 30 × 30 μm representative field of cells is shown. Scale bars, 5 μm. **C,** Competition analysis of the Δ*3eff* mutant complemented with different TseP variants. Competition assays were repeated once. **D,** Enzymatic activity of the TseP, TseP^C and amidase-inactive mutant TseP^{H339A}. The error bars indicate the mean ± standard deviation of three biological replicates.

Figure 4. TseP homologs showed the same *in vitro* PG-hydrolysis activity with the TseP. A, Maximum-likelihood phylogeny of TseP homologs. Phylogeny was constructed using the IQ-tree web server with bootstrap 1000 times. Proteins tested in this study are highlighted in red. **B,** Amidase activity analysis of TseP homologs. **C,** *In vitro* PG-hydrolysis activity of the TseP homologs. Products in **B** and **C** were analyzed through UPLC-QTOF MASS. **D,** GC contents of the *tseP* gene cluster and *AHA_1849* gene cluster. **E,** Summary of TseP^N and TseP^C homologs output by Foldseek Search server. **F,** Structure alignments of TseP and homologs WL1483_2262 and DSB67_24810.

Figure 5. Crystal structure of TseP^C. A, Enzymatic activity of TseP, TseP^C, and lysozyme in hydrolyzing purified *E. coli* PG. The error bars indicate the mean standard deviation of three biological replicates. **B,** The overall structure of the TseP^C. The small lobe and large lobe are shown in cyan and green, respectively, with the connecting loop depicted in red. **C,** Electrostatic potential maps of TseP^C with the Y720, R723, and K796 shown as a stick model. The electrostatic surface potentials are colored red for negative charges, blue for positive charges, and white for neutral residues. **D,** Structural comparison of TseP^C and lysozyme (PDB ID: 1LZC). The catalytic sites are shown as a stick model. **E,** *E. coli* PG digestion analysis of TseP^C and TseP$^{C-E663D}$ mutant. Error bars indicate the mean ± standard deviation of three biological replicates, and statistical significance was calculated using a two-tailed Student's *t*-test. ****, $P < 0.0001$.

Figure 6. Modification of surface charge enables TseP^C to kill Gram-positive bacteria. A, *B. subtilis* PG digestion analysis of TseP and TseP^C. PG was treated with 0.25 M NaOH for 12 h at 37 °C to remove cross-linked peptides and teichoic acid. The lysis percentage was calculated by detecting the changes of OD$_{600}$ during 1 h. The hen-egg lysozyme was used as a positive control. **B,** Electrostatic potential maps of the TseP^C and TseP^{C4+}. The active sites and the mutation region are highlighted in red circles. The negatively charged residues D604, E609, D841, and E845 in the mutation region are shown as a stick model and colored in green, and lysines are colored in cyan. **C,** *In vitro* PG-hydrolysis activity of the TseP^C and TseP^{C4+}. RBB-labelled *E. coli* PG was used as the substrate and the lysis percentage was detected by dye release. **D,** *B. subtilis* PG digestion analysis of TseP^C and TseP^{C4+}. Exponential phase *B. subtilis* cells (OD$_{600}$~1.0) were used as substrate. The lysis percentage was calculated by detecting the changes of OD$_{600}$ during 1 h with the enzyme concentration at 100 nM. **E,** Secretion analysis of TseP^C and TseP^{C4+} in the Δ*tseP* mutant. RpoB serves as an equal loading and autolysis control. RpoB and 3V5-tagged TseP^C proteins were detected using specific antibodies. **F,** Statistical analysis of *B. subtilis* cells in the competition assays. Error bars of statistical analysis in **A**, **C**, **D**, and **F** indicate the mean $\pm$ standard deviation of three biological replicates, and statistical significance was calculated using a two-tailed Student's *t*-test. ns, not significant; *, $P<0.05$; **, $P<0.01$; ****, $P<0.0001$.

Figure 7. Model of TseP dual functions and engineering. This schematic illustrates the dual amidase-lysozyme activities of TseP and demonstrates an effective engineering strategy to enhance T6SS and its effector capabilities. The T6SS can independently secrete both the N- and C-terminal domains, with multiple evidence suggesting an evolutionary fusion event. These domains interact directly with the upstream-encoded carrier protein VgrG2 for secretion. TseP^N, the N-terminal domain, functions as a Zn^{2+}-dependent amidase, while the C-terminal domain, TseP^C, exhibits lysozyme activities. However, in its native form, TseP^C does not lyse Gram-positive *B. subtilis* cells. Through structural-guided design, TseP^C was engineered to create TseP^{C4+} by altering its surface charge. This modification enables TseP^{C4+} to lyse *B. subtilis* cells without affecting its T6SS-dependent secretion. Consequently, the T6SS-equipped cell acquires the ability to lyse *B. subtilis* in a contact-independent manner. Given the diversity of TseP-like effectors and T6SS species, this approach holds significant potential for modulating interspecies competition and combating antimicrobial resistance.

A

Secreted protein | Cell lysates

B

pBAD | pTseP | pTseP^H339A

C

	10×diluted	10×diluted
ΔvasK		
WT		
pBAD		
pTseP		
pTseP^N		
pTseP^C		
pTseP^H19A		
pTseP^H23A		
pTseP^H109A		
pTseP^H339A		
pTseP^H359A		
pTseP^E539A		

Prey: *E. coli* MG1655 Killer: SSU strains

D

TseP Enzymatic

- TseP
- TseP^C
- TseP^H339A

x-axis: PG concentration (mg/ml)
y-axis: Activity (mg•μmol-1•h-1)

① T6SS can independently secrete TseP^N and TseP^C

② TseP^N exhibits Zn^{2+} dependent amidase activity

④ Structure-guided TseP^{C4+} engineering

⑤ T6SS secretes TseP^{C4+}

⑥ TseP^{C4+} lyses *B. subtilis*

③ No cell lysis

GC content = 49.23%

GC content = 39.67%

amidase

Table 1. Data collection and refinement statistics of TseP^C crystallization

	TseP^C- 8XCL
Data collection	
Space group	*C 2 2 21*
a, b, c(Å)	44.48 138.80 95.83
α, β, γ(°)	90.00 90.00 90.00
Wavelength(Å)	0.97852
Resolution(Å)	47.91 – 2.27(2.40 – 2.27)
CC$_{1/2}$	0.984(0.840)
Unique reflections	25776 (4304)
R_{meas}(%)^a	23.6(83.6)
Mean $I/\sigma(I)^a$	6.1(2.6)
Completeness(%)^a	95.9(99.4)
Multiplicity ^{a}	12.9(12.0)
Refinement	
Resolution(Å)	47.91 – 2.27
$R_{work}/R_{free}{}^b$	0.189 / 0.227
No.atoms	
Protein	1910
Water	123
Average B factors (Å^2)	
Protein	31.72
Water	37.40
R.m.s. deviations	
Bond lengths(Å)	0.007
Bond angles(°)	0.871
Ramachandran plot	
Favored (%)	97.50
Allowed (%)	2.50
Outliers (%)	0.00

^{a} The values in parentheses are for the outermost shell.

^{b} R_{free} is the R_{work} based on 5% of the data excluded from the refinement.

$R_{meas} = \sum_{hkl} \sqrt{n/(n-1)} \sum_{i=1}^{n} |I_i(hkl) - \langle I(hkl)\rangle| / \sum_{hkl} \sum_i I_i(hkl)$, where $\langle I(hkl)\rangle$ is the mean intensity of a set of equivalent reflections.

$R_{work} = \sum_{hkl} ||F_{obs}| - |F_{calc}|| / \sum_{hkl} |F_{obs}|$ where F_{obs} and F_{calc} are observed and calculated structure factors, respectively.

Supplemental Figure S1. TsiP inhibits both the amidase and lysozyme activities of TseP. The immunity

protein TsiP was incubated with TseP or TseP^C on ice for 12 h before being mixed with PG. Products were analyzed

by UPLC-QTOF-MS.

Supplemental Figure S2. The amidase activity of TseP is not required for either T6SS assembly or lysozyme function. A, Time-lapse imaging of VipA-sfGFP signals in the Δ*3eff* mutant complemented with different TseP amidase-inactive mutants. Each sample was captured every 10 s for 5 min and temporally color-coded. Color scale used to temporally color code the VipA-sfGFP signals is shown at the bottom. A 30 × 30 μm representative field of cells is shown. Scale bars, 5 μm. **B,** The maximum enzymatic activity of TseP and its variants. PG substrates (1.5 mg/ml) were treated with 10 nM TseP or its variants, respectively. Error bars indicate the mean ± standard deviation of three biological replicates, and statistical significance was calculated using a two-tailed Student's *t*-test. ns, not significant; *, $P < 0.05$; **, $P < 0.01$.

Supplemental Figure S3. Protein sequence analysis of the TseP homologs. Sequences were retrieved from

UniProt and aligned using Jalview. Alignment view was generated using ESPript 3 with default settings. Proteins

tested in this study are highlighted in red. The amidase active sites are marked with a black triangle at the bottom.

Supplemental Figure S4. Structural and mutational analyses of the C-terminal domain. A, Secondary structure analysis of TseP^C. Data was calculated using EMBL-EBI webs with PDBsum tool (https://www.ebi.ac.uk/thornton-srv/databases/pdbsum/). **B,** Electrostatic potential maps of the TseP^C and lysozyme. The electrostatic surface potentials were colored red for negative charges, blue for positive charges, and white for neutral residues. **C,** Digestion of *B. subtilis* PG with different TseP variants. Exponential phase *B. subtilis* cells (OD_{600}~1.0) were used as substrates, and the lysis percentage was calculated by detecting the changes in optical density at 600 nm. **D,** Statistical analysis of killer cells during competition assays for which the survival of prey cells is shown in Figure 6F. For **C** and **D**, error bars indicate the mean ± standard deviation of three biological replicates, and statistical significance was calculated using a two-tailed Student's *t*-test. ns, not significant; **, $P < 0.01$.

A

B

Active sites

-63.576   63.576
electrostatic potential maps

Lysozyme

Active sites

-62.631   62.631
electrostatic potential maps

TseP^C

C

Lysis of *B. subtilis*

D

Supplemental Figure S5. EF-hand domain has no effect on TseP activity. A, Protein sequence alignment (top)

and structure alignment (bottom) of the EF-hand domain of TseP to the known EF-hand domain-containing protein

$_{CD}$EFhd2 (PDB ID: 5H0P) (59). **B,** Statistical analysis of T6SS sheath assemblies in the Δ*3eff* mutant

complemented with different TseP variants. Error bars indicate the mean ± standard deviation of three biological

replicates, and statistical significance was calculated using a two-tailed Student's *t*-test. ns, not significant; **, *P*

< 0.01. **C,** Competition assays of the Δ*3eff* mutant complemented with different TseP variants against *E. coli*

MG1655. Competition assays were repeated twice.

Supplemental Figure S6. Expression and secretion of TseP variants. A, Secretion analysis of Hcp in the Δ*3eff*

mutant complemented with different TseP variants. RpoB serves as an equal loading and autolysis control. Hcp,

RpoB, and 3V5-tagged TseP proteins were detected using specific antibodies. **B,** Protein expression of TseP and

its variants in *E. coli* and SSU strains. RpoB serves as an equal loading control. RpoB and 3V5-tagged TseP

proteins were detected using specific antibodies.

Supplemental Figure S7. Purification of TseP variants and homologous proteins. A, Purification of His-TseP, His-TseP^{E663A}, His-SUMO-TseP^N, His-SUMO-TseP^C, and His-SUMO-TseP$^{C-E663A}$. The His-SUMO tag was removed by SUMO protease. Proteins were used for PG-digestion analysis in Figure 2A, 2D, and 2E. **B,** Purification of TseP proteins with or without EDTA. Proteins were used for PG-digestion analysis in Figure 2E. **C,** Purification of TseP^N variants with a His-SUMO tag. The His-SUMO tag was removed by SUMO protease. Proteins were used for PG-digestion analysis in Figure 2D. **D,** Purification of AHA_1849, PSPTO_5204, and their mutants. Proteins were used for PG-digestion analysis in Figure 4B and 4C. **E,** Purification of TseP^C and TseP$^{C-E663D}$ with an His-SUMO tag. The His-SUMO tag was removed by SUMO protease. Proteins were used for PG-lysis in Figure 5E. **F,** Purification of TseP^C variants with an His-SUMO tag. The His-SUMO tag was removed by SUMO protease. Proteins were used for PG-lysis in Figure 6C and Figure S4C. All these proteins were purified with Ni-NTA affinity chromatography column, eluted using imidazole, and analyzed via SDS-PAGE.

Supplementary Table S1. Strains and plasmids used in this study.

Strain	Genotype	Description	Source
E. coli			
BL21(DE3)	*F– ompT gal dcm lon hsdSB(rB–mB–) λ(DE3 [lacI lacUV5T7p07 ind1 sam7 nin5]) [malB+]K-12(λS)*	Strain used for protein expression	Lab stock
DH5α	*F- Φ80lacZΔM15 Δ(lacZYA-argF) U169 recA1 endA1 hsdR17 phoA supE44 thi-1 gyrA96 relA1 λ-*	Strain used for cloning	Invitrogen
T-fast	*F' proA+ B+ lacIq ΔlacZM15 / fhuA2 Δ(lac-proAB) glnV galK16 galE15 R(zgb-210::Tn10)TetR endA1 thi-1 Δ(hsdS-mcrB)5*	Strain used for cloning	TIANGEN
MG1655	*F-, lambda-, rph-1*	K-12 wild-type strain used as prey for T6SS killing	Lab stock
Aeromonas dhakensis SSU	WT	Parental strain	[1]
	ΔvasK	T6SS null, in-frame deletion of *vasK*	[1]
	ΔtseP	In-frame deletion of *tseP*	[1]
	Δ3eff, vipA-sfgfp	In-frame deletion of *tseP*, *tseI*, and *tseC*, and fusion expression of sfGFP protein and VipA protein.	This study

Plasmid

	Plasmid	Description	Source
pBAD24	pBAD24Kan	Arabinose inducible expression plasmid, kanamycin resistance	Lab stock
	pBAD24kan-VgrG2-TseP-V5	Arabinose inducible expression of VgrG2 and TseP	This study
	pBAD24kan-VgrG2-TseP^N-V5	Arabinose inducible expression of VgrG2 and TseP^N	This study
	pBAD24kan-VgrG2-TseP^C-V5	Arabinose inducible expression of VgrG2 and TseP^C	This study

	pBAD24kan-VgrG2-TseP^{H19A}-V5	Arabinose inducible expression of VgrG2 and TseP^{H19A}	This study
	pBAD24kan-VgrG2-TseP^{H23A}-V5	Arabinose inducible expression of VgrG2 and TseP^{H23A}	This study
	pBAD24kan-VgrG2-TseP^{H109A}-V5	Arabinose inducible expression of VgrG2 and TseP^{H109A}	This study
	pBAD24kan-VgrG2-TseP^{H339A}-V5	Arabinose inducible expression of VgrG2 and TseP^{H339A}	This study
	pBAD24kan-VgrG2-TseP^{H359A}-V5	Arabinose inducible expression of VgrG2 and TseP^{H359A}	This study
	pBAD24kan-VgrG2-TseP^{E539A}-V5	Arabinose inducible expression of VgrG2 and TseP^{E539A}	This study
	pBAD24kan-VgrG2-TseP^{EFhandmut}-V5	Arabinose inducible expression of VgrG2 and TseP^{EFhandmut}	This study
pBAD18	pBAD18Cm	Arabinose inducible expression plasmid, choloramphenicol resistance	Lab stock
pET28a		IPTG inducible expression plasmid, gentamycin resistant	Lab stock
	pET28a-His-SUMO-TseP	IPTG inducible expression of SUMO-TseP fusion with an N-terminal 6×His tag	Lab stock
	pET28a-His-SUMO-TseP^{E663A}	IPTG inducible expression of SUMO-TseP^{E663A} fusion with an N-terminal 6×His tag	[2]
	pET28a-His-SUMO-TseP^{N}	IPTG inducible expression of SUMO-TseP fusion with an N-terminal 6×His tag	This study
	pET28a-His-SUMO-TseP^{N-H19A}	IPTG inducible expression of SUMO-TseP^{N-H19A} fusion with an N-terminal 6×His tag	This study
	pET28a-His-SUMO-TseP^{N-H23A}	IPTG inducible expression of SUMO-TseP^{N-H23A} fusion with an N-terminal 6×His tag	This study
	pET28a-His-SUMO-TseP$^{N-H109A}$	IPTG inducible expression of SUMO-TseP$^{N-H109A}$ fusion with an N-terminal 6×His tag	This study
	pET28a-His-SUMO-TseP$^{N-H339A}$	IPTG inducible expression of SUMO-TseP$^{N-H339A}$ fusion with an N-terminal 6×His tag	This study
	pET28a-His-SUMO-TseP$^{N-H359A}$	IPTG inducible expression of SUMO-TseP$^{N-H359A}$ fusion with an N-terminal 6×His tag	This study
	pET28a-His-SUMO-TseP$^{N-E539A}$	IPTG inducible expression of SUMO-TseP$^{N-E539A}$ fusion with an N-terminal 6×His tag	This study
	pET28a-His-SUMO-TseP^{C}	IPTG inducible expression of SUMO-TseP^{C} fusion with an N-terminal 6×His tag	This study

pET28a-His-SUMO-TseP$^{C-E663D}$	IPTG inducible expression of SUMO-TseP$^{C-E663D}$ fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-TseP^{C4+}	IPTG inducible expression of SUMO-TseP^{C4+} fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-TseP$^{C-D604K}$	IPTG inducible expression of SUMO-TseP$^{C-D604K}$ fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-TseP$^{C-D841K}$	IPTG inducible expression of SUMO-TseP$^{C-D841K}$ fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-TseP$^{C-D841K/E845K}$	IPTG inducible expression of SUMO-TseP$^{C-D841K/E845K}$ fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-TsiP	IPTG inducible expression of SUMO-TsiP fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-VgrG2	IPTG inducible expression of SUMO-VgrG2 fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-AHA_1849	IPTG inducible expression of SUMO-AHA_1849 fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-AHA_1849^{N}	IPTG inducible expression of SUMO-AHA_1849^{N} fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-AHA_1849$^{N-H369A}$	IPTG inducible expression of SUMO- AHA_1849$^{N-H369A}$ fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-AHA_1849^{C}	IPTG inducible expression of SUMO-AHA_1849^{C} fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-AHA_1849$^{C-D689A}$	IPTG inducible expression of SUMO-AHA_1849$^{C-D689A}$ fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-PSPTO_5204	IPTG inducible expression of SUMO-PSPTO_5204 fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-PSPTO_5204^{N}	IPTG inducible expression of SUMO-PSPTO_5204^{N} fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-PSPTO_5204$^{N-H106A}$	IPTG inducible expression of SUMO-PSPTO_5204$^{N-H106A}$ fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-PSPTO_5204$^{N-H271A}$	IPTG inducible expression of SUMO-PSPTO_5204$^{N-H271A}$ fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-PSPTO_5204^{C}	IPTG inducible expression of SUMO-PSPTO_5204^{C} fusion with an N-terminal 6×His tag	This study
pET28a-His-SUMO-PSPTO_5204$^{C-E397A}$	IPTG inducible expression of SUMO-PSPTO_5204$^{C-E397A}$ fusion with an N-terminal 6×His tag	This study

Supplementary Table S2. The sequence identity of the N-terminal of TseP and homologs. Sequence identity was calculated using ClustalW web server. Colors are assigned from blue to red indicating low to high sequence similarity, respectively.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
TseP		41.8	88.2	88.7	47.1	12.4	14.3	12.1	13	12.1	13.3	12.6	12.3	12.5	16.7
NCTC12917_01703	41.8		42.9	41.1	60.6	14.9	13.7	13	14.7	13.9	13.3	10.9	10	14.4	17.4
WP3S18E02_26330	88.2	42.9		88.5	46.8	13	12.8	10.9	15.1	12	12.8	12.4	12	13.7	17
AHA_1849	88.7	41.1	88.5		47.4	12.5	11.6	12.6	13.3	11.7	13	13.1	15.7	13.1	17.1
PLESHI_03291	47.1	60.6	46.8	47.4		14.5	13.9	13.4	16.3	12.6	13.5	11.1	14.6	13.8	17.5
CY652_03435	12.4	14.9	13	12.5	14.5		15.6	12.8	13.5	42.1	13.2	17.4	13.5	12.2	14.9
BTO02_30765	14.3	13.7	12.8	11.6	13.9	15.6		14	13.7	13.3	18.4	13.3	13.1	34.9	12.8
PSPTO_5204	12.1	13	10.9	12.6	13.4	12.8	14		12.3	10.7	13.2	13.2	15.3	13.4	14.6
Dda3937_03232	13	14.7	15.1	13.3	16.3	13.5	13.7	12.3		13.3	17.2	14.2	14.4	17.4	18.1
EBL_c11670	12.1	13.9	12	11.7	12.6	42.1	13.3	10.7	13.3		13.7	12.3	10.7	11.6	12.8
SAMN04488136_12163	13.3	13.3	12.8	13	13.5	13.2	18.4	13.2	17.2	13.7		11.8	13.5	17.3	14.9
ABW06_23985	12.6	10.9	12.4	13.1	11.1	17.4	13.3	13.2	14.2	12.3	11.8		11.6	12.4	12.2
SAMN05216206_1427	12.3	10	12	15.7	14.6	13.5	13.1	15.3	14.4	10.7	13.5	11.6		15.9	16.3
C4K68_12090Tola_1753	12.5	14.4	13.7	13.1	13.8	12.2	34.9	13.4	17.4	11.6	17.3	12.4	15.9		13.8
Tola_1753	16.7	17.4	17	17.1	17.5	14.9	12.8	14.6	18.1	12.8	14.9	12.2	16.3	13.8	

Supplementary Table S3. The sequence identity of the C-terminal of TseP and homologs. Sequence identity was calculated using ClustalW web server. Colors are assigned from blue to red indicating low to high sequence similarity, respectively.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
TseP		10	11.8	12.4	13.6	48.1	48.9	49.8	20.7	24.2	19.8	17.4	10.1	19.3	12.1
NCTC12917_01703	10		69.2	18.7	11.4	9.5	11.4	11.4	11.8	9.5	11.4	11.4	14.7	12.3	17.1
WP3S18E02_26330	11.8	69.2		17.1	11.3	12.2	10.9	11.8	11.3	10.9	11.8	10	14.9	11.8	15.4
AHA_1849	12.4	18.7	17.1		12.4	10.9	10.4	14	11.4	11.4	10.9	12.4	15	12.4	13.5
PLESHI_03291	13.6	11.4	11.3	12.4		14.2	14	14.1	13.9	13.1	13.2	13	13.6	14.6	11.6
CY652_03435	48.1	9.5	12.2	10.9	14.2		46.4	49.8	21	23.6	19.8	20.6	11.4	21.5	11.6
BTO02_30765	48.9	11.4	10.9	10.4	14	46.4		50.6	21.3	22.6	20.7	20.4	10.1	22.7	9.8
PSPTO_5204	49.8	11.4	11.8	14	14.1	49.8	50.6		22.4	24.6	20.3	17	10.5	21	10.3
Dda3937_03232	20.7	11.8	11.3	11.4	13.9	21	21.3	22.4		58.5	26.4	21.3	10.1	33.9	9.4
EBL_c11670	24.2	9.5	10.9	11.4	13.1	23.6	22.6	24.6	58.5		25.1	23.3	10.1	35.2	8.5
SAMN04488136_12163	19.8	11.4	11.8	10.9	13.2	19.8	20.7	20.3	26.4	25.1		21.1	9.7	25.1	10.3
ABW06_23985	17.4	11.4	10	12.4	13	20.6	20.4	17	21.3	23.3	21.1		12.7	18.9	10.7
SAMN05216206_1427	10.1	14.7	14.9	15	13.6	11.4	10.1	10.5	10.1	10.1	9.7	12.7		11	50.4
C4K68_12090Tola_1753	19.3	12.3	11.8	12.4	14.6	21.5	22.7	21	33.9	35.2	25.1	18.9	11		11.6
Tola_1753	12.1	17.1	15.4	13.5	11.6	11.6	9.8	10.3	9.4	8.5	10.3	10.7	50.4	11.6	

References

[1] X. Liang, R. Moore, M. Wilton, M. J. Q. Wong, L. Lam, and T. G. Dong. Identification of divergent type VI secretion effectors using a conserved chaperone domain. Proc. Natl. Acad. Sci. U. S. A. 2015, 112(29):9106–9111.

[2] X. Liang, T.-T. Pei, Z.-H. Wang, W. Xiong, L.-L. Wu, P. Xu, S. Lin, and T. G. Dong. Characterization of lysozyme-like effector TseP reveals the dependence of type VI secretion system (T6SS) secretion on effectors in *Aeromonas dhakensis* strain SSU. Appl. Environ. Microbiol. 87(12):e00435-21.

[3] X. Liang, T.-T. Pei, H. Li, H.-Y. Zheng, H. Luo, Y. Cui, M.-X. Tang, Y.-J. Zhao, P. Xu, and T. Dong. VgrG-dependent effectors and chaperones modulate the assembly of the type VI secretion system. PLOS Pathog. 2021, 17(12):e1010116.