Going around the bend to understand the role of leg coalescence in metachronal swimming.

Nils B. Tack, Sara O. Santos, Monica M. Wilhelmus[*]
School of Engineering, Brown University, 345 Brook St., Providence, RI 02912, USA.
[*] Author for correspondence (mmwilhelmus@brown.edu)

Running title: Pleopod bending and coalescence

Summary statement: Shrimp swimming legs bend nearly horizontally and cluster together during
metachronal propulsion to reduce drag and improve the overall swimming performance.

Key words: Pleopod, bending, coalescence, shrimp, metachronal propulsion

Acknowledgements: We are grateful to Erika Tavares for performing the μ-CT scans of *P.*
vulgaris. We also gratefully acknowledge Adrian Herrera-Amaya for his insightful discussions
that strengthened the foundation of this work. We thank Marjorie Bradley for her assistance with
coordination and administrative support throughout the duration of this research. Funding for this
research was provided by the National Aeronautics and Space Administration (NASA) Rhode
Island Established Program to Stimulate Competitive Research (EPSCoR) Seed Grant.

Competing interests: The authors declare no competing interests.

Author contributions: Conceptualization: N.B.T, M.M.W; Methodology: N.B.T, S.O.S.,
M.M.W.; Software: N.B.T; Validation: N.B.T.; Formal analysis: N.B.T.; Investigation: N.B.T.,
M.M.W.; Resources: M.M.W.; Writing - original draft: N.B.T.; Writing - review & editing: N.B.T,
S.O.S, M.M.W; Visualization: N.B.T; Funding acquisition: M.M.W.

Funding: This project was supported by funding to M.M.W. from the National Aeronautics and
Space Administration (NASA) Rhode Island Established Program to Stimulate Competitive
Research (EPSCoR) Seed Grant.

Data Availability Statement: The shrimp swimming and morphological data and metadata files
reported in this investigation have been deposited in the Wilhelmus Lab Brown digital repository
(BDR). CAD files and custom Matlab scripts developed to compute kinematics and fluid flow data
have also been made available.

ABSTRACT

Many of the most abundant aquatic invertebrates display metachronal swimming by sequentially
beating closely spaced flexible appendages. Common biophysical mechanisms like appendage
spatial asymmetry and phase drive the success and performance of this locomotor mode, which is
generally explained by the need to maximize thrust production. However, the potential role of
these mechanisms in drag reduction, another important contributor to the overall swimming
performance, has yet to be evaluated. We present a comprehensive overview of the morphological,
functional and physical mechanisms promoting drag reduction during metachronal swimming by

exploring appendage differential bending and leg grouping (coalescence). We performed μ-CT
and *in-vivo* velocimetry measurements of shrimp (*Palaemonetes vulgaris*) to design a five-legged
robotic metachronal analog. This test platform enabled simultaneous flow and force measurements
to quantify the thrust and drag forces produced by flexible and stiff pleopods (legs) beating
independently or coalescing. We tested the hypothesis that coalescence and bending effectively
reduce drag during the recovery stroke (RS). The curved cross-section of the pleopods enables
passive asymmetrical bending during the RS to reduce their coefficient of drag by up to 75.8%
relative to stiff pleopods. Bending promotes physical interactions facilitating the coalescence of
three pleopods at any time during the RS to reduce drag such that the mean net thrust produced
during coalescence is increased by 30.2%. These improvements are explained by the production
of a weaker wake compared with stiff and non-coalescing pleopods. Our results describe
fundamental biological and physical components of metachronal propulsion that may aid the
development of novel bio-inspired underwater vehicles.
57
58

INTRODUCTION
Metachronal propulsion consists of sequentially beating closely spaced coordinated
appendages to generate thrust. It is employed by a formidable number of aquatic species living
across a wide scale spectrum, from microscopic protozoa to large lobsters. Its ubiquity across taxa,
including many of the major groups performing mass diel vertical migrations like copepods and
krill (Forward, 1988; Mauchline, 1998), suggests common biophysical mechanisms that drive
evolution, fitness, and ecology.
By coordinating several appendages metachronally rather than synchronously, metachronal
swimmers induce steadier flow to generate thrust more efficiently (Ford et al., 2019). The
antiplectic metachronal wave – propagating in the swimming direction – causes anterior
appendages to actuate after their posterior counterparts and promote beneficial tip vortex
interactions (Ford and Santhanakrishnan, 2021). The induced wake travels in the opposite direction
of the metachronal wave and can interact constructively with the downstream appendages to
further enhance thrust (Ford and Santhanakrishnan, 2021; Garayev and Murphy, 2021). These
mechanisms are particularly sensitive to appendage synchronicity (i.e., phase) and spacing (Alben
et al., 2010; Ford and Santhanakrishnan, 2021; Ford et al., 2019; Garayev and Murphy, 2021) for
which most metachronal swimmers fall within a narrow range (reviewed in (Byron et al., 2021)
Metachronal appendages must also achieve significant spatial or temporal asymmetry between the
power and recovery strokes to maximize thrust production (Herrera-Amaya and Byron, 2023; Lou
et al., 2022).
The biophysical commonalities among metachronal animals have been consistently found to
promote the highest average flux and fluid momentum, thereby maximizing performance during
forward swimming (Alben et al., 2010; Byron et al., 2021; Ford and Santhanakrishnan, 2021; Ford
et al., 2019; Herrera-Amaya and Byron, 2023; Khaderi et al., 2012). As such, metachronal
propulsion has been thought of as a strategy whose performance stems from maximizing thrust
production (Alben et al., 2010; Byron et al., 2021; Colin et al., 2020; Ford et al., 2019). However,
other major commonalities at the appendage and whole-system levels can be equally crucial in
reducing drag, which is another critical performance factor to consider in self-propelled systems.

At large scales, crustaceans have complex biramous swimming legs (pleopods) and fine setae that open and close during the power and recovery phases, respectively. Recent studies have highlighted the effect of dynamically modulating the effective surface area of the propulsors (the projected area perpendicular to the flow) in producing spatial asymmetry between the power and recovery strokes (Byron et al., 2021; Herrera-Amaya and Byron, 2023; Lou et al., 2022; Vogel, 2020). While this has been shown to effectively overcome both the resisting body drag and the drag on the appendages to achieve propulsion, the role of bending has yet to be evaluated.

Metachronal swimmers tend to keep their appendages relatively stiff during the power stroke but display significant spanwise bending to reduce the effective profile area, and thus any form drag, during the recovery stroke (Hessler, 1985; Johnson and Tarling, 2008). Ciliated microorganisms generate asymmetric waveforms by actively sliding adjacent microtubules within the propulsor (Dutcher, 2019). However, whether other larger metachronal organisms like crustaceans achieve active appendage spatial asymmetry using a similar mechanism is unknown. Passive asymmetrical bending resulting from the inherent structural or material properties of the appendages is more likely given the high mechanical stability and functional adaptation of crustacean exoskeletal elements through structural and compositional diversity (Fabritius et al., 2016; Raabe et al., 2005). By bending under fluid loading, flexible appendages can reduce their effective surface area and become more streamlined (Vogel, 2020). Self-reconfiguration not only lowers the drag coefficient but also reduces the stress on the structure proportionally to the resistive forces of the fluid (Alben et al., 2002; Gosselin et al., 2010). Also, inter-appendage spacing is not a physical limitation to motion, with flexibility promoting physical interactions, thus enabling a wider range of motions than their stiff counterparts (Brennen, 1974).

Most metachronal swimmers produce antiplectic waves (Byron et al., 2021; Colin et al., 2010; Daniels et al., 2021; Garayev and Murphy, 2021; Murphy et al., 2011; Sensenig et al., 2010; Tamm, 2014) that cause two or more adjacent appendages to coalesce such that their profiles can overlap. This is made possible by the proximity and relative phases of neighboring appendages (Byron et al., 2021; Murphy et al., 2011). As posterior appendages enter their recovery phase, they press against the anterior appendages that are transitioning from the power to the recovery stroke. This is particularly apparent in metachronal animals operating in the transitional to inertial flow regimes with Reynolds number, Re > 10 (Blake and Sleigh, 1974; Daniels et al., 2021; Garayev and Murphy, 2021; Murphy et al., 2011).

Here, we explore the role coalescence plays during forward swimming for the first time. Coalescence may leverage the interactions between multiple flexible appendages to further reduce the total effective surface area to the flow, thereby lessening interactions with the fluid and lowering drag during the appendage recovery stroke. While stiff appendages can still coalesce, by virtue of proximity and phase, interactions within a group might cause destructive collisions or incomplete contact causing flow instabilities. Flexibility presents distinct advantages in this context.

Appendage asymmetrical bending and coalescence are apparent from simple observations of freely swimming animals. However, we still need a systematic understanding of the relationship between flexibility, temporal coordination, and coalescence to explain how these mechanisms reduce drag to enhance propulsion. Recent computer fluid dynamics (CFD) models provided insights into how metachronally beating appendages can exploit a vortex-weakening mechanism

to produce less overall drag and improve the thrust-to-power ratio significantly (Lionetti et al.,
2023). However, experimental data about the underlying biophysical principles of drag reduction
in metachronal propulsion are scarce. It is experimentally challenging to measure the flow around
and among closely spaced appendages and disentangle the resulting thrust and drag force. Still,
several biological and robotic studies successfully identified how krill generate thrust and lift by
estimating the momentum of the wake (Ford and Santhanakrishnan, 2021; Ford et al., 2019;
Murphy et al., 2013). However, the contributions from individual and interacting legs and the
independent effects of appendage bending and coalescence on the swimming performance could
not be isolated. In addition, the recurrent simplification of appendage morphology as rigid rather
than flexible plates (Ford and Santhanakrishnan, 2021; Murphy et al., 2011; Ruszczyk et al., 2022)
might have also inadvertently undermined the significance of appendage bending as a major
contributor to drag reduction.
This investigation presents a comprehensive overview of the morphological, functional, and
physical mechanisms promoting drag reduction during metachronal swimming by establishing the
relationship between appendage flexibility, temporal coordination, and coalescence. Using μ-CT
measurements, we examined the morphology of marsh grass shrimp (*P. vulgaris*) pleopods to
explore the characteristics enabling passive asymmetrical bending. We combined in-vivo
kinematics and velocimetry experiments on free-swimming *P. vulgaris* to quantify bending and
coalescence and visualize their effect on the flow during the recovery stroke. We used this
biological data and leveraged our Pleobot (Santos et al., 2023) to design a scaled metachronal robot
with five morphologically accurate pleopod analogs fitted with flexible appendages capable of
asymmetrical bending. We quantified the thrust and drag forces of independent and coalescing
pleopods using simultaneous force and fluid flow measurements and evaluated the resulting wake.
We compared these results against rigid pleopods to answer the following three questions: Does
pleopod bending significantly reduce drag during the leg recovery phase? Given that shrimp group
up to three recovering pleopods, does coalescence turn the drag of several appendages to that of
only one? How complementary are bending and coalescence in optimizing drag reduction? We
provide evidence that reducing drag, rather than simply enhancing thrust, is a determinant factor
in the functional morphology and coordination of metachronal propulsors. Our results leverage
our understanding of the unifying biophysical principles of this locomotor mode that can be
applied to the design of novel bioinspired underwater robots.
161
162

MATERIALS AND METHODS
Animals
Marsh grass shrimp (*P. vulgaris*) (n = 13; body length = 3.11 ± 0.25 cm) were collected in June
2022 from Narragansett Bay (Rocky Point State Park, Warwick, RI, USA). The shrimp were
housed at room temperature (21°C) in a 38-liter aquarium. The capture and experimentations were
conducted in accordance with the laws of the State of Rhode Island.
169

Micro CT scan
A shrimp specimen was euthanized in 25% ethanol (seawater, 30 ppt) and transferred to a 70%
ethanol solution for 20 min until rigor mortis occurred. The left and right pleopods (from P1 to P5)

were amputated at the proximal end of the protopod. This was done to separate the pleopods to
avoid physical interactions and to enhance the clarity of the resulting high-resolution μ-CT scan.
The samples were preserved in 100% ethanol for five days and stained in a 1% solution of Iodine
in 100% ethanol for 24 h. They were then submerged in 98% Hexamethyldisilazane (HMDS) for
24 h and air-dried overnight. Chemical drying using HMDS produces similar results to critical-
point drying but allows larger samples to be processed (Alba-Tercedor and Alba-Alejandre, 2019;
Bhattacharya et al., 2020). The samples were finally mounted onto a custom-made 3D-printed
mount that fit the holding tray of the μ-CT scanner. A SkyScan 1276 high-resolution
microtomography (Bruker, Billerica, MA, USA), upgraded to receive a Hamamatsu microfocus
X-ray source (L10321-67, Bridgewater Township, NJ, USA) and a 4K camera (XIMEA
MH110XC, Lakewood, CO, USA) was used to perform high-resolution scans (see supplemental
materials for specific scanning parameters). We used the Bruker micro-CT's Skyscan software
(NRecon, DataViewer, CTAnalyser) for primary 3D reconstructions from cross-section images.
Volume rendering images were obtained with Slicer v.5.0.3. The 3D reconstruction was used to
extract transverse sections of the endopodite and exopodite of the ten legs of the specimen to
measure the radius of curvature of the chord, R, giving the chordwise curvature κ as its reciprocal
(see details in the supplementary materials). To account for differences in the width of the ramal
structures that proportionally impact the raw value of the radius of curvature, we normalized κ to
the width, W of the ramal structures (established from the above transverse sections) as $\kappa = W/R$.
192
Pleopod flexural stiffness
Five specimens were euthanized and all their pleopods were amputated at the proximal end of
the protopod and were used within one hour. The fresh protopod of each pleopod was fixed to a
rigid metal pin using cyanoacrylate glue by immobilizing the peduncle-ramal joint. In all cases,
either the endopodite or the exopodite of one leg in a pair was randomly amputated to avoid
mechanical interactions during testing. The second ramal structure was isolated similarly using the
opposing leg. The mounted pleopods were immersed in 30‰ salt water to maintain the natural
material properties and were placed on a precision balance (VWR-124B2, VWR International,
USA, precision = 0.1mg). We applied a series of point force loads at the appendage tip on the
ventral and dorsal sides, corresponding to the power and recovery strokes, respectively.
Specifically, we applied the point force at the tip of the membranous part of the rami rather than
the tip of the projecting distal setae. This is because the setae have different material properties
and spread out, thus letting the pin through. Five replicates were obtained on both sides of each
structure and two pictures were taken for each replicate, starting with the unloaded state and then
the loaded state. We analyzed each image of the deflected pleopod unit using MATLAB R2022a
(MathWorks, Natick, MA, USA) to determine its effective beam length and deflection. Beam
length (L, in m) was measured as the distance from the peduncle-ramal joint to the point of applied
force. Deflection (δ, in m) was measured as the mean vertical distance traveled by the tip of the
appendage from its initial reference location for each set of five replicates. Overall spanwise
flexural stiffness EI was measured using the beam equation as:
213
214
$$EI = \frac{FL^3}{3\delta} \qquad (1)$$

where F is the product of the mass reported on the balance readout and the gravitational
acceleration (in Newtons). This equation measures the flexural stiffness over the entire beam
length and assumes that the beam is homogeneous. Because the equation applies only to small
displacements, we removed any measurements for pleopod displacement $\delta > 0.08L$. This is
because if δ is large relative to L, the beam is bent down so far that its coordinates in the x-
dimension while loaded are significantly different from its coordinates while unloaded, thus
invalidating measurements(Combes and Daniel, 2003). Eq. 1 is valid for our measurements
because even in the most extreme cases (where δ is near $0.08L$), the error in coordinate introduced
by length changes in the x-dimension was less than 0.3% (equivalent to the measurement error).
225
Flow visualization
Free-swimming experiments were performed in a 9.5-l aquarium. Individual shrimp were
brought near the center of the tank upstream of the field of view and within the focal plane of the
camera using a beaker. After the flow stabilized, the shrimp were allowed to exit the beaker and
swim freely through the field of view, reaching constant speed by the time they entered it. We
performed bright-field PIV by seeding the water with 10 μm particles (Dantec Dynamics,
Skovlunde, Denmark) and back-lighting the animals with an LED illuminator (M530L2-C3, Thor
Labs, Newton, NJ, USA) coupled with a collimating Fresnel lens (focal distance = 10 cm). We
recorded images from the lateral view (corresponding to the left side of the animals) using a high-
speed digital video camera (Fastcam Nova R2, Photron, Tokyo, Japan) at 2000 frames s^{-1} at a
resolution of 2048 × 1472 pixels. The camera was fitted with a 60 mm macro lens (Nikon, AF
Micro Nikkor, Nikon, Japan) set at f/5.6, yielding a depth of focus for PIV analyses DoF ≈ 57.1
μm (see supplementary materials). Fluid velocity vectors were calculated from sequential images
analyzed using the DaVis 10 software package (LaVision, Göttingen, Germany). Image pairs were
analyzed with three passes of overlapping interrogation windows (75%) of decreasing size of 64
× 64 pixels to 48 × 48 pixels. All frames were used for the analysis, yielding a separation between
frames (dt) of 5×10^{-4} s. Velocity fields were input to a custom program in MATLAB 2022a
(MathWorks, Natick, MA, USA) that computed the corresponding vorticity fields.
244
Kinematics and morphometrics
Pleopod kinematics data were extracted from image sequences using a custom program in
MATLAB. The profiles of individual pleopods (from P1 to P5) of the left side of each shrimp were
digitized by manually fitting a Bezier curve controlled by two handles. This method accurately
matches the deformation of the pleopods during both the power and recovery stroke. Because the
protopod is rigid, it was digitized as a straight segment, while the endopodite was digitized as a
Bezier curve. We only digitized the endopodite because the exopodite abducts during the power
stroke and rotates out of plane. In contrast, the endopodite remains aligned with the focal plane.
Every 10 frames were used for analysis, yielding a separation between frames (dt) of 5×10^{-3} s.
Pleopod spanwise bending during swimming was quantified using the curvature along the ramal
structures for each leg. Points along the pleopod of length L were specified as fractional distance
from the peduncle-ramal joint. The radius of curvature R was measured at all points along the
pleopod Bezier profiles. We used curvature, κ (the reciprocal of R), normalized to pleopod length
L.

The α angle, defining the pleopod beat amplitude, was calculated by measuring the angle
between the protopod and the line passing through the proximal joint of the protopods of P1 and
P5 (aligned in the direction of swimming). The β angle is traditionally measured between the
protopodite and the line passing through the proximal and distal sections of the ramal structures.
However, because the endopodite and exopodites bend, we instead measured β between the
protopod and the line tangent to the proximal section of the endopodite (corresponding to 20% of
the total length of the endopodite) to account only for the movement due to the peduncle-ramal
joint. A Fast Fourier transform of the α angle over time was used to determine the sequence-
averaged pleopod beat frequency.
Pleopod coalescence was quantified for each pleopod as the ratio of pleopod grouping duration
during a beat (when in contact with at least one adjacent pleopod) to the duration of the
corresponding recovery stroke. Coalescence was discretized using three categories: leading (when
the pleopod in a coalescing group is facing the flow), confined (when the pleopod is surrounded
by an anterior and posterior pleopod), and trailing (when the pleopod in a group is posterior the
others). The phase φ between two consecutive pleopods was measured as the time between the
start of the power stroke of a corresponding pleopod relative to that of its posterior neighbor (the
latter propagating the metachronal wave earlier) and was normalized to its beat period (calculated
as $1/f$ where f is the beat frequency). The ratio of pleopod spacing (B) to pleopod length (L) was
calculated as the distance between the proximal protopodite joint of a corresponding pleopod and
that of its posterior neighbor, divided by the sum of its protopodite and exopodite lengths. Because
P5 does not have a posterior neighbor, B_{P5} was measured as the distance between P4 and P5.
280
Robotic analog
We used the mean leg morphometrics (n = 10 shrimp) and kinematics data for the α and β
angles of one representative shrimp to design and operate a 20× scaled 3D printed five-legged
shrimp robotic analog (Fig. 1, Table 1). The kinematics data from a single specimen demonstrating
horizontal steady swimming was used to drive the robotic leg motions with no destructive
interference between pleopods, as averaging data from multiple specimens masked individual
variations and caused physical contact issues (see supplementary materials). The design and
actuation of the legs are based on the *Pleobot* (Santos et al., 2023), whose kinematics and
hydrodynamics have been validated experimentally against biological data. Here, we fitted the
robot with flexible ramal structures (Practi- Shim™, thickness = 0.0254 mm), whose chord profile
was heat-shaped using the same curvature profile of shrimp pleopods to achieve differential
stiffness (see supplementary methods). Note that because the P1 endopodite is sexually dimorphic,
atrophied, and not involved in propulsion in *P. vulgaris*, the robotic P1 pleopod did not include an
endopodite. The viscosity of the fluid and the beat frequency of the legs were adjusted to match
the Re of the appendages of the five-legged robot to that of the marsh grass shrimp used in in-vivo
experiments ($Re_{robot} \approx Re_{shrimp} = 1720$, see supplementary materials). A 3:2 glycerin-water mix
with density of $\rho = 1160$ kg m^{-3} and dynamic viscosity, $\mu = 0.0114$ kg m^{-1} s^{-1}, was used in the
experiments.
We performed high-speed 2D PIV and force measurements synchronously to temporally match
the kinematics and PIV sequences to the resulting instantaneous forces produced by and acting
upon the pleopods. Measurements were acquired after at least ten consecutive metachronal waves

to ensure the flow was fully established. Forces were measured by a 6-axis Nano 17 force
transducer (ATI Industrial Automation, MI, USA) from which the robot was suspended. The
transducer was interfaced to a data acquisition (DAQ) system (NI USB-6210, National Instrument,
USA, TX) outputting data to a custom MATLAB script that converted raw voltages to force data.
The sensor bias was measured before each experimental trial and subtracted from the force data to
eliminate the potential effects of sensor drift over time. The robot was initially tested in air to
measure the inertial effects during leg motions to account for the added mass during the
experiments. However, because of the low beat frequency and the fact that the pleopods have low
mass by virtue of being made of thin plastic shims, the forces due to inertia were too small to be
distinguishable from the static noise produced by the force transducer under its minimum
measurement threshold (uncertainty = 2.25 mN). Therefore, we considered added mass effects to
be negligible in the experiments.
We used high-speed two-dimensional PIV to compute velocity fields around the robotic legs.
Experiments were carried out in a 210-liter aquarium measuring $119 \times 43 \times 41$ cm (internal length
$\times$ width $\times$ height). The robot was suspended above the surface of the fluid with its pleopods fully
submerged. Lateral recordings were acquired by a high-speed, high-resolution digital video camera
(Fastcam Nova R3, Photron, Tokyo, Japan) at 500 or 750 frames per second (4096 pixels $\times$ 2304
pixels). Seeding particles (10 μm Dantec Dynamics, Skovlunde, Denmark) were illuminated by
two adjacent coplanar vertical laser sheets (532 nm, 4000 mW, OptoEngine LLC, Optotronics
LLC, Mead, CO, USA) to ensure a uniform particle density field along the entire length of the
robot. Both laser sheets pointed upward along the medial plane of the robot (vertical plane passing
through the tips of the endopodites).
Several leg configurations were tested to isolate the effects pleopod asymmetrical bending and
coalescence have on the fluid flows and forces. For instance, we tested individual pleopods (P1–
P5) fitted with flexible or stiff appendages to determine the thrust and drag generated in both cases.
Each pleopod was tested independently by keeping all the other legs horizontally, out of the way
and influence of the flow produced by the selected pleopod. To investigate the effects of
coalescence, we performed five-legged and three-legged experiments. The five-legged cases were
analogous to the free-swimming marsh grass shrimp while the three-legged cases aimed at
precisely reproducing a unique coalescing group of three adjacent pleopods. This was done to
eliminate the effects of the other legs not involved in the corresponding coalescing group. We
report data for the P2-P3-P4 coalescing group. To test the effect of coalescence on the flow and
forces specifically, we compared the results to the instantaneous forces produced by the same, but
independent, legs, time-synchronized to the corresponding metachronal wave of the coalescing
group. The theoretical thrust when coalescence is not occurring was calculated as the cumulative
axial force produced by the individual legs. We also tested cases when the metachronal wave was
phase-shifted by 180 degrees. This unnatural kinematics caused leg contacts during the power
stroke, but complete separation of the pleopods during the recovery stroke. Lastly, to explore the
compounded effect of bending and coalescence, we tested the above leg configurations with
flexible and stiff ramal appendages.
Fluid velocity fields were computed using the DaVis 10.2 software package (LaVision,
Göttingen, Germany). Image pairs were analyzed with three passes of overlapping interrogation

windows (75%) with decreasing size from 64 × 64 pixels to 48 × 48 pixels. No smoothing of the
velocity data was performed.
The force coefficient C_F, whose thrust and drag coefficients (C_T and C_D, respectively) were of
interest in this investigation, was calculated using the axial forces oriented in the swimming
direction, corresponding here with the body axis of the five-legged robot. In this context, the axial
forces generated by individual legs contribute entirely to thrust (in the swimming direction) or
drag (opposite to the swimming direction). The cycle-averaged coefficients were computed over
8–10 consecutive cycles (based on availability of complete cycles at the beginning and end of the
video sequence) from the instantaneous force magnitude T, as:

$$C_F = \frac{T}{\frac{1}{2}\rho U^2 A} \tag{2}$$

where ρ is the density of the 3:2 glycerin-water mix, U is the cycle-average appendage tip velocity,
and A is the stroke-phase dependent total surface area of the appendage. A was maximal during
the power stroke as the exopodite extended fully (see Fig. 1), thus combining the surface area of
the endopodite and exopodite in the calculation of the force coefficients during the power stroke.
During the recovery stroke when the exopodite adducts and overlaps completely with the smaller
endopodite, the total surface area of the pleopod is reduced to the surface of the exopodite only.
The coefficients are calculated using the total surface area of the pleopods rather than their
effective surface area – as if both pleopod types were stiff. This is to highlight the expected
difference in the drag the stiff and bending pleopods produce (due to different effective surface
area) despite having the same kinematics, shape, size and total surface area. U was calculated
assuming the tip of the appendage moves along an arc along the imaging plane during a beat as:

$$U = 2\pi L f \left(\frac{2\theta}{360}\right) \tag{3}$$

where L is the total length of the pleopod (including protopodite and exopodite), f is the beat
frequency, and θ is the appendage stroke amplitude, in degrees, obtained from the α angle maxima
and minima over several consecutive beats. The α angle was measured for each leg relative to the
swimming direction by measuring the leg kinematics using the DLTdv8 package for MATLAB.
This was done by tracking landmarks on the protopodite, such as the distal joint screw. The slight
orientation bias of the rostro-caudal axis relative to horizontal (bias < 1°) was accounted for in the
α computation.

Statistics
All statistical tests were performed using MATLAB 2023b. For biological measurements of
the pleopod chordwise curvature, the rami were pooled by type for the only specimen available
(for a total of 8 endopodites and 10 exopodite). Note that the sexually dimorphic and atrophied P1
endopodite were excluded from this analysis because they differ substantially from the other legs
and are not involved in swimming. Because of the low sample count and the sample distributions
were not normally distributed (One-sample Kolmogorov-Smirnov test, p-value < 0.001), we
compared the differences between the endopodite and exopodite using a Mann-Whitney U-test.

Similarly, the flexural stiffness of the anterior and posterior faces of the endopodite and exopodite
(n = 5 shrimp), was compared using a *U*-test for each leg. The relationship between pleopod
bending and the angular velocity ω of the pleopod defining its overall motion relative to the fluid
was examined separately for the power and recovery strokes. For the recovery stroke, we found
that the relationship between the local curvature κ/L along the ramus of length L pleopod angular
velocity is best described by a power fit ($\kappa/L = a\omega^b + c$) while the power stroke is best modeled as
a linear fit ($\kappa/L = a\omega^b + c$). For robotics experiments, the mean thrust and drag coefficients for each
consecutive beat cycle (for individual legs fitted with flexible or stiff ramal appendages) were
computed for the power and recovery strokes, respectively, to calculate the cycle-average mean
coefficients (from 8 to 10 consecutive cycles). We performed *t*-tests to compare the mean thrust
and drag coefficients of individual flexible pleopods with their stiff counterparts (P1 through P5).
Differences were considered significant at *p*-values < 0.05. All the kinematics parameters and force
data are reported as mean ± standard deviation (s.d.).
399
400
RESULTS
Pleopod morphology and asymmetrical bending
Cross sections of all the pleopods of one representative *P. vulgaris* obtained from reconstructed
μ-CT scans show that the posterior face of the ramal structures — endopodite and exopodite — is
concave, with pronounced chordwise curvature (Fig. 2A). Pooling the data by ramus type revealed
no statistical difference in the normalized curvature *W/R* of the endopodites and exopodites (8
endopodites and 10 exopodites, *U*-test, $p = 0.083$, Fig. 2) with $W/R_{endopodite} = 1.26 \pm 0.21$ and
$W/R_{exopodite} = 1.11 \pm 0.14$ (see Table 1 for *W/R* of individual pleopod). Note that the chordwise
curvature of the left and right P1 endopodites were excluded from analysis because they are
sexually dimorphic, atrophied, and not involved in propulsion. Interestingly, we observed that the
anterior face of the endopodites and exopodites was relatively planar. While this characteristic
may play a role in the bending properties of the rami, we determined that chordwise curvature was
the dominant feature enabling asymmetrical bending. Only the curved profile was considered in
the design of the robotic appendages for ease of fabrication and to simplify the model. The rami
of each robotic pleopod were shaped using the normalized curvature of the rami of the respective
leg measured in *P. vulgaris* (Table 1).
The pronounced posterior chordwise curvature of the rami caused the legs to remain stiff
during the power stroke but bend almost horizontally during the recovery stroke (Fig. 3A). Flexural
stiffness measurements of freshly dissected *P. vulgaris* pleopods (Fig. 3B,C) indicate that the
concave posterior face of the endopodites and exopodites stiffens the rami during the power stroke
but renders them more flexible during the recovery stroke. The rami are, on average, 1.7 times
stiffer when a point force is applied posteriorly (corresponding to the power stroke) compared with
the anterior face (analogous to the recovery stroke), thus resulting in pronounced spatial
asymmetry (*t*-tests, $0.008 < p < 0.222$; n = 5 shrimp; Fig. 3, Table S1). This mechanism is
reminiscent of construction measuring tapes, whose chordwise curvature is, coincidentally, similar
to that of shrimp to keep the tape straight when measuring but flexible to wind it in the casing.
Stiffness measurements were obtained for the membranous part of the rami and excluded the setae.

Pronounced pleopod spanwise bending occurs exclusively during the recovery stroke, as seen with elevated local curvature propagating from the proximal to the distal sections of the rami (Fig. 4A, Fig. S4). The rami do not bend along a singular point. Instead, the inflection point induced in the proximal section at the beginning of the recovery stroke travels distally over time until the end of the recovery phase. The rami remain nearly horizontal during the first half of the recovery phase (during the acceleration phase of the recovering pleopod) but eventually unfurl completely in preparation for the following power stroke when it needs to remain stiff (Fig S5). We found that the rami bent passively — rather than actively — due to the resistive forces of the flow acting on the anterior face during the recovery stroke. While we could not measure the strain the drag forces generated, leading to pleopod deformation, we could infer the resistive effects of the flow via the instantaneous angular velocity. This is because at Re ≈ 1700, the drag acting on the rami due to the rotation of the pleopod about the α joint scales with the square of the velocity of the appendage (the linear velocity at any point along the span of the rami), thus causing strain (i.e., bending) due to hydrodynamic loading to scale with the pleopod angular velocity. By measuring the instantaneous local curvature at a fixed point along the rami during a beat (we selected $0.3L$, n = 6 shrimp), we found that spanwise bending of the rami was proportional to the angular velocity of the pleopod. It was particularly significant during the recovery stroke when the local curvature scaled as a power relationship with pleopod angular velocity ($\kappa/L = a\omega^{b} + c$, Fig. 4B, Fig. S4). In contrast, the rami bent only very little with increasing pleopod angular velocity during the power stroke, which followed a linear trend in this case ($\kappa/L = a\omega + b$, Fig. 4B, Fig. S4). This is because the pleopod must remain stiff to generate maximum thrust.

Individual, isolated robotic pleopods fitted with flexible rami behaved similarly and had comparable fluid-structure interactions to their rigid counterparts during the power stroke (Fig. 5A,C). This is because, in both cases, the pleopods remained stiff to sustain the largest surface area and generate maximum thrust. For each pleopod, flexible and stiff rami generated identical power stroke (PS) tip vortices and had equivalent mean thrust coefficients (t-tests, $0.09 < p < 0.58$, Fig. 5E,F). Both cases shed the PS tip vortex from the pleopod tip (Fig. 5B,D). The most notable differences between the flexible and stiff cases emerged during the recovery stroke. Because the flexible rami bend almost horizontally, they allowed the power stroke wake to circulate unimpeded posteriorly (Fig. 5B). In contrast, the stiff rami disrupted the majority of the PS wake which prevented most of the wake from circulating rearward, causing a significant increase in the drag experienced by the leg. By virtue of bending nearly horizontally, the flexible appendages interacted only marginally with the PS wake and had a reduced effective surface area to the flow compared to the stiff rami. This resulted in a significant reduction of the drag coefficient C_D of P2 through P5 by 41.0, 66.4, 75.8, and 42.6% respectively (t-tests, $p < 0.05$ for P2–P5 and $p = 0.08$ for P1; Fig. 5F). While the flexible rami induced a dominantly rearward wake, the stiff rami shed a recovery stroke (RS) tip vortex carrying forward momentum (Fig. 5C). We did not find evidence that this flow induced fluid-structure interactions during the upcoming power stroke that impacted either the anterior flow along the pleopod or the thrust coefficient significantly. Note that all the pleopods have a positive C_F during the second half of the RS, which corresponds to the deceleration phase of the stroke. Because the robot was tethered, this caused the posterior flow to stagnate and push against the pleopod in the swimming direction, thus inducing thrust. Additionally, unlike the posterior pleopods, P1 and P2 finish and initiate a beat with a significant

positive C_F. This is largely because of the interplay between the effects of the posterior thrust-
producing stagnating flow and the drag-inducing role of the interactions with the PS wake and
induced flow. Because P1 and P2 have a smaller maximum α ($\alpha < 90°$) than P3–P5 they produce
a wake with a less pronounced posterior component which is largely overcome by the forces
induced by the posterior stagnating flow. Note how the initial CF converges to 0 from the anterior
to the posterior pleopods as their respective wake and induced flow becomes progressively more
horizontal. While the flexible rami induced a dominantly rearward wake, the stiff rami shed a
recovery stroke (RS) tip vortex carrying forward momentum (Fig. 5C). We did not find evidence
that this flow induced fluid-structure interactions during the upcoming power stroke that impacted
either the anterior flow along the pleopod or the thrust coefficient significantly.
481
Pleopod coalescence
While pleopod bending plays a crucial role in decreasing the overall drag of each pleopod
during the recovery stroke, it also enhances leg coalescence. Time series of coalescing pleopods
in free-swimming *P. vulgaris* show that flexible pleopods can wrap around one another and
conform to the shape of the surrounding pleopods to enhance coalescence (Fig. 6A, Fig. S5).
Another benefit of flexibility compared with stiff rami lies in the significant reduction in
potentially destructive physical interactions that would occur upon colliding with one another.
Because of the length and proximity of the legs and the phase relative to one another, up to three
pleopods form a tightly sealed coalescing group. Three-legged coalescence emerges from the
specific pleopod-separation-to-length ratio, $B/L = 0.364 \pm 0.014$ (n = 6 shrimp, all pleopods
pooled) and inter-pleopod phase lag $\varphi = 0.18 \pm 0.02$ t/T which causes several legs to perform their
recovery strokes simultaneously (Fig. 6D).
The role of each leg changes during a coalescence cycle. Here, we use P3 as an example. As
P3 initiates its coalescence cycle, it leads the P3-P4-P5 group, thus facing the flow directly. Upon
reaching the anterior P2, it becomes confined between P2 (anterior) and P4 (posterior) which
shields P3 from anterior and posterior flows. The coalescence group becomes P2-P3-P4. Finally,
as P4 separates from the P2-P3-P4 group, P3 trails the P1-P2-P3 group, thus exposing it to posterior
flow. P1 and P2 are exceptions since they only have one posterior and one anterior neighbor,
respectively (Fig. 7).
We use the coalescence ratio (CR), to quantify the duration of leg coalescence for a given
pleopod relative to the duration of its recovery stroke. P2, P3, and P4, which have one adjacent
anterior and posterior pleopods during coalescence, have a CR > 0.9, with $CR_{P2} = 0.95 \pm 0.05$,
$CR_{P3} = 0.92 \pm 0.04$, and $CR_{P4} = 0.93 \pm 0.03$ (n = 6 shrimp, Fig. 5C). The majority of the recovery
stroke is dominated by pleopod coalescence. More importantly, each of these pleopods is confined
between two adjacent legs for 43.6 ± 4.7, 34.8 ± 4.8, and $28.6 \pm 4.2\%$ of their recovery stroke,
respectively, thus guarding individual legs from significant interactions with the surrounding fluid
for about 1/3 of the recovery stroke. P1 and P5, which lead and trail the coalescing group,
respectively since they only have one neighboring appendage, still contribute to coalescence for
about 2/3 of their recovery stroke, given $CR_{P1} = 0.76 \pm 0.08$ and $CR_{P5} = 0.60 \pm 0.06$.*P. vulgaris*
displays more than one metachronal wave at a given time, therefore several coalescence cycles
occur within the span of a complete metachronal wave propagating from P5 to P1 (Fig. 6D). This
is because the anterior legs finish their coalescence cycles while the posterior legs initiate a new

metachronal wave that begins a new coalescence cycle almost immediately after. This further
enhances the overall influence of pleopod coalescence during swimming.
Our robotics data emphasized how inter-pleopod phase and coalescence promotes the most
consistent rearward flux and fluid momentum to maximize thrust (Fig. 7, 8, Fig. S6). When this
arrangement breaks down, such as when artificially reversing the metachronal wave to induce
pleopod divergence during their recovery stroke, the wake becomes more turbulent, the thrust less
consistent, and individual pleopod wakes are discernable (Fig. 7, Fig. S6). To investigate how
coalescence influences force production, particularly during the phase when the pleopods are
closely grouped during their recovery stroke, we tested a three-legged configuration (P2-P3-P4,
Fig. 8) that replicated the conditions within a coalescing group. We compared the instantaneous
forces produced during the coalescence phase by the same, but independent, legs, time-
synchronized to the corresponding metachronal wave of three-legged case. We found that three
coalescing pleopods generate 30.2% more thrust than independent pleopods during the
corresponding coalescence phase (Fig. 8C).
528
529
DISCUSSION
The sequential beating of coordinated appendages makes metachronal swimming an efficient
mode of locomotion by producing steadier fluid flows than a single appendage to achieve the
highest average flux and fluid momentum, thereby maximizing swimming performance. Several
biological adaptations such as appendage spatial asymmetry and phase, and physical phenomena
promoting inter-appendage constructive flow interactions have been identified as the main
mechanisms of efficient thrust production. We found that other functional traits such as the passive
bending of the swimming appendages and coalescence through metachronal coordination are also
crucial in reducing drag; another determinant factor of performance in self-propelled systems.
Using biological and bioinspired robotics data, we provide a comprehensive overview of the
underlying mechanisms of pleopod asymmetrical bending and coalescence to explain their role in
reducing drag during forward swimming. In particular, we explore their individual and joint
contributions to enhancing propulsion.
543
The role of pleopod passive bending
Appendage bending is a ubiquitous functional feature of the recovery stroke in many
metachronal swimmers like copepods, ctenophores, krill, and polychaete worms. This contrasts
with the power stroke during which the appendages remain relatively stiff to increase the surface
area and maximize thrust. Rowing appendages are often designed, either with an articulation or
with asymmetric bending rigidity, such that they flex posteriorly (i.e., during the recovery stroke)
but not anteriorly (i.e., power stroke) (Hessler, 1985; Johnson and Tarling, 2008). In *P. vulgaris*,
similar to krill, the ramal joint modulating the β angle between the rami and the protopodite can
partially contribute to spatial asymmetry between the power and recovery strokes (Murphy et al.,
2011; Santos et al., 2023). However, we found that the spatial asymmetry of the pleopods is
primarily driven by the differential flexural stiffness along the posterior and anterior directions
(Fig. 3). The rami are, on average, 1.7 times more flexible in the posterior direction —

corresponding to the recovery stroke when the resistive force of the flow acts upon the anterior
face.
What is the source of the observed asymmetry? Studies on ciliated microorganisms indicate
that asymmetric waveforms are produced by actively sliding adjacent microtubules within the
propulsor (Dutcher, 2019). This, however, is not applicable to metachronal crustaceans whose
biramous pleopod rami are monolithic structures (Fig. 2, Fig. S1). To date, there is no
morphological evidence suggesting the presence of structures (i.e., muscles, microtubules) capable
of inducing controlled spanwise bending of the rami via active processes. Contrary to organisms
living in the viscous regime, larger organisms can tune their structural (i.e., geometry) and material
properties to the inertial forces they experience to achieve passive deformation under strain (Koehl,
1984; Vogel, 2020; Wainwright and Koehl, 1976). Doing so eliminates the need for active control,
thus greatly simplifying actuation. Detailed μ-CT scans of the pleopods revealed that the posterior
face was concave, with normalized chordwise curvature $\kappa = 1.26\pm0.21$ and 1.11 ± 0.40 for the
endopodites and exopodites respectively (Fig. 2). In plants, such chordwise geometry enables
different compliances under bending with opposite orientations, with the structure bending toward
the concave face (Guzmán et al., 2016; Wei et al., 2022), analogous to the recovery stroke in *P.*
vulgaris. This mechanism is reminiscent of construction site measuring tapes whose chordwise
curvature — which coincidentally have similar values to *P. vulgaris* — gives them strength when
measuring with the concave surface facing upward and resisting gravity but high flexibility for
winding in the casing. The curved geometry of the pleopod thus emerges as a passive strategy to
effectively achieve spatial asymmetry. The pleopods can bend during the recovery stroke when
the hydrodynamic forces resisting the motion overcome the inherently weak stiffness given by its
concave shape. In contrast, this geometry stiffens the pleopods during the power stroke when
resistive forces act in the opposite direction. These effects are further illustrated by the pronounced
change in local spanwise curvature with pleopod angular velocity we observed only during the
recovery stroke (Fig. 4). For Re >1000, drag scales with the square of the velocity of the
appendages (Batchelor, 2000), making the strain due to hydrodynamic loading (i.e., bending)
inherently linked with the motion of the pleopod during a beat. As such, by passively bending
under fluid loading, the flexible pleopods can autonomously reduce their projected surface area to
the flow and become more streamlined during the recovery stroke (Vogel, 2020).
Self-reconfiguration reduces the stress on the pleopod proportionally to the resistive fluid
forces (Alben et al., 2002; Gosselin et al., 2010). We found that bending lowers the drag coefficient
of individual pleopods by 41.0 to 75.8% depending on the pleopod, compared to their stiff
counterpart (Fig. 5). The source of this substantial drop in the induced drag is twofold: 1) bending
reduces the effective surface area to the flow, and 2) bending mitigates fluid-structure interactions,
notably with the rearward-traveling power stroke wake. Because the pleopods work at Re >1000
in *P. vulgaris*, presenting a smaller profile in the flow lowers drag-inducing inertial effects and
their nearly horizontal orientation further trades inertial effects for much weaker shear forces
(Gosselin et al., 2010). Drag reduction is enhanced by limiting interactions with the power stroke
wake that circulates nearly unimpeded posteriorly during the recovery stroke (Fig. 5A–D). For
comparison, stiff pleopods systematically disrupt the power stroke wake whose kinetic energy
induces substantially higher drag, equivalent in most cases to the forces produced during the power
stroke (Fig. 5E). Note that being asymmetrically flexible does not affect performance during the

power stroke as we found no difference in thrust production between the flexible and stiff
treatments. The overall instantaneous force patterns during a beat are consistent with data obtained
with our previous krill analog (Santos et al., 2023), but the contrast between the power and
recovery strokes is stronger because we accounted for pleopod asymmetrical bending. By
maximizing the thrust to drag ratio via spatial asymmetry between the power and recovery phases,
shrimp can effectively overcome the resisting appendage and body drag to achieve propulsion
(Byron et al., 2021; Herrera-Amaya and Byron, 2023; Lou et al., 2022; Vogel, 2020). Producing
less overall drag during the recovery phase reduces the power needed to actuate the pleopod, thus
promoting economical swimming and potentially lowering the cost of transport (COT) (Lionetti
et al., 2023).
609
The role of pleopod coalescence
The phase-dependent grouping of several closely spaced adjacent appendages —
coalescence — is intrinsic to antiplectic metachronal waves (Alben et al., 2010; Byron et al., 2021;
Ford and Santhanakrishnan, 2021). We found that the mean inter-pleopod phase $\varphi = 0.18 \pm 0.02$
t/T and pleopod $B/L = 0.36 \pm 0.01$ of *P. vulgaris* promotes the tight grouping of three leg pairs
during their recovery phase and the complete separation of the legs during their power stroke (Figs.
6, 7). Many metachronal organisms display strikingly similar parameters and kinematics that fall
within relatively narrow ranges — $0.2 < B/L < 0.65$ and $20 < \varphi < 25\%$ (Byron et al., 2021; Garayev
and Murphy, 2021; Murphy et al., 2011). These commonalities are often explained in the context
of the power stroke for which there is a need to synchronize the swimming appendages and
circulate the wake in a manner that promotes efficient force production (Ford and
Santhanakrishnan, 2021; Ford et al., 2019; Murphy et al., 2013). Our robotics data support this
idea that the particular inter-pleopod phase and configuration promotes the highest rearward
average flux and fluid momentum to maximize thrust (Fig. 7, Fig. S6). When this arrangement
breaks down, such as when artificially reversing the metachronal wave to induce pleopod
divergence during their recovery stroke, the wake becomes more turbulent, the thrust less
consistent, and individual pleopod wakes are discernable (Fig. 7, Fig. S6). A recent numerical
study about ctenophores found that appendage tip vortex interactions between recovering adjacent
appendages are equally as important in reducing drag and can improve the thrust-to-power ratio
by up to 55% (Lionetti et al., 2023). As such, by promoting inter-pleopod interactions, coalescence
can have a significant impact on the performance of metachronal propulsion. This potential
tradeoff between pleopod separation and grouping during the different stages of a beat likely
explains why metachronal appendages beat within such narrow phase and *B/L* ranges.
The importance of coalescence in metachronal propulsion is highlighted by its overall
pervasiveness during the recovery stroke. Coalescence was observed spanning over 90% of the
recovery phase of the legs (Fig. 6). Note, however, that while all the legs display coalescence, their
position and role within a coalescing group change sequentially from leading, to being confined,
to trailing the group as the metachronal wave propagates. Consequently, the pleopods are exposed
to changing flow conditions during their beat, which coalescence leverages to reduce the overall
drag.
Pleopods P2–P4 (which have anterior and posterior neighbors) are completely confined within
a group and as such are not directly subject to the flow (Figs. 6AB, 7) for about one third of their

recovery stroke. To investigate how this affects force production, we tested a three-legged
configuration (P2-P3-P4, Fig. 8) that replicated the conditions within a natural coalescing group.
We chose this configuration to eliminate the contribution from the legs external to the coalescing
group of three legs, thus strictly isolating the effects due to coalescence. Contrary to our
expectations that net drag would be produced during the recovery phase of the pleopods, we found
that instead, only net thrust is generated throughout the entire metachronal wave. We compared
these results with the time-synchronized force data from the same three independently tested
pleopods and found that even without coalescence, only net thrust is produced during a
metachronal wave and fluctuates similarly over time (Fig. 8).

The inter-pleopod phase and spatial asymmetry are sufficient to ensure net thrust is always
produced (Ford and Santhanakrishnan, 2021), albeit not steadily. We must inform the reader that
because the robot was tethered, the results potentially slightly overestimate the forces, particularly
thrust. Tethering affects the flow by artificially reducing the system's speed to zero (Murphy et
al., 2013), thus causing water to stagnate posteriorly at the end of the pleopod recovery stroke, thus
partially enhancing the reactive forces of the flow in the thrust direction (swimming direction)
(Santos et al., 2023). Nonetheless, coalescing pleopods produce 30.2% more mean thrust during
the coalescence phase than their independent counterparts. Given that the main difference between
these two cases resides in the strong inter-pleopod interactions only during the recovery phase of
the pleopods, coalescence emerges as an effective strategy to mitigate drag and improve the overall
performance. This is due in part to the strong contact between the pleopods' surfaces forming a
tight seal, shielding the coalescing group from flow instabilities that would otherwise develop
between the legs (Fig. 7, Fig. S5). We observed that the recovery stroke of P5 (initiating the
metachronal wave) does not shed a wake in the free stream when coalescence occurs. Instead, the
posterior bound vortex is consistently circulated behind the next anterior pleopod as the
coalescence group propagates to P1. We observed that the traveling recovery stroke bound vortex
was trapped between the pleopods and the much stronger inferior, rearward-traveling power stroke
wake into which it dissipates at the end of the metachronal wave (Fig 7). This mechanism is
consistent with the vortex-weakening mechanism found in ctenophores that harnesses fluid-
structure interactions among closely spaced appendages to reduce drag (Lionetti et al., 2023).

Overlapping the profile of several appendages via coalescence can reduce the cumulative
effective surface area of multiple appendages arranged in series. But does coalescence turn the
drag of three otherwise separate pleopods to that of only one? As discussed above, we found that
coalescence does not induce net drag at all, which is a great departure from individual legs that all
produced net drag during their recovery phase (Figs. 5, 8). We found that this is primarily due to
the combined effects of the phase and the resulting coalescence. The inter-pleopod phase
intrinsically synchronizes the forces produced by the pleopods such that net thrust is always
generated. However, because it is naturally inducing coalescence — coalescence cannot exist
without the appropriate phase — we could not strictly isolate the independent contribution of
pleopod grouping (i.e., overlapping profiles) and the three phases of coalescence to lowering the
drag. Future experiments designed to measure the instantaneous force output by each leg within a
group could help clarify how coalescence affects their performance relative to their independent
counterparts.

Pleopod bending and coalescence are complementary

As discussed above, pleopod coalescence is intrinsically linked with the *B/L* ratio, the inter-pleopod phase, and beat amplitude. Coalescence occurs by virtue of the spatial and temporal synchronization of several appendages inducing contact when two or more adjacent appendages are oriented in opposite directions (i.e., posteriorly during the power stroke and anteriorly during the recovery stroke). Thus, coalescence can occur even in the absence of appendage bending. However, having flexible pleopods leverages coalescence by enhancing structure-structure and fluid-structure interactions. In addition to the mechanical and hydrodynamic benefits of bending discussed above, because the appendages can deform, they are not as limited by the available space between them and physical interactions as their stiff counterparts, thus enabling wider stroke amplitude (Brennen, 1974). This gain in the plasticity of the motions of the pleopods signifies that physical interactions are not detrimental and might instead be instrumental in promoting constructive physical interactions, such as when the legs coalesce.

While stiff rami can still coalesce, they achieve only partial spanwise contact, often resulting in strong physical interactions that disrupt the geometry of other pleopods during the recovery stroke. These undesired effects induce a turbulent wake and cause significant disturbances in the instantaneous forces produced during coalescence (Fig.7, S6C). In contrast, spanwise bending enables the precise overlapping of several pleopods such that they conform to each others shape and form a tight seal within a coalescing group to prevent inter-pleopod flow (Fig. 6,7, S5). This reduces the impact of the recovery stroke on the overall wake and promotes more consistent thrust production (Fig. S6A). However, one potential challenge the close contact of adjacent pleopod poses is the need to overcome capillary adhesion when separating the trailing pleopod from a coalescing group. Because of the surface tension of water, a thin layer of water is still confined between two adjacent legs, which applies a force acting at the interface between the fluid and the wetted surface. At intermediate Re, the capillary effects can be significant (Butler and Vella, 2020; Lambert, 2013), potentially rendering coalescence less effective.at the moment of pleopod separation. We found that having flexible pleopods solves this problem. The trailing and confined pleopods in a coalescing group (see Fig. 6 for reference) slide against one another and the rami of the trailing leg unfurl progressively along their span during the later phase of the recovery stroke due to direct physical interaction with the anterior leg (Fig. 6, S5). This mechanism is particularly noticeable in Fig. 4A where the instantaneous local spanwise curvature maximum travels from the proximal to the distal section of the rami and is greatest near the tip during the late phase of the recovery stroke. Unlike two rigid flat plates that induce the highest capillary effect when pulled apart in opposite directions, unfurling the trailing pleopod localizes the capillary force at the point of separation rather than the entire surface (Butler and Vella, 2020; Lambert, 2013), thus reducing the force required to separate from the coalescing group. Pleopod bending is crucial to enhance structure-structure and fluid-structure interactions and leverages coalescence from a simple byproduct of inter-pleopod phase, separation and amplitude, to a mechanism optimized for minimizing disruptive hydrodynamic effects, thereby improving overall swimming performance.

While *P. vulgaris* displays perfectly overlapping pleopods during coalescence, appendage contact is not always present in other organisms like ctenophores (Colin et al., 2020; Herrera-Amaya and Byron, 2023) and ciliates (Blake, 2001; Blake and Sleigh, 1974; Sleigh, 1989) whose appendage Re is substantially lower (Re < 100). Still, it is interesting that all the metachronal

swimmers studied to date display some form of flexible appendages that can bend during their recovery stroke, leading to the formation of an array of somewhat overlapping appendages. At higher Re, this configuration combines the reduced total effective surface area of individual appendages to the flow due to bending and the masking effect from the anterior appendage to reduce drag (Alben et al., 2002; Barsu et al., 2016; Gosselin et al., 2010). By enabling close physical contact between the pleopods, shrimp mitigate parasitic flow features that would develop within a loose pleopod cluster. At low Re where viscous forces become more important, the need to prevent inter-appendage flow is not as crucial given these instabilities dissipate quickly. However, achieving appendage overlap during their recovery stroke by maintaining adequate appendage spacing, phase, and amplitude, may serve in reducing surface drag (i.e., friction) and can promote fluid interactions that can reduce drag and improve the thrust-to-power ratio (Lionetti et al., 2023).

CONCLUSIONS

Metachronal swimming is commonly found across many taxa and a wide range of scales, suggesting common biophysical mechanisms driving its success. The performance of metachronal propulsion is generally perceived from the need to maximize thrust production. However, we found that asymmetrically flexible appendages and coalescence (major commonalities among these organisms) also enhance the overall swimming performance by significantly reducing drag. Using experimental methods, we present a comprehensive overview of the underlying morphological, functional, and physical mechanisms to quantify the drag-reduction capabilities of pleopod flexibility and coalescence. We found that the curved cross-sectional profile of the pleopods enabled passive asymmetrical bending during the recovery stroke to reduce the coefficient of drag of each pleopod by 41 to 75.8% relative to stiff pleopods. In the context of metachronal swimming when up to three pleopod pairs interact, coalescence reduces the overall drag of the group such that the mean net thrust produced during the coalescence phase is increased by 30.2%. Nonetheless, because we could not resolve the forces for each pleopod within a coalescing cluster during the transition from leading, to confinement, to trailing the group, the question whether the drag of three overlapping pleopods becomes that of only one remains open. Future robotics experiments, including untethered swimming, might provide the answer and help quantify the energetic benefit of pleopod bending and coalescence to demonstrate their role in achieving high swimming performance and efficiency. Our current result improves our understanding of the fundamental biological and physical components of metachronal propulsion that may aid the development of novel bio-inspired underwater metachronal vehicles.

REFERENCES

Alba-Tercedor, J. and Alba-Alejandre, I. (2019). Comparing micro-CT results of insects with classical anatomical studies: The European honeybee (*Apis mellifera* Linnaeus, 1758) as a benchmark (Insecta: Hymenoptera, Apidae). *Microscopy* **3**, 12–15.

Alben, S., Shelley, M. and Zhang, J. (2002). Drag reduction through self-similar bending of a flexible body. *Nature* **420**, 479–481.

Alben, S., Spears, K., Garth, S., Murphy, D. and Yen, J. (2010). Coordination of multiple
appendages in drag-based swimming. *J. R. Soc. Interface* **7**, 1545–1557.

Barsu, S., Doppler, D., Jerome, J. J. S., Rivière, N. and Lance, M. (2016). Drag
measurements in laterally confined 2D canopies: Reconfiguration and sheltering effect. *Phys.*
Fluids **28**, 107101.

Batchelor, G. K. (2000). *An introduction to fluid dynamics*. Cambrige University Press.
Cambridge, UK: Cambridge university press.

Bhattacharya, R., Saha, S., Kostina, O., Muravnik, L. and Mitra, A. (2020). Replacing
critical point drying with a low-cost chemical drying provides comparable surface image quality
of glandular trichomes from leaves of *Millingtonia hortensis* L. f. in scanning electron
micrograph. *Appl. Microsc.* **50**, 15.

Blake, J. (2001). Fluid Mechanics of Ciliary Propulsion. In *Computational Modeling in*
Biological Fluid Dynamics (ed. Fauci, L. J.) and Gueron, S.), pp. 1–51. New York, NY:
Springer.

Blake, J. R. and Sleigh, M. A. (1974). Mechanics of ciliary locomotion. *Biol. Rev.* **49**, 85–125.

Brennen, C. (1974). An oscillating-boundary-layer theory for ciliary propulsion. *J. Fluid Mech.*
65, 799–824.

Butler, M. D. and Vella, D. (2020). Detachment in capillary adhesion: the relative roles of
tilting and separation. *IMA J. Appl. Math.* **85**, 673–702.

Byron, M. L., Murphy, D. W., Katija, K., Hoover, A. P., Daniels, J., Garayev, K., Takagi,
D., Kanso, E., Gemmell, B. J. and Ruszczyk, M. (2021). Metachronal motion across scales:
current challenges and future directions. *Integr. Comp. Biol.* **61**, 1674–1688.

Colin, S. P., Costello, J. H., Hansson, L. J., Titelman, J. and Dabiri, J. O. (2010). Stealth
predation and the predatory success of the invasive ctenophore *Mnemiopsis leidyi*. *Proc. Natl.*
Acad. Sci. **107**, 17223–17227.

Colin, S. P., Costello, J. H., Sutherland, K. R., Gemmell, B. J., Dabiri, J. O. and Du Clos, K.
T. (2020). The role of suction thrust in the metachronal paddles of swimming invertebrates. *Sci.*
Rep. **10**, 17790.

Combes, S. A. and Daniel, T. L. (2003). Flexural stiffness in insect wings II. Spatial
distribution and dynamic wing bending. *J. Exp. Biol.* **206**, 2989–2997.

Daniels, J., Aoki, N., Havassy, J., Katija, K. and Osborn, K. J. (2021). Metachronal
swimming with flexible legs: A kinematics analysis of the midwater polychaete *Tomopteris*.
Integr. Comp. Biol. **61**, 1658–1673.

Dutcher, S. K. (2019). Asymmetries in the cilia of *Chlamydomonas*. *Philos. Trans. R. Soc. B*
Biol. Sci. **375**, 20190153.

Fabritius, H.-O., Ziegler, A., Friák, M., Nikolov, S., Huber, J., Seidl, B. H. M., Ruangchai,
S., Alagboso, F. I., Karsten, S., Lu, J., et al. (2016). Functional adaptation of crustacean
exoskeletal elements through structural and compositional diversity: a combined experimental
and theoretical study. *Bioinspir. Biomim.* **11**, 055006.

Ford, M. P. and Santhanakrishnan, A. (2021). On the role of phase lag in multi-appendage
metachronal swimming of euphausiids. *Bioinspir Biomim* **16**, 066007.

Ford, M. P., Lai, H. K., Samaee, M. and Santhanakrishnan, A. (2019). Hydrodynamics of
metachronal paddling: effects of varying Reynolds number and phase lag. *R. Soc. Open Sci.* **6**,
191387.

Forward, R. B. (1988). Diel vertical migration: Zooplankton photobiology and behaviour.
Oceanogr. Mar. Biol. Annu. Rev. **26**, 361–393.

Garayev, K. and Murphy, D. W. (2021). Metachronal swimming of Mantis Shrimp:
Kinematics and interpleopod vortex interactions. *Integr. Comp. Biol.* **61**, 1631–1643.

Gosselin, F., de LANGRE, E. and Machado-Almeida, B. A. (2010). Drag reduction of flexible
plates by reconfiguration. *J. Fluid Mech.* **650**, 319–341.

Guzmán, J. E. V., Hernández-Badillo, C. and Zenit, R. (2016). Experimental study of the
deflections of curved plates exposed to pulsating cross-flows. *Acta Mech.* **227**, 3621–3637.

Herrera-Amaya, A. and Byron, M. L. (2023). Omnidirectional propulsion in a metachronal
swimmer. *PLOS Comput. Biol.* **19**, e1010891.

Hessler, R. R. (1985). Swimming in Crustacea. *Earth Environ. Sci. Trans. R. Soc. Edinb.* **76**,
115–122.

Johnson, M. L. and Tarling, G. A. (2008). Influence of individual state on swimming capacity
and behaviour of Antarctic krill Euphausia superba. *Mar. Ecol. Prog. Ser.* **366**, 99–110.

Khaderi, S. N., Den Toonder, J. M. J. and Onck, P. R. (2012). Fluid flow due to collective
non-reciprocal motion of symmetrically-beating artificial cilia. *Biomicrofluidics* **6**,.

Koehl, M. A. R. (1984). How Do Benthic Organisms Withstand Moving Water? *Am. Zool.* **24**,
57–70.

Lambert, P. ed. (2013). *Surface Tension in Microsystems: Engineering Below the Capillary*
Length. Berlin, Heidelberg: Springer.

Lionetti, S., Lou, Z., Herrera-Amaya, A., Byron, M. L. and Li, C. (2023). A new propulsion
enhancement mechanism in metachronal rowing at intermediate Reynolds numbers. *J. Fluid*
Mech. **974**, A45.

Lou, Z., Herrera-Amaya, A., Byron, M. L. and Li, C. (2022). Hydrodynamics of metachronal
motion: Effects of spatial asymmetry on the flow interaction between adjacent appendages.p.
Proceedings of the ASME 2022 Fluids Engineering Division Summer Meeting.

Mauchline, J. (1998). Behaviour. In *The biology of calanoid copepods*, pp. 400–455. London,
UK: Academic Press: London.

Murphy, D. W., Webster, D. R., Kawaguchi, S., King, R. and Yen, J. (2011). Metachronal
swimming in Antarctic krill: gait kinematics and system design. *Mar. Biol.* **158**, 2541–2554.

Murphy, D. W., Webster, D. R. and Yen, J. (2013). The hydrodynamics of hovering in
Antarctic krill. *Limnol. Oceanogr. Fluids Environ.* **3**, 240–255.

Raabe, D., Sachs, C. and Romano, P. (2005). The crustacean exoskeleton as an example of a
structurally and mechanically graded biological nanocomposite material. *Acta Mater.* **53**, 4281–
4292.

Ruszczyk, M., Webster, D. R. and Yen, J. (2022). Trends in stroke kinematics, Reynolds
number, and swimming mode in shrimp-like organisms. *Integr. Comp. Biol.* icac067.

Santos, S., Tack, N., Su, Y., Cuenca-Jiménez, F., Morales-Lopez, O., Gomez-Valdez, P. A.
and Wilhelmus, M. M. (2023). Pleobot: a modular robotic solution for metachronal swimming.
Sci. Rep. **13**, 9574.

Sensenig, A. T., Kiger, K. T. and Shultz, J. W. (2010). Hydrodynamic pumping by serial gill
arrays in the mayfly nymph *Centroptilum triangulifer*. *J. Exp. Biol.* **213**, 3319–3331.

Sleigh, M. A. (1989). Ciliary propulsion in protozoa. *Sci. Prog. 1933-* **73**, 317–331.

Tamm, S. L. (2014). Cilia and the life of ctenophores. *Invertebr. Biol.* **133**, 1–46.

Vogel, S. (2020). *Life in Moving Fluids: The Physical Biology of Flow-Revised and Expanded*
Second Edition. princeton, NJ, USA: Princeton university press.

Wainwright, S. A. and Koehl, M. A. R. (1976). The nature of flow and the reaction of benthic
cnidaria to it. In *Coelenterate Ecology and Behavior* (ed. Mackie, G. O.), pp. 5–21. Boston, MA:
Springer US.

Wei, A., Guo, Z. and Guo, F. (2022). Unveiling the Mechanism Behind the Asymmetric
Bending Compliance of Thin-Walled U-Shaped Strips: A Study Inspired by Plant Leaves. *Acta*
Mech. Solida Sin. **36**, 1–10.

List of figures

Figure 1. Biological and robotic models.

Figure 2. Chordwise curvature of the pleopods.

Figure 3. Differential flexural stiffness of the pleopods of *P. vulgaris*.

Figure 4. Passive asymmetrical bending of a representative pleopod (P3) during a beat.

Figure 5: Flexible ramal structure generate less drag than their stiff counterparts.

Figure 6. Pleopod coalescence.

Figure 7. Time series of the velocity and vorticity fields around pleopod P3 of a free-swimming marsh grass shrimp and the five-legged robot.

Figure 8. Comparison of the instantaneous thrust produced by the flexible coalescing pleopod group P4-P3-P2 and the cumulative forces generated by the same, but individual pleopods.

894
Figure 1. Biological and robotic models. **(A)** Representative free-swimming marsh grass shrimp
(*P. vulgaris*). **(B)** The tethered bioinspired 20× robot with five flexible appendages replicates
pleopod bending and coalescence, and measures forces. We used pleopods representative of the
left side of the animal since the kinematics data was obtained from the lateral views of free-
swimming shrimp exposing the left pleopods. Pleopods are identified from P1 (anterior) to P5
(posterior) for reference. The α, β, and γ angles are shown for reference. The horizontal gray line
is the water surface.
902
903
904
905

Figure 2. Chordwise curvature of the pleopods. (A) Reconstructed µ-CT scan of the right P2
pleopod of a representative specimen of *P. vulgaris* (anterior view). Cross sections show the
chordwise curvature, *W/R* of the posterior face of the exopodite and endopodite, with *W* as the
width and R the radius of curvature (see inset). Note the flat anterior face. **(B)** Mean normalized
chordwise curvature of the posterior face of the endopodites and exopodites (n = 10 legs).
Curvature was measured from chordwise cross sections halfway through the length of each ramus.
Endopodites and exopodites have comparable chordwise curvature (*U*-test, p = 0.083). Note that
the chordwise curvature of the left and right P1 endopodites were excluded because they are
sexually dimorphic, atrophied, and not involved in propulsion.

Figure 3. Differential flexural stiffness of the pleopods of *P. vulgaris*. (A) Differential stiffness keeps the pleopods stiff during the power stroke but causes significant bending during the recovery stroke. **(B)** Posterior and **(C)** anterior measurements of pleopod flexural stiffness. Pleopod deflection due to a point force F_y (applied by a pin) was emphasized here to illustrate the effect of differential stiffness. Note that the force was applied at the tip of the membranous section of the rami rather than the tip of the projecting distal setae. **(D)** The concave posterior face stiffens the rami when a force is applied posteriorly (corresponding to the power stroke). **(E)** In contrast, an anterior force causes buckling, thus enabling the rami to bend. The endopodites **(F)** and exopodites **(G)** are, on average, 1.7 times more flexible during the recovery vs. the power strokes (*t*-tests, $0.008 < p < 0.222$; n = 5 shrimp). The white dot in the violin plots indicates the mean, and the horizontal black line is the median. Shading represents the kernel density estimate for the data. Asterisks indicate statistical differences ($p < 0.05$) between the anterior and posterior faces of the same pleopod.

Figure 4. Passive asymmetrical bending of a representative pleopod (P3) during a beat. (A)
Local curvature along the normalized length of the flexible P3 exopodite during a full beat cycle.
Note that curvature is normalized to the ramus length L. No significant spanwise bending was
observed during the power stroke. This is highlighted by the blue P3 instantaneous profiles that
show that P3 remained stiff during this phase. In contrast, a wave of curvature travels from the
proximal to the distal section of the ramus during the recovery stroke, as seen with the substantial
increase in local curvature over the length of the ramus, over time. The red P3 profiles show how
this increased local curvature travels and increases distally as the pleopod unfurls in preparation
for a new power stroke. **(B)** Local spanwise curvature at $0.3L$ of P3 as a function of angular velocity
during a full beat (n = 6 shrimp). The local curvature increases with the angular velocity of the
pleopod during the recovery stroke following a power relationship ($\kappa/L = 5.84\times10^{-7}\omega1.66 + 0.04$,
$R^2 = 0.78$). In contrast, the rami bend only very little with increased pleopod angular velocity
during the power stroke which is best modeled using a linear fit ($\kappa/L = 1.39\times10^{-5}\omega + 0.04$, $R^2 =$
0.122).

Figure 5: Flexible ramal structure generates less drag than their stiff counterparts.
Instantaneous velocity and vorticity fields of P3 fitted with flexible ramal appendages **(A–B)** and stiff appendages **(C–D)**. Flexible and stiff ramal appendages generate comparable flow during the power stroke (PS) such as an anterior bound vortex and a tip vortex **(A, C)**. Both appendage types produce a posterior bound vortex and a PS tip vortex that sheds upon initiating the recovery stroke (RS) **(B,D)**. Note that the stiff appendages also generate an RS tip vortex. This vortex sheds at the end of the RS, producing a noticeable RS forward wake that is completely absent with the flexible appendages **(A,C)**. The flexible appendages generate a PS wake oriented downward and rearward while it is mostly oriented downward with stiff appendages because they disrupt the PS wake by recovering into it. Every three vectors were plotted for clarity. **(E)** Cycle average force coefficient C_F for flexible and stiff P1–P5 (decomposed as the thrust coefficient C_T when positive and drag coefficient C_D when negative). The data is averaged for several consecutive beat cycles ($n_{flex} \leq 9$ and $n_{stiff} \leq 10$). The colored background indicates the power (blue) and recovery (red) strokes. Shading indicates the standard deviation. Time is normalized to the beat period as t/T. **(F)** Comparison of the mean C_T (Thrust) and C_D (Drag) of the flexible and stiff ramal appendages. Letters indicate significant differences between force types within a species (*t*-tests, $p < 0.05$).

Figure 6. Pleopod coalescence. (A) Up to three adjacent pleopods coalesce during their recovery stroke as a metachronal wave travels forward. **(B)** When pleopods form a coalescing group, the anterior leg leads the movement while the posterior pleopod trails. The pleopod within a coalescing group is confined between the anterior and posterior legs surrounding it. **(C)** The coalescence ratio defines the proportion of the recovery stroke during which pleopods coalesce. P2–P4 coalesce > 90% of the time during their recovery stroke, switching from leading to trailing the group (n = 6 shrimp). **(D)** Pleopod coalescence and kinematics (visualized through the α amplitude of the five pleopods P1-P5) were time-averaged for one full metachronal wave (n = 6 shrimp). The metachronal wave period (T_{wave}) is the time between the beginning of the power stroke of P5 (initiating the wave) and the end of the recovery stroke of P1 (ending the wave), after all the other pleopods have completed their cycle. For P2–P4 with anterior and posterior neighbors, coalescence starts when a recovering posterior pleopod contacts the posterior face, followed by an anterior pleopod contacts the anterior face upon finishing its power stroke. The color code of the pleopod coalescence bars was applied to the α amplitude curves (from light to dark shades from P5 to P1 respectively). Errors bars and shading indicate the standard deviation for coalescence and α kinematics, respectively. Because several metachronal waves overlap within the duration of a complete metachronal wave, several coalescence cycles also take place. The coalescence taking place before and after the main coalescence cycle shown in panel **(D)** is represented using shaded coalescence bars.

Figure 7. Time series of the velocity and vorticity fields around pleopod P3 of a free-swimming marsh grass shrimp and the five-legged robot. (A) *P. vulgaris* perform leg kinematics inducing pleopod coalescence during the recovery phase of each pleopod as a metachronal wave propagates. During the recovery stroke (RS) (t/T_{wave} = 0.51), P3 leads the coalescing group including P4 and P5. A shear layer forms along its anterior face while a tip vortex forms behind P5 (trailing the group). Note that P3 sheds its tip vortex at the end of its power stroke. P3 becomes confined between P2 and P4 (t/T_{wave} = 0.68) and is no longer subject to any flow. Instead, the leading P2 faces the flow. P5's tip vortex moves behind P4. P3 eventually trails the coalescing group and receives P4's tip vortex (t/T_{wave} = 0.71). At the end of its recovery stroke (t/T_{wave} = 79), P3 passes its tip vortex to P2 and sheds no visible wake. **(B)** The five-legged robot fitted with flexible rami and displaying natural kinematics (NK) generates an analogous wake with the same backward orientation and overall similar flow features during the recovery phase of the legs as they coalesce. **(C)** The five-legged robot fitted with stiff rami generates a turbulent wake when the recovering pleopods intercept the power stroke wake. Contrary to flexible pleopods, the

stiff pleopods generate individual wakes with forward and downward components. **(D)** The five-
legged robot fitted with flexible rami but performing reversed kinematics (RK) to prevent
coalescence during the recovery phase of the pleopods generates a backward wake that also has a
strong downward component. Note that the RS wake of each pleopod interacts with the strong PS
flow generated by the pleopods that coalesced during their power phase. The converging flow
induces a downward wake. **(E)** The five-legged robot fitted with flexible rami and performing
reversed kinematics produced wakes with the most forward-oriented components out of all the
above cases. This is because each pleopod shed a posterior bound vortex and a tip vortex at the
end of their individual RS. Yellow arrows represent the overall direction of the pleopod wakes.
For clarity, every two vectors are plotted for *P. vulgaris* and every 3 vectors are plotted for the
robot.

1071

Figure 8. Comparison of the instantaneous thrust produced by the flexible coalescing pleopod group P4-P3-P2 and the cumulative forces generated by the same, but individual pleopods. (A) Instantaneous forces produced by independent legs, time-synchronized to the corresponding metachronal wave when the same three legs beat together in the three-legged robotic configuration. The theoretical thrust when coalescence is not occurring is estimated as the cumulative axial force produced by the legs (black line). Overall, net thrust is produced even when coalescence does not occur. Colored shading indicates the standard deviation for at least 8 consecutive beat cycles. Background grey shading indicates the coalescence phase. **(B)** Instantaneous thrust produced by a coalescing group of three pleopods. Coalescence promotes greater net thrust during the coalescing phase (grey shaded background). Note that the cumulative forces from non-coalescing independent pleopods induces a temporal offset in the force. Kinematics silhouettes of the coalescing group are provided for t/T_{wave} = 0.2, 0.4, 0.6, and 0.8 for reference. Blue profiles indicate the pleopods are performing their power stroke (PS) and red profiles indicate their recovery stroke (RS). **(C)** Coalescing pleopods generate 30.2% more thrust than the same but independent actuating pleopods during the coalescence phase.

List of tables

Table 1. Comparative leg morphometrics of *P. vulgaris* and the robotic analog.

Table 1. Comparative leg morphometrics of *P. vulgaris* and the robotic analog. Mean protopodite, endopodite, and exopodite lengths for *P. vulgaris* are reported as mean ± s.d. (n = 10 shrimp). Endopodite and exopodite chordwise curvature was calculated from cross-sections of three-dimensionally reconstructed CT-scan of the 10 legs of one specimen as the mean of the left and right appendages.

Model	Parameters	P1	P2	P3	P4	P5
P. vulgaris	mean protopodite length (mm)[a]	2.12± 0.11	2.45± 0.15	2.36±0.18	2.08±0.11	1.71±0.11
	mean endopodite length (mm)	N/A	2.26±0.19	2.45±0.25	2.44±0.17	2.06±0.13
	normalized endopodite chordwise curvature[b]	0.61[c]	1.27	1.31	1.03	1.44
	mean exopodite length (mm)	2.56±0.26	2.58±0.25	2.89±0.22	2.93±0.22	2.60±0.25
	normalized exopodite chordwise curvature[b]	1.27	1.24	1.07	1.00	0.99
	pleopod spacing (mm)[d]	N/A	1.64±0.17	1.8±0.10	1.84±0.13	1.76±0.17
Robot	scale	20	20	20	20	20
	protopodite length (mm)	42.4	49.0	47.3	41.5	34.3
	endopodite length (mm)	N/A	45.1	49.0	48.9	41.1
	normalized endopodite chordwise curvature[e]	N/A	1.27	1.31	1.03	1.44
	endopodite chordwise radius (mm)	N/A	7.7	7.7	9.3	5.4
	endopodite surface area (mm²)	N/A	323.54	358.96	341.63	233.85
	exopodite length (mm)	51.2	51.6	57.8	58.5	52.0
	normalized exopodite chordwise curvature[e]	1.27	1.24	1.07	1.00	0.99
	exopodite chordwise radius (mm)	7.9	10.0	13.1	14.1	12.8
	exopodite surface area (mm²)	380.47	473.15	596.99	609.09	487.29
	leg spacing[d] (mm)	N/A	32.7	36.1	36.9	35.3

[a]Measured between the proximal and ramal joints.
[b]Based on the mean of the CT scan of the left and right pleopods of one shrimp.
[c]The P1 endopodite is sexually dimorphic and atrophied in marsh grass shrimp and is not involved in swimming. It was excluded in the robot.
[d]Pleopod spacing reported as the spacing between the particular leg and its anterior neighbor; Measured at the center of two adjacent protopodite proximal joints.
[e]The robotic endopodites and exopodites have the same chordwise curvature as that found in marsh grass shrimp.