1
2 **Control of tongue movements by the Purkinje cells of the cerebellum**
3
4

5 Paul Hage, Mohammad Amin Fakharian, Alden M. Shoup, Jay S. Pi, Ehsan Sedaghat-Nejad, Simon P.
6 Orozco, In Kyu Jang, Vivian Looi, Toren Arginteanu, and Reza Shadmehr
7
8 Laboratory for Computational Motor Control, Dept. of Biomedical Engineering, Johns Hopkins School of
9 Medicine, Baltimore, Maryland 21205, USA
10
Correspondence: Reza Shadmehr, Johns Hopkins School of Medicine, 410 Traylor Building, 720 Rutland
Ave., Baltimore, MD 21205. Email: shadmehr@jhu.edu

Acknowledgements: The work was supported by grants from the National Science Foundation (CNS-
1714623), and the NIH (R01-EB028156, R37-NS128416).

Author contributions: PH and RS conceived experiments. PH, MAF, AMS, JSP, ESN, and SPO collected the
data. PH, MAF, IKJ, VL, and RS analyzed the data. PH made the figures and performed statistical analysis.
RS wrote the manuscript.

Competing interests: None.

Data availability: https://osf.io/WDXU4/

ORCID ID: Paul Hage 0000-0002-7006-9877, Mohammad Amin Fakharian 0000-0002-7741-8689, Alden
Shoup 0000-0003-1206-6587, Simon Orozco 0000-0001-6092-7228, Jay Pi 0000-0003-4036-2877, Ehsan
Sedaghat-Nejad 0000-0001-6732-1296, In Kyu Jang 0000-0002-1832-3743, Vivian Looi 0000-0002-9392-
4178, Toren Arginteanu, 0000-0002-6029-7323, Reza Shadmehr 0000-0002-7686-2569.

Abstract

To quantify the cerebellum's contributions to control of the tongue, we trained head-fixed marmosets
to make dexterous movements, harvesting food from small tubes that were placed orthogonal to the
mouth. We identified the lingual regions in lobule VI of the vermis and recorded from hundreds of
Purkinje cells (P-cells), each in sessions where the subject produced thousands of licks. Most movements
aimed for one of the small tubes, while other movements groomed the mouth area. To quantify
contributions of a P-cell to control of the tongue, we relied on the fact that in a small fraction of the
licks, the input from the inferior olive produced a complex spike (CS), which then briefly but completely
silenced the P-cell. When the movements were targeting a tube, the CS rates increased during
protraction for both ipsilateral and contralateral targets, thus identifying the preferred axis of motion in
the olivary input, termed CS-on. However, for grooming movements this modulation was absent. We
compared the tongue's trajectory in the targeted movement that had experienced the CS with
temporally adjacent targeted licks that had not. When the SS suppression occurred during protraction,
the tongue exhibited hypermetria, and when the suppression took place during retraction, the tongue
exhibited slowing. These effects amplified when two P-cells were simultaneously suppressed. Therefore,
CS-induced suppression of P-cells in the lingual vermis disrupted the forces that would normally
decelerate the tongue as it approached the target, demonstrating a specialization in stopping the
movement. Because the CS-on direction tended to align with the direction of downstream forces
produced during P-cell suppression, this suggests that for targeted licks, the olivary input defined an axis
of control for the P-cells.

Significance statement

During dexterous licking, a CS-induced suppression of P-cells in the lingual vermis inhibited the forces
that would otherwise retract the tongue, resulting in hypermetria during protraction and slowing during
retraction. Because the direction of these forces aligned with the direction of motion specified by the
olivary input, a pattern that is also present for P-cells in the oculomotor region of the cerebellum, the
results imply a general computation for P-cells during control of targeted movements.

Introduction

We use our tongue for verbal communication and mastication, skills that require the brain to control more than 100 muscles simultaneously (1). Damage to the cerebellum disrupts these movements, resulting in abnormal muscle activation patterns (2) that bear a resemblance to ataxia of the arm (3). Indeed, life without a cerebellum in humans (4), or inactivation of the deep cerebellar nuclei in mice (5), do not eliminate tongue movements. Rather, the movements become inaccurate. For example, if the activities of Purkinje cells (P-cell) are disrupted via silencing of molecular layer interneurons, the tongue's trajectory becomes erratic and the subject is no longer able to efficiently harvest liquid rewards (6). Thus, the role of the cerebellum in control of the tongue may be similar to its function during control of the limbs and the eyes: stopping the movement on target. But how might the cerebellum achieve this?

In primates, stimulation of the fastigial nucleus moves the tongue predominantly in the ventral-dorsal axis, while stimulation of the dentate nucleus moves it mainly in the lateral-medial axis (7). Notably, tongue muscles are most readily activated through stimulation of the fastigial nucleus (as compared to the other cerebellar nuclei) (8), suggesting that the P-cells that are located in the vermis play a prominent role in control of the tongue. Unfortunately, there are no published reports of P-cell activity in the vermis during targeted tongue movements in any species, but more is known regarding activity of P-cells in Crus I and Crus II regions (in rodents). For example, before start of a bout, many P-cells in Crus I and Crus II show an increase in their simple spikes (SS), while a smaller number show a decrease (9). Once the licking begins, the SS rates as a population are phase-locked to the rhythm of the lick, with rate maxima occurring near lick onset (5). Complex spikes (CSs) also exhibit their highest rates during protraction (9, 10). The challenge is to understand the relationship between the activities of P-cells and control of the tongue.

Here, we explored this question in marmosets, as they not only have a long tongue (21 mm long, 8 mm wide) (11), but they are also skillful in using it, burrowing into small tree holes, extracting sap and insects. We trained head-fixed marmosets to make thousands of tongue movements to harvest food from small tubes that were placed at 90° to the mouth. We then relied on the fact that the inferior olive not only transmits unexpected sensory events to the cerebellum (12–15), it also transmits information regarding the direction of the planned movement (15, 16). The result is a complex spike that briefly but completely suppresses the P-cell (17, 18), resulting in a movement (19), or disruption of the ongoing movement (20, 21). Thus, the CS events provide an opportunity to ask whether the resulting P-cell suppression is followed by a change in control of the tongue. Indeed, SS suppression disrupted the deceleration phase as the tongue approached the target, as well as the retraction phase as it returned to the mouth.

Regardless of whether the target of the movement was ipsilateral or contralateral to the location of the recording, as a population the CS rates were highest during protraction, termed CS-on. Critically, the CS-on direction of the P-cells, extension of the tongue, coincided with the direction of downstream forces that their SS suppression induced on the tongue's trajectory, protraction. Thus, the encoding of the sensorimotor space by the climbing fiber input coincided with the downstream effects that the SS suppression had on control of the tongue. Because a similar phenomenon is present among P-cells that contribute to control of the eyes (15, 21), our results suggest that across disparate types of movements, the inferior olive imposes a consistent architecture of control upon the cerebellum.

Results

We tracked the eyes and the tongue of head-fixed marmosets (n=3) as they performed visually guided saccades in exchange for food (Fig. 1A). The subjects performed a sequence of saccades, at the end of which we delivered an increment of food (slurry mixture of apple sauce and monkey chow). This food was presented via either the left or the right tube for 50-300 consecutive trials, and then switched tubes. Because the amounts were small (0.015-0.02 mL), and the tubes were located at ±90° with respect to the mouth, harvesting was effortful (22), requiring skillful movements toward a target that was just large enough to accommodate the tongue (4.4 mm diameter tube). As a result, the subjects naturally chose to work for a few consecutive saccade trials (n=6.1±0.02 successful trials per work period), allowing the food to accumulate, then stopped making saccades to targets and initiated their harvest via a bout of licking (n=22.03 ± 0.04 licks per harvest period, Fig. 1B).

The tongue movements were of two general types: in the task relevant licks the subjects aimed for the tube, whereas in the task irrelevant licks the subjects groomed their mouth (Fig. 1A). We further divided the task relevant licks into two subtypes, those that touched the edge of the tube and harvested the food that was near the tip (Fig. 1A, labeled 2&4), and those that penetrated the tube and harvested the food that was deeper (Fig. 1A, labeled 1&5). During a typical 2-3 hour recording session, the subjects performed n=4401±11 task relevant licks (i.e., aimed at a tube, mean±SEM), and n=1310±4 task irrelevant licks (Supplementary Fig. 1). We tracked the motion of the tongue in the horizontal plane using DeepLabCut (23), and combined MRI and CT image guided procedures (24) to place heptodes and silicon probes in lobule VI of the vermis. Over the course of 3.5 years, we recorded from n=284 P-cells (Fig. 1C, Supplementary Fig. 2). A neuron was identified as a definitive P-cell (n=230) because of the presence of complex spikes (CS). In addition, we included data from putative P-cells (n=54) for which we could not isolate the CSs but the neuron was located in the P-cell layer and exhibited 0 ms synchronous simple spike (SS) interactions with other confirmed P-cells (15, 16, 25).

Fig. 1D illustrates the activities of two P-cells. One P-cell increased its SS activity before bout onset, earlier when the tongue targeted the ipsilateral tube, while another P-cell decreased its SS activity, earlier for the contralateral tube (with respect to the site of recording). As the bout began, the CS rates appeared to increase briefly in both P-cells while the SS rates were modulated in a rhythmic pattern that appeared related to tongue kinematics. We focused on three periods during each lick: protraction acceleration period (Fig. 1D, a-b), protraction deceleration period (b-c), and retraction acceleration period (c-d). We then quantified the relationship between the spikes that were generated by a given P-cell and its downstream effects on the motion of the tongue.

The CS-induced SS suppression resulted in overshooting of the tongue during protraction, and slowed return during retraction

As a population, the n=230 confirmed P-cells exhibited CS rates that increased near the onset of protraction (Figs. 2A), peaking around the time the tongue touched the tube, and then decreased below baseline near retraction onset. The rate increases during protraction were larger for ipsilateral licks (within cell difference, protraction period, mean±SEM: 0.077±0.024, t(229)=26.7, p=1.9E-72). Thus, as a population, for both target directions, the CS-on was protraction of the tongue, whereas the CS-off direction was retraction.

As expected, the olivary input suppressed the P-cell's SS production (Fig. 2B, left: all P-cells,
right: single P-cell), lasting an average of 14.8±0.44 ms during licking (time to 85% recovery of SS rate,
mean±SEM). However, the CS events were rare: a CS occurred in only 5.45±0.01% of the licks during
protraction acceleration period (mean±SEM), 9.28±0.01% of the licks during protraction deceleration
period, and 6.59±0.009% of the licks during retraction acceleration period (Fig. 2C). We collected a large
number of licks per neuron (4401±11 licks, Fig. 2D) and then asked whether the CS-induced SS
suppression affected the motion of the tongue.

For each P-cell we considered triplets of consecutive licks $\{n-1, n, n+1\}$ in which all three
licks were of the same type, i.e., contacted the same part of the tube (edge or inner). We then selected
the subset of triplets in which there was a CS at only a single period in lick n, but no CS during any period
in the two neighboring licks $n-1$, and $n+1$. For example, consider tongue trajectories in which there
was a CS in one P-cell during the acceleration period of protrusion (Fig. 3A). This acceleration period was
brief (49.9±0.019 ms), during which the SS activity was normally rising and nearly identical in licks $n-1$
and $n+1$ (Fig. 3B, 1st row), but suppressed during lick n. We measured the change in SS activity by
comparing lick n with licks $n-1$ and $n+1$ and plotted the results of each comparison in Fig. 3B, green
& blue solid lines (labeled suppressed, top row, right column). As a control, we performed a
bootstrapping procedure in which we generated a pseudo data set for each P-cell by randomly assigning
the CS label to a lick and comparing it with its two temporally adjacent neighbors. As expected, in trials
in which there was a CS, the SSs were suppressed during the acceleration period of protraction, but then
rapidly recovered.

To ask whether there were any effects of this brief SS suppression on tongue kinematics, we
measured the distance of the tongue tip to the mouth, and also its angle with respect to the midline,
and then compared the SS suppressed trajectories with the neighboring licks in which a CS was absent.
The effects appeared consistent among the P-cells (Fig. 3A, red colors): following SS suppression, the
tongue was slightly displaced toward the target. Statistical testing, however, which relied on
bootstrapping procedure to compute 95% confidence intervals (CI), demonstrated that the changes
were within the error bounds. That is, the CS-induced suppression during the acceleration period of
protraction had no significant effects on tongue trajectory (Fig. 3C, measured via distance to the mouth,
and its angle, at peak protraction speed and peak displacement).

However, as the protraction proceeded, the effects of SS suppression increased. If the
suppression occurred during the deceleration period of protraction (103.6±0.04 ms), the tongue
exhibited hypermetria (Fig. 3E, second row), producing increased displacement and increased angle of
the tongue's trajectory (Figs. 3F, displacement: 0.37±0.002 mm, 95%CI = [-0.45 0.26] mm, angle:
3.35±0.02 deg, 95%CI = [-2.89 2.30] deg). Notably, the effects were similar regardless of whether lick n
was compared to the previous lick ($n-1$), or the subsequent lick ($n+1$) (Fig. 3E, second row, blue and
green traces). Moreover, the effects were consistent across the P-cells (Fig. 3D, red colors, also
Supplementary Figure 3, left panel), but larger for contralateral licks (within cell difference, ipsilateral
minus contralateral, at lick endpoint, displacement mean±SEM: -0.15±0.04 mm, t(229)=-3.97, p=9.6E-05,
angle: -1.41±0.37, t(229)=-3.79, p=9.69E-05).

These results hinted that SS suppression prevented normal deceleration, producing hypermetria
in the direction of the target. If this interpretation is valid, then a similar suppression during the
retraction period should produce forces that are again in the direction of protraction, now impeding the

tongue's return. That is, if SS suppression during protraction sped the outward movement, then the
same suppression during retraction should slow the inward movement.
As retraction began, the population CS activity (Fig. 2A) had fallen below baseline, i.e., opposite
of the activity during protraction. Yet, if the CS occurred during the acceleration period of retraction
(78.3±0.02 ms), the resulting suppression was again an outward displacement of the tongue (Fig. 3G).
Indeed, comparison of lick n with $n-1$ or $n+1$ revealed a consistent effect: an increased distance and
angle of the tongue to the mouth (Fig. 3I, displacement: 1.29±0.002 mm, 95%CI = [-0.71 0.69] mm,
angle: 4.70±0.02 deg, 95%CI = [-3.18 3.21] deg). These effects were present for ipsilateral and
contralateral licks (Fig. 3H), larger for the contralateral licks (within cell difference, ipsilateral minus
contralateral, at lick endpoint, displacement: -0.37±0.09 mm, t(229)=-3.97, p=4.8E-05, angle: -2.06±0.50,
t(229)=-4.1, p=2.71E-05), and consistent across the P-cells (Fig. 3G, red colors). The effect of suppression
on lick trajectory for each P-cell is shown in Supplementary Figure 3, right panel. Thus, the CS-induced SS
suppression during retraction slowed the return of the tongue.
In summary, when a CS-induced SS suppression occurred during the deceleration period of the
tongue's protrusion, the tongue overextended toward the target. When this suppression occurred
during retraction, the tongue's return was slowed. These effects were present for targets on both the
ipsilateral and contralateral sides, but greater when the target was contralateral to the recording. Thus,
it appeared that SS suppression disrupted the motor commands that would normally stop the tongue
during protraction and return the tongue during retraction.
197
Perturbations scaled when pairs of P-cells were simultaneously suppressed
Our dataset included n=298 pairs of simultaneously recorded P-cells. This provided the opportunity to
test whether near simultaneous suppression of 2 P-cells had a greater effect on tongue kinematics as
compared to when only one of the two P-cells was suppressed.
As before, we collected triplets of consecutive licks $\{n-1, n, n+1\}$ where all the licks were of
the same type (directed toward the same part of the tube), but only lick n had a CS. We divided the
triplets based on whether a CS was present in only one of the P-cells, or both P-cells, then computed
trajectory differences between licks n and $n-1$, as well as licks n and $n+1$. Finally, for each pair of P-
cells we averaged $n-(n-1)$ and $n-(n+1)$ to increase statistical power (because there were far
fewer licks in which both P-cells experienced a CS during the same period of the movement).
We found that if the CS occurred in both P-cells during the deceleration phase of protraction,
there was significantly greater displacement of the tongue, and its angle, as compared to when only one
of the P-cells was suppressed (Fig. 4A, angle: t(248) = -3.7167, p = 2.5E-04, displacement: t(248) = -
5.5471,p = 7.43E-08). Similarly, when the suppression occurred during retraction, the return phase of
the movement experienced a greater slowing in the case of 2 P-cells as compared to a single P-cell (Fig.
4B, angle: t(192) = 6.34, p = 1.6tE-09, displacement: t(192) = 6.50, p = 6.79E-10). Thus, near
simultaneous suppression of 2 P-cells roughly doubled the kinematic effects, further disrupting
deceleration and return of the tongue.
216
Control studies
The fact that a CS was present during a given period in lick n may have been because earlier in the
tongue's trajectory there was an event (for example, an error), that affected that movement, increasing

the likelihood of a CS, and resulting in compensatory movements that followed the CS. To check for this,
for each period during which we observed a CS we considered the tongue's trajectory in the same lick
but during the period preceding the CS. For example, for the licks in which there was a CS in the
protraction deceleration period we focused on the acceleration period of the same movement. By
comparing the lick in which the CS had occurred with its two neighbors, we found that in the period
before the CS had occurred tongue kinematics remained within the 95% confidence intervals of chance:
distance to mouth and angle of the tongue at peak speed were not different than chance (Fig. 3F,
displacement: -0.11±0.001 mm, 95%CI = [-0.22 0.23] mm, angle: 0.19±0.01 deg, 95%CI = [-1.36 1.45]
deg).
Next, we checked the tongue movements in which there was a CS in the retraction period. We
compared the tongue's trajectory in the protraction period of the same lick at peak protraction speed,
and at peak protraction displacement. We found that before the CS had occurred, at the onset of the
retraction the distance to mouth and tongue angle were not different than chance (Fig. 3I,
displacement: 0.01±0.002 mm, 95%CI = [-0.37 0.35] mm, angle: 0.68±0.02 deg, 95%CI = [-3.05 2.80]
deg). Moreover, during the preceding protraction in the same lick, at peak tongue speed distance and
angle were again not different than chance (displacement: -0.66±0.003 mm, 95%CI = [-0.70 0.74] mm,
angle:-2.16±0.02 deg, 95%CI = [-3.35 3.58] deg). Yet, if during the return phase a CS was present, the
movement was slowed (Fig. 3I, displacement: 1.29±0.002 mm, 95%CI = [-0.71 0.69] mm, angle:
4.70±0.02 deg, 95%CI = [-3.18 3.21] deg). That is, the tongue's trajectory before the CS was not
significantly different than the neighboring licks, but after the CS-induced SS suppression the trajectories
diverged.
As a further control, we considered movements in which there was a CS event just before
protrusion onset (Supplementary Figure 4). In these movements, the SS rates were suppressed, but
importantly, the rates recovered around 10 ms after lick onset (Supplementary Figure 4, 1st row).
Critically, despite the presence of a CS just before the lick had started, the SS rates during the entire
protrusion and retraction periods were indistinguishable from licks that had not. The trajectory of the
tongue as measured via distance and angle remained within the 95% confidence interval bounds of
chance (Supplementary Figure 4, 3rd and 4th rows).

Effect of SS suppression remained consistent across P-cells regardless of SS modulation

We had observed that for nearly every P-cell, during both protraction and retraction, the CS-induced SS
suppression was followed by an extension of the tongue (Figs. 3D & 3G). This consistency was surprising
in light of the diversity that was present in the SS patterns: before the onset of the bout, some P-cells
had increased their SS rates with respect to baseline, while others had decreased (Fig. 1D). Did the
kinematic effects of SS suppression differ in these two groups?
In our data set of n=142 of P-cells with SSs, most cells (n=123) increased their SS rates before
the onset of the bout, but a minority exhibited a decrease (n=31) (Figs. 5A and 5B). We separated the P-
cells into SS increasers and decreasers and found that despite the differences in their SS patterns, the CS
rates increased in both groups near bout onset (Fig. 5C, peak CS firing rate change from baseline, SS
increasers: 0.15±0.04 Hz, SS decreasers: 0.26±0.10 Hz). We then compared tongue trajectories in
triplets of consecutive licks $\{n-1, n, n+1\}$, finding that during protraction, following SS suppression in
the deceleration period, in both groups of P-cells the tongue trajectory was displaced away from the

mouth, exhibiting a greater distance and angle (Fig. 5D, displacement: 0.38±0.005 mm, 95%CI = [-0.1
0.049] mm, angle: 3.72±0.05 deg, 95%CI = [-0.62 0.43] deg, SS decreasers: displacement: 0.45±0.04 mm,
95%CI = [0.04 -0.06] mm, angle: 4.28±0.54 deg, 95%CI = [0.42 -0.45] deg). Similarly, during retraction, SS
suppression in both groups of P-cells produced a slowing of the tongue's return, resulting in a greater
distance and angle (Fig. 5E, displacement: 1.25±0.01 mm, 95%CI = [-0.12 0.12] mm, angle: 4.45±0.05
deg, 95%CI = [-0.53 0.56] deg, SS decreasers: displacement: 1.04±0.04 mm, 95%CI = [0.07 -0.07] mm,
angle: 3.48±0.17 deg, 95%CI = [0.37 -0.35] deg).
269   Thus, despite the differences in the patterns of SS activity during licking among the P-cells, the
270   downstream effect of CS-induced SS suppression was consistently an extension of the tongue.
271
272   *The population SS rates encoded parameters of tongue kinematics*
273   In order to organize the P-cells into a population, we relied on the fact that the downstream effects of
274   SS suppression, for both increasers and decreasers, in both ipsilateral and contralateral directions, were
275   to produce forces that extended the tongue. Thus, we combined all cells and computed a population
276   response during various phases of tongue movements.
277   Across the cells, the CS rates peaked at approximately the timepoint of maximum protraction
velocity (Figs. 6A, left column), exhibiting a greater rate for movements toward the ipsilateral side
(ipsilateral: 0.21±0.02 Hz, contralateral: 0.16±0.02 Hz, within cell difference, average CS rate, ipsilateral
minus contralateral, $t(229)=28.1$, $p=2.5E-76$). In contrast, the CS rates declined below baseline before
the onset of retraction (Fig. 6B, left column). Thus, the CS-on direction across the P-cells was
protraction, whereas CS-off was retraction, suggesting that the CS-on direction aligned with the
downstream effects of SS suppression.
284   To analyze the SS activities as a population, we had to consider the fact that during a bout, as
the SS rates modulated about a mean, the mean was not stationary (Fig. 5A). Rather, the mean SS rates
rose or fell before bout onset, then drifted back toward the values before start of the bout, reaching
pre-bout levels at bout ended. Thus, to quantify modulation of SS rates, for each P-cell we considered a
2 sec moving window to compute the running average of its firing rate, then computed the SS rates with
respect to this mean. The 2 sec window was chosen because it was roughly an order of magnitude larger
than the duration of a typical lick.
291   If a suppression in SS rates produced downstream forces that pulled the tongue outwards
toward the target, then during deceleration the SS rates should increase, thus commanding forces that
would stop the motion of the tongue. Indeed, the SS rates peaked near protraction peak speed, and
were larger for contralateral licks (Fig. 6A, right column, ipsilateral: 11.58±1.75 Hz, contralateral:
12.91±1.95 Hz, within cell difference, $t(156)=13.6$, $p=2.9E-28$). The rates began declining at retraction
onset and reached a minimum near retraction peak speed (Fig. 6B). To test if this modulation varied
with tongue kinematics, we quantified lick vigor, defined as the peak speed of a given protraction with
respect to the speed expected for a lick of the same amplitude (22, 26) (Supplementary Figure 5). High
vigor licks exhibited greater SS rates at peak protraction speed (Fig. 6D, within cell difference, high vigor
minus low vigor, 2.92±0.60 Hz, $t(156)=14.1$, $p=1.2E-29$).
301   Given that SS suppression during protraction led to increased extension, if we interpret the
increase in SS rates as commanding downstream forces to induce retraction, then the increased SS rates
during high vigor licks is consistent with the idea that the P-cells are firing to stop the protrusion. If true,

then protractions that have the same peak speed, but a longer deceleration, should not only exhibit
greater SS rates at peak speed, but also a longer duration of activity above baseline. Indeed, we found
that licks that had similar peek speed but longer decelerations tended to also have a longer period of SS
rate increase (Fig. 6F, time to baseline crossing, high duration: 126.80±7.68 ms, low duration:
86.02±7.23 ms, paired t-test, t(156)=7.84, p=6.75E-13).
All of these results were for tongue movements in which the target was a small tube. Would the
same pattern be present for movements that did not require such precision? To answer this question,
we turned to the task irrelevant, grooming licks. These licks tended to be slower, with a peak speed that
was roughly half the speed of the task relevant movements (Fig. 6E, light and dark shaded regions, task
relevant: 290.5±3.6 mm/s, task irrelevant: 153.8±2.5 mm/s). Remarkably, during protraction of the
grooming licks the CS rates were essentially at baseline: the CS rates were an order of magnitude
smaller than when the movements were task relevant (Fig. 6F, left, paired t-test, reward vs grooming,
combined directions, t(229)=-3.27, p=0.00126).
Similarly, the SS rates during protraction were much smaller during the task irrelevant
movements as compared to the movements that were task relevant (Fig. 6E, right, paired t-test, reward
vs grooming, combined directions, t(156)=8.6, p=8.44E-15). In sharp contrast, during retraction of the
task irrelevant licks, both the CS and the SS rates were modulated below baseline by amounts that were
roughly comparable to the rates of the task relevant licks. That is, the fundamental difference in the P-
cell activity between the tube directed and grooming licks was in the protraction phase, but not
retraction. This suggests that the difference in the engagement of the P-cells during these two kinds of
licks was unlikely to be due to kinematics (i.e., velocity).
In summary, the CS-on direction across the P-cells was protraction, whereas CS-off was
retraction. As a result, the CS-on direction in the population aligned with the downstream effects of SS
suppression, i.e., protrusion. The SS rates peaked during protraction, and were greater when the
movement had greater speed or required a greater period of deceleration. In contrast, grooming licks
exhibited little or no CS and SS modulation during protraction.
330
Discussion
What does the cerebellum contribute to control of tongue movements? Here, we found that when an
olivary input arrived at a P-cell, the resulting suppression prevented the normal deceleration of the
tongue as it approached the target, resulting in hypermetria. When the same suppression occurred
during retraction, the tongue's return was slowed. These results were consistent for both ipsilateral and
contralateral targets. Notably, if the suppression was present in pairs of simultaneously recorded P-cells,
the effects were magnified. Thus, regardless of whether the P-cell suppression occurred during
protraction or retraction, the downstream effects were forces that pulled the tongue outwards. Because
during unperturbed movements the SS rates reached their maximum at peak velocity of protraction, we
infer that the consequence of this activity was to decelerate the tongue. This implies that the
contributions of the cerebellum to control of tongue movements are similar to that of the eyes (27, 28)
and the limbs (29): stopping the movement on target.
Like the SS rates, the CS rates peaked near protraction peak velocity, then fell below baseline
before the onset of retraction. Thus, for both ipsilateral and contralateral movements, the "CS-on
action" across the P-cells was protrusion, while the "CS-off action" was retraction. Notably, the

downstream effects of SS suppression were to extend the tongue. Thus, there was a correspondence
between the vector that described the CS-on action, and the vector that described the effects of SS
suppression. This fact is notable because the same principle holds for P-cells in the oculomotor region of
the cerebellum during saccadic eye movements (21): the olivary input to an oculomotor P-cell is most
active when a saccade is planned in direction CS-on, and the downstream effects of that P-cell's SS
suppression is to pull the eyes also in direction CS-on. Thus, for both eye movements and tongue
movements, the olivary input dictates a coordinate system with which to decode the downstream
contributions of a P-cell to control of that movement (30, 31).
Notably, not all tongue movements engaged the cerebellum. Rather, when the purpose of the
movement was to groom the face, during the protraction phase the CS rates remained at baseline while
the SS modulations were reduced. Similarly, during task relevant saccades both the CS and the SS rates
were strongly modulated, whereas during task irrelevant saccades the CS rates were at baseline and the
SS modulations were damped (15, 16). Thus, the strong engagement of the inferior olive and the P-cells
during precision tongue movements were missing during grooming licks.
360
The origin of climbing fiber activity at movement onset
We were surprised that the CS rates peaked not after the tongue contacted the tube, but around the
time that the protraction began. This observation reproduced findings of Welsh et al. (10) in rats, who
recorded from the lateral cerebellum and found that the CS rates peaked near the onset of tongue
protrusion, even when the tongue was deafferented. Indeed, in many types of movements, including
walking (32), reaching (33–37), and moving the wrist (38), the CS rates peak near movement onset. For
example, in the oculomotor vermis, near saccade onset the olivary input informs the P-cells regarding
the direction of the upcoming movement (15). How might the inferior olive be involved in transmitting
movement information to the cerebellum?
A key observation is that the inferior olive not only receives input from the superior colliculus
(39–41), but that subthreshold stimulation of a region of the colliculus leads to CS production without
producing a movement (42). Thus, it is possible that the increased CS rates around the onset of
protraction reflect activities of regions that initiated the tongue movement, i.e., the motor cortex and
the superior colliculus (43). This prediction remains to be tested with simultaneous recordings from the
colliculus, motor cortex, and the cerebellum.
376
Preparing to initiate a tongue movement coincided with ramping activity in the P-cells
As the subject prepared to initiate a licking bout, the P-cells exhibited a ramping activity. Previous
reports have noted that during this period there is ramping activity in the tongue regions of the motor
cortex (44), as well as in the fastigial (44) and the dentate nuclei (45). Inhibiting the fastigial during the
ramping period disrupts planning of the movement and removes the selectivity that the motor cortical
cells exhibit for direction of tongue movements (44), while inhibiting the dentate disrupts the ramping
activity in the motor cortex (45). Similarly, exciting the P-cells in the vermis abolishes the ramping
activity in motor cortex during the delay period of a decision-making task (46). This implies that as one
prepares to initiate a movement, the rising activity in the motor cortex is controlled via a loop through
the cerebellum. Because most of the P-cells in the vermis exhibited increasing activity during the delay
period, we speculate that their role is to reduce the rate of rise in the motor cortex, effectively delaying
the onset of the bout.

Using the olivary input to infer P-cell contributions to control of behavior

Complex spikes are rare neural events, occurring at around once per second. They briefly and
completely suppress the SS rates, which induce downstream effects on the cerebellar nuclei (47),
potentially producing movements (19). However, the CS rates are modulated to encode the direction of
sensory prediction errors (12, 15, 48, 49). Thus, it was critical to test that the kinematic effects that we
measured following a CS-induced SS suppression were not a consequence of a feedback response to
kinematic deviations that occurred before the CS.

We did this by comparing triplets of temporally adjacent licks, finding that while the kinematics
preceding the CS event remained within the chance error bounds, the kinematics that followed were
robustly different than chance. Notably, the downstream effects of the CS-induced suppression were in
the same direction, i.e., extension of the tongue, regardless of whether the CS events occurred during
protraction or retraction. This is notable because the CS rates were maximum during protraction, and
minimum during retraction, yet their kinematic effects were the same.

The idea that the olivary input can alter ongoing movements was noted by Ebner and colleagues
during reaching movements (20), and subsequently observed during saccades (21). For example, during
saccades following the CS-induced SS suppression, the eyes are pulled in the CS-on direction of the P-
cell. This is consistent with the fact that optogenetic increase in the SS rates suppresses the cerebellar
nuclei (50), pulling the eyes approximately in direction CS+180 (51). While relying on the stochasticity in
the olivary input has the disadvantage of lacking a causal manipulation, it has the advantage of relating
the kinematic effects to CS-on properties of each P-cell, something that would not be possible with large
scale optogenetic stimulation.

Methods

Data were collected from three marmosets, *Callithrix Jacchus*, 2 male and 1 female, 350-370 g, subjects 125D (Mirza), 59D (Ramon), and 132F (Charlie), during a 3.5-year period. The marmosets were born and raised in a colony that Prof. Xiaoqin Wang has maintained at the Johns Hopkins School of Medicine since 1996. The procedures on the marmosets were approved by the Johns Hopkins University Animal Care and Use Committee in compliance with the guidelines of the United States National Institutes of Health.

Data acquisition

Following recovery from head-post implantation surgery, the animals were trained to make saccades to visual targets and rewarded with a mixture of applesauce and lab diet (24). Visual targets were presented on an LCD screen. Binocular eye movements were tracked at 1000 Hz using EyeLink in subject R and M, and 2000 Hz using VPIX in subject C. Tongue movements were tracked with a 522 frame/sec Sony IMX287 FLIR camera, with frames captured at 100 Hz.

We performed MRI and CT imaging on each animal and used the imaging data to design an alignment system that defined trajectories from the burr hole to various locations in the cerebellar vermis (24), including points in lobule VI and VII. We used a piezoelectric, high precision microdrive (0.5 micron resolution) with an integrated absolute encoder (M3-LA-3.4-15 Linear smart stage, New Scale Technologies) to advance the electrode.

We recorded from the cerebellum using quartz insulated 4 fiber (tetrode) or 7 fiber (heptode) metal core (platinum/tungsten 95/05) electrodes (Thomas Recording), and 64 channel checkerboard or linear high density silicon probes (M1 and M2 probes, Cambridge Neurotech). We connected each electrode to a 32 or 64 channel head stage amplifier and digitizer (RHD2132 and RHD2164, Intan Technologies, USA), and then connected the head stage to a communication system (RHD2000 Evaluation Board, Intan Technologies, USA). Data were sampled at 30 kHz and band-pass filtered (2.5 - 7.6 kHz).

Behavioral protocol

Each trial began with fixation of a center target after which a primary target appeared at one of 8 randomly selected directions at a distance of 5-6.5 deg. As the subject made a saccade to this primary target, that target was erased, and a secondary target was presented at a distance of 2-2.5 deg, also at one of 8 randomly selected directions. The subject was rewarded if following the primary saccade, it made a corrective saccade to the secondary target, landed within 1.5 deg radius of the target center, and maintained fixation for at least 200 ms. The food was provided in two small tubes (4.4 mm diameter), one to the left and the other to the right of the animal. A successful trial produced a food increment in one of the tubes and would continue to do so for 50-300 consecutive trials, then switch to the other tube. Because the food increment was small, the subjects naturally chose to work for a few consecutive trials, tracking the visual targets and allowing the food to accumulate, then stopped tracking and harvested the food via a licking bout. The subjects did not work while harvesting, and often fixated the tube. As a result, the behavior consisted of a work period of targeted saccades, followed by a harvest period of targeted licking, repeated hundreds of times per session.

We analyzed tongue movements using DeepLabCut(52). Our network was trained on 89 video recordings of each subject with 15-25 frames extracted and labeled from each recording. The network was built on the ResNet-152 prE-trained model, and then trained over 1.03×10^6 iterations with a batch size of 8, using a GeForce GTX 1080Ti graphics processing unit. A Kalman filter was further applied to

456   improve quality and smoothness of the tracking, and the output was analyzed in MATLAB to quantify
457   varying lick events and kinematics. We tracked the tongue tip and the edge of the food in the tube, along
458   with control locations (nose position and tube edges). We tracked all licks, regardless of whether they
459   were aimed toward a tube, or not.
460
Neurophysiological analysis
We used OpenEphys (53) for electrophysiology data acquisition, and then used P-sort (25) to identify the
simple and complex spikes in the heptodes and tetrodes recordings, and Kilosort and Phi (54) to identify
the spikes for the silicon probes. Simple and complex spike instantaneous firing rate were calculated
from peri-event time histograms with 1 ms bin size. We used a Savitzky–Golay filter (2nd order, 31
datapoints) to smooth the traces for visualization purposes. CS baseline firing rates were computed by
dividing the total number of spikes by the duration of the entire recording. SS baseline firing rates were
computed using two different methods depending on the analysis. For bout related responses, baseline
was defined as the average firing rate in a 300 ms window preceding bout onset by 700 ms, i.e. during
the [-1000 to -700] ms. However, to analyze the activities during individual licks, because the rates were
not stationary but gradually changing from the first to the last lick in the bout, baseline SS rates were
computed using the average firing within a sliding window of 2 seconds, consisting of roughly 5 licks.
473
Computing the kinematic effects of CS-induced SS suppression
For each P-cell we considered triplets of consecutive tubE-directed licks $\{n-1, n, n+1\}$, where all
three licks were of the same type, i.e., contacted the same part of the tube (edge or inner). We then
selected the subset of triplets in which there was a CS at only a single period in lick n, but no CS during
all periods in the two neighboring licks $n-1$, and $n+1$. We then compared tongue trajectories
between the lick that had a CS with the two neighboring licks, i.e., $n-(n-1)$ and $n-(n+1)$.
480
Statistical analysis
In order to compute the bounds for the null hypothesis, we used bootstrapping to compute 95%
confidence intervals. We shuffled the assignment of CS tags from the lick in which it had occurred to a
randomly assigned lick of the same type. Using this pseudo-data, we then selected triplets of
consecutive tubE-directed licks and computed trajectory differences among neighboring licks, averaging
$n-(n-1)$ and $n-(n+1)$. We computed this expected value for each cell, computed a mean across
all the cells, and then repeated the shuffling 30 times to compute 95% confidence intervals.
488

489
Figure 1. Marmosets produced dexterous tongue movements during recordings from lobule VI of the cerebellar
vermis. A. Subjects made saccades to visual targets and received a small amount of food as reward via one of two
tubes placed obliquely to the mouth. In the task relevant licks, they directed their tongue to the edge of the tube
to harvest food near the tip (trajectories 2, 4), or inside the tube to harvest food that was deeper (trajectories 1,
5). In task irrelevant licks, they groomed their face (trajectory 3). **B**. Subjects chose to work for consecutive trials,
making saccades and allowing the food to accumulate, then harvested the food in bouts of licking. **C**. We employed
silicon probes to record from lobule VI of the vermis. **D**. Simple and complex spikes (SS, CS) of two Purkinje cells,
aligned to bout onset and lick onset. A single lick was divided into acceleration period of protraction (a-b),
deceleration period of protraction (b-c), and acceleration period of retraction (c-d). Filled color regions indicate
tongue velocity.
500

501
Figure 2. CS rates increased with protraction and decreased with retraction. A. CS rates across the population
aligned to protraction onset, touch of the tube, and retraction onset. **B**. CS-induced SS suppression, averaged
across the subset of P-cells for which both the CS and the SS were isolated (left). Examples of SS and CS waveforms
for a single P-cell are shown at right. **C**. CS activity across all P-cells, aligned to protraction, touch, and retraction,
sorted based on CS rate for contralateral licks at lick onset. **D**. Percentage of licks with a CS during a specific period
of time for each neuron. **E**. Number of task relevant licks per neuron. Error bars are SEM.
508

509
Figure 3. CS-induced SS suppression produced hypermetria during protraction and slowing during retraction. A.
Suppression during the acceleration period of protraction (orange bar). Traces show average tongue trajectory for
each P-cell during suppressed and control licks. Ipsilateral licks are shown to the left and contralateral to the right.
Heatmap quantifies change in endpoint trajectory between suppressed and control licks for each cell. **B**. Top row:
SS rates for licks $\{n-1, n, n+1\}$, where only lick n experienced a CS. Filled color curves indicate tongue velocity.
Second row: trajectory of the tongue in lick n as compared to its two temporally neighboring licks. Trajectory is
measured via distance from tip of the tongue to the mouth and angle of the tip with respect to midline. The filled
region is 95% CI. **C**. Distance to mouth and angle in lick n as compared to neighboring licks. Shaded region is 95%
CI. **D-F**. Suppression during the deceleration period of protraction induces hypermetria. Same format as in part A-

C. **G-I.** Suppression during the acceleration period of retraction induces slowing. Same format as in part A-C. Error
bars are SEM.
521
522

523
Figure 4. Suppression of multiple P-cells scaled the perturbation to the tongue. **A**. Kinematic effects of CS-
induced P-cell suppression during the deceleration period of protraction (orange bar). Traces show change in
tongue trajectory in suppressed licks vs. control licks, measured via distance from tip of the tongue to the mouth
and angle of the tip with respect to midline. Gray shaded region is 95% CI. Brown filled region is tongue velocity. **B**.
Kinematic effects of P-cell suppression during the acceleration period of retraction (orange bar). Same format as in
part A. Error bars are SEM.
530

531
Figure 5. Effects of suppression remained consistent across P-cells regardless of SS modulation. A. Some of the P-
cells exhibited an increase in SS rates before bout onset, while others exhibited a decrease. L1 is first lick in the
bout, Ln is last lick. **B**. Change in SS rates for all P-cells (with respect to baseline), aligned to bout onset. **C**. Change
in complex spike rate for the same P-cells. **D**. Kinematic effects of SS suppression during the deceleration period of
protraction (orange bar). Shaded region is 95% CI. **E**. Kinematic effects of SS suppression during the acceleration
period of retraction (orange bar). Error bars are SEM.
538

539
Figure 6. SS rates were greater for high vigor and long deceleration licks. **However, little modulation was present**
during task irrelevant tongue movements. A. CS and SS rates across the population aligned to protraction peek
speed. As a population, the direction of CS-on was protraction. **B**. CS and SS rates aligned to retraction onset. **C**.
Left figure shows SS rates for high and low vigor protractions (same amplitude). Right figure shows within cell
difference in SS rates for high minus low vigor licks. Shaded regions are tongue speed for high and low vigor licks
(left) and change in speed (right). **D**. Left figure shows SS rates for protractions that long or short deceleration
periods but the same peak speed. Right figure shows within cell differences. Shaded regions are tongue speed for
long and short deceleration period licks (left) and change in speed (right). **E**. Modulation of CS and SS rates during
task relevant and task irrelevant licks. Light shaded region is tongue speed for task relevant movements, while dark
shaded region is tongue speed for task irrelevant movements. Error bars are SEM.
550

551
552 **Supplementary Figure 1.** The number of task relevant (tube directed) and task irrelevant (grooming) licks recorded
553 per neuron.
554

555
556 **Supplementary Figure 2**. The number of neurons recorded from each marmoset. CS only refers to P-cells for which
557 only the complex spikes were recorded. CS&SS refers to P-cells for which both the CS and SS were recorded. SS
558 only refers to putative P-cells for which only the SS were recorded.
559

560
561
Supplementary Figure 3. Kinematic effects of CS-induced P-cell suppression in Cartesian coordinates. Left plot
shows the change in the position of the tip of the tongue at the end of protraction in movements that experienced
a CS during the deceleration period of protraction, with respect to movements that did not experience a CS during
any period. Right plot shows the change in position of the tip of the tongue as measured at peak retraction
velocity, comparing movements that experienced a CS during the retraction acceleration period with movements
that did not experience a CS in any period. Each line is the average effect for a single P-cell. P-cell suppression
during protraction produced hypermetria, whereas the suppression during retraction resulted in slowing.
569
570

571
Supplementary Figure 4. There was little or no change in lick kinematics when the CS-induced SS suppression
occurred before lick onset. We chose the period following completion of one lick and the start of the next lick,
which was on average 30ms in duration, i.e., the period in which the tongue was in the mouth. Top row: SS rates,
aligned to lick onset, protraction peak speed, and retraction onset. Second row: change in SS rates. Third row:
change in tongue displacement. Fourth row: change in tongue angle. The filled region is 95% confidence interval.
Error bars are SEM.
578

579
580 **Supplementary Figure 5**. Peak velocity as a function of lick amplitude during protraction and retraction for task
581 relevant and task irrelevant licks in each subject. Error bars are SEM.
582

1.   W. J. M. Levelt, *Speaking: From Intention to Articulation* (MIT Press, 1993).

2.   M. Gentil, EMG analysis of speech production of patients with Friedreich disease. *Clin. Linguist.*
Phon. **4**, 107–120 (1990).

3.   T. Vilis, J. Hore, Central neural mechanisms contributing to cerebellar tremor produced by limb
perturbations. *J.Neurophysiol.* **43**, 279–291 (1980).

4.   F. Yu, Q. J. Jiang, X. Y. Sun, R. W. Zhang, A new case of complete primary cerebellar agenesis:
clinical and imaging findings in a living patient. *Brain* **138**, e353 (2015).

5.   J. L. Bryant, J. D. Boughter, S. Gong, M. S. LeDoux, D. H. Heck, Cerebellar cortical output encodes
temporal aspects of rhythmic licking movements and is necessary for normal licking frequency.
Eur. J. Neurosci. **32**, 41–52 (2010).

6.   M. A. Gaffield, J. M. Christie, Movement Rate Is Encoded and Influenced by Widespread, Coherent
Activity of Cerebellar Molecular Layer Interneurons. *J.Neurosci.* **37**, 4751–4765 (2017).

7.   J. P. Bowman, L. D. Aldes, Organization of the cerebellar tongue representation in the monkey.
Exp. Brain Res. **39**, 249–259 (1980).

8.   L. D. Aldes, J. P. Bowman, Representation of the tongue in the cerebellar nuclei of the monkey.
Exp. Neurol. **64**, 202–215 (1979).

9.   M. A. Gaffield, B. A. Sauerbrei, J. M. Christie, Cerebellum encodes and influences the initiation,
performance, and termination of discontinuous movements in mice. *eLife* **11**, e71464 (2022).

10.   J. P. Welsh, E. J. Lang, I. Suglhara, R. Llinas, Dynamic organization of motor control within the
olivocerebellar system. *Nature* **374**, 453–457 (1995).

11.   K. Kubota, S. Hayama, Comparative anatomical and neurohistological observations on the tongues
of pigmy and common marmosets. *Anat. Rec.* **150**, 473–485 (1964).

12.   D. J. Herzfeld, Y. Kojima, R. Soetedjo, R. Shadmehr, Encoding of error and learning to correct that
error by the Purkinje cells of the cerebellum. *Nat.Neurosci.* **21**, 736–743 (2018).

13.   C. Ju, *et al.*, Neurons of the inferior olive respond to broad classes of sensory input while subject to
homeostatic control. *J. Physiol.* **597**, 2483–2514 (2019).

14.   S. Tsutsumi, *et al.*, Purkinje Cell Activity Determines the Timing of Sensory-Evoked Motor Initiation.
Cell Rep. **33**, 108537 (2020).

15.   J. S. Pi, *et al.*, The olivary input to the cerebellum dissociates sensory events from movement plans.
Proc. Natl. Acad. Sci. **121**, e2318849121 (2024).

16.   E. Sedaghat-Nejad, J. S. Pi, P. Hage, M. A. Fakharian, R. Shadmehr, Synchronous spiking of
cerebellar Purkinje cells during control of movements. *Proc. Natl. Acad. Sci.* **119**, e2118954119
(2022).

17. Y. Sato, A. Miura, H. Fushiki, T. Kawasaki, Short-term modulation of cerebellar Purkinje cell activity
after spontaneous climbing fiber input. *J. Neurophysiol.* **68**, 2051–2062 (1992).

18. W. T. Thach, Somatosensory receptive fields of single units in cat cerebellar cortex. *J.Neurophysiol.*
30, 675–696 (1967).

19. V. Romano, *et al.*, Olivocerebellar control of movement symmetry. *Curr. Biol.* **32**, 654-670.e4
(2022).

20. M. L. Streng, L. S. Popa, T. J. Ebner, Climbing Fibers Control Purkinje Cell Representations of
Behavior. *J.Neurosci.* **37**, 1997–2009 (2017).

21. S. Z. Muller, *et al.*, Complex spikes perturb movements and reveal the sensorimotor map of
Purkinje cells. *Curr. Biol.* **33**, 4869-4879.e3 (2023).

22. P. Hage, *et al.*, Effort cost of harvest affects decisions and movement vigor of marmosets during
foraging. *eLife* **12**, RP87238 (2023).

23. A. Mathis, *et al.*, DeepLabCut: markerless pose estimation of user-defined body parts with deep
learning. *Nat. Neurosci.* **21**, 1281–1289 (2018).

24. E. Sedaghat-Nejad, *et al.*, Behavioral training of marmosets and electrophysiological recording
from the cerebellum. *J.Neurophysiol.* **122**, 1502–1517 (2019).

25. E. Sedaghat-Nejad, *et al.*, P-sort: an open-source software for cerebellar neurophysiology. *J.
Neurophysiol.* **126**, 1055–1075 (2021).

26. T. R. Reppert, *et al.*, Movement vigor as a trait-like attribute of individuality. *J.Neurophysiol.* **120**,
741–757 (2018).

27. Y. Kojima, F. R. Robinson, R. Soetedjo, Cerebellar fastigial nucleus influence on ipsilateral abducens
activity during saccades. *J.Neurophysiol.* **111**, 1553–1563 (2014).

28. C. Bourrelly, J. Quinet, L. Goffart, The caudal fastigial nucleus and the steering of saccades toward
a moving visual target. *J. Neurophysiol.* **120**, 421–438 (2018).

29. M. I. Becker, A. L. Person, Cerebellar Control of Reach Kinematics for Endpoint Precision. *Neuron*
103, 335–348 (2019).

30. T. J. H. Ruigrok, J. Voogd, Organization of projections from the inferior olive to the cerebellar nuclei
in the rat. *J. Comp. Neurol.* **426**, 209–228 (2000).

31. R. Apps, M. Garwicz, Anatomical and physiological foundations of cerebellar information
processing. *Nat.Rev.Neurosci.* **6**, 297–311 (2005).

32. T. M. Hoogland, J. R. De Gruijl, L. Witter, C. B. Canto, C. I. De Zeeuw, Role of Synchronous
Activation of Cerebellar Purkinje Cell Ensembles in Multi-joint Movement Control. *Curr. Biol.* **25**,
1157–1165 (2015).

33.   S. Kitazawa, T. Kimura, P. B. Yin, Cerebellar complex spikes encode both destinations and errors in
arm movements. *Nature* **392**, 494 (1998).

34.   A. L. Hewitt, L. S. Popa, T. J. Ebner, Changes in Purkinje cell simple spike encoding of reach
kinematics during adaption to a mechanical perturbation. *J.Neurosci.* **35**, 1106–1124 (2015).

35.   W. Heffley, *et al.*, Coordinated cerebellar climbing fiber activity signals learned sensorimotor
predictions. *Nat. Neurosci.* **21**, 1431–1441 (2018).

36.   M. J. Wagner, *et al.*, A neural circuit state change underlying skilled movements. *Cell* **184**, 3731-
3747.e21 (2021).

37.   D. J. Calame, M. I. Becker, A. L. Person, Cerebellar associative learning underlies skilled reach
adaptation. *Nat. Neurosci.* 1–12 (2023). https://doi.org/10.1038/s41593-023-01347-y.

38.   T. Ishikawa, *et al.*, Releasing dentate nucleus cells from Purkinje cell inhibition generates output
from the cerebrocerebellum. *PLoS.One.* **9**, e108774 (2014).

39.   Y. Kojima, R. Soetedjo, Change in sensitivity to visual error in superior colliculus during saccade
adaptation. *Sci.Rep.* **7**, 9566 (2017).

40.   Y. Kojima, R. Soetedjo, Elimination of the error signal in the superior colliculus impairs saccade
motor learning. *Proc.Natl.Acad.Sci.U.S.A* **115**, E8987–E8995 (2018).

41.   P. J. May, S. Warren, Y. Kojima, The superior colliculus projection upon the macaque inferior olive.
Brain Struct. Funct. (2024). https://doi.org/10.1007/s00429-023-02743-7.

42.   R. Soetedjo, Y. Kojima, A. F. Fuchs, How cerebellar motor learning keeps saccades accurate. *J.*
Neurophysiol. **121**, 2153–2162 (2019).

43.   A. Thomas, *et al.*, Superior colliculus bidirectionally modulates choice activity in frontal cortex. *Nat.*
Commun. **14**, 7358 (2023).

44.   Z. Gao, *et al.*, A cortico-cerebellar loop for motor planning. *Nature* **563**, 113–116 (2018).

45.   F. P. Chabrol, A. Blot, T. D. Mrsic-Flogel, Cerebellar Contribution to Preparatory Activity in Motor
Neocortex. *Neuron* **103**, 506–519 (2019).

46.   J. Zhu, H. Hasanbegović, L. D. Liu, Z. Gao, N. Li, Activity map of a cortico-cerebellar loop underlying
motor planning. *Nat. Neurosci.* 1–13 (2023). https://doi.org/10.1038/s41593-023-01453-x.

47.   T. Tang, T. A. Blenkinsop, E. J. Lang, Complex spike synchrony dependent modulation of rat deep
cerebellar nuclear activity. *eLife* **8**, e40101 (2019).

48.   J. F. Medina, S. G. Lisberger, Links from complex spikes to local plasticity and motor learning in the
cerebellum of awake-behaving monkeys. *Nat.Neurosci.* **11**, 1185–1192 (2008).

49.   Y. Yang, S. G. Lisberger, Role of Plasticity at Different Sites across the Time Course of Cerebellar
Motor Learning. *J. Neurosci.* **34**, 7077–7090 (2014).

50.   J. Chaumont, *et al.*, Clusters of cerebellar Purkinje cells control their afferent climbing fiber
683   discharge. *Proc.Natl.Acad.Sci.U.S.A.* **110**, 16223–16228 (2013).

684   51.   R. Soetedjo, G. D. Horwitz, Closed-Loop Optogenetic Perturbation of Macaque Oculomotor
685   Cerebellum: Evidence for an Internal Saccade Model. *J. Neurosci.* **44** (2024).

686   52.   A. Mathis, *et al.*, DeepLabCut: markerless pose estimation of user-defined body parts with deep
687   learning. *Nat. Neurosci.* **21**, 1281–1289 (2018).

688   53.   J. H. Siegle, *et al.*, Open Ephys: an open-source, plugin-based platform for multichannel
689   electrophysiology. *JNeural Eng* **14**, 045003 (2017).

690   54.   M. Pachitariu, N. Steinmetz, S. Kadir, M. Carandini, H. K. D, Kilosort: realtime spike-sorting for
691   extracellular electrophysiology with hundreds of channels. *bioRxiv* 061481 (2016).
692   https://doi.org/10.1101/061481.

693
694