Title: Multi-species genome-wide CRISPR screens identify conserved suppressors of cold-

induced cell death

Authors

Breanna Lam[1,*], Kathrin M. Kajderowicz[1,2,*], Heather R. Keys[1], Julian M. Roessler[1], Evgeni M.

Frenkel[1], Adina Kirkland[1], Punam Bisht[1], Mohamed A. El-Brolosy[1,3], Rudolf Jaenisch[1], George

W. Bell[1], Jonathan S. Weissman[1], Eric C. Griffith[4], Sinisa Hrvatin[1,†]

Affiliations

[1]Whitehead Institute for Biomedical Research and Department of Biology, Massachusetts

Institute of Technology, Cambridge, MA 02142, USA

[2]Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology,

Cambridge, MA 02142, USA

[3]Harvard Society of Fellows, Cambridge, MA, 02138, USA

[4]Department of Neurobiology, Harvard Medical School, Boston, MA 02115, USA

*These two authors contributed equally to this work

Corresponding author

[†]Sinisa Hrvatin (shrvatin@wi.mit.edu)

Summary

Cells must adapt to environmental changes to maintain homeostasis. One of the most striking

environmental adaptations is entry into hibernation during which core body temperature can

decrease from 37°C to as low at 4°C. How mammalian cells, which evolved to optimally function

within a narrow range of temperatures, adapt to this profound decrease in temperature remains

poorly understood. In this study, we conducted the first genome-scale CRISPR-Cas9 screen in

cells derived from Syrian hamster, a facultative hibernator, as well as human cells to investigate

the genetic basis of cold tolerance in a hibernator and a non-hibernator in an unbiased manner.

Both screens independently revealed glutathione peroxidase 4 (GPX4), a selenium-containing

enzyme, and associated proteins as critical for cold tolerance. We utilized genetic and

pharmacological approaches to demonstrate that GPX4 is active in the cold and its catalytic

activity is required for cold tolerance. Furthermore, we show that the role of GPX4 as a

suppressor of cold-induced cell death extends across hibernating species, including 13-lined

ground squirrels and greater horseshoe bats, highlighting the evolutionary conservation of this

mechanism of cold tolerance. This study identifies GPX4 as a central modulator of mammalian

cold tolerance and advances our understanding of the evolved mechanisms by which cells

mitigate cold-associated damage—one of the most common challenges faced by cells and

organisms in nature.

Introduction

Rapid temperature changes pose a challenge to all clades of life, including endotherms.

Homeotherms such as mammals routinely defend a core body temperature set-point, with

profound thermal deviations leading to organ dysfunction and death[1–4]. Indeed, several studies

have shown that while homeotherm-derived cells typically possess a capacity for mild cold

tolerance in culture, primary cultured cells and cell lines derived from a number of mammalian

species, including humans, exhibit high rates of cell death following prolonged severe cold

exposure[5–8]. Although considerable work has focused on understanding the cellular responses

and adaptations to an increased temperature (heat stress and heat-shock)[9–11], little is known

about how cell biological processes modulate cell sensitivity to decreased temperatures.

The mechanisms underlying cold-induced death remain poorly understood but have been

hypothesized to involve phase transition of cellular membranes, with the resulting membrane

damage leading to ion gradient disruption and irrecoverable cellular dysfunction[12–15]. However,

many warm-blooded animals, including some primates, have evolved the ability to enter torpor

and hibernation – states during which body temperature can decrease far below its homeostatic

set-point[16–18]. In many animals, *torpor* is a state of profoundly reduced metabolic rate and body

temperature lasting from hours to days, while *hibernation* is a seasonal behavior comprising

multiple bouts of *torpor* interrupted by periodic arousals to normal body temperature. During

torpor, the core body temperature of many small hibernators including the Syrian golden hamster

(*Mesocricetus auratus*), the 13-lined ground squirrel (*Ictidomys tridecemlineatus*), and the greater

horseshoe bat (*Rhinolophus ferrumequinum*), reaches 4-10°C[19–21]. Animals can remain at these

low temperatures for extended periods of time (up to several weeks), indicating that their cells

and tissues are either genetically predisposed to cold tolerance, or that they can adapt to tolerate

long-term cold exposure[19–22].

Although in principle the ability to tolerate cold temperatures could be conveyed by systemic

factors present in torpid or hibernating animals, cell lines derived from hibernating rodents

maintain high viability when exposed to 4°C for several days, indicating the presence of cell-

intrinsic mechanisms of cold tolerance[5–8]. Several studies have reported distinctive responses of

hibernator-derived cells in the cold. For example, Syrian hamster-derived epithelial-like hamster

kidney (HaK) cells maintain mitochondrial membrane potential and ATP production during cold

exposure[7], and both HaK and epithelial-like smooth muscle cells derived from the Syrian golden

hamster are able to prevent excess reactive oxygen species damage in response to cold

exposure[5]. Similarly, unlike human induced pluripotent stem cell (iPSC)-derived neurons, cultures

of iPSC-derived neurons from 13-lined ground squirrels retain microtubule stability in the cold[6].

Despite metabolic and cell biological differences between various hibernator and non-hibernator-

derived cells in the context of cold exposure, the underlying genetic modulators of cold tolerance

have yet to be systematically explored. Additionally, it is still unknown to what extent the pathways

that regulate cold tolerance in one hibernating species are conserved across other evolutionarily

distant hibernators and potentially even present in non-hibernators.

Here, we carry out the first unbiased genome-scale CRISPR-Cas9 screens in cells derived from

a cold-tolerant hibernator (Syrian hamster) as well as human (non-hibernator) cells to identify

genetic pathways required for long-term cold tolerance. Surprisingly, these screens independently

identify a common mechanism dependent on the selenocysteine-containing enzyme Glutathione

Peroxidase 4 (GPX4) as a key mediator of cellular cold tolerance. Employing genetic and

pharmacological approaches, we confirm these findings and demonstrate that increased GPX4

activity is sufficient to improve cold tolerance in human cells. We further show that functional

GPX4 is required for cold tolerance across several hibernating and non-hibernating mammals,

including in distantly related hibernating horseshoe bats (*Rhinolophus ferrumequinum*), and

propose that the GPX4 pathway may be a widespread, evolutionarily ancient metazoan

mechanism of cold tolerance across endotherms.

Results

Assay for measuring viability of cold-exposed cells

Our initial evaluation of the ability of human K562 leukemia cells to tolerate prolonged exposure

to extreme cold (4°C) temperatures led us to establish a new method for assessing viability of

cold-exposed cells. To avoid confounding changes in media composition or pH, we transferred

cultured cells into incubators cooled to 4°C with the CO_2 concentration calibrated to maintain

physiological pH (7.4). Cell survival was assessed based on exclusion of trypan blue, a

membrane-impermeable dye, immediately upon removal from 4°C. We observed a 27 ± 3%

survival rate for K562 cells following 24 hours at 4°C using this approach. However, recounting
live cells after a 24-hour rewarming at 37°C revealed a ~6-fold increase in live cells, inconsistent
with the reported 16-24 hour doubling rate of this cell line (**Figure 1b, Figure S1a**). Reasoning
that a substantial fraction of trypan blue-positive cells at 4°C remain alive and capable of cell
division and recovery upon rewarming, we carried out a time course of trypan blue staining shortly
after removal from 4°C (**Figure 1a**). We found that the fraction of trypan blue-positive cells
dramatically reduced following a brief rewarming period (30 minutes to 1 hour, either at room
temperature or at 37°C), consistent with the number of live cells observed after 24 hours of
rewarming (**Figure 1b,c, Figure S1a,b**). To determine whether this effect was isolated to K562
cells, we measured trypan blue-positive cells at 4°C or following a brief period of rewarming in
several human cell lines (HEK 293T, HeLa, and RPE1 cells), as well as Syrian hamster kidney
fibroblasts (BHK-21 cells) (**Figure S1c-f**)**. We consistently observed that, absent a rewarming
period, trypan blue staining significantly overestimates cold-induced cell death. We thus adopted
this brief rewarming period (~30-minutes) as a standard assay to measure cell viability of cold-
exposed cells in our subsequent studies.

Hibernator-derived cells show enhanced cold tolerance compared to human cells
Using our modified assay for cell viability after cold challenge, we examined the relative cold
tolerance of hibernator and non-hibernator cells. We tested four commonly used human cell lines
(HT1080, HeLa, RPE1, and K562), as well as two cell lines (BHK-21 and HaK) derived from
Syrian hamsters (*Mesocricetus auratus*), a hibernating mammal. Cells were placed at 4°C - the
lower end of the temperature range which Syrian hamsters reach during hibernation - and we
assessed cell survival after 1, 4, and 7 days of cold exposure. Consistent with prior reports[5,7,8],
hamster-derived cell lines maintained high levels of viability, whereas human cell lines showed
varying degrees of death (**Figures 1d-f**), with several exhibiting less than 50% survival following
7 days in the cold. Notably, we observed generally higher cell survival rates than previously
reported[5], likely owing to our modified method of measuring cell viability. Similar survival trends
were also observed using an orthogonal assay based on lactate dehydrogenase (LDH) release
from dead cells into the media (**Figure 1g**).

Genome-scale CRISPR screen in a hibernator-derived cell line identifies suppressors of
cold-induced cell death
To investigate the molecular pathways regulating cold resistance in hibernator-derived BHK-21
cells, we designed a genome-scale CRISPR-Cas9 screen. Taking advantage of a recent

chromosome-level assembly of the *Mesocricetus auratus* genome[23] and a modified CRISPOR-

based guide selection algorithm (Methods), we generated a library of 218,143 single guide RNAs

(sgRNAs) targeting all 21,473 annotated genes (~10 guides per gene), including 2,299 intergenic-

targeting and 250 non-targeting sgRNAs as negative controls. The pooled library was introduced

into BHK-21 cells via a lentiviral vector expressing the sgRNA along with Cas9, achieving ~1000-

fold library representation with a multiplicity of infection (MOI) < 1.

Considering that many mammalian species, including Syrian hamsters, undergo prolonged

periods of cold exposure (deep torpor) and rewarming (interbout arousals) during seasonal

hibernation, we sought to recapitulate these temperature changes in our screen. We thus exposed

BHK-21 cells to three cycles consisting of 4°C cold exposure (4 days) followed by rewarming at

37°C (2 days) ("4°C Cycling", **Figure 2a**). To isolate genes selectively required upon cold

exposure, we also analyzed BHK-21 cultures transduced in parallel that were maintained at 37°C

for four passages/cycles ("37°C Control", **Figure 2a**). For both paradigms, genomic DNA was

isolated from cells during each cycle/passage to monitor selective sgRNA depletion. To assess

our screen performance, we tested whether sgRNAs targeting previously identified human core

essential genes[24] were significantly depleted compared with nontargeting and intergenic sgRNAs

and found that we can robustly distinguish between these functional categories at all three cycles

and at both 37°C and 4°C (estimated p-value < 2.2e-16) (**Figure S2**).

To identify genes that selectively modify (suppress or potentiate) cold-induced cell loss, we

compared the guides depleted during three cycles of cold exposure and rewarming to controls at

37°C with matched population doublings. Among the 21,473 targeted genes, only one gene,

ribosomal protein S29 (Rps29)[25], was selectively required in control 37°C conditions and only nine

genes appeared selectively required during cold exposure (FDR < 0.1, log2 fold-change [Log2FC]

>1, **Figure 2b, Table S1, S2**). Four selectively required genes - *Dld*, *Lias*, *Lipt1*, and *Ybey* – are

involved in lipoylation and mitochondrial RNA processing[26–29]. Strikingly, five of the nine required

genes - *Gpx4*, *Eefsec*, *Pstk*, *Secisbp2*, and *Sepsecs* - represent known components of the

glutathione/Glutathione Peroxidase 4 (GPX4) antioxidant pathway, indicating that this pathway

functions as a potent suppressor of cold-induced cell death in hibernator-derived BHK-21 cells

(**Figure 2c**).

Gpx4 encodes a selenocysteine-containing lipid antioxidant enzyme that acts to suppress lipid

peroxidation via the glutathione-mediated reduction of lipid hydroperoxides to non-toxic lipid

alcohols[30]. GPX4 inhibition sensitizes cells to ferroptosis, a distinct form of programmed cell death

characterized by iron-dependent accumulation of lipid peroxidation[31]. Notably, cold exposure has

previously been associated with increased lipid peroxidation and ferroptosis in human cells[5].

Underscoring the requirement for Gpx4 activity, four of the eight remaining genes selectively

required upon cold exposure (*Eefsec, Secisbp2, Pstk, Sepsecs*) are directly or indirectly involved

in selenocysteine incorporation into cellular proteins and thus required for Gpx4 activity (**Figures**

2d-h)[32,33].

Given that our initial screen design involved repeated rewarming cycles, we could not exclude the

possibility that the Gpx4 pathway confers resistance to rewarming-associated stress rather than

cold tolerance per se. To address this issue, we took advantage of the high cold tolerance of BHK-

21 cells and included a second screen arm using similar methods in which cells were continuously

exposed to 4°C for 15 days (**Figure 2i**). Sequencing prior to and following 15 days of cold

exposure, followed by a brief rewarming to 37°C, confirmed significant depletion (FDR < 0.1,

Log2FC < -0.5) of sgRNAs targeting *Gpx4*, *Eefsec*, and *Secisbp2* (as well as a trend toward

depletion for *Sepsecs* and *Pstk*), confirming a critical role for the Gpx4 pathway in BHK-21 cell

prolonged cold tolerance (**Figures 2j,k, Table S2, S3**). Thus, two unbiased genome-scale screen

arms identified Gpx4 as a major suppressor of cold-induced cell death in hibernator-derived BHK-

21 cells in the context of both cyclical and continuous cold exposure.

Gpx4 catalytic activity is essential for hibernator cell survival during cold exposure

To further validate the role of GPX4 in cold-mediated cell death, we employed the top-ranked

sgRNA from our screen to generate clonal *Gpx4* knockout BHK-21 cells. We sorted individual

cells to obtain clonal knockout cell lines and maintained them in the presence of liproxstatin-1

since they showed stunted growth at 37°C (**Figure S4a**). We confirmed the absence of Gpx4 via

immunoblotting (**Figure 3a**). While wild-type (WT) BHK-21 cells maintained high cell viability at

4°C, all three *Gpx4* knockout lines exhibited complete cell death within 4 days of cold exposure

(****P < 0.0001) (**Figure 3a)**. Moreover, these differences reflected Gpx4 loss, as opposed to

off-target effects insofar as lentiviral re-expression of cytosolic hamster Gpx4 (****P < 0.0001),

but not a GFP control gene or a catalytically dead mutant form of Gpx4 (mGPX4) that has a

serine in the active site instead of a selenocysteine, restored cold tolerance of *Gpx4* knockout

lines (**Figures 3b,c).**

Together, these genetic experiments strongly implicate the catalytic activity of Gpx4 in BHK-21

cold tolerance. However, the constitutive nature of these interventions precluded a determination

of whether Gpx4 functions to actively oppose cell death during cold-exposure or whether loss of

its catalytic activity prior to cold exposure sensitizes cells to subsequent cold challenge. To

address this issue, we used RSL3 and ML162, two competitive small-molecule Gpx4 inhibitors[34–

36]. Acute treatment of cultured BHK-21 cells with these inhibitors during cold exposure yielded

dose-dependent increases in cell death. Indeed, by four days of treatment, both RSL3- and

ML162-treated BHK-21 cells display significantly increased cold-induced death compared to

untreated controls (****P < 0.0001) (**Figures 3d,e**). Importantly, relative to WT cells, *Gpx4*

knockout cells exhibited no significant increase in cold-induced cell death upon RSL3 treatment

following a short cold exposure (8 hours) (**Figure S3**), confirming the specificity of these inhibitors

under the test conditions. We therefore conclude that Gpx4 catalytic activity is required in BHK-

21 cells during the course of cold exposure to actively oppose cold-induced cell death.

Given that Gpx4 activity acts as a major endogenous suppressor of ferroptosis[31], we tested

whether the cold-induced cell death observed under Gpx4 inhibition occurred via the ferroptosis

pathway. To this end, we made use of two small molecule inhibitors of ferroptosis, ferrostatin-1,

a lipophilic antioxidant, and deferoxamine mesylate (DFO), a Fe^{2+} chelator. Both inhibitors

effectively rescued cold-induced cell death in the presence of the Gpx4 inhibitor RSL3 (**Figure**

3f), indicating that acute inhibition of Gpx4 activity in the cold drives cell death via ferroptosis.

Together, through a combination of genetic and pharmacological approaches, our studies

demonstrate that Gpx4 activity confers substantial cold resistance in hibernator-derived cells,

protecting cells from cold-induced ferroptosis.

Genome-scale CRISPR screen identifies determinants of cold sensitivity in human cells

In contrast to hibernator-derived cell lines, human cells display varying cold sensitivities in

culture, with some exhibiting pronounced intolerance to the cold (**Figure 1d-g**). Consistent with

prior reports[5,8,37], pharmacological experiments in several human cell lines (K562, HT1080,

RPE, and HeLa cells) confirmed that cold-induced cell-loss occurred primarily via ferroptosis,

with both antioxidant ferroptosis inhibitors (ferrostatin-1 and liproxstatin-1) and iron chelators

(DFO and 2'2'-pyridine) increasing the cold survival in all four cell lines in a dose-dependent

manner. By contrast, neither the Caspase inhibitor Z-VAD-FMK nor the necroptosis inhibitor

necrostatin-1 significantly affected cell viability (**Figure S5, S6**).

To gain further insight into the genetic modifiers of human cell cold sensitivity and resistance,

we designed and performed an additional CRISPR-Cas9 screen in cold-sensitive human K562

cells exposed to cold temperatures. A pooled lentiviral CRISPR-Cas9 library targeting 19,381

genes (~5 sgRNAs per gene) was delivered to K562 cells at an MOI of ~0.5. Transduced cells

were placed at 4°C for five days to allow for cold-induced cell death to occur in approximately 79

$\pm$ 1.76% of cells, followed by a three-day rewarming to 37°C to allow for amplification of

remaining cells (**Figure 4a**). This cold-rewarming cycle was repeated three times, with cells

collected after each cycle to monitor the relative enrichment and depletion of individual sgRNAs.

As in our prior screens, separate cultures transduced in parallel were maintained at 37°C for

comparative analysis. In addition, to discriminate between ferroptosis-dependent and -

independent regulators of cold tolerance, we conducted a parallel screen in cold-exposed K562

cells under continuous ferrostatin-1 treatment. As before, the efficiency and specificity of the

screens were assessed based on effective depletion of characterized core essential genes[24] as

well as the lack of depletion of negative control guides (250 nontargeting and 125 intergenic

sgRNAs) (**Figure S7**).

To our surprise, this screen did not identify any genes whose disruption significantly enhanced

K562 cold tolerance. By contrast, we identified 176 genes selectively required during cold

cycling relative to constant 37°C control conditions (FDR < 0.1, LFC < -0.1, and requiring >2

sgRNAs, **Figure 4b**). It is notable that K562 cold survival was sensitive to the targeting of larger

number of genes than the BHK-21 line, potentially indicating increased dependencies on

genetic pathways which are not essential in BHK-21 cells (**Figure S8, Table S6**). Among the

pathways identified, we honed in on the selenocysteine incorporation pathway and observed

GPX4 and the selenocysteine incorporation genes *PSTK*, *SEPSECS*, and *EEFSEC* within the

top-ranked K562 cold-protective genes, suggesting that the GPX4 pathway confers significant

resistance to the cold not only in cold-tolerant hibernator cells, but also in cold-sensitive human

cells (**Figures 4c-g**). Notably, we identified 11 genes that were no longer essential (Fold

enrichment > 0.50, FDR < 0.10) in the presence of Ferrostatin-1, indicating that these genes act

to suppress cold-induced ferroptosis (**Figures 4h-n, Table S4, S5**). Although multiple pathways

have been described as suppressors of ferroptosis in other contexts[38–42], it is notable that 5 of

11 identified genes represent known components of the GPX4 pathway.

To confirm these findings, we generated three clonal *GPX4* knockout K562 cell lines and

confirmed the loss of GPX4 via immunoblotting (**Figure 5a**). Similar to our findings in BHK-21

cells, K562 *GPX4* knockout clones exhibited stunted growth and were expanded in the

presence of liproxstatin-1 (**Figure S4b**). Importantly, GPX4 loss greatly increased cold-induced

cell death (****$P < 0.0001$) (**Figures 5a**), thus confirming an essential role for *GPX4* in human

cell cold tolerance. Similar effects were also observed upon acute pharmacologically inhibition

of GPX4 with RSL3 or ML162 at 4°C (****$P < 0.0001$) (**Figures 5b, c**). These effects were

dependent on GPX4 inhibition insofar as the cold tolerance of K562 *GPX4* knockout cells was

unaffected by RSL3 treatment (**Figure S9**). In addition, RSL3-driven cold-induced cell death

was rescued by either ferrostatin-1 or DFO treatment, indicating that RSL3-induced cell loss

occurs via the ferroptosis pathway (**Figure 5d**). Increased cold-induced death in the *GPX4*

knockout K562 cell lines could be rescued upon lentiviral expression of either the cytosolic

human (hs) or hamster (ma) GPX4, but not GFP or catalytically dead GPX4 variants (**Figure**

5e). Together, our genetic and pharmacological studies reveal that K562 cells, despite being

significantly less cold-tolerant than hamster BHK-21 cells, also rely on the GPX4 pathway to

suppress cold-induced ferroptosis.

GPX4 catalytic activity is limiting for K562 cell cold tolerance

Given that the GPX4 pathway is active during cold exposure and essential for cold survival in

both hamster BHK-21 and human K562 cells, it is somewhat surprising that these cell lines

exhibit markedly different cold tolerances. In this regard, it has previously been suggested[5] that

upon cold exposure human cells, in contrast to hibernator-derived cells, rapidly deplete

intracellular glutathione, an essential co-factor for GPX4 function, which could thus account for

their increased cold sensitivity. To test this idea, we supplemented K562 growth media with

either cell-permeable Glutathione reduced ethyl ester (GSH-MEE), N-acetyl-cysteine (NAC), or

L-Cystine, a glutathione precursor; however, these interventions failed to significantly increase

K562 cells' cold tolerance under our culture conditions (**Figure S10**). We therefore examined

the possibility that GPX4 protein levels in K562 cells are limiting for their cold tolerance. To this

end, we generated K562 lines overexpressing either human or hamster GPX4. Both of these

overexpression lines exhibited strikingly improved cold tolerance compared to the wild-type

K562 parental line and a GFP-overexpressing control cell line (****$P < 0.0001$) (**Figure 5f**.

Indeed, GPX4-overexpressing K562 cells displayed comparable cold tolerance to that of

hibernator-derived BHK-21 cells (**Figure 1d, g**). To confirm that increased GPX4 catalytic

activity was responsible for the improved cell cold tolerance, we generated additional K562 lines

overexpressing catalytically dead forms of hamster or human GPX4 in which the active site

selenocysteine was mutated to serine and found that these GPX4 mutant lines exhibited no

increase in cold tolerance (**Figure 5f**). Taken together, these results strongly suggest that GPX4

abundance serve as a key limiting determinant of K562 cell cold tolerance. Moreover, our

finding that human and hamster GPX4 overexpression confer comparable increases in K562

cold tolerance may indicate that the intrinsic activity of the hamster and human GPX4 enzymes

are similar.

GPX4 activity is broadly required for cold tolerance in primary cells across evolutionarily

distant mammalian species

Given that our studies had largely employed a small number of transformed cell lines, we

sought to extend our investigation to examine cold tolerance in primary cell types as well as

cells derived from additional hibernating species. To this end, we obtained primary and

immortalized fibroblasts from human, mouse, and rat as well as from Syrian hamster, 13-lined

ground squirrel (*Ictidomys tridecemlineatus*), and horseshoe bat (*Rhinolophus ferrumequinum*),

three distantly related mammalian hibernators, and characterized their cold tolerance. While

hibernator-derived fibroblasts exhibited robust, roughly uniform cold tolerance, we observed a

surprisingly variable degree of cold tolerance across non-hibernator primary cells (**Figure 6a**),

with human dermal fibroblasts exhibiting a remarkable 79 ± 4.93% cell viability after seven days

at 4ºC, comparable to that observed with hamster-derived BHK-21 cells.

To examine the role of GPX4 in primary cell cold resistance, we transduced cells with Cas9 and

sgRNAs targeting human *GPX4* to create a population of *GPX4* knockout human kidney

fibroblasts (**Figure 6b**). Consistent with our prior findings in cell lines, GPX4 loss resulted in

significantly decreased viability in the cold, which was largely rescued by ferrostatin-1 treatment

in fibroblast cells (**Figure 6b**). Indeed, RSL3 treatment significantly decreased the viability of

wild-type cold-exposed human kidney fibroblasts **(Figures 6c, d)**, and these effects were

dependent on GPX4 inhibition as the cold tolerance of human kidney fibroblast *GPX4* knockout

cells was largely unaffected by RSL3 treatment (**Figure S4c**).

To more widely examine the dependence of primary cell cold tolerance on GPX4 activity, we

obtained and tested fibroblasts derived from six mammalian species. We found the GPX4

activity widely contributes to cold tolerance across hibernators and non-hibernators was

dependent on GPX4, as RSL3-treated cells showed increased cold-induced cell death that was

largely rescued by ferrostatin-1 treatment (**Figures 6e, f**). We also extended these findings to a

non-fibroblast cell type, obtaining similar results with primary cortical neuronal cultures prepared

from neonatal mice and hamsters (**Figures 6g, h**). Taken together, our data indicate that

primary cell types are strongly sensitive to GPX4 loss in the cold and suggest that their

differential cold sensitivities may reflect different levels of GPX4-mediated cold tolerance.

Discussion

Hibernators can lower their body temperature to ~4°C for several days to weeks, indicating that

their cells and tissues possess the ability to survive extended periods of cold. Previous studies

have indicated that ferroptosis contributes to cold-induced cell death in cultured human

cells[5,8,37]; however, the genetic modifiers of cold sensitivity in hibernators and non-hibernators

have yet to be systematically explored. Here, we conducted a series of unbiased genome-wide

CRISPR-Cas9 screens in both hibernator- and non-hibernator-derived cells to investigate the

mechanisms controlling cellular cold tolerance. Our findings, further validated using stable

genetic knockout lines and pharmacological inhibitors, identify the GPX4 pathway as a critical

suppressor of cold-induced cell death. Indeed, overexpression studies in human K562 cells

suggest that GPX4 abundance can serve as a key limiting determinant of cellular cold tolerance.

Notably, Sone et al.[43] recently also identified GPX4 as a strong regulator of cold tolerance

through overexpression screening. These independent findings further support the current study

identifying GPX4 as a conserved suppressor of cold-induced cell death. The consistency of our

observations across diverse cell lines and primary cells, including cells derived from six different

mammalian species, argues that GPX4 serves as an essential and evolutionarily conserved

mechanism to protect cells from cold-induced ferroptotic cell death.

During the course of our studies, we made the surprising observation that cells stained with the

commonly employed membrane-impermeable dye trypan blue exhibit increased dye retention

immediately after cold exposure, an artifact not reflecting genuine cell death. Rather, cell

counting following brief rewarming yields more accurate results, in some cases reflecting

significantly higher cell viability than previously appreciated. Employing this modified approach

to measure cold-induced cell death across sixteen cell lines and primary cell types points to

more nuanced variations in cold survival across cell types. These differences are highlighted by

the pronounced contrast between the poor cold viability of human HT1080 cells and the robust

cold tolerance observed in human fibroblasts. The underlying basis of this transient cell

permeability for trypan blue following cold exposure remains unclear; however, it is noteworthy

that in *S. cerevisiae* heat and chemical insults have been found to induce a brief window of

membrane permeability to PI prior to membrane repair[44]. Likewise, uptake of propidium ions

across intact cell membranes has been previously observed in bacteria showing high

membrane potentials[45]. Our observations suggest that further investigation into the mechanisms

underlying transient membrane permeability in different cell types and stress conditions could

lead to improved methods for accurately measuring cell viability, ultimately enhancing our

understanding of cellular stress.

We report, to our knowledge, the first genome-scale hibernator CRISPR-Cas9 screen in cells

derived from Syrian hamster. To create a genome-scale CRISPR-Cas9 library, we implemented

a CRISPOR-based computational pipeline for sgRNA selection and benchmarked its success

by carrying out the hamster and human screens described here. Our validated sgRNA libraries

and associated algorithms should serve as a valuable resource to the hibernation community,

as well as other scientific communities developing CRISPR-based tools for non-model

organisms.

While our data point to GPX4 levels as a major determinant of cellular cold tolerance, whether

endogenous levels of functional Gpx4 across various cell types correlates with their cold

tolerance remains to be tested. It also remains possible that other factors such as GPX4

subcellular localization, the abundance of the cellular selenocysteine incorporation machinery,

the abundance of glutathione, as well as GPX4-pathway independent factors all play important

roles in limiting cold tolerance across various cell types. Although we did not see an increase in

cold cell viability upon treatment of K562 cells with GSH-MEE, it is notable that in our screens,

sgRNAs targeting genes involved in the glutathione biosynthesis (e.g. *GCLC*, *GSS*[46,47]) showed

increased depletion in cold-exposed K562 versus BHK-21 cells (**Figure S10, Figure S11**). This

suggest a reduced dependency on the conventional glutathione biosynthetic pathway in cold-

exposed BHK-21 cells. In addition, while our data points to a key cellular need to reduce cold-

associated lipid hydroperoxides, the mechanisms that drive peroxide accumulation in the cold

remain unclear. Specifically, the roles of reactive oxygen species (ROS) production,

polyunsaturated fatty acid (PUFA) levels, and free iron concentration should be further explored

to understand the specific drivers of cold-induced ferroptotic cell death. Such insights have the

potential to inform the development of more effective strategies for managing conditions

exacerbated by cold exposure, such as ischemic injuries and organ transplantation.

We limited our current investigations to cell culture paradigms. However, future *in vivo* studies in

hibernating organisms that naturally undergo profound drops in core body temperature could

yield further insight into the adaptive mechanisms that govern cellular survival during periods of

hibernation and torpor, while also allowing for the study of cold tolerance mechanisms in cell

types that cannot presently be maintained in culture. Similarly, a *Gpx4* conditional knockout

mouse line has been generated and used to show that *Gpx4* is essential for mitochondrial

integrity and neuronal survival in adult animals[48]; however, these animals have not yet been

used to evaluate the contribution of *Gpx4* to *in vivo* cellular and tissue cold tolerance.

The ability to hibernate and survive long-term cold exposure is present in many mammalian

species from rodents to bats to primates, raising the possibility that the ability to enter

hibernation and tolerate cold was an ancestral trait present in protoendotherms. Our observation

that a single pathway centered around GPX4 is required across diverse mammalian species to

protect cells from cold-induced cell death, raises a question whether GPX4 has a role in

protection from cold-induced ferroptosis in other cold-tolerant vertebrates including fish,

amphibians, reptiles, and birds that also enter torpor. Such studies will advance our

understanding of the evolved mechanisms by which cells mitigate cold-associated damage—

one of the most common challenges faced by cells and organisms in nature.

Acknowledgments

We thank Wei Li for generously sharing 13-lined ground squirrel (*Ictidomys tridecemlineatus*)

fibroblasts and Thomas P. Zwaka for sharing greater horseshoe bat (*Rhinolophus*

ferrumequinum) fibroblasts. We thank Amanda Chilaka and Sumeet Gupta of the Whitehead

Institute Genome Technology Core for high-throughput sequencing. We thank Aurora Lavin-

Peter, Aleksandar Markovski, Tara Thakurta, William S. Owens, Sheamin Khyeam, Karina

Smolyar, and Elena Assad for their technical support. We thank Whitney Henry, Matthew

Vander Heiden, Alison Ringel, Morgan Sheng, Myriam Heiman, and members of the Hrvatin

laboratory for providing feedback on the study. Schematics created with BioRender.com. This

material is based upon work supported by the National Science Foundation Graduate Research

Fellowship under Grant No. 2141064 and The Paul and Daisy Soros Fellowship for New

Americans (PSM). Research reported in this publication was supported by The G. Harold and

Leila Y. Mathers Charitable Foundation and the NIDDK/NIH under Award Number

DP2DK136123. The content is solely the responsibility of the authors and does not necessarily

represent the official views of the National Institutes of Health.

Author contributions

B.L., K.M.K., and S.H. conceived and designed the study. B.L. and K.M.K. designed, performed,

and analyzed experiments. B.L., K.M.K., H.R.K. designed and performed genome wide CRISPR

screens. A.K. assisted with the genome wide CRISPR screens. B.L. performed the genetic and

pharmacological studies to validate the screens. K.M.K. performed the experiments with primary

cells. E.M.F. and G.W.B. designed algorithms to select genome-wide Syrian hamster

(*Mesocricetus auratus*) CRISPR library. M.E.B. and P.B. immortalized mouse embryonic

fibroblasts and greater horseshoe bat (*Rhinolophus ferrumequinum*) fibroblasts. J.M.R., R.J.,

J.S.W., and E.C.G. advised on the study. B.L., S.H., and E.C.G led the writing of the manuscript

with contributions from other authors. E.C.G. and S.H. obtained funding for the research. All

authors approved and reviewed the manuscript.

Declaration of interests

The authors declare no competing interests.

Figures

Figure 1. Hibernator-derived cells exhibit increased cold resistance

a, Schematic of cold cell viability counting, consisting of 1 day cold exposure (4°C), followed by

a short rewarming at 37°C for 15, 30, or 60 minutes before trypan blue staining. **b,** Number of

viable K562 cells based on trypan blue staining after one day at 4°C and subsequent rewarming

for 24 hours at 37°C. Numbers are normalized to initial cell counts. Blue shaded regions indicate

4°C exposure and shaded pink regions indicate 37°C exposure. Red square indicates

calculated cell counts after one day at 4°C based on the viable cell number measured after 24

hour rewarming. **c**, Viability of K562 cells was assessed by trypan blue staining after incubation

at 37°C for 0, 15, 30, or 60 minutes following 24 hours at 4°C ($n = 4$). Cells incubated at 37°C

for 15, 30, and 60 minutes show a significant increase in cell counts compared to cells counted

without rewarming (n = 4 samples per condition, ***$P = 0.0007$, ****$P < 0.0001$). **d**, Viability of

hibernator (BHK-21, HaK)- and non-hibernator (HeLa, RPE1, HT1080, K562)-derived cell lines

at 4°C as measured by trypan blue staining. Hibernator-derived lines show significantly

increased cold resistance compared to lines derived from non-hibernators at 7 days 4°C ($n = 4$

samples per data point, ****P < 0.0001). **e, f**, Fluorescence images of hibernator- and non-

hibernator-derived cell lines after 4 days at 4°C. Cultures were stained with 1 µg/mL Hoechst

33342 and 1 µg/mL propidium iodide (PI) to distinguish live and dead cells. **g**, Viability of

hibernator- and non-hibernator-derived cell lines at 4°C as measured by LDH release (n = 4

samples per data point, ****P < 0.0001). All values show mean ± SEM, with significance

measured by one-way ANOVA adjusted for multiple comparisons by Dunnett's test. *P < 0.05;

**P < 0.01; ***P < 0.001; ****P < 0.0001; ns P > 0.05.

Figure 2. Unbiased CRISPR screens identify the GPX4 pathway as necessary for cold-induced survival in hibernator-derived BHK-21 cells

a, Schematic of CRISPR screen paradigm, consisting of four passages of control cells kept at 37°C and three cycles of cold exposure (4°C) for 4 days followed by rewarming (37°C) for 2 days. Yellow dots indicate points of sample collection. **b**, Volcano plot showing median log2 fold-change of genes after three cycles of cold exposure and rewarming (4°C) compared to cells following three passages at 37°C. Red dots indicate selectively required genes with a median log2 fold-change < -1 or > 1 and FDR < 0.1. **c**, Heatmap of the median log2FC of ferroptosis-

related genes after three cycles of cold exposure and rewarming (4°C) compared to three

passages at 37°C. **d**, Partial schematic of the selenocysteine incorporation pathway. Left:

Production of the GPX4 selenoprotein requires recoding of a UGA codon to the amino acid

selenocysteine (Sec). This process involves a cis-acting SECIS element within the *Gpx4* mRNA

3' UTR, SECIS binding protein 2 (SECISBP2), a specific eukaryotic elongation factor

(EEFSEC), and Sec-charged tRNA. Right: The Sec-charged tRNA is generated by the

combined action of Phosphoseryl-tRNA kinase (PSTK), Selenophosphate synthetase 2

(SEPHS2), and selenocysteine synthase (SEPSECS). **e-h**, Median log2 fold-change (log2FC) of

10 guides per targeted gene, showing guide depletion over three cycles of cold exposure and

rewarming. Significance between Cycle 1 versus Cycle 3 is measured by two-way ANOVA

adjusted for multiple comparisons by Dunnett's test. Significance between 37°C and 4°C for

each cycle is measured by two-way ANOVA adjusted for multiple comparisons by Bonferroni's

test. **e**, *Gpx4*, **f**, *Eefsec*, **g**, *Secisbp2,* **h**, *Pstk*. **i**, Schematic of CRISPR screen paradigm,

showing cells exposed to 4°C continuously for 15 days. Yellow dot indicates point of sample

collection. **j**, Volcano plot showing median log2 fold-change in abundance of guides targeting

the indicated genes after 15 days of 4°C exposure compared to one passage at 37°C. Red dots

indicate selectively required genes with a median log2 fold-change < -0.5 or > 0.5 and FDR <

0.10. **k**, Heatmap of the median log2FC in abundance of guides targeting ferroptosis-related

genes after 15 days of 4°C exposure compared to 37°C control cultures. *$P < 0.05$; **$P < 0.01$;

***$P < 0.001$; ****$P < 0.0001$; ns $P > 0.05$.

Figure 3. GPX4 activity is required for BHK-21 cell cold tolerance

a, Stable *Gpx4* knockout (KO) BHK-21 cell lines exhibit reduced cold tolerance. Left: Western

blot of wild-type (WT) and individual *Gpx4* KO clone lysates for GPX4 and β-actin loading

control. Right: Viability of *Gpx4* KO lines is significantly lower than WT BHK-21 cells at 7 days

4°C by trypan blue staining ($n = 4$, ****$P < 0.0001$), with complete cell death by four days at 4°C.

b, c, Reintroduction of wild-type Syrian hamster GPX4 (GPX4), but not a catalytically dead form

of GPX4 (mGPX4) rescues cold-induced cell death in two independent BHK-21 *Gpx4* KO clonal

cell lines. Left panels: Western blots for HA and GPX4 along with β-actin loading control. Right

panels: Expression of WT hamster GPX4 showed significantly higher cell viability at 7 days 4°C

compared to the corresponding parental *Gpx4* KO, GFP-, and mGPX4-expressing lines by

trypan blue staining ($n = 4$, ****$P < 0.0001$). **d, e**, Treatment with the GPX4 inhibitors RSL3 (**d**)

or ML162 (**e**) results in enhanced cold-induced death in BHK-21 cells by 4 days at 4°C as

measured by trypan blue exclusion ($n = 4$, ****$P < 0.0001$). **f**, Cold-induced BHK-21 cell death

upon RSL3 treatment occurs via ferroptosis. BHK-21 cells were placed at 4°C and treated with

RSL3 (1 µM) and/or the ferroptosis inhibitor ferrostatin-1 (Fer-1, 1 µM) or iron chelator DFO

(100 µM) for 4 days ($n = 4$). Treatment with RSL3 resulted in significantly lower cell viability than

no treatment as determined by one-way ANOVA adjusted for multiple comparisons by Tukey's

HSD (****$P < 0.0001$). All values show mean ± SEM, with significance measured by two-tailed t

test, unless otherwise indicated. *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.0001$; ns $P >$

0.05.

Figure 4. Genome-wide CRISPR screens identify GPX4 as a suppressor of cold-induced cell death in human cells

a, Schematic of CRISPR screen paradigm, consisting of three cycles of 5 days of cold exposure (4°C) interrupted by 3 day rewarming (37°C) periods. Yellow dots indicate points of sample collection. **b**, Volcano plot showing median log2 fold-change in abundance of guides targeting the indicated genes after three cycles of 4°C cold exposure compared to three passages at 37°C (Control). Red dots indicate selectively required genes with a median log2 fold-change < -

0.5 or > 0.5 and FDR < 0.1. **c-g**, Combined median log2 fold-change (log2FC) of 5 sgRNAs per

targeted gene, showing sgRNA depletion over three cycles of cold exposure. Significance

between Cycle 1 versus Cycle 3 is measured by two-way ANOVA adjusted for multiple

comparisons by Dunnett's test. Significance between 37°C and 4°C for each cycle is measured

by two-way ANOVA adjusted for multiple comparisons by Bonferroni's test. **c**, *GPX4*, **d**,

EEFSEC, **e**, *SECISBP2*, **f**, *PSTK*, **g**, *SEPSECS*. **h**, Volcano plot showing median log2 fold-

change in abundance of guides targeting the indicated genes after three cycles of 4°C cold

exposure in the presence or absence of 1 μM of ferrostatin-1. Red dots indicates selectively

required genes with a log2 fold-change < -0.5 or > 0.5 and FDR < 0.1. **i**, Heatmap of the median

log2FC in abundance of guides targeting ferroptosis-related genes after three cycles of cold

exposure (4°C) compared to 37°C control cultures with and without ferrostatin-1. **j-n**, Depletion

of 5 sgRNAs per gene over three cycles of cold exposure and rewarming with and without

ferrostatin-1 (1 uM) treatment. **j**, *GPX4* (***P* = 0.0047), **k**, *EEFSEC* (***P* = 0.0059), **l**, *SECISBP2*

(ns, *P* = 0.5457), **m**, *PSTK* (**P* = 0.0393), **n**, *SEPSECS* (***P* = 0.0029) as measured by two-

tailed t-test at Cycle 3. All values show mean ± SEM. **P* < 0.05; ***P* < 0.01; ****P* < 0.001; *****P*

< 0.0001; ns *P* > 0.05.

Figure 5. Endogenous GPX4 activity is limiting for K562 cell cold tolerance

a, Stable *GPX4* knockout (KO) K562 cell lines exhibit reduced cold tolerance. Left: Western

blotting of wild-type (WT) and individual *GPX4* KO clones for GPX4 and β-actin loading control.

Right: Viability of *GPX4* KO lines is significantly lower than WT K562 cells as measured by

trypan blue staining (n = 4, ****P < 0.0001), with complete cell death by one day at 4°C.

Significance measured by two-tail t-test at 7 days 4°C. **b, c**, Treatment with the GPX4 inhibitors

RSL3 (**b**) or ML162 (**c**) results in enhanced K562 cold-induced death by 4 days at 4°C (n = 4,

****P < 0.0001) as measured by two-tailed t-test. **d**, Cold- and RSL3-induced K562 cell death

occurs via ferroptosis. K562 cells were placed at 4°C and treated with RSL3 (1 µM) and/or the

ferroptosis inhibitor ferrostatin-1 (Fer-1, 1 µM) or iron chelator DFO (5 µM) for 4 days (n = 4).

Treatment with RSL3 resulted in significantly lower cell viability than no treatment as determined

by one-way ANOVA adjusted for multiple comparisons by Tukey's HSD (***P = 0.0002). **e**,

Reintroduction of wild-type human (hs) or hamster (ma) GPX4, but not catalytically dead forms

of GPX4 (Gpx4 U46S), rescues cold-induced cell death in a K562 *GPX4* KO clonal cell line (n =

4). Left: Western blot for GPX4 levels and β-actin loading control. Right: Expression of WT

human GPX4 and hamster GPX4 resulted in significantly higher cell viability compared to the

corresponding parental *GPX4* KO, GFP-, and mutGPX4-expressing lines by trypan blue

staining, as measured by one-way ANOVA adjusted for multiple comparisons by Dunnett's test

at 7 days 4°C (****P < 0.0001). **f**, Overexpression of wild-type human or hamster GPX4, but not

catalytically dead forms of GPX4 (mutGPX4), suppresses cold-induced cell death in a K562

cells (n = 4). Left: Western blot for GPX4, with β-actin loading control. Right: Expression of WT

human GPX4 and hamster GPX4 resulted in significantly higher cell viability compared to wild-

type, GFP-, and mutGPX4-expressing K562 lines by trypan blue staining, as measured by one-

way ANOVA adjusted for multiple comparisons by Dunnett's test at 7 days 4°C (****P < 0.0001).

All values show mean ± SEM. *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; ns P > 0.05.

Figure 6. GPX4 is required for cold tolerance across both several hibernator and non-

hibernator mammalian species and cell types

a, Viability of hibernator cells (Syrian hamster embryonic fibroblasts, Greater horseshoe bat

embryonic fibroblasts, and 13-lined ground squirrel embryonic fibroblasts) and non-hibernator

cells (SV40-immortalized mouse embryonic fibroblasts, human adult kidney fibroblasts, human

adult dermal fibroblasts, rat adult dermal fibroblasts, and mouse adult dermal fibroblasts)

exposed to 4°C as measured by trypan blue staining (n = 4, ****P < 0.0001). **b**, Human kidney

fibroblast *GPX4* knockout cells exhibit reduced cold tolerance compared to WT cells. Left:

Western blot of wild-type (WT) and *GPX4* KO cells for GPX4 and β-actin loading control. Right:

Viability of *GPX4* KO cells is significantly lower than WT cells at 4°C as measured by trypan

blue staining (n = 4, ****P < 0.0001). **c**, Gpx4 activity is essential for cold survival of primary

human kidney fibroblasts. Human kidney fibroblasts were placed at 4°C and left untreated or

treated with RSL3 (1 µM) and/or the ferroptosis inhibitor ferrostatin-1 (Fer-1, 1 µM) for 7 days (n

= 4, ****P < 0.0001). **d**, Representative fluorescence images of human kidney fibroblasts after 4

days at 4°C with no treatment, 1 µM Fer-1, 1 µM RSL3, or 1 µM Fer-1 and 1 µM RSL3. Cultures

were stained with Hoechst 33342 and propidium iodide (PI) to identify live cells. **e**, Gpx4 activity

is essential for cold survival in hibernator cells (Syrian hamster embryonic fibroblasts, Greater

horseshoe bat embryonic fibroblasts, and 13-lined ground squirrel embryonic fibroblasts) and

non-hibernator cells (SV40-immortalized mouse embryonic fibroblasts, human adult kidney

fibroblasts, human adult dermal fibroblasts, rat adult dermal fibroblasts, and mouse adult dermal

fibroblasts). Cells were placed at 4°C and left untreated, treated with RSL3 (1 µM), and/or

ferrostatin-1 (Fer-1, 1 µM) for 7 days (n = 4). **f**, Expanded Day 4 data from **e)** indicates that

Gpx4 activity is essential for fibroblast survival in the cold across several hibernator and non-

hibernator species. Cells were placed at 4°C and left untreated or treated with RSL3 (1 µM)

and/or ferrostatin-1 (Fer-1, 1 µM) for 7 days (n = 4, ****P < 0.0001). **g-h**, Gpx4 activity is

essential in **g)** mouse primary cortical neuron cultures and **h)** hamster primary cortical neuron

cultures. Cells were placed at 4°C and left untreated, treated with RSL3 (1 µM) and/or the

ferroptosis inhibitor ferrostatin-1 (Fer-1, 1 µM) for 1 or 4 days (n = 4). RSL3 treatment increased

cell death, which was rescued by ferrostatin-1. All values show mean ± SEM, with significance

measured by one-way ANOVA adjusted for multiple comparisons with Tukey's HSD. *P < 0.05;

**P < 0.01; ***P < 0.001; ****P < 0.0001; ns P > 0.05.

Tables

Table S1. Significant genes from Cycle 3 of genome-wide BHK-21 screen (median log2 fold-change < -1 or > 1 and FDR < 0.1)

Gene	Median Log2FC	FDR
Rps29	1.3794	0.017327
Dld	-1.7915	0.003094
Eefsec	-2.1607	0.000707
Gpx4	-2.2215	0.000707
Lias	-1.1881	0.011251
Lipt1	-1.1444	0.05562
Pstk	-2.3621	0.000707
Secisbp2	-1.1041	0.000707
Sepsecs	-2.3116	0.000707
Ybey	-1.5789	0.017492

Table S3. Significant genes from 15 days 4°C of genome-wide BHK-21 screen (median log2 fold-change < -0.5 or > 0.5 and FDR < 0.1)

Gene	Median Log2FC	FDR
LOC101827392	0.5431	0.066832
Eefsec	-0.81913	0.00165
Gpx4	-1.5133	0.00165
Secisbp2	-0.74996	0.00165

Table S4. Significant genes from Cycle 3, 4°C + ferrostatin-1 vs 4°C of genome-wide K562 screen (median log2 fold-change < -0.5 or > 0.5 and FDR < 0.1)

Gene	Median Log2FC	FDR
UBA3	0.59092	0.035754
PSTK	0.85375	0.000707
GPX4	1.3025	0.000707
CMIP	0.83519	0.09946
OXSM	0.82371	0.003094
HSCB	1.0999	0.060891
FTH1	1.4415	0.000707
SEPHS2	0.98322	0.000707
EEFSEC	0.91866	0.000707
FDXR	1.2588	0.000707
SEPSECS	0.9745	0.000707

Methods

Cell culture

Two hibernator-derived (BHK-21, HaK) and four non-hibernator-derived (HT1080, RPE1, HeLa, K562) cell lines were used. All cells were purchased from ATCC. K562 cells were cultured in Roswell Park Memorial Institute (RPMI) 1640 Medium (Gibco #11875093) supplemented with 10% fetal bovine serum (GeminiBio #100-106) and 1% penicillin/streptomycin. The remaining cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (Thermo Fisher Scientific #12430054) supplemented with 10% fetal bovine serum (GeminiBio #100-106) and 1% penicillin/streptomycin (Thermo Fisher Scientific #15140122).

Cell viability assay

Cells were seeded on 24-well culture plates at 50,000 cells per well and allowed to adhere for 24 hours at 37°C. Cells were then placed at 4°C with 5% CO_2 for varying time periods (1, 4, or 7 days), with a 30-minute rewarming at 37°C before assessing cell viability via trypan blue (TB) (Thermo Fisher Scientific #15250061) staining. To assess cell viability using the TB assay, cells were washed with PBS (Thermo Fisher Scientific #10010023), trypsinized and centrifuged at 300 x g for 3 minutes. The resulting pellet was resuspended in media, TB was added to a final concentration of 0.2%, and cells were manually counted using a hemocytometer.

Cell death assay (LDH)

The amount of LDH in the supernatant was measured using the LDH-Glo Cytotoxicity Assay (Promega J2380) following the manufacturer's protocol. The samples were mixed with reagents on microplates, and luminescence was measured after a 30-minute incubation at room temperature. The maximum amount of LDH in each was measured by fully lysing replicate wells with 2% Triton X-100 (Thermo Fisher Scientific #A16046.AE). Background LDH signal was measured from media-only wells and subtracted from sample values before normalization to fully lysed wells in order to determine the amount of cytotoxicity per sample.

Compound sources

ML162 (SML0521), RSL3 (SML2234), erastin (E7781), 2'2'-Bipyridyl (D216305), and necrostatin-1 (N9037°C) were obtained from MilliporeSigma. Ferrostatin-1 (S7243), Liproxstatin-1 (S7699), and z-VAD-FMK (S7023) were purchased from Selleck Chemicals LLC. Deferoxamine mesylate (ab120727) was purchased from Abcam.

Genome-wide library design

sgRNAs targeting the hamster genome build GCF_017639785.1 (BCM_Maur_2.0) were

designed based on NCBI Refseq gene annotations. Protein-coding regions of all gene exons

were filtered to retain only the most constitutively expressed exons and extended 20 nt on each

end. These regions were provided as input for CRISPOR[49] to pick and score all potential

guides. For each gene, guides were filtered by potential problem flags and then ranked by an

overall score derived from its efficiency (the Doench16/Fusi score), a custom motif penalty, a

fractional frameshift (from the Lindel score and the fractional distance to the beginning of the

region), and its specificity (derived from the MIT specificity score). For each gene, after the

guide with the best score was chosen, subsequent guides were also penalized by a location

score, adding a penalty for presence in the same exon and proximity to previously selected

guides for the same gene. If more than ten potential guides were identified for a gene, the ten

with the highest scores were selected.

For the human genome-wide library, we obtained a list of protein-coding genes in human based

on Ensembl release 98. The top five sgRNAs were picked for protein-coding genes and

controls, for a total of 102223 sgRNAs. Human intergenic sgRNAs were chosen by first

subsetting the list of all designed intergenic-region-targeted sgRNAs by Rank = 1, then requiring

the sgRNAs to target only one genomic locus. This list of 465 sgRNAs was ranked by MIT

specificity score in descending order, and the top 449 sgRNAs were chosen.

Library cloning

For the hamster library, 214,116 unique protospacer sequences targeting ~21,000 unique gene

symbols (> 99% with 10 sgRNAs/gene symbol), along with 2,299 intergenic-targeting and 250

nontargeting sequences were synthesized as an oligonucleotide pool (Agilent Technologies).

Separately, for the human library, 98,077 unique protospacer sequences targeting ~19,600

Ensembl transcript IDs (> 99.9% with 5 sgRNAs/gene), 449 intergenic-targeting sequences, 50

non-targeting sequences, and one sequence targeting the *AAVS1* safe harbor locus were

synthesized as an oligonucleotide pool (Agilent Technologies). A guanine nucleotide was

prepended to each 20-nucleotide protospacer sequence that began with A, C, or T. For both

hamster and human libraries, homology arms were prepended (5'-

TATCTTGTGGAAAGGACGAAACACC-3') and appended (5'-

GTTTAAGAGCTATGCTGGAAACAGCATAGC-3'). Additional adapters were pre- and appended

to the hamster library to enable subpool amplification if desired (subpool 1: 5'-

TCGGTGTATTGCTAGTGCGAACCCA-3' and 5'-ATCGTGTGAAAGGTGCCGCTATTGC-3';

subpool 2: 5'-GGTTCGTTCTACACATGGAAGCGGC-3' and 5'-

GAGGACTTCGAGTAGAACGCTGGCG-3'). Oligonucleotide pools were PCR amplified (forward

primer: 5'-TTTCTTGGCTTTATATATCTTGTGGAAAGGACGAAACACC-3'; reverse primer: 5'-

ATTTAAACTTGCTATGCTGTTTCCAGCATAGCTCTTAAAC-3') and cloned into

pLentiCRISPRv2-Opti (a gift from David Sabatini; Addgene plasmid # 163126;

http://n2t.net/addgene:163126; RRID:Addgene_163126). Briefly, 0.25 µL of a 100 nM pool was

PCR amplified in 50 µL reactions using Q5 HotStart DNA Polymerase (New England Biolabs)

for 10 cycles under the following conditions:

1 cycle   98 °C   2 minutes

10 cycles   98 °C   10 seconds

740   55-62 °C   15 seconds

741   72 °C   15 seconds

1 cycle   72 °C   2 minutes

1 cycle   4 °C   hold

All concentration steps were performed using a DNA Clean and Concentrator-5 kit

(ZymoResearch). Amplification from 8 gradient annealing reactions (55-62 °C) was assessed,

and successful reactions were pooled and concentrated. The vector was digested overnight at

37 °C with FastDigest Esp3I and FastAP (ThermoFisher Scientific), gel purified using a

Zymoclean gel DNA recovery kit (ZymoResearch), and concentrated. NEBuilder HiFi DNA

Assembly Master Mix (New England Biolabs) was used in 2 x 40-80 µL bulk assembly

reactions, for a combined total reaction volume of 160 µL containing 4 µg of vector and 160 ng

of PCR amplicon (hamster) or 120 µL total containing 3 µg of vector and 120 ng of PCR

amplicon (human). Each bulk reaction was distributed in 5 µL aliquots and incubated for 10

minutes at 52.2 °C, pooled, concentrated, introduced into Endura Electrocompetent DUO cells

(Lucigen) by electroporation, and plated on 8-16 LB agar with 75 µg/mL carbenicillin in 245 mm

x 245 mm square bioassay dishes. A dilution series was also plated to assess electroporation

efficiency. Cells were incubated overnight at 30 °C, collected, and DNA was isolated using a

ZymoPURE II Plasmid DNA Maxiprep kit (ZymoResearch). Plasmid from separate

electroporations was combined proportionally based on electroporation efficiency for a

combined total library coverage of > 50-fold for each library. Sequence representation in the

libraries was assessed as described below.

Lentivirus production for CRISPR-Cas9 screen

For large-scale virus production, HEK-293T (1.5×10^7) cells were seeded in T175 cm^2 flasks in

DMEM (Thermo Fisher Scientific #12430054) supplemented with 10% fetal bovine serum

(GeminiBio #100-106). Media was changed 24 hours later to 20 mL viral production medium:

IMDM (Thermo Fisher Scientific #1244053) supplemented with 20% heat-inactivated fetal

bovine serum (GeminiBio #100-106). Cells were transfected 8 hours later with a mix containing

76.8 µL Xtremegene-9 transfection reagent (MilliporeSigma #06365779001), 3.62 µg pCMV-

VSV-G (Addgene plasmid # 8454; http://n2t.net/addgene:8454; RRID:Addgene_8454)[50], 8.28

µg psPAX2 (a gift from Didier Trono; Addgene plasmid # 12260; http://n2t.net/addgene:12260;

RRID:Addgene_12260), and 20 µg sgRNA/Cas9 plasmid and Opti-MEM (Thermo Fisher

Scientific #11058021) to a final volume of 1 mL. Media was changed 16 hours later to 55 mL

fresh viral production medium. Viral supernatant was collected and filtered through a 0.45 µm

filter and aliquoted 48 hours post-transfection, then stored at -80°C until use.

CRISPR screen in K562 cells

K562 cells (3.9×10^8) were transduced with a pooled genome-wide lentiviral sgRNA library in a

Cas9-containing vector at MOI <1. The transduced cells were selected with 3 µg/mL puromycin

(Thermo Fisher Scientific #A1113803), and 1×10^8 cells were passaged every 48-72 hours at a

density of 6×10^5 cells/T175 cm^2 flask in 45 mL RPMI 1640 (Gibco #11875093) medium

supplemented with 10% fetal bovine serum (GeminiBio #100-106) and 1%

penicillin/streptomycin for the duration of the screen. At 5 days post-puromycin selection, 1×10^8

cells were exposed to the cold with or without 1 µM ferrostatin-1 (Selleck Chemicals #S7243).

K562 cells (1×10^8) were collected from the each of surviving populations after cold exposure

and/or rewarming as well as a matched untreated population. Genomic DNA was isolated using

the QIAmp DNA Blood Maxiprep kit (Qiagen # 51192), and high-throughput sequencing libraries

were prepared.

CRISPR screen in BHK-21 cells

BHK-21 cells (8.55×10^8) were transduced with a pooled genome-wide lentiviral sgRNA library in

a Cas9-containing vector at an MOI of 0.25. Transduced cells were selected with 2 µg/mL

puromycin (Thermo Fisher Scientific #A1113803), and 2.56×10^8 cells were passaged every 48-

72 hours at a density of 5×10^6 cells/15-cm dish in DMEM (Thermo Fisher Scientific #12430054)

medium supplemented with 10% fetal bovine serum (GeminiBio #100-106) and 1%

penicillin/streptomycin for the duration of the screen. At 6 days post-puromycin selection,

2.56×10^8 cells were exposed to the cold. Cells (2.4×10^8) were collected from each of the

surviving populations after cold exposure and/or rewarming as well as from a matched

untreated population. During the cycles, cells were placed at 4°C for 4 days, then rewarmed to

37°C for 24 hours, reseeded, and after a further 24 hours at 37°C were placed back at 4°C to

begin a subsequent cycle. Genomic DNA was isolated using the QIAmp DNA Blood Maxiprep

kit (Qiagen # 51192), and high-throughput sequencing libraries were prepared.

Sequencing library preparation

The QIAamp DNA Blood Maxiprep Kit (Qiagen # 51192) was used to extract genomic DNA

(gDNA) from cell pellets of 2.5-5x10^7 cells according to manufacturer's instructions with minor

modifications: QIAGEN Protease was replaced with 500 µL of 10 mg/mL Proteinase K

(MilliporeSigma # 3115879001) in water, and cells were lysed overnight; centrifugation was

performed for 2 minutes and 5 minutes after the first and second wash, respectively; 1 mL of

water preheated to 70 °C was used to elute gDNA, followed by centrifugation for 5 minutes.

gDNA was quantified using the Qubit dsDNA HS Assay kit (Thermo Fisher Scientific #Q32851).

PCR amplification of sgRNA sequences was performed in 50 µL reactions using ExTaq

Polymerase (Takara Bio #RR001B) with the following program:

1 cycle   95 °C 5 minutes

16, 24, or 28 cycles   95 °C 10 seconds

817   60 °C 15 seconds

818   72 °C 45 seconds

1 cycle   72 °C 5 minutes

1 cycle   4 °C   hold

Using the following primers:

Forward: 5'- AATGATACGGCGACCACCGAGATCTACACCCCACTGACGGGCACCGGA - 3'

Reverse: 5'- CAAGCAGAAGACGGCATACGAGATCnnnnnnTTTCTTGGGTAGTTTGCAGTTTT

- 3'

Where "nnnnnn" denotes the barcode used for demultiplexing.

Plasmid libraries were amplified for 16 cycles using 10 ng input per 50 µL reaction. gDNA was

initially amplified for 24 (human) or 28 (hamster) cycles in 50 µL test PCR reactions with 1, 3, 4,

5, or 6 µg input. An additional 20-75 reactions were performed using 3-6 µg per reaction

(sampling 150 µg or 300 µg gDNA total for human and hamster screens, respectively). Select-a-

Size DNA Clean and Concentrator (Zymo Research #D4080) or HighPrep PCR (MagBio

Genomics) was used to purify 95-100 µL of each pooled sample, which were eluted with 12-20

µL water and quantified using the Qubit dsDNA HS Assay kit prior to sequencing for 50 cycles

on an Illumina Hiseq 2500 or NovaSeq using the following primers:

Read 1 sequencing primer: 5'-

GTTGATAACGGACTAGCCTTATTTAAACTTGCTATGCTGTTTCCAGCATAGCTCTTAAAC - 3'

Index sequencing primer: 5'-

TTTCAAGTTACGGTAAGCATATGATAGTCCATTTTAAAACATAATTTTAAAACTGCAAACTAC

CCAAGAAA - 3'

CRISPR screen data analysis

Sequencing reads from the human screen were trimmed and mapped to the sgRNA library

using Bowtie version 1.0.0[51], allowing no mismatches, and counted. Sequencing reads from the

hamster screen were trimmed and mapped to the sgRNA library using a combination of

command line prompts and a custom Python function which generated a dataframe of counts

which perfectly (no permissibility for any base mismatches) matched sequenced reads to the full

guide library (code supplied on Github). Data from both human and hamster screens was

similarly analyzed using MAGeCK version 0.5.9.5[52] using a gene test false discovery rate (FDR)

threshold of 0.05, the FDR method for p-value adjustment, and the median as the gene-level

scoring metric.

PANTHER overrepresentation test was conducted on a set of 204 genes (FDR < 0.1) from the

third cycle of the human K562 CRISPR-Cas9 screen ['Analyzed List']. A list of genes expressed

in K562 cells at 37°C collected from bulk RNA sequencing was used as the ['Reference List']:

https://pantherdb.org/tools/compareToRefList.jsp. Fisher's Exact test type and false discovery

rate correction were utilized. Bulk RNA sequencing of K562 cells at 37°C was conducted by

generating a cDNA library prepared by depleting rRNA using the NEBNext rRNA Depletion Kit

(Human/Mouse/Rat) (New England Biolabs) and then the NEBNext Ultra II Directional RNA

Library Prep Kit (New England Biolabs). cDNA libraries were amplified using appropriate

multiplexed index primers for PCR.

Data was visualized using R version 4.2.1[53] corrplot package version 0.92[54] and base graphics,

and GraphPad Prism version 10.

Generation of stable CRISPR/Cas9-targeted cell lines

Homology arms were prepended (5'-TATCTTGTGGAAAGGACGAAACACC-3') and appended
(5'-GTTTAAGAGCTATGCTGGAAACAGCATAGC-3') to the top two ranked human or hamster
GPX4-targeting guides from the genome-wide screens. These guides were PCR amplified and
cloned into pLentiCRISPRv2-Opti (a gift from David Sabatini; Addgene plasmid # 163126;
http://n2t.net/addgene:163126; RRID:Addgene_163126). K562 and BHK-21 cells were then
infected with LentiCRISPRv2. Transduced cells were selected with 3 µg/mL puromycin (Thermo
Fisher Scientific #A1113803) for K562 cells and 2 µg/mL for BHK-21 cells beginning two days
after infection. After one passage, cells were sorted (BD FACSAria™ Fusion Flow Cytometer)
into 96-well plates, at one cell per well to generate clonal knockout cell lines. Cells were cultured
in media with 2.5 µM liproxstatin-1 (Selleck Chemicals #S7699) to prevent cell death. Western
blotting was used to confirm GPX4 loss in individual knockout clones prior to further use.

Human sgRNA:
5'- CGTGTGCATCGTCACCAACG - 3'
5'- CTTGGCGGAAAACTCGTGCA - 3'

Hamster sgRNA:
5'- CGTGTGCATCGTCACCAACG - 3'
5'- CTTGGCTGAGAATTCGTGCA - 3'

Generation of CRISPR/Cas9-targeted human kidney fibroblast pooled cells

Primary human kidney fibroblasts were infected with LentiCRISPRv2 with sgRNAs targeting
GPX4, using the top two ranked *GPX4*-targeting guides from the genome-wide human k562
screen. Transduced cells were selected with 1 µg/mL puromycin (Thermo Fisher Scientific
#A1113803) beginning four days after infection. Cells were cultured in media with 2.5 µM
liproxstatin-1 (Selleck Chemicals #S7699) to prevent cell death. Western blotting was used to
confirm GPX4 loss in individual knockout clones prior to further use.

Immuoblotting

Cells were plated at 6×10^5 cells per well in a 6-well plate and maintained overnight at 37°C.
Cells were then collected when wells were confluent. The media was aspirated, and cells were
washed with ice-cold PBS (Thermo Fisher Scientific #10010023). Cells were scraped in 100 µL

of RIPA buffer (150 mM NaCl, 1% Triton X-100, 0.5% Na-Dexycholate, 0.1% SDS, 50 mM Tris,

pH 8) with protease inhibitors (MilliporeSigma #04693159001) and collected in microcentrifuge

tubes. The contents were then vortexed for 30 minutes at 4°C and centrifuged at 16000 x g for

20 minutes at 4°C. Protein content of the samples was measured using the Pierce BCA Protein

Assay kit (Thermo Fisher Scientific #23225) and resuspended in 100 µL of RIPA buffer with

protein loading dye (10% SDS, 500 mM DTT, 50% Glycerol, 250 mM Tris-HCl and 0.5%

bromophenol blue dye, pH 6.8). Protein samples were boiled for 5 minutes at 95°C, vortexed,

and centrifuged at 16000 x g for 5 minutes before loading onto a gel. The gel was run at 120 V

for 2 hours before being transferred to a Polyvinylidene difluoride (PVDF) membrane (Thermo

Fisher Scientific #88518). The transfer cassette was run at 45 V for 2 hours. The membranes

were then transferred to 5% milk TBST (1X Tris-Buffered Saline, 0.1% Tween® 20 Detergent)

and shaken for 15 minutes before being washed in TBST. Primary antibodies recognizing GPX4

(Abcam #ab41787) or HA (Cell Signaling Technology #3724) were added at 1:1000 in 5% BSA

(MilliporeSigma #A9418) and incubated overnight at 4°C. The membrane was then washed 3

times in 1x TBST and 1:3000 anti-rabbit secondary antibody (Cell Signaling Technology #7074)

in 5% milk TBST was added. Membranes were incubated for 1 hour at room temperature and

washed 3x in 1x TBST. Membranes were then incubated in Pierce™ ECL Western Blotting

Substrate (Thermo Fisher Scientific #32106) for 5 minutes before imaging.

Generation of HA-GPX4 and HA-GFP overexpression constructs

To generate GPX4-overexpression lines, entire human *GPX4* (cytosolic; NM_001367832.1) and

hamster *Gpx4* were amplified from the cDNA of K562 and BHK-21 cells, respectively. The

cloned transcript fragments include the 3' UTR that contains the SECIS element necessary for

selenocysteine incorporation. The *GPX4* genomic DNA fragments were cloned into pLJM1

(Addgene plasmid # 19319; http://n2t.net/addgene:19319; RRID:Addgene_19319) and

confirmed via sequencing. To create GPX4 mutants, (U46S), non-overlapping primers were

used to create mutations via inverse PCR, as previously described[55], converting the UGA stop

codon to UCA, encoding serine. GFP was amplified from pLJM1-EGFP (Addgene plasmid #

19319; http://n2t.net/addgene:19319; RRID:Addgene 19319) with Gibson overhangs to enable

insertion into a lentivirus vector. HA-GPX4 and HA-GFP DNA fragments were then amplified by

PCR and cloned into blasticidin lentiviral vectors [gift from Whitney Henry at the Whitehead

Institute for Biomedical Research] using standard cloning methods, with point mutations in the

PAM and guide sequences to prevent targeting of exogenously expressed *GPX4*.

HEK-293T cells (5×10^5) were seeded in 2 wells of a 6-well plate in DMEM (Thermo Fisher

Scientific #12430054) supplemented with 10% fetal bovine serum (GeminiBio #100-106).

Media was changed 16 hours later to 2 mL per well of viral production medium: IMDM (Thermo

Fisher Scientific #1244053) supplemented with 20% heat-inactivated fetal bovine serum

(GeminiBio #100-106). Cells were transfected 8 hours later with a mix containing 4.22 µL

Xtremegene-9 transfection reagent (MilliporeSigma #06365779001), 181 ng pCMV-VSV-G

(Addgene plasmid # 8454; http://n2t.net/addgene:8454; RRID:Addgene_8454)[50], 414 ng

psPAX2 (a gift from Didier Trono; Addgene plasmid # 12260; http://n2t.net/addgene:12260;

RRID:Addgene_12260), and 1 µg HA-GPX4 or HA-GFP plasmid and Opti-MEM (Thermo Fisher

Scientific #11058021) to a final volume of 50 µL per well. Media was changed 16 hours later to

2.5 mL per well of fresh viral production medium. Viral supernatant was collected and aliquoted

48 hours post-transfection, then stored at -80°C until use.

K562 or BHK-21 cells (6×10^5, wild-type and GPX4 knockout clones) were transduced with the

lentiviral HA-tagged GPX4 or GFP in the presence of 10 µg/mL polybrene. After 8 hours of

incubation at 37°C, media was changed to virus-free media. After 48 hours, the transduced cells

were selected with blasticidin (Thermo Fisher Scientific #A1113903) at a concentration of 6

µg/mL for K562 cells and 4 µg/mL for BHK-21 cells. Overexpression of GPX4 and GFP was

confirmed via immunoblotting.

Primary cell culture

Three hibernator (13-lined ground squirrel postnatal dermal fibroblasts [gift from Wei Li at Duke

University], SV40-immortalized Greater horseshoe bat embryonic dermal fibroblasts [gift from

Rudolf Jaenisch at the Whitehead Institute for Biomedical Research], and Syrian golden

hamster embryonic primary dermal fibroblasts [isolated]) and five non-hibernator (mouse

embryonic SV40-immortalized dermal fibroblasts [gift from Jonathan Weissman at the

Whitehead Institute for Biomedical Research], adult human kidney fibroblasts [purchased from

ATCC], adult human dermal fibroblasts [purchased from ATCC], adult rat dermal fibroblasts

[purchased from ATCC], and adult mouse fibroblasts [purchased from ATCC]) primary cells

were cultured in Dulbecco's modified Eagle's medium (Thermo Fisher Scientific #12430054)

supplemented with 10% fetal bovine serum (GeminiBio #100-106) and 1%

penicillin/streptomycin (Thermo Fisher Scientific #15140122). Cells were seeded on 24-well

culture plates at ~50,000 cells per well and allowed to adhere for 24 hours at 37°C. Cells were

then placed at 4°C with 5% CO_2 for varying time periods (1, 4, or 7 days), with a 30-minute

rewarming at 37°C before assessing cell viability via trypan blue (TB) (Thermo Fisher Scientific

#15250061) staining. To assess cell viability using the TB assay, cells were washed with PBS

(Thermo Fisher Scientific #10010023), trypsinized and centrifuged at 300 x g for 3 minutes. The

resulting pellet was resuspended in media, TB was added to a final concentration of 0.2%, and

cells were manually counted using a hemocytometer.

Primary neuron isolation and culture

Primary Syrian golden hamster and C57BL/6 mouse cortical neurons were isolated at postnatal

day 0. The day before isolation, standard tissue culture treated plates were coated with 3 µg/mL

Poly-L-Ornithine (MilliporeSigma #P4957) and warmed at 37°C for 24 hours. The morning of

isolation, wells were washed with 1X PBS (Thermo Fisher Scientific #10010023). To isolate

neurons, pups were placed on ice for 2-5 minutes until movement ceased. Afterwards, they

were washed with 10% ethanol, decapitated, and their heads placed in a 10x dissociation media

bath (10.16 g $MgCl_2$ hexahydrate and 11.915 g HEPES in 450 mL HBSS brought to a pH of 7

using NaOH with subsequent addition of 1.182 g of kynurenic acid heated to 65% while stirring

until the solution was fully dissolved and cooled to room temperature and adjusted to a pH of

7.2. The dissociation media was then diluted from 10X to 1X in HBSS and filtered through a

0.22-µm vacuum filter). Dissected brains were placed in fresh dissociation media, where the

cortices were dissected out and transferred to 50 mL Falcon tubes on ice containing 1X

dissociation media. Afterwards, the dissociation media was removed from the cortices and

replaced with papain solution (5 mL 1x dissociation media, 5-7 grains of L-Cyteine, and 172 µL

Worthington papain [crystalline suspension in 50mM sodium acetate, pH 4.5 incubated in

1.1mM EDTA, 0.067mM mercaptoethanol and 5.5mM cysteine-HCl at 37°C. Activates to 20

units per 1 mg protein]) that was preheated to 37°C for 30 minutes. The cortices were incubated

in the papain solution for 3-5 minutes at 37°C. Afterwards, the papain solution was replaced with

trypsin inhibitor solution (10 mg of trypsin inhibitor dissolved in 10 mL of 1X dissociation media,

preheated to 37°C) and incubated subsequently at 37°C for 3-5 minutes, repeated 3 times.

Afterwards, the trypsin inhibitor solution was carefully removed and replaced with 6 mL of

preheated complete neurobasal media (5 mL of GlutaMAX5, 10 mL of B-27, 5 mL of

penicillin/streptomycin in neurobasal media filtered through a 0.22-µm vacuum filter). Using a 5

mL pipette tip, corticies were triturated until no tissue clumps were present. The cortices were

then passed through a 100-µm cell strainer to remove large debris and centrifuged at 300 x g for

3 minutes before resuspension in 5 mL of complete neurobasal media before plating at a

density of 50,000 mixed cortical neuron isolated cells per well of a 24 well plate. Subsequently,

50% of the media was replaced every second day with fresh 37°Ç pre-warmed media.

1003

Statistical analyses and software information

Data are generally plotted as mean ± S.E.M. unless otherwise indicated. No statistical methods

were used to predetermine sample sizes. Unless otherwise indicated, all replication numbers in

the figure legends (n) indicate biological replicates. Statistical significance was determined using

a two-tailed Student's T-test, one-way ANOVA, or two-way ANOVA using Prism 10 software

(GraphPad Software) unless otherwise indicated. Statistical significance was set at $p \leq 0.05$

unless otherwise indicated. Figures were finalized in Adobe Illustrator 2024.

1011

1012 **Supplemental information**

1013 Document S1. Figures S1-S11

1014 Table S2. Excel file containing hamster screen data, related to Figure 2

1015 Table S5. Excel file containing human screen data, related to Figure 4

1016 Table S6. Excel file containing GO_Biological Processes, related to Figure S8

1017

References

1. Cheshire, W.P. (2016). Thermoregulatory disorders and illness related to heat and cold
stress. Autonomic Neuroscience *196*, 91–104. https://doi.org/10.1016/j.autneu.2016.01.001.

2. Physiology, Thermal Regulation - StatPearls - NCBI Bookshelf
https://www.ncbi.nlm.nih.gov/books/NBK499843/.

3. Tan, C.L., and Knight, Z.A. (2018). Regulation of body temperature by the nervous system.
Neuron *98*, 31–48. https://doi.org/10.1016/j.neuron.2018.02.022.

4. Morrison, S.F., and Nakamura, K. (2019). Central Mechanisms for Thermoregulation. Annu
Rev Physiol *81*, 285–308. https://doi.org/10.1146/annurev-physiol-020518-114546.

5. Hendriks, K.D.W., Joschko, C.P., Hoogstra-Berends, F., Heegsma, J., Faber, K.-N., and
Henning, R.H. (2020). Hibernator-Derived Cells Show Superior Protection and Survival in
Hypothermia Compared to Non-Hibernator Cells. IJMS *21*, 1864.
https://doi.org/10.3390/ijms21051864.

6. Ou, J., Ball, J.M., Luan, Y., Zhao, T., Miyagishima, K.J., Xu, Y., Zhou, H., Chen, J., Merriman,
D.K., Xie, Z., et al. (2018). iPSCs from a Hibernator Provide a Platform for Studying Cold
Adaptation and Its Potential Medical Applications. Cell *173*, 851-863.e16.
https://doi.org/10.1016/j.cell.2018.03.010.

7. Hendriks, K.D.W., Lupi, E., Hardenberg, M.C., Hoogstra-Berends, F., Deelman, L.E., and
Henning, R.H. (2017). Differences in mitochondrial function and morphology during cooling
and rewarming between hibernator and non-hibernator derived kidney epithelial cells.

8. Anegawa, D., Sugiura, Y., Matsuoka, Y., Sone, M., Shichiri, M., Otsuka, R., Ishida, N.,
Yamada, K., Suematsu, M., Miura, M., et al. (2021). Hepatic resistance to cold ferroptosis in
a mammalian hibernator Syrian hamster depends on effective storage of diet-derived α-
tocopherol. Commun Biol *4*, 796. https://doi.org/10.1038/s42003-021-02297-6.

9. Park, H.G., Han, S.I., Oh, S.Y., and Kang, H.S. (2005). Cellular responses to mild heat
stress. CMLS, Cell. Mol. Life Sci. *62*, 10–23. https://doi.org/10.1007/s00018-004-4208-7.

10. Richter, K., Haslbeck, M., and Buchner, J. (2010). The Heat Shock Response: Life on the
Verge of Death. Molecular Cell *40*, 253–266. https://doi.org/10.1016/j.molcel.2010.10.006.

11. Karunanithi, S., and Brown, I.R. (2015). Heat shock response and homeostatic plasticity.
Front. Cell. Neurosci. *9*. https://doi.org/10.3389/fncel.2015.00068.

12. Kruuv, J., Glofcheski, D., Cheng, K.H., Campbell, S.D., Al-Qysi, H.M., Nolan, W.T., and
Lepock, J.R. (1983). Factors influencing survival and growth of mammalian cells exposed to
hypothermia. I. Effects of temperature and membrane lipid perturbers. J Cell Physiol *115*,
179–185. https://doi.org/10.1002/jcp.1041150212.

13. Hochachka, P.W. (1986). Defense strategies against hypoxia and hypothermia. Science
231, 234–241. https://doi.org/10.1126/science.2417316.

14. Rule, G.S., Frim, J., Thompson, J.E., Lepock, J.R., and Kruuv, J. (1978). The effect of
membrane lipid perturbers on survival of mammalian cells to cold. Cryobiology *15*, 408–414.
https://doi.org/10.1016/0011-2240(78)90059-7.

15. Zachariassen, K.E. (1991). Hypothermia and cellular physiology. Arctic Med Res *50 Suppl*
6, 13–17.

16. Heller, H.C., and Hammel, H.T. (1972). CNS control of body temperature during hibernation.
Comp Biochem Physiol A Comp Physiol *41*, 349–359. https://doi.org/10.1016/0300-
9629(72)90066-7.

17. Sunagawa, G.A., and Takahashi, M. (2016). Hypometabolism during Daily Torpor in Mice is
Dominated by Reduction in the Sensitivity of the Thermoregulatory System. Sci Rep *6*,
37011. https://doi.org/10.1038/srep37011.

18. Geiser, F. (2004). Metabolic rate and body temperature reduction during hibernation and
daily torpor. Annu Rev Physiol *66*, 239–274.
https://doi.org/10.1146/annurev.physiol.66.032102.115105.

19. Chayama, Y., Ando, L., Tamura, Y., Miura, M., and Yamaguchi, Y. (2016). Decreases in
body temperature and body mass constitute pre-hibernation remodelling in the Syrian
golden hamster, a facultative mammalian hibernator. R Soc Open Sci *3*, 160002.
https://doi.org/10.1098/rsos.160002.

20. Park, K.J., Jones, G., and Ransome, R.D. (2000). Torpor, arousal and activity of hibernating
Greater Horseshoe Bats (Rhinolophus ferrumequinum). Functional Ecology *14*, 580–588.
https://doi.org/10.1046/j.1365-2435.2000.t01-1-00460.x.

21. Cooper, S.T., Richters, K.E., Melin, T.E., Liu, Z., Hordyk, P.J., Benrud, R.R., Geiser, L.R.,
Cash, S.E., Simon Shelley, C., Howard, D.R., et al. (2012). The hibernating 13-lined ground
squirrel as a model organism for potential cold storage of platelets. Am J Physiol Regul
Integr Comp Physiol *302*, R1202–R1208. https://doi.org/10.1152/ajpregu.00018.2012.

22. Heldmaier, G., Ortmann, S., and Elvert, R. (2004). Natural hypometabolism during
hibernation and daily torpor in mammals. Respiratory Physiology & Neurobiology *141*, 317–
329. https://doi.org/10.1016/j.resp.2004.03.014.

23. Harris, R.A., Raveendran, M., Lyfoung, D.T., Sedlazeck, F.J., Mahmoud, M., Prall, T.M.,
Karl, J.A., Doddapaneni, H., Meng, Q., Han, Y., et al. (2022). Construction of a new
chromosome-scale, long-read reference genome assembly for the Syrian hamster,
Mesocricetus auratus. Gigascience *11*, giac039.
https://doi.org/10.1093/gigascience/giac039.

24. Hart, T., Tong, A.H.Y., Chan, K., Van Leeuwen, J., Seetharaman, A., Aregger, M.,
Chandrashekhar, M., Hustedt, N., Seth, S., Noonan, A., et al. (2017). Evaluation and Design
of Genome-Wide CRISPR/SpCas9 Knockout Screens. G3 (Bethesda) *7*, 2719–2727.
https://doi.org/10.1534/g3.117.041277.

25. Mirabello, L., Macari, E.R., Jessop, L., Ellis, S.R., Myers, T., Giri, N., Taylor, A.M., McGrath,
K.E., Humphries, J.M., Ballew, B.J., et al. (2014). Whole-exome sequencing and functional
studies identify RPS29 as a novel gene mutated in multicase Diamond-Blackfan anemia
families. Blood *124*, 24–32. https://doi.org/10.1182/blood-2013-11-540278.

26. Summer, S., Smirnova, A., Gabriele, A., Toth, U., Fasemore, A.M., Förstner, K.U., Kuhn, L.,
Chicher, J., Hammann, P., Mitulović, G., et al. (2020). YBEY is an essential biogenesis
factor for mitochondrial ribosomes. Nucleic Acids Res *48*, 9762–9786.
https://doi.org/10.1093/nar/gkaa148.

27. Ni, M., Solmonson, A., Pan, C., Yang, C., Li, D., Notzon, A., Cai, L., Guevara, G., Zacharias,
L.G., Faubert, B., et al. (2019). Functional Assessment of Lipoyltransferase-1 Deficiency in
Cells, Mice, and Humans. Cell Rep *27*, 1376-1386.e6.
https://doi.org/10.1016/j.celrep.2019.04.005.

28. Xu, G., Li, W., Zhao, Y., Fan, T., Gao, Q., Wang, Y., Zhang, F., Gao, M., An, Z., and Yang,
Z. (2024). Overexpression of Lias Gene Alleviates Cadmium-Induced Kidney Injury in Mice
Involving Multiple Effects: Metabolism, Oxidative Stress, and Inflammation. Biol Trace Elem
Res *202*, 2797–2811. https://doi.org/10.1007/s12011-023-03883-x.

29. Douce, R., Bourguignon, J., Neuburger, M., and Rébeillé, F. (2001). The glycine
decarboxylase system: a fascinating complex. Trends Plant Sci *6*, 167–176.
https://doi.org/10.1016/s1360-1385(01)01892-1.

30. Weaver, K., and Skouta, R. (2022). The Selenoprotein Glutathione Peroxidase 4: From
Molecular Mechanisms to Novel Therapeutic Opportunities. Biomedicines *10*, 891.
https://doi.org/10.3390/biomedicines10040891.

31. Seibt, T.M., Proneth, B., and Conrad, M. (2019). Role of GPX4 in ferroptosis and its
pharmacological implication. Free Radical Biology and Medicine *133*, 144–152.
https://doi.org/10.1016/j.freeradbiomed.2018.09.014.

32. Varlamova, E.G., Goltyaev, M.V., Novoselov, S.V., Novoselov, V.I., and Fesenko, E.E.
(2013). Selenocysteine biosynthesis and mechanism of incorporation into growing proteins.
Mol Biol *47*, 488–495. https://doi.org/10.1134/S0026893313040134.

33. French, R., and Simonovic, M. (2012). Synthesis and decoding of selenocysteine and
human health. Croatian medical journal *53*, 535–550.
https://doi.org/10.3325/cmj.2012.53.535.

34. Chen, T., Leng, J., Tan, J., Zhao, Y., Xie, S., Zhao, S., Yan, X., Zhu, L., Luo, J., Kong, L., et
al. (2023). Discovery of Novel Potent Covalent Glutathione Peroxidase 4 Inhibitors as Highly
Selective Ferroptosis Inducers for the Treatment of Triple-Negative Breast Cancer. J. Med.
Chem. *66*, 10036–10059. https://doi.org/10.1021/acs.jmedchem.3c00967.

35. Shintoku, R., Takigawa, Y., Yamada, K., Kubota, C., Yoshimoto, Y., Takeuchi, T., Koshiishi,
I., and Torii, S. (2017). Lipoxygenase-mediated generation of lipid peroxides enhances
ferroptosis induced by erastin and RSL3. Cancer Science *108*, 2187–2194.
https://doi.org/10.1111/cas.13380.

36. Sui, X., Zhang, R., Liu, S., Duan, T., Zhai, L., Zhang, M., Han, X., Xiang, Y., Huang, X., Lin,
H., et al. (2018). RSL3 Drives Ferroptosis Through GPX4 Inactivation and ROS Production
in Colorectal Cancer. Front. Pharmacol. *9*. https://doi.org/10.3389/fphar.2018.01371.

37. Hattori, K., Ishikawa, H., Sakauchi, C., Takayanagi, S., Naguro, I., and Ichijo, H. (2017).
Cold stress-induced ferroptosis involves the ASK1-p38 pathway. EMBO Rep *18*, 2067–
2078. https://doi.org/10.15252/embr.201744228.

38. Doll, S., Freitas, F.P., Shah, R., Aldrovandi, M., Da Silva, M.C., Ingold, I., Goya Grocin, A.,
Xavier Da Silva, T.N., Panzilius, E., Scheel, C.H., et al. (2019). FSP1 is a glutathione-
independent ferroptosis suppressor. Nature *575*, 693–698. https://doi.org/10.1038/s41586-
019-1707-0.

39. Bersuker, K., Hendricks, J.M., Li, Z., Magtanong, L., Ford, B., Tang, P.H., Roberts, M.A.,
Tong, B., Maimone, T.J., Zoncu, R., et al. (2019). The CoQ oxidoreductase FSP1 acts
parallel to GPX4 to inhibit ferroptosis. Nature *575*, 688–692. https://doi.org/10.1038/s41586-
019-1705-2.

40. Hu, Q., Wei, W., Wu, D., Huang, F., Li, M., Li, W., Yin, J., Peng, Y., Lu, Y., Zhao, Q., et al.
(2022). Blockade of GCH1/BH4 Axis Activates Ferritinophagy to Mitigate the Resistance of
Colorectal Cancer to Erastin-Induced Ferroptosis. Front Cell Dev Biol *10*, 810327.
https://doi.org/10.3389/fcell.2022.810327.

41. Ma, T., Du, J., Zhang, Y., Wang, Y., Wang, B., and Zhang, T. (2022). GPX4-independent
ferroptosis—a new strategy in disease's therapy. Cell Death Discov. *8*, 1–8.
https://doi.org/10.1038/s41420-022-01212-0.

42. Zhang, W., Dai, J., Hou, G., Liu, H., Zheng, S., Wang, X., Lin, Q., Zhang, Y., Lu, M., Gong,
Y., et al. (2023). SMURF2 predisposes cancer cell toward ferroptosis in GPX4-independent

manners by promoting GSTP1 degradation. Molecular Cell *83*, 4352-4369.e8. https://doi.org/10.1016/j.molcel.2023.10.042.

43. Sone, M., Mitsuhashi, N., Sugiura, Y., Matsuoka, Y., Maeda, R., Yamauchi, A., Okahashi, R., Yamashita, J., Sone, K., Enju, S., et al. (2023). Identification of genes supporting cold resistance of mammalian cells: lessons from a hibernator. Preprint at bioRxiv, https://doi.org/10.1101/2023.12.27.573489 https://doi.org/10.1101/2023.12.27.573489.

44. Davey, H.M., and Hexley, P. (2011). Red but not dead? Membranes of stressed Saccharomyces cerevisiae are permeable to propidium iodide. Environ Microbiol *13*, 163–171. https://doi.org/10.1111/j.1462-2920.2010.02317.x.

45. Kirchhoff, C., and Cypionka, H. (2017). Propidium ion enters viable cells with high membrane potential during live-dead staining. Journal of Microbiological Methods *142*, 79–82. https://doi.org/10.1016/j.mimet.2017.09.011.

46. Li, F.-J., Long, H.-Z., Zhou, Z.-W., Luo, H.-Y., Xu, S.-G., and Gao, L.-C. (2022). System Xc - /GSH/GPX4 axis: An important antioxidant system for the ferroptosis in drug-resistant solid tumor therapy. Front Pharmacol *13*, 910292. https://doi.org/10.3389/fphar.2022.910292.

47. Forman, H.J., Zhang, H., and Rinna, A. (2009). Glutathione: Overview of its protective roles, measurement, and biosynthesis. Mol Aspects Med *30*, 1–12. https://doi.org/10.1016/j.mam.2008.08.006.

48. Yoo, S.-E., Chen, L., Na, R., Liu, Y., Rios, C., Remmen, H.V., Richardson, A., and Ran, Q. (2012). Gpx4 ablation in adult mice results in a lethal phenotype accompanied by neuronal loss in brain. Free Radic Biol Med *52*, 1820–1827. https://doi.org/10.1016/j.freeradbiomed.2012.02.043.

49. Concordet, J.-P., and Haeussler, M. (2018). CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. Nucleic Acids Res *46*, W242–W245. https://doi.org/10.1093/nar/gky354.

50. Stewart, S.A., Dykxhoorn, D.M., Palliser, D., Mizuno, H., Yu, E.Y., An, D.S., Sabatini, D.M., Chen, I.S.Y., Hahn, W.C., Sharp, P.A., et al. (2003). Lentivirus-delivered stable gene silencing by RNAi in primary cells. RNA *9*, 493–501. https://doi.org/10.1261/rna.2192803.

51. Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biol *10*, R25. https://doi.org/10.1186/gb-2009-10-3-r25.

52. Li, W., Xu, H., Xiao, T., Cong, L., Love, M.I., Zhang, F., Irizarry, R.A., Liu, J.S., Brown, M., and Liu, X.S. (2014). MAGeCK enables robust identification of essential genes from genome-scale CRISPR/Cas9 knockout screens. Genome Biol *15*, 554. https://doi.org/10.1186/s13059-014-0554-4.

53. R Core Team (2022). R: A Language and Environment for Statistical Computing | BibSonomy. R Foundation for Statistical Computing.

54. Wei T, S.V. (2021). R package "corrplot": Visualization of a Correlation Matrix (Version 0.92). https://github.com/taiyun/corrplot.

1192   55. Silva, D., Santos, G., Barroca, M., and Collins, T. (2017). Inverse PCR for Point Mutation
1193   Introduction. In PCR: Methods and Protocols Methods in Molecular Biology., L. Domingues,
1194   ed. (Springer), pp. 87–100. https://doi.org/10.1007/978-1-4939-7060-5_5.

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226 **Supplement**

1227

1228

Supplement 1. Permeability to trypan blue changes rapidly upon cell rewarming

a, Number of viable K562 cells based on trypan blue staining after one day at 4°C and

subsequent rewarming for 24 hours at 37°C. Numbers are normalized to initial cell counts. Dots

indicate viable cell number based on trypan blue staining of cells after incubation at room

temperature for 15, 30, or 60 minutes. Blue shaded regions indicate 4°C exposure and shaded

pink regions indicate 37°C exposure. Red square indicates calculated cell counts after one day

at 4°C based on the viable cell number measured after 24 hour rewarming. **b**, Viability of K562

cells was assessed by trypan blue staining after incubation at room temperature for 0, 15, 30, or

60 minutes following 24 hours at 4°C (n = 4). Cells incubated at room temperature for 15

minutes show a significant increase in cell counts compared to cells counted immediately (**P =

0.0016). Cells incubated at room temperature for 30 minutes show a significant increase in cell

counts compared to a 15-minute incubation (**P = 0.0023), while no significant difference in

viability was observed between cells incubated for 30 or 60 minutes (ns, P = 0.4059). **c-f**,

Viability of cells was assessed by trypan blue staining after incubation at 37°C for 0, 15, 30, or

60 minutes following 24 hours at 4°C (n = 4). **c**, HEK293T, **d**, RPE1, **e**, HeLa, **f**, BHK-21. No

significant difference in cell viability was observed between cells incubated for 30 or 60 minutes

for HEK293T (ns, P = 0.3122), RPE1 (ns, P = 0.5137), HeLa (ns, P = 0.9735), and BHK-21 (ns,

P = 0.9998) cells. All values show mean ± SEM, with significance determined by one-way

ANOVA adjusted for multiple comparisons by Tukey's HSD. *P < 0.05; **P < 0.01; ***P < 0.001;

****P < 0.0001; ns P > 0.05.

1249

1250

Supplement 2. Depletion of Core Essential Genes in Genome-Wide BHK-21 Screens

a-d, Genes ranked by median fold-change (log2) in genome-wide BHK-21 screens. **a**, after

three cycles of cold exposure and rewarming (Cycle 3 4°C), **b**, matched constant 37°C control

condition (Cycle 3 37°C), **c**, after 15 days of 4°C exposure (15 Days 4°C), **d**, matched constant

37°C control condition (15 Days 37°C). Core essential genes[24] indicated in red are positioned

below based on gene rank to demonstrate their depletion in each screen condition. **e-f**, Boxplots

showing log2 fold change in representation for the population of control sgRNAs (gray; n = 250)

or sgRNAs targeting core essential genes[24] (blue; n = 4635) over **e)** three cycles of cold

exposure and rewarming (4°C) or **f)** constant 37°C control conditions. The line within each box

represents the median, the bounds of each box represent the first and third quartiles, and the

whiskers extend to the furthest data point within 1.5 times the interquartile range. A two-sided

Kolmogorov-Smirnov test was used to test the difference between each pair of control/essential-

gene-targeting sgRNA distributions (estimated p-value < 2.2e-16 for all six pairs in e and f).

1264

1265

1266

Supplement 3. RSL3 treatment has no effect on the viability of cold-exposed *Gpx4* KO BHK-21 cells.

a-d, Wild-type BHK-21 cells **(a)** and three independent *Gpx4* KO BHK-21 clonal lines **(b-d)** were treated with RSL3 (1 μM) and ferrostatin-1 (Fer-1, 1 μM) as indicated for 24 hours at 4°C (*n* = 4) prior to trypan blue staining. Wild-type BHK-21 cells show a significant decrease in cell viability when treated with RSL3 compared to no treatment (**P = 0.0013), whereas *Gpx4* KO lines show no significant changes in viability (ns, P > 0.05). All values show mean ± SEM, with significance measured by one-way ANOVA adjusted for multiple comparisons by Dunnett's test. *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; ns P > 0.05.

Supplement 4. Growth of BHK21 and K562 *GPX4* knockout cells and effects of RSL3

treatment on human kidney fibroblast *GPX4* knockout cells

a, Viable K562 cells recorded as percentage relative to start based on trypan blue staining over

the course of 3 days at 37°C. Cell growth is significantly decreased in *GPX4* K562 KO clones

compared to WT K562 cells. Supplementation of liproxstatin-1 (2.5 µM) increases cell viability in

GPX4 KO cells and not WT cells at 37°C. **b**, Viable BHK-21 cells recorded as percentage

relative to start based on trypan blue staining over the course of 2 days. Cell growth is

significantly decreased in *Gpx4* BHK-21 KO cells compared to WT BHK-21 cells.

Supplementation of liproxstatin-1 (2.5 µM) increases cell viability in *Gpx4* KO cell and not WT

cells at 37°C. **c**, Human kidney GPX4 KO cells were placed at 4°C and left untreated or treated

with RSL3 (1 µM) for 4, and 7 days (n = 4). Treatment with RSL3 does not confer significant

additional death (n=4 per timepoint and condition) as measure by two-tailed t-test. All values

show mean ± SEM, with significance measured by one-way ANOVA adjusted for multiple

comparisons with Tukey's HSD unless otherwise specified. *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$;

****$P < 0.0001$; ns $P > 0.05$.

1292

1293

1294

1295

1296

1297

1298

1299

1300

Supplement 5. Ferroptosis inhibitors and iron chelators increase cold cell viability in a

dose-dependent manner

a-h, K562 **(a-d)** and HT1080 **(e-h)** cells were treated with varying concentrations of the

ferroptosis inhibitors, ferrostatin-1 and liproxstatin-1, and iron chelators, deferoxamine and 2'2'-

pyridine, for four days at 4°C prior to assaying cell viability by trypan blue staining (n = 3). All

values show mean ± SEM.

1307

1308

1309

Supplement 6. Cells derived from non-hibernators undergo cold-induced ferroptotic cell

death

a, K562 cells were treated with ferrostatin-1 (1 µM) (****$P < 0.0001$), liproxstatin-1 (1 µM) (****P

< 0.0001), Z-VAD-FMK (1 µM) (ns, $P = 0.9760$), or necrostatin-1 (1 µM) (ns, $P = 0.9835$) for 4

days at 4°C prior to trypan blue staining (n = 4). **b**, HT1080 cells treated with ferrostatin-1 (1 µM)

(****$P < 0.0001$), liproxstatin-1 (1 µM) (****$P < 0.0001$), Z-VAD-FMK (1 µM), or necrostatin-1 (1

µM) for 4 days at 4°C prior to trypan blue staining (n = 4). **c**, HeLa cells treated with ferrostatin-1

(1 µM) (****$P < 0.0001$), liproxstatin-1 (1 µM) (****$P < 0.0001$), Z-VAD-FMK (1 µM) (ns, $P >$

0.9999), or necrostatin-1 (1 µM) (ns, $P = 0.9987$) for 10 days at 4°C prior to trypan blue staining

(n = 4). **d**, RPE1 cells treated with ferrostatin-1 (1 µM) (****$P < 0.0001$), liproxstatin-1 (1 µM)

(****$P < 0.0001$), Z-VAD-FMK (1 µM) (ns, $P = 0.9291$), or necrostatin-1 (1 µM) (ns, $P > 0.9999$)

for 10 days at 4°C prior to trypan blue staining (n = 4). **e**, K562 cells treated with deferoxamine

mesylate (5 µM) (***$P = 0.0002$) or 2'2'-pyridine (10 µM) (****$P < 0.0001$) for 4 days at 4°C prior

to trypan blue staining (n = 3). **f**, HT1080 cells treated with deferoxamine mesylate (100 µM)

(****$P < 0.0001$) or 2'2'-pyridine (5 µM) (***$P = 0.0002$) for 4 days at 4°C prior to trypan blue

staining (n = 3). **g**, HeLa cells treated with deferoxamine mesylate (5 µM) (****$P < 0.0001$) or

2'2'-pyridine (5 µM) (****$P < 0.0001$) for 10 days at 4°C prior to trypan blue staining (n = 3). **h**,

RPE1 cells treated with deferoxamine mesylate (100 µM) (****$P < 0.0001$) or 2'2'-pyridine (5

µM) (***$P < 0.0001$) for 10 days at 4°C prior to trypan blue staining ($n = 3$). All values show

mean ± SEM, with significance measured one-way ANOVA adjusted for multiple comparisons

by Dunnett's test. *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.0001$; ns $P > 0.05$.

1331

1332

1333

Supplement 7. Depletion of Core Essential Genes in Genome-Wide K562 Screens

a-d, Genes ranked by median fold-change (log2) in genome-wide K562 screens. **a**, after three

cycles of cold exposure (Cycle 3 4°C), **b**, matched constant 37°C control condition (Cycle 3

37°C), **c**, three cycles of cold exposure with 1 µM ferrostatin-1 (Cycle 3 4°C + Fer-1), **d**,

matched constant 37°C control condition with 1 µM ferrostatin-1 (Cycle 3 37°C + Fer-1). Core

essential genes[24] (red bars) are positioned below based on gene rank to demonstrate their

depletion in each screen condition. **e-f**, Boxplots showing the log2 fold change in representation

for the population of control sgRNAs (gray; n = 500) or sgRNAs targeting core essential genes[24]

(blue; n = 3219) over **e)** three cycles of cold exposure (4°C) or **f)** constant 37°C control

conditions. The line within each box represents the median, the bounds of each box represent

the first and third quartiles, and the whiskers extend to the furthest data point within 1.5 times

the interquartile range. A two-sided Kolmogorov-Smirnov test was used to test the difference

between each pair of control/essential-gene-targeting sgRNA distributions (estimated p-value <

2.2e-16 for all six pairs in e and f).

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

Supplement 8. Top 20 enriched pathways at Cycle 3 (4°C vs. 37°C) in K562 cells

a, Graphical representation of the top 20 enriched, differentially expressed gene sets (204

genes; FDR < 0.1) from the genome-scale CRISPR-Cas9 screen in K562 cells (Cycle 3 4°C vs.

3 passages at 37°C). GO_Biological Processes Panther overrepresentation test was used to

determine enriched gene sets. Functional annotation analysis of the selectively required genes

identified pathways related to translational readthrough, selenocysteine incorporation, protein

insertion into the ER, glycosylation, fatty acid metabolism, and mitochondrial respiration. A full

list of pathways is provided in Table S6.

1369

1370

1371

1372

Supplement 9. RSL3 treatment has no effect on the viability of cold-exposed *GPX4* KO

K562 cells.

a-d, Wild-type K562 cells **(a)** and three independent *GPX4* KO K562 clonal lines **(b-d)** were

treated with RSL3 (1 µM) and ferrostatin-1 (Fer-1, 1 µM) as indicated for 8 hours at 4°C (*n* = 4)

prior to trypan blue staining. Wild-type K562 cells show a significant decrease in cell viability

when treated with RSL3 compared to no treatment (*P = 0.0213), whereas *GPX4* KO lines show

no significant changes in viability (ns, *P* > 0.05). All values show mean ± SEM, with significance

measured by one-way ANOVA adjusted for multiple comparisons by Dunnett's test. *P < 0.05;

**P < 0.01; ***P < 0.001; ****P < 0.0001; ns *P* > 0.05.

1382

Supplement 10. Cell permeable glutathione and glutathione precursors do not increase cold cell viability

a, K562 cells were placed at 37°C and treated with cell permeable glutathione GSH-MEE (1 µM) and/or the glutathione synthesis inhibitor buthionine sulfoximine (BSO, 1 µM) for 1 day (n = 3). Treatment with BSO resulted in increased cell death (**P = 0.0018) that was rescued by GSH-MEE (ns, P = 0.9961). **b,** Treatment with GSH-MEE has no effect on K562 cold-induced death

after 3 days at 4°C as measured by trypan blue staining (n = 3, 5 µM; ns, P = 0.1558). **c,** K562

cells were placed at 37°C and treated with ferroptosis inducer Erastin (10 µM) and N-

acetylcysteine (NAC, 10 µM) for 1 day (n = 3). Treatment with Erastin resulted in increased cell

death (***P = 0.0006) that was rescued by NAC (ns, P = 0.1091). **d,** Treatment with N-

acetylcysteine does not increase cold cell viability in K562 cells after 2 days at 4°C as measured

by trypan blue staining (n = 3, 10 µM; *P = 0.0352). **e,** Treatment with L-Cystine does not

increase cold cell viability in K562 cells after 2 days at 4°C as measured by trypan blue staining

(n = 3, 100 µM; ns, P = 0.9843). All values show mean ± SEM, with significance measured by

one-way ANOVA adjusted for multiple comparisons by Dunnett's test. *P < 0.05; **P < 0.01;

***P < 0.001; ****P < 0.0001; ns P > 0.05.

1401

1402

1403

1404
1405
Supplement 11. Glutathione biosynthesis genes show increased depletion in cold-
exposed K562 cells
a,c, Log2 fold-change (log2FC) of 10 guides per targeted gene in BHK-21 genome-wide screen,
showing guide depletion over three cycles of cold exposure and rewarming. **a**, *Gclc*, **c**, *Gss*. **b**,
d, Log2 fold-change (log2FC) of 5 guides per targeted gene in K562 genome-wide screen,
showing significant guide depletion over three cycles of cold exposure. **b**, *GCLC*, **d**, *GSS*.
Significance between Cycle 1 versus Cycle 3 is measured by two-way ANOVA adjusted for
multiple comparisons by Dunnett's test. Significance between 37°C and 4°C for each cycle is
measured by two-way ANOVA adjusted for multiple comparisons by Bonferroni's test. *P < 0.05;
**P < 0.01; ***P < 0.001; ****P < 0.0001; ns P > 0.05.
1416