Intracellular lipopolysaccharide regulates ER remodeling upon bacterial infection

Authors

Yi-Lin Cheng[1,2,3,]*, João Mello-Vieira[1,2], Adriana Covarrubias-Pinto[1], Alexis Gonzalez[1], Santosh Kumar Kuncha[1,2,4], Chun Kew[1,2], Kaiyi Zhang[5], Muhammad Awais Afzal[6], Nour Diab[5], Sophia Borchert[7], Wenbo Chen[1,2], Timothy Chun Huang[3,8], Mathias Walter Hornef[5], Christian A. Hübner[6], Michael Hensel[7], Ivan Dikic[1,2,9,10,11,]*

[1]Institute of Biochemistry II, Faculty of Medicine, Goethe University, Frankfurt, Germany

[2]Buchmann Institute for Molecular Life Sciences, Goethe University, Frankfurt, Germany

[3]Institute of Basic Medical Sciences, College of Medicine, National Cheng Kung University, Tainan, Taiwan

[4]Altos Labs Inc, Bay Area Institute, Redwood City, CA, USA

[5]Institute of Medical Microbiology, RWTH University Hospital, Aachen, Germany

[6] Institute of Human Genetics, Jena University Hospital, Friedrich Schiller University, Jena, Germany

[7]Abt. Mikrobiologie, Universität Osnabrück, Osnabrück, Germany.

[8]School of Medicine, College of Medicine, National Cheng Kung University, Tainan, Taiwan

[9]Max Planck Institute of Biophysics, Frankfurt, Germany

[10]Fraunhofer Institute for Molecular Biology and Applied Ecology (IME),

[11]Lead contact

*Correspondence: dikic@biochem2.uni-frankfurt.de;

yilincheng@gs.ncku.edu.tw

 Graphical Abstract

In Brief

Cytosolic LPS binds to the ER-phagy receptor FAM134B, triggering ER fragmentation, and subsequently, the formation of FAM134B-positive vacuoles for bacterial survival.

Highlights

- Intracellular LPS induces FAM134B-mediated ER fragmentation.
- Lipid A of LPS binds to FAM134B.
- O-antigen of LPS triggers oligomerization of FAM134B and membrane fragmentation.
- FAM134B-positive *Salmonella*-containing vacuoles are not acidified, providing a niche for *Salmonella* survival.
- Outer membrane vesicles are required for cytosolic LPS delivery to activate FAM134B-mediated ER fragmentation.

SUMMARY

The remodeling of endoplasmic reticulum (ER) is finely regulated by selective autophagy, known as ER-phagy. However, whether Gram-negative bacteria, which are known to manipulate the host membranous system, regulate ER-phagy remains elusive. We find that bacterial lipopolysaccharide (LPS) released upon infection with *Salmonella* Typhimurium colocalizes with FAM134B, a reticulon-like protein that operates as an ER-phagy receptor. Pulldown assays reveal that LPS binds directly to FAM134B. The interaction occurs via lipid A of LPS and positively charged amino acid residues on the amphipathic helices and C-terminal region of FAM134B. Upon binding, O-Antigen of LPS promotes FAM134B oligomerization and ER fragmentation in *in vitro* liposome membrane remodeling assays. Delivery of LPS into the cytoplasm via transfection significantly increases the number of FAM134B–LC3B-decorated ER fragments, which are recruited to *Salmonella*-containing vacuoles. Both intracellular and extracellular bacteria use outer membrane vesicles (OMVs) to deliver LPS into the cytosol for FAM134B activation and ER fragmentation. Taken together, our results show that Gram-negative bacteria subvert the host ER remodeling system to create bacteria-containing vacuoles that provide a niche for bacterial survival.

INTRODUCTION

Multiple intracellular Gram-negative bacteria have been identified to manipulate host membrane system, including the endoplasmic reticulum (ER), which is the most abundant membranous system in the mammalian cells, to create a niche for bacteria survival (Escoll et al., 2016). However, whether there is a common component of Gram-negative bacteria participating in the regulation of ER membranes is unclear. The selective autophagy of the ER (ER-phagy) has recently been identified as the master mechanism for ER remodeling and turnover (Gubas and Dikic, 2022), enabling lysosomal degradation of different ER components, which are engulfed into autophagosomes through the interaction of LC3B with ER-phagy receptors (Chino and Mizushima, 2020; Gubas and Dikic, 2022). Eight mammalian ER-phagy receptors have been identified, including the FAM134 family (Khaminets et al., 2015; Kumar et al., 2021; Reggio et al., 2021), RTN3 (Grumati et al., 2017), Sec62 (Fumagalli et al., 2016), CCPG1 (Smith et al., 2018), ATL3 (Chen et al., 2019), TEX264 (An et al., 2019; Chino et al., 2019), and CALCOCO1(Nthiga et al., 2020), which facilitate the degradation of different ER subdomains and respond to different types of stress conditions. The FAM134 family and RTN3L, in particular, are able to induce significant ER remodeling. They contain two hairpin-like domains (transmembrane domains; TM) anchored into the lipid bilayer and connected by an amphipathic helix (AH) linker, as well as another carboxy-terminal AH, which is connected to the second TM. Together they form a so-

called reticulum homology domain (RHD), whose integrity appears critical for ER-
phagy (Bhaskara et al., 2019; Grumati *et al.*, 2017; Khaminets *et al.*, 2015). The RHD
of FAM134B, the best-studied ER-phagy receptor, can sense highly positive
membrane curvature, thereby triggering the formation of FAM134B clusters that
provide an additional force for ER membrane budding and scission (Bhaskara *et al.*,
2019; Gonzalez et al., 2023; Jiang et al., 2020; Siggel et al., 2021).

LPS is the main component of the outer membrane of Gram-negative bacteria. It
is composed of a conserved lipid A moiety, the inner core and outer core
oligosaccharides, and O-antigen (O-Ag), the outermost part composed of a variable
number of repeating sugar residues. In recent decades, most studies have focused on
the extracellular role of LPS, which is specifically sensed on the plasma membrane by
the Toll-like receptor 4 (TLR4)/myeloid differentiation factor 2 (MD2) co-receptor
complex, that acts in conjugation with CD14 to trigger signaling cascades for the
expression of inflammatory genes (Park et al., 2009). The release of LPS by
intracellular *Salmonella* to cytosolic vesicles was reported by Portillo and colleagues
as early as 1997, however the targeted proteins and downstream effects are still
elusive (Garcia-del Portillo et al., 1997). More recently, human caspase-4 (mouse
homologue caspase-11) and caspase-5 have been identified as innate immune
receptors that specifically recognize intracellular LPS (Shi et al., 2014). The lipid A
moiety of LPS directly binds to the amino (N)-terminal caspase-activation and
recruitment domain (CARD) of these caspases, triggering their oligomerization and
activation (Shi *et al.*, 2014). Activated caspase-4/-11 further cleave gasdermin D to
liberate its N-terminal domain, which has membrane-pore-forming activity at the
plasma membrane and thereby induces pyroptosis (Aglietti et al., 2016; Ding et al.,
2016). Moreover, LPS on cytosolic *Salmonella* is also recognized by guanylate-binding
protein (GBP) 1, which initiates the assembly of a GBP coat for caspase-4 recruitment
and activation to trigger non-canonical inflammasome signaling (Santos et al., 2020).

Not only intracellular Gram-negative bacteria – after their escape from vacuoles -
expose LPS to the cytosol, but also extracellular Gram-negative bacteria by having
their LPS-enriched outer membrane vesicles (OMVs) internalized into the cell (Vanaja
et al., 2016). OMVs bud from Gram-negative bacterial cell envelopes and have a
diameter of 20 nm to 250 nm. The bacterial envelop is composed of the cytoplasmic
membrane and the outer membrane, which enclose the periplasmic space containing
a peptidoglycan layer and periplasmic proteins that stabilize the membrane through
various crosslinks (Juodeikis and Carding, 2022; Schwechheimer and Kuehn, 2015;
Silhavy et al., 2010). Upon infection, environmental stress factors released by the host
disturb the stability of those crosslinks and increase OMV budding. OMVs are
internalized into the host cell via endocytosis, and LPS is released from early

endosomes into the cytosol, where it activates caspase-11 (Vanaja *et al.*, 2016) and
possibly exerts other effects.
Here we describe a novel function for intracellular LPS mediated by binding to
FAM134B ER-phagy receptor. Both *Salmonella* infection and LPS transfection induced
FAM134B-mediated ER fragmentation. *Salmonella* LPS colocalized with FAM134B as
determined by immunofluorescence experiments, and biochemical analysis showed
that lipid A of LPS directly bound to FAM134B via electrostatic forces. Although lipid A
was sufficient for FAM134B binding, oligomerization of FAM134B and the following
membrane fragmentation required the presence of O-Ag of LPS. A part of the
fragmented FAM134B-positive ER was recruited by bacteria to form the *Salmonella*-
containing vacuole. We also found that OMVs are responsible for the delivery of LPS
from both intracellular and extracellular bacteria into the cytosol. Taken together, this
study identified FAM134B as a new intracellular target of LPS, enabling ER membrane
remodeling and formation of the survival niche.
133
134 **RESULTS**
135
LPS colocalizes with FAM134B and triggers FAM134B-mediated ER
fragmentation
To investigate whether intracellular Gram-negative bacteria activate ER-phagy,
we analyzed the protein turnover of several ER-phagy receptors, including FAM134B,
CCPG1, Sec62, RTN3L, and TEX264, during *Salmonella enterica* serovar
Typhimurium SL1344 infection (Figure S1A). The expression of FAM134B increased
upon infection while there were no significant changes for other ER-phagy receptors
(Figures S1A). We observed a slight decrease in RTN3L but it did not colocalize with
LC3B after *Salmonella* infection (Figure S1B). This result implies that *Salmonella*
mainly affects FAM134B-mediated ER-phagy, which is usually associated with
transcriptional up-regulation of FAM134B (Chino and Mizushima, 2020). In HeLa TREx
FLAG-HA-FAM134B-inducible cells, *Salmonella* strongly induced colocalization of
FAM134B puncta with LC3B puncta, indicating induction of FAM134B-mediated ER
fragmentation upon infection (Figure 1A). To our surprise, we found LPS puncta
(arrowheads) colocalizing with FAM134B which was not associated with bacteria
(arrows), indicated by nucleic acid staining with 4',6-diamidino-2-phenylindole (DAPI),
either in wild-type (WT) FAM134B or LC3B-interacting region mutant (mtLIR)
FAM134B (Figures 1A and S1C). Moreover, endogenous FAM134B, also colocalized
with *Salmonella* LPS after infection (Figure S1D). Quantitative analysis by confocal
quantitative image cytometer CQ1 showed that 40% of FAM134B puncta colocalized
with LPS after *Salmonella* infection (Figures 1B and 1C). The number of

FAM134B/LC3B-positive puncta after *Salmonella* infection was significantly increased
in WT FAM134B-expressing HeLa cells, but not in mtLIR FAM134B-expressing HeLa
cells, indicating *Salmonella* infection caused induction of FAM134B-mediated ER-
phagy and ER fragmentation (Figures 1D and S1E). Co-immunoprecipitation (Co-IP)
assays revealed that *Salmonella* enhanced interaction of FAM134B with LC3B-II in the
presence or absence of Bafilomycin A1 (BafA1) (Figure 1E). Our previous study
defined ER fragments as puncta containing FAM134B, LC3B-II and ER proteins that
do not directly bind LC3B (Gonzalez *et al.*, 2023). To clarify whether LPS can trigger
FAM134B-mediated ER fragmentation, we transfected *Salmonella* LPS in HeLa cells
followed by immunostaining and confocal microscopy analysis. The results showed
that LPS colocalized with FAM134B puncta and the number of FAM134B/LC3B-
positive puncta increased in presence of LPS (Figures 1 F and 1G). Moreover,
transfection of *Salmonella* LPS also triggered FAM134B fragments delivered to
lysosomes, indicating by LAMP1 (Figure S1F). Thus, *Salmonella* LPS initiated
FAM134B-mediated ER fragmentation.

LPS binds to FAM134B via electrostatic force
Since LPS colocalized with FAM134B in the confocal images, we hypothesized
that LPS binds directly to FAM134B. Therefore, we carried out a series of streptavidin
pull down assays to clarify the mode of interaction (Figure 2A). We used biotin-
conjugated *E. coli* O111:B4 LPS immobilized on streptavidin sepharose to study its
interaction with either HA-FAM134B or HA-RTN3L (Figure 2B). We detected that biotin-
LPS strongly pulled down HA-FAM134B in a dose-dependent manner, but not HA-
RTN3L (Figure 2B). The binding of LPS to FAM134B was LIR motif-independent
(Figure S2A). Pulldown assays after transfection of biotin-LPS into cells expressing
FLAG-HA-FAM134B or FLAG-HA-RTN3L showed similar results (Figure S2B).
Importantly, upon *Salmonella* infection in cells expressing FLAG-HA-FAM134B, LPS
was also able to interact with FAM134B (Figure 2C). No significant interaction between
RTN3L and LPS was detected upon *Salmonella* infection (Figure 2C). To further clarify
which segment of LPS is required for binding to FAM134B, we performed a competition
assay. WT LPS (also called smooth (S)-LPS), rough LPS, including Ra-LPS (LPS
without O-Ag) and Re-LPS (LPS containing lipid A and three 2-Keto-3-deoxy-octonate
(KDO)), and lipid A, were preincubated with lysates containing FLAG-HA-FAM134B,
followed by incubation with biotin-LPS-immobilized streptavidin sepharose. We
analyzed whether the preincubation with different LPS variants can block the binding
of biotin-LPS to FAM134B (Figures 2D and 2E). The results reveal that all kinds of LPS
significantly reduced the binding ability of biotin-LPS to FAM134B, except Muramyl
dipeptide (MDP), which is another component of the bacteria cell wall, demonstrating
that lipid A is sufficient for FAM134B binding (Figure 2E). Furthermore, polymyxin B, a

cationic peptide antibiotic, which interacts with lipid A via ionic and hydrophobic forces,
inhibited the binding of biotin-LPS to FAM134B in a dose-dependent manner (Figure
2F).

Next, we characterized how FAM134B binds to LPS. Since lipid A contains
multiple negatively charged groups (Adams et al., 2014), we hypothesized that the
interaction might occur through electrostatic forces. We analyzed each domain of
FAM134B and found that two cytoplasmic amphipathic helices, the linker and C-
terminal region of the amphipathic helices (AHL and AHC), and the C-terminal region
(Cterm), are highly solvent-accessible and enriched with positively charged residues,
including arginine (R) and lysine (K) (Figure 2G). According to this analysis, we
mutated R or K residues to glutamic acid (E) in these three FAM134B regions (Figure
2H). We then evaluated the interactions of these mutants with biotin-LPS. Each mutant
FAM134B construct showed significantly reduced binding to biotin-LPS, and binding
was completely abolished when mutations in all three regions were combined (Figure
2I). Transfection of FAM134B TRIPLEmt (mutations in all three regions) robustly
blocked colocalization of FAM134B with LPS and LC3B (Figures 2J and 2K). These
results were also confirmed by co-immunoprecipitation experiments in which
FAM134B TRIPLEmt-HA co-precipitated significantly less LPS and LC3B-II compared
to FAM134B WT-HA (Figure 2L). Moreover, *in vitro* streptavidin pulldown assays
showed that biotin-LPS pulled down FAM134C and FAM134B with similar efficiency
whereas FAM134A interacted significantly weaker (Figure S2C). This result could be
explained by the fact that FAM134B and FAM134C have a high sequence similarity,
from which the sequence of FAM134A clearly deviates (Figure S2D-S2F).

FAM134B oligomerization is triggered by O-Ag of LPS
Oligomerization of FAM134B is reported to drive ER membrane scission followed
by ER-phagy induction (Jiang *et al.*, 2020). Especially, caspases also undergo
oligomerization after binding with LPS (Shi *et al.*, 2014). We therefore hypothesized
that LPS binds to FAM134B to induce its oligomerization. S-protein-FLAG-streptavidin
binding peptide (SFB)-FAM134B was expressed and purified from HEK293T cells
followed by incubation with S-LPS or lipid A. In order to analyze oligomeric species of
FAM134B, we performed blue native PAGE, which enforces migration of native protein
complexes or oligomers based solely on their molecular weight (Wittig et al., 2006),
followed by immunoblotting with FLAG. S-LPS significantly increased the presence of
FAM134B oligomers of approximately 720 kDa (Figure 3A). To our surprise, lipid A did
not show a similar effect, although it played a major role in binding to FAM134B,
suggesting that O-Ag is indispensable for the oligomerization of FAM134B. To test this
idea, we infected HeLa cells expressing SFB-FAM134B with an isogenic set of

Salmonella mutant strains with defined defects in the O-antigen biosynthesis, including
the *rfbP* (O-antigen transferase) and *rfaL* (O-antigen ligase) mutant strains lacking O-
Ag, the *rfaG* (glucosyltransferase I) mutant strain lacking O-Ag and outer core, as well
as the *rfaH* (transcriptional activator) mutant strain with reduced polymeric O-Ag
(Figures 3B and 3C) (Zenk et al., 2009). After infection, SFB-FAM134B was purified
and blue native PAGE analysis was performed. The results showed that FAM134B
oligomers in the range of ~720 kDa to ~1048 kDa were significantly reduced after
infection with any of the O-Ag-defective strains compared to WT and complemented
strains (Figure 3B).
FAM134B clustering would further induce membrane curvature and remodeling
(Bhaskara *et al.*, 2019). To further prove that O-Ag regulates membrane remodeling
driven by FAM134B, we conducted a liposome-remodeling assay in which incubation
with purified FAM134B drives membrane fragmentation (Bhaskara *et al.*, 2019). We
first determined the suitable lipid-to-protein ratio (LPR) for the examination of the effect
of LPS on FAM134B-triggered membrane fragmentation (Figures S3A and S3B). LPR
40:1 triggered strong liposome remodeling, indicated by a significant reduction of the
liposome diameter compared to the control group of liposomes only (Empty). LPR
100:1 had a moderate effect, resulting in a slightly reduced liposome size (Figures S3A
and S3B). Thus, we used LPR 100:1 to study the action of LPS (Figures 3D-3G). S-
LPS significantly enhanced the membrane remodeling effect of FAM134B compared
to liposomes incubated with FAM134B alone in a dose-dependent manner (Figures 3D
and 3E). Notably, LPS alone did not change the size of liposomes compared to the
control sample, demonstrating that LPS cannot fragment membrane by itself, but
enhances the action of FAM134B on membranes (Figures 3D and 3E). Compared to
S-LPS, Ra-LPS and Re-LPS, which lack the O-Ag, had less pronounced effects on the
size of liposomes, i.e. on the ability of FAM134B to induce membrane fragmentation
(Figures 3F and 3G). To further verify the role of O-Ag in ER fragmentation in cells, we
performed co-localization analysis of HA-FAM134B and LC3B after infection of WT, the
isogenic mutant strains without O-Ag, and the two complemented strains of ∆*rfaL* and
∆*rfaG* (Figures 3H and S3C). The number of FAM134B/LC3B-positive puncta was
significantly reduced upon infection with O-Ag mutant strains, ∆*rfbP,* ∆*rfaL,* ∆*rfaG,* and
∆*rfaH* strains*,* in comparison to the WT strain. On contrary, complemented strains of
∆*rfaL* and ∆*rfaG* induced a similar number of FAM134B/LC3B-positive puncta as the
WT strain (Figures 3H and S3C). The bacteria count per cell did not show significant
differences among the groups (Figure 3I and S3C). Compared to S-LPS, transfection
of Ra-LPS, Re-LPS, and lipid A, which are the LPS without O-Ag, did not increase the
number of HA-FAM134B/LC3B-positive puncta (Figures S3D and S3E). The other two
cell wall components of bacteria, Pam3CSK4 and MDP, did not affect puncta formation

(Figures S3D and S3E). These results demonstrated that O-Ag of LPS triggered
oligomerization of FAM134B and FAM134B-mediated membrane remodeling.
272

FAM134B is recruited to *Salmonella*-containing vacuoles and promoted bacteria
growth

To further investigate the fate and function of FAM134B after infection, we
measured the FAM134B-mediated ER-phagy flux by mCherry-EGFP-FAM134B WT or
LIR mutant-inducible HeLa cells after *Salmonella* infection for 2 hr and 4 hr (Figures
S4A and S4B). Owing to the instability of EGFP in acidified environment, only the red
fluorescence of mCherry would be detected when FAM134B is delivered to lysosomes
(Chino *et al.*, 2019; Jiang *et al.*, 2020; Khaminets *et al.*, 2015). *Salmonella* strongly
induced formation of red FAM134B puncta 4 hr post-infection in mCherry-eGFP-
FAM134B WT HeLa cells, but not in cells expressing the LIR mutant (Figures S4A and
S4B). Interestingly, FAM134B positive for both GFP and mCherry was recruited to
bacteria producing vacuoles (Figures S4A-1). In cells with highly proliferative bacteria,
which usually occur after bacteria escape from *Salmonella*-containing vacuoles (SCVs)
into cytosol, double-color FAM134B was also recruited to bacteria (Figure S4C). In
Salmonella-infected intestinal tissue sections of 5 days-old suckling mice, FAM134B
was also recruited to *Salmonella* (Figure S4D). These structures were positive for
calnexin and REEP5, as well (Figure S4E and S4F). Moreover, in FAM134B knockout
(KO) mouse bone-marrow-derived macrophages (BMDMs), ER-surrounded
Salmonella were significantly reduced compared to WT BMDMs (Figures S5A-S5C).
These results suggest that *Salmonella* sequesters FAM134B, possibly to recruit ER
membrane to the vacuole.

According to previous studies, invasive *Salmonella* can survive in the
endolysosomal membrane-derived SCVs or be cleared after sequestration in
xenophagosomes (LaRock et al., 2015; Madan et al., 2012; Perrin et al., 2004;
Ravenhill et al., 2019; Wild et al., 2011). We decided to study whether the FAM134B-
positive structures belong to one of these two compartments. We observed that
bacteria decorated with FAM134B-HA, colocalized with LC3B but not with the
endolysosomal marker LAMP1 (Figure 4A). Quantitative analysis indicated that around
50% LC3B-positive *Salmonella*-containing compartments were FAM134B-positive
after *Salmonella* infection for 2 hr, and around 40 % at 4 hr and 6 hr post-infection, but
very few LAMP1-positive SCVs were FAM134B-positive (Figure 4B). Indeed, almost
all FAM134B/LC3B-positive vacuoles were lysotracker-negative, indicating lack of
acidification of the compartment (Figures 4C and 4D). The xenophagosomes without
FAM134B expression showed to be either lysotracker-positive or -negative, which is
due to the different stages of autophagy flux (Figures 4C and 4D). Depletion of

FAM134B significantly increased the percentage of lysotracker-positive
xenophagosomes (Figures 4E and 4F). In the interactome of FAM134B during
Salmonella infection, we identified multiple bacterial factors, including outer membrane
proteins, slyB and ompA, STM1249, turfA, steA, as well as sopB, which was reported
to block lysosome fusion with SCVs (Bakowski et al., 2010) (Figures S6). Nevertheless,
whether these bacterial proteins contribute to FAM134B-mediated bacterial survival
requires further investigation. To confirm the role of FAM134B in bacterial growth, we
performed a plating assay to measure colony-forming units (CFU) in FAM134B-WT or
FAM134B-LIR mutant inducible cells after *Salmonella* infection. Induction of WT
FAM134B, but not LIR mutant FAM134B increased CFU compared to non-inducible
group (Figure 4G). Depletion of FAM134B either in FAM134B knockdown HeLa cells,
FAM134B KO mouse embryonic fibroblasts (MEFs), and FAM134B KO BMDMs
reduced *Salmonella* multiplication (Figures 4H, 4I, and S5C). These results indicate
that recruitment of FAM134B enables the formation of a compartment without
acidification leading to bacterial survival.
323

OMVs are required for LPS delivery to induce FAM134B-mediated ER remodeling
Transmission electron microscopy (TEM) studies revealed that LPS secreted by
Salmonella is presented on vesicular structures that resemble OMVs generated by
bacteria (Garcia-del Portillo *et al.*, 1997). OMVs, which are enriched in LPS on the
surface, are also responsible for delivery of LPS from extracellular bacteria to the host
cell cytosol via endocytosis (Vanaja *et al.*, 2016). Hence, we investigated whether OMV
production is required for FAM134B-mediated ER fragmentation. OMV production is
governed by various mechanisms that detach the outer membrane from the underlying
peptidoglycan and then allow outward budding to form vesicles (Juodeikis and Carding,
2022; Schwechheimer and Kuehn, 2015). For example, the outer membrane protein A
(OmpA) is the major outer membrane porin in *E. coli* and *S. enterica* and is linked to
peptidoglycan. Lack of OmpA results in decreased stability of the outer membrane and
increased release of OMVs, which has been previously reported for *S. enterica*
serovars Typhimurium (Deatherage et al., 2009) and Typhi (Nevermann et al., 2019),
and was confirmed for the mutant strains used in our study by ultrastructural analyses
(Figure S7). Another example is TolAB, which encodes components of the multi-
subunit Tol-Pal complex that controls the constriction of the cell envelope during cell
division. Mutant strains lacking TolAB show increased OMV production due to defects
in partitioning the outer membrane during cell elongation and septation (Szczepaniak
et al., 2020). DegS is also involved in OMV production. It is part of a sensor system
responding to periplasmic stress, which is, for example, induced by accumulation of
misfolded secretory proteins. If the stress response system is absent, the accumulation

of misfolded proteins in the periplasm induces increased OMV production to remove toxic aggregates (Nevermann *et al.*, 2019). On the contrary, PagL is a deacylase that removes one lipid from hexa- or hepta-acylated lipid A of LPS (Kawasaki et al., 2004). This modification alters topology of LPS in the outer membrane, resulting in increased membrane curvature and OMV release (Elhenawy et al., 2016). We therefore established *Salmonella* strains mutated in the genes *pagL*, *ompA*, *tolAB*, and *degS,* and obtained the images of OMVs production by scanning electron microscopy (SEM) and TEM followed by measurement of OMVs amount and size distribution (Figure S7). Compared to the WT strain, the ∆*ompA*, ∆*tolAB*, and ∆*degS* strains showed increased production of OMVs. Conversely, no vesicles could be detected with the ∆*pagL* strain (Figure S7C), which is in line with previous findings showing that deletion of *pagL* results in decreased generation of OMVs by intracellular *Salmonella* (Elhenawy *et al.*, 2016).

Mutation of *ompA*, *tolAB*, and *degS*, associated with increased OMVs production, augmented the number of FAM134B/LC3B-positive puncta to varying degrees (Figures 5A, 5B, and S7D). Especially, mutation of *tolAB* strongly increased puncta formation compared to the WT group. Conversely, the number of FAM134B/LC3B-positive puncta decreased when *pagL* was mutated (Figures 5A and S7D). The LPS levels in the cell lysates and the amount of LPS interacting with FAM134B positively correlated with the number of FAM134B/LC3B-positive puncta (Figures 5A and 5C). The bacterial counts for the different isogenic mutant strains did not show significant change at 2 hr post-infection (Figure 5B). To further explore whether OMVs can directly activate FAM134B-mediated ER fragmentation, we purified OMVs from *Salmonella* and treated HeLa TREx FLAG-HA-FAM134B cells with these OMVs. After treatment with *Salmonella* OMVs, we observed colocalization of LPS and HA-FAM134B. The number of FAM134B/LC3B-positive puncta also increased (Figures 5D and 5E). To differentiate whether FAM134B mainly bound to shed LPS or the LPS on OMVs, we stained OMVs by Vybrant DiO dye after purification and then treated the cells. Confocal microscopy showed that FAM134B/LPS-positive puncta did not colocalize with Vybrant DiO-positive OMVs (Figure S8A). Furthermore, the O-Ag mutant strains-derived OMVs lost the ability to trigger FAM134B-mediated ER fragmentation (Figure 5F). We further tested whether extracellular bacteria, more specifically Enteropathogenic *Escherichia coli* (EPEC), can induce a similar phenomenon. In addition to direct exposure, transwells with 0.45 μm polycarbonate membranes were used to prevent bacteria passage but allow OMVs transition. The results reveal that EPEC induced similar amount of FAM134B/LC3B-positive puncta in direct infection and transwell system (Figures 5G and 5H). The *E. coli* strain without O-Ag, BL21, was used as the negative control, showing no significant formation of FAM134B puncta (Figures 5I and S8B).

384   The number of LC3B-decorated FAM134B puncta also increased in the cells with
385   EPEC-derived OMVs, which were pre-stained by Vybrant DiO after purification.
386   However, BL21-derived OMVs did not cause same effect (Figures 5I and S8B). These
387   results suggest that intracellular and also extracellular bacteria utilize OMVs for the
388   delivery of LPS to activate FAM134B-mediated ER fragmentation.
389
DISCUSSION
Here we report that FAM134B, an ER-phagy receptor, interacts with LPS to induce
ER membrane fragmentation and to form vacuoles that surround *Salmonella* for
bacterial survival. The interaction between LPS and FAM134B was mainly mediated
by lipid A and positively charged amino acid residues in FAM134B. Besides, O-Ag of
LPS was responsible for oligomerization of FAM134B and the subsequent membrane
fragmentation, although it remains to be investigated how the molecular dynamics are
regulated in detail.
We have previously shown that the wedge-shaped membrane inclusions formed
by the RHD of FAM134B trigger membrane curvature followed by membrane
fragmentation (Bhaskara *et al.*, 2019). Our recent study (Gonzalez *et al.*, 2023)
uncovered that ubiquitination of the RHD domain finely regulates its membrane
remodeling ability, promotes large-scale membrane deformation, increases receptor
cluster size, and enhances ER-phagy flux. In the present study, we found that
mutations of multiple lysine residues of FAM134B, which are also sites for
ubiquitination, significantly inhibited the binding ability to LPS (Figure 2). Thus, LPS
might play a similar role as ubiquitination for FAM134B oligomer formation and
membrane fragmentation.
The fact that LPS can directly bind to FAM134B to induce ER membrane
fragmentation (Figure 3) and ER-phagy supports a model in which *Salmonella* hijacks
ER-phagy and prevents FAM134B-positive autophagosomes from traveling to
lysosomes. SopB, a *Salmonella* virulence factor that we found to bind to FAM134B,
could possibly play a role in this scenario as it inhibits autophagosome-lysosome
fusion so that the FAM134B-positive vacuoles cannot be acidified, allowing the
bacteria to multiply (Figures 4 and S6) (Bakowski *et al.*, 2010). However, this needs
to be investigated further. Based on the knockdown assay, we speculate that without
FAM134B, bacteria cannot sequester enough ER membrane, resulting in a vacuole
that is more susceptible to xenophagy. Recently, Gatica *et al.* showed that *Salmonella*
SopF inhibits amino acid starvation-induced FAM134B-mediated ER-phagy in
HEK293T cells, suggesting *Salmonella* effectors may play controversial roles in
different conditions. Nevertheless, this work did not test the role of SopF in
Salmonella or LPS-induced ER-phagy (Gatica et al., 2024).

In addition to vacuole formation, this mechanism may also serve other purposes,
since ER remodeling participates in different physiological functions, such as vesicular
trafficking, the unfolded protein response, ER-associated degradation, ER tubule
formation, ER-lysosome associated degradation, and ER-organelle contact-site
formation (Gubas and Dikic, 2022). ER shaping and remodeling enables human cells
to deal with ER stress, which is triggered by live Gram-positive bacteria through the
innate sensor stimulator of interferon genes (STING), while simultaneously inducing
autophagy (Moretti et al., 2017). Upon ER stress induction, ER proteins were found to
co-fractionate with LC3B, suggesting that ER-phagy is activated to balance ER stress
and maintain homeostasis (Moretti *et al.*, 2017). It will be interesting to further
investigate whether FAM134B-mediated ER-remodeling is involved in other
pathogenic or immune responses upon Gram-negative bacterial infection.

Our immunofluorescence experiments showed that the FAM134B puncta mainly
colocalized with LPS that was not localized on OMVs (Figure S8A). Although we cannot
yet draw any conclusions about the properties of the LPS released from OMVs, studies
on the binding between LPS and guanylate binding protein-1 (GBP1) provide us with
some clues for future investigations (Kutsch et al., 2020). GBP1 can bind directly to
lipid A moieties of LPS micelles, which form in aqueous solution (Santos et al., 2003),
or to LPS on the bacterial surface. It then undergoes polymerization, which is mediated
by the O-Ag (Kutsch *et al.*, 2020). Similarly, we found that binding of FAM134B to LPS
was mediated by lipid A moieties and O-Ag was required for oligomerization of
FAM134B. Moreover, FAM134B did not only bind to secreted LPS, which might be
present in micellar form, but was also recruited to bacteria, which mirrors the finding
for GBP1. Furthermore, we observed that LPS puncta appeared at a quite early stage
of infection, when only few bacteria were present. How these bacteria release LPS is
still unclear. Nevertheless, several studies have shown that GBPs and the immunity-
related GTPase family member B10 (IRGB10) may play a crucial role in the release of
LPS into the cytosol. GBPs were shown to be recruited to intracellular pathogens and
lyse the pathogen-containing vacuoles, resulting in the release of bacteria into the
cytosol and LPS being detectable therein (Meunier et al., 2014). GBPs further recruit
IRGB10 to the bacterial surface, disrupting the bacterial cell membrane and releasing
bacterial ligands, including LPS (Man et al., 2016). Although we found in this study that
the ability of different *Salmonella* isogenic mutant strains to produce OMVs correlates
with intracellular LPS levels (Figures 5A and 5D), the mechanism for release of LPS
from the intracellular OMVs remains unknown.

In summary, our work reveals a novel LPS-binding protein, FAM134B, which
undergoes oligomerization after binding to LPS, thereby triggering ER fragmentation.
The fragmented ER membranes are recruited to *Salmonella* to form the niche for

bacterial survival. Overall, this study provided new mechanistic insights into the pathways by which Gram-negative bacteria induce ER-remodeling and establish its proliferative niche.

Acknowledgments

We thank Prof. Dr. Volkhard A. J. Kempf at the Institute for Medical Microbiology and Infection Control, Hospital of the Goethe University for the kind gift of the EPEC O127:H6 strain. We thank the staff at the Quantitative Proteomics Unit at Goethe University for support with mass spectrometry and the Frankfurt Center for Advanced Microscopy (FCAM) for access to microscopes. This research is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), project number 259130777-SFB 1177, the European Union (ERC, ER-REMODEL, 101055213), the HMWK-funded cluster grant ENABLE, and grants from Else Kroener Fresenius Stiftung and Dr. Rolf M. Schwiete Stiftung to ID. M.H. was supported by the DFG by project P8 in SFB1557 and HE1964/23-1 in priority program SPP 2225 'Exit Strategies'. C.A.H. was supported by the DFG HU 800/14-1. Y.L.C. was supported by funding through a National Science and Technology Council, project NSTC 112-2320-B-006-012-MY2 and project 108-2917-I-564-021.

Author contribution

Conceptualization, Y.L.C. and I.D.; Methodology, Y.L.C., S.K.K., K.C., A.C-P, A.G., C.K., J.M.V., M.A.A., S.B., M.W.H.,C.A.H., M.H., N.D., and W.C.; Investigation, Y.L.C., S.K.K., A.C-P, A.G., J.M.V., M.A.A., C.K. and K.C.; Writing – Y.L.C., S.K.K., K.C., A.C-P, A.G., C.K., M.W.H., M.H., T.C.W., C.A.H. and I.D.; Funding Acquisition, I.D.; Resources, I.D., M.W.H., C.A.H. and M.H.; Supervision, I.D.

Declaration of interests

The authors declare no competing interests.

Supplemental information

Document S1. Figures S1–S8

FIGURE LEGENDS

Figure 1. *Salmonella* LPS activates FAM134B-mediated ER fragmentation.

(A) Immunofluorescence staining of HA, *Salmonella* LPS, and LC3B in HeLa TREx inducible cell lines expressing FLAG-HA-FAM134B that were infected with SL1344 for

2 hr. DAPI staining was used for cell nuclear and bacteria DNA staining. Scale bar, 10 μm.

(B and C) Quantification of HA and *Salmonella* LPS staining in HeLa TREx inducible cell lines expressing FLAG-HA-FAM134B infected with SL1344 for 2 hr. The number of LPS-positive HA-FAM134B puncta per cell (B) and the percentage of LPS-positive HA-FAM134B puncta in all HA-FAM134B puncta (C) were quantified using CellPathfinder software.

(D) Number of LC3B-positive HA-FAM134B puncta in HeLa TREx inducible cell lines expressing wild type (WT) or LIR mutant (mtLIR) FLAG-HA-FAM134B infected with SL1344 for 2 hr. Representative images are shown in Figure S1C.

(E) Co-IP of endogenous LC3B-II with HA in HeLa TREx inducible cells without doxycycline (Dox) treatment or with FLAG-HA-FAM134B WT or FLAG-HA-FAM134B mtLIR after Dox treatment. The cells were infected with SL1344 for 2 hr and pretreated with 200 ng/ml Bafilomycin A1 (BafA1) for 3 hr. GAPDH was used as an internal control of input.

(F and G) Immunofluorescence staining and quantification of HA, *Salmonella* LPS, and LC3B in HeLa TREx inducible cell lines expressing FLAG-HA-FAM134B transfected with 1 μg/ml *Salmonella* LPS for 18 hr by lipofectamine 2000. Lipo, lipofectamine only. (F) Arrows indicate the puncta with colocalization of HA-FAM134B and *Salmonella* LPS, and arrowheads indicate the puncta with colocalization of HA-FAM134B and LC3B. Scale bar, 10 μm. (G) Quantification of LC3B-positive HA-FAM134B puncta per cell.

(B and C) Data presented as mean + standard error of the mean (sem), n=3, > 400 cells. Differences were statistically analyzed by an unpaired two-tailed Student's t-test

(D and G) Solid bars of boxes indicate the medians. Boxes represent interquartile range from 25th to 75th percentile, and whiskers indicates 10th to 90th percentile. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test. Data were collected 60 cells from three independent biological replicates.

**p < 0.01, ***p < 0.001. ns, not significant.

Figure 2. LPS bind to FAM134B

(A) Schematic representation of streptavidin pulldown assays.

(B) Streptavidin pulldown assays that determine the binding of the indicated amounts of biotin-conjugated *E. coli* O111:B4 LPS and HA-tagged FAM134B or RTN3L in FLAG-HA-FAM134B- or FLAG-HA-RTN3L-expressing HeLa TREx cell lysates. GAPDH was used as an internal control of input.

(C) Co-IP of LPS with HA in HeLa TREx inducible cell lines without Dox treatment or expressing FLAG-HA-FAM134B or FLAG-HA-RTN3L by Dox treatment, which were infected with SL1344 for 2 hr. GAPDH was used as an internal control of input.

(D) Schematic representation of the LPS structure and the segments of different mutant variants of LPS, including lipid A, Re-LPS, Ra-LPS, and smooth (S)-LPS. P, phosphate; KDO, 3-deoxy-D-*manno*-octulosonic acid; Hep, L-*glycero*-D-*manno*-heptose; PEtn, phosphatidyl-ethanolamine; Glc, D-glucose; Gal, D-galactose; GlcNac, N-acetyl-D-glucosamine.

(E) Unlabeled S-LPS, Ra-LPS, Re-LPS, lipid A, and muramyl dipeptide (MDP) were used for competition with the binding of biotin-conjugated *E. coli* O111:B4 LPS to HA-tagged FAM134B in Dox-induced FLAG-HA-FAM134B HeLa TREx cell lysates.

(F) Indicated concentrations of polymyxin B (Poly B) were used to inhibit the binding of biotin-conjugated *E. coli* O111:B4 LPS to HA-tagged FAM134B in Dox-induced FLAG-HA-FAM134B HeLa TREx cell lysates.

(G) Schematic representation of the full length FAM134B sequence, which is composed of a reticulum homology domain (RHD), N-terminal (Nter, grey) and C-terminal regions (Cter, grey). LC3-interacting region consisting of 4 amino acids (LIR, pink) is located in the C-terminal region. The RHD consists of two transmembrane segments TM12 and TM34 (green), which are connected by a 60-residue linker, and two additional terminal segments. The C-terminal fragment of the RHD (AHC, yellow) and the linker-helix (AHL, yellow) form conserved amphipathic helices. The lower table shows the number of positively charged residues (arginine, R; Lysine, K) and the percentage of solvent accessible in every segment of FAM134B.

(H) Substitution of amino acids (R or K to glutamic acid, E) in segment AHL (AHLmt), AHC (AHCmt), and C-terminal (CTERmt). DOUBLEmt 1, DOUBLEmt 2, and TRIPLEmt indicate combination as shown in the table.

(I) Streptavidin pulldown assays that determine the binding of the indicated amount of

biotin-conjugated *E. coli* O111:B4 LPS to HA-tagged wild type or different mutant

FAM134B transfected in HEK293T cells. GAPDH was used as an internal control of

input.

(J and K) Immunofluorescence staining and quantification of HA and *Salmonella* LPS,

and LC3B in FAM134B WT-HA or FAM134B TRIPLEmt-HA transfected HeLa cells

infected with SL1344 for 2 hr. Scale bar, 10 μm. (J) Quantification of number of LC3B-

positive HA-FAM134B puncta per cell (K). Solid bars of boxes indicate the medians.

Boxes represent interquartile range from 25th to 75th percentile, and whiskers indicates

10th to 90th percentile. Differences were statistically analyzed by one-way ANOVA and

Tukey's multiple comparison test. ***$p < 0.001$. ns, not significant. Data were collected

60 cells from three independent biological replicates.

(L) Co-IP of LPS and endogenous LC3B-II in FAM134B WT-HA or FAM134B

TRIPLEmt-HA transfected HeLa cells, which were infected with SL1344 for 2 hr.

GAPDH was used as an internal control of input. The bands at ~ 22kD.

Figure 3. O-Ag of LPS induces FAM134B oligomerization and membrane

fragmentation

(A) Size analysis of S protein-FLAG-Streptavidin binding peptide (SFB)-FAM134B

oligomers after incubation with indicated amount of *Salmonella* S-LPS or lipid A by

blue native page and immunoblotting of FLAG.

(B) Size analysis of SFB-FAM134B oligomers purified from cell lysates infected with

NCTC12023 WT or O-Ag mutant (Δ) and complemented (compl.) strains. Upper panel

shows lower exposure (Low exp) blot and lower blot shows higher exposure (high exp)

blot. GAPDH was used as an internal control of input.

(C) Schematic representation of LPS structure of different O-Ag mutant strains,

including the strains mutant for *rfbP, rfaL*, *rfaG,* and *rfaH.*

(D-E) Representative images (D) and quantitative results (E) of remodeled

proteoliposomes (scale bars, 100 nm) taken from negative stain Transmission Electron

Microscopy (nsTEM). Empty liposomes (Empty) were added FAM134B-GST

recombinant protein at a lipid to protein ratio (LPR) 100:1 without (LPR 100:1) or with

0.5 mg/ml LPS (LPR 100:1+LPS 0.5) or 1 mg/ml LPS (LPR 100:1+ LPS 1). Incubation

of liposome with 1 mg/ml LPS was taken as negative control to check the effect of LPS
on liposome. The diameter of the representative proteoliposome with average size for
each group showed by dotted line or arrow.

(F-G) Representative images (F) and quantitative results (E) of different LPS, including
1 mg/ml of S-LPS, Ra-LPS, and Re-LPS, remodeled proteoliposomes (LPR 100:1,
scale bars, 100 nm). The diameter of the representative proteoliposome with average
size for each group showed by dotted line or arrow.

(E and G) The size distributions (n=300 each) from nsTEM images were measured
and showed by violin plots, showing a central boxplot (black lines indicate median with
interquartile range) with mirrored histograms on both sides (G).

(H and I) Immunofluorescence of HA, *Salmonella* LPS, and LC3B in HeLa TREx
inducible cell line expressing FLAG-HA-FAM134B infected with NCTC12023 WT or
LPS O-Ag mutant (Δ) and complemented (compl.) strains for 2 hr (H). DAPI staining
was used for cell nuclear and bacterial DNA staining. Scale bar, 10 μm. Quantification
of LC3B-positive HA-FAM134B puncta per cell (I) and *Salmonella* count per cell (J).
Data were collected 60 cells from three independent biological replicates.
Representative images are shown in Figure S3C.

(E, G, H, I) Differences were statistically analyzed by one-way ANOVA and Tukey's
multiple comparison test. *$p<0.05$, **$p<0.01$, ***$p < 0.001$, ns not significant.
606
Figure 4. FAM134B-positive *Salmonella*-containing vacuoles are unacidified,
promoting *Salmonella* survival

(A-B) Immunofluorescence staining of HA, LC3B, and LAMP1, indicating SCV, in HeLa
TREx inducible cell line expressing FLAG-HA-FAM134B infected with SL1344 for 4 hr
(A). DAPI was used for cell nuclear and bacterial DNA staining. Scale bar, 10 μm. The
percentage of FAM134B and LC3B double-positive *Salmonella* in all LC3B-positive
Salmonella, as well as the percentage of FAM134B and LAMP1 double-positive
Salmonella in all LAMP1-positive *Salmonella* was quantified using CellPathfinder
software (B). Data presented as mean + sem, n=3, > 400 cells.

(C-D) Immunofluorescence staining of HA and LC3B, and lysotracker staining in HeLa
TREx inducible cell line expressing FLAG-HA-FAM134B infected with SL1344 for 4 hr
(C). DAPI staining was used for cell nuclear and bacterial DNA staining. Scale bar, 10
μm. The percentages of lysotracker-positive or negative *Salmonella* to the indicating

markers *Salmonella* were quantified (D). Data were collected 60 cells from three independent biological replicates.

(E-F) Immunofluorescence staining of LC3B and lysotracker staining in HeLa cells transfected with control siRNA or FAM134B siRNA infected with SL1344 for 4 hr (E). DAPI was used for cell nuclear and bacterial DNA staining. Scale bar, 10 μm. The percentage of lysotracker-positive *Salmonella* to LC3B-positive *Salmonella* (F)*.* Data were collected 60 cells from three independent biological replicates.

(G) Colony-forming assay was performed to quantify the bacteria count in HeLa TREx inducible cells expressing FLAG-HA-FAM134B WT or FLAG-HA-FAM134B mtLIR infected with SL1344 for 1 hr and 8 hr. The fold increase of bacteria multiplication was calculated by normalizing bacteria count at 8 hr to 1 hr post-infection.

(H-I) Colony-forming assay was performed to quantify the bacteria count in HeLa cell transfected with control siRNA or FAM134B siRNA (H) or FAM134B WT MEF and FAM134B KO MEF infected with SL1344 100 moi at the indicated time points.

(B, D, F, G) Data represented as mean + sem. Differences were statistically analyzed by unpaired t-test (F and G) or by two-way ANOVA (H and I). **$p<0.01$, ***$p < 0.001$, ns, not significant.

Figure 5. LPS for FAM134B-mediated fragmentation is delivered by OMV

 (A-B) Immunofluorescence staining of HA, *Salmonella* LPS, and LC3B in cells infected with different mutant strains, including Δ*ompA*, Δ*tolAB*, and Δ*degS*, and Δ*pagL* strains. DAPI was used for cell nuclear and bacterial DNA staining. Scale bar, 10 μm. Quantification of LC3B-positive HA-FAM134B puncta (A) and *Salmonella* count per cell (B). Representative images are shown in Figure S7D.

(C) Co-IP of LPS with HA in cells infected with NCTC12023 WT or indicated mutant strains for 2 hr. GAPDH was used as an internal control of input.

(D-E) Immunofluorescence staining of HA, *Salmonella* LPS, and LC3B in cells treated with purified SL1344 OMVs for 6 hr (D). DAPI was used for cell nuclear DNA staining. Scale bar, 10 μm. Quantification of LC3B-positive HA-FAM134B puncta cell per cell (E).

(F) Immunofluorescence staining of HA and LC3B in cells treated with purified OMVs form NCTC12023 WT or indicated mutant strains for 6 hr. DAPI was used for cell nuclear DNA staining. Scale bar, 10 μm. Quantification of LC3B-positive HA-FAM134B puncta per cell (F).

(G-H) Immunofluorescence staining of HA and LC3B in cells infected by EPEC or BL21 directly or with 0.45 μm transwell for 4 hr (G). DAPI was used for cell nuclear DNA staining. Scale bar, 10 μm. Quantification of LC3B-positive HA-FAM134B puncta cell per cell (H).

(I) Immunofluorescence staining of HA and LC3B in cells treated with purified OMVs form EPEC or BL21 for 6 hr. Quantification of LC3B-positive HA-FAM134B puncta cell per cell with OMVs, stained by DiO Vybrant. Representative images are shown in Figure S8B.

HeLa TREx inducible cell lines expressing FLAG-HA-FAM134B was used for all experiments in Fig. 5. (A, B, E, F, H, and I) Solid bars of boxes indicate the medians. Boxes represent interquartile range from 25^{th} to 75^{th} percentile, and whiskers indicates 10^{th} to 90^{th} percentile. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test. ***$p < 0.001$, ns, not significant. Data were collected 60 cells from two (A, B and H) three independent (E, F and I) biological replicates.

STAR METHODS
LEAD CONTACT AND MATERIALS AVAILABILITY
Further information and requests for reagents may be directed to and will be fulfilled by the Lead Contact Ivan Dikic (ivan.dikic@biochem2.de). All unique/ stable reagents generated in this study are available from the LEAD Contact with a completed materials transfer agreement (MTA).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mammalian cell culture

HeLa and HEK293T cells were obtained from American Type Culture Collection (ATCC). WT and Fam 134b KO MEFs and BMDMs were isolated from respective mice genotypes. Animal experiments are approved by the Thüringer Landesamt für Lebensmittelsicherheit und Verbraucherschutz (TLV) (license number UKJ 22-019). MEFs were isolated and cultured from E13.5 stage embryos and SV40 large T antigen plasmid was used to immortalize MEFs. To culture BMDMs, femur and tibia bones

were isolated from 12-16 week old WT and Fam134b KO mice after euthanasia. The
bones were cleaned, sterilized, and the bone marrow was flushed out with sterile PBS.
It was centrifuged and re-suspended with BMDM culture media (DMEM supplemented
with 10% fetal bovine serum, 1% penicillin/streptomycin and 20 ng/mL of murine
macrophage colony-stimulating factor), and plated in petri dishes. Fresh media was
added on day 2. By day 5, differentiated macrophages were attached and the debris
were removed, the cells were washed with 1x PBS, collected, counted, and plated for
the experiments. Flp-In HeLa TREx cell lines (Invitrogen) were used to generate
inducible cell lines by cloning the cDNAs of FAM134B, FAM134B mtLIR (DDFELLD
LIR substituted by seven alanine residues), and RTN3L into MSCV iTAP N-FLAG-HA
retroviral vector. Then, 2 μg/ml puromycin and 15 μg/ml blasticidin were treated for
selection and maintenance of stable cells. Cells were cultured in Dulbecco's Modified
Eagle's Medium (DMEM) supplemented with 10% fetal bovine serum, 1%
penicillin/streptomycin at 37°C with 5% CO_2. BMDMs were additionally supplemented
with 20 ng/mL of murine macrophage colony-stimulating factor (Miltenyi BioTech).
701
702 Microbe strains
703
704 *S.* Typhimurium strains used in this study. The following isogenic mutant and
705 complemented NCTC12023 strains were established by Prof. Dr. Michael Hensel
706 (Zenk *et al.*, 2009).
707
1. SL1344 (WT)
2. NCTC12023 (WT)
3. NCTC12023 strains with deletion on *ompA, tolAB, degS* or *pagL.*
Mutant strains of *S. enterica* serovar Typhimurium strain NCTC12023 were
generated by λ Red-mediated homologous recombination basically as described
before (Datsenko and Wanner, 2000), using pWRG730 as helper plasmid for
controlled expression of Red functions (Hoffmann et al., 2017). The 60-mer
oligonucleotide sets as listed in Table X were used to amplify the *aph* targeting
cassette using pKD13 (for *tolA*, *degS*, or *ompA*) or pKD4 (for *pagL*) as template DNA.
Proper insertion of *aph* cassettes into target genes was confirmed by colony PCR
using gene-specific check primers flanking deletions, and universal K1 Red Del.
Confirmed mutated alleles were moved into fresh strain background using P22
transduction. The effects of mutations on outer membrane integrity were analysed
by SEM and nsTEM of strains grown overnight on LB agar (Fig. S7)
4. NCTC12023 strains with deletion on *rfbP, rfaL, rfaG,* or *rfaH*.
Mutant of *rfbP* (O-Ag transferase; NP_461027; Δ*rfbP*) and *rfaL* (O-Ag ligase;

NP_462613; Δ*rfaL*) are defective in the addition of the O-Ag to the outer core. Mutant of *rfaG* (glucosyltransferase I; NP_462622; Δ*rfaG*) caused outer core cannot be added to inner core. Mutant of *rfaH* (transcriptional activator; NP_ 462862; Δ*rfaH*), which is the gene regulates the stability of the long mRNA for LPS biosynthesis genes, also reduced polymeric O-Ag.

5. NCTC12023 strains with complementation of deletion of *rfaL* and *rfaG*

E. coli strains used in this study.

1. NEB Turbo

2. EPEC O127:H6 (A kind gift from Prof. Dr. Volkhard A. J. Kempf)

3. BL21

Chemically competent NEB Turbo cells were used for plasmid amplification.

METHOD DETAILS

Bacterial infection

S. Typhimurium and *E. coli* were cultured overnight and then diluted at ratio of 1 to 33 in fresh LB broth. The diluted cultures were then grown at 37°C for 3 h. For infection, the resuspended bacteria were adjusted to OD600 of 1 and added to HeLa cells or MEFs at a MOI of 100, and BMDMs at a MOI of 10 without antibiotics. After 30 min of infection, media were replaced by medium with 100 μg/ml gentamicin to remove extracellular bacteria. Samples were collected at the indicated time points after infection. For colony forming assay, which was performed as previously described (Steele-Mortimer, 2008), after 10 min of *S.* Typhimurium infection, the cells were washed by PBS twice to removed extracellular bacteria and incubated in antibiotics-free DMEM for 20 min. Then, the media were changed to DMEM containing 50 μg/ml gentamicin for 40 min and replaced by DMEM containing 5 μg/ml gentamicin. The cells were lysed with lysis buffer (1% Triton X-100 and 0.1% SDS in PBS) at indicated time points and serial diluted by PBS and plated onto LB plates. After 24 h incubation at 37°C, the colonies were calculated.

Immunofluorescence staining and confocal imaging

Cells which seeded in the 6-well plates with cover slips fixed by 4% paraformaldehyde in PBS for 15 min at room temperature. The buffer with 0.1% saponin, 1% bovine serum albumin (BSA), 0.05% NaN3 in PBS was used for permeabilization and blocking for 1 h at room temperature. Samples were then probed

by indicated primary antibodies as well as Alexa 488, Alexa 546, and Alexa 633
fluorescently labelled secondary antibodies (Invitrogen). After DAPI staining for
nucleus and bacteria DNA, the samples were fixed by mounting medium. Confocal
imaging was done using a Zeiss LSM 780 confocal microscope and analyzed by Zeiss
Zen (black edition) software. For high-throughput confocal imaging, cells were seeded
in the 96-well black plates with glass bottom followed by immunofluorescence staining
and imaging by CellVoyager CQ1 (Yokogawa) system and quantified using
CellPathfinder software.

Immunoprecipitation and western blotting

For immunoprecipitation, cells were lysed by the lysis buffer containing 50 mM
Tris-HCl (pH 7.5),150 mM NaCl, 5 mM EDTA, and 1% NP-40 with protease inhibitor
cocktail (Sigma Aldrich). Lysates were incubated with 15 µl anti-HA agarose beads
(SantaCruz Biotechnology) overnight at 4°C with end-to-end rotation. Beads were
washed three times with wash buffer containing 50 mM Tris-HCl (pH 7.5), 400 mM
NaCl, 5 mM EDTA). Proteins were eluted with 2X loading dye by boiling for 10 min
followed by western blotting. For western blotting, cells were lysed by the lysis buffer
containing 50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 5 mM EDTA, and 1% Triton X-100
with protease inhibitor cocktail (Sigma Aldrich). After protein quantification, samples
were boiled with loading dyes for 10 min. Samples were separated by running SDS-
PAGE with Tris-Glycine gels followed by transferred to PVDF membranes, which were
then blocked with 5% milk in TBST. After blocking, the membranes were sequentially
hybridized with primary and HRP (horseradish peroxidase) conjugated secondary
antibodies. the chemiluminescence method was performed to detect proteins
expression.

Transfection of siRNA
Cells were seeded in the 6-well plates on a day before experiments. FAM134B
siRNA and Scramble siRNA (Eurofins) were transfection by lipofectamine RNAiMAX
(Invitrogen) as manufacturer's instruction. After 48 hr culture, cells were reseeded for
the following experiments.

Streptavidin pulldown assay

To immobilize LPS onto streptavidin agarose beads, the indicated amount of
biotin-conjugated *E. coli* O111:B4 LPS (InvivoGen) were added to 10 µl of streptavidin
agarose beads and incubated at 4°C with rotation for 1 hr. The beads were then

washed 3 times with the lysis buffer containing 50 mM Tris-HCl (pH 7.5), 150 mM NaCl,
5 mM EDTA, and 1% NP-40 to remove unconjugated ligands. Inducible TREx HeLa
cells of FLAG-HA-FAM134B and FLAG-HA-RTN3 induced by treatment of 1 µg/ml
doxycycline for 24 hr or HEK cells transfected with indicated plasmids were lysed and
collected in the lysis buffer supplemented with a protease inhibitor mixture. Cell lysates
were precleared by incubating with streptavidin Sepharose at 4°C with rotation for 1 hr.
Supernatants were transferred to a new microtube and incubated with biotin-LPS-
immobilized streptavidin agarose beads overnight at 4°C with rotation. The beads were
then washed three times with wash buffer containing 50 mM Tris-HCl (pH 7.5), 400
mM NaCl, 5 mM EDTA and eluted in 2X loading dye by boiling for 10 min followed by
western blotting to detect the binding proteins. For competition assay, 50 µg of
unlabeled S-LPS, Ra-LPS, Re-LPS, lipid A, and MDP were first incubated with pre-
cleared lysates at 4°C for 1 hr with rotation before adding biotinylated LPS-immobilized
streptavidin beads. For examination of polymyxin B effect, the indicated concentrations
of polymyxin B were first added to biotinylated LPS-immobilized streptavidin beads and
incubate with rotation for 0.5 hr. After twice washes by lysis buffer, the pre-cleared
lysates were added.
817
818   FAM134B protein purification
819
For purification of FAM134B protein from HEK293T cells, the plasmid of S protein-
FLAG-Streptavidin binding peptide (SFB)-FAM134B was transfected to HEK293T cells.
Cells were lysed with the lysis buffer containing 50 mM Tris-HCl, pH 7.4,150 mM NaCl,
0.5 mM EDTA, and 1% TritonX-100. After centrifuge at 17,000 g, 15 min, the lysates
were collected and incubated with streptavidin Sepharose (Cytiva) at constant agitation
at 4 °C, overnight. After washing three time with the buffers (50 mM Tris, pH=7.4,150
mM NaCl, 0.5 mM EDTA) containing 0.5 % NP-40, 0.3 % NP-40 and 0.1 % NP-40,
respectively, the buffers were removed and the SFB-FAM134B protein on the
Sepharose was further eluted by incubated with 2 mM D-biotin dissolved in the buffer
containing 50 mM Tris-HCl, pH 7.4,150 mM NaCl, 0.5 mM EDTA, and 0.1% NP-40 for
1 hr at 4°C with constant agitation at 800 rpm. The eluted SFB-FAM134B was collected
for blue native page analysis.
Recombinant GST-FAM134B was purified from *E. coli* for Liposome-remodeling
assay as previous described (Gonzalez *et al.*, 2023). Briefly, Fam134B-GST plasmids
were transformed into BL21DE3 strain of *E. coli*, and a single colony was used to grow
the primary culture in the presence of appropriate antibiotic (ampicillin). The overnight
grown primary cultures were used to inoculate secondary cultures and allowed to grow
up to a OD of 0.6 by shaking at 200 rpm and 37 °C, followed by induction of the cultures

using 0.5 mM of IPTG for 3 hours. The cells were harvested, suspended in ice cold
PBS and lysed using sonication, centrifuged (10,000 xg) to separate the supernatant
and pellet. The resultant supernatant was further fractioned by ultracentrifugation at
80,000g for 100 minutes at 4 °C. The pellet fractions were dissolved in PBS buffer
containing dodecyl β-D-maltoside (DDM; 0.05%), followed by affinity purification using
glutathione-Sepharose TM4 Fast Flow columns (GE Healthcare). The protein was
eluted from the column using PBS containing 10 mM reduced glutathione and 0.05%
DDM. The fractions containing Fam134B-GST protein were concentrated and buffered
exchanged with storage buffer (50 mM HEPES pH 7.5, 150 mM NaCl) containing
0.0075%DDM.
848
Blue native page

Blue native page analysis was performed as previous described with slight
modifications (Jiang *et al.*, 2020; Wittig *et al.*, 2006). SFB-FAM134B protein was
incubated with indicated amount of S-LPS or lipid A at 37°C for 2 hr. Then, the mixture
was incubated with 2 μl 50% glycerol and 3 μl 0.1% Coomassie blue G-250 for 15 min
at room temperature followed by loading to 4%–12% gradient native PAGE gel (Bio-
Rad), running in blue cathode/anode buffer (Invitrogen) for 2 hr at 150 V in an ice bath.
The gels were further transfer to PVDF membranes by electroblotting followed by
detection with anti-FLAG antibody.

Liposome preparation

Liposomes were prepared as previous described (Bhaskara *et al.*, 2019). In brief,
two kinds of lipids,1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC, Avanti Polar
Lipids Inc.) and 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE, Avanti Polar
Lipids Inc.), were dissolved in a mixture of chloroform and methanol (4:1) in a glass
flask to obtain the molar ratio of 0.8:0.2 (DOPC:DOPE). The organic solvent was then
evaporated to get a dry lipid film, which was then hydrated with liposome buffer (50
mM HEPES, 150 mM NaCl buffer at pH 7.4) to a final concentration 15 mg/ml lipid
solution. The liposomes were formed by sonication in an ultrasound bath followed by
flash-freeze in liquid nitrogen and slowly thaw in room temperature for 10 times cycles.
The hydrated liposomes were further extruded by a lipid extruder (Avanti Polar Lipids
Inc.) with 200 nm polycarbonate membranes (Avanti Polar Lipids) for 21 times.

Liposome-remodeling assay

Liposome-remodeling assay was performed as previous described with minor
modifications (Bhaskara *et al.*, 2019). Liposomes and the purified FAM134B-GST

protein were mixed at an indicated lipid-to-protein ratio (LPR) and incubated for 18 h at 22 °C in 30 µL of liposome buffer with constant agitation (600 rpm). To investigate the effect of different LPS, FAM134B-GST protein was first added to liposome at LPR 100:1 for 1 h, then added LPS to the proteoliposome and incubated for 17 h at 22 °C. All the samples were assessed by negative-stain transmission electron microscopy. For imaging, samples were first diluted to a final lipid concentration of 1 mg/ml by liposome buffer. 5 µl of each diluted sample was deposited to the carbon-coated copper grids (SPI Supplies), which were already glow-discharged for 20 s at 15 mA and 0.38 mbar vacuum. After 1 minute incubation and washes with buffer for twice, 1% uranyl formate was subsequently used to stain the grids for 10 seconds at room temperature. A 120 kV Tecnai Spirit Biotwin electron microscope (FEI) equipped with a 4k × 4k CCD detector (US4000-1, Gatan) was used to taken 10 micrographs for each sample. Image J software was used to measure the sizes (diameters) of 300 proteoliposomes for each sample.

OMV isolation

0.5 L overnight grown bacterial cultures were harvested for OMV isolation. Bacterial cells were first removed by centrifugation (12,000 g for 15 min), the supernatant was filtered through 0.45 µm PVDF membrane, the filtered broth concentrated to 50 ml (Vivaflow 200, 100,000 MWCO PES, Sartorius), and the OMVs were pelleted by ultracentrifugation at 60,000 rpm (369548.3 g) for 1 hr at 4 °C. The OMVs were washed with PBS and pelleted again by ultracentrifugation. The OMVs were resuspended in 0.5 ml PBS. In the cases which needed to do immunofluorescence labeling, Vybrant DiO cell labeling dye (ThermoFisher Scientific) was added at 1:50 and incubated at 37 °C for 20 min. After labeling, the excess dye was removed and washed by PBS three times at 3000 rpm (956 g) by Micron 10 kDa centrifugal filter units (Millipore) in a benchtop centrifuge. The OMVs were further sterilized by syringe filters (0.22 um) and measured concentration by bicinchoninic acid assay (BCA) method. OMVs were added to the cells at concentration of 25 µg/ml.

Quantification and statistical analysis

GraphPad Prism 8 was used to do statistical analysis. Comparisons of two variables were analyzed by unpaired and two-tailed Student's t tests. For three or more sets of data, statistics were performed by one-way ANOVA with Tukey's multiple comparison post test. * indicates $p < 0.05$, ** indicates $p < 0.01$, *** indicates $p < 0.001$. N represents the number of independent biological replicates. Data are represented as mean ± sem.

REFERENCES

Adams, P.G., Lamoureux, L., Swingle, K.L., Mukundan, H., and Montano, G.A.
(2014). Lipopolysaccharide-induced dynamic lipid membrane reorganization:
tubules, perforations, and stacks. Biophys J *106*, 2395-2407.
10.1016/j.bpj.2014.04.016.

Aglietti, R.A., Estevez, A., Gupta, A., Ramirez, M.G., Liu, P.S., Kayagaki, N., Ciferri,
C., Dixit, V.M., and Dueber, E.C. (2016). GsdmD p30 elicited by caspase-11
during pyroptosis forms pores in membranes. Proc Natl Acad Sci U S A *113*,
7858-7863. 10.1073/pnas.1607769113.

An, H., Ordureau, A., Paulo, J.A., Shoemaker, C.J., Denic, V., and Harper, J.W.
(2019). TEX264 Is an endoplasmic reticulum-resident ATG8-interacting protein
critical for ER remodeling during nutrient stress. Mol Cell *74*, 891-908 e810.
10.1016/j.molcel.2019.03.034.

Bakowski, M.A., Braun, V., Lam, G.Y., Yeung, T., Heo, W.D., Meyer, T., Finlay, B.B.,
Grinstein, S., and Brumell, J.H. (2010). The phosphoinositide phosphatase
SopB manipulates membrane surface charge and trafficking of the
Salmonella-containing vacuole. Cell Host Microbe *7*, 453-462.
10.1016/j.chom.2010.05.011.

Bhaskara, R.M., Grumati, P., Garcia-Pardo, J., Kalayil, S., Covarrubias-Pinto, A.,
Chen, W., Kudryashev, M., Dikic, I., and Hummer, G. (2019). Curvature
induction and membrane remodeling by FAM134B reticulon homology domain
assist selective ER-phagy. Nat Commun *10*, 2370. 10.1038/s41467-019-
10345-3.

Chen, Q., Xiao, Y., Chai, P., Zheng, P., Teng, J., and Chen, J. (2019). ATL3 Is a
tubular ER-phagy receptor for GABARAP-mediated selective autophagy. Curr
Biol *29*, 846-855 e846. 10.1016/j.cub.2019.01.041.

Chino, H., Hatta, T., Natsume, T., and Mizushima, N. (2019). Intrinsically disordered
protein TEX264 mediates ER-phagy. Mol Cell *74*, 909-921 e906.
10.1016/j.molcel.2019.03.033.

Chino, H., and Mizushima, N. (2020). ER-phagy: quality control and turnover of
endoplasmic reticulum. Trends Cell Biol *30*, 384-398.
10.1016/j.tcb.2020.02.001.

Datsenko, K.A., and Wanner, B.L. (2000). One-step inactivation of chromosomal
genes in Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A
97, 6640-6645. 10.1073/pnas.120163297.

Deatherage, B.L., Lara, J.C., Bergsbaken, T., Rassoulian Barrett, S.L., Lara, S., and

946   Cookson, B.T. (2009). Biogenesis of bacterial membrane vesicles. Mol
947   Microbiol *72*, 1395-1407. 10.1111/j.1365-2958.2009.06731.x.

948 Ding, J., Wang, K., Liu, W., She, Y., Sun, Q., Shi, J., Sun, H., Wang, D.C., and Shao,
949   F. (2016). Pore-forming activity and structural autoinhibition of the gasdermin
950   family. Nature *535*, 111-116. 10.1038/nature18590.

951 Elhenawy, W., Bording-Jorgensen, M., Valguarnera, E., Haurat, M.F., Wine, E., and
952   Feldman, M.F. (2016). LPS Remodeling Triggers Formation of Outer
953   Membrane Vesicles in Salmonella. mBio *7*. 10.1128/mBio.00940-16.

954 Escoll, P., Mondino, S., Rolando, M., and Buchrieser, C. (2016). Targeting of host
955   organelles by pathogenic bacteria: a sophisticated subversion strategy. Nat
956   Rev Microbiol *14*, 5-19. 10.1038/nrmicro.2015.1.

957 Fumagalli, F., Noack, J., Bergmann, T.J., Cebollero, E., Pisoni, G.B., Fasana, E.,
958   Fregno, I., Galli, C., Loi, M., Solda, T., et al. (2016). Translocon component
959   Sec62 acts in endoplasmic reticulum turnover during stress recovery. Nat Cell
960   Biol *18*, 1173-1184. 10.1038/ncb3423.

961 Garcia-del Portillo, F., Stein, M.A., and Finlay, B.B. (1997). Release of
962   lipopolysaccharide from intracellular compartments containing Salmonella
963   typhimurium to vesicles of the host epithelial cell. Infect Immun *65*, 24-34.
964   10.1128/iai.65.1.24-34.1997.

965 Gatica, D., Alsaadi, R.M., Hamra, R.E., Mueller, R., Miyazaki, M., Sad, S., and
966   Russell, R.C. (2024). The ER-phagy receptor FAM134B is targeted by
967   *Salmonella* Typhimurium to promote infection. bioRxiv,
968   2024.2003.2028.587227. 10.1101/2024.03.28.587227.

969 Gonzalez, A., Covarrubias-Pinto, A., Bhaskara, R.M., Glogger, M., Kuncha, S.K.,
970   Xavier, A., Seemann, E., Misra, M., Hoffmann, M.E., Brauning, B., et al.
971   (2023). Ubiquitination regulates ER-phagy and remodelling of endoplasmic
972   reticulum. Nature. 10.1038/s41586-023-06089-2.

973 Grumati, P., Morozzi, G., Holper, S., Mari, M., Harwardt, M.I., Yan, R., Muller, S.,
974   Reggiori, F., Heilemann, M., and Dikic, I. (2017). Full length RTN3 regulates
975   turnover of tubular endoplasmic reticulum via selective autophagy. Elife *6*.
976   10.7554/eLife.25555.

977 Gubas, A., and Dikic, I. (2022). ER remodeling via ER-phagy. Mol Cell *82*, 1492-
978   1500. 10.1016/j.molcel.2022.02.018.

979 Hoffmann, S., Schmidt, C., Walter, S., Bender, J.K., and Gerlach, R.G. (2017).
980   Scarless deletion of up to seven methyl-accepting chemotaxis genes with an
981   optimized method highlights key function of CheM in Salmonella Typhimurium.
982   PLoS One *12*, e0172630. 10.1371/journal.pone.0172630.

Jiang, X., Wang, X., Ding, X., Du, M., Li, B., Weng, X., Zhang, J., Li, L., Tian, R., Zhu,

Q., et al. (2020). FAM134B oligomerization drives endoplasmic reticulum
membrane scission for ER-phagy. EMBO J *39*, e102608.
10.15252/embj.2019102608.

Juodeikis, R., and Carding, S.R. (2022). Outer membrane vesicles: biogenesis,
functions, and issues. Microbiol Mol Biol Rev *86*, e0003222.
10.1128/mmbr.00032-22.

Kawasaki, K., Ernst, R.K., and Miller, S.I. (2004). 3-O-deacylation of lipid A by PagL,
a PhoP/PhoQ-regulated deacylase of Salmonella typhimurium, modulates
signaling through Toll-like receptor 4. J Biol Chem *279*, 20044-20048.
10.1074/jbc.M401275200.

Khaminets, A., Heinrich, T., Mari, M., Grumati, P., Huebner, A.K., Akutsu, M.,
Liebmann, L., Stolz, A., Nietzsche, S., Koch, N., et al. (2015). Regulation of
endoplasmic reticulum turnover by selective autophagy. Nature *522*, 354-358.
10.1038/nature14498.

Kumar, D., Lak, B., Suntio, T., Vihinen, H., Belevich, I., Viita, T., Xiaonan, L.,
Vartiainen, A., Vartiainen, M., Varjosalo, M., and Jokitalo, E. (2021). RTN4B
interacting protein FAM134C promotes ER membrane curvature and has a
functional role in autophagy. Mol Biol Cell *32*, 1158-1170. 10.1091/mbc.E20-
06-0409.

Kutsch, M., Sistemich, L., Lesser, C.F., Goldberg, M.B., Herrmann, C., and Coers, J.
(2020). Direct binding of polymeric GBP1 to LPS disrupts bacterial cell
envelope functions. EMBO J *39*, e104926. 10.15252/embj.2020104926.

LaRock, D.L., Chaudhary, A., and Miller, S.I. (2015). Salmonellae interactions with
host processes. Nat Rev Microbiol *13*, 191-205. 10.1038/nrmicro3420.

Madan, R., Rastogi, R., Parashuraman, S., and Mukhopadhyay, A. (2012).
Salmonella acquires lysosome-associated membrane protein 1 (LAMP1) on
phagosomes from Golgi via SipC protein-mediated recruitment of host
Syntaxin6. J Biol Chem *287*, 5574-5587. 10.1074/jbc.M111.286120.

Man, S.M., Karki, R., Sasai, M., Place, D.E., Kesavardhana, S., Temirov, J., Frase,
S., Zhu, Q., Malireddi, R.K.S., Kuriakose, T., et al. (2016). IRGB10 liberates
bacterial ligands for sensing by the AIM2 and caspase-11-NLRP3
inflammasomes. Cell *167*, 382-396 e317. 10.1016/j.cell.2016.09.012.

Meunier, E., Dick, M.S., Dreier, R.F., Schurmann, N., Kenzelmann Broz, D., Warming,
S., Roose-Girma, M., Bumann, D., Kayagaki, N., Takeda, K., et al. (2014).
Caspase-11 activation requires lysis of pathogen-containing vacuoles by IFN-
induced GTPases. Nature *509*, 366-370. 10.1038/nature13157.

Moretti, J., Roy, S., Bozec, D., Martinez, J., Chapman, J.R., Ueberheide, B.,
Lamming, D.W., Chen, Z.J., Horng, T., Yeretssian, G., et al. (2017). STING

1022  senses microbial viability to orchestrate stress-mediated autophagy of the
1023  endoplasmic reticulum. Cell *171*, 809-823 e813. 10.1016/j.cell.2017.09.034.

1024 Nevermann, J., Silva, A., Otero, C., Oyarzun, D.P., Barrera, B., Gil, F., Calderon, I.L.,
1025  and Fuentes, J.A. (2019). Identification of genes involved in biogenesis of
1026  outer membrane vesicles (OMVs) in *Salmonella enterica* Serovar Typhi. Front
1027  Microbiol *10*, 104. 10.3389/fmicb.2019.00104.

1028 Nthiga, T.M., Kumar Shrestha, B., Sjottem, E., Bruun, J.A., Bowitz Larsen, K.,
1029  Bhujabal, Z., Lamark, T., and Johansen, T. (2020). CALCOCO1 acts with
1030  VAMP-associated proteins to mediate ER-phagy. EMBO J *39*, e103649.
1031  10.15252/embj.2019103649.

1032 Park, B.S., Song, D.H., Kim, H.M., Choi, B.S., Lee, H., and Lee, J.O. (2009). The
1033  structural basis of lipopolysaccharide recognition by the TLR4-MD-2 complex.
1034  Nature *458*, 1191-1195. 10.1038/nature07830.

1035 Perrin, A.J., Jiang, X., Birmingham, C.L., So, N.S., and Brumell, J.H. (2004).
1036  Recognition of bacteria in the cytosol of Mammalian cells by the ubiquitin
1037  system. Curr Biol *14*, 806-811. 10.1016/j.cub.2004.04.033.

1038 Ravenhill, B.J., Boyle, K.B., von Muhlinen, N., Ellison, C.J., Masson, G.R., Otten,
1039  E.G., Foeglein, A., Williams, R., and Randow, F. (2019). The Cargo Receptor
1040  NDP52 Initiates Selective Autophagy by Recruiting the ULK Complex to
1041  Cytosol-Invading Bacteria. Mol Cell *74*, 320-329 e326.
1042  10.1016/j.molcel.2019.01.041.

1043 Reggio, A., Buonomo, V., Berkane, R., Bhaskara, R.M., Tellechea, M., Peluso, I.,
1044  Polishchuk, E., Di Lorenzo, G., Cirillo, C., Esposito, M., et al. (2021). Role of
1045  FAM134 paralogues in endoplasmic reticulum remodeling, ER-phagy, and
1046  Collagen quality control. EMBO Rep *22*, e52289. 10.15252/embr.202052289.

1047 Santos, J.C., Boucher, D., Schneider, L.K., Demarco, B., Dilucca, M., Shkarina, K.,
1048  Heilig, R., Chen, K.W., Lim, R.Y.H., and Broz, P. (2020). Human GBP1 binds
1049  LPS to initiate assembly of a caspase-4 activating platform on cytosolic
1050  bacteria. Nat Commun *11*, 3276. 10.1038/s41467-020-16889-z.

1051 Santos, N.C., Silva, A.C., Castanho, M.A., Martins-Silva, J., and Saldanha, C. (2003).
1052  Evaluation of lipopolysaccharide aggregation by light scattering spectroscopy.
1053  Chembiochem *4*, 96-100. 10.1002/cbic.200390020.

1054 Schwechheimer, C., and Kuehn, M.J. (2015). Outer-membrane vesicles from Gram-
1055  negative bacteria: biogenesis and functions. Nat Rev Microbiol *13*, 605-619.
1056  10.1038/nrmicro3525.

1057 Shi, J., Zhao, Y., Wang, Y., Gao, W., Ding, J., Li, P., Hu, L., and Shao, F. (2014).
1058  Inflammatory caspases are innate immune receptors for intracellular LPS.
1059  Nature *514*, 187-192. 10.1038/nature13683.

Siggel, M., Bhaskara, R.M., Moesser, M.K., I, D.I., and Hummer, G. (2021).
FAM134B-RHD protein clustering drives spontaneous budding of asymmetric
membranes. J Phys Chem Lett *12*, 1926-1931. 10.1021/acs.jpclett.1c00031.

Silhavy, T.J., Kahne, D., and Walker, S. (2010). The bacterial cell envelope. Cold
Spring Harb Perspect Biol *2*, a000414. 10.1101/cshperspect.a000414.

Smith, M.D., Harley, M.E., Kemp, A.J., Wills, J., Lee, M., Arends, M., von Kriegsheim,
A., Behrends, C., and Wilkinson, S. (2018). CCPG1 Is a non-canonical
autophagy cargo receptor essential for ER-phagy and pancreatic ER
proteostasis. Dev Cell *44*, 217-232 e211. 10.1016/j.devcel.2017.11.024.

Steele-Mortimer, O. (2008). Infection of epithelial cells with Salmonella enterica.
Methods Mol Biol *431*, 201-211. 10.1007/978-1-60327-032-8_16.

Szczepaniak, J., Press, C., and Kleanthous, C. (2020). The multifarious roles of Tol-
Pal in Gram-negative bacteria. FEMS Microbiol Rev *44*, 490-506.
10.1093/femsre/fuaa018.

Vanaja, S.K., Russo, A.J., Behl, B., Banerjee, I., Yankova, M., Deshmukh, S.D., and
Rathinam, V.A.K. (2016). Bacterial outer membrane vesicles mediate cytosolic
localization of LPS and caspase-11 activation. Cell *165*, 1106-1119.
10.1016/j.cell.2016.04.015.

Wild, P., Farhan, H., McEwan, D.G., Wagner, S., Rogov, V.V., Brady, N.R., Richter, B.,
Korac, J., Waidmann, O., Choudhary, C., et al. (2011). Phosphorylation of the
autophagy receptor optineurin restricts Salmonella growth. Science *333*, 228-
233. 10.1126/science.1205405.

Wittig, I., Braun, H.P., and Schagger, H. (2006). Blue native PAGE. Nat Protoc *1*,
418-428. 10.1038/nprot.2006.62.

Zenk, S.F., Jantsch, J., and Hensel, M. (2009). Role of Salmonella enterica
lipopolysaccharide in activation of dendritic cell functions and bacterial
containment. J Immunol *183*, 2697-2707. 10.4049/jimmunol.0900937.

1087

Figure 1 Cheng et al

Figure 2 Cheng et al

Figure 3 Cheng et al

Figure 4 Cheng et al

Figure 5 Cheng et al

Supplementary information

Intracellular lipopolysaccharide regulates ER remodeling upon bacterial infection

Authors

Yi-Lin Cheng, João Mello-Vieira, Adriana Covarrubias-Pinto, Alexis Gonzalez, Santosh Kumar Kuncha, Chun Kew, Kaiyi Zhang, Muhammad Awais Afzal,   Nour Diab, Sophia Borchert, Wenbo Chen, Timothy Chun Huang, Mathias Walter Hornef, Christian A. Hübner, Michael Hensel, Ivan Dikic

Figure S1

Figure S1. *Salmonella* infection mainly activates FAM134B-mediated ER fragmentation, related to Figure 1

(A) HeLa cells were infected with SL1344 for the indicated time points followed by analyzing the protein expression of indicated ER-phagy receptors, and LC3B by western blotting. GAPDH was used as an internal control of input.

(B) Immunofluorescence staining of FLAG and LC3B in HeLa TREx inducible cell lines expressing FLAG-HA-RTN3L, which were infected with SL1344 for 2 hr. DAPI was used for cell nuclear and bacterial DNA staining. Scale bar, 10 μm.

(C) Immunofluorescence staining of HA, *Salmonella* LPS, and LC3B in HeLa TREx inducible cell lines expressing LIR mutant (mtLIR) FLAG-HA-FAM134B infected with SL1344 for 2 hr. DAPI was used for cell nuclear and bacterial DNA staining. Scale bar, 10 μm.

(D) Immunofluorescence staining of FAM134B and *Salmonella* LPS in HeLa cells infected by SL1344 for 2 hr. Arrows indicate the puncta with colocalization of FAM134B and *Salmonella* LPS.

(E) Immunofluorescence staining of HA and LC3B in HeLa TREx inducible cell lines expressing wild type (WT) or mtLIR FLAG-HA-FAM134B infected with SL1344 for 2 hr. DAPI was used for cell nuclear and bacterial DNA staining. Scale bar, 10 μm.

(F) Immunofluorescence staining of HA, LAMP1, and *Salmonella* LPS in HeLa TREx inducible cell lines expressing FLAG-HA-FAM134B, which were transfected with 1 μg/ml *Salmonella* LPS for 18 hr by lipofectamine 2000. Lipo, lipofectamine only.

Figure S2

Figure S2. LPS bind to FAM134B and FAM134C, but not RTN3L and FAM134A, related to Figure 2

(A) Co-IP of LPS with HA in HeLa TREx inducible cell lines without Dox treatment or expressing FLAG-HA-FAM134B WT or FLAG-HA-FAM134B mtLIR with Dox treatment infected with SL1344 100 moi for 2 hr. GAPDH was used as an internal control of input.

(B) Streptavidin pulldown assays for determining the binding of the transfected biotin-conjugated *E. coli* O111:B4 LPS (10 µg) to HA-tagged FAM134B or RTN3L in FLAG-HA-FAM134B- or FLAG-HA-RTN3L-expressing HeLa TREx cell lines. GAPDH was used as an internal control of input.

(C) Streptavidin pulldown assays determining the binding of the biotin-conjugated *E. coli* O111:B4 LPS to HA-tagged FAM134A, FAM134B, or FAM134C in HA-FAM134A-, HA-FAM134B-, or HA-FAM134C-transfected HeLa cell lysates. GAPDH was used as an internal control of input.

(D-F) Phylogenetic tree of FAM134 proteins: Phylogenetic tree of FAM134A, B and C was generated by using ClustalW followed by analyzing the data through the iTOL server (https://itol.embl.de/). Representative sequences were derived from the FASTA for the following organism across phylum Chordata. Dr: *Danio rerio*, Xt: *Xenopus tropicalis*, Gg: *Gallu gallus*, Rn: *Rattus norvegicus*, Mm: *Mus musculus* and Hs: *Homo sapiens*

Figure S3

Figure S3. S-LPS trigger membrane fragmentation, related to Figure 3

(A-B) Representative images (A) of remodeled proteoliposomes (scale bars, 100 nm) taken from nsTEM. Empty liposomes (Empty) were added to FAM134B-GST recombinant protein at a lipid to protein ratio (LPR) 100:1 (LPR 100:1) or 40:1 (LPR 40:1). The size distributions (n=300 each) from nsTEM images were measured and presented by violin plots, showing a central boxplot (black lines indicate median with interquartile range) with mirrored histograms on both sides (B). Differences

were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test. ***p < 0.001. ns, not significant.

(C) Immunofluorescence of HA, *Salmonella* LPS, and LC3B in HeLa TREx inducible cell line expressing FLAG-HA-FAM134B were infected with NCTC12023 WT or LPS O-Ag mutant (Δ) and complemented (compl.) strains for 2 hr. DAPI staining was used for cell nuclear and bacterial DNA staining. Scale bar, 10 μm.

(D-E) Immunofluorescence staining of HA and LC3B in HeLa TREx inducible cell line expressing FLAG-HA-FAM134B, transfected with 1 μg/ml S-LPS, Ra-LPS, Re-LPS, Lipid A, Pam3CSK4, and MDP for 18 hr by lipofectamine 2000 (D). Lipo, lipofectamine only. DAPI was used for cell nuclear DNA staining. Scale bar, 10 μm. Quantification of LC3B-positive HA-FAM134B puncta per cell (E). Solid bars of boxes indicate the medians. Boxes represent interquartile range from 25^{th} to 75^{th} percentile, and whiskers indicate 10^{th} to 90^{th} percentile. Differences were statistically analyzed by one-way ANOVA and Tukey's multiple comparison test. ***p < 0.001. ns, not significant. Data were collected 60 cells from three independent biological replicates.

Figure S4

Figure S4. *Salmonella* **activates FAM134B-mediated ER-phagy and ER fragmentation, forming bacteria-containing vacuoles positive for FAM134B and ER markers, related to Figure 4**

(A-C) HeLa TREx inducible cell lines expressing mCherry-eGFP-FAM134B WT or mCherry-eGFP-FAM134B mtLIR were infected with SL1344 for 2 hr or 4hr (A). The inset of square 1 showed the *Salmonella* surrounded by mCherry and eGFP double positive-FAM134B (A). Quantification of mCherry-positive and eGFP-negative puncta (red) and double-positive (yellow) puncta per cell (B). The cell with hyperproliferating *Salmonella* showed bacteria surrounded by mCherry and eGFP double positive-FAM134B (C). DAPI staining was used for cell nuclear and bacterial DNA staining. Scale bar, 10 μm.

(D) Immunofluorescence staining of FAM134B (red) in *Salmonella*-infected intestinal tissue section obtained at day 4 post-infection. *Salmonella* GFP staining was used for detecting GFP plasmid carried by *Salmonella* (white). DAPI was used for nuclear DNA staining. Scale bar, 20 μm.

(E-F) Immunofluorescence staining of HA and Calnexin (E) or REEP5 (F) in HeLa TREx inducible cell lines expressing FLAG-HA-FAM134B, which were infected with SL1344 for 4 hr. DAPI was used for nuclear and bacterial DNA staining. Scale bar, 10 μm.

Figure S5

Figure S5., FAM134B promotes ER-surrounded *Salmonella*-containing vacuoles formation and bacterial growth in BMDM, related to Figure 4.

(A-C) Immunofluorescence staining of REEP5 (ER marker) and *Salmonella* LPS in WT BMDMs or FAM134B KO BMDMs, which were infected with SL1344 for 2hr (2hpi) or 4 hr (4hpi). *Salmonella* LPS staining was used to indicate bacteria. Quantification of immunofluorescence analysis represented as (B) number of vacuoles stained with LPS and REEP5 per cell, (C) percentage of *Salmonella* (LPS) positive to REEP5 from total *Salmonella*. Data are mean ± sem of the number of ER-positive *Salmonella*-containing vacuoles (LPS+ and REEP+) and % of these vacuoles (LPS+ and REEP5+) from total *Salmonella* (LPS+) per cell (2 h: $n = 34$ for the FAM134B KO, $n = 30$ for the WT and 4h: $n = 36$ for FAM134B KO and $n = 29$, Kruskal–Wallis/Dunn's post-test). **p<0.01, **p<0.0001, ****p<0.0001 ns, not significant.

(D) Colony-forming assay was performed to quantify the bacteria count in WT or FAM134B KO BMDMs followed by infection with SL1344 10 moi for the indicated timepoints. The growth fold change at 4 hr was normalized to 2 hr, error bars represent standard error of the mean. *p<0.05. Data were collected from three pairs of WT and FAM134B KO mice.

Figure S6

Figure S6. FAM134B-interacting *Salmonella* proteins obtained from the interactome of FAM134B, related to Figure 4

(A) MS workflow for the analysis of HA–FAM134B co-immunoprecipitated from cell lysates following *Salmonella* infection (2 and 4h) and Control (non-infected cells, blue) or from mock (HA empty, light blue). HF, Q Exactive HF (Ultra-High-Field Orbitrap) mass spectrometer; LFQ, label free quantification; 3xR 3 biological replicates.

(B) Table showing the list of *Salmonella* effectors found in the HA-FAM134B co-immunoprecipitation. Significant and enriched hits are colored in red (significant values were $\log_2$ enrichment > 1.0 and $-\log_{10}$ p value > 1.3). Data are means ± s.d. of $n = 3$ independent experiments.

(C) Heat map comparing the interaction of Salmonella effector proteins with WT FAM134B after 2 or 4 hours of infections (immuno-isolated using HA beads). Interaction partners with $\log_2$ enrichment > 1.0 and $-\log_{10}$ p value > 1.3 were plotted. $n = 3$ independent experiments, one-tailed unpaired Student's test.

(D) Annotation enrichment analysis of the *Salmonella* effector proteins interacting with FAM134B. Bars represent significantly enriched gene ontology biological process (GOBP), gene ontology cellular component (GOCC) for the identified *Salmonella* proteins.

Figure S7

Figure S7. Formation of OMV by various *Salmonella* strains and OMV size distribution, related to Figure 5.

(A-C) STM WT, or mutant strains deficient in *pagL*, *tolAB*, *ompA*, or *degS* were grown on filter paper placed on LB agar (A), or overnight on LB agar to allow formation of colonies (B). (A) Cells were fixed, dehydrated, and coated with 10 nm gold and prepared for SEM imaging on JEOL IT-200 and Zeiss Auriga systems. (B) For TEM micrographs, samples were prepared using negative stain. The study was performed as triplicates, 20 bacteria per strain were recorded. Cultures were grown statically on filter papers on LB agar. The experiment was performed as duplicates, at least ten images were generated from each strain. Representative OMV are indicated by arrowheads. Scale bars, 500 nm. (C) Size distribution of OMV was determined using morphometric analyses in FIJI. For this, the areas of OMV of at least six bacteria per strains were quantified and the frequency of various OMV size populations was plotted as histogram. The WT strain possessed OMVs of 918 nm^2 on average, and OMVs up to 2,000 nm^2 were also detected. No vesicles could be detected for $\Delta pagL$. The other three mutant strains showed increased production of OMVs compared to WT strain, and the average sizes of OMVs were 1,611 nm^2 in $\Delta tolAB$, 3,153 nm^2 in $\Delta ompA$, and 590 nm^2 in $\Delta degS$.

(D) Immunofluorescence staining of HA, *Salmonella* LPS, and LC3B in cells infected with different mutant strains, including $\Delta ompA$, $\Delta tolAB$, and $\Delta degS$, and $\Delta pagL$ strains. DAPI was used for cell nuclear and bacterial DNA staining. Scale bar, 10 μm. Quantification of LC3B-positive HA-FAM134B puncta and *Salmonella* count per cell are shown in Figure 5A and 5B.

Figure S8

A

B

Figure S8. FAM134B colocalized with LPS, which is not associated with OMVs, related to Figure 5

(A) *Salmonella* OMVs were stained by DiO Vybrant dye first, then applied to HeLa TREx inducible cell lines expressing FLAG-HA-FAM134B for 4 hr followed by Immunofluorescence staining of HA and *Salmonella* LPS. DAPI was used for nuclear DNA staining. Scale bar, 10 μm.

(B) EPEC or BL21 OMVs were stained by DiO Vybrant dye first, then applied to HeLa TREx inducible cell lines expressing FLAG-HA-FAM134B for 4 hr followed by Immunofluorescence staining of HA and LC3B.

METHOD DETAILS

Mouse model for *S.* Typhimurium infection
Salmonella enterica subsp. enterica serovar Typhimurium ATCC14028 (NCTC12023) carrying a green fluorescent protein (GFP) expression plasmid (pGFP, AmpR, kindly provided by Brendan Cormack, Stanford, USA) were cultured in Luria Bertani (LB) broth overnight at 37°C. The overnight culture was diluted 1:10 in fresh LB and incubated at 37°C until reaching the logarithmic phase (OD600 approximately 0.5). Bacteria were washed and adjusted to OD600 0.55–0.60 containing approximately 1.5–2.0×10^8 CFU/ml. One-day-old C57BL/6 animals were infected orally with 10^2 CFU *S.* Typhimurium in a volume of 1 µl PBS. The administered inoculum was confirmed by serial dilution and plating. At day 4 post-infection, the small intestine tissue was collected and fixed in 4% PFA.

HA-FAM134B interactome analysis in cells infected with *Salmonella* and sample preparation for MS

HeLa TREx cells were induced with doxycycline to express HA-FLAG FAM134B. After 16 h, cells were infected with *Salmonella* for 2 and 4 hours. After the infection with *Salmonella*, cells were lysed with 1% Triton X-100 in lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, Roche EDTA-free protease inhibitor cocktail and NEM), followed by incubation with HA-beads (Sigma-Aldrich) overnight at 4 °C on a rotating platform. Protein-bound beads were washed three times with lysis buffer and three times in the same buffer without detergents before the elution. Samples were eluted with 25 µl SDC buffer (2% sodium deoxycholate and 50 mM Tris-HCl pH 8.5) and heated to 95 °C for 5 min. Additionally, 25 ul of 50 mM Tris-HCl pH 8.5 containing 1 mM TCEP, 4 mM chloroacetamide were added to the samples in order to alkylate and reduce the proteins and incubated to 95 °C for 5 min. We then added 500 ng trypsin in 25 µl 50 mM Tris-HCl (pH 8.5) to each sample and incubated overnight at 37 °C. The reaction was stopped with 150 µl of 1% TFA in isopropanol. Peptides were cleaned up using SDB-RPS stage tips (Sigma-Aldrich). After one wash with 1% TFA in isopropanol and one wash with 0.2% TFA in water, peptides were eluted in 80% acetonitrile plus 1.25% ammonia. Eluted peptides were dried and processed for LC–MS.

LC–MS analysis

Dried peptides were resolved in 2% acetonitrile containing 0.1% TFA and loaded to the 35-cm, 75-µm ID fused-silica column, which packed in house with 1.9-µm C18 particles (Reprosil pur, Dr. Maisch), then analyzed on a QExactive HF mass spectrometer coupled to an easy nLC 1200 (Thermo Fisher Scientific). An integrated column oven (Sonation) was used to maintain the column at 50 °C. Peptides were eluted using a non-linear gradient of 4–28% acetonitrile over 45 min followed by sprayed into the mass spectrometer with a nanoFlex ion source (Thermo Fisher Scientific). Full-scan MS spectra (300–1,650 *m/z*) were acquired in a profile mode with resolution of 60,000 at *m/z* 200, a maximum injection time of 20 ms and an automatic gain control (AGC) target value of 3 × 10^6. Using a 1.4-Th window, up to 15 of the most intense peptides per full scan were isolated for fragmentation by higher energy collisional dissociation with normalized collision energy of 28.

MS/MS spectra were acquired at a resolution of 30,000, a maximum injection time of 45 ms and an AGC target value of 1×10^5 in centroid mode. Ions with unassigned charge states, single charged ions, and ions with a charge state of more than four were excluded for fragmentation. To minimize the acquisition of fragment spectra representing already acquired precursors, dynamic exclusion was set to 20 s.

MS data processing

MaxQuant (v1.6.17.0) was used to process MS raw data. Acquired spectra were searched against the human 'one sequence per gene' database (Taxonomy ID 9606) downloaded from UniProt (12 March 2020; 20,531 sequences), and a collection of 244 common contaminants ('contaminants.fasta' provided with MaxQuant v1.6.17.0) using the Andromeda search engine integrated into MaxQuant (v1.6.17.0). Protein identifications were obtained by filtering at false discovery rates below 1% for both peptide spectrum matches and proteins using a target–decoy strategy (Elias and Gygi, 2007). Protein quantification and data normalization relied on the MaxLFQ algorithm (Cox et al., 2014). The MS proteomics data have been deposited to the ProteomeXchange Consortium through the PRIDE partner repository (Perez-Riverol et al., 2022) with the dataset identifiers PXD053709 (Fig. S6). For protein assignment, the UniProt human database v2019 including a list of common contaminants was introduced to correlate with spectra. Tryptic specifications and default settings for mass tolerances in MS and MS/MS spectra were applied for searches. The match-between-run feature was used with a time window of 1 min. For further analysis, Perseus (v2.0.7.0) was applied to obtain the proteins with significant changed between groups after filtering contaminants, reverse entries, and proteins that were only identified by a modified peptide.

Establishment of mCherry-EGFP-FAM134B inducible cell lines

mCherry-EGFP-FAM134B WT and mCherry-EGFP-FAM134B mtLIR constructs were cloned into pcDNA5/FRT/TO vector by GATEWAY technology, followed by co-transfection with recombinase pOG44 into Flp-In HeLa TREx cells. Then, 400 µg/ml hygromycin and 15 µg/ml blasticidin were treated for selection and maintenance of stable cells. Induction was performed with 2 µg/ml doxycycline for 24 hr.

Analysis of OMV production by SEM

For ultrastructural analyses, liquid cultures were incubated overnight. Circles of Whatman Nucleopore membrane filters (pore size 0.1 µm) were punched out of using a biopsy punch. The filters were placed shiny side up on an agar plate, and 2 µl of bacterial cultures were transferred to the center of filters. The agar plates were incubated overnight at 37 °C. The filters were fixed for 15 min in a 1.5 ml reaction tube containing 2.5% GA in PBS. The fixative was removed and washed three times with PBS. An increasing EtOH series was used to dehydrate the samples. This involved 10%, 30%, 50%, 70%, 90% EtOH for 10 min each, with the last three dilution levels

applied twice. Subsequently, the samples were incubated in 100% EtOH for 15 min and critical point drying was performed. After the samples were glued onto SEM stubs, they were coated with a 10 nm thick gold layer in order to inhibit charging during microscopy. Afterwards, the samples were imaged using a JEOL IT-200 for overview images, and with a Zeiss Auriga 60 for high-resolution images.

Analysis of OMV production using negative stain TEM

For high-resolution ultrastructural analyses, *Salmonella* cultures were grown on agar plates, then bacterial material was picked up with an inoculation loop and resuspended in 500 µl PBS. Here, the samples were carefully inverted to inhibit detachment of OMVs. Then, 5 µl of the bacterial suspension was transferred to a glow-discharged EM copper slot grid with Formvar charcoal coating and the cells were allowed to adhere for 5 min. The suspension was carefully removed with a filter paper and subsequently washed three times with 5 µl MilliQ. For negative stain, 1% uranyl acetate (UA) was applied to the grid for 1 min, and the excess UA was then removed with a filter paper. Afterwards, the samples were imaged by the Zeiss LEO912 and the JEOL JEM 2100Plus.

REFERENCES

Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M. (2014). Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. Mol Cell Proteomics *13*, 2513-2526. 10.1074/mcp.M113.031591.

Elias, J.E., and Gygi, S.P. (2007). Target-decoy search strategy for increased confidence in large-scale protein identifications by mass spectrometry. Nat Methods *4*, 207-214. 10.1038/nmeth1019.

Perez-Riverol, Y., Bai, J., Bandla, C., Garcia-Seisdedos, D., Hewapathirana, S., Kamatchinathan, S., Kundu, D.J., Prakash, A., Frericks-Zipper, A., Eisenacher, M., et al. (2022). The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. Nucleic Acids Res *50*, D543-D552. 10.1093/nar/gkab1038.

KEY RESOURCES TABLE

Intracellular lipopolysaccharide regulates ER remodeling upon bacterial infection

Authors

Yi-Lin Cheng, João Mello-Vieira, Adriana Covarrubias-Pinto, Alexis Gonzalez, Santosh Kumar Kuncha, Chun Kew, Kaiyi Zhang, Muhammad Awais Afzal, Nour Diab, Sophia Borchert, Wenbo Chen, Timothy Chun Huang, Mathias Walter Hornef, Christian A. Hübner, Michael Hensel, Ivan Dikic

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
HA	Roche	Cat# 11867423001; RRID:AB_390918
GAPDH	Cell signaling	Cat# 2118 RRID:AB_561053
Salmonella LPS	Abcam	Cat# ab8274 RRID:AB_306423
LC3B	MBL Life Science	Cat# PM036 RRID:AB_2274121
LC3B	MBL Life Science	Cat# M152-3 RRID:AB_1279144
Biotin	Santa Cruz	Cat# sc-101339 RRID:AB_1119609
FLAG	Sigma Aldrich	Cat# F3165 RRID:AB_259529
LAMP1	Abcam	Cat# ab24170 RRID:AB_775978
FAM134B	Proteintech	Cat# 21537-1-AP RRID:AB__2878879
CCPG1	Polyclonal rabbit, affinity purified with N-term peptide, gift from Simon Wilkinson	N/A
Sec62	Novusbio	Cat# NBP1-84045 RRID:AB_11037283
RTN3L	TebuBio	Cat# PA2256 RRID:AB_2665372
TEX264	Novus	Cat# NBP1-89866 RRID:AB_11009420
Calnexin	Abcam	Cat# ab22595 RRID:AB_2069006
REEP5	Santa Cruz	Cat# sc-393508
GAPDH	Cell signaling	Cat#2118S, RRID: AB_561053
GFP (for Salmonella-GFP)	Abcam	Cat#ab13970 RRID:AB_300798
Chemicals		
Biotinylated ultrapure LPS from *E. coli* 0111:B4	InvivoGen	Cat# tlrl-lpsbiot
LPS from *Salmonella enterica* serotype typhimurium	Sigma-Aldrich	Cat# L2262

LPS (rough strains) from *Salmonella enterica* serotype typhimurium TV119 (Ra mutant)	Sigma-Aldrich	Cat# L6016
LPS (rough strains) from *Salmonella enterica* serotype typhimurium SL1181 (Re mutant)	Sigma-Aldrich	Cat# L9516
Lipid A, monophosphoryl *Salmonella enterica* serotype minnesota Re 595 (Re mutant), Ready Made Solution	Sigma-Aldrich	Cat# SBR00012
Muramyldipeptide	InvivoGen	Cat# tlrl-mdp
Pam3CSK4	InvivoGen	Cat# tlrl-pms
Polymyxin B sulfate salt	Sigma-Aldrich	Cat# P4932-1MU
Recombinant DNA		
pcDNA3.1-FAM134B-WT-HA	Dikic Lab	N/A
pcDNA3.1-FAM134B-AHLmt-HA	This study	N/A
pcDNA3.1-FAM134B-AHCmt-HA	This study	N/A
pcDNA3.1-FAM134B-CTERmt-HA	This study	N/A
pcDNA3.1-FAM134B-DOUBLEmt1	This study	N/A
pcDNA3.1-FAM134B-DOUBLEmt2	This study	N/A
pcDNA3.1-FAM134B-TRIPLEmt	This study	N/A
pDEST-pMH-SFB-FAM134B	Dikic Lab	N/A
pGEX6P1-FAM134B-GST	Dikic Lab	N/A
Experimental Models: Cell Lines		
HeLa	ATCC	Cat# CCL-2; RRID: CVCL_0030
HEK293T	ATCC	Cat# CRL-3216; RRID: CVCL_0063
HeLa TREx Cells	Invitrogen	Cat# R71407; RRID:CVCL_D587
MEF WT	Christian Hübner	N/A
MEF FAM134BKO	Christian Hübner	N/A
BMDM WT	Christian Hübner	N/A
BMDM FAM134BKO	Christian Hübner	N/A
Experimental Models: Organisms/Strains		
Salmonella Typhimurium SL1344	Dikic Lab	N/A

Salmonella Typhimurium NCTC12023	Michael Hensel	N/A
Salmonella Typhimurium MvP887 (Δ*rfbP*)	Michael Hensel	N/A
Salmonella Typhimurium MvP886 (Δ*rfaL*)	Michael Hensel	N/A
Salmonella Typhimurium MvP885 (Δ*rfaG*)	Michael Hensel	N/A
Salmonella Typhimurium MvP759 (Δ*rfaH*)	Michael Hensel	N/A
Salmonella Typhimurium P3313 (*rfaL* complemented strain)	Michael Hensel	N/A
Salmonella Typhimurium P3314 (*rfaG* complemented strain)	Michael Hensel	N/A
Salmonella Typhimurium MvP2462 (Δ*ompA*)	Michael Hensel	N/A
Salmonella Typhimurium MvP2464 (Δ*tolAB*)	Michael Hensel	N/A
Salmonella Typhimurium MvP2555 (Δ*degS*)	Michael Hensel	N/A
Salmonella Typhimurium MvP544 (Δ*pagL*)	Michael Hensel	N/A
Enteropathogenic *Escherichia coli* O127:H6	Volkhard A. J. Kempf	N/A
Escherichia coli BL21	New England Biolabs	N/A
Escherichia coli Turbo	New England Biolabs	N/A
Software and Algorithms		
ClustalW	(Larkin et al., 2007)	https://www.ebi.ac.uk/Tools/msa/clustalw2/
iTOL server	(Letunic and Bork, 2021)	https://itol.embl.de/
GraphPad Prism 8.0	GraphPad Software	https://www.graphpad.com/
Zen (black edition)	Zeiss	https://www.micro-shop.zeiss.com/en/us/softwarefinder/software-categories/zen-black/
MaxQuant	(Tyanova et al., 2016a)	https://maxquant.net/maxquant/
Perseus	(Tyanova et al., 2016b)	https://maxquant.net/perseus/
Oligonucleotides (5' – 3 ')		
tolA Del13 For: GAGTAACAGGCGAACAGTTTTTTGGGAACCGAGAGTGTCAATTCCGGGGATCCGTCGACC-3'	This study	Michael Hensel
tolB Del13 Rev: TTCTTTAATTCCTTTAGTAATCAATTAATTATTATCACAGTGTAGGCTGGAGCTGCTTCG	This study	Michael Hensel
tolA DelCheck For:	This study	Michael Hensel

CGTTAGCGTCCATACCAGTG		
pagL Del For: AAATAACTATTGACATTGAAATGGT GGTGGAGTGTATATGGTGTAGGCT GGAGCTGCTTC	This study	Michael Hensel
pagL Del Rev: CTTCAGCCAGCAACTCGCTAATTGT TATTCAACTTCAGAACATATGAATAT CCTCCTTAG	This study	Michael Hensel
pagL DelCheck For: GTGACATAACAGAAGTGTGG	This study	Michael Hensel
degS Del13 For: TGCTGCTGCCGTTCCCTTTTTTAAC GACGCCTCCATCATGATTCCGGGGA TCCGTCGACC	This study	Michael Hensel
degS Del13 Rev: CTAGTCGTCGAACGACGCGGGCAAAAG CCGCGTCGTTTTATGTAGGCTGGAGCT GCTTCG	This study	Michael Hensel
degS DelCheck For: GAAATGCGCAAAGTGATGGC	This study	Michael Hensel
ompA Del13 For: ATGGCGTATTTTGGATGATAACGAGGCG CAAAAAATGAAAATTCCGGGGATCCGTC GACC	This study	Michael Hensel
ompA Del13 Rev: CCCCGCGACGCGGGGTTTTTTATCAGA CGGAAACTTAAGCTGTAGGCTGGAGCT GCTTCG	This study	Michael Hensel
ompA DelCheck For: AGCCGTAGATATCGGTAGAG	This study	Michael Hensel
K1 Red Del: CAGTCATAGCCGAATAGCCT	This study	Michael Hensel