# A systematic review quantifying host feeding patterns of *Culicoides* species responsible for pathogen transmission

Emma L Fairbanks[1,*], Michael J Tildesley[1], and Janet M Daly[2]

[1]The Zeeman Institute for Systems Biology & Infectious Disease Epidemiology Research, Mathematics Institute and School of Life Sciences, University of Warwick, Coventry, CV4 7AL, UK
[2]One Virology, Wolfson Centre for Global Virus Research, School of Veterinary Medicine and Science, University of Nottingham, Loughborough. LE12 5RD, UK
[*]Corresponding author: emma-louise.fairbanks@warwick.ac.uk

## Abstract

*Culicoides* biting midges are significant vectors of various pathogens, impacting both human and animal health globally. Understanding their host feeding patterns is crucial for deepening our understanding of disease transmission dynamics and developing effective control strategies. While several studies have identified the sources of blood meals in *Culicoides*, a quantitative synthesis of their host preferences and the factors influencing these behaviours is lacking. A systematic literature search focused on gathering data on (1) host selection and (2) host preference. For reviewing host selection we focused on studies reporting the identification of blood meal sources in individual *Culicoides*. When reviewing host preference we focused on studies comparing the number of *Culicoides* caught on or nearby different host species at the same location. Analysis revealed that some *Culicoides* species exhibit fixed host preferences, consistently feeding on specific hosts such as cattle and horses, while others display more opportunistic feeding behaviours. Notable variations were observed across different geographic regions. The findings indicate that host availability significantly influences *Culicoides* feeding patterns. This study highlights the complexity of host selection in *Culicoides* biting midges, which has implications for disease transmission. The variability in feeding behaviours underscores the need for regional assessments to inform targeted vector control strategies.

**Key words:** Host preference, host availability, host selection, vector-borne disease, blood meal analysis, disease transmission, arbovirus, vector

# 1 Background

*Culicoides* need a blood meal in order to develop eggs and produce offspring. While feeding on hosts they may act as vectors in the transmission of pathogens. Vector-borne diseases transmitted by *Culicoides* can severely impact human and animal health, as well as economies.

The World Health Organization (WHO) established four criteria for the recognition of a vector in the 1960s: (1) detection of the pathogen in unfed, field-collected insects; (2) demonstration of infection of the insect with the pathogen by feeding on an infectious host or artificial substitute; (3) demonstration of successful transmission of the pathogen from the insect to a susceptible host during feeding; (4) field evidence of association in space and time between the insects and susceptible hosts [1]. The fourth criterion is commonly explored by analysing the blood meals of field caught vectors; if the vectors are feeding on the susceptible hosts then this criterion is met.

Which hosts *Culicoides* feed on is influenced by availability (whether a host is present) and preference (what the vector prefers to feed on). For some vectors, availability has more of an influence on host selection, often referred to as opportunistic feeding. Alternatively, vectors more influenced by host preference are said to have fixed feeding patterns. These feeding behaviours are often explored by analysing the proportion of blood meals which originate from each host. For opportunistic vector species the proportion from each host type can be variable within small geographical regions, whereas the proportions are more consistent for vector species with fixed feeding behaviour.

Host selection can significantly impact disease transmission dynamics. If a vector that is capable of transmitting a pathogen regularly feeds on a species that is susceptible to infection with it, the capacity for disease transmission is much higher than if they rarely feed on that host species. It has long been understood that host selection plays a significant role in vectorial capacity, which estimates the number of new infectious cases that result from a vector feeding on a single infectious host on a single day [2]. Integral to this is the duration of the extrinsic incubation period (EIP), which refers to how long after feeding on an infectious host *Culicoides* become infectious, the probabilities of vector-to-host and host-to-vector transmission, the life expectancy of *Culicoides* and — of relevance here — the frequency with which the *Culicoides* species under consideration takes a blood meal from the host species. The proportion of blood meals from humans, referred to as the human blood index (HBI), is sometimes used to inform mathematical models for disease transmission [3–6].

While previous reviews have summarised the hosts identified in *Culicoides* blood meals [7, 8], there is a notable lack of quantitative reviews on host selection and the factors influencing feeding patterns. This study aims to fill this gap by systematically analysing host selection and host preferences of *Culicoides* species responsible for pathogen transmission. We conducted a comprehensive systematic literature search across multiple databases, focusing on studies that identified blood meal sources at the species level or compared the number of *Culicoides* attracted to different host types. Eligible studies were included based on predefined criteria to ensure robustness and relevance.

# 2 Methods

## 2.1 Systematic literature search

A systematic search (limited to the title and abstract) was performed (by E. L. Fairbanks) using PubMed, bioRvix, medRvix, arXiv, vectorbase, Paperity, Pubmed central and CAB direct on 19/09/2023. The search term was ("culicoides" OR "biting midges") AND ("human blood index" OR "HBI" OR "host preference" OR "trophic preference" OR "meal preference" OR "blood preference" OR "blood meal" OR "blood-meal" OR "meal source" OR "blood source" OR "host blood" OR "anthropophilic index" OR "blood fed" OR "forage" OR "host use" OR "host associations"), adapted from a search term used to identify the HBI of Anopheles malaria vectors [9].

Duplicate studies were removed and the title and abstracts of the remaining articles were read and kept if they referenced the collection of *Culicoides* and were not review articles. The full text of articles which passed this criterion were read. Studies were eligible for data extraction for the blood meal review, to inform feeding ratios, if they identified the blood meals at the individual *Culicoides* level. Studies were eligible for data extraction for the host preference review if they compared the attractiveness of at least two host species by collection of *Culicoides* on or in close proximity to the hosts in the field.

## 2.2 Data extraction and analysis

Studies that reported *C. pallidipennis* were recorded as *C. imicola*, due to the renaming of this vector. Species within the Obsoletus and Variipennis complexes were pooled for analysis because studies which use morphology for species identification would not be able to determine the specific species. The Variipennis complex consists of five subspecies: *C. variipennis*, *C. sonorensis*, *C. occidentalis*, *C. australis* and *C. albertensis*. The Obsoletus complex consists of three species: *C. obsoletus*, *C. scoticus* and *C. montanus*. Other members of the Obsoletus group, *C. chiopterus* and *C. dewulfi*, can be identified morphologically and were therefore analysed separately.

Data analysis was performed using Rstudio [10]. Figures were generated using the *ggplot2* package in Rstudio [10, 11].

### 2.2.1 Blood meal review

For the quantitative blood-meal review, data extracted from each article included: country, location (including latitude and longitude where reported), years sampling took place, trapping methods, *Culicoides* species identification methods, blood meal identification method and, for each *Culicoides* species or complex, the number of identified blood meals and how many where attributed to each host type.

Some studies included collections at more than one location; data from these were extracted separately. If the results reported were for pooled locations these were recorded as one data

set with the mid-point between the locations as the latitude and longitude. If a blood meal was identified as from a particular host species at a location, it was assumed to be present at that location. Therefore, *Culicoides* species that were not reported to have fed on the host at the location were recorded as having 0% of the blood meal coming from the host type. This was included in calculations of the weighted mean, weighted 95% confidence interval and range of the percentage of blood meals from each host type that is susceptible to a disease for which the *Culicoides* species/complex is a competent vector. Sheep and goats were not considered to be susceptible to vesicular stomatitis virus [12].

These summary statistics were calculated for all host species and *Culicoides* species/complex combinations for which a pathogen was identified as infecting both the host and *Culicoides* species and evidence was found of the *Culicoides* species feeding on the host (the host was identified as a blood meal in at least one data point). Results were aggregated and summarised by *Culicoides* species/complex, host species and region (Americas, Asia, Africa, Australia or Europe).

The mean was weighted by the total number of the *Culicoides* species caught for each data point. To estimate the uncertainty associated with the observed proportions of host species in vector blood meals, bootstrap confidence intervals were calculated. This non-parametric approach involved resampling the original data with replacement to create numerous simulated samples. The process was repeated 1,000 times for each combination of vector species, host species and region.

For each bootstrap sample, the proportion of blood meals derived from a specific host relative to the total number of meals analysed was calculated. This step generated a distribution of mean percentages for each host species, from which the $2.5^{th}$ and $97.5^{th}$ percentiles were determined. These percentiles represent the lower and upper bounds of the 95% confidence interval, respectively, providing an estimate of the range within which the true mean percentage likely falls.

### 2.2.2   Host preference review

For the quantitative host-preference review, data extracted included: country, co-ordinates (latitude and longitude, if available), years of collection, trapping methods, *Culicoides* species identification methods, *Culicoides* species, host attractant and the number of the *Culicoides* species caught on the host attractant.

For each study we compared the ratio of the number of each *Culicoides* species caught on each combination of host pairs. Where more than one study investigated a *Culicoides* species preferences for a host pair the mean and range of these ratios was calculated. These results were then compared between host preference studies and to results from the vector blood meal review.

If trapping was performed for more than one distance from the host, data were extracted for the closest distance to the host. This is because as the distance from hosts increases, the

127 number of *Culicoides* caught decreases [13]. The closest location to the host is assumed to
128 most accurately reflect the number biting the host.

# 3 Results

A total of 532 articles were found during the systematic search from Pubmed, bioRvix,
Paperity, Pubmed central and CAB direct. Of these 141 were duplicates. No articles were
found on medRvix, arXiv or vectorbase. Of the 391 unique articles, 308 were excluded during
a first screening of their titles for being reviews or not including collection of *Culicoides*. After
reading the full texts of the remaining 78 articles which could be retrieved a further 23 were
removed for not collecting *Culicoides* or reporting the number of *Culicoides* found to have
fed on each host species. During the screening, five titles were identified that qualified for a
full text read but could not be accessed. In total, data were extracted from 55 articles; 45
analysed in the blood meal review and 11 analysed in the host preference review with one
article containing data used in both reviews. The process of the literature search is described
by the PRISMA flow diagram [14] in Figure 1.

## 3.1 Blood meal review

All data generated from the blood meal systematic search are provided in Supplementary
file 1. Overall, 514 data points were extracted from 45 studies including 124 locations in
26 countries [15–59]. Of these studies, 53% only gathered data from one location Figure 2.
The most common trapping method was light-baited suction traps, used at 77% of locations.
Data inspection revealed no significant differences in blood meal composition across trap-
ping methods, suggesting that the choice of trapping technique did not substantially bias the
results. Polymerase chain reaction (PCR) was used to identify the host from which blood
meals where taken in 87% of studies and was used in all studies conducted and reported after
1992. Alternative methods for identifying blood meals included enzyme-linked immunosor-
bent assay (ELISA) in 4% and precipitin tests in 9% of studies. Of the 45 studies, 24 used
morphology alone to identify *Culicoides* species, whereas 16 also used PCR and 1 also used
PCR and mass spectrometry. Four studies did not specify how the species of *Culicoides* was
identified. The most common geographical area analysed in studies was Europe (n=19),
followed by Africa (n=8), the Americas (n=8), Asia (n=8) and Australia (n=2). There has
been a rapid and substantial increase in the number of studies reported since 2007, especially
in Europe, the Americas and Asia, possibly due to rising concerns around *Culicoides*-borne
pathogens.
159
160 After filtering for disease vectors that fed on competent hosts, 355 data points were analysed.
161 The pathogens and corresponding vectors found to have fed on each host are listed by host
162 species in Table 1.
163
164 There are some pathogens to which vertebrate hosts listed in 1 are known to be susceptible,
165 but these are not included in Table 1. Pathogens are only ascribed to a vertebrate host if
166 there was evidence that a *Culicoides* species/complex associated with transmission of the

Figure 1: PRISMA flow diagram describing the process of the systematic search.

pathogen fed on the susceptible host in the identified blood meals. For example, birds and sloths are known to be susceptible to Oropouche virus [60], goats and buffalo to Tibet orbivirus [61], buffalo, antelope, alpaca (camelid), giraffe, crocodile, rhinoceros and rodents to Shuni virus [62, 63], rodents, sloths, non-human primates, marsupials, cats, anteaters, shuckand raccoons to Leishmania [64–67], rodents and non-human primates to mansonella [68], zebra to equine encephalosis virus [69], sheep and goats for Aino virus [70], elephants, giraffe, swine and camelids to bovine ephemeral fever [71], giraffe to Schmallenberg virus [72], birds to Thimiri virus, as well as camelids to Akabane virus [73] and vesicular stomatitis virus [12].

Some *Culicoides* species have also been associated with transmission of pathogens for which there was no evidence that they had fed on susceptible hosts in the identified blood meals. For example, *C. pallidicornis*, *C. poperinghensis* and *C. impunctatus* for avian haemoparasites [50, 74, 75], *C. paraensis*, *C. zuluensis* and *C. nivosus* for epizootic haemorragic disease virus [76], *C. pycnostictus* and *C. bedfordi* for bluetongue virus [77], *C. leucostictus* for bluetongue virus and epizootic haemorrhagic disease virus [76, 77], *C. endermania* for bluetongue virus, epizootic haemorrhagic disease virus and equine encephalosis virus [78, 79], *C. mahasarakhamense* for Leishmania [40], *C. austeni*, *C. grahamii* and *C. milnei* for mansonella [80] and *C. circumscriptus* for Akabane virus [81].

Figure 2: Summary of data collected from 45 studies for the blood meal review. (a) The number of sites in each study, (b) The method of catching *Culicoides* at each site, (c) the method of identifying the host source of blood meals, (d) the method of identifying *Culicoides* species and (e) the number of studies capturing *Culicoides* each year by geographical region.

| Host | Pathogens | Vectors |
| --- | --- | --- |
| Antelope | BEFV [71], BTV [82], EHDV [83] | *C. imicola* [71, 84, 85], *C. magnus* [76, 77], *C. stellifer* [35, 86] |
| Boar/pig | BEFV [87], VSV [12] | *C. imicola* [71], Variipennis complex [88, 89] |
| Bird | Avian haemoparasites [90], SHUV [62, 91] | *C. arakawae* [92, 93], *C. circumscriptus* [94], *C. crepuscularis* [95], *C. festivipennis* [94, 96, 97], *C. fulvus* [23], *C. kibunensis* [50, 94, 98], Obsoletus complex [94], *C. oxystoma* [23], *C. pictipennis* [50, 74, 94], *C. punctatus* [94], *C. semimaculatus* [50], Variipennis complex [99] |
| Buffalo | AKAV [100], BEFV [71], BTV [101], EHDV [102], leishmania [103], SBV [104] | *C. actoni* [105], *C. asiana* [106], *C. brevitarsis* [71, 76, 107, 108], *C. dewulfi* [109, 110], *C. imicola* [84, 85, 111], *C. lupicaris* [112], Obsoletus complex [111, 113–115], *C. oxystoma* [36, 76, 106, 116, 117], *C. peregrinus* [36, 76, 118, 119], *C. punctatus* [76, 106, 115, 120, 121], *C. shortti* [105] |
| Camelid | BTV [122], EHDV [123], SBV [124] | Obsoletus complex [111, 113–115] |
| Canine | BTV [125], AHSV [126], leishmania [127] | *C. actoni* [105], *C. crepuscularis* [95, 128], *C. furens* [129], *C. imicola* [84, 130], *C. gulbenkiani* [77], *C. oxystoma* [36, 76, 116], *C. peregrinus* [36, 118, 119], *C. stellifer* [86], Variipennis complex [131–134] |
| Cattle | AINOV [70], AKAV [135], BEFV [71], BTV [101], DOUV [70], EHDV [102], leishmania [103], PEAV [70], SBV [136], SHAV [70], SHUV [137], TIBOV [61], TINV [70], VSV [138] | *C. actoni* [105], *C. bolitinos* [76, 139], *C. brevitarsis* [70, 71, 76, 107, 108], *C. chiopterus* [109], *C. coarctatus* [71], C. cornutus [76], *C. crepuscularis* [95, 128], *C. dewulfi* [109, 110], *C. festivipennis* [140], *C. fordae* [36], *C. foxi* [141], *C. fulvus* [142], *C. furens* [129], *C. haematopotus* [8], *C. imicola* [70, 71, 84, 85, 143], *C. impunctatus* [144], *C. insignis* [141, 145, 146], *C. jacobsoni* [70, 106, 147, 148], *C. kingi* [149, 150], *C. lupicaris* [112], *C. magnus* [76, 77], *C. milnei* [45, 77], *C. newsteadi* [85, 151], Obsoletus complex [111, 113–115], *C. onderstepoortensis* [76], *C. orientalis* [152], *C. oxystoma* [36, 76, 106, 116, 117], *C. peregrinus* [36, 76, 118, 119], *C. pulicaris* [85, 153, 154], *C. punctatus* [76, 106, 115, 120, 121], *C. pusillus* [155, 156], *C. shortti* [105], *C. stellifer* [8, 35, 86], Variipennis complex [8, 88, 89, 99, 131–134, 157, 158] |

| | | |
|---|---|---|
| Deer | BEFV [87], BTV [159], EHDV [102], SBV [160], VSV [161] | *C. chiopterus* [109], *C. debilipalpis* [95, 162], *C. imicola* [71, 84, 85], *C. impunctatus* [144], *C. haematopotus* [8], *C. insignis* [146], *C. newsteadi* [85, 151], Obsoletus complex [113–115], *C. pulicaris* [85, 153, 154], *C. punctatus* [76, 115, 120], *C. pusillus* [155, 156], *C. spinosus* [163], *C. stellifer* [8, 35, 86], Variipennis complex [88, 89, 133, 134, 157, 158], *C. venustus* [8, 35] |
| Donkey | AHSV [126], EEV [69] | *C. bolitinos* [164, 165], *C. imicola* [130], *C. leucostictus* [166], Obsoletus complex [167], *C. onderstepoortensis* [165] |
| Elephant | BTV [168], SBV [72] | *C. dewulfi* [109, 110], Obsoletus complex [111, 113–115] |
| Goat | AKAV [135], BEFV [87], BTV [101], EHDV [102], leishmania [103], SABOV [70]. SBV [136] | *C. chiopterus* [109], *C. fordae* [36], *C. fulvus* [142], *C. imicola* [71, 84, 85, 111, 143], *C. lupicaris* [112], *C. newsteadi* [85, 151], Obsoletus complex [111, 113–115], *C. oxystoma* [36, 76, 106, 116, 117], *C. peregrinus* [36, 76, 118], *C. pulicaris* [85, 153, 154], *C. punctatus* [76, 106, 115, 120, 121], *C. schultzei* [81, 169], *C. shortti* [105], Variipennis complex [8, 131–134, 157, 158] |
| Hippopotamus | BEFV [71] | *C. imicola* [71] |
| Horse | AHSV [126], AKAV [100], EEV [69, 170], PEAV [70], SHUV [171], VSV [172] | *C. bolitinos* [164, 165], *C. imicola* [70, 130], *C. leucostictus* [166], Obsoletus complex [167], *C. oxystoma* [106, 116, 117], *C. pulicaris* [167], *C. punctatus* [106, 121], *C. schultzei* [81], *C. stellifer* [8], Variipennis complex [8, 88, 89, 99] |
| Human | Leishmania [103], mansonella [68], OROV [173], VSV [172] | *C. furens* [129, 174], *C. huffi* [36, 40], *C. insignis* [141], *C. oxystoma* [36], *C. paraensis* [175], *C. peregrinus* [36, 118], *C. stellifer* [8] |
| Other primate | OROV [60] | *C. paraensis* [175] |
| Rabbit/Hare | BUTV [70], leishmania [176], | Variipennis complex [70, 131, 132] |
| Raccoon | VSV [161] | *C. stellifer* [8] |
| Rhinoceros | AHSV [177], AKAV [177], BTV [168, 177], EHDV [177] | *C. imicola* [84, 85, 130], *C. punctatus* [76, 106, 120, 121] |
| Rodent | BTV [178], OROV [179], VSV [172] | C. paraensis [175], C. stellifer [8, 86] |
| Sheep | AKAV [135], BEFV [87], BTV [101], EHDV [180], SBV [136] | *C. actoni* [105], *C. chiopterus* [109], *C. cornutus* [76], *C. dewulfi* [109, 110], *C. festivipennis* [140], *C. fulvus* [142], *C. imicola* [71, 84, 85, 111], *C. impunctatus* [144], *C. lupicaris* [112], *C. magnus* [76, 77], *C. milnei* [45, 77], *C. newsteadi* [85, 151], *C. peregrinus* [76, 119], *C. pulicaris* [85, 153, 154], *C. punctatus* [76, 106, 115, 120, 121], *C. stellifer* [35, 86], Obsoletus complex [111, 113–115], *C. oxystoma* [76, 106, 116, 117], Variipennis complex [8, 133, 134, 157] |

| Zebra | AHSV [126] | *C. leucostictus* [166] |

Table 1: Host species, pathogens and corresponding vectors identified in 45 articles describing blood meals. Pathogens considered were African horse sickness virus (AHSV), Aino virus (ANIOV), Akabane virus (AKAV), avian haemoparasites, bluetongue virus (BTV), bovine ephemeral fever virus (BEFV), Buttonwillow virus (BUTV), Douglas virus (DOUV), epizootic haemorrhagic disease virus (EHDV), equine encephalosis virus (EEV), Leishmania, Mansonella, Oropouche virus (OROV), Peaton virus (PEAV), Sabo virus (SABOV), Schmallenberg virus (SBV), Shamonda virus (SHAV), Shuni virus (SHUV), Tibet orbivirus (TIBOV), Tinaroo virus (TINV) and vesicular stomatitis virus (VSV).

Table 2 presents the total number of the *Culicoides* species caught and the weighted mean, weighted 95% confidence interval and range of the blood index for susceptible hosts which they were found to have taken a blood meal from (Table 1) with data for at least 4 locations for each region. These summary statistics are provided for all *Culicoides* species and corresponding susceptible hosts in Table 1 for each region in Supplementary file 2.

Bluetongue virus has been recovered from wild rodents, however their role as virus reservoirs and potential implication in transmission is unclear [178]. Six studies reported *Culicoides* species susceptible to bluetongue virus feeding on rodents in the Americas and Europe (*C. crepuscularis*, *C. stellifer*, Obsoletus complex, *C. punctatus* and *C. debilipalpis*). Considering studies with more than four of the *Culicoides* species with blood meals identified, we found that less than 3% of these vector species had fed on rodents. This is lower than most susceptible host species, including antelope, buffalo, camelids, cattle, deer, elephants, goats, rhinoceros and sheep. Canids were fed on by competent bluetongue virus *Culicoides* species in nine locations, however similarly to rodents, the percentage of blood meals belonging to this host species is relatively low.

*Culicoides* species exhibit diverse host preferences, ranging from highly specific to opportunistic feeding behaviours. Some species demonstrate a strong preference for birds, such as *C. crepuscularis*, *C. festivipennis*, and *C. circumscriptus*, with weighted means for the percentage of blood meals taken from birds $\geq 95\%$. Conversely, other species show a clear mammalian preference, exemplified by the Variipennis complex and *C. punctatus*, with weighted means for the percentage of blood meals from non-mammalian hosts $\leq 5\%$. However, several species display more opportunistic host selection behaviours. The Obsoletus complex, *C. kibunensis*, and *C. pictipennis*, for instance, exhibit a broad range in the percentage of blood meals from birds across different locations, indicating flexibility in host choice.

Within mammalian-feeding species, further preferences are observed. Some species show a fixed feeding behaviour for specific mammals, such as *C. debilipalpis*, with weighted mean and range for the percentage of blood meals from deer of 95.5% and 91.7–100%, respectively. In contrast, other species demonstrate more opportunistic feeding within the mammalian class. Members of the Variipennis and Obsoletus complexes, for example, have been observed feeding on a variety of mammals including cattle, deer, and horses, suggesting a broader host range within mammals.

While dogs have been reported to become infected with African horse sickness virus [126], their role in the transmission cycle may be more complex. It is possible that dog infections occur through non-vector routes, such as ingestion of infected meat from horse carcasses [126]. We identified two studies which reported African horse sickness virus in the competent *Culicoides* vector species *C. imicola* and *C. oxystoma*, with 1% and 9% having fed on canines, respectively. This is substantially lower than percentages of blood meals belonging to horses for these species.

Table 2: The total number of each *Culicoides* species with blood meals identified, the number of locations they were collected and the weighted mean, weighted 95% confidence interval and range of the percentage of blood meals from each susceptible host they were found to feed on given the *Culicoides* species was caught in at least 4 locations in a region.

| Host | Area | Species | Locations | Total *Culicoides* | Weighted mean | Weighted 95% CI | Range |
|------|------|---------|-----------|-------------------|---------------|-----------------|-------|
| Bird | Americas | *C. crepuscularis* | 6 | 43 | 95.3 | [91.7,98.3] | [90.0,100] |
| Bird | Europe | *C. festivipennis* | 7 | 24 | 100 | [100,100] | [100,100] |
| Bird | Europe | *C. circumscriptus* | 5 | 19 | 94.7 | [85.0,100] | [75.0,100] |
| Bird | Europe | *C. kibunensis* | 4 | 133 | 55.3 | [29.3,81.8] | [12.5,92.9] |
| Bird | Europe | *C. pictipennis* | 4 | 19 | 26.3 | [6.7,76.7] | [6.7,100] |
| Bird | Europe | *C. punctatus* | 4 | 555 | 2.3 | [0.2,6.9] | [0.2,13.6] |
| Cattle | Americas | Variipennis complex | 4 | 360 | 50.8 | [23.3,79.6] | [11.7,100] |
| Cattle | Asia | *C. peregrinus* | 9 | 32 | 78.1 | [61.1,91.7] | [25.0,100] |
| Cattle | Asia | *C. shortti* | 4 | 12 | 75.0 | [55.0,100] | [40.0,100] |
| Cattle | Asia | *C. oxystoma* | 8 | 85 | 24.7 | [15.6,39.5] | [15.6,100] |
| Cattle | Australia | *C. brevitarsis* | 5 | 285 | 99.6 | [99.3,99.9] | [99.3,100] |
| Cattle | Europe | *C. dewulfi* | 5 | 28 | 92.9 | [87.3,98.0] | [83.3,100] |
| Cattle | Europe | *C. lupicaris* | 7 | 17 | 88.2 | [76.2,100] | [66.7,100] |
| Cattle | Europe | *C. chiopterus* | 10 | 97 | 85.6 | [71.3,97.8] | [44.4,100] |
| Cattle | Europe | *C. punctatus* | 12 | 578 | 83.0 | [75.5,87.0] | [27.3,100] |
| Cattle | Europe | *C. pulicaris* | 8 | 114 | 76.3 | [49.8,91.3] | [9.1,100] |
| Cattle | Europe | *C. newsteadi* | 6 | 15 | 73.3 | [46.7,100] | [20.0,100] |
| Cattle | Europe | Obsoletus complex | 14 | 443 | 37.7 | [17.5,59.6] | [1.7,100] |
| Deer | Americas | *C. debilipalpis* | 6 | 89 | 95.5 | [93.7,97.1] | [91.7,100] |
| Deer | Americas | *C. stellifer* | 6 | 3594 | 84.1 | [70.9,90.3] | [10.5,100] |
| Deer | Americas | *C. pusillus* | 4 | 26 | 46.2 | [37.5,68.8] | [37.5,100] |
| Deer | Americas | Variipennis complex | 4 | 354 | 21.8 | [2.3,41.2] | [2.3,100] |
| Deer | Europe | Obsoletus complex | 6 | 163 | 35.0 | [4.4,66.7] | [2.4,100] |
| Deer | Europe | *C. punctatus* | 5 | 52 | 25.0 | [2.6,61.1] | [2.6,100] |
| Horse | Africa | *C. imicola* | 6 | 306 | 38.9 | [29.3,63.9] | [29.3,100] |
| Horse | Europe | Obsoletus complex | 11 | 404 | 48.8 | [27.6,68.0] | [2.4,100] |
| Horse | Europe | *C. punctatus* | 8 | 588 | 10.9 | [2.5,30.2] | [1.7,100] |
| Goat | Asia | *C. peregrinus* | 4 | 11 | 36.4 | [27.1,64.6] | [25.0,100] |

Figure 3: Summary of data collected from 11 studies for the host preference review. (a) The method of catching *Culicoides* in each study, (b) the method of identifying *Culicoides* species and (c) the number of studies capturing *Culicoides* each year by country.

## 3.2  Host preference review

Overall, 252 data points were extracted from 11 studies including locations in 8 countries [13, 57, 181–189]. Each study only collected data from one location (Figure 3). *Culicoides* collection methods included box traps (such as drop traps or stables used to trap insects for collection by aspiration), aspiration directly from host species, sticky traps attached to host species and net sweeping close to host species. Of the 11 studies, 8 used morphology alone to identify *Culicoides* species, whereas 2 also used PCR and 1 did not specify the method. The most common geographical regions analysed in studies was Europe (n=7), followed by Africa (n=1), the Americas (n=1), Asia (n=1) and Australia (n=1). Most studies were performed in the Netherlands (n=4). Like the blood meal analysis there has been a substantial increase in the number of studies in recent years.

Table 3 shows the results from host-preference studies for host pairs with at least two studies comparing their preferences for a *Culicoides* species. This result is provided for all host pairs

investigated for each *Culicoides* species in Supplementary file 3.

The mean ratios of the numbers of *Culicoides* caught on two host species varied from 0.48–91. Our analysis reveals varying degrees of consistency in host preference ratios across studies. For instance, the cattle-to-horse preference ratios for *C. chiopterus*, *C. impunctatus*, and *C. pallidicornis* show relatively low variability (ranges of 5.65-5.85, 2.25-2.50, and 4.82-4.83, respectively). In contrast, other species exhibit markedly higher variability in host preferences; for example, the cattle-to-sheep ratio for *C. achrayi* ranges from 1.32 to 100.5 across different studies. The level of variability does not appear to be associated with the location. When *Culicoides* species that were investigated multiple times at the same location, some studies found similar catch ratios between the hosts and others did not.

In this review, two studies reported results for all subspecies of the Obsoletus group pooled. These studies are reported separately to the Obsoletus complex. We observe that results for comparing cattle to sheep (Table 3) are not consistent between the Obsoletus group and Obsoletus complex, with the Obsoletus group showing less preference for cattle. This might be explained by the wide range of results found for *C. dewulfi*, suggesting it has very opportunistic feeding behaviours.

Table 4 shows the summary statistics for host pairs and corresponding *Culicoides* species with more than four data points in the blood meal review (included in Table 2). We observe that the trends from the host preference review generally agree with the blood meal review, in terms of the host most commonly selected. For *C. punctatus* the blood meal review found the mean percentage of blood meals belonging to cattle and horses was 83.0% (95% CI: [75.5,87.0]) and 10.9% (95% CI: [2.5,30.2]), respectively. However, the host preference review did not find evidence of a preference for cattle with the ratio of the number of vectors found on cattle compared to horses ranging from 0.88 to 2.59.

Table 3: The mean and range of ratios of the number of vectors caught for host attractants for *Culicoides* species and host pairs investigated in at least 2 host preference studies. Studies marked with a † show the same study site was used for both studies. All data summarised in this table was from Europe.

| Host pair | Species | Studies | Total *Culicoides* | Mean ratio | Ratio range |
|---|---|---|---|---|---|
| Cattle:Horse | C. fascipennis | 2 † | 102 | 15.12 | [1.21,29.00] |
| Cattle:Horse | C. chiopterus | 2 | 4883 | 5.75 | [5.65,5.85] |
| Cattle:Horse | C. pallidicornis | 2 † | 204 | 4.83 | [4.82,4.83] |
| Cattle:Horse | C. achrayi | 2 † | 1855 | 2.44 | [1.79,3.10] |
| Cattle:Horse | C. impunctatus | 2 † | 20 | 2.38 | [2.25,2.50] |
| Cattle:Horse | C. pulicaris | 2 † | 429 | 1.96 | [0.49,3.44] |
| Cattle:Horse | C. stigma | 2 † | 237 | 1.80 | [0.85,2.75] |
| Cattle:Horse | C. punctatus | 2 † | 3841 | 1.74 | [0.88,2.59] |
| Cattle:Horse | Obsoletus complex | 3 | 7398 | 0.96 | [0.02, 1.66] |
| Cattle:Horse | C. dewulfi | 3 | 5451 | 0.63 | [0.01,1.18] |
| Cattle:Horse | C. heliophilus | 2 † | 36 | 0.48 | [0.14,0.82] |
| Cattle:Sheep | C. achrayi | 4 | 984 | 42.05 | [1.32,100.5] |
| Cattle:Sheep | C. pulicaris | 3 | 666 | 30.01 | [12.53,55.00] |
| Cattle:Sheep | C. punctatus | 4 | 3792 | 22.55 | [1.68,50.65] |
| Cattle:Sheep | C. chiopterus | 4 | 7566 | 14.14 | [1.24,24.21] |
| Cattle:Sheep | C. dewulfi | 5 | 8976 | 5.91 | [0.33,13.66] |
| Cattle:Sheep | Obsoletus complex | 5 | 15509 | 5.30 | [2.40,11.43] |
| Cattle:Sheep | C. impunctatus | 3 | 82 | 2.79 | [1.25,3.75] |
| Cattle:Sheep | Obsoletus group | 2 | 5471 | 1.03 | [1.00,1.06] |
| Horse:Sheep | Obsoletus complex | 2 | 2304 | 91.31 | [2.13,180.50] |
| Horse:Sheep | C. dewulfi | 2 | 2141 | 36.79 | [6.91,66.67] |

Table 4: Comparison of host selection ratios between host preference studies and blood meal analyses for *Culicoides* species. For the host preference studies the mean ratio of the number of vectors caught on different host attractants is given. For the blood meal studies the ratio of the means of the percentage of blood meals from each host type is given. Only comparisons with $\geq 4$ blood meal data points per host are included.

| Host pair | Species | Region | Host preference review | Blood meal review | | |
|---|---|---|---|---|---|---|
| | | | Mean ratio | Mean % host 1 | Mean % host 2 | Ratio |
| Cattle:Goat | C. peregrinus | Asia | 43.86 | 78.1 | 36.4 | 2.15 |
| Cattle:Horse | C. punctatus | Europe | 1.74 | 83.0 | 10.9 | 7.61 |
| Cattle:Horse | Obsoletus complex | Europe | 0.96 | 37.7 | 48.8 | 0.77 |

## 4 Discussion

Our results indicate a complex pattern of host preference among different *Culicoides* species, reflecting both species-specific and regional variations in host selection. This systematic review revealed that while some *Culicoides* species demonstrated a strong preference for specific host types, others appeared more opportunistic, suggesting that host availability might play a significant role in their feeding behaviour. Awareness of this variability is crucial for understanding the dynamics of disease transmission and vector management.

The data collected in this review provide evidence of association in space and time between the insects and susceptible hosts, supporting their potential roles as vectors of pathogens and demonstrating one of the WHO criteria for vector recognition [1]. For some of the pathogens considered in this study not all of the other criteria are met. For example, transmission to hosts from infectious vectors has not been demonstrated for bovine ephemeral fever virus [190, 191].

Studies have shown that sensory structures of *Culicoides* species (palps and antennae) can determine whether they prefer feeding on birds or mammals [192, 193] or are opportunistic feeders [194]. Studies have also found evidence of *Culicoides* feeding on other insects including common fruit flies (*Drosophila melanogaster*) [21] and *Aedes*, *Culex* and *Anopheles* mosquitoes [195]. If *Culicoides* are regularly feeding on other insects which feed on the blood of vertebrate hosts, for example mosquitoes, the origin of the blood meals present may reflect the foraging behaviours of the mosquitoes, rather than of the *Culicoides*.

Our results mostly agree with Martínez-de la Puente et al. [7], who reviewed literature to find that most European *Culicoides* species feed on multiple vertebrate species, but present clear preferences for birds or mammals. However, our results for the Obsoletus group found the mean percentage of blood meals that came from birds in locations where birds were present was 30%, with a maximum of 50% in one study. This contradicts results from Martínez-de la Puente et al. [7] which suggested that all members of the Obsoletus group and complex were

mammalphilic. Hopken et al. [38] found that many American *Culicoides* species were more opportunistic feeders than was thought prior to the study which identified blood meals of *Culicoides* caught in North America.

In our blood meal review we included data points for all competent *Culicoides* species and susceptible hosts at a location. This approach considers that in different settings *Culicoides* may feed on different hosts. Previous studies have suggested that the obsoletus complex prefers cattle to horses [184], however our review found that where horses are present, horse blood constitutes the highest percentage of the blood meals. These results do not necessarily contradict each other, but show the importance of considering the other host species in a setting and how the impact of this may vary for each species.

This variation in the origin of blood meals for some species was also observed in the host preference review. While similar results were found when comparing the number of a *Culicoides* species caught on or nearby some host pairs, these were often not consistent between studies, including studies at the same location. Although trends observed in the blood meal and host preference reviews generally agreed, this was not always the case. This highlights the need for a deeper understanding of the interaction between host preference and host availability for opportunistic feeders and its possible impact on disease transmission.

Host preference has been shown to be inherited for *Aedes* [196] and *Anopheles* [197] mosquitoes, with offspring more likely to select the same host as generations before them. If this trend is also true for *Culicoides* then this could contribute to the variations in results found in the host preference. If some hosts are usually less present at the location of the trial, the host preferences of local vectors could be more skewed towards host species that are more abundant in the area. More investigation into the innate host preferences of *Culicoides* spp. is required, as well as knowledge of the host distributions around trial sites to verify the reliability of these results.

The number of *Culicoides* collected from light traps near hosts is less than the number collected by aspiration directly from the hosts [13, 198]. Elbers and Gonzales [13] showed that the number of *Culicoides* caught by net sweeping significantly declined with increasing distance from hosts. In contrast, the number of *Culicoides* caught increased with increasing distance of hosts from the light trap.

Other viruses than those reported have been associated with *Culicoides* (including Ingwavuma virus, Jatobal virus, Manzanilla virus, Iquitos virus, Madre de Dios virus, Pintupo virus, Sango virus and Sathuperi virus [70]). However transmission has not been linked to specific species, often due to testing pooled *Culicoides* species.

# 5  Conclusion

Our findings suggest that future research should explore the drivers of host selection in more detail. We recommend the data on host selection needs to be assessed for a range of set-

tings and at different locations in order to understand the foraging behaviours of *Culicoides* species. When using these data to parameterise mathematical models for disease transmission, the influence on host selection due to the variation in the settings in which hosts are found should be considered. Parameterising a model using data from one study is likely unreliable. Additionally, blood meal analysis with the distributions of host species usually found at a location are potentially more reliable than host preference studies comparing the number of *Culicoides* found on or near a host if one of the hosts is not usually present at the location.

These insights are instrumental for vector control strategies. Understanding that some vectors have fixed preferences helps in targeting control measures more effectively, focusing on the hosts that are most likely to be bitten in a particular setting. However, the adaptability in host preference observed in some species also warns against a one-size-fits-all approach to vector management. It emphasises the need for regional assessments of vector behaviour and host interactions to tailor interventions appropriately.

**Data availability**   Supplementary file 1 contains all data gathered during the blood-meal systematic search. Supplementary file 2 contains summary statistics for the blood meal review for all *Culicoides* species and corresponding susceptible hosts. Supplementary file 3 contains results for all host pairs investigated for each *Culicoides* species in the host-preference study.

**Conflict of interest**   The authors declare no conflict of interest.

**Ethical approval**   The authors confirm that the ethical policies of the journal, as noted on the journal's author guidelines page, have been adhered to. No ethical approval was required.

**Author contributions**
**ELF:** Conceptualisation, Methodology, Software, Programming, Validation, Formal analysis, Investigation, Data curation, Writing - Original Draft, Writing - Review & Editing, Visualisation, Project administration, Funding acquisition; **MJT:** Methodology, Writing - Review & Editing, Visualisation, Project administration, Funding acquisition; **JMD:** Writing - Original Draft, Writing - Review & Editing, Visualisation, Funding acquisition

**Financial support**   The authors were supported by the Horserace Betting Levy Board (vet/prj/809). MJT is funded on a joint BBSRC/EEID grant (BB/T004312/1).

# References

[1] World Health Organisation. *Arboviruses and human disease : report of a WHO scientific group*, 1966. Available at: https://iris.who.int/handle/10665/40664 [Accessed 13-July-2024].

[2] C Garrett-Jones. The human blood index of malaria vectors in relation to epidemiological assessment. *Bull World Health Organ*, 30(2):241, 1964.

[3] EL Davis, TD Hollingsworth, and MJ Keeling. An analytically tractable, age-structured model of the impact of vector control on mosquito-transmitted infections. *PLOS Comput Biol*, 20(3):e1011440, 2024. doi: 10.1371/journal.pcbi.1011440.

[4] N Chitnis, T Smith, and R Steketee. A mathematical model for the dynamics of malaria in mosquitoes feeding on a heterogeneous host population. *J Biol Dyn*, 2(3): 259–285, 2008. doi: 10.1080/17513750701769857.

[5] V Ermert, AH Fink, AE Jones, and AP Morse. Development of a new version of the Liverpool Malaria Model. I. Refining the parameter settings and mathematical formulation of basic processes based on a literature review. *Malar J*, 10:35, 2011. doi: 10.1186/1475-2875-10-35.

[6] MG Basáñez, K Razali, A Renz, and D Kelly. Density-dependent host choice by disease vectors: epidemiological implications of the ideal free distribution. *Trans R Soc Trop Med Hyg*, 101(3):256–269, 2007. doi: 10.1016/j.trstmh.2006.08.009.

[7] J Martínez-de la Puente, J Figuerola, and R Soriguer. Fur or feather? feeding preferences of species of *Culicoides* biting midges in Europe. *Trends Parasitol*, 31(1):16–22, 2015. doi: 10.1016/j.pt.2014.11.002.

[8] BL McGregor, PT Shults, and EG McDermott. A review of the vector status of North American *Culicoides* (Diptera: Ceratopogonidae) for bluetongue virus, epizootic hemorrhagic disease virus, and other arboviruses of concern. *Curr Trop Med Rep*, 9 (4):130–139, 2022. doi: 10.1007/s40475-022-00263-8.

[9] Y Wang, N Chitnis, and EL Fairbanks. Optimizing malaria vector control in the Greater Mekong Subregion: a systematic review and mathematical modelling study to identify desirable intervention characteristics. *Parasit Vector*, 17(1):162, 2024. doi: 10.1186/s13071-024-06234-4.

[10] RStudio Team. *Rstudio: Integrated Development Environment for R*. RStudio, PBC., Boston, MA, 2020. URL `http://www.rstudio.com/`.

[11] H Wickham, W Chang, L Henry, TL Pedersen, K Takahashi, C Wilke, K Woo, H Yutani, D Dunnington, and RStudio. *Create Elegant Data Visualisations Using the Grammar of Graphics*, 2022. URL `https://ggplot2.tidyverse.org`.

[12] Beverly Schmitt. Vesicular stomatitis. *Vet Clin Food Anim Pract*, 18(3):453–459, 2002. doi: 10.1016/S0749-0720(02)00031-2.

[13] ARW Elbers and JL Gonzales. *Culicoides* (Diptera: Ceratopogonidae) abundance is influenced by livestock host species and distance to hosts at the micro landscape scale. *Insects*, 14(7):637, 2023. doi: 10.3390/insects14070637.

[14] A Liberati, DG Altman, J Tetzlaff, C Mulrow, PC Gøtzsche, JPA Ioannidis, M Clarke, PJ Devereaux, J Kleijnen, and D Moher. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Ann Intern Med*, 151(4):W65–94, 2009. doi: 10.7326/0003-4819-151-4-200908180-00136.

[15] BL McGregor and A Lewis. Host associations of *Culicoides* biting midges in Northeastern Kansas, USA. *Animals*, 13(15):2504, 2023. doi: 10.3390/ani13152504.

[16] S Kar, B Mondal, A Pal, R Harsha, and A Mazumdar. Blood meal analysis of *Culicoides* species associated with livestock in West Bengal, India. *Med Vet Entomol*, 36 (4):503–510, 2022. doi: 10.1111/mve.12588.

[17] A Vasić, N Zdravković, D Aniță, J Bojkovski, M Marinov, A Mathis, M Niculaua, EL Oșlobanu, I Pavlović, D Petrić, et al. Species diversity, host preference and arbovirus detection of *Culicoides* (Diptera: Ceratopogonidae) in south-eastern Serbia. *Parasit Vector*, 12(1):61, 2019. doi: 10.1186/s13071-019-3292-3.

[18] KE Sloyer, C Acevedo, SM Wisely, and ND Burkett-Cadena. Host associations of biting midges (Diptera: Ceratopogonidae: *Culicoides*) at deer farms in Florida, USA. *J Med Entomol*, 60(3):518–526, 2023. doi: 10.1093/jme/tjad036.

[19] KE Sloyer, C Acevedo, AE Runkel, and ND Burkett-Cadena. Host associations of biting midges (Diptera: Ceratopogonidae: *Culicoides*) near sentinel chicken surveillance locations in Florida, USA. *J Am Mosq Control Assoc*, 35(3):200–206, 2019. doi: 10.2987/19-6834.1.

[20] ME England, P Pearce-Kelly, VA Brugman, S King, S Gubbins, F Sach, CJ Sanders, NJ Masters, E Denison, and S Carpenter. *Culicoides* species composition and molecular identification of host blood meals at two zoos in the UK. *Parasit Vector*, 13(1):139, 2020. doi: 10.1186/s13071-020-04018-0.

[21] D Slama, N Haouas, H Mezhoud, H Babba, and E Chaker. Blood meal analysis of *Culicoides* (Diptera: Ceratopogonidae) in central Tunisia. *PLoS One*, 10(3):e0120528, 2015. doi: 10.1371/journal.pone.0120528.

[22] Z Kasičová, A Schreiberová, A Kimáková, and A Kočišová. Blood meal analysis: host-feeding patterns of biting midges (Diptera, Ceratopogonidae, *Culicoides* Latreille) in Slovakia. *Parasite*, 28, 2021. doi: 10.1051/parasite/2021058.

[23] S Sunantaraporn, T Hortiwakul, K Kraivichian, P Siriyasatien, and N Brownell. Molecular identification of host blood meals and detection of blood parasites in *Culicoides* Latreille (Diptera: Ceratopogonidae) collected from Phatthalung province, Southern Thailand. *Insects*, 13(10):912, 2022. doi: 10.3390/insects13100912.

[24] P Jomkumsing, A Surapinit, T Saengpara, and P Pramual. Genetic variation, DNA barcoding and blood meal identification of *Culicoides* Latreille biting midges (Diptera: Ceratopogonidae) in Thailand. *Acta Tropica*, 217:105866, 2021. doi: 10.1016/j.actatropica.2021.105866.

[25] BL McGregor, T Stenn, KA Sayler, EM Blosser, JK Blackburn, SM Wisely, and ND Burkett-Cadena. Host use patterns of *Culicoides* spp. biting midges at a big game preserve in Florida, USA, and implications for the transmission of orbiviruses. *Med Vet Entomol*, 33(1):110–120, 2019. doi: 10.1111/mve.12331.

[26] A Tomazatos, H Jöst, J Schulze, M Spînu, J Schmidt-Chanasit, D Cadar, and R Lühken. Blood-meal analysis of *Culicoides* (Diptera: Ceratopogonidae) reveals a broad host range and new species records for Romania. *Parasit Vector*, 13(1):79, 2020. doi: 10.1186/s13071-020-3938-1.

[27] LPC Carvalho, AMP Júnior, PFM de Paulo, GS Silva, G da Silva Costa, MT de Souza Freitas, FAC Pessoa, and JF Medeiros. DNA-based blood meal analysis of *Culicoides* (Diptera: Ceratopogonidae) species from Jamari National Forest, Southwestern Amazon, Brazil. *Acta Tropica*, 221:106025, 2021. doi: 10.1016/j.actatropica.2021.106025.

[28] SB Lassen, SA Nielsen, and M Kristensen. Identity and diversity of blood meal hosts of biting midges (Diptera: Ceratopogonidae: *Culicoides* Latreille) in Denmark. *Parasit Vector*, 5:143, 2012. doi: 10.1186/1756-3305-5-143.

[29] L Hadj-Henni, T De Meulemeester, J Depaquit, P Noël, A Germain, R Helder, and D Augot. Comparison of vertebrate cytochrome b and prepronociceptin for blood meal analyses in *Culicoides. Front Vet Sci*, 2:15, 2015. doi: 10.3389/fvets.2015.00015.

[30] B Gomontean, K Vaisusuk, W Chatan, K Wongpakam, P Sankul, L Lachanthuek, Mintara, I Thanee, and P Pramual. Diversity, abundance and host blood meal analysis of Culicoides Latreille (Diptera: Ceratopogonidae) from cattle pens in different land use types from Thailand. *Insects*, 14(7):574, 2023. doi: 10.3390/insects14070574.

[31] MA González, D Bravo-Barriga, MA Rodríguez-Sosa, J Rueda, E Frontera, and PM Alarcón-Elbal. Species diversity, habitat distribution, and blood meal analysis of haematophagous dipterans collected by CDC-UV light traps in the Dominican Republic. *Pathogens*, 11(7):714, 2022. doi: 10.3390/pathogens11070714.

[32] K Kamyingkird, S Choocherd, W Chimnoi, N Klinkaew, C Kengradomkij, P Phoosangwalthong, N Thammasonthijarern, K Pattanatanang, T Inpankaew, J Phasuk, and B Nimsuphan. Molecular identification of *Culicoides* species and host preference blood meal in the African horse sickness outbreak-affected area in Hua Hin district, Prachuap Khiri Khan province, Thailand. *Insects*, 14(4):369, 2023. doi: 10.3390/insects14040369.

[33] MA Riddin, Gert Johannes Venter, K Labuschagne, and Martin H Villet. Bloodmeal analysis in *Culicoides* midges collected near horses, donkeys and zebras in the Eastern Cape, South Africa. *Med Vet Entomol*, 33(4):467–475, 2019. doi: 10.1111/mve.12381.

[34] SB Lassen, SA Nielsen, H Skovgård, and M Kristensen. Molecular identification of bloodmeals from biting midges (Diptera: Ceratopogonidae: *Culicoides* Latreille) in Denmark. *Parasitol Res*, 108:823–829, 2011. doi: 10.1007/s00436-010-2123-4.

[35] BL McGregor, KE Sloyer, KA Sayler, O Goodfriend, JMC Krauer, C Acevedo, X Zhang, D Mathias, SM Wisely, and ND Burkett-Cadena. Field data implicating *Culicoides stellifer* and *Culicoides venustus* (Diptera: Ceratopogonidae) as vectors of epizootic hemorrhagic disease virus. *Parasit Vector*, 12:258, 2019. doi: 10.1186/s13071-019-3514-8.

[36] N Songumpai, C Promrangsee, P Noopetch, P Siriyasatien, and K Preativatanyou. First evidence of co-circulation of emerging *Leishmania martiniquensis*, *Leishmania orientalis*, and *Crithidia* sp. in *Culicoides* biting midges (Diptera: Ceratopogonidae), the putative vectors for autochthonous transmission in Southern Thailand. *Trop Med Infect Dis*, 7(11):379, 2022. doi: 10.3390/tropicalmed7110379.

[37] MR Van der Saag, X Gu, MP Ward, and PD Kirkland. Development and evaluation of real-time pCR assays for bloodmeal identification in *Culicoides* midges. *Med Vet Entomol*, 30(2):155–165, 2016. doi: 10.1111/mve.12163.

[38] MW Hopken, BM Ryan, KP Huyvaert, and AJ Piaggio. Picky eaters are rare: DNA-based blood meal analysis of *Culicoides* (Diptera: Ceratopogonidae) species from the United States. *Parasit Vector*, 10(1):169, 2017. doi: 10.1186/s13071-017-2099-3.

[39] J Snyman, LP Snyman, K Labuschagne, GJ Venter, and M Venter. The utilisation of CytB and COI barcodes for the identification of bloodmeals and *Culicoides* species (Diptera: Ceratopogonidae) reveals a variety of novel wildlife hosts in South Africa. *Acta Tropica*, 219:105913, 2021. doi: 10.1016/j.actatropica.2021.105913.

[40] S Sunantaraporn, A Thepparat, A Phumee, S Sor-Suwan, R Boonserm, G Bellis, and P Siriyasatien. *Culicoides* Latreille (Diptera: Ceratopogonidae) as potential vectors for *Leishmania martiniquensis* and *Trypanosoma* sp. in northern Thailand. *PLoS Negl Trop Dis*, 15(12):e0010014, 2021. doi: 10.1371/journal.pntd.0010014.

[41] S Talavera, F MUÑOZ-MUÑOZ, M Verdún, N Pujol, and N Pagès. Revealing potential bridge vectors for BTV and SBV: a study on *Culicoides* blood feeding preferences in natural ecosystems in Spain. *Med Vet Entomol*, 32(1):35–40, 2018. doi: 10.1111/mve.12263.

[42] C Garros, L Gardes, X Allene, I Rakotoarivony, E Viennet, S Rossi, and T Balenghien. Adaptation of a species-specific multiplex PCR assay for the identification of blood meal source in *Culicoides* (Ceratopogonidae: Diptera): applications on Palaearctic biting midge species, vectors of orbiviruses. *Infect Genet Evol*, 11(5):1103–1110, 2011. doi: 10.1016/j.meegid.2011.04.002.

[43] MT Bakhoum, M Fall, MT Seck, L Gardes, AG Fall, M Diop, I Mall, T Balenghien, T Baldet, G Gimonneau, C Garros, and J Bouyer. Foraging range of arthropods with veterinary interest: new insights for Afrotropical *Culicoides* biting midges (Diptera: Ceratopogonidae) using the ring method. *Acta tropica*, 157:59–67, 2016. doi: 10.1016/j.actatropica.2016.01.023.

[44] C Ninio, D Augot, JC Delecolle, B Dufour, and J Depaquit. Contribution to the knowledge of *Culicoides* (Diptera: Ceratopogonidae) host preferences in France. *Parasitol Res*, 108:657–663, 2011. doi: 10.1007/s00436-010-2110-9.

[45] AR Walker and FG Davies. A preliminary survey of the epidemiology of bluetongue in Kenya. *J Hyg (Lond)*, 69(1):47–60, 1971. doi: 10.1017/s0022172400021239.

[46] S Díaz-Sánchez, A Hernández-Jarguín, A Torina, IG Fernández de Mera, A Estrada-Peña, M Villar, F La Russa, V Blanda, J Vicente, S Caracappa, C Gortazar, and J de la Fuente. Biotic and abiotic factors shape the microbiota of wild-caught populations of the arbovirus vector *Culicoides imicola*. *Insect Mol Biol*, 27(6):847–861, 2018. doi: 10.1111/imb.12526.

[47] D Santiago-Alarcon, P Havelka, E Pineda, G Segelbacher, and HM Schaefer. Urban forests as hubs for novel zoonosis: blood meal analysis, seasonal variation in *Culicoides* (Diptera: Ceratopogonidae) vectors, and avian haemosporidians. *Parasitol*, 140(14): 1799–1810, 2013. doi: 10.1017/S0031182013001285.

[48] JH Calvo, B Berzal, C Calvete, MA Miranda, R Estrada, and J Lucientes. Host feeding patterns of *Culicoides* species (Diptera: Ceratopogonidae) within the Picos de Europa National Park in northern Spain. *Bull Entomol Res*, 102(6):692–697, 2012. doi: 10.1017/S0007485312000284.

[49] A Bobeva, P Zehtindjiev, M Ilieva, D Dimitrov, A Mathis, and S Bensch. Host preferences of ornithophilic biting midges of the genus *Culicoides* in the Eastern Balkans. *Med Vet Entomol*, 29(3):290–296, 2015. doi: 10.1111/mve.12108.

[50] D Santiago-Alarcon, P Havelka, HM Schaefer, and G Segelbacher. Bloodmeal analysis reveals avian *Plasmodium* infections and broad host preferences of *Culicoides* (Diptera: Ceratopogonidae) vectors. *Plos One*, 7(2):e31098, 2012. doi: 10.1371/journal.pone.0031098.

[51] J Mart'inez-de la Puente, J Martínez, M Ferraguti, A Morales-de la Nuez, N Castro, and J Figuerola. Genetic characterization and molecular identification of the bloodmeal sources of the potential bluetongue vector *Culicoides obsoletus* in the Canary Islands, Spain. *Parasit Vector*, 5:147, 2012. doi: 10.1186/1756-3305-5-147.

[52] L Xue, W Lin, X Yuanyuan, L Jiangxin, D Qinglang, and H Xiaohui. Molecular identification of gastric blood source of *Culicoides* in Ganshui Town, Qijiang, Chongqing, China. *Chin J Zoonoses*, 37(12):1064–1070, 2021. doi: 10.3969/j.issn.1002-2694.2021.00.155.

[53] E Videvall, S Bensch, M Ander, J Chirico, R Sigvald, and R Ignell. Molecular identification of bloodmeals and species composition in *Culicoides* biting midges. *Med Vet Entomol*, 27(1):104–112, 2013. doi: 10.1111/j.1365-2915.2012.01038.x.

[54] A Voller, DE Bidwell, et al. The enzyme-linked immunosorbent assay (ELISA) test for the identification of blood-meals of haematophagous insects. *Bull Entomol Res*, 76 (2):321–330, 1986. doi: 10.1017/S0007485300014796.

[55] A Blackwell and W Mordue. Identification of bloodmeals of the Scottish biting midge, *Culicoides impunctatus*, by indirect enzyme-linked immunosorbent assay (ELISA). *Medical Vet Entomol*, 8(1):20–24, 1994. doi: 10.1111/j.1365-2915.1994.tb00378.x.

[56] MB Nathan. Transmission of the human filarial parasite *Mansonella ozzardi* by *Culicoides phlebotomus* (Williston)(Diptera: Ceratopogonidae) in coastal north Trinidad. *Bull Entomol Res*, 71(1):97–106, 1981. doi: 10.1017/S0007485300051063.

[57] BH Kay, PFL Boreham, AL Dyce, and HA Standfast. Blood feeding of biting midges (Diptera: Ceratopogonidae) at Kowanyama, Cape York Peninsula, North Queensland. *Aust J Entomol*, 17(2):145–149, 1978. doi: 10.1111/j.1440-6055.1978.tb02222.x.

[58] Y Braverman, PFL Boreham, and R Galum. The origin of blood meals of female *Culicoides pallidipennis* trapped in a sheepfold in Israel. *J Med Entomol*, 8(4):379–381, 1971. doi: 10.1093/jmedent/8.4.379.

[59] EM Nevill and Dora Anderson. Host preferences of *Culicoides* midges (Diptera: Ceratopogonidae) in South Africa as determined by precipitin tests and light trap catches. 1972.

[60] D Romero-Alvarez and LE Escobar. Emergent viruses in America: The case of Oropouche virus. *Int J Infect Dis*, 73:98, 2018. doi: 10.1016/j.ijid.2018.04.3644.

[61] J Wang, H Li, Y He, Y Zhou, A Xin, D Liao, and J Meng. Isolation of Tibet orbivirus from culicoides and associated infections in livestock in Yunnan, China. *Virol J*, 14 (1):105, 2017. doi: 10.1186/s12985-017-0774-9.

[62] J Steyn, P Motlou, Charmaine Van E, M Pretorius, VI Stivaktas, J Williams, LP Snyman, PE Buss, B Beechler, A Jolles, E Perez-Martin, JG Myburgh, J Steyl, and M Venter. Shuni virus in wildlife and nonequine domestic animals, South Africa. *Emerg Infect Dis*, 26(7):1521–1525, 2020. doi: 10.3201/eid2607.190770.

[63] BM McIntosh. Susceptibility of some African wild rodents to infection with various arthropod-borne viruses. *Trans R Soc Trop Med Hyg*, 55(1):63–68, 1961. doi: 10.1016/0035-9203(61)90041-4.

[64] A Özbilgin, İ Çavuş, A Yıldırım, and C Gündüz. Do the rodents have a role in transmission of cutaneous leishmaniasis in Turkey? *Mikrobiyol Bul*, 52(3):259–272, 2018. doi: 10.5578/mb.66828.

[65] R Lainson, RR Braga, AAA De Souza, MM Povoa, EAY Ishikawa, and FT Silveira. Leishmania (Viannia) shawi sp. n., a parasite of monkeys, sloths and procyonids in Amazonian Brazil. *Ann Parasitol Hum Comp*, 64(3):200–207, 1989. doi: 10.1051/parasite/1989643200.

[66] A Montoya, LP de Quadros, M Mateo, L Hernández, R Gálvez, G Alcántara, R Checa, MÁ Jiménez, C Chicharro, I Cruz, and G Miró. Leishmania infantum infection in Bennett's Wallabies (Macropus rufogriseus rufogriseus) in a Spanish wildlife park. *J Zoo Wildl Med*, 47(2):586–593, 2016. doi: 10.1638/2014-0216.1.

[67] I Azami-Conesa, M Teresa Gómez-Muñoz, and RA Martínez-Díaz. A systematic review (1990–2021) of wild animals infected with zoonotic Leishmania. *Microorganisms*, 9(5): 1101, 2021. doi: 10.3390/microorganisms9051101.

[68] O Mediannikov and S Ranque. Mansonellosis, the most neglected human filariasis. *New Microbes New Infect*, 26:S19–S22, 2018. doi: 10.1016/j.nmni.2018.08.016.

[69] R Williams, DH Du Plessis, and W Van Wyngaardt. Group-reactive ELISAs for detecting antibodies to African horsesickness and equine encephalosis viruses in horse, donkey, and zebra sera. *J Vet Diagn Invest*, 5(1):3–7, 1993. doi: 10.1177/104063879300500102.

[70] F Sick, M Beer, H Kampen, and K Wernike. *Culicoides* biting midges—underestimated vectors for arboviruses of public health and veterinary importance. *Viruses*, 11(4):376, 2019. doi: 10.3390/v11040376.

[71] PJ Walker and E Klement. Epidemiology and control of bovine ephemeral fever. *Vet Res*, 46:124, 2015. doi: 10.1186/s13567-015-0262-4.

[72] FM Molenaar, SA La Rocca, M Khatri, J Lopez, F Steinbach, and A Dastjerdi. Exposure of Asian elephants and other exotic ungulates to Schmallenberg virus. *PLoS One*, 10(8):e0135532, 2015. doi: 10.1371/journal.pone.0135532.

[73] R Saidi, F DOĞAN, VS Ataseven, and Y Ergün. Antibody detection against Akabane (AKA) and Bluetongue (BT) viruses in Algerian dromedary camels. *Turk J Vet Anim Sci*, 44(1):142–145, 2020. doi: 10.3906/vet-1905-44.

[74] R Žiegytė, R Bernotienė, and V Palinauskas. *Culicoides segnis* and *Culicoides pictipennis* biting midges (Diptera, Ceratopogonidae), new reported vectors of *Haemoproteus* parasites. *Microorganisms*, 10(5):898, 2022. doi: 10.3390/microorganisms10050898.

[75] R Žiegytė, V Palinauskas, R Bernotienė, TA Iezhova, and G Valkiūnas. *Haemoproteus minutus* and *Haemoproteus belopolskyi* (Haemoproteidae): complete sporogony in the biting midge *Culicoides impunctatus* (Ceratopogonidae), with implications on epidemiology of haemoproteosis. *Exp Parasitol*, 145:74–79, 2014. doi: 10.1016/j.exppara.2014.07.014.

[76] G Savini, A Afonso, P Mellor, I Aradaib, H Yadin, M Sanaa, W Wilson, F Monaco, and M Domingo. Epizootic haemorrhagic disease. *Res Vet Sci*, 91(1):1–17, 2011. doi: 10.1016/j.rvsc.2011.05.004.

[77] JT Paweska, GJ Venter, and PS Mellor. Vector competence of South African *Culicoides* species for bluetongue virus serotype 1 (BTV-1) with special reference to the effect of temperature on the rate of virus replication in *C. imicola* and *C. bolitinos*. *Med Vet Entomol*, 16(1):10–21, 2002. doi: 10.1046/j.1365-2915.2002.00334.x.

[78] SL Vigil, MG Ruder, D Shaw, J Wlodkowski, K Garrett, M Walter, and JL Corn. Apparent range expansion of *Culicoides* (Hoffmania) *insignis* (Diptera: Ceratopogonidae) in the southeastern United States. *J Med Entomol*, 55(4):1043–1046, 2018. doi: 10.1093/jme/tjy036.

[79] JT Paweska and GJ Venter. Vector competence of *Culicoides* species and the seroprevalence of homologous neutralizing antibody in horses for six serotypes of equine encephalosis virus (EEV) in South Africa. *Med Vet Entomol*, 18(4):398–407, 2004. doi: 10.1111/j.0269-283X.2004.00524.x.

[80] WL Nicholas. The bionomics of *Culicoides austeni*, vector of *Acanthocheilonema* perstans in the rainforest of the British Cameroons, together with notes on *C. grahamii* and other species which may be vectors in the same area. *Ann Trop Med Parasitol*, 47 (2):187–206, 1953. doi: 10.1080/00034983.1953.11685560.

[81] SB Dağalp, B Dik, Ft Doğan, TA Farzani, VS Ataseven, G Acar, İ Şahinkesen, and A Özkul. Akabane virus infection in Eastern Mediterranean Region in Turkey: *Culicoides* (Diptera: Ceratopogonidae) as a possible vector. *Trop Anim Health Prod*, 53 (2):231, 2021. doi: 10.1007/s11250-021-02661-y.

[82] WO Neitz. The blesbuck (*Damaliscus albifrons*) as a carrier of heartwater and blue tongue. *J S Afr Vet Assoc*, 4(1):24–26, 1933.

[83] GA Chalmers, HN Vance, and GJ Mitchell. An outbreak of epizootic hemorrhagic disease in wild ungulates in Alberta. 1964.

[84] GJ Venter. *Culicoides* spp.(Diptera: Ceratopogonidae) as vectors of bluetongue virus in South Africa–A review. 2015. doi: 10.12834/VetIt.505.2436.2.

[85] M Quaglia, C Foxi, G Satta, G Puggioni, R Bechere, M De Ascentis, SG d'Alessio, M Spedicato, A Leone, M Pisciella, O Portanti, L Teodori, L Di Gialleonardo, C Cammá, G Savini, and M Goffredo. Culicoides species responsible for the transmission of epizootic haemorrhagic disease virus (EHDV) serotype 8 in Italy. *Veterinaria Italiana*, 59(1), 2023. doi: 10.12834/VetIt.3347.22208.1.

[86] GR Mullen, RH Jones, Y Braverman, and KE Nusbaum. Laboratory infections of *Culicoides debilipalpis* and *C. stellifer* (Diptera: Ceratopogonidae) with bluetongue virus. *Prog Clin Biol Res*, 178:239–243, 1985.

[87] JY Yeh and YJ Ga. Role of cervids in the epidemiology of bovine ephemeral fever virus infection in the Republic of Korea: A cross-sectional retrospective study. *Vet Med Sci*, 9(1):301–306, 2023. doi: 10.1002/vms3.970.

[88] P Rozo-Lopez, B Londono-Renteria, and BS Drolet. Venereal transmission of vesicular stomatitis virus by *Culicoides sonorensis* midges. *Pathogens*, 9(4):316, 2020. doi: 10.3390/pathogens9040316.

[89] P Rozo-Lopez, B Londono-Renteria, and BS Drolet. Impacts of infectious dose, feeding behavior, and age of *Culicoides sonorensis* biting midges on infection dynamics of vesicular stomatitis virus. *Pathogens*, 10(7):816, 2021. doi: 10.3390/pathogens10070816.

[90] GF Bennett, M Whiteway, and C Woodworth-Lynas. A host-parasite catalogue of the avian haematozoa. 1982.

[91] G Valkiūnas and TA Iezhova. Insights into the biology of Leucocytozoon species (Haemosporida, Leucocytozoidae): why is there slow research progress on agents of leucocytozoonosis? *Microorganisms*, 11(5):1251, 2023. doi: 10.3390/microorganisms11051251.

[92] M Inumaru, K Nakamura, T Odagawa, Mo Suzuki, K Murata, and Y Sato. The first detection of avian haemosporidia from *Culicoides* biting midges in Japan, with notes on potential vector species and the transmission cycle. *Vet Parasitol Reg Stud Reports*, 39:100840, 2023. doi: 10.1016/j.vprsr.2023.100840.

[93] CY Yu, JS Wang, and CC Yeh. *Culicoides arakawae* (Diptera: Ceratopogonidae) population succession in relation to leucocytozoonosis prevalence on a chicken farm in Taiwan. *Vet Parasitol*, 93(2):113–120, 2000. doi: 10.1016/s0304-4017(00)00362-9.

[94] R Bernotienė, R Žiegytė, G Vaitkutė, and G Valkiūnas. Identification of a new vector species of avian haemoproteids, with a description of methodology for the determination of natural vectors of haemosporidian parasites. *Parasit Vector*, 12(1):307, 2019. doi: 10.1186/s13071-019-3559-8.

[95] ME Becker, J Roberts, ME Schroeder, G Gentry, and LD Foil. Prospective study of epizootic hemorrhagic disease virus and bluetongue virus transmission in captive ruminants. *J Med Entomol*, 57(4):1277–1285, 2020. doi: 10.1093/jme/tjaa027.

[96] R Žiegytė, V Palinauskas, and R Bernotienė. Natural vector of avian *Haemoproteus asymmetricus* parasite and factors altering the spread of infection. *Insects*, 14(12):926, 2023. doi: 10.3390/insects14120926.

[97] J Martínez-de la Puente, J Martinez, J Rivero-de Aguilar, J Herrero, and S Merino. On the specificity of avian blood parasites: revealing specific and generalist relationships between haemosporidians and biting midges. *Mol Ecol*, 20(15):3275–3287, 2011. doi: 10.1111/j.1365-294X.2011.05136.x.

[98] R Žiegytė, E Platonova, E Kinderis, A Mukhin, V Palinauskas, and R Bernotienė. *Culicoides* biting midges involved in transmission of haemoproteids. *Parasit Vector*, 14:27, 2021. doi: 10.1186/s13071-020-04516-1.

[99] TWR Möhlmann, J Oymans, PJ Wichgers Schreur, CJM Koenraadt, J Kortekaas, and CBF Vogels. Vector competence of biting midges and mosquitoes for Shuni virus. *PLoS Negl Trop Dis*, 12(12):e0006993, 2018. doi: 10.1371/journal.pntd.0006609.

[100] PD Kirkland. Akabane and bovine ephemeral fever virus infections. *Vet Clin Food Anim Pract*, 18(3):501–514, 2002. doi: 10.1016/s0749-0720(02)00026-9.

[101] T Sohail, T Yaqub, T Abbas, M Rabbani, J Nazir, SM Maqbool, S Yaqub, M Habib, N Mukhtar, M Shahbaz, MY Zahoor, and MZ Shabbir. Seroprevalence of Bluetongue virus in small and large ruminants in Punjab province, Pakistan. *Acta Trop*, 189:22–29, 2019. doi: 10.1016/j.actatropica.2018.09.020.

[102] L Jiménez-Cabello, S Utrilla-Trigo, G Lorenzo, J Ortego, and E Calvo-Pinilla. Epizootic hemorrhagic disease virus: Current knowledge and emerging perspectives. *Microorganisms*, 11(5):1339, 2023. doi: 10.3390/microorganisms11051339.

[103] NR Bhattarai, G Van der Auwera, S Rijal, A Picado, N Speybroeck, B Khanal, S De Doncker, ML Das, B Ostyn, C Davies, M Coosemans, D Berkvens, M Boelaert, and JC Dujardin. Domestic animals and epidemiology of visceral leishmaniasis, Nepal. *Emerg Tnfect Dis*, 16(2):231–237, 2010. doi: 10.3201/eid1602.090623.

[104] AK Azkur, H Albayrak, A Risvanli, Z Pestil, E Ozan, O Yılmaz, S Tonbak, A Cavunt, H Kadı, HC Macun, D Acar, E Özenç, S Alparslan, and H Bulut. Antibodies to Schmallenberg virus in domestic livestock in Turkey. *Trop Anim Health Prod*, 45: 1825–1828, 2013. doi: 10.1007/s11250-013-0415-2.

[105] S Kar, B Mondal, A Pal, and A Mazumdar. Molecular identification of *Culicoides oxystoma* and *Culicoides actoni* vectors of bluetongue virus. *Med Vet Entomol*, 37(3): 534–541, 2023. doi: 10.1111/mve.12651.

[106] T Yanase, T Kato, Y Hayama, H Shirafuji, M Yamakawa, and S Tanaka. Oral susceptibility of Japanese *Culicoides* (Diptera: Ceratopogonidae) species to Akabane virus. *J Med Entomol*, 56(2):533–539, 2019. doi: 10.1093/jme/tjy201.

[107] WT Tay, PJ Kerr, and LS Jermiin. Population genetic structure and potential incursion pathways of the bluetongue virus vector *Culicoides brevitarsis* (Diptera: Ceratopogonidae) in Australia. *PLoS One*, 11(1):e0146699, 2016. doi: 10.1371/journal.pone.0146699.

[108] TD St George, HA Standfast, and DH Cybinski. Isolations of Akabane virus from sentinel cattle and *Culicoides brevitarsis*. *Australian Veterinary Journal*, 54(12):558–561, 1978. doi: 10.1111/j.1751-0813.1978.tb02412.x.

[109] AI Paslaru, A Mathis, P Torgerson, and E Veronesi. Vector competence of pre-alpine *Culicoides* (Diptera: Ceratopogonidae) for bluetongue virus serotypes 1, 4 and 8. *Parasit Vector*, 11:466, 2018. doi: 10.1186/s13071-018-3050-y.

[110] LD Rasmussen, C Kirkeby, R Bødker, B Kristensen, TB Rasmussen, GJ Belsham, and A Bøtner. Rapid spread of Schmallenberg virus-infected biting midges (*Culicoides* spp.) across Denmark in 2012. *Transbound Emerg Dis*, 61(1):12–16, 2014. doi: 10.1111/tbed.12189.

[111] N Pagès, S Talavera, M Verdún, N Pujol, M Valle, A Bensaid, and J Pujols. Schmallenberg virus detection in *Culicoides* biting midges in Spain: First laboratory evidence for highly efficient infection of *Culicoides* of the Obsoletus complex and *Culicoides imicola*. *Transbound Emerg Dis*, 65(1):e1–e6, 2018. doi: 10.1111/tbed.12653.

[112] P Romon, M Higuera, JC Delécolle, T Baldet, G Aduriz, and A Goldarazena. Phenology and attraction of potential *Culicoides* vectors of bluetongue virus in Basque Country (northern Spain). *Vet Parasitol*, 186(3-4):415–424, 2012. doi: 10.1016/j.vetpar.2011.11.023.

[113] R Meiswinkel, T Baldet, R De Deken, W Takken, JC Delécolle, and PS Mellor. The 2006 outbreak of bluetongue in northern Europe—the entomological perspective. *Prev Vet Med*, 87(1–2):55–63, 2008. doi: 10.1016/j.prevetmed.2008.06.005.

[114] S Carpenter, HL Lunt, D Arav, GJ Venter, and PS Mellor. Oral susceptibility to bluetongue virus of *Culicoides* (Diptera: Ceratopogonidae) from the United Kingdom. *J Med Entomol*, 43(1):73–78, 2006. doi: 10.1093/jmedent/43.1.73.

[115] M Larska, L Lechowski, M Grochowska, and JF Żmudziński. Detection of the Schmallenberg virus in nulliparous *Culicoides obsoletus/scoticus* complex and *C. punctatus*—the possibility of transovarial virus transmission in the midge population and of a new vector. *Vet Microbiol*, 166(3-4):467–473, 2013. doi: 10.1016/j.vetmic.2013.07.015.

[116] M Fall, AG Fall, MT Seck, J Bouyer, M Diarra, T Balenghien, C Garros, MT Bakhoum, O Faye, T Baldet, and G Gimonneau. Circadian activity of *Culicoides oxystoma* (Diptera: Ceratopogonidae), potential vector of bluetongue and African horse sickness viruses in the Niayes area, Senegal. *Parasitol Res*, 114:3151–3158, 2015. doi: 10.1007/s00436-015-4534-8.

[117] H Kurogi, K Akiba, Y Inaba, and M Matumoto. Isolation of Akabane virus from the biting midge *Culicoides oxystoma* in Japan. *Vet Microbiol*, 15(3):243–248, 1987. doi: 10.1016/0378-1135(87)90078-2.

[118] S Kaewmee, C Mano, T Phanitchakun, R Ampol, T Yasanga, U Pattanawong, A Junkum, P Siriyasatien, PA Bates, and N Jariyapan. Natural infection with *Leishmania* (Mundinia) *martiniquensis* supports *Culicoides peregrinus* (Diptera: Ceratopogonidae) as a potential vector of leishmaniasis and characterization of a Crithidia sp. isolated from the midges. *Front Microbiol*, 14:1235254, 2023. doi: 10.3389/fmicb.2023.1235254.

[119] P Banerjee, A Sarkar, and A Mazumdar. Effect of substrate salinity and pH on life history traits of the bluetongue virus vector *Culicoides peregrinus*. *Bull Entomol Res*, 113(6):829–837, 2023. doi: 10.1017/S0007485323000512.

[120] R Meiswinkel, F Scolamacchia, M Dik, J Mudde, E Dijkstra, IJK Van Der Ven, and ARW Elbers. The Mondrian matrix: *Culicoides* biting midge abundance and seasonal incidence during the 2006–2008 epidemic of bluetongue in the Netherlands. *Med Vet Entomol*, 28(1):10–20, 2014. doi: 10.1111/mve.12013.

[121] T Yanase, Y Hayama, H Shirafuji, T Tsutsui, and Y Terada. Surveillance of *Culicoides* biting midges in northern Honshu, Japan, during the period of Akabane virus spread. *J Vet Med Sci*, 81(10):1496–1503, 2019. doi: 10.1292/jvms.19-0303.

[122] CA Batten, B Harif, MR Henstock, S Ghizlane, L Edwards, C Loutfi, CAL Oura, and M El Harrak. Experimental infection of camels with bluetongue virus. *Res Vet Sci*, 90 (3):533–535, 2011. doi: 10.1016/j.rvsc.2010.07.013.

[123] GM Cosseddu, B Doumbia, M Scacchia, C Pinoni, A Di Provvido, A Polci, K Isselmou, A Di Gennaro, M Spedicato, I Carmine, G Savini, A Capobianco Dondona, F Iapaolo, F Valleriani, AB El Mamy, Y Barry, and F Monaco. Sero-surveillance of emerging viral diseases in camels and cattle in Nouakchott, Mauritania: an abattoir study. *Trop Anim Health Prod*, 53:195, 2021. doi: 10.1007/s11250-021-02636-z.

[124] C Schulz, M Beer, and B Hoffmann. Schmallenberg virus infection in South American camelids: Field and experimental investigations. *Vet Microbiol*, 180(3-4):171–179, 2015. doi: 10.1016/j.vetmic.2015.08.024.

[125] CAL Oura and M El Harrak. Midge-transmitted bluetongue in domestic dogs. *Epidemiol Infect*, 139(9):1396–1400, 2011. doi: 10.1017/S0950268810002396.

[126] SJ Dennis, AE Meyers, II Hitzeroth, and EP Rybicki. African horse sickness: A review of current understanding and vaccine development. *Viruses*, 11(9):844, 2019. doi: 10.3390/v11090844.

[127] F Dantas-Torres. The role of dogs as reservoirs of Leishmania parasites, with emphasis on Leishmania (Leishmania) infantum and Leishmania (Viannia) braziliensis. *Vet Parasitol*, 149(3-4):139–146, 2007. doi: 10.1016/j.vetpar.2007.07.007.

[128] ME Becker, WK Reeves, SK Dejean, MP Emery, EN Ostlund, and LD Foil. Detection of bluetongue virus RNA in field-collected *Culicoides* spp. (Diptera: Ceratopogonidae) following the discovery of bluetongue virus serotype 1 in white-tailed deer and cattle in Louisiana. *J Med Entomol*, 47(2):269–273, 2014. doi: 10.1603/me09211.

[129] JJ Ríos-Tostado, Hi Castillo-Ureta, EH Torres-Montoya, JI Torres-Avendaño, V Olimón-Andalón, CE Romero-Higareda, G Silva-Hidalgo, and JM Zazueta-Moreno. Molecular detection of *Leishmania* (*L.*) *mexicana* (Kinetoplastida: Trypanostomatidae) DNA in *Culicoides furens* (Diptera: Ceratopogonidae) from an area with autochthonous canine leishmaniasis in Northwestern Mexico. *Acta Parasitol*, 66:1055–1058, 2021. doi: 10.1007/s11686-021-00335-1.

[130] PS Mellor and C Hamblin. African horse sickness. *Vet Res*, 35(4):445–466, 2004. doi: 10.1051/vetres:2004021.

[131] T Becvar, B Vojtkova, P Siriyasatien, J Votypka, D Modry, P Jahn, P Bates, S Carpenter, P Volf, and J Sadlova. Experimental transmission of Leishmania (Mundinia) parasites by biting midges (Diptera: Ceratopogonidae). *PLoS Pathogens*, 17(6):e1009654, 2021. doi: 10.1371/journal.ppat.1009654.

[132] V Seblova, J Sadlova, B Vojtkova, J Votypka, S Carpenter, PA Bates, and P Volf. The biting midge *Culicoides sonorensis* (Diptera: Ceratopogonidae) is capable of developing late stage infections of *Leishmania enriettii*. *PLoS Negl Trop Dis*, 9(9):e0004060, 2015. doi: 10.1371/journal.pntd.0004060.

[133] WL Kramer, RH Jones, FR Holbrook, TE Walton, and CH Calisher. Isolation of arboviruses from *Culicoides* midges (Diptera: Ceratopogonidae) in Colorado during an epizootic of vesicular stomatitis New Jersey. *J Med Entomol*, 27(4):487–493, 1990. doi: 10.1093/jmedent/27.4.487.

[134] R Morales-Hojas, M Hinsley, IM Armean, R Silk, LE Harrup, A Gonzalez-Uriarte, E Veronesi, L Campbell, D Nayduch, C Saski, WJ Tabachnick, P Kersey, S Carpenter, and M Fife. The genome of the biting midge *Culicoides sonorensis* and gene expression analyses of vector competence for bluetongue virus. *BMC Genomics*, 19:624, 2018. doi: 10.1186/s12864-018-5014-1.

[135] Y_ Inaba, H Kurogi, and T Omori. Akabane disease: epizootic abortion, premature birth, stillbirth and congenital arthrogryposis-hydranencephaly in cattle, sheep and goats caused by Akabane virus. 1975. doi: 10.1111/j.1751-0813.1975.tb09397.x.

[136] V Gülyaz. A brief review of Schmallenberg virus. *J Etlik Vet Microbiol*, 38(2):59–61, 2011.

[137] OR Causey, GE Kemp, CE Causey, and VH Lee. Isolations of Simbu-group viruses in Ibadan, Nigeria 1964–69, including the new types Sango, Shamonda, Sabo and Shuni. *Ann Trop Med Parasitol*, 66(3):357–362, 1972. doi: 10.1080/00034983.1972.11686835.

[138] JR Mohler. *Vesicular stomatitis of horses and cattle*. Number 662. US Department of Agriculture, 1918.

[139] GJ Venter, JT Paweska, AA Van Dijk, PS Mellor, and WJ Tabachnick. Vector competence of *Culicoides bolitinos* and *C. imicola* for South African bluetongue virus serotypes 1, 3 and 4. *Med Vet Entomol*, 12(4):378–385, 1998. doi: 10.1046/j.1365-2915.1998.00116.x.

[140] D Muz, B Dik, and MN Muz. The investigation of *Culicoides* (Diptera: Ceratopogonidae) species and bluetongue virus and Schmallenberg virus in Northwest Türkiye. *Trop Anim Health Prod*, 55(1):39, 2023. doi: 10.1007/s11250-023-03454-1.

[141] JMM Rebêlo, BL Rodrigues, MCA Bandeira, JLP Moraes, RS Fonteles, and SRF Pereira. Detection of *Leishmania amazonensis* and *Leishmania braziliensis* in *Culicoides* (Diptera, Ceratopogonidae) in an endemic area of cutaneous leishmaniasis in the Brazilian Amazonia. *J Vector Ecol*, 41(2):303–308, 2016. doi: 10.1111/jvec.12227.

[142] Y Fujisawa, T Homat, A Thepparat, T Changbunjong, K Sutummaporn, S Kornmatitsuk, and B Kornmatitsuk. DNA barcode identification and molecular detection of bluetongue virus in *Culicoides* biting midges (Diptera: Ceratopogonidae) from western Thailand. *Acta Trop*, 224:106147, 2021. doi: 10.1016/j.actatropica.2021.106147.

[143] VH Lee. Isolation of viruses from field populations of *culicoides* (Diptera: Ceratopogonidae) in Nigeria. *J Med Entomol*, 16(1):76–79, 1979. doi: 10.1093/jmedent/16.1.76.

[144] DM Jennings and PS Mellor. The vector potential of British *Culicoides* species for bluetongue virus. *Vet Microbiol*, 17(1):1–10, 1988. doi: 10.1016/0378-1135(88)90074-0.

[145] MM Ayala, F Díaz, MV Micieli, GR Spinelli, and MM Ronderos. Rapid and efficient detection by PCR of *Culicoides insignis* (Diptera: Ceratopogonidae), the main vector of bluetongue virus (BTV) in the Neotropical region. *J Med Entomol*, 59(4):1211–1216, 2022. doi: 10.1093/jme/tjac065.

[146] BL McGregor, D Erram, BW Alto, JA Lednicky, SM Wisely, and ND Burkett-Cadena. Vector competence of Florida *Culicoides insignis* (Diptera: Ceratopogonidae) for epizootic hemorrhagic disease virus serotype-2. *Viruses*, 13(3):410, 2021. doi: 10.3390/v13030410.

[147] YL Duan, ZX Yang, G Bellis, and L Li. Isolation of Tibet orbivirus from *Culicoides jacobsoni* (Diptera, Ceratopogonidae) in China. *Parasit Vector*, 14(1):432, 2021. doi: 10.1186/s13071-021-04899-9.

[148] YL Duan, L Li, G Bellis, ZX Yang, and HC Li. Detection of bluetongue virus in *Culicoides* spp. in southern Yunnan Province, China. *Parasit Vector*, 14:68, 2021. doi: 10.1186/s13071-020-04518-z.

[149] B Hakima, HS Hwang, and KY Lee. Molecular identification of *Culicoides* (Diptera: Ceratopogonidae) species in Algeria. *Acta Trop*, 202:105261, 2020. doi: 10.1016/j.actatropica.2019.105261.

[150] PS Mellor, R Osborne, and DM Jennings. Isolation of bluetongue and related viruses from *Culicoides* spp. in the Sudan. *Epidemiol Infect*, 93(3):621–628, 1984. doi: 10.1017/s0022172400065190.

[151] C Foxi, G Delrio, G Falchi, MG Marche, G Satta, and L Ruiu. Role of different *Culicoides* vectors (Diptera: Ceratopogonidae) in bluetongue virus transmission and overwintering in Sardinia (italy). *Parasit Vector*, 9:440, 2016. doi: 10.1186/s13071-016-1733-9.

[152] E Mukhopadhyay, S Hazra, GK Saha, and D Banerjee. Altitudinal variation and bioclimatic variables influencing the potential distribution of *Culicoides orientalis* Macfie, 1932, suspected vector of bluetongue virus across the North Eastern Himalayan belt of Sikkim. *Acta Trop*, 176:402–411, 2017. doi: 10.1016/j.actatropica.2017.09.008.

[153] DV Nolan, S Carpenter, J Barber, PS Mellor, JF Dallas, AJ Mordue, and SB Piertney. Rapid diagnostic PCR assays for members of the *Culicoides obsoletus* and *Culicoides pulicaris* species complexes, implicated vectors of bluetongue virus in Europe. *Vet Microbiol*, 124(1-2):82–94, 2007. doi: 10.1016/j.vetmic.2007.03.019.

[154] J Kęsik-Maliszewska, M Larska, ÁB Collins, and J Rola. Post-epidemic distribution of Schmallenberg virus in *Culicoides* arbovirus vectors in Poland. *Viruses*, 11(5):447, 2019. doi: 10.3390/v11050447.

[155] ES Farias, JF Almeida, JW Pereira-Silva, LS Coelho, CM Ríos-Velásquez, SLB Luz, and FAC Pessoa. Diversity of biting midges *Culicoides* (Diptera: Ceratopogonidae), potential vectors of disease, in different environments in an Amazo-

nian rural settlement, Brazil. *Rev Soc Bras Med Trop*, 53:e20200067, 2020. doi: 10.1590/0037-8682-0067-2020.

[156] BA Mullens and CE Dada. Spatial and seasonal distribution of potential vectors of hemorrhagic disease viruses to peninsular bighorn sheep in the Santa Rosa mountains of southern California. *J Wildl Dis*, 28(2):192–205, 1992. doi: 0.7589/0090-3558-28.2.192.

[157] BL McGregor, D Erram, C Acevedo, BW Alto, and ND Burkett-Cadena. Vector competence of *Culicoides sonorensis* (Diptera: Ceratopogonidae) for epizootic hemorrhagic disease virus serotype 2 strains from Canada and Florida. *Viruses*, 11(4):367, 2019. doi: 10.3390/v11040367.

[158] N De Regge, M Madder, I Deblauwe, B Losson, C Fassotte, J Demeulemeester, F Smeets, M Tomme, and AB Cay. Schmallenberg virus circulation in *Culicoides* in Belgium in 2012: field validation of a real time RT-PCR approach to assess virus replication and dissemination in midges. *PloS One*, 9(1):e87005, 2014. doi: 10.1371/journal.pone.0087005.

[159] RA Vosdingh, DO Trainer, and BC Easterday. Experimental bluetongue disease in white-tailed deer. *Can J Comp Med Vet Sci*, 32(1):382, 1968.

[160] E Laloy, E Bréard, C Sailleau, C Viarouge, A Desprat, S Zientara, F Klein, J Hars, and S Rossi. Schmallenberg virus infection among red deer, France, 2010–2012. *Emerg Infect Dis*, 20(1):131, 2014. doi: 10.3201/eid2001.130411.

[161] LARS Karstad and RP Hanson. Vesicular stomatitis in deer. *Am J Vet Res*, 18(66): 162–166, 1957.

[162] KE Sloyer, ND Burkett-Cadena, A Yang, JL Corn, SL Vigil, BL McGregor, SM Wisely, and JK Blackburn. Ecological niche modeling the potential geographic distribution of four *Culicoides* species of veterinary significance in Florida, USA. *PLoS One*, 14(2): e0206648, 2019. doi: 10.1371/journal.pone.0206648.

[163] KE Smith, DE Stallknecht, CT Sewell, EA Rollor, GR Mullen, and RR Anderson. Monitoring of *Culicoides* spp. at a site enzootic for hemorrhagic disease in white-tailed deer in Georgia, USA. *J Wildl Dis*, 32(4):627–642, 1996. doi: 0.7589/0090-3558-32.4. 627.

[164] R Meiswinkel, M Baylis, and K Labuschagne. Stabling and the protection of horses from *Culicoides bolitinos* (Diptera: Ceratopogonidae), a recently identified vector of African horse sickness. *Bull Entomol Res*, 90(6):509–515, 2000. doi: 10.1017/s0007485300000626.

[165] GJ Venter, D Groenewald, E Venter, KG Hermanides, and PG Howell. A comparison of the vector competence of the biting midges, *Culicoides (Avaritia) bolitinos* and *C.(A.) imicola*, for the Bryanston serotype of equine encephalosis virus. *Med Vet Entomol*, 16 (4):372–377, 2002. doi: 10.1046/j.1365-2915.2002.00385.x.

[166] M Goffredo, G Savini, M Quaglia, U Molini, V Federici, M Catalani, O Portanti, V Marini, MA Florentius, A Pini, and M Scacchia. Orbivirus detection from *Culicoides* collected on African horse sickness outbreaks in Namibia. *Veterinaria Italiana*, 51(1): 17–23, 2015. doi: 10.12834/VetIt.275.1016.2.

[167] M Baylis, HEL Hasnaoui, H Bouayoune, J Touti, and PS Mellor. The spatial and seasonal distribution of African horse sickness and its potential *culicoides* vectors in Morocco. *Med Vet Entomol*, 11(3):203–212, 1997. doi: 10.1111/j.1365-2915.1997. tb00397.x.

[168] J Caballero-Gómez, D Cano Terriza, J Pujols, E Martínez-Nevado, MD Carbonell, R Guerra, J Recuero, P Soriano, J Barbero, and I García-Bocanegra. Monitoring of bluetongue virus in zoo animals in Spain, 2007–2019. *Transbound Emerg Dis*, 69(4): 1739–1747, 2022. doi: 10.1111/tbed.14147.

[169] SM Al-Busaidy and PS Mellor. Epidemiology of bluetongue and related orbiviruses in the Sultanate of Oman. *Epidemiol Infect*, 106(1):167–178, 1991. doi: 10.1017/ s0950268800056533.

[170] BJ Erasmus, TF Adelaar, JD Smit, G Lecatsas, and T Toms. The isolation and characterization of equine encephalosis virus. 1970.

[171] TP Motlou, J Williams, and M Venter. Epidemiology of shuni virus in horses in South Africa. *Viruses*, 13(5):937, 2021. doi: 10.3390/v13050937.

[172] GJ Letchworth, LL Rodriguez, and J Del Cbarrera. Vesicular stomatitis. *Vet J*, 157 (3):239–260, 1999. doi: 10.1053/tvjl.1998.0303.

[173] CR Anderson, L Spence, WG Downs, and THG Aitken. Oropouche virus: a new human disease agent from Trinidad, West Indies. *Am J Trop Med Hyg*, 10(4):574–578, 1961.

[174] Robert C Lowrie Jr and Christian Raccurt. *Mansonella ozzardi* in Haiti II. Arthropod vector studies. *Am J Trop Med Hyg*, 30(3):598–603, 1981. doi: 10.4269/ajtmh.1981. 30.598.

[175] LHM Feitoza, LPC de Carvalho, LR da Silva, ACA Meireles, FGF Rios, GS Silva, PFM de Paulo, FAC Pessoa, JF de Medeiros, and GR Julião. Influence of meteorological and seasonal parameters on the activity of *Culicoides paraensis* (Diptera: ceratopogonidae), an annoying anthropophilic biting midge and putative vector of Oropouche virus in Rondônia, Brazilian Amazon. *Acta Trop*, 243:106928, 2023. doi: 10.1016/j.actatropica. 2023.106928.

[176] MV Ortega-García, FrJ Salguero, A Rodríguez-Bertos, I Moreno, N García, T García-Seco, G Luz Torre, L Domínguez, and M Domínguez. A pathological study of Leishmania infantum natural infection in European rabbits (Oryctolagus cuniculus) and Iberian hares (Lepus granatensis). *Transbound Emerg Dis*, 66(6):2474–2481, 2019. doi: 10.1111/tbed.13305.

[177] C Fischer-Tenhagen, C Hamblin, S Quandt, and K Frölich. Serosurvey for selected infectious disease agents in free-ranging black and white rhinoceros in Africa. *J Wildl Dis*, 36(2):316–323, 2000. doi: 10.7589/0090-3558-36.2.316.

[178] R Du Toit. Insect vectors and virus diseases. *J S Afr Vet Assoc*, 26(4):263–268, 1955.

[179] FP Pinheiro, AP Travassos da Rosa, JF Travassos da Rosa, and G Bensabath. An outbreak of Oropouche virus diease in the vicinity of santarem, para, barzil. *Tropenmed Parasitol*, 27(2):213–223, 1976.

[180] TH Noon, SL Wesche, D Cagle, DG Mead, EJ Bicknell, GA Bradley, S Riplog-Peterson, D Edsall, and C Reggiardo. Hemorrhagic disease in bighorn sheep in Arizona. *J Wildl Dis*, 38(1):172–176, 2002. doi: 10.7589/0090-3558-38.1.172.

[181] E Viennet, C Garros, L Gardes, I Rakotoarivony, X Allene, R Lancelot, D Crochet, C Moulia, T Baldet, and T Balenghien. Host preferences of Palaearctic *Culicoides* biting midges: implications for transmission of orbiviruses. *Med Vet Ento*, 27(3):255–266, 2013. doi: 10.1111/j.1365-2915.2012.01042.x.

[182] S Kar, B Mondal, J Ghosh, SM Mazumdar, and A Mazumdar. Host preference of bluetongue virus vectors, *Culicoides* species associated with livestock in West Bengal, India: Potential relevance on bluetongue epidemiology. *Acta Tropica*, 235:106648, 2022. doi: 10.1016/j.actatropica.2022.106648.

[183] ARW Elbers and R Meiswinkel. *Culicoides* (Diptera: Ceratopogonidae) host preferences and biting rates in the Netherlands: comparing cattle, sheep and the black-light suction trap. *Vet Parasitol*, 205(1-2):330–337, 2014. doi: 10.1016/j.vetpar.2014.06.004.

[184] ARW Elbers and R Meiswinkel. *Culicoides* (Diptera: Ceratopogonidae) and livestock in the Netherlands: comparing host preference and attack rates on a Shetland pony, a dairy cow, and a sheep. *J Vector Ecol*, 40(2):308–317, 2015. doi: 10.1111/jvec.12169.

[185] GM Thompson, S Jess, AW Gordon, and AK Murchie. Sticky-trapping biting midges (*Culicoides* spp.) alighting on cattle and sheep: effects of trap colour and evidence for host preference. *Parasitol Res*, 113:3085–3094, 2014. doi: 10.1007/s00436-014-3974-x.

[186] BA Mullens and CE Dada. Insects feeding on desert bighorn sheep, domestic rabbits, and Japanese quail in the Santa Rosa mountains of southern California. *J Wildl Dis*, 28(3):476–480, 1992. doi: 10.7589/0090-3558-28.3.476.

[187] T Ayllón, AM Nijhof, W Weiher, B Bauer, X Allène, and PH Clausen. Feeding behaviour of *Culicoides* spp.(Diptera: Ceratopogonidae) on cattle and sheep in northeast Germany. *Parasit Vector*, 7:34, 2014. doi: 10.1186/1756-3305-7-34.

[188] ARW Elbers, JL Gonzales, and R Meiswinkel. Comparing *Culicoides* biting rates in horses and cattle in The Netherlands: potential relevance to African horse sickness epidemiology. *Entomol Exp Appl*, 166(7):535–544, 2018. doi: 10.1111/eea.12684.

[189] M Fall, AG Fall, MT Seck, J Bouyer, M Diarra, R Lancelot, G Gimonneau, C Garros, MT Bakhoum, O Faye, T Baldet, and T Balenghien. Host preferences and circadian rhythm of *Culicoides* (Diptera: Ceratopogonidae), vectors of African horse sickness and bluetongue viruses in Senegal. *Acta tropica*, 149:239–245, 2015. doi: 10.1016/j.actatropica.2015.06.012.

[190] IM Mackerras, MJ Mackerras, and FM Burnet. Experimental studies of ephemeral fever in Australian cattle. Common Wealth of Australia, Council Sci. *Ind Res Bull*, (136), 1940.

[191] JE Stokes, KE Darpel, S Gubbins, S Carpenter, MM Fernández de Marco, LM Hernández-Triana, AR Fooks, N Johnson, and C Sanders. Investigation of bovine ephemeral fever virus transmission by putative dipteran vectors under experimental conditions. *Parasit Vector*, 13(1):597, 2020. doi: 10.1186/s13071-020-04485-5.

[192] Y Braverman, K Frish, M Reis, and KY Mumcuoglu. Host preference of *Culicoides* spp. from Israel based on sensory organs and morphometry (Diptera: Ceratopogonidae). 2012. doi: 10.1127/entom.gen/34/2012/97.

[193] D Augot, L Hadj-Henni, SE Strutz, D Slama, C Millot, J Depaquit, and JM Millot. Association between host species choice and morphological characters of main sensory structures of culicoides in the palaeartic region. *PeerJ*, 5:e3478, 2017. doi: 10.7717/peerj.3478.

[194] E Isberg, Y Hillbur, and R Ignell. Comparative study of antennal and maxillary palp olfactory sensilla of female biting midges (Diptera: Ceratopogonidae: *Culicoides*) in the context of host preference and phylogeny. *J Med Entomol*, 50(3):485–492, 2013. doi: 10.1603/me12235.

[195] Y Ma, J Xu, Z Yang, X Wang, Z Lin, W Zhao, Y Wang, X Li, and H Shi. A video clip of the biting midge *Culicoides anophelis* ingesting blood from an engorged *Anopheles* mosquito in Hainan, China. *Parasit Vector*, 6(1):326, 2013. doi: 10.1186/1756-3305-6-326.

[196] LG Mukwaya. Genetic control of feeding preferences in the mosquitoes *Aedes* (Stegomyia) *simpsoni* and *aegypti*. *Physiol Entomol*, 2(2):133–145, 1977. doi: 10.1111/j.1365-3032.1977.tb00091.x.

[197] Ar Ulloa García, JI Arredondo Jiménez, I Fernández Salas, MH Rodríguez, and L González Cerón. Innate host selection in *Anopheles vestitipennis* from southern Mexico. *J Am Mosq Control Assoc*, 20(4):337–341, 2004.

[198] AC Gerry, VSI Monteys, JOM Vidal, O Francino, and BA Mullens. Biting rates of *Culicoides* midges (Diptera: Ceratopogonidae) on sheep in northeastern Spain in relation to midge capture using UV light and carbon dioxide-baited traps. *J Med Entomol*, 46(3):615–624, 2014. doi: 10.1603/033.046.0329.