1 **On the use of kinship and familiarity associated social information in mediating**
2 **_Drosophila melanogaster_ oviposition decisions**
3

4 Emily Rakosy[1,2], Sanduni Talagala[1] & Tristan A.F. Long[1]*
5
6 [1.] Department of Biology, Wilfrid Laurier University, 75 University Avenue West, Waterloo,
7 ON, Canada
8 [2.] Department of Cell & Systems Biology, University of Toronto Mississauga, 3359 Mississauga
9 Road, Mississauga, ON, Canada
10 * Corresponding author. Email: tlong@wlu.ca ORCID: 0000-0002-8708-2728

11
12 ***Abstract***
13

Decisions about where to lay one's eggs are of great importance, as the specific conditions in
one's offspring developmental environment may be critical to their subsequent survival and
lifetime reproductive success. Information produced by conspecifics can potentially be useful to
an individual's oviposition decision-making process as this "social information" may provide an
energetically cheaper means of assessing site suitability rather than acquiring it personally.
However, not all conspecific public information may be equally beneficial, and cues produced by
kin may be especially valuable as they might signal uniquely suitable microenvironments, as well
as suggesting other fitness advantages resulting from improved foraging success and/or a
decreased risk of competition/cannibalism amongst kin compared to sites where unrelated
conspecifics are located. Using the fruit fly, _Drosophila melanogaster_, we explored whether public
information use is associated with kin-based egg-laying decisions. Kinship is potentially
recognized in several ways, including cues produced by environmentally-associated gut
microbiota of the egg-laying flies, and so we also explored whether there were biases in how focal
females interacted with cues from individuals that differed in both their genetic relatedness, as
well as their environmental "familiarity." In a series of inter-connected assays, we examined the
egg-laying behaviour of females that interacted with a choice of potential egg-laying substrates
that differed in the manner of their prior conspecific exposure. Not only do female flies lay more
eggs on sites that had cues left behind by conspecifics compared to unexposed substrates, but they
also exhibited strong biases in their oviposition patterns that was consistent with being sensitive to
the kinship status of the prior substrate occupants. Furthermore, while females did not apparently
discriminate between sites differing in prior egg abundance, they did exhibit bias in their egg-
laying based on the population of origin of the demonstrator female. The basis of the kinship
categorization by ovipositing females appears to be based on phenotypes that reflect true genetic
relatedness that are neither mimicked nor disrupted by the conspecific's developmental
environment. These results highlight the potential usefulness of _D. melanogaster_ as a model to
understand the evolution of social behaviour in the expression of decision-making.

40
41 ***Keywords***: Social information, _Drosophila melanogaster_, oviposition, conspecifics, kin recognition,
42 microenvironments, behaviour, copying, public information
43

Competing interests: The authors declare no competing interests.
Ethics: This study did not require approval from an ethics committee.
Supporting Data: All data analysed in this manuscript are archived in Borealis, the Canadian
Dataverse Repository: https://doi.org/10.5683/SP3/N7XBOJ

Introduction

In species where there is little-to-no post-natal parental care, decisions about where and

when an individual should lay their eggs are likely very important, as there are potentially dire

fitness outcomes for one's offspring if unsuitable oviposition sites are chosen (Thorpe 1945,

Heard 1994, Lihoreau et al. 2016). In order to make better decisions, using cues produced by

conspecifics (aka "social information" *sensu* Valone and Templeton 2002) may be of value to the

individual as they may provide meaningful information about potential egg-laying sites that can

be obtained at a lower energetic cost than having to explore and evaluate alternate options

oneself. Social information can be valuable in reducing an individual's uncertainty about their

environment, and may help them locate areas that are rich in important resources such as food,

shelter, desirable mates, increased foraging success, or are at a lower risk of predation/parasitism,

any of which may benefit the mimicking individual and/or their offspring (Sarin and Dukas

2009, Blanchet et al. 2010, Grüter and Leadbeater 2014). Furthermore, the energy savings of

using social information over private information can also be redirected into increased

provisioning and/or production of offspring, thereby enhancing the benefits of its use in making

decisions (Golden and Dukas 2014).

While fruit flies, *Drosophila melanogaster*, have a long and storied use as a model in studies of

genetics, development, behaviour and evolution (*see* Weiner 2000, Brookes 2001), they are also

emerging as a powerful model in which to study social learning and the use of social information

(Leadbeater 2009, Chen and Sokolowski 2022). *D. melanogaster* has been found to use the visual,

gustatory, tactile, and olfactory cues produced by conspecifics in a number of different decision-

making contexts. For instance, flies use social cues when engaged in spatial learning tasks to

improve the speed that a "safe zone" is detected (Foucaud et al. 2013). Virgin female flies that are

exposed to deposits left by conspecific demonstrators exhibit lower post-mating egg-laying rates

than unexposed females, presumably because they are better able to anticipate the forthcoming

availability of oviposition sites and the potential degree of future larval competition (Fowler et al.

2022). Mate choice can involve social information as female fruit flies are more likely to mate

with males exhibiting similar characteristics as a male they've observed successfully mating with

another female (Mery et al. 2009, Monier et al. 2018, Nöbel et al. 2018, Nöbel et al. 2022, *but see*

Belkina et al. 2021) presumably because mate choice is often costly to females (*see* Reynolds and

Gross 1990), and copying the decisions made by conspecifics represents an economical adaptive

strategy (Monier et al. 2018). Important information about environmental conditions can also be

obtained indirectly by flies interacting with conspecifics. In a study by Kacsoh et al. (2015) female

flies decreased their oviposition rate in response to exposure to parasitic wasps, and this defensive

change in egg-laying behaviour was adopted by inexperienced flies housed with the wasp-

exposed females, with cues about environmental risk communicated visually using the

demonstrator flies' wings. Thus, while *D. melanogaster* has not been considered a "social" species

(Chen and Sokolowski 2022), it is evident that fruit flies can, and do, use social information to

inform their decisions.

One important context where social information seems to be used extensively by *D.*

melanogaster females is in selecting suitable oviposition sites (Battesti *et al.* 2012, Durisko et al. 2014,

Sarin and Dukas, 2009, Golden and Dukas 2014, Malek and Long 2020, Moreira-Soto et al.

2024). In the wild, there may be considerable heterogeneity in the availability and quality of

oviposition sites that are available to female *D. melanogaster* (Reaume and Sokolowki 2006,

Markow 2015), and the choice of oviposition environment can have profound consequences for

their offspring's fitness (*e.g.* Prasad et al. 2003, Klepsatel et al. 2020). *D. melanogaster* females may

gain insight on the suitability of oviposition sites through direct observation of other females

ovipositing (*as in* Sarin and Dukas, 2009 and Battesti *et al.* 2012), the presence of larvae (*e.g.*

Golden and Dukas, 2014) and/or the presence of chemical cues produced by conspecifics (*e.g.*

Venu et al. 2014, Lin et al. 2015, Duménil et al. 2016, Moreira-Soto et al. 2024). Laying one's

eggs at sites already chosen by conspecifics may also be of benefit to the offspring's success due to

enhanced communal digestion via secreted enzymes (Gregg, et al. 1990, Sakaguchi and Suzuki

2013), increased beneficial microbial activity (Gilbert 1980, Wertheim et al. 2002a, Wertheim et

al. 2002b), and/or improved group foraging (Dombrovski et al. 2017, Khodaei et al. 2020, Shoot

et al. 2024). The reliance upon social cues by ovipositing *D. melanogaster* is not without potential

pitfalls. First, by outsourcing one's information gathering to others, there is the risk of mimicking

other's bad decisions. For instance, in a study by Golden and Dukas (2014) where females were

given a choice of laying their eggs on a site where nutrition was poor but contained social cues

(the presence of larvae) or a site where the media contained 3 times more nutrition but lacked

conspecific cues, the ovipositing females treated them as being of equal quality, even though any

offspring laid on the former patches did not have access to sufficient nutrients to survive.

Similarly, Duménil et al. (2016) observed that the presence of a mated female demonstrator on

yeast-free media made this undesirable oviposition site equally-valued as yeasted media (that has

not been exposed to conspecifics) to later ovipositing females. Furthermore, laying one's eggs

near those of others can have additional challenges arising from competition for limited resources

(De Jong 1976, Joshi et al. 1996, Golden and Dukas 2014), increased exposure to toxic waste

products (Borash et al. 1998), and the threat of cannibalism at many developmental stages

(Vijendravarma et al. 2013, Ahmed et al. 2015, Yang 2018). However, some of these risks may

be mediated if females are able to oviposit at sites containing kin, as *D. melanogaster* larvae are

more co-operative (Khodaei and Long 2019), and less cannibalistic (Khodaei and Long 2020,

Fisher et al. 2021) towards close relatives, compared to how they engage with unrelated

conspecifics. Thus, if females were able to differentiate social information produced by kin from

those generated by other demonstrators, they might be able to achieve greater fitness benefits by

selectively copying the choices made by relatives. Additionally, as there can be extensive intra-

specific genetic variation *D. melanogaster* for nutritional preferences and dietary tolerances (*e.g.*

Toshima et al. 2014, Havula et al. 2022), preferentially laying eggs near those of close relatives

may increase the likelihood of selecting the best micro-habitat for the success of one's own

offspring.

When considering the possibility that individuals might differentially use social cues

produced by relatives and non-relatives, one must also consider what (if any) cues could indicate

kinship (or could be inferred to be reliable proxies of kinship). In *D. melanogaster* such information

could conceivably be perceived from cuticular hydrocarbons (CHCs) and/or gut bacteria left

behind by conspecifics. CHCs are long-chain hydrocarbons that are synthesized in insect

oenocyte cells which become part of the cuticle's waxy coating along with lipids, aldehydes,

alcohols, esters and fatty acids which helps prevent water loss (*reviewed in* Richard and Hunt 2013,

Chung and Caroll 2015). These fat molecules also play an additional important role in olfactory

communication related to species recognition and mate choice decisions (*reviewed in* Singer 1998,

Ferveur 2005, Scott et al. 2011, Chung and Caroll 2015). CHCs may provide accurate

information about kinship, as in addition to differences between species (Moreira-Soto et al.

2024) there also exists substantial intra-specific genetic variation within *D. melanogaster* populations

for these phenotypes (Foley et al. 2007, Dembeck et al. 2015, Heys et al. 2018). It has been

suggested that assortative mating patterns may be mediated by the degree of similarity of CHC

chemical profiles of males and females (Haerty et al. 2002, Robinson et al. 2012). Recent work

has revealed that female flies transfer their own CHCs (along with male-specific compounds

acquired during mating) to their laid eggs (Moreira-Soto et al. 2024), which could conceivably be

used as signals of kinship by subsequent ovipositing females.

In addition to CHC chemical cues, olfaction-based decisions may be influenced by signals

associated with differences in an individual's bacterial community (Sharon et al. 2010, Najarro et

al. 2015). While the *D. melanogaster* microbiome is relatively simple compared to other hosts

(Broderick and Lemaitre 2012), there can be substantial variation between individuals/families

from the same population in the composition of their microbial communities (Wang and

Staubach 2018, Heys et al. 2018). Individuals acquire some of their microbial symbionts from

their mothers, as like CHCs, bacteria are transferred by ovipositing host mothers to the surface of

their eggs, and the subsequent consumption of the chorion by hatched larvae helps establish their

own gut flora (Bakula 1969, Guilhot et al. 2023). Thus, differences in cues produced by bacterial

communities could conceivably and independently be used to infer kinship (Lizé et al. 2014,

Lewis et al. 2014). Importantly, CHC phenotypes and/or microbiotic communities in *D.*

melanogaster can also be influenced by the flies' developmental environment (Sharon et al. 2010,

Fedina et al. 2012, Lizé et al. 2014, Venu et al. 2014, Heys et al. 2018, García-Roa et al. 2022,

Pavković-Lučić et al. 2022), raising the possibility that recognition of "kin" may alternately be the

recognition of similarity of environmental history ("familiarity"), or some combination of the two

factors. In a previous study that examined whether kinship mediated the magnitude and

consequences of male-male competition in adult flies, it was found that reductions in male-

induced harm were *only* seen in groups of males that were both closely related to each other *and*

'familiar' by having developed in the same larval environment (Le Page et al. 2017). Lizé et al.

(2014) described how females were more likely to mate with a familiar male, while males' ability

to strategically avoid investing into mates with a sibling was impeded if she had developed in the

same environment. Similarly, Heys et al. (2018) noted that the ability of fruit flies of both sexes to

perceive the kinship status of their mate appeared to be dependent on the state of the mate's

microbiome – with females only acting differently towards kin when those individuals had

developed in the same media type (and had an intact microbiota), while a male's increased

investment into mating with an unrelated female only occurred if she had been grown in a

different environment (or whose microbiota had been disrupted with the use of antibiotics). The

complicated nature of the interaction between developmental environment and relatedness was

also explored by García-Roa et al. (2022) who found that flies developing in different

environments exhibited both different CHC and microbial profiles, and that related flies had

microbiotic communities of similar diversity (but not composition), and that diversity was

correlated with CHC phenotypic profiles. Together these studies highlight that socially-available

kinship cues (or proxies thereof) may involve a number of independent and inter-related olfactory

signals, whose expression may depend not only on relatedness but also on the specific nature of

the developmental environment.

With these ideas and findings in mind, in this study we set out to examine whether female

flies showed bias in their use of social information cues left behind by conspecifics when selecting

ovipositing sites, and would respond differently to cues produced by kin vs non-kin.

Simultaneously, we also explored whether a female's response to potential kinship cues were

conditional on the conspecific's developmental environment. Overall, the goal of this research is

to better characterize copying behavior in *D. melanogaster* and to gain better insight into the factors

that influence these oviposition site decisions.

Materials and Methods

Population Protocols and Fly Maintenance

All fruit flies used in our assays were obtained from one of three associated laboratory *D.*

melanogaster populations: *Ives* (hereafter "IV"), "IV-*bw*", and "IV-*bwD*". The IV population was

derived from a sample of mated females collected in 1975 from Amherst MA, USA sample

population (Rose 1984). The IV-*bw* population was created by introgressing the brown-eye

recessive allele, *bw*[1], into the genetic background of the IV population via repeated rounds of

back-crossing, to create a competitor population (Khodaei and Long 2020). The IV-*bwD*

population was also created in the same fashion, by introducing the *bwD* allele (a dominant allele

that also produces a brown-eyed phenotype) into an IV genetic background. The IV-*bw* and IV-

bwD populations have been subject to period rounds of additional back-crossing to the IV

population in an attempt to minimize their divergence from the base population, with the last

session occurring approximately 2 years prior to the start of this experiment. These three

populations are each maintained at a census size of ~3500 adults per generation, incubated at

25°C incubation temperature with 60% humidity on a 12L:12D hour diurnal light cycle (Martin

and Long 2015). Flies are maintained in standard *Drosophila* vials (95mm H x 28.5mm OD)

containing 10mL of banana/agar/killed-yeast food, dusted with live yeast (Rose 1984, Shoot et

al. 2024). Populations are cultured on a 14-day cycle, wherein every fortnight adult flies are

removed from their natal vials, mixed *en masse* under light CO_2 anaesthesia and distributed into

vials containing new media for up to 18h, before being removed and the eggs laid are then

trimmed by hand to a density of approximately 100 eggs per vial (Martin and Long 2015).

Experiment 1: Test of effect of conspecific relatedness and developmental environment on egg-laying behavioural

decisions.

In our first experiment, we collected adult male and female flies as virgins (within 8h of

eclosion from their pupae) from the IV population, and housed them individually. We then

created 50 families by randomly combining single males and females into "mini-egg" chambers

(Kartell 733/4 polyethylene 20 mL sample vials; 74.5mm H x 24.8mm OD; *see* Figure S1a)

whose lid contained juice/agar media (Sullivan et al. 2000) and was used as a "dish" for

ovipositing. We left these chambers in the incubator for ~18h, before removing the adult flies

and counting the number of eggs that had been laid overnight. We then evenly split each family's

eggs among 2 different vials each containing 10ml of media, and then added a sufficient number

of similarly-aged eggs obtained from the IV-*bw* population so that each vial ultimately contained

~100 eggs apiece. We returned these pairs of vials to the incubator where they were allowed to

develop for the next 9 days when we began collecting virgin females for use in the experiment.

For each pair of vials, we haphazardly designated one as the "focal" vial, from which we

collected 3 virgin females: a IV female ("the focal female"), a second IV female (the focal female's

sister, raised in the same vial), and one brown-eyed IV-*bw* female (an unrelated female, raised in

the same vial as the focal female). From the other (non-focal) vial in the pair, we collected 2 virgin

females: an IV female (the focal female's sister, raised in a different vial) and one brown-eyed IV-

bw female (an unrelated female, raised in a different vial than the focal female). We kept these

females individually for up to 48h in test-tubes containing media to further verify their virgin

status (by the absence of any fertilized eggs). Next, we introduced two similarly-aged adult IV

males into the focal female's vial (so that all her offspring would express the wild-type red-eye

phenotype). At the same time we transferred all other females into their own mini-egg chambers

whose dish lids contained standard banana media containing 3 IV-*bwD* males so that all offspring

produced by demonstrators would express a *brown-eyed* eye phenotype and to ensure that any

male-specific compounds transferred to the dish surface would be consistent across all dishes (Lin

et al. 2015, Moreira-Soto et al. 2024). Females and males were left overnight to ensure sufficient

time for mating to occur. On the following morning all non-focal flies were removed from the

mini-egg chambers, and sets of four dishes were affixed using adhesive putty to the bottom of a

larger chamber, (hereafter "arena"), which consisted of an inverted plastic box (KIS Omni box,

20.3 x 15.9 x 9.6cm) modified by the addition of mesh vent holes to box's outer edges (*see* Figure

S1b, Young et al. 2018). A fifth dish (that had never been exposed to flies) was also placed in the

arena before it was sealed shut, and the (lightly anesthetized) focal female was introduced via an

access hole. Thus the five dishes inside each arena represent available oviposition sites for the

focal female that potentially differ in their social information: The 'related/same' dish contains

cues produced by a sister that developed in the same vial environment as the focal IV female; the

'unrelated/same' dish contains cues from an unrelated IV-*bw* female that developed in the same

vial environment as the focal IV female; the related/different' dish contains cues produced by a

sister that developed in a different vial environment than the focal IV female; the

'unrelated/different' dish contains cues from an unrelated IV-*bw* female that developed in a

different vial environment as the focal IV female; and finally the 'control/unexposed' dish

contains no social information, as it had no prior exposure to flies (*see* Figure S1c). The arenas

were housed in a quiet, well-lit room, and were left undisturbed for 20 minutes following the

introduction of the focal female. The arenas were then observed every 20 min for ~4.5h (starting

at 10:30am), and for 4h on the next day (beginning at 9:30am) for a total of 26 observation

sessions. During each observation session, the arenas were watched for a 5s time period, allowing

for the identification of which of the 5 dishes (if any) the focal IV female was located. The identity

of the different treatments was unknown to the observers. One all observations were complete,

we unsealed the arena, discarded the focal female and transferred the media in each dish into

vials containing 10m of media. These labelled vials were sealed and returned to the incubator for

2 weeks. At that time, all eclosed adults were removed, and their eye colours were tallied. Red-

eyed flies were the offspring of the focal female, while brown eye-flies ($wt+/bwD$ or bw_1/bwD)

were produced by one of the conspecific females.

Experiment 2: Test of effect of conspecific egg production on focal female egg-laying behavioural decisions.

Based on the egg-laying patterns observed in Experiment 1, we performed a series of

complementary assays designed to test alternative hypotheses for the focal flies' oviposition

biases. Thus, in this experiment we set out to examine whether the number of eggs laid by

conspecifics on a substrate was associated with the number of eggs laid by a focal female. We

began by collecting mated IV-*bw* female flies from the lab's stock population and placed either 1,

2, or 0 individuals overnight into mini-egg chambers containing media in their dish lid. The next

morning, one of each type of the dishes was affixed to the bottom of an arena (*see* Figure S1d).

Next, we added a single mated IV female to each arena and left them undisturbed for 22h,

before transferring the media in each dish to a vial containing 10ml of media. We returned those

vials to the incubator and counted the number of wild-type and brown-eye offspring that had

eclosed 12 days later.

Experiment 3: Test of egg-laying biases of focal females towards cues from flies originating from the IV versus the

IV-bw populations.

In our third experiment we explored the alternate hypothesis that the egg-laying patterns

observed in Experiment 1 might reflect preferences of the IV-originating focal female to lay eggs

on substrates that had come into contact with any other IV flies (regardless of their degree of

immediate genetic consanguinity) and/or the avoidance of substrates that had come in contact

with any IV-*bw* flies. For this assay we set up replicate vials containing 5 mated IV and 5 mated

IV-*bw* females, that were allowed to oviposit for 18h before being discarded, and the number of

eggs laid standardized to 100 apiece. These vials were placed in the incubator, and we collected a

virgin red-eyed and a virgin brown-eyed female from each vial as they eclosed (thus both flies had

experienced the same developmental environment). Each female was then placed into their own

mini egg chamber with 2 IV-*bwD* males overnight before being discarded, and the dish affixed to

the inside of an arena, along with an unexposed/control dish (*see* Figure S1e). An individual

mated IV female obtained from the lab stock population was introduced to each of the arenas,

and her location in the arena was periodically observed every 20 minutes over the course of 5h

(for a total of 15 sessions). The arena was left undisturbed overnight, and the following morning

media was removed from the dishes and transferred to vials containing 10ml of media. These

vials were incubated for 12 days, and the number of offspring with different eye colours was

recorded.

Experiment 4: Revisiting the effect of conspecific relatedness on egg-laying behavioural decisions.

Based on the outcomes of our second and third experiments, we set about directly

comparing the egg-laying behaviour of focal females towards substrates that had been exposed to

their sibling or to a entirely unrelated IV-derived female. As in experiment 1 we collected the

focal female's parents from the IV population, mated them, placed them into mini-egg chambers

overnight, then counted and split their eggs amongst 2 vials, with additional similarly-aged IV-bw

eggs added to bring the density to a total of 100 eggs apiece. From each of the pairs of vials we

collected virgin females as they eclosed and kept them individually. One of the vials was

arbitrarily designated as that which yielded the focal female (who was then housed with 2 IV

males from the stock population overnight), and a female from the other vial (*i.e.* her sister raised

in a different environment) was housed with 2 IV-*bwD* males overnight in a mini egg chamber.

We also haphazardly chose one of the IV females collected from one of the other pairs of vials

(thus an unrelated female, raised in a different environment) and housed her 2 IV-*bwD* males

overnight in a mini egg chamber. These dishes (along with a control/unexposed dish) were

placed into an arena (*see* Figure S1f), into which we added the mated focal females. Her location

was periodically observed every 20 minutes over the course of 5h (for a total of 15 sessions). Next,

the chamber was left undisturbed overnight and the following morning, we transferred the media

from each dish into a new vial containing 10ml of media which were incubated for an additional

12 days. The eye colour of all adult flies present in these vials were tallied and recorded.

Statistical Analyses

All data analyses were conducted in the R statistical computing environment (version 4.0.3, R

Core Team, 2020). In all of our experiments we compared the total number of visits by focal

females to the five different lids in each arena using generalized linear models (GLMs), with

quasipoisson error distributions, where the independent factor was the lid treatment category.

We determined whether the treatment means were significantly different from each other with

likelihood-ratio chi-square tests using the *Anova* function in the *car* package (Fox and Weisberg

2011), followed by Tukey HSD test executed by the *glht* function in the *multcomp* package

(Hothorn et al. 2008) if necessary to determine the specific location of differences between

treatments. For experiment 1 we also set out to determine if there was any difference in patch

visitation that varied with relatedness and/or familiarity treatments, by using the same response

data (excluding observations made on the control lid, as this treatment by definition had received

no prior conspecific contact) which was analyzed using a 2-way GLM with quasipoisson errors.

The degree of kinship, the source, and their interaction were the independent factors in this

model. The significance of these effects was determined using a likelihood-ratio chi-square test

using the *Anova* function.

For all of our experiments we (separately) analyzed the total number of red-eyed and brown-

eyed adults that successfully eclosed from the different lids in each arena using GLMs, with

quasipoisson error distributions, where the independent factor was the lid treatment category.

For each we determined whether the treatment means were significantly different from each

other using likelihood-ratio chi-square tests implemented using the *Anova* function followed by

Tukey HSD tests if necessary to determine the specific location of differences between

treatments. As above for experiment 1 data we also set out to determine if there was any

difference in offspring number associated with relatedness and/or familiarity treatments, by

excluding data collected from the control lid treatment, and analyzing the rest of the data set

using a 2-way GLM with quasipoisson errors. The degree of kinship, the vial source, and their

interaction were the independent factors in this model. The significance of these effects was

determined using a likelihood-ratio chi-square test using the *Anova* function.

RESULTS

Experiment 1: Test of effect of conspecific relatedness and developmental environment on egg-laying behavioural

decisions.

In our first experiment, we compared the cumulative number of visits that *D. melanogaster*

focal females made to five different dishes, as well as the number of offspring that eclosed as

adults from each of those dishes. Our analysis found that those females exhibited heterogeneity in

their dish associations (GLM: LRχ^2 =17.8, df=4, p=0.001), and while they visited the

control/unexposed dishes less frequently than the demonstrator dishes, within these four

treatment groups, they did not express any significant bias (Figure S2). Offspring production

patterns indicated that the females' oviposition patterns were also heterogeneous (GLM: LRχ^2

=110.08, df=4, p=4.91x10^{-23}), with significantly more red-eyed focal offspring eclosing from

those dishes that had previously been in contact with a related demonstrator individual than from

either the control/unexposed dish or either of the dishes whose prior occupant was an unrelated

individual (Figure 1). Our focused two-way GLM confirmed that the amount of oviposition

activity on a dish was significantly associated with the demonstrator's kinship (LRχ^2 =78.94,

df=1, p=6.40x10^{-19}), and there was no significant effect of developmental environment (LRχ^2

=1.97, df=1, p=0.16), and the interaction between kinship and developmental environment was

also not significant (LRχ^2 =0.176, df=1, p=0.67). We also noted that the number of

demonstrator offspring were not homogeneous across the four treatment groups, with more

brown-eyed offspring eclosing from vials whose prior occupant was an unrelated female (IV-*bw*,

mated to a IV-*bwD* male) than those that had previously been exposed to an unrelated female.

Experiment 2: Test of effect of conspecific egg production on focal female egg-laying behavioural decisions.

In our second experiment, we set out to test the alternate hypothesis that the patterns of

red-eyed offspring laid by our focal females was influenced by the abundance of *brown-eyed* eggs

already present on the dishes. In this assay either zero, one or two demonstrator females was

housed on a dish prior to being offered as a potential oviposition site for a mated focal female.

More demonstrators on a dish yielded more *brown-eyed* offspring: zero *brown-eyed* offspring eclosed

from the control dishes, and two demonstrator females produced, on average more *brown-eyed*

offspring than a single demonstrator female (GLM: LRχ^2 =19.97, df=1, p=7.88x10^{-16}; Figure

2a). Despite the dramatic differences in the number of demonstrator eggs between the three

dishes, there was no significant difference in the number of *red-eyed* focal offspring that eclosed

from the three dishes (GLM: LRχ^2 =3.69, df=2, p=0.157; Figure 2b), but there was a trend

towards *more* offspring eclosing from those dishes that had greater amounts of demonstrator

exposure/offspring.

Experiment 3: Test of egg-laying biases of focal females towards cues from flies originating from the IV versus the

IV-bw populations.

In our third experiment, we set out to test the alternate hypothesis that the patterns of *red-*

eyed offspring laid by our focal females was influenced by the population of origin of the

demonstrator females, and not their degree of kinship. This involved comparing the cumulative

number of visits that *D. melanogaster* focal females made to three different dishes (a

control/unexposed dish, a dish whose demonstrator was an unrelated IV female or a dish whose

demonstrator was an unrelated IV-*bw* female), as well as the number of offspring that eclosed as

adults from each of those dishes. Our analysis found that focal females exhibited heterogeneity in

their dish associations (GLM: LRχ^2 =92.27, df=2, p=9.17 x10^{-21}) with flies associating with the

demonstrator dishes more frequently than with the control/unexposed dishes; but no significant

difference in the mean number of visits to the two demonstrator dishes (Figure S3). The

distribution of focal female offspring was heterogenous across the three dish types (GLM: LRχ^2

=96.1, df=2, p=1.39 x10^{-21}). Focal females laid more eggs on the unrelated IV dish than they did

either on the control/unexposed dish, or the unrelated *IV-bw* dish. The mean number of *red-eyed*

offspring that were collected from the *IV-bw* dish was not significantly different from the number

collected from the control/unexposed dish (Figure 3b). We also noted that the number of

demonstrator offspring were significantly different from each other, with more brown-eyed

offspring eclosing from vials whose prior occupant was an unrelated IV female (mated to a IV-

bwD male) than those that that had previously been exposed to an unrelated IV-*bw* female (also

mated to a IV-*bwD* male) (GLM: LRχ^2 =366.14, df=2, p=3.11 x10^{-80}; Figure 3a).

Experiment 4: Revisiting the effect of conspecific relatedness on egg-laying behavioural decisions.

In our fourth experiment, we set out (again) to compare the patterns of dish visitation and

oviposition by focal females on three different dishes (a control/unexposed dish, a dish whose

demonstrator was an unrelated IV female, or a dish whose demonstrator was an unrelated IV

female). The focal females exhibited bias on their dish visitation rates (GLM: $LR\chi^2$ =12.86, df=2,

p=0.0016), as they were more frequent visitors to the two types of demonstrator dishes than the

control/unexposed dishes (but did not differ significantly between these two treatment types;

Figure S4). The distribution of focal female offspring was heterogenous across the three dish

types (GLM: $LR\chi^2$ =180.24, df=2, p=7.26x10^{-40}). Focal females laid significantly more eggs on

the related IV dish than they on the unrelated IV dish, and both demonstrator dishes yielded

more focal female offspring than did the control/unexposed dishes (Figure 4). There was no

significant difference in the number of demonstrator offspring eclosing from the related IV dishes

than from the unrelated IV dishes (GLM: $LR\chi^2$ = 2.59, df=1, p=0.11; Figure S5).

DISCUSSION

The decision where to oviposit may be very important, especially in these species where

there is no post-laying parental care (Prasad et al. 2003, Klepsatel et al. 2020). Fruit flies,

Drosophila melanogaster, like many other species, use social information in their decision-making

(Leadbeater 2009, Chen and Sokolowski 2022). While there are benefits of laying one's eggs in

proximity to those of others, there are also risks – which might be mediated by the degree of

kinship between conspecifics. In this study, we set out to examine the possibility that female flies

might selectively use cues produced by close relatives, presumably to increase their offspring's

chance of success. In doing so, we simultaneously explored the factors that might influence how

flies might infer kinship, as well as investigate alternate hypotheses about social information

usage. Our assays confirm the extensive use of social cues by *D. melanogaster* females, and

furthermore collectively suggest that they are capable to kinship discrimination and are capable

of distinguishing between conspecifics from different populations. These results add to the

growing body of knowledge about the sophisticated social information use by this model species

(Chen and Sokolowski 2022).

In all assays focal females associated more with those substrates that had come into

contact with conspecifics than with the unexposed/control dishes (Figures S2, S3 & S4), clearly

illustrating the importance of social cues in their decision-making, and consistent with numerous

previous studies (*e.g.* Battesti *et al.* 2012, Duménil et al. 2016, Malek and Long 2020). However,

not all conspecific cues elicited the same responses. In the first assay, we compared the egg-laying

patterns of focal females on dishes exposed to a range of different demonstrators. By splitting sets

of full sibling eggs between two different vials to develop, and collecting both adult relatives and

non-relatives from each of the pairs of vials, we set out to test if and how *D. melanogaster* females

might distinguish kin cues. The focal females laid significantly more eggs on those dishes where

the demonstrator female was a sister, but did not appear to differentiate between the cues of

demonstrator sisters originating from the same or a different vial environment (Figure 1a). These

results suggest not only that ovipositing females selectively use social information, and favour cues

produced by their relatives, but furthermore that kinship cues were not adopted by non-kin

demonstrators developing in the same vial, nor disrupted in those kin developing in a different

vial. Determining the specific cues being used by focal females is beyond the scope of this study as

demonstrator females transfer their gut bacteria, CHCs (and male-specific compounds acquired

during copulation) to the surface of their eggs and the media (Bakula 1969, Guilhot et al. 2023,

Moreira-Soto et al. 2024). The expression these olfactory cues can be influenced by both an

individual's genotype and/or their developmental environment (Sharon et al. 2010, Fedina et al.

2012, Lizé et al. 2013, Venu et al. 2014, Heys et al. 2018, García-Roa et al. 2022, Pavković-

Lučić et al. 2022). While the absence of significant differences associated with the developmental

environment (Figure 1b), could suggest that "familiarity" (and by extension the microbiotic cues

that could be used as proxies of kinship) are not as important as "relatedness", it should be noted

that in our experimental design, the conditions of the two vial environments were similar (same

media, same larval density), which may have minimized the potential for the development of

differences in these cues. Interestingly, Malek and Long (2020) observed that when focal females

developed in a protein-rich vial environment, they oviposited more on dishes where the

demonstrator from the same population had also developed on similar media, but when the focal

females developed on standard media, they laid more on those dishes where the demonstrator

came from a different *D. melanogaster* population but had also developed on standard media than

they did on dishes where the demonstrator was from the same population but had developed on

high-protein media. Thus, one should not necessarily conclude that our results indicate that

CHCs are the only way in which egg kinship was/can be recognized by the focal females in this

species, and future studies exploring this idea across a range of divergent environments are likely

to expand our understanding of how social information is produced and interpreted in *D.*

melanogaster.

While the observed pattern of focal female oviposition is consistent with kin-associated

biases, other processes could also have resulted in the differential egg-laying outcomes, and so in

order to validate or rule-out these alternate hypotheses we conducted several complementary

assays. We had been initially struck with the distinctive pattern of offspring laid by the

demonstrator females (Figure 1b), wherein there were more *brown-eyed* conspecific offspring

collected from dishes where the demonstrator was unrelated than when the demonstrator was

related. Perhaps our focal females weren't deciding to lay their eggs on dishes that contained

their kin's offspring, but instead were avoiding dishes where there were already lots of eggs (and a

heightened risk of competition and/or cannibalism)? We explored this possibility in our second

assay, where focal females could choose between dishes with dramatically different number of

eggs (Figure 2a). If females were avoiding dishes where there were too many conspecific eggs

already present, we would expect to see more eggs laid on the "one female" treatment than on

the "two females" treatment. In fact, we observed no significant difference in focal female

oviposition bias between the two groups (and if anything, they often laid *more* eggs on the latter

group's dishes; Figure 2b). Previous work by Golden and Dukas (2014) found that female flies

given a choice between a control and a demonstrator dish (which had been exposed to 5 or 20

larvae), did not appear to discriminate against the demonstrator dishes that possessed cues

indicating greater competition during offspring development. Thus, we are fairly confident that

the differences in focal female egg-laying patterns observed in experiment one were not due to

differences in the quantity of demonstrator cues present on a dish. As a side note, we suggest that

the differences in the number of demonstrator offspring observed in that assay is a side effect of

the way in which our demonstrator females were collected (*see* Figure S6 *caption for details*).

The second alternate hypothesis we explored was that the biased pattern of oviposition

made by the focal females in the first experiment on the two "related" dishes, was not due to the

perceived kinship of the demonstrators, but instead was because those demonstrators originated

from the same population (IV), and the demonstrators for the two "unrelated" dishes were from a

different population (IV-*bw*). While IV and IV-*bw* are maintained following the same protocols,

and IV-*bw* is periodically back-crossed to IV, perceptible differences in the cues present in these

isolated populations (or their microbiotic communities) may have arisen via drift and/or

selection. Thus in our third experiment, we compared egg-laying behaviour for focal females

exposed to (unrelated) IV or IV-*bw* demonstrator females. We found that focal females laid more

eggs on the former dishes than they did on the latter (and virtually none on the

control/unexposed dishes). This suggests that at least some of the patterns of oviposition seen in

our first experiment may not necessarily be the result of recognition of kinship *per se*, but instead

with cue(s) that differ between the IV and IV-*bw* populations. Thus, based on the insights gained

in our third assay, we set out to re-examine the social information use of *D. melanogaster* females in

our fourth assay where females were given a choice of oviposition sites that either had no prior

exposure (control), a demonstrator sister (raised in a different vial), or an unrelated demonstrator

female (also from the IV population and raised in a different vial). We observed that focal females

still biased their egg-laying towards the dishes that had previously been in contact with their

sisters, albeit to a lesser extent than seen in experiment 1. This provides additional support to the

hypothesis that female *D. melanogaster* are both capable of distinguishing between cues produced

by kin and non-kin, and that these cues are used in their social decision-making. In doing so, they

may be trying to obtain the potential benefits of developing in a group (Gilbert 1980, Gregg, et

al. 1990, Sakaguchi and Suzuki 2013, Wertheim et al. 2002a, Wertheim et al. 2002b,

Dombrovski et al. 2017, Shoot et al. 2024), while reducing the potential risks from conspecifics

(Khodaei and Long 2019, Khodaei and Long 2020, Fisher et al. 2021).

Collectively, our results illustrate the subtle and plastic nature of social information use by

female *D. melanogaster*: Given a choice, a mated female *D. melanogaster* will lay more eggs on a dish

that has previously been in contact with a sister than one exposed to a mated female from the

same population; will lay more eggs on a dish if it has been previously exposed to an unrelated

mated female from the same population than one exposed to a mated female from a different

population; and will lay more eggs on a dish where the previous occupant was a mated female

from the same population (and raised in a similar environment) than on a previously unoccupied

dish (i.e. the unexposed/control dish). At the same time, a mated female *D. melanogaster* does not

appear to show any bias in their egg laying towards dishes that differ in the number of eggs (and

their associated biochemical cues) present, nor do they show any bias between dishes that had

previously been occupied by a sister raised in the same environment or a different vial. Together,

these pattens illuminate that *D. melanogaster* females are sensitive to differences in the cues

produced by conspecifics and add to the growing appreciation of social decision-making in this

model species.

REFERENCES

Ahmad, M., Chaudhary, S.U., Afzal, A.J. and Tariq, M., 2015. Starvation-induced dietary
behaviour in *Drosophila melanogaster* larvae and adults. *Scientific reports*, *5*(1), p.14285.
Bakula, M., 1969. The persistence of a microbial flora during postembryogenesis of *Drosophila*
melanogaster. *Journal of invertebrate pathology*, 14(3), pp.365-374.
Battesti, M., Moreno, C., Joly, D. and Mery, F., 2012. Spread of social information and
dynamics of social transmission within *Drosophila* groups. *Current biology*, 22(4), pp.309-313.
Battesti, M., Moreno, C., Joly, D. and Mery, F., 2015. Biased social transmission in *Drosophila*
oviposition choice. *Behavioral Ecology and Sociobiology*, *69*, pp.83-87.
Belkina, E.G., Shiglik, A., Sopilko, N.G., Lysenkov, S.N. and Markov, A.V., 2021. Mate choice
copying in *Drosophila* is probably less robust than previously suggested. *Animal*
Behaviour, *176*, pp.175-183.
Blanchet, S., Clobert, J. and Danchin, E., 2010. The role of public information in ecology and
conservation: an emphasis on inadvertent social information. *Annals of the New York Academy*
of Sciences, *1195*(1), pp.149-168.
Borash, D.J., Gibbs, A.G., Joshi, A. and Mueller, L.D., 1998. A genetic polymorphism
maintained by natural selection in a temporally varying environment. *The American*
Naturalist, *151*(2), pp.148-156.
Broderick, N.A. and Lemaitre, B., 2012. Gut-associated microbes of *Drosophila melanogaster*. *Gut*
microbes, *3*(4), pp.307-321.
Brookes, M., 2002. *Fly: The unsung hero of twentieth-century science*. Harper Collins.
Chen, M. and Sokolowski, M.B., 2022. How social experience and environment impacts
behavioural plasticity in *Drosophila*. *Fly*, *16*(1), pp.68-84.
Chung, H. and Carroll, S.B., 2015. Wax, sex and the origin of species: dual roles of insect
cuticular hydrocarbons in adaptation and mating. *BioEssays*, *37*(7), pp.822-830.
De Jong, G., 1976. A model of competition for food. I. Frequency-dependent viabilities. *The*
American Naturalist, *110*(976), pp.1013-1027.
Dembeck, L.M., Böröczky, K., Huang, W., Schal, C., Anholt, R.R. and Mackay, T.F., 2015.
Genetic architecture of natural variation in cuticular hydrocarbon composition in *Drosophila*
melanogaster. elife, *4*, p.e09861.
Durisko, Z., Anderson, B. and Dukas, R., 2014. Adult fruit fly attraction to larvae biases
experience and mediates social learning. *Journal of Experimental Biology*, *217*(7), pp.1193-
1197.
Dombrovski, M., Poussard, L., Moalem, K., Kmecova, L., Hogan, N., Schott, E., Vaccari, A.,
Acton, S. and Condron, B., 2017. Cooperative behavior emerges among *Drosophila*
larvae. *Current Biology*, *27*(18), pp.2821-2826.
Duménil, C., Woud, D., Pinto, F., Alkema, J.T., Jansen, I., Van Der Geest, A.M., Roessingh, S.
and Billeter, J.C., 2016. Pheromonal cues deposited by mated females convey social
information about egg-laying sites in *Drosophila melanogaster*. *Journal of chemical ecology*, *42*,
pp.259-269.
Fedina, T.Y., Kuo, T.H., Dreisewerd, K., Dierick, H.A., Yew, J.Y. and Pletcher, S.D., 2012.
Dietary effects on cuticular hydrocarbons and sexual attractiveness in *Drosophila*. *PloS one*,
7(12), p.e49799.
Ferveur, J.F., 2005. Cuticular hydrocarbons: their evolution and roles in *Drosophila* pheromonal
communication. Behavior genetics, *35*, pp.279-295.

Fisher, A.M., Le Page, S., Manser, A., Lewis, D.R., Holwell, G.I., Wigby, S. and Price, T.A.,
2021. Relatedness modulates density-dependent cannibalism rates in *Drosophila*. *Functional*
Ecology, *35*(12), pp.2707-2716.
Foley, B., Chenoweth, S.F., Nuzhdin, S.V. and Blows, M.W., 2007. Natural genetic variation in
cuticular hydrocarbon expression in male and female *Drosophila melanogaster*. *Genetics*, *175*(3),
pp.1465-1477.
Foucaud, J., Philippe, A.S., Moreno, C. and Mery, F., 2013. A genetic polymorphism affecting
reliance on personal versus public information in a spatial learning task in *Drosophila*
melanogaster. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1760), p.20130588.
Fowler, E.K., Leigh, S., Rostant, W.G., Thomas, A., Bretman, A. and Chapman, T., 2022.
Memory of social experience affects female fecundity via perception of fly deposits. *BMC*
biology, *20*(1), p.244.
Fox, J., and Weisberg, S., 2011. An {R} Companion to applied regression, 2nd edition.
Thousand Oaks CA: Sage.
García-Roa, R., Domínguez-Santos, R., Pérez-Brocal, V., Moya, A., Latorre, A. and Carazo, P.,
2022. Kin recognition in *Drosophila*: rearing environment and relatedness can modulate gut
microbiota and cuticular hydrocarbon odour profiles. *Oikos*, *2022*(4), p.e08755.
Golden, S. and Dukas, R., 2014. The value of patch-choice copying in fruit flies. *PloS one*, *9*(11),
p.e112381.
Gregg, T.G., McCrate, A., Reveal, G., Hall, S. and Rypstra, A.L., 1990. Insectivory and social
digestion in Drosophila. *Biochemical genetics*, *28*, pp.197-207.
Grueter, C. and Leadbeater, E., 2014. Insights from insects about adaptive social information
use. *Trends in ecology & evolution*, *29*(3), pp.177-184.
Guilhot, R., Xuéreb, A., Lagmairi, A., Olazcuaga, L. and Fellous, S., 2023. Microbiota
acquisition and transmission in *Drosophila* flies. *Iscience*, *26*(9).
Havula, E., Ghazanfar, S., Lamichane, N., Francis, D., Hasygar, K., Liu, Y., Alton, L.A.,
Johnstone, J., Needham, E.J., Pulpitel, T. and Clark, T., 2022. Genetic variation of
macronutrient tolerance in *Drosophila melanogaster*. *Nature communications*, *13*(1), p.1637.
Haerty, W., Jallon, J.M., Rouault, J., Bazin, C. and Capy, P., 2002. Reproductive isolation in
natural populations of *Drosophila melanogaster* from Brazzaville (Congo). *Genetica*, *116*,
pp.215-224.
Heard, S.B., 1994. Imperfect oviposition decisions by the pitcher plant mosquito (*Wyeomyia*
smithii). *Evolutionary Ecology*, *8*, pp.493-502.
Heys, C., Lizé, A., Colinet, H., Price, T.A., Prescott, M., Ingleby, F. and Lewis, Z., 2018.
Evidence that the microbiota counteracts male outbreeding strategy by inhibiting sexual
signaling in females. *Frontiers in Ecology and Evolution*, *6*, p.29.
Heys, C., Lizé, A., Lewis, Z. and Price, T.A., 2020. *Drosophila* sexual attractiveness in older males
is mediated by their microbiota. *Microorganisms*, *8*(2), p.168.
Hothorn, T., Bretz, F., & Westfall, P. 2008. Simultaneous inference in general parametric
models. *Biometrical Journal*. *50*(3), 346–363.
Joshi, A., Knight, C.D. and Mueller, L.D., 1996. Genetics of larval urea tolerance in *Drosophila*
melanogaster. *Heredity*, *77*(1), pp.33-39.
Kacsoh, B.Z., Bozler, J., Ramaswami, M. and Bosco, G., 2015. Social communication of
predator-induced changes in *Drosophila* behavior and germ line physiology. Elife, *4*,
p.e07423.

Klepsatel, P., Knoblochová, D., Girish, T.N., Dircksen, H. and Gáliková, M., 2020. The
influence of developmental diet on reproduction and metabolism in *Drosophila*. *BMC*
evolutionary biology, *20*, pp.1-15.
Khodaei, L. and Long, T.A.F., 2019. Kin recognition and co-operative foraging in *Drosophila*
melanogaster larvae. *Journal of evolutionary biology*, 32(12), pp.1352-1361.
Khodaei, L., Newman, T., Lum, S., Ngo, H., Maoloni, M. and Long, T.A.F. 2020. On the
expression of co-operative feeding behaviour in 3rd instar *Drosophila melanogaster* larvae!
bioRxiv: p.678631.
Khodaei, L. and Long, T.A.F, 2020. Kin recognition and egg cannibalism by *Drosophila*
melanogaster larvae. *Journal of Insect Behavior*, *33*(1), pp.20-29.
Leadbeater, E., 2009. Social learning: what do *Drosophila* have to offer? *Current Biology*, *19*(9),
pp.R378-R380.
Lefranc, A. and Bundgaard, J., 2000. The influence of male and female body size on copulation
duration and fecundity in *Drosophila melanogaster*. *Hereditas*, 132(3), pp.243-247.
LePage, S., Sepil, I., Flintham, E., Pizzari, T., Carazo, P. and Wigby, S., 2017. Male relatedness
and familiarity are required to modulate male-induced harm to females in *Drosophila*.
Proceedings of the Royal Society B: Biological Sciences, *284*(1860), p.20170441.
Lewis, Z., Heys, C., Prescott, M. and Lizé, A., 2014. You are what you eat: gut microbiota
determines kin recognition in *Drosophila*. *Gut microbes*, *5*(4), pp.541-543.
Lihoreau, M., Poissonnier, L.A., Isabel, G. and Dussutour, A., 2016. *Drosophila* females trade off
good nutrition with high-quality oviposition sites when choosing foods. *Journal of*
Experimental Biology, *219*(16), pp.2514-2524.
Lin, C.C., Prokop-Prigge, K.A., Preti, G. and Potter, C.J., 2015. Food odors trigger *Drosophila*
males to deposit a pheromone that guides aggregation and female oviposition decisions.
elife, *4*, p.e08688.
Lizé, A., McKay, R. and Lewis, Z., 2014. Kin recognition in *Drosophila*: the importance of
ecology and gut microbiota. *The ISME journal*, *8*(2), pp.469-477.
Long, T.A., Pischedda, A., Stewart, A.D. and Rice, W.R., 2009. A cost of sexual attractiveness to
high-fitness females. *PLoS biology*, 7(12), p.e1000254.
Malek, H.L. and Long, T.A.F. 2020. On the use of private versus social information in
oviposition site choice decisions by *Drosophila melanogaster* females. *Behavioural ecology*.
31(3):739-749.
Markow, T.A., 2015. The secret lives of *Drosophila* flies. *elife*, *4*, p.e06793.
Martin, E.S. and Long, T.A.F., 2015. Are flies kind to kin? The role of intra-and inter-sexual
relatedness in mediating reproductive conflict. *Proceedings of the Royal Society B: Biological*
Sciences, *282*(1821), p.20151991.
Mery, F., Varela, S.A., Danchin, É., Blanchet, S., Parejo, D., Coolen, I. and Wagner, R.H.,
2009. Public versus personal information for mate copying in an invertebrate. *Current*
Biology, *19*(9), pp.730-734.
Monier, M., Nöbel, S., Isabel, G. and Danchin, E., 2018. Effects of a sex ratio gradient on female
mate-copying and choosiness in *Drosophila melanogaster*. *Current Zoology*, *64*(2), pp.251-258.
Moreira-Soto, R.D., Khallaf, M.A., Hansson, B.S. and Knaden, M., 2024. How conspecific and
allospecific eggs and larvae drive oviposition preference in *Drosophila*. *Chemical Senses*, *49*,
p.bjae012.
Najarro, M.A., Sumethasorn, M., Lamoureux, A. and Turner, T.L., 2015. Choosing mates
based on the diet of your ancestors: replication of non-genetic assortative mating in
Drosophila melanogaster. *PeerJ*, *3*, p.e1173.

Nöbel, S., Danchin, E. and Isabel, G., 2018. Mate-copying for a costly variant in *Drosophila*
*melanogaste*r females. *Behavioral Ecology*, *29*(5), pp.1150-1156.
Nöbel, S., Monier, M., Fargeot, L., Lespagnol, G., Danchin, E. and Isabel, G., 2022. Female
fruit flies copy the acceptance, but not the rejection, of a mate. *Behavioral Ecology*, *33*(5),
pp.1018-1024.
Pavković-Lučić, S., Trajković, J., Miličić, D., Anđelković, B., Lučić, L., Savić, T. and Vujisić, L.,
2022. "Scent of a fruit fly": Cuticular chemoprofiles after mating in differently fed
Drosophila melanogaster (Diptera: Drosophilidae) strains. *Archives of Insect Biochemistry and*
Physiology, *109*(3), p.e21866.
Partridge, L., Ewing, A. and Chandler, A., 1987. Male size and mating success in *Drosophila*
melanogaster: the roles of male and female behaviour. *Animal Behaviour*, *35*(2), pp.555-562.
Prasad, N.G., Shakarad, M., Rajamani, M. and Joshi, A., 2003. Interaction between the effects
of maternal and larval levels of nutrition on pre-adult survival in *Drosophila melanogaster*.
Evolutionary Ecology Research, *5*(6), pp.903-911.
R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.
Reaume, C.J. and Sokolowski, M.B., 2006. The nature of *Drosophila melanogaster*. *Current Biology*,
16(16), pp.R623-R628.
Reynolds, J.D. and Gross, M.R., 1990. Costs and benefits of female mate choice: is there a lek
paradox?. *The American Naturalist*, *136*(2), pp.230-243.
Richard, F.J. and Hunt, J.H., 2013. Intracolony chemical communication in social insects.
Insectes sociaux, *60*, pp.275-291.
Robinson, S.P., Kennington, W.J. and Simmons, L.W., 2012. Assortative mating for relatedness
in a large naturally occurring population of *Drosophila melanogaster*. *Journal of Evolutionary*
Biology, *25*(4), pp.716-725.
Rose, M.R., 1984. Laboratory evolution of postponed senescence in *Drosophila*
melanogaster. *Evolution*, *38*(5), pp.1004-1010.
Sarin, S. and Dukas, R., 2009. Social learning about egg-laying substrates in fruitflies. *Proceedings*
of the Royal Society B: Biological Sciences, *276*(1677), pp.4323-4328.
Scott, D., Shields, A., Straker, M., Dalrymple, H., Dhillon, P.K. and Harbinder, S., 2011.
Variation in the male pheromones and mating success of wild caught *Drosophila melanogaster*.
PLoS one, *6*(8), p.e23645.
Sharon, G., Segal, D., Ringo, J.M., Hefetz, A., Zilber-Rosenberg, I. and Rosenberg, E., 2010.
Commensal bacteria play a role in mating preference of *Drosophila melanogaster*. *Proceedings of*
the National Academy of Sciences, *107*(46), pp.20051-20056.
Shoot, T.T., Miller, N.Y. and Long, T.A.F, 2024. Plasticity in expression of fruit fly larval
feeding clusters in response to changes in food quality and distribution. *Insectes Sociaux*, *71*,
pp.1-11.
Singer, T.L., 1998. Roles of hydrocarbons in the recognition systems of insects. *American Zoologist*,
38(2), pp.394-405.
Sullivan, W., Ashburner, M., & Hawley, R. 2000. *Drosophila* protocols. Cold Spring Harbor, NY:
Cold Spring Harbor Laboratory Press.
Toshima, N., Hara, C., Scholz, C.J. and Tanimura, T., 2014. Genetic variation in food choice
behaviour of amino acid-deprived *Drosophila*. *Journal of insect physiology*, *69*, pp.89-94.
Thorpe, W.H., 1945. The evolutionary significance of habitat selection. *The Journal of Animal*
Ecology, *14*(2), pp.67-70.

Valone, T.J. and Templeton, J.J., 2002. Public information for the assessment of quality: a
widespread social phenomenon. *Philosophical Transactions of the Royal Society of London. Series B:*
Biological Sciences, *357*(1427), pp.1549-1557.
Venu, I., Durisko, Z., Xu, J. and Dukas, R., 2014. Social attraction mediated by fruit flies'
microbiome. *Journal of Experimental Biology*, *217*(8), pp.1346-1352.
Vijendravarma, R.K., Narasimha, S. and Kawecki, T.J., 2013. Predatory cannibalism in
Drosophila melanogaster larvae. *Nature communications*, *4*(1), p.1789.
Wang, Y. and Staubach, F., 2018. Individual variation of natural *D. melanogaster*-associated
bacterial communities. *FEMS microbiology letters*, *365*(6), p.fny017.
Weiner, J., 2000. *Time, love, memory: A great biologist and his quest for the origins of behavior*. Vintage.
Wertheim, B., Dicke, M. and Vet, L.E., 2002a. Behavioural plasticity in support of a benefit for
aggregation pheromone use in *Drosophila melanogaster*. *Entomologia Experimentalis et Applicata*,
103(1), pp.61-71.
Wertheim, B., Marchais, J., Vet, L.E.M. and Dicke, M., 2002. Allee effect in larval resource
exploitation in *Drosophila*: an interaction among density of adults, larvae, and micro-
organisms. *Ecological Entomology*, *27*(5), pp.608-617.
Yang, D., 2018. Carnivory in the larvae of *Drosophila melanogaster* and other *Drosophila*
species. *Scientific Reports*, *8*(1), p.15484.
Young, Y., Buckiewicz, N. and Long, T.A.F., 2018. Nutritional geometry and fitness
consequences in *Drosophila suzukii*, the Spotted-Wing *Drosophila*. *Ecology and evolution*, *8*(5),
pp.2842-2851.
733

ACKNOWLEDGEMENTS

We would like to thank Simran Mann, Lukas Ghiglione, and Harleen Taneja of the #DrosLife

Lab for their fly-pushing, behavioral observations and camaraderie. Natasha B. Gallo provided

helpful feed-back and constructive comments. Ruby Lindsay and Michael Steeleworthy of the

Wilfrid Laurier University Library are thanked for their help with data archiving. T.A.F.L. was

funded with a Natural Sciences and Engineering Research Council Discovery grants (RGPIN-

2016-06133 and RGPIN-2022-03988). This work was conducted at Wilfrid Laurier University,

which exists on the traditional territory of the Neutral, Anishnawbe, and Haudenosaunee

peoples.

FIGURES

Figure 1. Boxplots illustrating data collected in the first experiment of a) the total number of wild-type (*red-eyed*) adult flies that eclosed from eggs laid by the focal *D. melanogaster* females on 5 different media dishes, b) the total number of *brown-eyed* adult flies that eclosed from those same 5 media dishes, which had been laid by the prior conspecific occupants on the dish. The boxes enclose the middle 50% of data (the inter-quartile range, IQR), with the thick horizontal line representing the location of median. Data points > ±1.5*IQR are designated as outliers. Whiskers extend to largest/smallest values that are not outliers, indicated as closed circles. The results of a Tukey HSD post-hoc tests comparing group mean is indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level. The control/unexposed dishes by definition had not had any prior exposure to a female capable of producing *brown-eyed* offspring, and thus those dishes never yielded any offspring with that phenotype.

Figure 2. Boxplots illustrating data collected in the second experiment of a) the number of
brown-eyed adult flies that eclosed from dishes that had been exposed to 0, 1 or 2 conspecific
females, b) the number of *wild-type* (red-eyed) adult flies that eclosed from the same dishes that
had been laid by the focal female *D. melanogaster.* Boxplot components as in Figure 1. The results
of a Tukey HSD post-hoc test comparing group mean is indicated by letters, where groups that
do not share the same letter are considered statistically different at the α=0.05 level. The
control/unexposed dishes by definition had not had any prior exposure to a female capable of
producing *brown-eyed* offspring, and thus those dishes never yielded any offspring with that
phenotype
754

Figure 3. Boxplots illustrating data collected in third experiment of a) the number of *brown-eyed* adult flies that eclosed from dishes that had been exposed to either an unrelated IV female (mated to a IV-*bwD* male), a unrelated IV-*bw* female (mated to a IV-*bwD* male), or neither b) the number of *wild-type* (red-eyed) adult flies that eclosed from the same dishes that had been laid by the focal female *D. melanogaster.* Boxplot components as in Figure 1. The control/unexposed dishes by definition had not had any prior exposure to a female capable of producing *brown-eyed* offspring, and thus those dishes never yielded any offspring with that phenotype. The results of a Tukey HSD post-hoc test comparing group mean is indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level.

756
757

Figure 4. Boxplots illustrating data collected in the fourth experiment of the number of *wild-type* (red-eyed) adult flies that had been laid by the focal female *D. melanogaster* that eclosed from dishes that had been exposed to either a related IV female (mated to a IV-*bwD* male), an unrelated IV female (mated to a IV-*bwD* male), or neither. Boxplot components as in Figure 1. The results of a Tukey HSD post-hoc test comparing group mean is indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level.

760 **SUPPLEMENTARY FIGURES**

Figure S1. Drawings illustrating **a)** the "mini-egg" chambers used to collect eggs and/or create media that exhibited social cues. **b)** an observation arena into which the focal ovipositing female was introduced and monitored. At the bottom of the chamber the lids that exhibited different kinds of social information were affixed. **c)** the arrangement of dishes in the first experiment. In this experiment there were four lids that exhibited social information and the 'control/unexposed' dish that contained no social information (as it had no prior exposure to flies). **d)** the arrangement of the two dishes with social information and the control dish in the second experiment. **e)** the arrangement of the two dishes with social information and the control dish in the third experiment. **f)** the arrangement of the two dishes with social information and the control dish in the fourth experiment.

Figure S2. Boxplots illustrating the cumulative number of observations made across 26 sessions in the first experiment in which a focal female *D. melanogaster* was observed on the surface of one of 5 different media dishes present in the arena in the first experiment. The boxes enclose the middle 50% of data (the inter-quartile range, IQR), with the thick horizontal line representing the location of median. Data points > ±1.5*IQR are designated as outliers. Whiskers extend to largest/smallest values that are not outliers, indicated as closed circles. The results of a Tukey HSD post-hoc test comparing group mean is indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level.

Figure S3. Boxplots illustrating the cumulative number of observations made across 26 sessions in the third experiment in which a focal female *D. melanogaster* was observed on the surface of one of 3 different media dishes present in the arena that had previously been exposed to either an unrelated IV female (mated to a IV-*bwD* male), an unrelated IV-*bw* female (mated to a IV-*bwD* male), or neither. Boxplot components as in Figure S2. The results of a Tukey HSD post-hoc test comparing group mean is indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level.

Figure S4. Boxplots illustrating the cumulative number of observations made across 26 sessions in the fourth experiment in which a focal female *D. melanogaster* was observed on the surface of one of 3 different media dishes present in the arena that had previously been exposed to either a related IV female (mated to a IV-*bwD* male), an unrelated IV female (mated to a IV-*bwD* male), or neither. Boxplot components as in Figure S2. The results of a Tukey HSD post-hoc test comparing group mean is indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level.

Figure S5. Boxplots illustrating the number of *brown-eyed* adult flies that eclosed from dishes that had been exposed to either an related IV female (mated to a IV-*bwD* male), a unrelated IV female (mated to a IV-*bwD* male), or neither in the fourth experiment. Boxplot components as in Figure S2. The control/unexposed dishes my definition had not had any prior exposure to a female capable of producing *brown-eyed* offspring, and thus those dishes never yielded any offspring with that phenotype.

Early Eclosion &
Larger Body Size

Later Eclosion &
Smaller Body Size

Eclosion time/Body size

Figure S6. Schematic diagram depicting hypothetical distribution of female *D. melanogaster* eclosion times in a mixture of eggs collected at the same time from the IV and the IV-*bw* populations (where we assume are randomly sampled, and that both populations have identical distributions, means and variances). While both subset means are the same, as there is a greater number of IV-*bw* eggs in the vial than IV eggs, the larger sample size of the former will likely contain individuals exhibiting a wider range of phenotypes than the latter. The unanticipated consequence of this being that when sampling for virgin female flies, there is a greater probability of colleting a suitable IV-*bw* before encountering a suitable IV female. As body size is negatively correlated with eclosion time (Partridge et al. 1987), and positively correlated with female fecundity (Lefranc and Bundgaard 2000, Long et al. 2009), it is conceivable that the IV-*bw* and the IV females collected and used as demonstrators differed in their capacity to produce offspring, resulting in the conspecific patterns seen in Figure 1b and 3b. It is worth noting that in our fourth experiment, where only *red-eyed* females were collected from vials containing both IV and IV-*bw* eggs that there was no significant difference in the number of demonstrator offspring that were collected (Figure S5), which is consistent with the hypothesis described above.

769