1 **On the use of kinship and familiarity associated social information in mediating**
2 ***Drosophila melanogaster* oviposition decisions.**
3
4 Emily Rakosy[1,2]*, Sanduni Talagala[1] & Tristan A.F. Long[1]
5
6 [1.] Department of Biology, Wilfrid Laurier University, Waterloo, Ontario, Canada
7 [2.] Department of Cell & Systems Biology, University of Toronto Mississauga, Ontario, Canada
8 * Corresponding author. Email: emily.rakosy@mail.utoronto.ca ORCID: 0000-0002-4491-3704
9

Abstract

11
Decisions where an individual lays their eggs are important, as the choice may affect their
offspring's survival and lifetime reproductive success. Information produced by conspecifics can
potentially be useful in decision-making as this "social information" may provide an energetically
cheaper means of assessing oviposition site suitability rather than acquiring it personally.
However, as not all public information may be equally beneficial, cues produced by kin may be
especially valuable as they might signal suitable microenvironments, and are associated with
other fitness advantages resulting from improved foraging success and/or a decreased risk of
competition/cannibalism compared to sites where unrelated conspecifics are located. Using the
fruit fly, *Drosophila melanogaster*, we explored whether public information use is associated with kin-
based egg-laying decisions. Kinship is potentially recognized in several ways, including
environmentally-associated proxy cues, so we explored whether there were biases in how focal
females interacted with cues from conspecifics that differed in both genetic relatedness, and
environmental "familiarity." In a series of inter-connected assays, we examined the behaviour of
focal females that interacted with a choice of potential egg-laying substrates that differed in the
manner of their prior conspecific exposure, and counted the offspring that eclosed from these
different substrates. Sites that had exhibited cues produced by conspecific demonstrators were
visited more, and yielded more focal offspring compared to unexposed substrates. Furthermore,
patterns of bias in offspring production were consistent with ovipositing females exhibiting
sensitivity to the kinship status of the prior substrate's occupants. The basis of the kinship
categorization by ovipositing females appears to be based on phenotypes that reflect true genetic
relatedness, but the nature of the social information can be affected by other factors. These
results further highlight the potential usefulness of *D. melanogaster* as a model to understand the
evolution of social behaviour in the expression of decision-making.
35
36 ***Keywords*:** Social information, *Drosophila melanogaster*, oviposition, conspecifics, kin recognition,
37 microenvironments, behaviour, copying, public information
38
39 ***Competing interests*:** The authors declare no competing interests.
40 ***Ethics*:** This study did not require approval from an ethics committee.
41 ***Supporting Data*:** All data analysed in this manuscript are archived in Borealis, the Canadian
42 Dataverse Repository: https://doi.org/10.5683/SP3/N7XBOJ

Introduction

In species where there is little-to-no post-natal parental care, decisions about where and when an individual should lay their eggs are likely very important, as there are potentially dire fitness outcomes for one's offspring if unsuitable oviposition sites are chosen (Thorpe 1945, Heard 1994, Lihoreau et al. 2016). In order to make better decisions, using cues produced by conspecifics (aka "social information" *sensu* Valone and Templeton 2002) may be of value to the individual as they may provide meaningful information about potential egg-laying sites that can be obtained at a lower energetic cost than having to explore and evaluate alternate options oneself. Social information can be valuable in reducing an individual's uncertainty about their environment, and may help them locate areas that are rich in important resources such as food, shelter, desirable mates, increased foraging success, or are at a lower risk of predation/parasitism, any of which may benefit the mimicking individual and/or their offspring (Sarin and Dukas 2009, Blanchet et al. 2010, Grüter and Leadbeater 2014). Furthermore, the energy savings of using social information over private information can also be redirected into increased provisioning and/or production of offspring, thereby enhancing the benefits of its use in making decisions (Golden and Dukas 2014).

While fruit flies, *Drosophila melanogaster*, have a long and storied use as a model in studies of genetics, development, behaviour and evolution (Weiner 2000, Brookes 2001), they are also emerging as a powerful model in which to study social learning and the use of social information (Leadbeater 2009, Chen and Sokolowski 2022). *D. melanogaster* has been found to use the visual, gustatory, tactile, and olfactory cues produced by conspecifics in a number of different decision-making contexts. For instance, flies use social cues when engaged in spatial learning tasks to improve the speed that a "safe zone" is detected (Foucaud et al. 2013). Virgin female flies that are exposed to deposits left by conspecific demonstrators exhibit lower post-mating egg-laying rates

than unexposed females, presumably because they are better able to anticipate the forthcoming

availability of oviposition sites and the potential degree of future larval competition (Fowler et al.

2022). Mate choice can involve social information use as female fruit flies are more likely to mate

with males exhibiting similar characteristics as a male they've observed successfully mating with

another female (Mery et al. 2009, Monier et al. 2018, Nöbel et al. 2018, Nöbel et al. 2022).

While the robustness of this phenomenon is debatable (Belkina et al. 2021) it may still be

advantageous because mate choice is often costly to females (Reynolds and Gross 1990), and

copying the decisions made by conspecifics represents an economical adaptive strategy (Monier

et al. 2018). Important information about environmental conditions can also be obtained

indirectly by flies interacting with conspecifics. In a study by Kacsoh et al. (2015) female flies

decreased their oviposition rate in response to exposure to parasitic wasps, and this defensive

change in egg-laying behaviour was adopted by inexperienced flies housed with the wasp-

exposed females, with cues about environmental risk communicated visually using the

demonstrator flies' wings. Thus, while *D. melanogaster* has not been considered a "social" species

(Chen and Sokolowski 2022), it is evident that fruit flies can, and do, use social information to

inform their decisions.

One important context where social information seems to be used extensively by *D.*

melanogaster females is in selecting suitable oviposition sites (Battesti *et al.* 2012, Durisko et al. 2014,

Sarin and Dukas, 2009, Golden and Dukas 2014, Malek and Long 2020, Moreira-Soto et al.

2024). In the wild, there may be considerable heterogeneity in the availability and quality of

oviposition sites that are available to female *D. melanogaster* (Reaume and Sokolowki 2006,

Markow 2015), and the choice of oviposition environment can have profound consequences for

their offspring's fitness (Prasad et al. 2003, Klepsatel et al. 2020). *D. melanogaster* females may gain

insight on the suitability of oviposition sites through direct observation of other females

ovipositing (Sarin and Dukas, 2009 and Battesti *et al.* 2012), the presence of larvae (Golden and

Dukas, 2014) and/or the presence of chemical cues produced by conspecifics (Venu et al. 2014,

Lin et al. 2015, Duménil et al. 2016, Moreira-Soto et al. 2024). The reliance upon social cues by

ovipositing *D. melanogaster* is not without potential pitfalls. First, by outsourcing one's information

gathering to others, there is the risk of mimicking other's bad decisions. For instance, in a study

by Golden and Dukas (2014) when female flies were given a choice of laying their eggs on a site

where nutrition was poor but contained social cues (the presence of larvae) or a site where the

media contained 3 times more nutrition but lacked conspecific cues, the ovipositing females

treated them as being of equal quality, even though any offspring laid on the former patches did

not have access to sufficient nutrients to survive. Similarly, Duménil et al. (2016) observed that

the presence of a mated female demonstrator on yeast-free media made this undesirable

oviposition site equally-valued as yeasted media (that has not been exposed to conspecifics) to

later ovipositing females. In *D. melanogaster* the widespread use of social information may arise

because there can be significant benefits of laying one's eggs at sites already chosen by

conspecifics as offspring growth and development may be enhanced through communal digestion

via secreted enzymes (Gregg, et al. 1990, Sakaguchi and Suzuki 2013), through increased local

beneficial microbial activity (Gilbert 1980, Wertheim et al. 2002a, Wertheim et al. 2002b),

and/or via improved larval group foraging (Dombrovski et al. 2017, Khodaei et al. 2020, Shoot

et al. 2024). However, laying one's eggs near those of conspecifics can also pose risks arising from

the competition for limited resources (De Jong 1976, Joshi et al. 1996, Golden and Dukas 2014),

increased exposure to toxic waste products (Borash et al. 1998), and the threat of cannibalism at

many developmental stages (Vijendravarma et al. 2013, Ahmed et al. 2015, Yang 2018).

Crucially, some of these dangers may potentially be mediated if females are able to oviposit at

sites containing kin, as *D. melanogaster* larvae are more co-operative (Khodaei and Long 2019),

and less cannibalistic (Khodaei and Long 2020, Fisher et al. 2021) towards close relatives,

compared to how they engage with unrelated conspecifics. Additionally, as there can be extensive

intra-specific genetic variation *D. melanogaster* for nutritional preferences and dietary tolerances

(Toshima et al. 2014, Havula et al. 2022), preferentially laying eggs near those of close relatives

may increase the likelihood of selecting the best micro-habitat for the success of one's own

offspring. Thus, if females were able to differentiate social information produced by kin from

those generated by other demonstrators, they might be able to achieve greater fitness benefits

(and avoiding some of the risks) by selectively copying the choices made by relatives.

When considering the possibility that individuals might differentially use social cues

produced by relatives versus non-relatives, one must also consider what (if any) cues could

indicate kinship (or could be inferred to be reliable proxies of kinship). In *D. melanogaster* such

information could conceivably be perceived from cuticular hydrocarbons (CHCs) and/or gut

bacteria left behind by conspecifics. CHCs are long-chain hydrocarbons that are synthesized in

insect oenocyte cells and become part of the cuticle's waxy coating, where they help prevent

water loss, as well as also playing an important role in olfactory communication related to species

recognition and mate choice decisions (Singer 1998, Ferveur 2005, Scott et al. 2011, Richard

and Hunt 2013, Chung and Caroll 2015, Moreira-Soto et al. 2024). CHCs may provide accurate

information about kinship as there exists a substantial genetic variation underlying these

phenotypic traits within *D. melanogaster* populations (Foley et al. 2007, Dembeck et al. 2015, Heys

et al. 2018). Recent work has revealed that female flies transfer their own CHCs (along with

male-specific compounds acquired during mating) to their laid eggs (Moreira-Soto et al. 2024),

which could conceivably be used as signals of potential kinship by subsequent ovipositing females.

In addition to CHC cues, olfaction-based decisions may be influenced by signals

associated with differences in an individual's bacterial community (Sharon et al. 2010, Najarro et

al. 2015). While the *D. melanogaster* microbiome is relatively simple compared to other hosts

(Broderick and Lemaitre 2012), substantial variation exists between individuals/families from the

same population in their microbial community composition (Wang and Staubach 2018, Heys et

al. 2018). Individuals acquire some of their microbial symbionts from their mothers, who transfer

bacteria to the surface of their eggs, and their larvae's subsequent consumption of their chorion

establishes their offspring's own gut flora (Bakula 1969, Guilhot et al. 2023). Thus, differences in

cues produced by bacterial communities could conceivably (and independently) be used to infer

kinship (Lizé et al. 2014, Lewis et al. 2014). An important and potentially complicating factor is

that CHC phenotypes and/or microbiotic communities in *D. melanogaster* can also be influenced

by the flies' developmental environment (Sharon et al. 2010, Fedina et al. 2012, Lizé et al. 2014,

Venu et al. 2014, Heys et al. 2018, García-Roa et al. 2022, Pavković-Lučić et al. 2022). This

raises the possibility that the phenomenon of "kin recognition" may be based on inherited cues of

relatedness, on the recognition of similarity of environmental histories ("familiarity"), or some

combination of these two factors, and that furthermore under some circumstances that these cues

might not provide contradictory or unreliable information about kinship. In a previous study that

examined whether kinship mediated the magnitude and consequences of male-male competition

in adult flies, it was found that reductions in male-induced harm were *only* seen in groups of males

that were both closely related to each other *and* 'familiar' by having developed in the same larval

environment (Le Page et al. 2017). Lizé et al. (2014) described how females were more likely to

mate with a familiar male, while males' ability to strategically avoid investing into mates with a

sibling was impeded if she had developed in the same environment. Similarly, Heys et al. (2018)

noted that the ability of fruit flies of both sexes to perceive the kinship status of their mate

appeared to be dependent on the state of the mate's microbiome – with females only acting

differently towards kin when those individuals had developed in the same media type (and had an

intact microbiota), while a male's increased investment into mating with an unrelated female only

occurred if she had been grown in a different environment (or whose microbiota had been

disrupted with the use of antibiotics). The complicated nature of the interaction between

developmental environment and relatedness was also explored by García-Roa et al. (2022) who

found that flies developing in different environments exhibited both different CHC and microbial

profiles, that related flies had microbiotic communities of similar diversity (but not composition),

and that diversity was correlated with CHC phenotypic profiles. Together these studies highlight

that socially-available kinship cues (or proxies thereof) may involve a number of independent and

inter-related olfactory signals, whose expression may depend not only on relatedness but also on

the specific nature of the developmental environment. As such, any study – such as ours – that

explores the potential role of kin recognition in decision making, must explore how individuals

respond to cues produced by demonstrators in which relatedness and familiarity are

independently manipulated.

With these ideas and findings in mind, in this study we set out to examine whether female

flies showed bias in their use of social information cues left behind by conspecifics when selecting

ovipositing sites, and would differ their use of egg-laying substrates that had previously been in

contact with either kin or non-kin demonstrator females. Simultaneously, we also explored

whether a female's response to potential kinship cues were conditional on the conspecifics'

"familiarity", by concurrently manipulating the developmental environment from which the

demonstrator females were obtained. Overall, the goal of this research is to better characterize

copying behavior in *D. melanogaster* and to gain better insight into the factors that influence these

oviposition site decisions.

Materials and Methods

Population Protocols and Fly Maintenance

All fruit flies used in our assays were obtained from one of three associated laboratory *D. melanogaster* populations: *Ives* (hereafter "IV"), "IV-*bw*", and "IV-*bwD*". The IV population was derived from a sample of mated females collected in 1975 from Amherst MA, USA sample population (Rose 1984). The IV-*bw* population was created by introgressing the brown-eye recessive allele, bw^1, into the genetic background of the IV population via 10 rounds of back-crossing, to create a competitor population that has >99.9% of the IV genetic background (Khodaei and Long 2020). The IV-*bwD* population was also created in the same fashion, by introducing the *bwD* allele (a dominant allele that also produces a brown-eyed phenotype) into an IV genetic background. The IV-*bw* and IV-*bwD* populations have been subject to periodic rounds of additional back-crossing to the IV population in an attempt to minimize their divergence from the base population, with the last session occurring approximately 50 generations prior to the start of this experiment. These three populations are each maintained at a census size of ~3500 adults per generation, incubated at 25°C incubation temperature with 60% humidity on a 12L:12D hour diurnal light cycle (Martin and Long 2015). Flies are maintained in standard *Drosophila* vials (95mm H x 28.5mm OD) containing 10mL of banana/agar/killed-yeast food, dusted with live yeast (Rose 1984, Shoot et al. 2024). Populations are cultured on a 14-day cycle, wherein every fortnight adult flies are removed from their natal vials, mixed *en masse* under light CO_2 anaesthesia and distributed into vials containing new media for up to 18h, before being removed and the eggs laid are then trimmed by hand to a density of approximately 100 eggs per vial (Martin and Long 2015).

Experiment 1: Test of the effects of conspecific relatedness and developmental environment on focal female behaviour and offspring production decisions.

209
210

In our first experiment, we collected adult male and female flies as virgins (within 8h of

eclosion from their pupae) from the IV population, and housed them individually. We then

created an initial 50 fly "families" by randomly combining single pairs of males and females into

"mini-egg" chambers (Kartell 733/4 polyethylene 20 mL sample vials; 74.5mm H x 24.8mm

OD; Fig S1a) whose lid contained juice/agar media (Sullivan et al. 2000) and was used as a

"dish" for ovipositing. We left these chambers in the incubator for ~18h, before removing the

adult flies and counting the number of eggs that had been laid overnight. We then evenly split

each family's eggs among 2 different vials each containing 10ml of media, and then added a

sufficient number of similarly-aged eggs obtained from the IV-*bw* population so that each vial

ultimately contained ~100 eggs apiece. We returned these pairs of vials to the incubator where

they were allowed to develop for the next 9 days, at which point we began collecting virgin

females for use in the experiment. For each pair of vials, we haphazardly designated one vial as

the "focal" vial, from which we collected 3 virgin females: a IV female ("the focal female"), a

second IV female (the focal female's sister, raised in the same vial), and one brown-eyed IV-*bw*

female (an unrelated "familiar" female, who had developed in the same vial as the focal female).

From the other ("non-focal") vial in the pair, we collected 2 virgin females: an IV female (the

focal female's sister, raised in a different "unfamiliar" developmental environment) and one

brown-eyed IV-*bw* female (an unrelated female, raised in environment that was "unfamiliar" to

the focal female). In this way we were able to obtain all possible combinations of demonstrator

females that differed in their genetic relatedness and their potential environmental familiarity.

We kept these females individually for up to 48h in test-tubes containing media to further verify

their virgin status (by the absence of any fertilized eggs). Next, we introduced two similarly-aged

adult IV males into the focal female's vial (so that all her offspring would express the wild-type

red-eye phenotype). At the same time, we transferred all other females into their own mini-egg

chambers whose dish lids contained standard banana media containing 3 IV-*bwD* males so that

all offspring produced by demonstrators would express a *brown-eyed* eye phenotype and to ensure

that any male-specific compounds transferred to the dish surface would be consistent across all

dishes (Lin et al. 2015, Moreira-Soto et al. 2024). Females and males were left overnight to

ensure sufficient time for mating to occur, and ovipositing to begin. On the following morning all

non-focal flies were removed from the mini-egg chambers, and sets of four dishes were affixed

using adhesive putty to the bottom of a larger chamber, (hereafter "arena"), which consisted of

an inverted plastic box (KIS Omni box, 20.3 x 15.9 x 9.6cm) modified by the addition of mesh

vent holes to box's outer edges (Fig S1b, Young et al. 2018). A fifth dish (that had never been

exposed to flies) was also placed in the arena before it was sealed shut, and the (lightly

anesthetized) focal female was introduced via an access hole. Thus the five dishes inside each

arena represent available oviposition sites for the focal female that potentially differ in their social

information: The 'related/same' dish contains cues produced by a sister that developed in the

same vial environment as the focal IV female; the 'unrelated/same' dish contains cues from an

unrelated IV-*bw* female that developed in the same vial environment as the focal IV female; the

related/different' dish contains cues produced by a sister that developed in a different vial

environment than the focal IV female; the 'unrelated/different' dish contains cues from an

unrelated IV-*bw* female that developed in a different vial environment as the focal IV female; and

finally the 'control/unexposed' dish contains no social information, as it had no prior exposure to

flies (Fig S1c). In total we established 36 replicate arenas, which were housed in a quiet, well-lit

room. Arenas were haphazardly rotated when placed on the counter to randomize any potential

directional room effects, and were left undisturbed for 20 minutes following the introduction of

the focal female. The arenas were then observed every 20 min for ~4.5h (starting at 10:30am),

and for 4h on the next day (beginning at 9:30am) for a total of 26 observation sessions. During

each observation session, the arenas were watched for a 5s time period, allowing for the

identification of which of the 5 dishes (if any) the focal IV female was located. The identity of the

different treatment dishes were unknown to the observers. Once all observations were complete,

we unsealed the arena, discarded the focal female and transferred the media in each dish into

vials containing 10m of media. These labelled vials were sealed and returned to the incubator for

2 weeks. At that time, all eclosed adults were removed, and their eye colours were tallied. Red-

eyed flies were the offspring of the focal female, while brown eye-flies (*wt+/bwD* or *bw$_1$/bwD*)

were produced by one of the conspecific females. We used the number of eclosed adult offspring

as a proxy for the egg number laid, in this and subsequent experiments, as in the IV population

at moderate larval densities, there is very high egg-to-adult survivorship (Khodaei et al. 2020).

Experiment 2: Test of the effect of conspecific offspring abundance on focal female
offspring production decisions.
Based on the offspring production patterns observed in Experiment 1, we performed a

series of complementary assays designed to test alternative hypotheses for the focal flies' potential

oviposition biases. Thus, in this experiment, we set out to examine whether the number of

offspring produced by conspecifics on a substrate was associated with the number offspring

subsequently producd on the same dish by a focal female. We began by collecting mated IV-*bw*

female flies from the lab's stock population and placed either 1, 2, or 0 individuals overnight into

mini-egg chambers containing media in their dish lid. The next morning, one of each type of the

dishes was affixed to the bottom of an arena (Fig S1d). Next, we added a single mated IV female

to each of the 48 replicate arenas and left them undisturbed for 22h, before transferring the

media in each dish to a vial containing 10ml of media. We returned those vials to the incubator

and counted the number of wild-type and brown-eye offspring that had eclosed 12 days later.

Experiment 3: Test of offspring production and behavioural biases of focal females towards cues from flies originating from the IV versus the IV-bw populations.

In our third experiment we explored the alternate hypothesis that the offspring production patterns observed in Experiment 1 might reflect preferences of the IV-originating focal female for substrates that had been in contact with any other IV flies (regardless of their degree of immediate genetic consanguinity) and/or the avoidance of substrates that had come in contact with any IV-*bw* flies. For this assay we set up replicate vials containing 5 mated IV and 5 mated IV-*bw* females, that were allowed to oviposit for 18h before being discarded, and the number of eggs laid standardized to 100 apiece. These vials were placed in the incubator, and we collected a virgin red-eyed and a virgin brown-eyed female from each vial as they eclosed (thus both flies had experienced the same developmental environment). Each female was then placed into their own mini egg chamber with 2 IV-*bwD* males overnight before being discarded, and the dish affixed to the inside of an arena, along with an unexposed/control dish (Fig S1e). An individual mated IV female obtained from the lab stock population was introduced to each of the arenas, and her location in the arena was periodically observed every 20 minutes over the course of 5h (for a total of 15 sessions). In total we established 72 replicate arenas, which were left undisturbed overnight, and the following morning media was removed from the dishes and transferred to vials containing 10ml of media. These vials were incubated for 12 days, and the number of offspring with different eye colours was recorded.

Experiment 4: Revisiting the effect of conspecific relatedness on offspring production and behavioural decisions.

Based on the outcomes of our second and third experiments, we set about to directly compare the behaviour and offspring production of focal females towards substrates that had been exposed to their sibling or to an unrelated IV-derived female. As in experiment 1 we

collected the focal female's parents from the IV population, mated them, placed them into mini-

egg chambers overnight, then counted and split their eggs amongst 2 vials, with additional

similarly-aged IV-*bw* eggs added to bring the density to a total of 100 eggs apiece. From each of

the pairs of vials we collected virgin females as they eclosed and kept them individually. One of

the vials was arbitrarily designated as that which would yield the focal female (who was collected

as a virgin, then housed with 2 IV males from the stock population overnight), and virgin IV

female from the other vial (*i.e.* the focal female's sister raised in a different environment) was

housed with 2 IV-*bwD* males overnight in a mini egg chamber. We also haphazardly chose one of

the virgin IV females collected from one of the other sets of vials (thus an unrelated female, raised

in a different environment) and housed her with 2 IV-*bwD* males overnight in a mini egg

chamber. These dishes (along with a control/unexposed dish) were placed into an arena (Fig

S1f), into which we added the mated focal females. In each of the 84 replicate arenas, the

location of the focal female was periodically observed every 20 minutes over the course of 5h (for

a total of 15 sessions). Next, the chamber was left undisturbed overnight and the following

morning, we transferred the media from each dish into a new vial containing 10ml of media

which were incubated for an additional 12 days. The eye colour of all adult flies present in these

vials were tallied and recorded.

Statistical Analyses
All data analyses were conducted in the R statistical computing environment (version 4.0.3, R

Core Team, 2020). In all of our experiments we compared the frequencies of visits by focal

females to the five different lids in each arena. As these data were zero-inflated (determined using

diagnostic functions in the *DHARMa* package (Hartig 2022)), we constructed zero-inflated

binomial mixed models using the *glmmTMB* function in the *glmmTMB* package (Brooks 2017)

where the independent fixed factor was the dish treatment category, and the arena that

contained them was included as a random effect. We determined whether the treatment means

were significantly different from each other with likelihood-ratio chi-square tests using the *Anova*

function in the *car* package (Fox and Weisberg 2011), which was followed, if necessary, by post-

hoc tests with Tukey method p-value adjustments executed using the *emmeans* function in the

emmeans package (Lenth 2022) to determine the specific location of differences between

treatments. For experiment 1 we also used a two-way, zero-inflated *glmmTMB* model to

determine if there was any difference in patch visitation that varied with relatedness and/or

familiarity treatments, by using the same response data (excluding observations made on the

control dish, as this treatment by definition had received no prior conspecific contact). The

demonstrator female's degree of kinship, their environmental origin, and the interaction between

these two factors were the independent fixed factors in this GLMM model, and arena was the

random effect. The significance of these effects was determined using a likelihood-ratio chi-

square test using the *Anova* function.

For all of our experiments we (separately) analyzed the total number of red-eyed and brown-

eyed adults that successfully eclosed from the different lids in each arena using zero-inflated

glmmTMB GLMMs, with poisson error distributions, where the independent fixed factor was the

dish treatment category, and arena was the random effect. For each model we determined

whether the treatment means were significantly different from each other using likelihood-ratio

chi-square tests implemented using the *Anova* function, and post-hoc tests were executed, if

necessary, using the *emmeans* function, with Tukey method p-value adjustments, to determine the

specific location of differences between treatments. As above for experiment 1 data, we also set

out to determine if there was any difference in offspring number associated with relatedness

and/or familiarity treatments, by excluding data collected from the control dish treatment, and

analyzing the rest of the data set using a 2-way zero-inflated *glmmTMB* GLMM with poisson

errors. The demonstrator female's degree of kinship, their environmental origin, and the

interaction between these two factors were the independent fixed factors in this model, and arena

was included as a random effects. The significance of these effects was determined using a

likelihood-ratio chi-square test using the *Anova* function, and, if necessary, post-hoc tests were

conducted using the *emmeans* function, with Tukey method p-value adjustments, to determine the

specific location of differences between treatments.

In experiment 2 we also compared the frequencies of dishes that yielded zero focal offspring

using the *chisq.test* function to conduct a Pearson's Chi-squared test, then examined the specific

differences between cells in the contingency table using the *chisq.posthoc.test* function in the

chisq.posthoc.test package (Ebbert, 2019).

Results

Experiment 1: Test of the effects of conspecific relatedness and developmental
environment on focal female behaviour and offspring production decisions.
In our first experiment, we compared the cumulative number of visits that *D. melanogaster*

focal females made to five different dishes, as well as the number of offspring that eclosed as

adults from each of those dishes. Our analysis found that those females exhibited heterogeneity in

their dish associations (LRχ^2 =20.12, df=4, p=0.0005), and while they visited the

control/unexposed dishes less frequently than the demonstrator dishes, within these four

treatment groups, they did not express any significant bias (Fig S2). Offspring production

patterns indicated that the females' oviposition patterns were also heterogeneous (Fig 1a; LRχ^2

=123.92, df=4, p=7.77x10^{-26}), with significantly more red-eyed focal offspring eclosing from

those dishes that had previously been in contact with a related demonstrator individual than from

either the control/unexposed dish or either of the dishes whose prior occupant was an unrelated

individual, and the greatest number of offspring eclosing from those dishes that had been exposed

to a related conspecific, raised in a different vial. Our focused two-way GLMM confirmed that

the amount of oviposition activity on a dish was significantly associated with the demonstrator's

kinship (LRχ^2 =84.01, df=1, p=4.91x10^{-20}) with more offspring eclosing from 'related' dishes

($\bar{x}$=4.97; 95% CI: 3.57, 6.92) than 'unrelated' dishes ($\bar{x}$=0.19; 95% CI: 0.08, 0.42), there was a

small but significant effect of developmental environment (LRχ^2 =9.38, df=1, p=0.0021), with

slightly more offspring eclosing from 'different' dishes ($\bar{x}$=1.08; 95% CI: 0.57, 1.84) than 'same'

dishes ($\bar{x}$=0.93; 95% CI: 0.52, 1.67), while the interaction between kinship and developmental

environment was not significant (LRχ^2 =1.20, df=1, p=0.27). We also noted that the number of

demonstrator offspring were not homogeneous across the four treatment groups (Fig 1b), with

more brown-eyed offspring eclosing from dishes whose prior occupant was a unrelated female

(IV-*bw*, mated to a IV-*bwD* male; $\bar{x}$=14.49; 95% CI: 13.03, 16.10) than those that had previously

been exposed to a related female (IV, mated to a IV-*bwD* male; $\bar{x}$=6.83; 95% CI: 5.95, 7.83;

LRχ^2 =147.93, df=1, p=4.90x10^{-34}), and fewer offspring eclosing from vials where the

demonstrator originated from the same vial ($\bar{x}$=8.55; 95% CI: 7.53, 9.71) than if they were

collected from a different vial ($\bar{x}$=11.57; 95% CI: 10.28, 13.02; LRχ^2 =30.83, df=1, p=2.82x10^{-8}

), yet there was no significant interaction between these two factors (LRχ^2 =0.001, df=1,

p=0.975).

Experiment 2: Test of the effect of conspecific offspring abundance production on focal
female offspring production decisions.
403   In our second experiment, we set out to test the alternate hypothesis that the biased

patterns of *red-eyed* offspring produced by our focal females was influenced by the abundance of

brown-eyed conspecific offspring already present on the dishes. In this assay either zero, one or two

demonstrator females were exposed to a dish prior to being offered as a potential oviposition site

to a mated focal female. More demonstrators on a dish yielded more *brown-eyed* offspring: zero

brown-eyed offspring eclosed from the control dishes, and two demonstrator females produced, on

average more *brown-eyed* offspring ($\overline{x}$=13.53; 95% CI: 11.2, 16.3) than a single demonstrator

female ($\overline{x}$ =8.68; 95% CI: 7.0, 10.8; LRχ^2 =36.14, df=1, p=1.84x10^{-9}; Fig 2a). Despite the

dramatic differences in the number of demonstrator offspring that eclosed between the three

dishes, there was no significant difference in the number of *red-eyed* focal offspring that eclosed

from the three dishes (LRχ^2 =1.99, df=2, p=0.371; Fig 2b). Interestingly, dishes where there had

been two demonstrator females has been present were significantly *less* likely to yield zero focal

female offspring (8/48) than those that had been exposed to one demonstrator females (25/48) or

those control dishes that lacked any demonstrator exposure (32/48; χ^2 =13.43, df=2, p=0.0012).

Experiment 3: Test of offspring production and behavioural biases of focal females
towards cues from flies originating from the IV versus the IV-bw populations.
420   In our third experiment, we set out to test the alternate hypothesis that the patterns of *red-*

eyed offspring produced by our focal females was influenced by the population of origin of the

demonstrator females, and not their degree of kinship. This involved comparing the cumulative

number of visits that *D. melanogaster* focal females made to three different dishes (a

control/unexposed dish, a dish where the demonstrator was an unrelated IV female, or a dish

whose demonstrator was an unrelated IV-*bw* female), as well as the number of offspring that

eclosed as adults from each of those dishes. Our analysis found that focal females exhibited

heterogeneity in their dish associations (LRχ^2 =144.8, df=2, p=3.61 x10^{-32}) with the

control/unexposed dishes receiving significantly fewer visits ($\overline{x}$=0.05; 95% CI: 0.03, 0.07), than

either the dishes that had been exposed to the unrelated IV demonstrator ($\bar{x}$=0.31; 95% CI: 0.25,

0.36) or the dishes that had been exposed to the unrelated IV-*bw* demonstrator ($\bar{x}$=0.25; 95% CI:

0.21, 0.31), with no significant difference in visitation rates to the two demonstrator dishes

(t=21.96, df=1, p=0.07; Fig S3). The distribution of focal female offspring was also heterogenous

across the three dish types (Fig 3b; LRχ^2 =139.27, df=2, p=5.72 x10^{-31}). More focal females

offspring eclosed from the unrelated IV dishes ($\bar{x}$=8.84; 95% CI: 7.45, 10.49) than they from

either the control/unexposed dishes ($\bar{x}$=2.32; 95% CI: 1.49, 3.61), or the unrelated *IV-bw* dishes

($\bar{x}$=3.09; 95% CI: 2.44, 3.91). The mean number of *red-eyed* offspring that were collected from the

IV-bw dish was not significantly different from the number collected from the control/unexposed

dish (t=1.31, df=1, p=0.39). We also noted that the number of eclosed demonstrator offspring

were significantly different between dish types (Fig 3a; LRχ^2 =89.42, df=1, p=3.82 x10^{-20}). with

more brown-eyed offspring eclosing from vials whose prior occupant was an unrelated IV female

(mated to a IV-*bwD* male; $\bar{x}$=12.33; 95% CI: 10.83, 14.05) than those that that had previously

been exposed to an unrelated IV-*bw* female (also mated to a IV-*bwD* male; $\bar{x}$=6.53; 95% CI:

5.55, 7.69).

Experiment 4: Revisiting the effect of conspecific relatedness on offspring production and
behavioural decisions.

In our fourth experiment, we set out (again) to compare the patterns of dish visitation and

offspring production by focal females on three different dishes (a control/unexposed dish, a dish

whose demonstrator was a related IV female, or a dish where the demonstrator was an unrelated

IV female). The focal females exhibited bias on their dish visitation rates (Fig S4; LRχ^2 =12.81,

df=2, p=0.0017), as they were significantly fewer visitations to the control/unexposed dishes

($\bar{x}$=0.05; 95% CI: 0.03, 0.07), than to related IV demonstrator dishes ($\bar{x}$=0.08; 95% CI: 0.06,

0.12) or to the unrelated IV dishes ($\bar{x}$=0.11; 95% CI: 0.08, 0.14). There was no significant

difference in visitation rates between the two demonstrator groups (t=1.33, df=1, p=0.38). The

distribution of focal female offspring was heterogenous across the three dish types (Fig 4; GLM:

LRχ^2 =368.28, df=2, p=1.07x10^{-80}). Significantly more focal female offspring eclosed from the

related IV dishes ($\overline{x}$=11.46; 95% CI: 9.96, 13.19) than from the unrelated IV dishes ($\overline{x}$=7.47;

95% CI: 6.43, 8.68), and both demonstrator dishes yielded more focal female offspring than did

the control/unexposed dishes ($\overline{x}$=0.48; 95% CI: 0.33, 0.69). There was a small, but statistically

significant difference in the number of demonstrator offspring eclosing from the related IV dishes

($\overline{x}$=11.3; 95% CI: 10.32, 12.4) and from the unrelated IV dishes ($\overline{x}$=10.1; 95% CI: 9.14, 11.1),

LRχ^2 = 6.59, df=1, p=0.01; Fig S5).

Discussion

The decision where to oviposit may be very important, especially in those species where

there is no post-laying parental care (Prasad et al. 2003, Klepsatel et al. 2020). Fruit flies,

Drosophila melanogaster, like many other species, use social information in their decision-making

(Leadbeater 2009, Chen and Sokolowski 2022). While there are benefits of laying one's eggs in

proximity to those of others, there are also risks – both of which might be mediated by the degree

of kinship between conspecifics. In this study, we set out to examine the possibility that female

flies might selectively use cues produced by close relatives, presumably to increase their

offspring's chance of success. In doing so, we simultaneously explored the factors that might

influence how flies might infer kinship, as well as investigate alternate hypotheses about social

information usage. Our assays confirm the extensive use of social cues by *D. melanogaster* females,

and furthermore collectively suggest that they are capable of kinship discrimination and are able

of distinguishing between conspecifics from different populations. These results add to the

growing body of knowledge about the sophisticated social information use by this model species

(Chen and Sokolowski 2022).

In all assays focal females associated more with those substrates that had come into

contact with conspecifics than with the unexposed/control dishes (Figs S2, S3 and S4), clearly

illustrating the importance of social cues in their decision-making, and consistent with numerous

previous studies (*e.g.* Battesti *et al.* 2012, Duménil et al. 2016, Malek and Long 2020). However,

not all conspecific cues elicited the same responses. In the first assay, we compared the offspring

production patterns of focal females on dishes exposed to a range of different demonstrators. By

splitting sets of full sibling eggs between two different vials to develop, and collecting both adult

relatives and non-relatives from each of the pairs of vials, we set out to test if and how *D.*

melanogaster females might distinguish kin cues. We observed that dishes that had been in previous

contact with a related demonstrator female yielded considerably more focal female offspring that

those where the demonstrator was non kin. Interestingly, dishes where the demonstrator came

from the same vial as the focal vial produced fewer focal female offspring than those when the

demonstrator had developed trh same vial, however the size of this difference was much smaller,

and there was no significant interaction between demonstrator kinship and origin factors (Fig 1a).

These results suggest that ovipositing females are selectively use social information, and largely

favouring cues produced by their relatives, while perceived environmental familiarity does not

enhance oviposition site use. Determining the specific cues being used by focal females is beyond

the scope of this study as demonstrator females transfer their gut bacteria, CHCs (and male-

specific compounds acquired during copulation) to the surface of their eggs and the media

(Bakula 1969, Guilhot et al. 2023, Moreira-Soto et al. 2024). The expression of these olfactory

cues can be influenced by both an individual's genotype and/or their developmental

environment (Sharon et al. 2010, Fedina et al. 2012, Lizé et al. 2013, Venu et al. 2014, Heys et

al. 2018, García-Roa et al. 2022, Pavković-Lučić et al. 2022). While the pattern of significant

differences associated with the developmental environment (Fig 1a) could suggest that

"familiarity" (and by extension the microbiotic cues that could be used as proxies of kinship) are

unimportant, and potentially misleading, cues of "relatedness", it should be noted that in our

experimental design, the conditions of the two vial environments were similar (same media, same

larval density), which may have minimized the potential for the development of differences in

these cues. Interestingly, Malek and Long (2020) observed that a focal female's offspring

production patterns depended on the developmental environment of *both* the focal female and the

demonstrator. In one assay, they observed focal females who developed in a protein-rich vial

environment, oviposited more on dishes where the demonstrator females (that came from the

same population) who had also developed on similar protein-rich media. However, in a second

assay when the focal females developed on standard media, these flies laid fewer eggs on dishes

where the demonstrator came from the same population, but had instead developed on protein-

rich media, than they did on dishes where the demonstrator came from a different *D. melanogaster*

population but had *also* developed on standard media. Thus, one should not necessarily conclude

that our results indicate that inherited cues are the only way in which egg kinship was/can be

recognized by the focal females in this species, and future studies exploring this idea across a

range of divergent environments are likely to expand our understanding of how social

information is produced and interpreted in *D. melanogaster*.

While the observed pattern of focal female offspring production is consistent with kin-

associated biases, other processes could also have resulted in the differential outcomes, and so in

order to validate or rule-out these alternate hypotheses we conducted several complementary

assays. We had been initially struck with the distinctive pattern of offspring production by the

demonstrator females (Fig 1b), wherein there were more *brown-eyed* conspecific offspring collected

from dishes where the demonstrator was unrelated than when the demonstrator was related.

Perhaps our focal females weren't deciding to lay their eggs on dishes that contained their kin's

offspring, but instead were avoiding dishes where there were already lots of eggs (and a

heightened risk of competition and/or cannibalism)? We explored this possibility in our second

assay, where focal females could choose between dishes with dramatically different number of

demonstrator offspring (Fig 2a). If females were avoiding dishes where there were too many

conspecific eggs already present, we would expect to see more focal offspring eclosing on the

"one female" treatment than on the "two females" treatment. In fact, we observed no significant

difference in focal female offspring production between the two groups (and if anything, there

were fewer cases where we observed *zero* focal female offspring produced when there were more

demonstrators; Fig 2b). Previous work by Golden and Dukas (2014) found that female flies given

a choice between a control and a demonstrator dish (which had been exposed to 5 or 20 larvae),

did not appear to discriminate against the demonstrator dishes that possessed cues indicating

greater competition during offspring development. Thus, we are fairly confident that the

differences in focal female offspring production patterns observed in experiment one were not

due to differences in the quantity of demonstrator cues present on a dish. As a side note, we

suggest that the differences in the number of demonstrator offspring observed in that assay is a

side effect of the way in which our demonstrator females were collected (*see* Fig S6 *caption for*

details).

The second alternate hypothesis we explored was that the biased pattern of oviposition

made by the focal females in the first experiment on the two "related" dishes, was not due to the

perceived kinship of the demonstrators, but instead was because those demonstrators originated

from the same population (IV), and the demonstrators for the two "unrelated" dishes were from a

different population (IV-*bw*). While IV and IV-*bw* are maintained following the same protocols,

and IV-*bw* is periodically back-crossed to IV, perceptible differences in the cues present in these

isolated populations (or their microbiotic communities) may have arisen via drift and/or

selection. Thus in our third experiment, we compared the behaviour and offspring production of

focal females exposed to (unrelated) IV or IV-*bw* demonstrator females. We found that focal

females produced more offspring on the former dishes than they did on the latter (and virtually

none on the control/unexposed dishes). This suggests that at least some of the patterns of

offspring production seen in our first experiment may not necessarily be the result of recognition

of kinship *per se*, but instead with other cue(s) that differ between the IV and IV-*bw* populations.

Thus, based on the insights gained in our third assay, we set out to re-examine the social

information use of *D. melanogaster* females in our fourth assay where females were given a choice

of oviposition sites that either had no prior exposure (control), a demonstrator sister (raised in a

different vial), or an unrelated demonstrator female (also from the IV population and raised in a

different vial). We observed that focal females still biased their offspring towards the dishes that

had previously been in contact with their sisters, albeit to a lesser extent than seen in experiment

1. This provides additional support to the hypothesis that female *D. melanogaster* are both capable

of distinguishing between cues produced by kin and non-kin, and that these cues are used in their

social decision-making. In doing so, they may be trying to obtain the potential benefits of

developing in a group (Gilbert 1980, Gregg, et al. 1990, Sakaguchi and Suzuki 2013, Wertheim

et al. 2002a, Wertheim et al. 2002b, Dombrovski et al. 2017, Shoot et al. 2024), while reducing

the potential risks from conspecifics (Khodaei and Long 2019, Khodaei and Long 2020, Fisher et

al. 2021).

570   Collectively, our results illustrate the subtle and plastic nature of social information use by

female *D. melanogaster*: Given a choice, a mated female *D. melanogaster* will be more likely to

produce offspring on a dish that has previously been in contact with any conspecific, compared

to an socially unexposed surface, there will be more offspring production associated with

substrates that have been in previous contact with an unrelated mated female from the same

population than one exposed to a mated female from a different population, and that focal if a

substrate has been exposed to a mated sister, that site will yield more focal offspring than if the

demonstrator is an unrelated individual. At the same time, mated female *D. melanogaster* do not

appear to show any bias in their egg production against dishes that differ in the number of

conspecific eggs (and their associated biochemical cues) present. Together, these pattens

illuminate that *D. melanogaster* females are sensitive to differences in the cues produced by

conspecifics and add to the growing appreciation of social decision-making in this model species.

References

Ahmad, M., Chaudhary, S.U., Afzal, A.J. and Tariq, M., 2015. Starvation-induced dietary
behaviour in *Drosophila melanogaster* larvae and adults. *Scientific reports*, *5*(1), p.14285.
Bakula, M., 1969. The persistence of a microbial flora during postembryogenesis of *Drosophila*
melanogaster. *Journal of Invertebrate Pathology*, 14(3), pp.365-374.
Battesti, M., Moreno, C., Joly, D. and Mery, F., 2012. Spread of social information and
dynamics of social transmission within *Drosophila* groups. *Current Biology*, 22(4), pp.309-313.
Battesti, M., Moreno, C., Joly, D. and Mery, F., 2015. Biased social transmission in *Drosophila*
oviposition choice. *Behavioral Ecology and Sociobiology*, *69*, pp.83-87.
Belkina, E.G., Shiglik, A., Sopilko, N.G., Lysenkov, S.N. and Markov, A.V., 2021. Mate choice
copying in *Drosophila* is probably less robust than previously suggested. *Animal*
Behaviour, *176*, pp.175-183.
Blanchet, S., Clobert, J. and Danchin, E., 2010. The role of public information in ecology and
conservation: an emphasis on inadvertent social information. *Annals of the New York Academy*
of Sciences, *1195*(1), pp.149-168.
Borash, D.J., Gibbs, A.G., Joshi, A. and Mueller, L.D., 1998. A genetic polymorphism
maintained by natural selection in a temporally varying environment. *The American*
Naturalist, *151*(2), pp.148-156.
Broderick, N.A. and Lemaitre, B., 2012. Gut-associated microbes of *Drosophila melanogaster*. *Gut*
Microbes, *3*(4), pp.307-321.
Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Casper W. Berg, C.W.,
Nielsen, A., Skaug, H.J., Maechler, M. and Bolker, B.M., 2017. glmmTMB Balances
Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed
Modeling. *The R Journal*, *9*(2), 378-400.
Brookes, M., 2002. *Fly: The unsung hero of twentieth-century science*. Harper Collins.
Chen, M. and Sokolowski, M.B., 2022. How social experience and environment impacts
behavioural plasticity in *Drosophila*. *Fly*, *16*(1), pp.68-84.
Chung, H. and Carroll, S.B., 2015. Wax, sex and the origin of species: dual roles of insect
cuticular hydrocarbons in adaptation and mating. *BioEssays*, *37*(7), pp.822-830.
De Jong, G., 1976. A model of competition for food. I. Frequency-dependent viabilities. *The*
American Naturalist, *110*(976), pp.1013-1027.
Dembeck, L.M., Böröczky, K., Huang, W., Schal, C., Anholt, R.R. and Mackay, T.F., 2015.
Genetic architecture of natural variation in cuticular hydrocarbon composition in *Drosophila*
melanogaster. *eLife*, *4*, p.e09861.
Durisko, Z., Anderson, B. and Dukas, R., 2014. Adult fruit fly attraction to larvae biases
experience and mediates social learning. *Journal of Experimental Biology*, *217*(7), pp.1193-
1197.
Dombrovski, M., Poussard, L., Moalem, K., Kmecova, L., Hogan, N., Schott, E., Vaccari, A.,
Acton, S. and Condron, B., 2017. Cooperative behavior emerges among *Drosophila*
larvae. *Current Biology*, *27*(18), pp.2821-2826.
Duménil, C., Woud, D., Pinto, F., Alkema, J.T., Jansen, I., Van Der Geest, A.M., Roessingh, S.
and Billeter, J.C., 2016. Pheromonal cues deposited by mated females convey social
information about egg-laying sites in *Drosophila melanogaster*. *Journal of Chemical Ecology*, *42*,
pp.259-269.
Ebbert, D., 2019. chisq.posthoc.test: A Post Hoc Analysis for Pearson's Chi-Squared Test for
Count Data. R package version 0.1.2. https://CRAN.R-
project.org/package=chisq.posthoc.test

Fedina, T.Y., Kuo, T.H., Dreisewerd, K., Dierick, H.A., Yew, J.Y. and Pletcher, S.D., 2012.
Dietary effects on cuticular hydrocarbons and sexual attractiveness in *Drosophila*. *PloS one*,
7(12), p.e49799.
Ferveur, J.F., 2005. Cuticular hydrocarbons: their evolution and roles in *Drosophila* pheromonal
communication. *Behavior genetics*, *35*, pp.279-295.
Fisher, A.M., Le Page, S., Manser, A., Lewis, D.R., Holwell, G.I., Wigby, S. and Price, T.A.,
2021. Relatedness modulates density-dependent cannibalism rates in *Drosophila*. *Functional
Ecology*, *35*(12), pp.2707-2716.
Foley, B., Chenoweth, S.F., Nuzhdin, S.V. and Blows, M.W., 2007. Natural genetic variation in
cuticular hydrocarbon expression in male and female *Drosophila melanogaster*. *Genetics*, *175*(3),
pp.1465-1477.
Foucaud, J., Philippe, A.S., Moreno, C. and Mery, F., 2013. A genetic polymorphism affecting
reliance on personal versus public information in a spatial learning task in *Drosophila
melanogaster*. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1760), p.20130588.
Fowler, E.K., Leigh, S., Rostant, W.G., Thomas, A., Bretman, A. and Chapman, T., 2022.
Memory of social experience affects female fecundity via perception of fly deposits. *BMC
Biology*, *20*(1), p.244.
Fox, J., and Weisberg, S., 2011. An {R} Companion to applied regression, 2nd edition.
Thousand Oaks CA: Sage.
García-Roa, R., Domínguez-Santos, R., Pérez-Brocal, V., Moya, A., Latorre, A. and Carazo, P.,
2022. Kin recognition in *Drosophila*: rearing environment and relatedness can modulate gut
microbiota and cuticular hydrocarbon odour profiles. *Oikos*, *2022*(4), p.e08755.
Golden, S. and Dukas, R., 2014. The value of patch-choice copying in fruit flies. *PloS one*, *9*(11),
p.e112381.
Gregg, T.G., McCrate, A., Reveal, G., Hall, S. and Rypstra, A.L., 1990. Insectivory and social
digestion in Drosophila. *Biochemical Genetics*, *28*, pp.197-207.
Grueter, C. and Leadbeater, E., 2014. Insights from insects about adaptive social information
use. *Trends in Ecology & Evolution*, *29*(3), pp.177-184.
Guilhot, R., Xuéreb, A., Lagmairi, A., Olazcuaga, L. and Fellous, S., 2023. Microbiota
acquisition and transmission in *Drosophila* flies. *iScience*, *26*(9).
Havula, E., Ghazanfar, S., Lamichane, N., Francis, D., Hasygar, K., Liu, Y., Alton, L.A.,
Johnstone, J., Needham, E.J., Pulpitel, T. and Clark, T., 2022. Genetic variation of
macronutrient tolerance in *Drosophila melanogaster*. *Nature Communications*, *13*(1), p.1637.
Haerty, W., Jallon, J.M., Rouault, J., Bazin, C. and Capy, P., 2002. Reproductive isolation in
natural populations of *Drosophila melanogaster* from Brazzaville (Congo). *Genetica*, *116*,
pp.215-224.
Heard, S.B., 1994. Imperfect oviposition decisions by the pitcher plant mosquito (*Wyeomyia
smithii*). *Evolutionary Ecology*, *8*, pp.493-502.
Heys, C., Lizé, A., Colinet, H., Price, T.A., Prescott, M., Ingleby, F. and Lewis, Z., 2018.
Evidence that the microbiota counteracts male outbreeding strategy by inhibiting sexual
signaling in females. *Frontiers in Ecology and Evolution*, *6*, p.29.
Heys, C., Lizé, A., Lewis, Z. and Price, T.A., 2020. *Drosophila* sexual attractiveness in older males
is mediated by their microbiota. *Microorganisms*, *8*(2), p.168.
Hartig, F., 2022. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. R package version 0.4.5. https://CRAN.R-
project.org/package=DHARMa
Joshi, A., Knight, C.D. and Mueller, L.D., 1996. Genetics of larval urea tolerance in *Drosophila
melanogaster*. *Heredity*, *77*(1), pp.33-39.

Kacsoh, B.Z., Bozler, J., Ramaswami, M. and Bosco, G., 2015. Social communication of
predator-induced changes in *Drosophila* behavior and germ line physiology. *eLife*, *4*,
p.e07423.
Klepsatel, P., Knoblochová, D., Girish, T.N., Dircksen, H. and Gáliková, M., 2020. The
influence of developmental diet on reproduction and metabolism in *Drosophila*. *BMC*
Evolutionary Biology, *20*, pp.1-15.
Khodaei, L. and Long, T.A.F., 2019. Kin recognition and co-operative foraging in *Drosophila*
melanogaster larvae. *Journal of Evolutionary Biology*, 32(12), pp.1352-1361.
Khodaei, L., Newman, T., Lum, S., Ngo, H., Maoloni, M. and Long, T.A.F. 2020. On the
expression of co-operative feeding behaviour in 3rd instar *Drosophila melanogaster* larvae!
bioRxiv: p.678631.
Khodaei, L. and Long, T.A.F, 2020. Kin recognition and egg cannibalism by *Drosophila*
melanogaster larvae. *Journal of Insect Behavior*, *33*(1), pp.20-29.
Leadbeater, E., 2009. Social learning: what do *Drosophila* have to offer? *Current Biology*, *19*(9),
pp.R378-R380.
Lefranc, A. and Bundgaard, J., 2000. The influence of male and female body size on copulation
duration and fecundity in *Drosophila melanogaster*. *Hereditas*, 132(3), pp.243-247.
Lenth, R.V., 2022. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package
version 1.7.2. https://CRAN.R-project.org/package=emmeans
LePage, S., Sepil, I., Flintham, E., Pizzari, T., Carazo, P. and Wigby, S., 2017. Male relatedness
and familiarity are required to modulate male-induced harm to females in *Drosophila*.
Proceedings of the Royal Society B: Biological Sciences, *284*(1860), p.20170441.
Lewis, Z., Heys, C., Prescott, M. and Lizé, A., 2014. You are what you eat: gut microbiota
determines kin recognition in *Drosophila*. *Gut microbes*, *5*(4), pp.541-543.
Lihoreau, M., Poissonnier, L.A., Isabel, G. and Dussutour, A., 2016. *Drosophila* females trade off
good nutrition with high-quality oviposition sites when choosing foods. *Journal of*
Experimental Biology, *219*(16), pp.2514-2524.
Lin, C.C., Prokop-Prigge, K.A., Preti, G. and Potter, C.J., 2015. Food odors trigger *Drosophila*
males to deposit a pheromone that guides aggregation and female oviposition decisions.
eLife, *4*, p.e08688.
Lizé, A., McKay, R. and Lewis, Z., 2014. Kin recognition in *Drosophila*: the importance of
ecology and gut microbiota. *The ISME Journal*, *8*(2), pp.469-477.
Long, T.A., Pischedda, A., Stewart, A.D. and Rice, W.R., 2009. A cost of sexual attractiveness to
high-fitness females. *PLoS Biology*, 7(12), p.e1000254.
Malek, H.L. and Long, T.A.F. 2020. On the use of private versus social information in
oviposition site choice decisions by *Drosophila melanogaster* females. *Behavioural Ecology*.
31(3):739-749.
Markow, T.A., 2015. The secret lives of *Drosophila* flies. *eLife*, *4*, p.e06793.
Martin, E.S. and Long, T.A.F., 2015. Are flies kind to kin? The role of intra-and inter-sexual
relatedness in mediating reproductive conflict. *Proceedings of the Royal Society B: Biological*
Sciences, *282*(1821), p.20151991.
Mery, F., Varela, S.A., Danchin, É., Blanchet, S., Parejo, D., Coolen, I. and Wagner, R.H.,
2009. Public versus personal information for mate copying in an invertebrate. *Current*
Biology, *19*(9), pp.730-734.
Monier, M., Nöbel, S., Isabel, G. and Danchin, E., 2018. Effects of a sex ratio gradient on female
mate-copying and choosiness in *Drosophila melanogaster*. *Current Zoology*, *64*(2), pp.251-258.

Moreira-Soto, R.D., Khallaf, M.A., Hansson, B.S. and Knaden, M., 2024. How conspecific and
allospecific eggs and larvae drive oviposition preference in *Drosophila*. *Chemical Senses*, *49*,
p.bjae012.
Najarro, M.A., Sumethasorn, M., Lamoureux, A. and Turner, T.L., 2015. Choosing mates
based on the diet of your ancestors: replication of non-genetic assortative mating in
Drosophila melanogaster. *PeerJ*, *3*, p.e1173.
Nöbel, S., Danchin, E. and Isabel, G., 2018. Mate-copying for a costly variant in *Drosophila*
melanogaster females. *Behavioral Ecology*, *29*(5), pp.1150-1156.
Nöbel, S., Monier, M., Fargeot, L., Lespagnol, G., Danchin, E. and Isabel, G., 2022. Female
fruit flies copy the acceptance, but not the rejection, of a mate. *Behavioral Ecology*, *33*(5),
pp.1018-1024.
Pavković-Lučić, S., Trajković, J., Miličić, D., Anđelković, B., Lučić, L., Savić, T. and Vujisić, L.,
2022. "Scent of a fruit fly": Cuticular chemoprofiles after mating in differently fed
Drosophila melanogaster (Diptera: Drosophilidae) strains. *Archives of Insect Biochemistry and*
Physiology, *109*(3), p.e21866.
Partridge, L., Ewing, A. and Chandler, A., 1987. Male size and mating success in *Drosophila*
melanogaster: the roles of male and female behaviour. *Animal Behaviour*, *35*(2), pp.555-562.
Prasad, N.G., Shakarad, M., Rajamani, M. and Joshi, A., 2003. Interaction between the effects
of maternal and larval levels of nutrition on pre-adult survival in *Drosophila melanogaster*.
Evolutionary Ecology Research, *5*(6), pp.903-911.
R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.
Reaume, C.J. and Sokolowski, M.B., 2006. The nature of *Drosophila melanogaster*. *Current Biology*,
16(16), pp.R623-R628.
Reynolds, J.D. and Gross, M.R., 1990. Costs and benefits of female mate choice: is there a lek
paradox?. *The American Naturalist*, *136*(2), pp.230-243.
Richard, F.J. and Hunt, J.H., 2013. Intracolony chemical communication in social insects.
Insectes sociaux, *60*, pp.275-291.
Robinson, S.P., Kennington, W.J. and Simmons, L.W., 2012. Assortative mating for relatedness
in a large naturally occurring population of *Drosophila melanogaster*. *Journal of Evolutionary*
Biology, *25*(4), pp.716-725.
Rose, M.R., 1984. Laboratory evolution of postponed senescence in *Drosophila*
melanogaster. *Evolution*, *38*(5), pp.1004-1010.
Sarin, S. and Dukas, R., 2009. Social learning about egg-laying substrates in fruitflies. *Proceedings*
of the Royal Society B: Biological Sciences, *276*(1677), pp.4323-4328.
Scott, D., Shields, A., Straker, M., Dalrymple, H., Dhillon, P.K. and Harbinder, S., 2011.
Variation in the male pheromones and mating success of wild caught *Drosophila melanogaster*.
PLoS one, *6*(8), p.e23645.
Sharon, G., Segal, D., Ringo, J.M., Hefetz, A., Zilber-Rosenberg, I. and Rosenberg, E., 2010.
Commensal bacteria play a role in mating preference of *Drosophila melanogaster*. *Proceedings of*
the National Academy of Sciences, *107*(46), pp.20051-20056.
Shoot, T.T., Miller, N.Y. and Long, T.A.F, 2024. Plasticity in expression of fruit fly larval
feeding clusters in response to changes in food quality and distribution. *Insectes Sociaux*, *71*,
pp.1-11.
Singer, T.L., 1998. Roles of hydrocarbons in the recognition systems of insects. *American Zoologist*,
38(2), pp.394-405.
Sullivan, W., Ashburner, M., & Hawley, R. 2000. *Drosophila* protocols. Cold Spring Harbor, NY:
Cold Spring Harbor Laboratory Press.

Toshima, N., Hara, C., Scholz, C.J. and Tanimura, T., 2014. Genetic variation in food choice
behaviour of amino acid-deprived *Drosophila*. *Journal of insect physiology*, *69*, pp.89-94.
Thorpe, W.H., 1945. The evolutionary significance of habitat selection. *The Journal of Animal*
Ecology, *14*(2), pp.67-70.
Valone, T.J. and Templeton, J.J., 2002. Public information for the assessment of quality: a
widespread social phenomenon. *Philosophical Transactions of the Royal Society of London. Series B:*
Biological Sciences, *357*(1427), pp.1549-1557.
Venu, I., Durisko, Z., Xu, J. and Dukas, R., 2014. Social attraction mediated by fruit flies'
microbiome. *Journal of Experimental Biology*, *217*(8), pp.1346-1352.
Vijendravarma, R.K., Narasimha, S. and Kawecki, T.J., 2013. Predatory cannibalism in
Drosophila melanogaster larvae. *Nature Communications*, *4*(1), p.1789.
Wang, Y. and Staubach, F., 2018. Individual variation of natural *D. melanogaster*-associated
bacterial communities. *FEMS Microbiology Letters*, *365*(6), p.fny017.
Weiner, J., 2000. *Time, love, memory: A great biologist and his quest for the origins of behavior*. Vintage.
Wertheim, B., Dicke, M. and Vet, L.E., 2002a. Behavioural plasticity in support of a benefit for
aggregation pheromone use in *Drosophila melanogaster*. *Entomologia Experimentalis et Applicata*,
103(1), pp.61-71.
Wertheim, B., Marchais, J., Vet, L.E.M. and Dicke, M., 2002. Allee effect in larval resource
exploitation in *Drosophila*: an interaction among density of adults, larvae, and micro-
organisms. *Ecological Entomology*, *27*(5), pp.608-617.
Yang, D., 2018. Carnivory in the larvae of *Drosophila melanogaster* and other *Drosophila*
species. *Scientific Reports*, *8*(1), p.15484.
Young, Y., Buckiewicz, N. and Long, T.A.F., 2018. Nutritional geometry and fitness
consequences in *Drosophila suzukii*, the Spotted-Wing *Drosophila*. *Ecology and Evolution*, *8*(5),
pp.2842-2851.
796

Acknowledgements

We would like to thank Simran Mann, Lukas Ghiglione, and Harleen Taneja of the #DrosLife

Lab for their fly-pushing, behavioral observations and camaraderie. Natasha B. Gallo and two

anonymous reviewers provided helpful feed-back and constructive comments. Ruby Lindsay and

Michael Steeleworthy of the Wilfrid Laurier University Library are thanked for their help with

data archiving. This work was conducted at Wilfrid Laurier University, which exists on the

traditional territory of the Neutral, Anishnawbe, and Haudenosaunee peoples.

FIGURES

Figure 1. More focal female offspring produced on dishes where the prior occupant was a related female, while more conspecific offspring were produced on dishes where the female was unrelated. Boxplots illustrating data collected in from the 36 replicate arenas in the first experiment of a) the total number of wild-type (*red-eyed*) adult flies that eclosed from eggs laid by the focal *D. melanogaster* females on 5 different media dishes, b) the total number of *brown-eyed* adult flies that eclosed from those same 5 media dishes, which had been laid by the prior conspecific occupants on the dish. The boxes enclose the middle 50% of data (the inter-quartile range, IQR), with the thick horizontal line representing the location of median. Data points $> \pm 1.5 *$IQR are designated as outliers. Whiskers extend to largest/smallest values that are not outliers, indicated as closed circles. The results of Tukey HSD post-hoc tests comparing group mean are indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level. The control/unexposed dishes by definition had not had any prior exposure to a female capable of producing *brown-eyed* offspring, and thus those dishes never yielded any offspring with that phenotype.

Figure 2. No evidence that focal female offspring production on dishes is inhibited by increased abundance of conspecific offspring. Boxplots illustrating the data collected from the 48 replicate arenas in the second experiment of a) the number of *brown-eyed* adult flies that eclosed from dishes that had been exposed to 0, 1 or 2 conspecific females, b) the number of *wild-type* (red-eyed) adult flies that eclosed from the same dishes that had been laid by the focal female *D. melanogaster*. Boxplot components as in Figure 1. The results of Tukey HSD post-hoc tests comparing group mean are indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level. The control/unexposed dishes by definition had not had any prior exposure to a female capable of producing *brown-eyed* offspring, and thus those dishes never yielded any offspring with that phenotype.

Figure 3. Focal female offspring production is higher on dishes where the unrelated demonstrator originates from the same (IV) population than a different (IV-bw) population. Boxplots illustrating data collected from the 72 replicate arenas in third experiment of a) the number of *brown-eyed* adult flies that eclosed from dishes that had been exposed to either an unrelated IV female (mated to a IV-*bwD* male), a unrelated IV-*bw* female (mated to a IV-*bwD* male), or neither b) the number of *wild-type* (red-eyed) adult flies that eclosed from the same dishes that had been laid by the focal female *D. melanogaster*. Boxplot components as in Figure 1. The control/unexposed dishes by definition had not had any prior exposure to a female capable of producing *brown-eyed* offspring, and thus those dishes never yielded any offspring with that phenotype. The results of Tukey HSD post-hoc tests comparing group mean are indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level.

Figure 4. More focal female offspring produced on dishes where the prior (IV-originating) occupant was a related female. Boxplot illustrating data collected from the 84 replicate arenas in the fourth experiment of the number of *wild-type* (red-eyed) adult flies that had been laid by the focal female *D. melanogaster* that eclosed from dishes that had been exposed to either a related IV female (mated to a IV-*bwD* male), an unrelated IV female (mated to a IV-*bwD* male), or neither. Boxplot components as in Figure 1. The results of a Tukey HSD post-hoc test comparing group mean is indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level.

819
820

822   **SUPPORTING INFORMATION**

Figure S1. Drawings illustrating **a)** the "mini-egg" chambers used to collect eggs and/or create media that exhibited social cues. **b)** an observation arena into which the focal ovipositing female was introduced and monitored. At the bottom of the chamber the lids that exhibited different kinds of social information were affixed. **c)** the arrangement of dishes in the first experiment. In this experiment there were four lids that exhibited social information and the 'control/unexposed' dish that contained no social information (as it had no prior exposure to flies). **d)** the arrangement of the two dishes with social information and the control dish in the second experiment. **e)** the arrangement of the two dishes with social information and the control dish in the third experiment. **f)** the arrangement of the two dishes with social information and the control dish in the fourth experiment.

Figure S2. Focal females visited demonstrator-exposed dishes more frequently than to control/unexposed dishes. Boxplots illustrating the cumulative number of observations made across 26 sessions in the first experiment in which a focal female *D. melanogaster* was observed on the surface of one of 5 different media dishes present in the 36 replicate arenas. The boxes enclose the middle 50% of data (the inter-quartile range, IQR), with the thick horizontal line representing the location of median. Data points $> \pm1.5*$IQR are designated as outliers. Whiskers extend to largest/smallest values that are not outliers, indicated as closed circles. The results of a Tukey HSD post-hoc test comparing group mean is indicated by letters, where groups that do not share the same letter are considered statistically different at the $\alpha=0.05$ level.

Figure S3. Focal females visited demonstrator-exposed dishes more frequently than to control/unexposed dishes. Boxplots illustrating the cumulative number of observations made across 26 sessions in the third experiment in which a focal female *D. melanogaster* was observed on the surface of one of 3 different media dishes present in the 72 replicate arenas that had previously been exposed to either an unrelated IV female (mated to a IV-*bwD* male), an unrelated IV-*bw* female (mated to a IV-*bwD* male), or neither. Boxplot components as in Figure S2. The results of a Tukey HSD post-hoc test comparing group mean is indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level.

Figure S4. Focal females visited demonstrator-exposed dishes more frequently than to control/unexposed dishes. Boxplots illustrating the cumulative number of observations made across 26 sessions in the fourth experiment in which a focal female *D. melanogaster* was observed on the surface of one of 3 different media dishes present in the 84 replicate arenas that had previously been exposed to either a related IV female (mated to a IV-*bwD* male), an unrelated IV female (mated to a IV-*bwD* male), or neither. Boxplot components as in Figure S2. The results of a Tukey HSD post-hoc test comparing group mean is indicated by letters, where groups that do not share the same letter are considered statistically different at the α=0.05 level.

Figure S5. Number of conspecific offspring collected from demonstrator-exposed dishes. Boxplots illustrating the number of *brown-eyed* adult flies that eclosed from dishes that had been exposed to either a related IV female (mated to a IV-*bwD* male), an unrelated IV female (mated to a IV-*bwD* male), or neither in the fourth experiment. There was a small (Cliff's Delta effect size=0.15) but statistically significant difference in the mean number of offspring collected from these dishes. Boxplot components as in Figure S2. The control/unexposed dishes my definition had not had any prior exposure to a female capable of producing *brown-eyed* offspring, and thus those dishes never yielded any offspring with that phenotype.

Figure S6. Schematic diagram depicting hypothetical distribution of female *D. melanogaster* eclosion times in a mixture of eggs collected at the same time from the IV and the IV-*bw* populations (where we assume are randomly sampled, and that both populations have identical distributions, means and variances). While both subset means are the same, as there is a greater number of IV-*bw* eggs in the vial than IV eggs, the larger sample size of the former will likely contain individuals exhibiting a wider range of phenotypes than the latter. The unanticipated consequence of this being that when sampling for virgin female flies, there is a greater probability of colleting a suitable IV-*bw* before encountering a suitable IV female. As body size is negatively correlated with eclosion time (Partridge et al. 1987), and positively correlated with female fecundity (Lefranc and Bundgaard 2000, Long et al. 2009), it is conceivable that the IV-*bw* and the IV females collected and used as demonstrators differed in their capacity to produce offspring, resulting in the conspecific patterns seen in Figure 1b and 3b. It is worth noting that in our fourth experiment, where only *red-eyed* females were collected from vials containing both IV and IV-*bw* eggs that there was only a small difference in the number of demonstrator offspring that were collected (Figure S5), which is consistent with the hypothesis described above.

831