Extensive haplotype diversity in a butterfly colour pattern supergene is fuelled by
incomplete recombination suppression

Rishi De-Kayne[1]*, Ian J. Gordon[2], Reinier F. Terblanche[3], Steve Collins[4], Kennedy Saitoti
Omufwoko[5], Dino J. Martins[6,7], Simon H. Martin[8]*

1. Department of Ecology and Evolutionary Biology, University of California Santa Cruz, 95060,
USA
2. Centre of Excellence in Biodiversity and Natural Resource Management, University of
Rwanda, Huye, Rwanda
3. Department of Conservation Ecology & Entomology, University of Stellenbosch, Stellen-
bosch, South Africa
4. African Butterfly Research Institute, Nairobi, Kenya
5. Department of Ecology and Evolutionary Biology, Princeton University, Princeton, USA
6. Turkana Basin Institute, Stony Brook University, Stony Brook NY 11794 USA
7. Mpala Research Centre, Nanyuki, Kenya.
8. Institute of Ecology and Evolution, University of Edinburgh, Edinburgh EH9 3FL, UK

* Corresponding Authors

Abstract:

Supergenes can evolve when recombination-suppressing mechanisms like inversions promote co-inheritance of alleles at two or more polymorphic loci that affect a complex trait. Theory shows that such genetic architectures can be favoured under balancing selection or local adaptation in the face of gene flow, but they can also bring costs associated with reduced opportunities for recombination. These costs may in turn be offset by rare 'gene flux' between inverted and ancestral haplotypes, with a range of possible outcomes. We aimed to shed light on these processes by investigating the BC supergene, which underlies three distinct wing colour morphs in *Danaus chrysippus*, a butterfly known as the African monarch, African queen and plain tiger. Using whole-genome resequencing data from 174 individuals, we first confirm the effects of BC on wing colour pattern: background coloration is associated with SNPs in the promoter region of *yellow*, within an inversionted part of the supergene, while forewing tip pattern is most likely associated with a copy-number-variable part of the same supergene. We then show that haplotype diversity within the supergene is surprisingly extensive: there are at least six divergent haplotype groups that experience suppressed recombination with respect to each other. Despite high divergence between these haplotype groups, we identify an unexpectedly large number of natural recombinant haplotypes. These evidently arose through crossovers between adjacent inversion 'modules' as well as through double crossovers within inversions. Furthermore, we show that at least one of the established haplotype groups probably arose through recombination between two pre-existing ones. Moreover, on at least two occasions, double crossovers within an inversion have led to the transfer of alleles for dark colouration in the promoter of *yellow* onto a different haplotype background. Overall, our findings paint a picture of dynamic evolution of supergene haplotypes, fuelled by incomplete recombination suppression.

Introduction

Genomic architectures that promote co-inheritance of certain combinations of alleles at multiple loci can be beneficial if they maintain adaptive allele combinations in the face of genetic mixing [1]. Such genetic architectures are referred to as supergenes, because they facilitate the inheritance of complex phenotypes in a simple Mendelian fashion, (reviewed in [2]). Supergenes are often associated with chromosomal inversions, which have the unique feature of suppressing recombination between haplotypes with distinct inversion orientations while allowing free recombination between haplotypes with the same orientation. This effectively divides the population into two subpopulations over a defined portion of the genome [3]. Inversion supergenes have been found to underpin trait polymorphisms under balancing selection, such as alternative life-history and reproductive strategies [4–8] and mimetic coloration [9]. Inversions can also be favoured during local adaptation in the face of gene flow between two distinct environments if they maintain locally adapted combinations that differentiate ecotypes [10–12]. Here we include these cases of locally adapted inversions under the umbrella term 'supergenes', provided they still act to maintain allele combinations that would otherwise be broken down through gene flow and recombination. Genomic studies of species that exhibit local adaptation frequently uncover inversions underpinning differences between ecotypes [13–16], and 'bottom-up' analyses have identified signatures consistent with locally adapted inversions even when the specific traits they may be associated with are unknown [17–19]. This suggests that supergene architectures may be ubiquitous in nature, highlighting a need to better understand their evolutionary dynamics.

Recent theoretical and empirical work suggests that supergenes may evolve over time in their structure, composition, and effects on phenotype and/or fitness. First, supergene structure can evolve through recurrent chromosomal rearrangements. Studies across multiple systems show that supergenes often comprise several rearrangements, implying stepwise growth in the region of suppressed recombination (e.g. in *Heliconius* butterflies [20], *Danaus* butterflies [21], and *Solenopsis* fire ants [6]). In addition to allowing the incorporation of additional co-adapted alleles at other loci, subsequent rearrangements of the supergene region can cause recombination suppression between more than two distinct haplotypes [21]. Second, even without physical expansion, the repertoire of traits affected by an inversion supergene may expand over time as additional alleles become established at other loci within the region of recombination suppression [11,17]. The fitness consequences of a supergene could also change if selective pressures change over time or across space, potentially limiting the value of a supergene in a changing environment or during dispersal into a new environment [22]. Even in a stable environment, inversion supergenes may be subject to accumulation of increased mutational load compared to the rest of the genome due to their reduced opportunities for recombination and reduced effective population size (N_e) (due to the effective subdivision of the population in that part of the genome) [23–25]. Some theoretical models

suggest that this may lead to heterokaryotype advantage through sheltering of recess-
ive deleterious mutations (associative overdominance) [23–25], or failure of locally ad-
apted inversions to reach high frequency [26].

Another process that contributes to the evolution of inversion supergenes is rare recom-
bination in heterokaryotypes, which results in some degree of gene flux between haplo-
types. While single crossovers within inversions are usually strongly suppressed due to
the production of unbalanced chromosomes [27], large inversions may allow for double-
crossovers, which result in the exchange of genetic material between arrangements, ef-
fectively creating a mosaic haplotype. In addition, gene flux of short fragments can oc-
cur through non-crossover gene conversion, which has been shown experimentally in
Drosophila to occur at least as often within inversion heterokaryotypes as in syntenic re-
gions [28–30]. Population genomic analyses in a range of different species support the
existence of considerable gene flux within inversions [31–34], and suggest that a drift-
flux equilibrium can be reached [3]. Gene flux provides a means by which supergenes
may avoid some of the costs of recombination suppression described above, potentially
facilitating their long-term persistence [6,23,34]. In some supergenes, sequence diver-
gence suggests persistence of polymorphism for millions of years, even through mul-
tiple speciation events [21,34]. However, the role of gene flux in supergene temporal dy-
namics remains under-explored. A compelling example of recombination creating a third
distinct haplotype is seen in the supergene that controls reproductive morphology and
behaviour in the ruff [5,34]. On the other hand, a mimicry supergene in *Papilio* butter-
flies shows phylogenetic relationships consistent with wholesale 'allelic turnover', in
which new haplotypes arise (possibly via recombination) and replace ancestral ones
[35]. Case studies such as these provide valuable insights into the range of processes
contributing to supergene evolution.

Danaus chrysippus, a butterfly known as the African monarch, African queen, and plain
tiger, presents an opportunity to investigate the contributions of the above processes to
the evolution of a large, ancient supergene. Like other milkweed butterflies, *D. chrysip-*
pus has bright warning patterns that advertise its toxicity. Across its range, it is divided
into several parapatric morphs with distinct warning patterns that meet in a broad hybrid
zone in eastern central Africa (Fig. 1A) [36,37]. Phenotypic differences in two forewing
traits are controlled by the 'BC supergene' on chromosome 15, which links at least two
colour patterning loci, the 'B' locus which affects whether background colour is dark
(dominant *B* allele) or pale (recessive *b* allele); and the 'C' locus which determines
whether the apical black tip and white band is present (recessive *c* allele), or absent
(dominant *C* allele) [21,38]. Previous genomic comparisons of a limited number of popu-
lations revealed the existence of three divergent haplotype groups (also called 'alleles')
at the BC supergene, which were named according to the morphs in which they are
found: 'orientis' (*Bc*, Southern Africa), 'klugii' (*bC*, East Africa) and 'chrysippus' (*bc*,

West Africa, North Africa, Mediterranean, Asia) [38]. Note that while a distinct morph 'al-
cippus' is found in West Africa, it only differs in its hindwing phenotype, controlled by a
locus on a different chromosome, but shares its forewing phenotype and corresponding
BC supergene haplotype with the chrysippus morph. Chromosome-scale assemblies of
each of the three haplotypes revealed that, rather than comprising a single large inver-
sion, the BC supergene is underpinned by a complex, modular rearrangement including
several inversions and a large copy-number variable (CNV) region [21]. This complex
architecture helps to explain how recombination is suppressed between more than two
divergent haplotype groups. Phylogenetic analysis showed that the rearrangements
probably occurred in a stepwise manner, beginning several million years ago, and that
the supergene has persisted through multiple speciation events [21]. Intriguingly, an ad-
ditional rearrangement has recently occurred and risen to high frequency in a small re-
gion of East Africa: a fusion of chr15 to the female-specific W chromosome [38,39]. Be-
cause crossing-over is limited to males in Lepidoptera, this fusion represents an addi-
tional mechanism of recombination suppression, effectively creating a new female-spe-
cific sub-group of the chrysippus haplotype [38]. There is also evidence for rare recom-
bination within the supergene based on phenotypes in crossing experiments, and a pu-
tative recombinant haplotype assembly from a hybrid-zone individual [21,40]. Taken to-
gether, the above findings suggest that, despite its age, the BC supergene continues to
evolve, raising questions about the full extent of diversity at this locus and the role of re-
combination in shaping this diversity.

Here we investigate haplotype diversity and evolution of the BC supergene using gen-
omic data from 174 *D. chrysippus* individuals representing 14 regions across Africa and
Southern Europe. We first confirmed that the supergene links loci controlling two distinct
wing colour pattern traits. We then describe haplotype diversity and identify additional
divergent haplotype groups beyond those previously described. Comparable levels of
genetic diversity in each haplotype group and across most of the discrete structural
modules of the supergene argues against ongoing allelic turnover, and instead suggest
a long-term polymorphism, probably driven by local adaptation. Perhaps most in-
triguingly, we find abundant evidence of recombination and gene flux between haplo-
type groups, with crossovers having occurred both between adjacent inversions and
within individual inversions, leading to exchange of functional colour pattern alleles.
These results support a role for somewhat rare recombination in the diversification and
long-term persistence of supergene haplotypes.

Results:

Widespread sampling confirms that the BC supergene is associated with forewing colour variation and is the main axis of genetic variation

We analysed resequenced genomes of 174 individual butterflies spanning the three core *D. chrysippus* forewing morphs, 'chrysippus' (note that for our purposes, this also includes form 'alcippus', which differs only in its hindwing), 'orientis', and 'klugii', as well as intermediate morphs (Fig. 1A; S2 Table). Our sampling includes areas of monomorphism for each morph: western Africa, South Africa and southeastern Kenya, respectively. We also sampled the hybrid zone (sampled in Rwanda and central Kenya, where all three forewing morphs and intermediates are found), and North Africa and the Mediterranean (where two of the three morphs, and intermediates are found). Twelve sequenced individuals were bred in captivity from parents of mixed or unknown origins and were not assigned to any geographic group. Pairwise F_{ST} between three monomorphic locations in South Africa (TSW population, orientis morph), Nigeria (NGA population, chrysippus morph) and Eastern Kenya (WAT population, klugii morph), is close to zero genome-wide, with the exception of a few large peaks, including the largest on chromosome 15 (chr15), as described previously (Fig. 1B; S2 Table)[38]. Genome-wide principal components analysis (PCA) for all autosomes excluding chr15 further supports minimal genetic structure outside of the BC supergene, with the exception that samples from the small island of St. Helena and north of the Saraha cluster separately from the rest of sub-saharan Africa (Figure A in S1 Text).

Figure 1. **Broad geographic distribution of *Danaus chrysippus* morphs and our continent-wide sampling of individuals spans wing pattern variation that is under-pinned largely by variation on chromosome 15. A.** The geographic distribution of the three main *D. chrysippus* forewing morphs; chrysippus, orientis, and klugii as well as putative heterozygotes (data from manual curation of GBIF record and other scientific collections, following the approach in [37]; left) and sampled locations of *D. chrysippus* individuals used for genomic analysis in the present study (right). Morphs and their corresponding inferred genotypes at the B and C loci are shown below. Note that for our purposes the forewing morph 'chrysippus' includes the west-African morph 'alcippus', which has the same forewing phenotype but differs in its hindwing phenotype. **B.** Genome-wide patterns of F_{ST}

between morphs (smoothed across discrete 50kb windows) highlighting significant differentiation along
the length of the chr15 supergene region. **C-D.** Genome-wide associations between SNP variation and **C.**
background colouration and **D.** forewing band presence/absence (both N=172) reflecting significant asso-
ciations between this region and wing-pattern variation. Significant associations, as identified via permuta-
tion tests (p < 0.05), are represented by black points.

Genome-wide association (GWAS) analysis confirms that the BC supergene region on
chr15 is associated with two forewing traits: the pale/dark background wing colour (also
known as the B locus; Fig. 1C), and the presence/absence of the apical black tip and
white band (also known as the C locus; Fig. 1D). In addition to a large peak of associ-
ation in the supergene, there are several peaks of association with pale/dark variation
on other chromosomes, suggesting that other loci also contribute to this trait. A repeat
of the GWAS using only samples from the hybrid zone (NYA, NRB, MPL populations,
N=87) recapitulates the same general result (Figure B in S1 Text), confirming that the
observed associations are not driven by correlated selection on other traits that follow a
similar geographic distribution. The cluster of SNPs most strongly associated with back-
ground wing colour fall between two genes: *yellow*, which is known to be involved in
melanin synthesis and coloration in other insects [41], and the *achaete-scute complex*
protein T3-like, which has been shown to be involved in wing scale development across
Lepidoptera [42]. The SNPs most strongly associated with the presence or absence of
the forewing black tip were found to fall within the copy-number-variable (CNV) region of
the supergene (Figure C in S1 Text). These findings suggest that the B and C loci are
distinct, and that the supergene is favoured because it maintains locally adapted com-
binations of alleles at these loci, and probably others among the ~150 genes it encom-
passes, in the face of gene flow.

Extensive genetic variation at the BC supergene
We previously described three divergent haplotype groups (sometimes called 'alleles')
of the BC supergene, corresponding to the three forewing morphs, maintained by sev-
eral inversions, intra-chromosomal translocations, and copy-number differences that
suppress recombination [21,38]. We therefore expected that diploid genotypes across
the BC supergene would form six distinct genetic clusters corresponding to the three
homozygous and three heterozygous states. However, as we describe below, several
lines of evidence suggest that there are additional divergent haplotypes, and possible
recombinants. First, both a distance-based neighborNet network and principal compon-
ents analysis for the complete supergene (excluding the CNV region) show more com-
plex genetic structuring than we expected (Fig. 2A, Figure A in S1 Text). Three clusters
representing homozygous genotypes for the three previously-described haplotype
groups are identifiable (Fig. 2A), and include individuals previously matched to these
genotypes [21]. One large cluster of individuals is intermediate between the known
chrysippus and klugii homozygotes (Fig. 2A) and includes known heterozygotes

between chrysippus and klugii from crosses and suspected wild heterozygotes based on wing patterns. However, numerous other individuals are dispersed across the network at varying distances between the three homozygous clusters. We therefore hypothesised that in addition to heterozygotes among the three known haplotype groups, our sampling may include additional divergent haplotype groups and/or recombinants among the three common haplotypes.

Figure 2. Genetic clustering and ancestry painting suggest additional divergent haplotype groups, and recombinants. A. NeighborNet network for unphased diploid genotypes across the BC supergene (excluding the CNV region). Each tip represents one diploid individual, and the network is constructed based on average pairwise genetic distances considering both haplotypes in each individual. We therefore expect 'heterozygous' individuals carrying two distinct haplotypes to be at intermediate positions in the network. The single individual represented by a black dot is the *Danaus melanippus* outgroup. **B.** Map of localities for sequenced individuals. Pie charts in panels A and B represent inferred ancestry components for each diploid individual from Admixture analysis with k=4 source populations (See Figures D-H in S1 Text for plots with other values of k). Note that for highly sampled localities, an arbitrary subset of sequenced individuals is shown in panel B. **C.** Ancestry painting across the central

portion of chr15 including the BC supergene for 27 representative individuals, including homozygotes,
heterozygotes and putative recombinants. The 27 selected individuals are indicated in panel A. See Fig-
ures I and J in S1 Text for ancestry painting for all individuals. The CNV region is excluded from the plot
for convenience as it cannot be reliably genotyped (represented as a white gap).

Given the complex relationships observed, we attempted to naively identify distinct hap-
lotype groups using Admixture [43] analysis, in which each individual is modelled in
terms of its ancestry components from k source populations. Because recombination
should occur freely among haplotypes with the same structural arrangement, and
should only be suppressed among those with different arrangements, the supergene re-
gion should comprise a set of semi-isolated sub-populations. We predicted that "hetero-
zygous" individuals (i.e. carrying two divergent haplotypes) will appear as admixed, with
approximately 50% ancestry contributions from two source populations, while homozy-
gotes should be assigned 100% ancestry from a single source. All models with k=3
(matching our *a priori* hypothesis) and above captured the three previously-described
haplotype groups corresponding to the klugii, chrysippus and orientis morphs. These
are represented in clusters of homozygous (100% ancestry) individuals from eastern,
western and southern Africa, respectively, while most hybrid-zone individuals appear
heterozygous, with 50% ancestry proportions (Figures E-H in S1 Text). However, as de-
scribed below, the models with k=4 to k=6, which all have lower cross-validation error
than k=3 (Figure D in S1 Text), provide compelling evidence for the existence of addi-
tional divergent haplotype groups that had not been previously described.

We have chosen to present the ancestry assignments according to the model with k=4
sources in Fig. 2 (see pie charts in Fig. 2A and 2B), because our sampling included ho-
mozygotes for all four putative haplotype groups. The fourth source population is in-
ferred to contribute ~100% ancestry to two South African individuals, and ~50% ances-
try to several others. We therefore infer that a fourth arrangement of the supergene oc-
curs in southeast Africa. This haplotype group is cryptic in the sense that it is associated
with a forewing phenotype indistinguishable from that of orientis. Hereafter, we refer to
this as the 'karamu' haplotype group.

In the model with k=5 sources (Figure G in S1 Text), the fifth cluster captures the previ-
ously described neo-W fusion - a lineage of chr15 haplotypes that arose recently in East
Africa when a copy of chr15 from the chrysippus haplotype group fused to the female-
limited W sex chromosome [38,39]. These females are each assigned ~50% ancestry
from this source, consistent with W being haploid in females. We previously showed
that the neo-W-chr15 fusion occurred recently and spread to high frequency in southern
Kenya over the past 2,200 years [38]. The high sequence similarity and complete lack
of recombination of the neo-W explains how it is identifiable as a distinct genetic cluster
despite its recent separation from the standard chrysippus haplotype group.

In the model with *k*=6 sources (Figure H in S1 Text), several individuals from North
Africa and the Mediterranean are assigned to a sixth distinct cluster, suggesting yet an-
other divergent haplotype group. We hypothesise that, similar to the karamu haplotype
from southeast Africa, another distinct arrangement of chr15 likely occurs north of the
Sahara. However, subsequent analyses (described below) suggest that none of our se-
quenced individuals are homozygous for this haplotype, limiting our ability to further in-
vestigate its status.

Ancestry painting confirms additional haplotype groups, as well as recombinants
To investigate fine-scale haplotype ancestry and recombination across the BC su-
pergene region, we applied two ancestry painting methods. These aim to assign ances-
try for phased-inferred haplotypes based on pre-defined source populations, using
either a hidden markov model [44] or defined windows of SNPs (see Methods for de-
tails). We used populations WAT, TSA and NGA as reference sets representing the klu-
gii, orientis and chrysippus haplotype groups, due to their monomorphic clustering (Fig.
2B). In addition, we used the two individuals from the SPA population showing 100%
karamu ancestry to represent this fourth haplotype group (Fig. 2B). Both ancestry paint-
ing methods agree strongly with our inferences from the Admixture analysis. All four
haplotype groups occur as intact haplotypes in most individuals, with large numbers of
both homozygotes (individuals 1-11 in Fig. 2C) and heterozygotes (individuals 12-18 in
Fig. 2C). See Figures I and J in S1 Text for all individuals. Heterozygotes are particu-
larly common in the hybrid-zone (populations NYA, MPL and NRB in Rwanda and cent-
ral Kenya, Figures I and J in S1 Text).

Ancestry painting also identifies individuals with mosaic ancestry consistent with recom-
bination between the divergent haplotypes. Most of these putative recombinant haplo-
types appear to be simple chimeras that result from a single crossover located between
two adjacent inversions (see for example individuals 19-21 in Fig. 2C). However, there
are also instances of more fine-scale mosaic ancestry consistent with recombination
within inversions (see for example individuals 22-25 in Fig. 2C). While gene conversion
can lead to the transfer of small haplotype tracts within inversions (tract lengths vary
across taxa but estimates suggest they are typically < 4kbp in length; [45]), this process
cannot explain the scale of mosaic ancestry tracts we observe (tens to hundreds of kb).
Instead, these patterns are consistent with occasional double crossovers, which allow
genetic exchange within the inversions without the formation of unbalanced gametes
(see Discussion). Some of these recombinant haplotype patterns appear multiple times
in unrelated individuals, implying that they may be increasing in frequency in the popula-
tion (see for example individuals 23-24 in Fig. 2C). Unsurprisingly, most recombinant
haplotypes are found in the hybrid zone (Figures I and J in S1 Text). However, we were
surprised to find two distinct recombinant haplotypes present in the genomes from the

small island of St. Helena (STH population), where only one of the six sequenced indi-
viduals did not carry a recombinant haplotype (Figures I and J in S1 Text).

Ancestry painting failed to assign ancestry to one of the haplotypes in some individuals
from North Africa and the Mediterranean (see for example individuals 26-27 in Fig. 2C).
These are individuals that were assigned to the 6th cluster in the admixture analysis
with k=6 sources (Figure H in S1 Text). This supports the existence of an additional di-
vergent haplotype of the BC supergene that occurs north of the Sahara. Other members
of this population carry a chrysippus-like haplotype, with some evidence for recombina-
tion with orientis (seen in individuals 25-27 in Fig. 2C). Because no individual is homo-
zygous for the additional divergent haplotype, we were unable to include it as a source
for ancestry painting, and it was not included in our subsequent analyses.

In summary, our findings suggest that there are (at least) six divergent haplotype broups
of the BC supergene that experience suppressed recombination with respect to each
other. Three were previously assembled and found to have distinct arrangements
(chrysippus, klugii and orientis); one results from fusion of chr15 to the W chromosome
(neo-W-chrysippus); and two are newly identified here based on genetic clustering, but
probably also represent distinct structural arrangements (karamu and the unnamed hap-
lotype from North Africa and the Mediterranean) (Table 1).

Table 1. Six divergent haplotype groups of the BC supergene

Haplotype Name	Geographic region(s)	Background colour (B locus genotype)	Forewing tip and band (C locus genotype)	Conventional name of morph	Origin
chrysippus	West Africa, North Africa & Mediterranean, Asia	Pale (*b*/*b*)	Present (*c*/*c*)	chrysippus, alcippus	Ancient
klugii	East Africa	Pale (*b*/*b*)	Absent (*C*/-)	klugii	Ancient
orientis	Southern Africa	Dark (*B*/-)	Present (c/c)	orientis	Ancient
karamu	Southeast Africa	Dark (*B*/-)	Present (c/c)	orientis	Recombination: klugii + orientis
neo-W-chrysippus	East Africa (Hybrid Zone)	Pale (*b*/*b*)	Present (*c*/*c*)	N/A (never homozygous)	W-fusion: chrysippus
Unnamed	North Africa & Mediterranean	Unknown (no homozygotes)	Unknown (no homozygotes)	Unknown (no homozygotes)	Ancient

Diversity and divergence patterns are consistent with long-term polymorphism
One possible explanation for the coexistence of multiple haplotypes associated with the
same wing phenotype in the same population (e.g. karamu and orientis haplotypes in
the SPA population) is that we are witnessing an ongoing 'allelic turnover' event [35], in
which a new haplotype is replacing an older one. If a new haplotype arose through a
novel structural rearrangement, it is expected to have undergone a bottleneck of N=1 at
its conception [3], which would result in dramatically reduced diversity immediately fol-
lowing its origin. Older haplotypes should have had the opportunity to recover diversity
following their initial bottleneck, but are nevertheless still expected to have lower di-
versity than the genomic background level because inversions effectively divide the
species into sub-populations with lower effective population size across the inverted re-
gion (even for the ancestral orientation) [3]. We therefore expected that all haplotype
groups may show reduced within-group diversity within the inverted regions, but we hy-
pothesised that a more pronounced reduction may be observed in those with derived in-
version orientations: orientis and chrysippus [21] and possibly the newly identified
karamu haplotype. Consistent with expectations, diversity is lower within the inverted re-
gions of chr15 compared to collinear regions in all four haplotype groups (Fig. 3b). The

extent of diversity reduction is notably similar in the four groups, with the exception that orientis has lower diversity in inverted region 1.1, which is known to be uniquely inverted in orientis. No haplotype group has diversity approaching zero, as would be expected under rapid, recent allelic turnover, so these patterns are more consistent with a long-term polymorphism at the BC supergene.

Figure 3. **The supergene region has reduced genetic diversity in all haplotype groups and elevated genetic differentiation between them, but no evidence for a very recent origin of any haplotype. A.** A graphical representation of the structural variation across the chrysippus, orientis, and klugii haplotypes described in [21]. The supergene evolved through rearrangements of four main regions (indicated with different colours, regions with an inverted orientation relative to the ancestral state are indicated with *). Regions 1.1, 1.2, 2, and 4 remain in single-copy and can be reliably aligned and genotyped. **B-E.** Population summary statistics computed in non-overlapping 50kb windows for representative populations of chrysippus (NGA population), orientis (TSW), klugii (WAT) and karamu (two homozygous individuals from the SPA), plotted across chr15 (excluding the CNV region). Note that the reference genome is from a klugii haplotype, hence regions 1 and 2 are immedi-

ately adjacent. Statistics plotted are: **B.** Nucleotide diversity (π) **C.** absolute pairwise divergence (d_{XY}), **D.**
net pairwise divergence (d_a), and **E.** pairwise differentiation (F_{ST}).

Genetic divergence and differentiation between haplotype groups are also consistent
with relatively ancient origins. Elevated F_{ST} and net divergence (d_a) are restricted to the
rearranged parts of the chromosome (Fig. 3), consistent with recombination suppres-
sion being restricted to the supergene (note that regions 1.1 and 4 are only inverted in
orientis and are collinear in klugii and chrysippus). Similar levels of differentiation are
seen between these three previously-described haplotype groups across most of the
supergene (excluding region 1.1), which may indicate that the arrangements all oc-
curred around the same time, although this pattern would also be expected under muta-
tion-drift-flux equilibrium [3]. However, divergence and differentiation are both notably
lower between klugii and the newly-described karamu haplotype group across most of
the supergene, consistent with either a more recent divergence between these haplo-
types, or a higher rate of gene flux between them. This is intriguing, because these hap-
lotypes are associated with distinct wing phenotypes. This raises the possibility that
gene flux between haplotype groups has allowed the exchange of functional alleles.

Recombination between supergene haplotypes has exchanged wing colour alleles
We set out to test whether the evolution of the karamu haplotype may have arisen
through recombination between haplotype groups. Specifically, we applied topology
weighting [46] using TWISST2 to ask whether there are tracts in which karamu is more
closely related to orientis than to klugii. This revealed that karamu clusters broadly with
klugii throughout the length of the supergene, but there are multiple narrow tracts at
which it clusters more closely with orientis, particularly in inverted region 2 (Fig. 4A). An-
cestry painting using Loter [44] confirms that one such narrow tract is a 20kb region
containing the gene *yellow* and part of its promoter, including nine of the ten SNPs most
strongly associated with wing background colouration according to our GWAS (Fig. 4B).
This therefore suggests that incorporation of the orientis-like allele at *yellow* (i.e. the *B*
allele at the B locus) through recombination causes the karamu haplotype to produce a
dark wing colour phenotype matching that of orientis, despite otherwise being more klu-
gii-like in its overall ancestry. We hypothesise that a similar allelic exchange at the C
locus causes karamu individuals to share the forewing black tip phenotype of orientis,
but we are unable to test this hypothesis, as this trait maps to the CNV region in which
genotyping and ancestry assignment are unreliable.

Figure 4. Two independent cases of recombination within the BC supergene with phenotypic consequences. A. Topology weightings across chr15 showing how the karamu haplotype is related to the klugii and orientis haplotypes. Upper panel shows three possible rooted genealogical topologies. Second panel shows weights for each topology along the chromosome, smoothed with a 20kb span. Arrows above the plot indicate the locations of inversions. Third panel shows unsmoothed topology weightings across a 1.5 Mb region corresponding to Inversion 2. **B.** Ancestry painting from Loter [44] across a 100 kb region within Inversion 2 showing ancestry tracts for two homozygous karamu individuals

compared to two representative individuals homozygous for the orientis and klugii haplotypes. Coding re-
gions are indicated below the plot, with the candidate gene for background coloration *yellow* indicated.
Green triangles represent the top 10 SNPs for background colour in our GWAS (Fig. 1C). There is evid-
ence for recombination throughout the supergene region, and specifically in the vicinity of *yellow*, consist-
ent with the hypothesis that orientis ancestry at this locus (i.e. the *B* allele) is associated with darker col-
ouration in karamu individuals. **C** and **D.** A second example of recombination in the promoter of *yellow*.
Plots are as described for panels A and B, except showing relationships between two individuals from
North Africa & Mediterranean with a chrysippus-like haplotype (according to Admixture and ancestry
painting), but dark background colouration. Again, orientis ancestry in the promoter region of *yellow* sug-
gests that recombination allowed the transfer of the *B* allele into a different genetic background, causing
darker wing colouration.

We identified a second probable case of recombination affecting background coloura-
tion. Several individuals from North Africa and the Mediterranean are homozygous for
chrysippus-like haplotypes according to chromosome painting (see for example indi-
vidual 25 in Fig. 2C), but have dark background colouration similar to that seen in ori-
entis. We again tested for fine-scale mosaic ancestry in two of these individuals using
topology weighting and ancestry painting. A very similar pattern to the karamu case is
observed: these individuals cluster strongly with chrysippus throughout the supergene,
but carry narrow orientis-like tracts (Fig. 4C), including a 10kb tract in the promoter re-
gion of *yellow* encompassing all ten SNPs most strongly associated with dark back-
ground colouration (Fig. 4D). This represents a distinct event in which orientis-like al-
leles (i.e. the *B* allele at the B locus) have been incorporated into a different genetic
background through recombination within an inversion, leading to an altered phenotype.

Historical gene flux also occurred between the major supergene haplotypes
Based on the extensive evidence for recombination observed, we hypothesised that
even the three originally described haplotype groups (orientis, klugii and chrysippus)
may have been historically shaped by gene flux. To explore this possibility, we first ran
another topology weighting analysis to examine fine-scale relationships between these
three groups. This generally agrees with the supergene structure: in each rearrange-
ment, the predominant genealogy clusters groups according to whether they carry the
rearranged or ancestral arrangement (Figure L in S1 Text). However, there is hetero-
geneity in these relationships, and occasionally, complete switches to a different rela-
tionship within each rearranged region (Figure L in S1 Text). These patterns are con-
sistent with historical gene flux through double-crossover events. However, considering
the high absolute divergence observed between these three haplotype groups (Fig. 3),
there is no compelling evidence for very recent exchange of large haplotype blocks
through double-crossovers.

Finally, we attempted to quantify gene flux between the three major haplotype groups
by fitting isolation-with-migration (IM) models, which are conventionally applied to model

species divergence with gene flow. Here haplotype groups are modelled as panmictic
populations that became isolated at some point in the past, and gene flux between the
distinct haplotype groups is modelled as 'migration' [33]. IM models were fitted using
gIMble [47], with separate models for each rearranged region and each pair of haplo-
type groups, because not all regions are inverted between all pairs. For example, re-
gions 1.1 and 4 are inverted in orientis only, so we expect approximately free recombin-
ation between klugii and chrysippus in these regions. Of the eight pairwise tests in
which the regions are inverted, three had best-fitting models without gene flux, while the
remaining five had estimated effective migration rates (m_e) ranging from 2.42x10^{-11} to
3.09x10^{-6} (this equates to a range of 3.02x10^{-5} to 4.89 effective migrants per generation
(M_e) assuming $M_e = m_e 4N_{e\ (recipient)}$; Fig. 5; Table 2). Higher rates are generally seen
between klugii and orientis, but there is no apparent trend with the size of the re-
arranged region. Of the four tests that involved collinear regions, two had the highest
estimated effective migration rates, and two had low or no inferred gene flux. This is
probably because these latter two collinear tracts (specifically region 1.2 between
chrysippus and orientis, and region 2 between klugii and chrysippus) are nevertheless
physically translocated with respect to each other, probably impairing correct meiotic
pairing. Taking these results together, we conclude that there is considerable evidence
for history of gene flux in at least some of the rearranged regions of the supergene, but
our estimated rates of gene flux should be interpreted with caution given the small size
of the regions analysed and resulting sampling noise and reduced inference power (we
note that one model failed to optimise), and the modelling restriction of unidirectional
gene flux.

Figure 5. glMble analysis highlights gene flow between specific supergene module alleles. A. A graphical representation of structural variation across the chrysippus, orientis, and klugii haplotypes replotted with data from [21] (regions with an inverted orientation relative to the ancestral state are indicated with *). **B.** Coalescent modelling between chrysippus and klugii individuals (top), chrysippus and orientis individuals (middle), and klugii and orientis individuals (bottom) highlighting which of the three model types, divergence only, divergence with gene flow from population A to B, and divergence with gene flow from B to A had the lowest composite likelihood for each region and comparison. Box borders indicate whether regions being compared are collinear (dotted line) or inverted (solid line) between morphs. * indicates the one case where the model failed to optimise.

Table 2. Summary of glMble results highlighting the best fitting models for each region and haplotype pair combination.

Inversion	Test	Model type	Ne population A	Ne population B	Ne ancestral	Divergence time	m_e	Composite likelihood
1.1	chrysippus - klugii	IM BA	4.03E+05	4.48E+05	1.76E+06	2.47E+06	3.09E-06	-439547.3587
1.1	chrysippus - orientis	DIV	4.02E+05	3.10E+05	1.11E+06	9.30E+05	-	-551959.4922
1.1	klugii - orientis	IM BA	5.78E+05	1.34E+05	2.77E+06	4.25E+06	7.20E-07	-605170.8143
1.2	chrysippus - klugii	DIV	1.71E+05	1.95E+05	1.05E+06	7.43E+05	-	-283401.4511
1.2	chrysippus - orientis	DIV	1.99E+05	2.50E+05	1.11E+06	9.62E+05	-	-321748.6401
1.2	klugii - orientis	IM AB	8.00E+04	2.19E+05	1.99E+06	2.27E+06	1.04E-06	-409100.246
2	chrysippus - klugii	IM AB	3.18E+05	2.73E+05	8.76E+05	9.46E+05	3.80E-11	-565003.353
2	chrysippus - orientis	IM AB	3.58E+05	3.12E+05	8.97E+05	9.60E+05	2.42E-11	-596504.91
2	klugii - orientis	DIV	1.97E+05	2.02E+05	8.66E+05	6.67E+05	-	-716365.7526
4	chrysippus - klugii	IM AB	3.94E+05	6.43E+05	1.59E+06	2.42E+06	2.71E-06	-399370.2631
4	chrysippus - orientis	IM AB	7.10E+05	5.84E+05	1.16E+06	7.62E+05	2.49E-11	-492519.5942
4	klugii - orientis	IM BA	7.60E+05	3.14E+05	2.04E+06	3.83E+06	8.79E-07	-493600.4502

Populations tested are listed in the order A-B. DIV = and divergence with no migration; IM AB = isolation with unidirectional migration from A to B; IM BA = from B to A. Divergence times are in generations.

Discussion:

Structural variants that suppress recombination have been found to contribute to local adaptation and the maintenance of complex phenotypic polymorphisms. However, to understand how these variants persist and change over time, it is necessary to go beyond simply associating genotype and phenotype, and to investigate how sequences evolve within these genomic regions. Here we show that the BC supergene of *Danaus chrysippus*, which underpins wing colour pattern variation, displays surprising haplotype diversity, with a greater number of divergent haplotype groups than there are described phenotypic morphs. Counterintuitively, this diversity is partly fuelled by recombination. Our findings suggest that supergene evolution is a highly dynamic process, raising a number of questions that should be considered in future structural variant research.

A paradox of our study is that we report evidence for both historical and recent recombination in a genomic region that was first identified because of its role in suppressing crossing over. In reality, the realised rate of recombination in the BC supergene must be low, otherwise we would not have been able to describe the highly diverged haplotype groups, nor to identify recombinants between them. Indeed, although a considerable fraction of our sequenced individuals carry haplotypes showing evidence for recent recombination, many of the recombination breakpoints are shared among individuals, implying that the observed recombinant haplotypes stem from a smaller number of recombination events. The modular structure of the supergene means that single crossovers between adjacent inversions can generate modular ancestry mosaics. However, the more complex mosaic ancestries of some haplotypes imply double crossovers within inversions. It is probably rare for two crossovers to occur within a single inversion of just a few megabases, due to crossover interference, especially considering that chromosome map lengths of butterflies in the same family tend to average around 50 cM [48,49], implying an average of one crossover per meiosis per bivalent. However, we note that once a double-crossover event has occurred, transferring and effectively un-inverting genetic material between inversion haplotypes, a more complex fine-scale ancestry mosaic can emerge through unrestricted single crossovers in subsequent generations. This process, repeated over time, is expected to lead to more gene flux near the centre of inversions than near the breakpoints, giving rise to a characteristic "suspension bridge" pattern of divergence between inversion haplotypes, as is seen for example in *Drosophila melanogaster* inversion 3R [32]. The lack of such a pattern in any of the *D. chrysippus* BC inversions requires further investigation. In addition to double crossovers, there is probably a considerable contribution to gene flux from non-crossover gene conversion, as has also been described in *Drosophila* [28]. However, as noted, the resulting flux is insufficient to cause homogenisation of allele frequencies as seen in the remainder of the genome, where $F_{ST} \sim 0$. The similar levels of divergence observed between three of the haplotype groups across the inversions suggests that a point of

mutation-drift-flux equilibrium has been reached [3].

Our findings do not provide direct evidence of the potential costs or benefits of recom-
bination within the supergene region. The simplest models of supergene evolution imply
that recombinant haplotypes have reduced fitness due to broken associations between
co-adapted or locally adapted alleles. One piece of indirect evidence in support of this
idea is the observed excess of recombinant haplotypes on the small island population of
St. Helena, which we believe to be recently bottlenecked, and therefore subject to
greater drift and less efficient purifying selection than mainland populations. On the
other hand, it is also likely that certain recombinant haplotypes are occasionally more fit
than the prevailing haplotypes, either because they provide new allelic combinations as-
sociated with a fit phenotype [5,22,34], or because they allow purging of deleterious
mutations [23]. Our finding that several haplotypes with the same complex ancestry mo-
saics appear in multiple individuals may imply a process of allelic turnover, in which an
existing 'allele' (or haplotype group) is in the process of being replaced by a fitter recom-
binant one. Indeed, the complex pattern of relationships among even the established
haplotype groups suggests that recombination and allelic turnover may have occurred
repeatedly throughout the evolutionary history of the BC supergene. As a result, it is
likely that none of the established haplotype groups we describe here closely resemble
the 'original' haplotypes captured when the rearrangements first occurred. A process of
dynamic turnover has been suggested in a different supergene system in *Papilio* butter-
flies [35], and may prove with further work to be the norm for supergenes, as it is for sex
chromosomes in some taxa [50].

A puzzling feature of the BC supergene is the apparent long-term persistence of mul-
tiple haplotype groups with (apparently) the same phenotypic effects. Specifically, the
newly described karamu haplotype found in eastern parts of South Africa is associated
with a colour pattern indistinguishable from that of the orientis haplotype found through-
out South Africa. Although we initially hypothesised that this may represent a case of
ongoing allelic turnover (in which the karamu haplotype may eventually replace the ori-
entis haplotype), this is not supported by the similarly high level of diversity seen in both
haplotype groups, implying that neither has experienced a recent or ongoing selective
sweep. An alternative explanation is that both of these haplotypes may be adapted to
different geographic regions by influencing traits other than colour pattern. It is certainly
possible that the BC supergene region, which contains approximately 150 protein-cod-
ing genes, contributes to other important ecological traits. Simulations have shown that
such adaptations may continue to accumulate after inversions have become established
[11]. It is notable that the karamu haplotype appears to have arisen through recombina-
tion between the southern orientis and eastern klugii haplotypes, and appears to be
most common in a geographic region intermediate between these two (though our
sampling is too sparse to be sure). It is therefore plausible that this haplotype confers an

adaptive advantage by combining an optimal warning pattern for southern Africa with
other environmental adaptations suited to east Africa. A similar scenario may explain
the existence of a dark-coloured morph in North Africa andn the Mediterranean with a
haplotype that is highly similar to that of the pale-coloured chrysippus morph, but carries
orientis-like variants at the promoter of *yellow* that result in dark colouration. In this
case, it may be that recombination allowed for allelic replacement at one locus while
maintaining the broader locally-adapted haplotype. Further investigation into the eco-
logy and fitness of different genotypes and phenotypes is required to test these hypo-
theses.

Another factor that may shape haplotype diversity is vulnerability of inversions to the ac-
cumulation of deleterious mutations. Both theoretical and empirical studies have shown
that inversions can accumulate excess deleterious load as a result of their lower effect-
ive population size compared to the rest of the genome and their reduced opportunities
for recombination [3,23,24]. This may drive turnover if new or recombinant haplotypes
have lower mutational load. It has also been suggested that load may promote poly-
morphism through associative overdominance, in which heterokaryotypes carrying dif-
ferent sets of deleterious recessive alleles have increased fitness relative to homokaryo-
types due to masking of the deleterious recessives [23]. While examples of heterokaryo-
type advantage are known [24], it is difficult to prove that associative overdominance is
the cause, especially given that the conditions under which it is likely to evolve are
highly restrictive [25]. In *D. chrysippus*, there is no compelling evidence for heterozygote
advantage: the three most common haplotype groups each occur in large regions of
monomorphism, and broad clines suggest that polymorphism in the hybrid zone reflects
a balance between selection (local adaptation) and extensive dispersal [37] rather than
heterokaryotype advantage. Further investigation is needed to specifically rule out het-
erokaryotype advantage, especially for the newly identified karamu haplotype and the
other divergent haplotype found north of the Sahara, which are only known from poly-
morphic locations. It is worth noting that the recent fusion of chr15 (chrysippus haplo-
type) to the W chromosome in East Africa [38] represents the origin of yet another diver-
gent haplotype that is only present in the heterozygous state (butterfly females are ZW
and do not undergo recombination), and therefore would be sheltered from the effects
of deleterious recessives.

Finally, it is highly probable that the unusual physical structure of chromosome 15 is at
least part of the explanation for the excessive diversity at the BC supergene. Most not-
ably, the chromosome carries a large copy-number-variable region which began to grow
around 7.5 MYA, and now comprises over a third of the chromosome in the klugii haplo-
type [21]. We hypothesise that this region, which is also rich in transposable elements,
creates fertile ground for the emergence of new divergent haplotypes through two
mechanisms of recombination suppression [21]: It could directly prevent crossing over

by impeding proper meiotic chromosome pairing in individuals with different comple-
ments of copies, and it could indirectly promote recombination suppression by propagat-
ing new inversions through ectopic recombination. Whether the physical characteristics
of this chromosome also facilitated its recent fusion to the W chromosome in East Africa
remains to be determined. The tendency of this one chromosome to be especially prone
to rearrangement echoes a similar pattern seen in the avian chromosome 4, which has
undergone both ancient rearrangements (possibly associated with an ancient su-
pergene [51]), and more recent fusions and further rearrangement in the formation of
neo-sex chromosomes [52–54]. An interesting feature that is shared by the *D. Chrysip-*
pus BC supergene, the *Heliconius numata* P supergene [24] and the fireant social su-
pergene [6], is the presence of multiple adjacent inversions that appear to share break-
points. This modular organisation means that crossovers that occur between two inver-
sions are not suppressed, allowing the distinct modules of the supergene to evolve
somewhat independently. Modular arrangements may simply be a side-effect of 'reuse'
of inversion breakpoints (e.g. in repeat-rich regions), but it is plausible that supergenes
with this structure may be more likely to persist long-term if it avoids some of the costs
of complete recombination suppression.

In conclusion, we have shown that a supergene with an apparently simple phenotypic
association shows unexpected diversity at the haplotype level. Our findings bring to light
the nuanced relationship between supergenes and recombination, in which incomplete
recombination suppression can fuel haplotype diversification, and possibly support long-
term persistence. Further work is needed to understand the selective and ecological
drivers underlying the observed diversity. Our study adds to a growing body of work re-
vealing the indirect effects of structural variation on the evolution of surrounding se-
quences [24,33,55,56]. Further work across a diverse range of taxa, along with popula-
tion genetic modelling will help to uncover the generality of these phenomena.

Methods:

Data collection, whole genome sequencing and genotyping

We analysed whole genome sequence data (Illumina, 150 bp paired-end) from 174 but-terflies, including 158 wild-caught butterflies representing each of the main colour morphs across Africa and the hybrid zone (Fig. 1A, S2 Table) and 12 captively reared individuals. Samples collection was conducted with the permission of landowners and under appropriate permits where relevant: NACOSTI/P15/3290/3607, NACOSTI/P15/2403/3602, National Commission for Science and Technology, Kenya; MINEDUC/S&T/459/2017, Ministry of Education, Rwanda; MPB5667 Mpumalanga Parks and Tourism Agency, South Africa; FAUNA0615202, Northern Cape Department of Environment and Nature Conservation, South Africa; EMDEP006/17, Environment & Natural Resources Directorate, St Helena Government; Permit 604 (2017), Council of Fuerteventura, Spain.

A subset of individuals had been sequenced previously [21,38,57]. For those newly se-quenced in this study, the same protocols were followed. Briefly, genomic DNA was ex-tracted from ethanol-preserved tissue using the Qiagen DNEasy Blood and Tissue Kit (Qiagen, Redwood City, CA). Illumina library preparation and sequencing (150bp, paired-end) was performed by Novogene (Cambridge, UK), using a Novaseq 6000 in-strument.

Reads were mapped to the Dchry2.2 reference genome (ENA accession GCA_916720795.1; [57]) using BWA mem v0.7.17 [58] with default parameters, and PCR duplicates were removed using PicardTools v2.11.1 (https://broadinstitute.git-hub.io/picard/). Indel realignment was then carried out using GATK v.3.8 [59]. To identify any substantial variation in read depth across the dataset, mean read depth was calculated using Mosdepth [60]. One individual was found to have substantially higher read depth (SM18W01) and as a result the bam file for this individual was randomly downsampled to 30% of its original size using SAMtools ([61]; samtools view -s 0.30) to eliminate biases in genotyping. Genotyping was carried out using BCFtools [62], with genotypes on the Z chromosome (chr1) called by defining ploidy based on known sex (i.e. females as haploid and males as diploid). Genotypes were then filtered using BCFtools to remove any positions called fewer than twice and to leave only genotypes with an individual depth of >= 7 and with genotype quality of >=30. Additional filtering re-moved SNPs with >10% missing data (corresponding to a haploid count of 313 across autosomes and 224 on the Z chromosome).

Population structure, diversity and divergence

Genomic variation across our dataset, spanning different regions of the genome, was assessed by producing PCAs from different sets of SNPs using PLINK [63]. Two SNP sets were used, representing (i) the autosomal regions of the genome excluding chr15

24   24

as well as the Z sex chromosome chr1 (14,548,985 SNPs), and (ii) rearranged regions
1, 2, and 4 of the BC supergene on chr15, described in [21] (173,998 SNPs). Genome-
wide nucleotide diversity (π), F_{ST}, and d_{XY} were computed in non-overlapping 50kb win-
dows across the genome using the script *popgenWindows.py* (github.com/si-
monhmartin/genomics_general release 0.2) and net divergence (d_a) was calculated
from this output (calculated as $d_a = d_{XY} - (\pi_{P1} + \pi_{P2})/2$ where P1 and P2 represent Popula-
tion 1 and Population 2, respectively).

Genome-wide association
The association between SNP variation and phenotypic variation was analysed using a
genome-wide association analysis implemented in PLINK across all chromosomes.
Phenotypes of scanned wings from 172 individuals were scored following [37] based on
whether the background colour was pale (1), intermediate (2), or dark (3), and whether
the forewing band was absent (1), partial (2), or present (3) (wings were not available to
phenotype for two individuals resulting in their omission from the GWAS). Empirical p-
values were computed using the default adaptive permutation method implemented in
PLINK. To confirm that geographic sampling structure alone does not underpin the pat-
terns of association observed across the genome we re-ran the GWAS using only the
87 individuals from the hybrid zone for which we had phenotypes (populations NYA,
NRB, and MPL).

Visualising variation in the BC supergene using NeighborNet
To visualise genetic relationships and clustering across the BC supergene, we gener-
ated a phylogenetic network based on pairwise genetic distances among diploid indi-
viduals using the Neighbor-Net algorithm [64], implemented in SplitsTree [65]. Pairwise
distances were computed using the script distMat.py (github.com/simonhmartin/genom-
ics_general release 0.2), which computes the average pairwise distance between the
four pairs of sequence haplotypes for each pair of diploid individuals. This analysis was
computed using modules 1, 2 and 4 of the supergene region.

Admixture
We aimed to naively determine the number of distinct haplotype groups of the su-
pergene by modelling each individual in terms of its ancestry from a fixed number of
source populations. To this end, we ran Admixture v1.3.0 [43] on a concatenated data-
set of loci from across the supergene modules 1, 2, and 4 (i.e. excluding the CNV re-
gion). Admixture was run specifying a range of clusters from k=3 to k=12 (Figure D in
S1 Text) and 20-fold cross-validation error (--cv=20) was computed.

Phasing
Phasing was carried out to assist with ancestry painting using two consecutive tools,

Whatshap v1.1 [66], a read-based phasing software, and Shapeit4 v4.2.2 [67] a statist-
ical phasing software using default parameters.

Ancestry painting and additional phase correction
To visualise genetic clustering of phased haplotypes, we used two approaches for an-
cestry painting along the chromosome. Both approaches require defined reference indi-
viduals, which were selected based on the Admixture analysis above. Reference indi-
viduals used for each analysis are described in the Results.

The first approach was Loter [44], which uses a hidden Markov model to assign ances-
try and identify breakpoints along the genome. First, phased genotypes were filtered us-
ing filterGenotypes.py (github.com/simonhmartin/genomics_general release 0.2) to re-
move any sites at which more than 60% of individuals are heterozygous, as such sites
are likely to be caused by mis-mapping and could be prone to break tracts of ancestry.
Haploid genotypes were output as counts of the minor allele (0 or 1), and Loter was run
using the python API in a wrapper script (loter_wrapper.py), with options rate_vote=0,
nb_bagging=20. Adjacent SNPs with identical ancestry scores were concatenated into
tracts to reduce the output file size.

A second approach for ancestry painting that we call 'distPaint' was applied to confirm
the Loter results, while providing a coarser visualisation at the whole-chromosome
scale. This approach uses a sliding-window along the genome and assigns ancestry for
each haplotype based on genetic distance (d_{XY}) from the reference sets of individuals. A
window size of 200 SNPs was used. Ancestry was assigned if this difference was signi-
ficantly lower for one reference population according to a Wilcoxon Rank Sum test
($p<=0.01$), otherwise ancestry was set to undefined. This was implemented using a cus-
tom python script distPaint.py (github.com/simonhmartin/genomics_general).

Initial visualisation of the ancestry painting from both approaches revealed obvious
cases of phasing errors (Figures K in S1 Text). Such errors could lead to false-positive
inference of recombination events. For example, say an individual is phased into two
haplotypes, that are then painted with ancestries "c-c-c-c-k" and "k-k-k-k-c" (where c
and k represent tracts of chromosome assigned to reference populations chrysippus
and klugii, respectively). This could either represent a case of an individual carrying two
recombinant haplotypes in which both recombination events happened to occur at the
same genomic region, or it could be a case of a single phasing error in an individual that
in fact caries two common haplotypes (c-c-c-c-c and k-k-k-k-k). The latter is of course
far more likely. We therefore applied a final heuristic phase correction approach to the
inferred ancestries which attempts to maximise similarity among haplotypes at the
whole-chromosome scale, thereby minimising inferred recombination events (Figure K
in S1 Text). For each diploid individual, this heuristic approach iterates over each an-

cestry-painted tract in the chromosome (e.g. from Loter) and switches the phase if this
would cause an increase in similarity of the two resulting haplotypes to one or more
other haplotypes in the full dataset (considering all ancestry blocks up to and including
the focal one). Because we are interested in identifying common shared haplotypes, we
consider the average similarity of the top three most similar haplotypes in the rest of the
dataset. The process is performed left-to-right and then right-to-left for each individual,
and then repeated again starting from the first individual, up to a total of 20 iterations for
Loter outputs and 100 iterations for distPaint.py outputs (the latter have fewer intervals,
allowing for more rapid computation). This algorithm is implemented in the script
phasepaint.py (https://github.com/simonhmartin/phasepaint).

Topology weighting
To complement the ancestry painting described above, we analysed changes in rela-
tionships among supergene haplotypes across the chromosome (as an indicator of pos-
sible recombination), using topology weighting [46], implemented in TWISST2 (https://
github.com/simonhmartin/twisst2). This approach infers genealogies and where they
change along the genome, and outputs a 'weighting' for each possible topology for the
relationships between defined groups of individuals in each genomic region. The input
variants file was further filtered to remove sites with more than 75% heterozygotes
(which are likely genotyping errors that could interfere with tree inference) and sites with
missing data for more than 50% of individuals. An outgroup was needed for polarisation,
for which we used a single *Danaus melanippus* individual (S2 Table), following the
alignment and genotyping procedures described above.

Modelling gene flux using gIMble
To detect and quantify gene flux between supergene haplotypes, we fitted isolation-
with-migration models for individuals homozygous for the chrysippus, klugii, orientis al-
leles using the gIMble framework [47]. Under the IM model, gene-flux between su-
pergene haplotypes is modelled as migration (gene flow) between panmictic popula-
tions. Six klugii individuals (WAT population), seven orientis individuals (TSW popula-
tion), and six chrysippus (NGA population) were included in the analysis (S2 Table).
Since gIMble carries out pairwise tests this resulted in six pairwise tests for each of the
4 regions tested. A VCF containing only data from these 19 individuals was first prepro-
cessed using *gimble preprocess*. Next, intergenic bed files were produced by using
bedtools intersect and bedtools subtract to extract and then exclude genic regions using
the reference annotation file. *gimble parse* was then run for each combination of su-
pergene region (i.e. regions 1.1, 1.2, 2, and 4, Fig. 3A) and pairwise comparison, fol-
lowed by *gimble blocks*, *gimble windows*, *gimble info*, and *gimble tally*. *gimble optimize*
was then run for each combination of region and pairwise test specifying each of three
models, DIV - a divergence only model, IM_AB an isolation with migration model where
gene flow occurs from population A into B, and IM_BA an isolation with migration model

where gene flow occurs from population B into A, resulting in a total of 36 models.
gimble query was then run to extract model summary information, including composite
likelihood, for each model allowing us to determine the best fitting of the three models
for each of the 12 region and pair combinations (full results are reported in https://git-
hub.com/RishiDeKayne/Danaus_WGS/blob/main/FINAL_GIMBLE_RESULTS_3morph-
s.xlsx). For detailed commands and model parameters see https://github.com/
RishiDeKayne/Danaus_WGS/blob/main/Danaus_WGS_commands.txt.

Acknowledgements:
We thank the following people for assistance with sampling, breeding and acquisition of
permissions: Rwanda - Constantin Sibomana and Jody Garbe; Kenya - Piera Ireri, Ivy
Ng'Iru, Godfrey Etelej, David Smith, Anna Orteu, Jenny York, Owen McMillan, Valarie
McMillan; South Africa - Jeremy Dobson; Ghana - Oskar Brattstrom; Spain (Canary Is-
lands) - Yeray Monasterio, David Smith; Tunisia - Roger Vila ; Italy: Richard ffrench-
Constant; St. Helena: David Pryce. We also thank Brian Charlesworth, Deborah Char-
lesworth, David Smith, Richard ffrench-Constant, Chay Graham, Thomas Decroly, Alex-
ander Mackintosh, Domink Laetsch and Konrad Lohse for valuable input on this work.

Data availability:
All raw sequencing reads are available via the European Nucleotide Archive (ENA) -
Sample accession numbers are provided in S2 Table. Commands and scripts for bioin-
formatics analyses can be found at https://github.com/RishiDeKayne/Danaus_WGS/.

Funding:
This work was supported by a The Royal Society (grants URF\R1\180682, RGF\EA\
181071, URF\R\231034 to S.H.M), the Swiss National Science Foundation (grants
P2BEP3_195567, P500PB_211005 to R.D.K) and the National Geographic Society
(grant WW-138R-17 to I.J.G).

References:

1. Charlesworth B, Charlesworth D. Selection of new inversions in multi-locus genetic systems. Genet Res . 1973;21: 167–183.

2. Thompson MJ, Jiggins CD. Supergenes and their role in evolution. Heredity . 2014;113: 1–8.

3. Charlesworth B. The effects of inversion polymorphisms on patterns of neutral genetic diversity. Genetics. 2023;224. doi:10.1093/genetics/iyad116

4. Mérot C, Llaurens V, Normandeau E, Bernatchez L, Wellenreuther M. Balancing selection via life-history trade-offs maintains an inversion polymorphism in a seaweed fly. Nat Commun. 2020;11: 670.

5. Küpper C, Stocks M, Risse JE, Dos Remedios N, Farrell LL, McRae SB, et al. A supergene determines highly divergent male reproductive morphs in the ruff. Nat Genet. 2016;48: 79–83.

6. Yan Z, Martin SH, Gotzek D, Arsenault SV, Duchen P, Helleu Q, et al. Evolution of a supergene that regulates a trans-species social polymorphism. Nat Ecol Evol. 2020;4: 240–249.

7. Li J, Cocker JM, Wright J, Webster MA, McMullan M, Dyer S, et al. Genetic architecture and evolution of the S locus supergene in Primula vulgaris. Nat Plants. 2016;2: 16188.

8. Kim K-W, Bennison C, Hemmings N, Brookes L, Hurley LL, Griffith SC, et al. A sex-linked supergene controls sperm morphology and swimming speed in a songbird. Nat Ecol Evol. 2017;1: 1168–1176.

9. Komata S, Kajitani R, Itoh T, Fujiwara H. Genomic architecture and functional unit of mimicry supergene in female limited Batesian mimic Papilio butterflies. Philos Trans R Soc Lond B Biol Sci. 2022;377: 20210198.

10. Kirkpatrick M, Barton N. Chromosome inversions, local adaptation and speciation. Genetics. 2006;173: 419–434.

11. Schaal SM, Haller BC, Lotterhos KE. Inversion invasions: when the genetic basis of local adaptation is concentrated within inversions in the face of gene flow. Philos Trans R Soc Lond B Biol Sci. 2022;377: 20210200.

12. Berdan EL, Barton NH, Butlin R, Charlesworth B, Faria R, Fragata I, et al. How chromosomal inversions reorient the evolutionary process. J Evol Biol. 2023;36: 1761–1782.

13. Lowry DB, Willis JH. A widespread chromosomal inversion polymorphism contributes to a major life-history transition, local adaptation, and reproductive isolation. PLoS Biol. 2010;8. doi:10.1371/journal.pbio.1000500

14. Hager ER, Harringmeyer OS, Wooldridge TB, Theingi S, Gable JT, McFadden S, et al. A chromosomal inversion contributes to divergence in multiple traits between deer mouse ecotypes. Science. 2022;377: 399–405.

15. Huang K, Andrew RL, Owens GL, Ostevik KL, Rieseberg LH. Multiple chromosomal inver-
sions contribute to adaptive divergence of a dune sunflower ecotype. Mol Ecol. 2020;29:
2535–2549.

16. Kapun M, Fabian DK, Goudet J, Flatt T. Genomic Evidence for Adaptive Inversion Clines in
Drosophila melanogaster. Mol Biol Evol. 2016;33: 1317–1336.

17. Todesco M, Owens GL, Bercovich N, Légaré J-S, Soudi S, Burge DO, et al. Massive haplo-
types underlie ecotypic differentiation in sunflowers. Nature. 2020;584: 602–607.

18. Harringmeyer OS, Hoekstra HE. Chromosomal inversion polymorphisms shape the gen-
omic landscape of deer mice. Nat Ecol Evol. 2022;6: 1965–1979.

19. Morales HE, Faria R, Johannesson K, Larsson T, Panova M, Westram AM, et al. Genomic
architecture of parallel ecological divergence: Beyond a single environmental contrast. Sci
Adv. 2019;5: eaav9963.

20. Joron M, Frezal L, Jones RT, Chamberlain NL, Lee SF, Haag CR, et al. Chromosomal re-
arrangements maintain a polymorphic supergene controlling butterfly mimicry. Nature.
2011;477: 203–206.

21. Kim K-W, De-Kayne R, Gordon IJ, Omufwoko KS, Martins DJ, Ffrench-Constant R, et al.
Stepwise evolution of a butterfly supergene via duplication and inversion. Philos Trans R
Soc Lond B Biol Sci. 2022;377: 20210207.

22. Roesti M, Gilbert KJ, Samuk K. Chromosomal inversions can limit adaptation to new envir-
onments. Mol Ecol. 2022;31: 4435–4439.

23. Berdan EL, Blanckaert A, Butlin RK, Bank C. Deleterious mutation accumulation and the
long-term fate of chromosomal inversions. PLoS Genet. 2021;17: e1009411.

24. Jay P, Chouteau M, Whibley A, Bastide H, Parrinello H, Llaurens V, et al. Mutation load at a
mimicry supergene sheds new light on the evolution of inversion polymorphisms. Nat
Genet. 2021;53: 288–293.

25. Charlesworth B. The fitness consequences of genetic divergence between polymorphic
gene arrangements. Genetics. 2024;226. doi:10.1093/genetics/iyad218

26. Jay P, Aubier TG, Joron M. The interplay of local adaptation and gene flow may lead to the
formation of supergenes. Mol Ecol. 2024; e17297.

27. Sturtevant AH, Beadle GW. The Relations of Inversions in the X Chromosome of Droso-
phila Melanogaster to Crossing over and Disjunction. Genetics. 1936;21: 554–604.

28. Chovnick A. Gene conversion and transfer of genetic information within the inverted region
of inversion heterozygotes. Genetics. 1973;75: 123–131.

29. Crown KN, Miller DE, Sekelsky J, Hawley RS. Local Inversion Heterozygosity Alters Re-
combination throughout the Genome. Curr Biol. 2018;28: 2984–2990.e3.

30. Korunes KL, Noor MAF. Pervasive gene conversion in chromosomal inversion heterozy-
gotes. Mol Ecol. 2019;28: 1302–1315.

31. Corbett-Detig RB, Hartl DL. Population genomics of inversion polymorphisms in Drosophila
melanogaster. PLoS Genet. 2012;8: e1003056.

32. Kapun M, Mitchell ED, Kawecki TJ, Schmidt P, Flatt T. An Ancestral Balanced Inversion
Polymorphism Confers Global Adaptation. Mol Biol Evol. 2023;40. doi:10.1093/molbev/
msad118

33. Lundberg M, Mackintosh A, Petri A, Bensch S. Inversions maintain differences between mi-
gratory phenotypes of a songbird. Nat Commun. 2023;14: 452.

34. Hill J, Enbody ED, Bi H, Lamichhaney S, Lei W, Chen J, et al. Low Mutation Load in a Su-
pergene Underpinning Alternative Male Mating Strategies in Ruff (Calidris pugnax). Mol Biol
Evol. 2023;40. doi:10.1093/molbev/msad224

35. Palmer DH, Kronforst MR. A shared genetic basis of mimicry across swallowtail butterflies
points to ancestral co-option of doublesex. Nat Commun. 2020;11: 6.

36. Smith DAS, Owen DF, Gordon IJ, Lowis NK. The butterfly Danaus chrysippus (L.) in East
Africa: polymorphism and morph-ratio clines within a complex, extensive and dynamic hy-
brid zone. Zool J Linn Soc. 1997;120: 51–78.

37. Liu W, Smith DAS, Raina G, Stanforth R, Ng'Iru I, Ireri P, et al. Global biogeography of
warning coloration in the butterfly Danaus chrysippus. Biol Lett. 2022;18: 20210639.

38. Martin SH, Singh KS, Gordon IJ, Omufwoko KS, Collins S, Warren IA, et al. Whole-chromo-
some hitchhiking driven by a male-killing endosymbiont. PLoS Biol. 2020;18: e3000610.

39. Smith DAS, Gordon IJ, Traut W, Herren J, Collins S, Martins DJ, et al. A neo-W chromo-
some in a tropical butterfly links colour pattern, male-killing, and speciation. Proc Biol Sci.
2016;283. doi:10.1098/rspb.2016.0821

40. Smith DA. Evidence for autosomal meiotic drive in the butterfly Danaus chrysippus L.
Heredity . 1976;36: 139–142.

41. Wittkopp PJ, Vaccaro K, Carroll SB. Evolution of yellow gene regulation and pigmentation
in Drosophila. Curr Biol. 2002;12: 1547–1556.

42. Galant R, Skeath JB, Paddock S, Lewis DL, Carroll SB. Expression pattern of a butterfly
achaete-scute homolog reveals the homology of butterfly wing scales and insect sensory
bristles. Curr Biol. 1998;8: 807–813.

43. Alexander DH, Lange K. Enhancements to the ADMIXTURE algorithm for individual ances-
try estimation. BMC Bioinformatics. 2011;12: 246.

44. Dias-Alves T, Mairal J, Blum MGB. Loter: A Software Package to Infer Local Ancestry for a
Wide Range of Species. Mol Biol Evol. 2018;35: 2318–2326.

45. Chen J-M, Cooper DN, Chuzhanova N, Férec C, Patrinos GP. Gene conversion: mechan-
isms, evolution and human disease. Nat Rev Genet. 2007;8: 762–775.

46. Martin SH, Van Belleghem SM. Exploring Evolutionary Relationships Across the Genome
Using Topology Weighting. Genetics. 2017;206: 429–438.

47.   Laetsch DR, Bisschop G, Martin SH, Aeschbacher S, Setter D, Lohse K. Demographically
explicit scans for barriers to gene flow using gIMble. PLoS Genet. 2023;19: e1010999.

48.   Davey JW, Chouteau M, Barker SL, Maroja L, Baxter SW, Simpson F, et al. Major Improve-
ments to the Heliconius melpomene Genome Assembly Used to Confirm 10 Chromosome
Fusion Events in 6 Million Years of Butterfly Evolution. G3 . 2016;6: 695–708.

49.   Shipilina D, Näsvall K, Höök L, Vila R, Talavera G, Backström N. Linkage mapping and
genome annotation give novel insights into gene family expansions and regional recombin-
ation rate variation in the painted lady (Vanessa cardui) butterfly. Genomics. 2022;114:
110481.

50.   Vicoso B. Molecular and evolutionary dynamics of animal sex-chromosome turnover. Nat
Ecol Evol. 2019;3: 1632–1641.

51.   Mirarab S, Rivas-González I, Feng S, Stiller J, Fang Q, Mai U, et al. A region of suppressed
recombination misleads neoavian phylogenomics. Proc Natl Acad Sci U S A. 2024;121:
e2319506121.

52.   Sigeman H, Strandh M, Proux-Wéra E, Kutschera VE, Ponnikas S, Zhang H, et al. Avian
Neo-Sex Chromosomes Reveal Dynamics of Recombination Suppression and W Degener-
ation. Mol Biol Evol. 2021;38: 5275–5291.

53.   Sigeman H, Zhang H, Ali Abed S, Hansson B. A novel neo-sex chromosome in Sylvietta
brachyura (Macrosphenidae) adds to the extraordinary avian sex chromosome diversity
among Sylvioidea songbirds. J Evol Biol. 2022;35: 1797–1805.

54.   Dierickx EG, Sin SYW, van Veelen HPJ, Brooke M de L, Liu Y, Edwards SV, et al. Genetic
diversity, demographic history and neo-sex chromosomes in the Critically Endangered
Raso lark. Proc Biol Sci. 2020;287: 20192613.

55.   Huang K, Ostevik KL, Elphinstone C, Todesco M, Bercovich N, Owens GL, et al. Mutation
Load in Sunflower Inversions Is Negatively Correlated with Inversion Heterozygosity. Mol
Biol Evol. 2022;39. doi:10.1093/molbev/msac101

56.   Jeong H, Baran NM, Sun D, Chatterjee P, Layman TS, Balakrishnan CN, et al. Dynamic
molecular evolution of a supergene with suppressed recombination in white-throated spar-
rows. Elife. 2022;11. doi:10.7554/eLife.79387

57.   Singh KS, De-Kayne R, Omufwoko KS, Martins DJ, Bass C, ffrench-Constant R, et al. Gen-
ome assembly of Danaus chrysippus and comparison with the Monarch Danaus plexippus.
G3 . 2021. doi:10.1093/g3journal/jkab449

58.   Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler transform.
Bioinformatics. 2010;26: 589–595.

59.   Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Auwera GAV der, et al.
Scaling accurate genetic variant discovery to tens of thousands of samples. bioRxiv. 2017;
201178.

60.   Pedersen BS, Quinlan AR. Mosdepth: quick coverage calculation for genomes and ex-
omes. Bioinformatics. 2018;34: 867–868.

61. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Align-
ment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.

62. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of
SAMtools and BCFtools. Gigascience. 2021;10. doi:10.1093/gigascience/giab008

63. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: a tool
set for whole-genome association and population-based linkage analyses. Am J Hum
Genet. 2007;81: 559–575.

64. Bryant D, Moulton V. Neighbor-net: an agglomerative method for the construction of phylo-
genetic networks. Mol Biol Evol. 2004;21: 255–265.

65. Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. Mol Biol
Evol. 2006;23: 254–267.

66. Martin M, Patterson M, Garg S, Fischer SO, Pisanti N, Klau GW, et al. WhatsHap: fast and
accurate read-based phasing. bioRxiv. 2016. p. 085050. doi:10.1101/085050

67. Delaneau O, Zagury J-F, Robinson MR, Marchini JL, Dermitzakis ET. Accurate, scalable
and integrative haplotype estimation. Nat Commun. 2019;10: 5436.