Emissions of climate-altering species from open vegetation fires in the

Mediterranean region - A review on methods and data

Rabia Ali Hundal[1,2,3], Saurabh Annadate[1,2,3], Rita Cesari,[4] Alessio Collalti[5,6] , Michela Maione[1,2], Paolo

Cristofanelli[1,]*

[1] Institute of Atmospheric Sciences and Climate (ISAC), National Research Council, 73100, I-40129 Bologna, Italy

[2] University of Urbino - Department of Pure and Applied Sciences, I-61029, Urbino, Italy

[3] Scuola Universitaria Superiore Pavia (IUSS), I-27100, Pavia, Italy

[4] Institute of Atmospheric Sciences and Climate (ISAC), National Research Council, 73100, Lecce, Italy

[5] Forest Modelling Lab., Institute for Agriculture and Forestry Systems in the Mediterranean (ISAFOM), National
Research Council of Italy, 06128, Perugia, Italy

[6]National Biodiversity Future Center (NBFC), 90133, Palermo, Italy
12

13
14 *Corresponding author: Paolo Cristofanelli, phone: +39 051 6399597, fax: +39 051 6399658,*
15 *p.cristofanelli@isac.cnr.it*
16

Abstract

The climate change over the Mediterranean region poses serious concerns about the role

of open vegetation fires in the emissions of climate-altering species. The aim of this work

is to review the current methodologies for quantifying the emissions of greenhouse gases

and black carbon from open vegetation fires, as well as the data provided by four state-

of-the-art inventories of emissions of carbon dioxide (CO_2), methane (CH_4), nitrous oxide

(N_2O) and black carbon (BC) in the Mediterranean region for the period 2003 - 2020.

A limited number of studies specifically addressed the quantification of emissions from

open fires in the Mediterranean region. Our data review of fire emissions in the

Mediterranean region, where "top-down" methods have not yet implemented, reveals

discrepancies across the four inventories examined (GFED v4.1s, GFAS v1.2, FINN v2.5,

and EDGAR v8.0). Among these, FINN v2.5 consistently reported the highest emissions,

while GFED v4.1s reported the lowest. We observed that the relative ranking of total

emissions between the inventories varied for the species considered (e.g. CO_2 vs. CH_4)

and that different proportions of emissions were attributed to the individual countries

included in the Mediterranean domain. We argued that these differences were related to

the different spatial resolutions of the input data used to detect the occurrence of fires,

the different approaches to calculating the amount of fuel available, and the emission

factors used.

The three inventories reporting wildfire emissions were consistent in identifying the

occurrence of peaks in the emissions for the years 2007, 2012 and 2017. We

hypothesized that La Niña events could partially contribute to triggering the occurrence

of these emission peaks.

To increase the accuracy and consistency of climate-altering emission data related to

open vegetation fires in the Mediterranean region, we recommend to integrate bottom-up

approaches with top-down inversion methods based on satellite and in-situ atmospheric

observations.

Keywords: Open Vegetation Fires, Greenhouse Gases, Black Carbon, Emission

Inventories, Mediterranean Region, Climate Change, Wildfire Emissions.

1. Introduction

Since the 19th century, anthropogenic activities have led to a continuous upward trend of greenhouse gas (GHG) emissions (Roelfsema et al., 2020), affecting various aspects of climate, including air and ocean temperatures, sea levels, extreme events related to heat waves and precipitation. In 2022, global anthropogenic GHG emissions reached a new high of 53.8 Gt of CO_2 equivalent, according to the latest version of the Emissions Database for Global Atmospheric Research (EDGAR v8.0, see Crippa et al., 2023). In 2015, the Paris Agreement established emission targets to keep the rise in global temperature within 1.5°C above pre-industrial levels (UNFCCC, 2015).

According to the latest World Meteorological Organization GHG bulletin (World Meteorological Organization, 2023), the three main GHGs, carbon dioxide (CO_2), methane (CH_4) and nitrous oxide (N_2O), have increased by 150%, 264% and 124% respectively in 2022 compared to 1750. Biomass burning related to open vegetation fires is a relevant source for all of them (Crutzen and Andreae, 1990; Shi et al., 2021). As suggested by Landry (2016), it should be noted that changes in the ecosystem dynamics due to climate change may add further uncertainty to the impact of wildfires on the decadal carbon cycle.

Open vegetation fires also emit dangerous pollutants such as particulate matter, carbon monoxide (CO), nitrogen oxides (NO_x), sulfur dioxide (SO_2), and volatile organic compounds, which can harm human health and ecosystems (Anav et al., 2024; Bowman and Johnston, 2005; De Marco et al., 2022). Particulate matter is also associated with major climatic impacts via aerosol-radiation and aerosol-cloud interactions (Intergovernmental Panel On Climate Change, 2023). Globally, biomass burning

contributes nearly 30-40% of the atmospheric black carbon (BC), a component of atmospheric aerosol particles resulting from the incomplete combustion of fossil fuels, wood (Bond et al., 2013). By absorbing solar radiation, BC, impacts the climate by heating the atmosphere, accelerating ice melt, and altering cloud properties (Kang et al., 2019, 2020; Klimont et al., 2017). According to Hamilton et al. (2018), reassessment of pre-industrial fire emissions has been shown to have a significant impact on anthropogenic aerosol forcing.

In Europe, emissions from open vegetation fires have been linked to climate variability and change, with anthropogenic influences being an important component (Fernandes et al., 2016; Song et al., 2009). Increasing aridity has been observed to push Mediterranean forest post-fire recovery towards open shrublands (Baudena et al., 2020), aggravating the frequency and intensity of fires in this region as predicted by non-stationary climate-fire models due to global warming (Turco et al., 2019). Wildfires and prescribed burns (controlled fires used for forest management, see Henderson et al. (2005)), the two main types of open vegetation fires that occur most frequently in Europe, both contribute to emissions of pollutants and GHGs. Wildfire intensity is driven by climate change, fuel accumulation, expanding human settlements, flammable vegetation, and prolonged burning (Bowman and Johnston, 2005). The Euro-Mediterranean southern region, where the climate is warmer and drier   is where Europe's wildfires are (and will be) most frequent (Lionello and Scarascia, 2018; Moreira et al., 2011; Noce et al., 2016).

Emissions from open vegetation fires are not as well characterised as other anthropogenic emissions due to the high variability in time and space, the different factors driving activity and the efficiency of emissions (Randerson et al., 2012), including the size

and the stage of the fires, the type of vegetation being burned, weather conditions, and

human intervention.

Recent studies have highlighted the importance of refining methodologies for emission

estimates from open vegetation fires. For instance, advances in satellite-based burned

area detection, together with the integration of detailed data on fuel types, meteorological

conditions and combustion phases have significantly improved accuracy of bottom-up

approaches (Scarpa et al., 2024). Furthermore, innovative observation-informed "top-

down" methods incorporating atmospheric chemical transport models have provided new

insights into emissions dynamics (Qu et al., 2021, Zhang et al., 2021). Emerging

research, such as the study by Pande et al. (2021), demonstrates the effectiveness of

integrating satellite data and GIS techniques to estimate biomass resources in semi-arid

regions, highlighting the utility for advanced geospatial methods for biomass burning.

Despite these advances, significant challenges remain, including biases in emission

factors, inconsistencies in detecting smaller fires and limited application of these

techniques in the Mediterranean region. These challenges underscore the need for

standardised, high-resolution methodologies to improve emission estimates and support

climate policies

The different existing methodologies to estimate open vegetation fire emissions can be

broadly classified as "bottom-up" or "top-down" (Kasischke and Penner, 2004).

The "bottom-up" approach is based on the combination of geophysical (e.g. type of

vegetation within a region, proportion of vegetation burned) and biophysical (emission

factors) information. The so-called "top-down" approach uses direct atmospheric

observations to quantify the net emissions. Traditionally, the "top-down" approach refers

to an energy based method. It uses the fire radiative power (FRP) measurements from

satellite remote sensing as a direct indicator of the biomass consumed by a fire (Roberts

et al., 2005). FRP is then multiplied by a predefined factor, known as the smoke emission

coefficient, to estimate pollutant's emission rates. Rather, in this review, we will refer to

the term "top-down" to indicate methods that do not directly use emission factors to

calculate emissions, but rather use observations in combination with inverse modeling

techniques (Bergamaschi et al., 2018).

To effectively estimate the impact of open vegetation fires on climate, it is essential to

have a thorough understanding of the relative performance of different approaches,

although each has strengths and limitations. Despite the critical role of fires as a source

of GHGs and BC, to our knowledge there is little specific information on open vegetation

fire emissions in the Mediterranean region. For this reason, this paper aims to review

existing methodologies to quantify GHG and BC emissions from open vegetation fires

and data provided by four state-of-the-art emission inventories GFEDv4.1s, GFASv1.2,

FINNv1.2 and EDGARv8.0 related to emissions of CO_2, CH_4, N_2O, and BC from open

vegetation fires over the Mediterranean region in the period 2003 - 2020.

Although global studies frequently address biomass-burning emissions, a comprehensive

regional analysis tailored to the entire Mediterranean region, which is particularly

susceptible to climate-driven alterations in fire dynamics, is still unavailable. This paper

provides a comparative evaluation of methodologies and data from major emission

inventories, focusing on their relative strengths and limitations. By addressing these gaps,

this review offers insights to improve emission quantification, particularly for climate

altering species, and supports future research and policy making in the Mediterranean

region

The paper is structured as follows: in Section 2, we review the methods used to calculate

the fire emissions, including both bottom-up and observation-informed "top-down"

(inverse modelling) methods, and report on use cases where available.. In Section 3, we

present the data review about emissions of climate-altering species in the Mediterranean

region related to open vegetation fires. Section 4 discusses the emissions estimates and

Section 5 reports our conclusions.

2. Methods used for estimating emissions of climate-altering species from

vegetation fires: an overview

In this section, we broadly describe two widely used approaches to quantify GHG and BC

emissions from open vegetation fires: the so-called "bottom-up" and observation-informed

"top-down" approaches.

2.1 Bottom-Up

In the bottom-up approach, emissions are derived based on activity data (e.g. area

burned, fuel consumption) and emission factors (i.e. a coefficient that describes the rate

at which a given activity releases a specific molecule into the atmosphere). This approach

is commonly used by environmental scientists and policymakers, offering transparency

and replicability in calculating emissions' contribution for specified areas.

A "bottom-up" system for quantifying open vegetation fire emissions consists of several

interrelated modules (see Figure SM1 in the Supplementary Material), each of which

provides specific information for a comprehensive emission estimate (van der Werf et al.,

2017). These modules include the detection of burned areas using satellite data such as

the Visible Infrared Imaging Radiometer Suite (VIIRS) (a sensor onboard the polar-

orbiting Suomi National Polar Orbiting Partnership; Suomi NPP, NOAA-21 and NOAA-21

weather satellites) or the Moderate Resolution Imaging Spectroradiometer (MODIS) (a

key instrument on board the Terra and Aqua satellites), the calculation of fuel

consumption based on biomass burned per unit area and the implementation of emission

factors to convert biomass into specific trace gas and aerosol emissions. Temporal

distribution modules consistently distribute monthly emissions into daily and hourly

values, taking into consideration fire spread rates and diurnal trends. Module outputs are

then aggregated into a geospatial framework that maps emissions across landscapes.

These components interact through data integration and module calibration to ensure

consistency with empirical data and remote sensing. The system provides a

comprehensive understanding of the wildfire impacts, including spatially explicit and

temporally resolved emissions estimates. By incorporating burned area, fuel

consumption, combustion completeness, emission variables, and temporal distribution,

the "bottom-up" system improves our ability to monitor and manage the environmental

and air quality impacts of wildfires. Volkova et al. (2019) enhanced Australia's GHG

emissions reporting from forest fires using a Tier 2 approach from the 2006

Intergovernmental Panel on Climate Change (IPCC) Guidelines. They refined the

emission estimate equations and incorporated detailed, country-specific data from

literature reviews and field measurements. This included improvements in parameters

such as the area burned, fuel mass, combustion factors, and emission factors.

Additionally, they transitioned to a more accurate model for dead organic matter recovery.

These refinements aimed to provide precise GHG emission estimates.

A few studies have used a bottom-up approach to estimate GHG, including CO_2, CH_4, and

N_2O, emissions from open vegetation fires in Europe.

A study by Rosa et al. (2011), estimated atmospheric emissions from wildfires in Portugal

from 1990 to 2000. The researchers used Landsat-based burnt area maps, land cover

maps, national forest inventory data, biometric models, and emissions factors from

literature. Emissions were calculated based on the area burnt, biomass loading,

combustion factor, and emission factor specific to land cover types. The results showed

a strong correlation between the area burnt and the emissions, with the combustion factor

for shrubs and emission factors identified as the primary sources of uncertainty compared

to other sectors, wildfire emissions constitute 1% to 9% of Portugal's total annual GHG

emissions.

Scarpa et al. (2024) employed a bottom-up approach to estimate GHG and particulate

matter emissions from rural and forest fires in Italy between 2007 and 2017. Their method

improved emission accuracy by integrating detailed data on burned areas, fuel types,

meteorological conditions and combustion phases, offering results that aligned well with

global and national inventories. According to this study, fire disturbance in broadleaf

forests, shrublands, and agricultural fuel types is the main source of vegetation fire

emissions in Italy, accounting for about 76 % of the total. They estimated average annual

emissions from open vegetation fire by 2346 Gg year^{-1} for CO_2 and 10.4 Gg year^{-1} for

CH_4.

2.2 Top-Down

The observation-informed "top-down" approach is a method in which atmospheric

observations are used to improve the emission estimates of the atmospheric species of

interest. This approach estimates emissions by adopting a framework that integrates

atmospheric observations, an atmospheric chemical transport model, and an inversion

algorithm. The approach is based on the relationship between atmospheric

concentrations or mole fractions with emissions, i.e. the so-called 'source-receptor

relationships' calculated by the chemical transport model. The top-down method uses a

"bottom-up" inventory as prior knowledge and optimises it to match the modelled

concentrations with observations. By leveraging the combined information from both

"bottom-up" estimates and observations, the inverse modelling approach significantly

reduces uncertainties in the posterior emission estimates (Rodgers, 2000). The accuracy

of these inverse modelling results and the spatial scale at which emission estimates are

obtained depend on the quality and spatial density of the measurements and the precision

of the transport model used in the analysis (Bergamaschi et al., 2018).

Modelled mixing ratios corresponding to the observations can be calculated as:

$$y^{mod} = Hx \tag{1}$$

where y^{mod} is a vector of the modelled mixing ratio, x is the unknown emission flux vector

and H is the source receptor relationship matrix. The emissions fluxes vector x includes

the variables to be optimised by the inversion, i.e. gridded emissions and other elements

such as background mixing ratio values.

Observed values can be expressed as:

$$y^{obs} = Hx + \varepsilon \qquad (2)$$

where y^{obs} is the observation vector and ε represents the observation error. However,

observations might not be sufficient to precisely constrain all components of the emission

vector. This situation leads to an ill-conditioned problem, requiring regularisation or the

introduction of supplementary information to obtain a meaningful solution. Typically, this

additional constraint is provided as a priori estimates of the unknowns, i.e. spatially

distributed information of the fluxes (x_b) , obtained from the bottom-up emission estimate.

Following Bayes' theorem, the most likely solution of the posterior emission flux (x)

minimises the difference between the observed and modelled mixing ratios. Assuming

that the observation and prior uncertainties follow Gaussian probability density functions,

this can be formulated as a cost function:

$$J(x) = \frac{1}{2}(x - x_b)^T B^{-1}(x - x_b) \; + \frac{1}{2}(Hx - y^{obs})^T R^{-1}(Hx - y^{obs}) \qquad (3)$$

In this equation, *B* represents the prior flux error covariance matrix, and *R* is the

observation error covariance matrix.

Several methods can be employed to find the optimum posterior state, defined as the

value of *x* for which eq. 3 has a minimum. Common approaches include the analytical

inversion (Stohl et al., 2009), Kalman filter (Dash et al., 2024), 3D-Var (Three-

Dimensional Variational Assimilation) and 4D-Var (Four-Dimensional Variational

Assimilation) (Meirink et al., 2006) optimisation techniques.

Inversion models have evolved rapidly in recent years, reducing uncertainties in emission
estimates. For a long time, the inversion methods were used to optimise the net emissions
of the species of interest. However, there have been advancements in the field to
accommodate the extraction of the sectorial contributions to the posterior total emissions.
In a commonly used and straightforward approach, net fluxes are partitioned to underlying
sector-based emissions by scaling fluxes based on the relative weight of sectors in a prior
inventory (Qu et al., 2021). This does not assume that the prior distribution of sectoral
emissions is correct, only that the relative allocation within a given grid cell is correct. A
similar approach accounts for emissions from different sectors having different prior error
standard deviations contributing differently to each sector (Shen et al., 2023). In another
approach by Zhang et al. (2021), the state vector is split into different emission sources,
such as wetland and non-wetland emissions. It allows the emission sector with higher
uncertainty to be treated differently than the well-known sector. In a similar approach by
Lu et al. (2021), the state vector is organised in three ways to obtain monthly, sectorial,
and provincial emissions according to the subnational boundaries for each year.
Assuming relative weights of different sectors in prior emissions imposes a correlation
between emission sectors that may not exist. Cusworth et al. (2021) discussed a method
that projects inverse CH_4 fluxes directly to emission sectors while accounting for
uncertainty structure and spatial resolution of prior fluxes and emissions. This method
can also be helpful in partitioning highly uncertain biospheric and anthropogenic
emissions to reconcile the global budget.

In more sophisticated approaches to get accurate sectorial estimates, the inversion
system is generally provided with some additional constraints. For instance, to constrain

CH$_4$ fluxes, additional trace gases can be employed to partition emissions. This method

becomes feasible when these trace gases are co-emitted from a specific source at a

characteristic ratio. For example, CO, the classical tracer for tracking smoke plumes from

fires, can be used to estimate methane emissions from activities such as biomass burning

(Heald et al., 2004). Similarly, ethane (C$_2$H$_6$) is emitted during fossil fuel production and

use, without significant emissions from biogenic sources used to quantify fossil fuel

emissions (Helmig et al., 2016; Peischl et al., 2013). Observations of isotopologues can

also be harnessed to apportion emissions through a similar approach. This method

involves evaluating the ratios of isotopologues emitted from different source types. Lan

et al. (2021) used CH$_4$ isotopologue measurements to better understand and differentiate

emissions from various sources, including biomass burning emissions. Analysing these

isotopic signatures provides valuable information on the sources and a more

comprehensive understanding of methane emissions. To our knowledge, for the

Mediterranean region, no "top-down" quantification based on inversion frameworks exists

for the emissions of climate-altering species from the open vegetation fires.

3. Quantification of open vegetation fire emissions over the Mediterranean basin:

a data review

We present a synthesis of the degree of agreement between different state-of-the-art

inventories estimating emissions of climate-altering species (CO$_2$, CH$_4$, N$_2$O and BC) from

open burning in the Mediterranean region. To this end, we compared emissions for the

whole area as well as for individual countries for four inventories (GFAS, GFED, FINN,

and EDGAR) from 2003 to 2020. Table 1 summarises key characteristics of the

considered inventories, including their acronyms, versions, producers, coverage details,

and resolutions. More specific details about each inventory are provided in the

supplementary material (Table SM1). Note that GFED, CAMS, and FINN include

anthropogenic and natural vegetation fires, whereas EDGAR only includes anthropogenic

emissions related to agricultural practices.

Table 1. Overview of fire emission inventories considered in the comparison.

Features	CAMS GFAS	GFED	FINN	EDGAR
Name	CAMS Global Fire Assimilation System	Global Fire Emissions Database Version 4.1	Fire INventory from NCAR	Emissions Database for Global Atmospheric Research Version 8.0
Acronym	GFAS	GFED	FINN	EDGAR
Version	1.2	4.1	2.5	8
Producer	Copernicus Atmosphere Monitoring Service	ORNL DAAC (Oak Ridge National Laboratory Distributed Active Archive Center)	National Center of Atmospheric Research (NCAR)	Joint Research Centre (JRC) in collaboration with the International Energy Agency
Spatial Coverage	Global	Global	Global	Global
Temporal Coverage	From 1 January 2003-present	June 1995 - present	2002-2023	Up to 2022
Spatial Resolution	0.1°x0.1°	0.25°x0.25°	0.1°x 0.1°	0.1°x0.1° and country level
Temporal Resolution	Daily/ Monthly/Yearly	Monthly	Daily	Annually

The Mediterranean region considered for aggregating emissions is shown in Figure 1. It

is based on HydroSHEDS (Hydrological data and maps based on Shuttle Elevation

Derivatives at multiple Scales) developed by the World Wildlife Fund (Lehner and Grill,

2013), a global dataset that provides detailed hydrographic information, like river

networks and watershed boundaries, in a consistent format for large scale applications.

The definition of the Mediterranean region is not unique and different definitions can be

adopted depending on the purposes and discipline (Giorgi, 2006; Lefèvre and Fady ,

2016; Meybeck et al., 2006; Schicker et al., 2010). In this case, we defined a domain

based on the extension of the catchment area of rivers flowing into the Mediterranean

Sea, including the lower part of the Nile catchment. This definition is a compromise

between other definitions and is consistent with Schicker et al. (2010).

Figure 1. The Mediterranean region domain (blue area in the map) selected to aggregate
emission fluxes from CAMS, GFED, FINN and EDGAR emission inventories.

The area of the Mediterranean that we have considered represents about 1.5% of the

global land surface. Using global CO_2 emissions from wildfires provided by Samborska et

al. (2024) as a reference, the percentage contribution of Mediterranean wildfires to global

wildfires emissions during the study period ranged from a minimum value of 0.18 - 0.32%

in 2012 to a maximum value of 0.24 - 0.45% in 2007, depending on the inventory

considered. It should be noted that these numbers refer to the domain as defined above

and may change depending on the spatial domain selected to represent the

Mediterranean region. For example, the relative values increase by including the whole

Iberian peninsula in the analysis as done by Gudmundsson et al. (2014) and Turco et al.

(2018).

Figure 2 reports the multi-annual variability of open fire emissions for CO_2, CH_4, N_2O, and

BC over the Mediterranean region from 2003 to 2020.

Figure 2. Annual emissions from open vegetation fires in the Mediterranean region (2003-2020):

CO_2 (a), CH_4 (b), N_2O (c) and BC (d). The different lines denote CAMS (red), GFED

(green), EDGAR (blue), and FINN (orange).

The GFED, CAMS, and FINN inventories are consistent in identifying temporal peaks in

the emissions, although there are differences in absolute values, with CAMS and FINN

often showing emission levels higher than GFED. EDGAR showed higher CO_2 emission

values than GFED for 10 out of 18 years. This is surprising, as EDGAR is expected to

represent a lower limit for emissions, as it only reports agriculture vegetation fires.

EDGAR does not capture the significant emission peaks observed in 2007, 2012 and

2017, probably because it only includes emissions related to agriculture and waste

burning.

In 2007, "extreme" fire danger conditions affected south-eastern Europe at the end of July

and August (Cesari et al., 2014). These conditions led to catastrophic events affecting

southern Italy and Greece, with record numbers of burnt areas. In 2007, the Rapid

Damage Assessment module of the European Forest Fires Information System reported

high burnt areas for Greece (271,516 ha) and Italy (153,753 ha) (European Commission

et al., 2008). The weather conditions that contributed to the wildfires included prolonged

heat waves with very high temperatures, prolonged droughts that preceded the events,

and strong winds that coincided with the ignition of the fire (Koutsias et al., 2012).

According to Schmuck (2013), in 2012 forest fires in the five southern EU countries

(Portugal, Spain, France, Italy, and Greece) burned a total area of 519,424 ha. This value

was well above the average for the previous 20 years (~ 400,000 ha) and among the

highest since 2000 (only 2003, 2005, and 2007 reported higher values of burnt areas, see

San-Miguel-Ayanz et al. (2022)). The winter and early spring were marked by a severe

drought in France, Spain and Italy with the period June to August hotter and drier than

normal (Schmuck, 2013).

In 2017, as reported by the European Drought Observatory, Italy experienced a severe

drought, followed by one of the driest springs in the last 60 years. During this period,

some regions received 80% less rainfall than normal. Two-thirds of Italy was affected by

these conditions (Joint Research Center, 2024) leading to a burnt area of 140,404 ha with

788 fires (Libertà et al., 2018). Interestingly, while 2007, 2012 and 2017 were years with

significant peaks in burned areas, the trend across these years shows a decrease in the

magnitude of these peaks. This decreasing trend is consistent with the behaviour

observed for emissions of key climate-altering species including CO_2, CH_4, N_2O and BC.

Figure 3 shows the percentage of contributions of different countries to the CO_2 emissions

from the Mediterranean region over the period 2003 - 2020. CO_2 has been presented as

a 'representative' case because it is the dominant GHG emitted by open burning and is a

major contributor to global warming and climate change. Results for CH_4, N_2O and BC

can be found in the supplementary material (Fig. SM2). It should be noted that EDGAR

was not included in this figure, as it only provides emission estimates for agricultural waste

burning and does not account for emissions from other large fires. After extracting the

Mediterranean region from each global inventory, we segregated national emissions

falling in the domain defined by Figure 1.

Figure 3. Annual CO$_2$ emissions-contribution for the top 5 emitting countries for CAMS/GFAS v1.2 (left column), GFED v4.1s(middle column) and FINN v2.5 (right column) for 2003 - 2020 in the Mediterranean region (Albania: ALB, Algeria: ALG; Egypt: EGY; France: FRA; Greece: GRE; Italy: ITA; Spain: ESP, Turkey: TUR; other countries: OTH).

An interesting question is the degree of agreement between the different inventories when

looking at individual countries included in the Mediterranean area. Here we discussed the

case for CO_2 emissions (Figure 3) as a representative example. Over the whole

investigation period, CAMS identified Algeria as the most important contributor for CO_2,

while both GFED and FINN indicated Italy. However, differences were found for the

different years. As an example, in 2012 which is characterised by the occurrence of a

peak in emissions, all inventories indicated Italy as a prominent contributor, but with

varying proportions: CAMS reported 24%, GFED 33% and FINN 39% of the total

emissions. A notable difference is that CAMS reported Spain as one of the top five

Mediterranean contributors for each year, while GFED reported Egypt and FINN reported

Albania. These discrepancies highlight the challenges of reconciling data from different

inventories and the need for further scrutiny of the methodologies used by each inventory

especially when looking at smaller spatial scales. It should be emphasised that the aim

of this country-level analysis is not to rank the contributors to emissions from open burning

of vegetation (this should take into account the different areas of the countries), but to

highlight the discrepancies in the estimates provided by the inventories here reviewed.

As suggested by Magi et al. (2012), through agricultural and waste burning, human

practices add further complexity to occurrence of vegetation fires, with human burning

practices in agriculture not necessarily following the temporal development of wildfires.

As EDGAR provides emission estimates for agricultural field burning and not those from

other large fires (e.g. biomass burning from Savannah and forests, see Petrescu et al.

(2020)), it will be used in the following to analyse the emissions of CO_2, CH_4, and N_2O

from the agriculture sector in the period 2003 - 2020 (Figure 4).

Figure 4. Time series of agriculture and waste burning emissions reported by EDGAR v.8 for different countries.

Table 2 illustrates the trends in emissions of CO_2, CH_4 and N_2O over the period 2003 to 2020 using EDGAR inventory. The trends are estimated using the Theil-Sen slope

estimator (Wilcox, 2010) and are presented for individual top 5 emission contributing

countries (France, Greece, Italy, Turkey, Egypt) and as an aggregated total for the

Mediterranean domain. The results show statistically significant downward tendencies for

most countries, with Turkey showing the largest decreases. The aggregated total

emissions reveal a reduction in CO_2 (-0.09 Tg year^{-1}), CH_4 (-0.16 Gg year^{-1}) and N_2O (-

4.2 Mg year^{-1}) emissions.

This reduction of the emissions from the European countries in the domain is consistent

with the implementation of directives defined to control agricultural and waste burning, as

highlighted by the United Nations Economic Commission for Europe and supported by

the European Union's sustainable waste management policies (United Nations

Environment Programme, 2024).

Table 2. Theil-Sen trend analysis of anthropogenic emissions (2003 - 2020) from the EDGAR inventory for Egypt ("EGY"), France ("FRA"), Greece ("GRC"), Italy ("ITA"), Turkey ("TUK"), and the whole study domain ("MED"). Asterisks (*, **, ***) represent statistical significance in confidence levels at $p<0.05$, $p<0.01$ and $p<0.001$, respectively. The 95% confidence intervals in the slopes are reported within brackets. Emissions for countries refer to the part of the national territory included in the study area shown in Figure 1.

Spatial region	CH_4 Trend (Gg yr^{-1})	CO_2 Trend (Tg yr^{-1})	N_2O Trend (Mg yr^{-1})
EGY	1.4×10^{-3} [-2.1×10^{-3}, 4.6×10^{-3}]	7.6×10^{-4} [-1.2×10^{-3}, 2.6×10^{-3}]	-0.0 [-0.1, 0.1]
FRA	-0.01 [-0.02, -0.01]***	-0.01 [-0.01, 0]***	-0.3 [-0.4, -0.2]***
GRC	- 0.03 [-0.05, -0.01]**	-0.02 [-0.03, -0.01]**	-0.8 [-1.2, -0.3]**
ITA	-0.04 [-0.07, -0.01]*	-0.02 [-0.04, 0]*	-1.0 [-1.9, -0.2]*
TUK	-0.05[-0.06, -0.03]***	-0.03 [-0.04, -0.02]***	-1.2 [-1.6, -0.8]***
MED	-0.16 [-0.21, -0.13]***	-0.09 [-0.21, -0.07]***	-4.2 [-5.4, -3.3]***

4. Discussion on emission estimates

Vegetation fires are a complex process associated with factors of different origins and occurring on different spatial and temporal scales, such as vegetation and landscape conditions, weather conditions, and human activities (Turco et al., 2013). It is thus difficult to quantify and univocally attribute the relative importance of these different processes to fire occurrences and related emissions. The data comparison reported in Section 3 showed that all three considered inventories that include wildfire contributions (i.e. GFED, GFAS and FINN) showed temporally consistent emission peaks for the years 2007, 2012 and 2017. Scarpa et al. (2024) identified these years as the most significant over the

period 2007 - 2017 in terms of GHG and particulate aerosol emissions from vegetation fires over Italy. Previous studies suggested that climatic processes, through their impact on fuel moisture, as well as on nature and availability of fuel, can represent a driving process for the inter-annual variability of regional fire patterns, especially in the Mediterranean region (Meyn et al., 2007). In particular, a ~5-year recurring maxima in the burned areas was observed by Turco et al. (2019) for the northeastern Iberian Peninsula related to summer precipitation and maximum temperature as well as antecedent climate conditions with a time lag of 1-2 years. From this perspective, it is interesting to note that all the years that are characterised by a peak in open vegetation fire emissions over the Mediterranean basin are featured by the occurrence of La Niña events. La Niña represents the "cold" phase of the El Niño Southern Oscillation (ENSO), a quasi-periodic (two to seven years) climate pattern affecting the equatorial Pacific circulation (Burton et al., 2020). According to NOAA, the magnitude of ENSO events is classified into categories such as "weak", "moderate", "strong", or "very strong". Based on this classification, the 2007 - 2008 event was categorized as "strong" ENSO, the 2011 - 2012 event as "moderate", and the 2017 - 2018 event as "weak" (Climate Prediction Center, 2024). Particularly cold phases of ENSO (i.e. La Niña) have occurred in May 2007 - May 2008, June 2011- May 2012 and June - December 2016 (Climate Prediction Center, 2024), i.e. at the same time as or in advance (with a lag time up to 1 year) of the observed peak fire seasons (Figure SM3 in the supplementary material).

ENSO is the primary cause of fluctuations in the terrestrial carbon cycle, particularly in the tropics (Cox et al., 2013): previous studies (Le Page et al., 2008; Zhai et al., 2016)

suggested a significant global influence of ENSO on the carbon budget by increased fire

activity and decreased terrestrial carbon sink capacity.

For the Asia-Pacific region, ENSO has been an important climate driver, but its influence

over North Atlantic Europe is quite challenging to establish (Ashok et al., 2007;

Ayarzagüena et al., 2018), despite previous research suggesting a robust climate

influence over Europe (Brönnimann, 2007).

The South Asian summer monsoon and ENSO are closely related through atmospheric

teleconnections. ENSO modulates the strength and intensity of monsoons with la Niña

events leading to stronger monsoons. This variability in the monsoon, in return, impacts

the Mediterranean climate through the monsoon-desert mechanism, affecting subsidence

and precipitation patterns in the region. In the years with a stronger monsoon, there is a

noticeable increase in the vertical advection of moist static energy and horizontal

advection of dry enthalpy over the Mediterranean region, leading to drier conditions

(Cherchi et al., 2014). Shaman (2014), showed decreased precipitation during La Niña

events over southern Mediterranean Europe in June - September and October -

December.

Thus, despite great caution should be exercised when discussing possible climatic

processes over the relatively short study period considered in this study, we argue that

ENSO may have played a role in determining part of the observed inter-annual variability

of GHG and BC emissions associated with open vegetation fires in the Mediterranean

region. With the aim of providing, also in the context of this review, indications to better

disentangle the possible relationship between ENSO and regional climate variability with

wildfire emissions, we calculated the anomalies of surface air temperature (2m

temperature) and total precipitation for the Mediterranean area. For this analysis we

utilised data from the ECMWF fifth generation reanalysis, i.e. ERA5 (Hersbach et al.,

2020), with a spatial resolution of 0.25° x 0.25°. The mean of ensemble members from

the dataset "ERA5 monthly averaged data on single levels from 1940 to present"

(Hersbach et al., 2023) provided by the Copernicus Climate Change Service was used.

Anomalies were calculated for the season May - August with respect to the means in the

referenced period. Years 2012 and 2017 were characterised by evident positive and

negative anomalies for temperature and precipitation, respectively (Fig. 5). No clear

anomaly signals were observed for 2007. These results did not change dramatically

according to the length of the seasons analysed (see Fig. SM4, in the Supplementary

Material). It should be noted that the 2007 fire season was mainly influenced by specific

large events that occurred over Italy and Greece in July and August (Cesari et al., 2014;

European Commission et al., 2008; Koutsias et al., 2012). It is therefore possible that our

Mediterranean-scale anomaly may not fully represent the weather variability occurring at

the sub-domain and sub-seasonal scales. Furthermore, it should be considered that, as

suggested by Turco et al. (2013), several other factors besides current weather conditions

may influence the occurrence of wildfires in the Mediterranean, including weather

conditions preceding (up to 2 years before) the actual fire season, as well as human

management, including fire suppression activities, which may affect fuel availability and

type/conditions. Finally, Zhang et al. (2019), pointed to the contrasting responses of La

Niña events in influencing the extratropical atmospheric circulation, adding another layer

of complexity in establishing a cause-and-effect relationship.

Figure 5. Surface temperature and precipitation anomalies from May to August over the period 2003 - 2020 for the Mediterranean domain as calculated from the ERA5 dataset.

It should be noted that our integrated basin perspective and the annual temporal resolution adopted to compare the emission inventory in this review paper, prevented us from considering specific emission variability that can occur among different source regions and on sub-seasonal time scales (Gudmundsson et al., 2014).

Despite their agreement on the diagnosis of peak years in emissions, there were notable differences in the quantification of total GHG and BC emissions in the Mediterranean for GFED, CAMS, and FINN. On average, FINN provides the highest emission estimates for all species considered, while GFED provides the lowest, even if it includes emissions from small fires. For CO_2 (CH_4), more consistent emission estimates are provided by FINN

(GFED) and CAMS. The differences in GHG and BC emission among GFED, CAMS

GFAS, and FINN might originate from variations in methodologies and input data sources.

GFED estimates burned area and emissions using satellite imagery, with a spatial

resolution of 0.25° x 0.25°, covering a range of fire types and vegetation. CAMS relies on

fire radiative power data from MODIS instruments to estimate emissions at a finer

resolution of 0.1° x 0,1°. FINN provides emissions estimates with high spatial resolution

(1 km^2), integrating fire detections from MODIS and Visible Infrared Imaging Radiometer

Suite (VIIRS). Thus, it can be argued that the higher spatial resolution of satellite data

used to obtain the burned area could partially explain the higher values of emissions.

However, it should be considered that each dataset applies different methods to calculate

the fuel amount as well as emission factors tailored to vegetation types and fire

characteristics. In particular, the differences in the emission factors can explain the

different relative agreement between the reviewed inventories when different species are

considered (e.g. CO_2 vs CH_4).

To assess the spread of the four emission inventories, we calculated the mean standard

deviation (σ) and the mean normalized standard deviation (coefficient of variation, v) for

the study period for the top three emitting countries and the entire study domain (Table

3). The σ parameter provides indication about the temporal variability of emissions during

the investigation period. The coefficient of variation (v) can be used as a measure of the

emission spread across the different inventories for the species.

Table 3. Average standard deviation (σ) and average normalised standard deviation (coefficient of variation, v) for Algeria ("ALG"), Greece ("GRC"), Italy ("ITA"), and the whole study domain ("MED") for different atmospheric species. σ is in Tg for CO_2 and in Gg for all other species.

Species	ALG		GRC		ITA		MED	
	σ	v	σ	V	σ	v	σ	v
CO_2	1.4	0.91	1.21	0.45	1.04	0.37	3.63	0.27
CH_4	1.9	0.76	2.3	0.54	3.34	0.45	8.72	0.34
N_2O	0.21	1.04	0.18	0.55	0.06	0.23	0.57	0.56
BC	0.36	0.6	0.27	0.41	0.4	0.39	0.68	0.19

Emissions for Algeria are highly variable across all species. Looking to the whole Mediterranean domain, BC emissions are the most consistent among the considered inventories, while N_2O emissions show the highest variability. Our analysis is consistent with recent scientific discussions highlighting the need to align emission inventories to achieve greater accuracy (Cowie et al., 2012; Yona et al., 2020). It is noteworthy that, based on the literature review presented in this work, no emission estimates from atmospheric inversion modelling exist so far for open vegetation fire emissions in the Mediterranean region. Working on the application of this methodology would represent a valuable contribution towards a better quantification (and reduction) of the uncertainties associated with the inventories used to constraint emission estimates in this work.

5. Conclusions

This work reviewed the current methodologies for quantifying climate-altering emissions

from open vegetation fires in the Mediterranean region, focusing on CO_2, CH_4, N_2O, and

BC. Based on our scientific literature review, limited specific efforts have been made to

investigate the role of open vegetation fires as a source of climate-altering species in the

Mediterranean. The few existing studies were based on a bottom-up approach, while

there are no results based on a top-down, observational methodology.

Our review revealed significant deviations in emission estimates from four widely used

inventories (GFED v4.1s, GFAS v1.2, FINN v2.5, and EDGAR v8.0), underlying the

constraints and limitations on the accounting of climate-altering emissions for this sector.

The differences in input data, fire detection, spatial/temporal resolution, and adopted

emission factors contribute to the variability in emission quantification between the

inventories. FINN v2.5, which uses the satellite data with the higher spatial resolution, is

the inventory that provides the highest emission estimates, while the relative differences

between the different inventories vary as a function of the species considered, suggesting

a role for the different emission factors used. Assuming that the variability among

inventories can be considered as a measure of underlying uncertainties in the total

emissions, the differences between the inventories suggested a need for further efforts in

improving estimates. This is also important when considering anthropogenic fire

emissions from the agricultural sector, in particular to determine the relative role of each

country's share in the Mediterranean region.

Inventories like GFED v4.1s, GFAS v1.2 and FINN v2.5 provide high-resolution temporal

data, making them valuable for numerical modeling and regional scale assessments.

Moreover, with respect to EDGAR (focused on agriculture and waste burning), they

provide more comprehensive data about fire emissions, including wildfires. However, their

coarse spatial resolution limits their applicability for specific applications like, e.g. city-

level modeling. Integrating these inventories with localised datasets, such as high-

resolution land-use maps and urban fire observations, is crucial to enhance their utility for

city-scale emissions modeling and air quality assessments. Despite these challenges, fire

inventories remain indispensable for understanding the atmospheric and climatic impacts

of fire emissions.

We hypothesised and discussed the possibility that the inter-annual variability in the

emissions can be related to global climate processes, like ENSO. In particular, a possible

relationship between the occurrence of La Niña events and the detected emission peaks

was discussed. However, the rather short period of time considered in this work, the

multiple factors, other than weather conditions that influence the occurrence of forest fires

in the Mediterranean, and the complex responses of the extratropical atmosphere to La

Niña, require further and more specific studies that go well beyond the objective of this

review.

This study highlights the importance of upgrading GHG emission inventories to better

reflect the impact of open fire emissions in the Mediterranean region. It highlights

significant differences in the reported emissions and underlines the importance of using

standardised, transparent, and verifiable methodologies for GHG accounting. The

differences between the inventories, especially when down-scaling to the portion of single

countries included in the investigated domain suggested further efforts in reducing

underlying uncertainties.

In general, we would recommend that future studies should focus on developing consistent methodologies for quantifying and reporting emissions, integrating satellite and ground-based measurement data through atmospheric inversion modelling as already done for other emission sectors. Policy makers should use improved inventories to develop targeted, evidence-based fire prevention and climate mitigation strategies, particularly for the Mediterranean region. This study highlights the need for accurate emissions data and its integration into policy frameworks to support effective climate change mitigation at regional and local levels.

Acknowledgments

We would like to thank the teams behind the GFED v4.1s, GFAS v1.2, FINN v2.5, and EDGAR v8.0 inventories for providing the crucial data that significantly contributed to our research. ERA5 data were provided by ECMWF through Copernicus Climate Change Service (C3S). We gratefully thank Annalisa Cherchi (CNR - ISAC) for the useful discussion about the role of ENSO in the Mediterranean region. We thank Riccardo Testolin (ISAFOM-CNR and NBFC, Italy) for providing the Mediterranean mask. We acknowledge the support from the University School for Advanced Studies IUSS Pavia ("Scuola Universitaria Superiore IUSS Pavia") and the Institute of Atmospheric Sciences and Climate (ISAC), National Research Council (CNR), Italy. We also acknowledge the project funded by the European Union – NextGenerationEU under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of December 16, 2021, rectified by Decree n.3175 of December 18, 2021 of the Italian Ministry of University and Research under award Number: Project code CN_00000033, Concession Decree No. 1034 of June 17, 2022 adopted by the Italian

Ministry of University and Research, CUP B83C22002930006, Project title "National

Biodiversity Future Centre - NBFC". During this research work, Rabia Ali Hundal and

Saurabh Annadate were supported by the Italian national inter-university PhD course in

Sustainable Development and Climate change (PhD SDC, link: www.phd-sdc.it).

Declaration of AI-assisted technologies in the writing process

During the preparation of this work the authors used DeepL (https://www.deepl.com/) in

order to check the English language. After using this tool, the authors reviewed and edited

the content as needed and take full responsibility for the content of the publication.

References

Anav, A., Sorrentino, B., Collalti, A., Paoletti, E., Sicard, P., Coulibaly, F., Manzini, J., Hoshika, Y., De Marco, A., 2024. Meteorological, chemical and biological evaluation of the coupled chemistry-climate WRF-Chem model from regional to urban scale. An impact-oriented application for human health. Environmental Research 257, 119401. https://doi.org/10.1016/j.envres.2024.119401

Ashok, K., Behera, S.K., Rao, S.A., Weng, H., Yamagata, T., 2007. El Niño Modoki and its possible teleconnection. Journal of Geophysical Research: Oceans 112. https://doi.org/10.1029/2006JC003798

Ayarzagüena, B., Ineson, S., Dunstone, N.J., Baldwin, M.P., Scaife, A.A., 2018. Intraseasonal Effects of El Niño–Southern Oscillation on North Atlantic Climate. Journal of Climate 31, 8861–8873. https://doi.org/10.1175/JCLI-D-18-0097.1

Baudena, M., Santana, V., Baeza, M., Bautista, S., Eppinga, M., Hemerik, L., Mayor, Á., Rodríguez-Mateos, F., Valdecantos, A., Vallejo, R., Vasques, A., Rietkerk, M., 2020. Increased aridity drives post-fire recovery of Mediterranean forests towards open shrublands. New Phytologist 225, 1500–1515. https://doi.org/10.1111/nph.16252

Bergamaschi, P., Danila, A., Weiss, R., Ciais, P., Thompson, R., Brunner, D., Levin, I., Meijer, Y., Chevallier, F., Janssens-Maenhout, G., Bovensmann, H., Crisp, D., Basu, S., Dlugokencky, E., Engelen, R., Gerbig, C., Günther, D., Hammer, S., Henne, S., Krol, M., 2018. Atmospheric monitoring and inverse modelling for verification of greenhouse gas inventories. https://doi.org/10.2760/759928

Bond, T.C., Doherty, S.J., Fahey, D.W., Forster, P.M., Berntsen, T., DeAngelo, B.J., Flanner, M.G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P.K., Sarofim, M.C., Schultz, M.G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S.K., Hopke, P.K., Jacobson, M.Z., Kaiser, J.W., Klimont, Z., Lohmann, U., Schwarz, J.P., Shindell, D., Storelvmo, T., Warren, S.G.,

Zender, C.S., 2013. Bounding the role of black carbon in the climate system: A scientific assessment. Journal of Geophysical Research: Atmospheres 118, 5380–5552. https://doi.org/10.1002/jgrd.50171

Bowman, D.M.J.S., Johnston, F.H., 2005. Wildfire Smoke, Fire Management, and Human Health. EcoHealth 2, 76–80. https://doi.org/10.1007/s10393-004-0149-8

Brönnimann, S., 2007. Impact of El Niño–Southern Oscillation on European climate. Reviews of Geophysics 45. https://doi.org/10.1029/2006RG000199

Burton, C., Betts, R.A., Jones, C.D., Feldpausch, T.R., Cardoso, M., Anderson, L.O., 2020. El Niño Driven Changes in Global Fire 2015/16. Front. Earth Sci. 8. https://doi.org/10.3389/feart.2020.00199

Cesari, R., D', M., Isidoro, N.A., Maurizi, A., Mircea, M., Monti, F., Pizzigalli, C., Tampieri, F., 2014. Modelling dispersion of smoke from wildfires in a Mediterranean area. IJEP 55, 219. https://doi.org/10.1504/IJEP.2014.065927

Cherchi, A., Annamalai, H., Masina, S., Navarra, A., 2014. South Asian Summer Monsoon and the Eastern Mediterranean Climate: The Monsoon–Desert Mechanism in CMIP5 Simulations. Journal of Climate 27, 6877–6903. https://doi.org/10.1175/JCLI-D-13-00530.1

Climate Prediction Center, 2024. El Niño / Southern Oscillation (ENSO). https://origin.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/enso.shtml (accessed 14 March 2024).

Cowie, A., Eckard, R., Eady, S., 2012. Greenhouse gas accounting for inventory, emissions trading and life cycle assessment in the land-based sector: a review. Crop Pasture Sci. 63, 284–296. https://doi.org/10.1071/CP11188

Cox, P.M., Pearson, D., Booth, B.B., Friedlingstein, P., Huntingford, C., Jones, C.D., Luke, C.M., 2013. Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability. Nature 494, 341–344. https://doi.org/10.1038/nature11882

Crippa, M., Guizzardi, D., Pagani, F., Banja, M., Muntean, M., Schaaf, E., Becker, W.E., Monforti-Ferrario, F., Quadrelli, R., Risquez, M.A., Taghavi-Moharamli, P., Köykkä, J., Grassi, G., Rossi, S., Melo, J., Oom, D., Branco, A., San-Miguel, J., Vignati, E., 2023. GHG emissions of all world countries [WWW Document]. JRC Publications Repository. https://doi.org/10.2760/953322

Crutzen, P.J., Andreae, M.O., 1990. Biomass Burning in the Tropics: Impact on Atmospheric Chemistry and Biogeochemical Cycles. Science 250, 1669–1678. https://doi.org/10.1126/science.250.4988.1669

Cusworth, D., Bloom, A., Ma, S., Miller, C., Bowman, K., Yin, Y., Maasakkers, J., Zhang, Y., Scarpelli, T., Qu, Z., Jacob, D., Worden, J., 2021. A Bayesian framework for deriving sector-based methane emissions from top-down fluxes. Communications Earth & Environment 2. https://doi.org/10.1038/s43247-021-00312-6

Dash, S., DeRosa, M.L., Dikpati, M., Sun, X., Mahajan, S.S., Liu, Y., Hoeksema, J.T., 2024. Ensemble Kalman Filter Data Assimilation Into Surface Flux Transport Model To Infer Surface Flows: An Observing System Simulation Experiment [WWW Document]. arXiv.org. URL https://arxiv.org/abs/2409.15233v1 (accessed 10.7.24).

De Marco, A., Sicard, P., Feng, Z., Agathokleous, E., Alonso, R., Araminiene, V., Augustatis, A., Badea, O., Beasley, J.C., Branquinho, C., Bruckman, V.J., Collalti, A., David-Schwartz, R., Domingos, M., Du, E., Garcia Gomez, H., Hashimoto, S., Hoshika, Y., Jakovljevic, T., McNulty, S., Oksanen, E., Omidi Khaniabadi, Y., Prescher, A.-K., Saitanis, C.J., Sase, H., Schmitz, A., Voigt, G., Watanabe, M., Wood, M.D., Kozlov, M.V., Paoletti, E., 2022. Strategic roadmap to assess forest vulnerability under air pollution and climate change. Global Change Biology 28, 5062–5085. https://doi.org/10.1111/gcb.16278

Joint Research Center, 2024. EDO Home - European Drought Observatory - JRC European Commission. https://edo.jrc.ec.europa.eu/edov2/php/index.php?id=1000 (accessed 26 March 2024).

European Commission, Institute for Environment and Sustainability (Joint Research Centre), Camia, A., San-Miguel-Ayanz, J., Kucera, J., 2008. Forest fires in Europe 2007. Publications Office of the European Union, LU. https://data.europa.eu/doi/10.2788/76413

Fernandes, P.M., Barros, A.M.G., Pinto, A., Santos, J.A., 2016. Characteristics and controls of extremely large wildfires in the western Mediterranean Basin. Journal of Geophysical Research: Biogeosciences 121, 2141–2157. https://doi.org/10.1002/2016JG003389

Giorgi, F., 2006. Climate change hot-spots. Geophysical Research Letters 33. https://doi.org/10.1029/2006GL025734

Gudmundsson, L., Rego, F.C., Rocha, M., Seneviratne, S.I., 2014. Predicting above normal wildfire activity in southern Europe as a function of meteorological drought. Environ. Res. Lett. 9, 084008. https://doi.org/10.1088/1748-9326/9/8/084008

Schmuck, G., 2013. Forest fires in Europe, Middle East and North Africa 2012. Publications Office, LU. https://data.europa.eu/doi/10.2788/58397

Hamilton, D.S., Hantson, S., Scott, C.E., Kaplan, J.O., Pringle, K.J., Nieradzik, L.P., Rap, A., Folberth, G.A., Spracklen, D.V., Carslaw, K.S., 2018. Reassessment of pre-industrial fire emissions strongly affects anthropogenic aerosol forcing. Nat Commun 9, 3182. https://doi.org/10.1038/s41467-018-05592-9

Heald, C.L., Jacob, D.J., Jones, D.B.A., Palmer, P.I., Logan, J.A., Streets, D.G., Sachse, G.W., Gille, J.C., Hoffman, R.N., Nehrkorn, T., 2004. Comparative inverse analysis of satellite (MOPITT) and aircraft (TRACE-P) observations to estimate Asian sources of carbon monoxide. Journal of Geophysical Research: Atmospheres 109. https://doi.org/10.1029/2004JD005185

Helmig, D., Rossabi, S., Hueber, J., Tans, P., Montzka, S.A., Masarie, K., Thoning, K., Plass-Duelmer, C., Claude, A., Carpenter, L.J., Lewis, A.C., Punjabi, S., Reimann, S., Vollmer, M.K., Steinbrecher, R., Hannigan, J.W., Emmons, L.K., Mahieu, E., Franco, B., Smale, D., Pozzer, A., 2016. Reversal of global atmospheric ethane and propane trends largely due to US oil and natural gas production. Nature Geosci 9, 490–495. https://doi.org/10.1038/ngeo2721

Henderson, M., Kalabokidis, K., Marmaras, E., Konstantinidis, P., Marangudakis, M., 2005. Fire and Society: A Comparative Analysis of Wildfire in Greece and the United States. Human Ecology Review 12, 169–182.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R.J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., Thépaut, J.-N., 2020 . The ERA5 global reanalysis. Quarterly Journal of the Royal Meteorological Society 146, 1999–2049. https://doi.org/10.1002/qj.3803

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J-N., 2023. ERA5 monthly averaged data on single levels from 1940 to present [dataset]. Copernicus Climate Change Service (C3S) Climate Data Store (CDS). https://doi.org/10.24381/cds.f17050d7

Intergovernmental Panel On Climate Change, 2023. Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, 1st ed. Cambridge University Press. https://doi.org/10.1017/9781009157896

Kang, S., Zhang, Q., Qian, Y., Ji, Z., Li, C., Cong, Z., Zhang, Y., Guo, J., Du, W., Huang, J., You, Q., Panday, A.K., Rupakheti, M., Chen, D., Gustafsson, Ö., Thiemens, M.H., Qin, D., 2019. Linking atmospheric pollution to cryospheric change in the Third Pole region: current progress and future prospects. National Science Review 6, 796–809. https://doi.org/10.1093/nsr/nwz031

Kang, S., Zhang, Y., Qian, Y., Wang, H., 2020. A review of black carbon in snow and ice and its impact on the cryosphere. Earth-Science Reviews 210, 103346. https://doi.org/10.1016/j.earscirev.2020.103346

Kasischke, E.S., Penner, J.E., 2004. Improving global estimates of atmospheric emissions from biomass burning. Journal of Geophysical Research: Atmospheres 109. https://doi.org/10.1029/2004JD004972

Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., Schöpp, W., 2017. Global anthropogenic emissions of particulate matter including black carbon. Atmospheric Chemistry and Physics 17, 8681–8723. https://doi.org/10.5194/acp-17-8681-2017

Koutsias, N., Arianoutsou, M., Kallimanis, A.S., Mallinis, G., Halley, J.M., Dimopoulos, P., 2012. Where did the fires burn in Peloponnisos, Greece the summer of 2007? Evidence for a synergy of fuel and weather. Agricultural and Forest Meteorology 156, 41–53. https://doi.org/10.1016/j.agrformet.2011.12.006

Lan, X., Basu, S., Schwietzke, S., Bruhwiler, L.M.P., Dlugokencky, E.J., Michel, S.E., Sherwood, O.A., Tans, P.P., Thoning, K., Etiope, G., Zhuang, Q., Liu, L., Oh, Y., Miller, J.B., Pétron, G., Vaughn, B.H., Crippa, M., 2021. Improved Constraints on Global Methane Emissions and Sinks Using δ13C-CH4. Global Biogeochemical Cycles 35, e2021GB007000. https://doi.org/10.1029/2021GB007000

Landry, J.-S., 2016. Non-deforestation fire vs. fossil fuel combustion: the source of CO_2 emissions affects the global carbon cycle and climate responses. Biogeosciences 13, 2137–2149. https://doi.org/10.5194/bg-13-2137-2016

Le Page, Y., Pereira, J.M.C., Trigo, R., da Camara, C., Oom, D., Mota, B., 2008. Global fire activity patterns (1996–2006) and climatic influence: an analysis using the World Fire Atlas. Atmospheric Chemistry and Physics 8, 1911–1924. https://doi.org/10.5194/acp-8-1911-2008

Lefèvre, F., Fady, B., 2016. Introduction to Mediterranean Forest Systems: Mediterranean Basin, in: Paine, T.D., Lieutier, F. (Eds.), Insects and Diseases of Mediterranean Forest Systems. Springer International Publishing, Cham, pp. 7–28. https://doi.org/10.1007/978-3-319-24744-1_2

Lehner, B., Grill, G., 2013. Global river hydrography and network routing: baseline data and new approaches to study the world's large river systems. Hydrological Processes 27, 2171–2186. https://doi.org/10.1002/hyp.9740

Libertà, G., Vivancos, T.A., Leray, T., Costa, H., San-Miguel-Ayanz, J., Branco, A., Durrant, T., Lana, F., Nuijten, D., Ahlgren, A.C., Löffler, P., Ferrari, D., De Rigo, D., Boca, R., Maianti, P., 2018. Forest fires in Europe, Middle East and North Africa 2017. Publications Office of the European Union. https://data.europa.eu/doi/10.2760/663443

Lionello, P., Scarascia, L., 2018. The relation between climate change in the Mediterranean region and global warming. Reg Environ Change 18, 1481–1493. https://doi.org/10.1007/s10113-018-1290-1

Lu, X., Jacob, D.J., Wang, H., Maasakkers, J.D., Zhang, Y., Scarpelli, T.R., Shen, L., Qu, Z., Sulprizio, M.P., Nesser, H., Bloom, A.A., Ma, S., Worden, J.R., Fan, S., Parker, R.J., Boesch, H., Gautam, R., Gordon, D., Moran, M.D., Reuland, F., Villasana, C.A.O., 2021. Methane emissions in the United States, Canada, and Mexico: Evaluation of national methane emission inventories and sectoral trends by inverse analysis of in situ (GLOBALVIEWplus CH₄ ObsPack) and satellite (GOSAT) atmospheric observations. https://doi.org/10.5194/acp-2021-671

Magi, B.I., Rabin, S., Shevliakova, E., Pacala, S., 2012. Separating agricultural and non-agricultural fire seasonality at regional scales. Biogeosciences 9, 3003–3012. https://doi.org/10.5194/bg-9-3003-2012

Meirink, J.F., Eskes, H.J., Goede, A.P.H., 2006. Sensitivity analysis of methane emissions derived from SCIAMACHY observations through inverse modelling. Atmospheric Chemistry and Physics 6, 1275–1292. https://doi.org/10.5194/acp-6-1275-2006

Meybeck, M., Dürr, H., Vörösmarty, C., 2006. Global coastal segmentation and its river catchment contributors: A new look at land-ocean linkage. Global Biogeochemical Cycles - GLOBAL BIOGEOCHEM CYCLE 20. https://doi.org/10.1029/2005GB002540

Meyn, A., White, P.S., Buhk, C., Jentsch, A., 2007. Environmental drivers of large, infrequent wildfires: the emerging conceptual model. Progress in Physical Geography: Earth and Environment 31, 287–312. https://doi.org/10.1177/0309133307079365

Moreira, F., Viedma, O., Arianoutsou, M., Curt, T., Koutsias, N., Rigolot, E., Barbati, A., Corona, P., Vaz, P., Xanthopoulos, G., Mouillot, F., Bilgili, E., 2011. Landscape – wildfire interactions in southern Europe: Implications for landscape management. Journal of Environmental Management 92, 2389–2402. https://doi.org/10.1016/j.jenvman.2011.06.028

Noce, S., Collalti, A., Valentini, R., Santini, M., 2016. Hot spot maps of forest presence in the Mediterranean basin. iForest - Biogeosciences and Forestry 9, 766. https://doi.org/10.3832/ifor1802-009

Pande, C.B., Moharir, K.N., Kumar Singh, S., Varade, A.M., Elbeltagi, A., Khadri, S.F.R., Choudhari, P., 2021. Estimation of crop and forest biomass resources in a semi-arid region using satellite data and GIS. Journal of the Saudi Society of Agricultural Sciences 20, 302–311. https://doi.org/10.1016/j.jssas.2021.03.002

Peischl, J., Ryerson, T.B., Brioude, J., Aikin, K.C., Andrews, A.E., Atlas, E., Blake, D., Daube, B.C., de Gouw, J.A., Dlugokencky, E., Frost, G.J., Gentner, D.R., Gilman, J.B., Goldstein, A.H., Harley, R.A., Holloway, J.S., Kofler, J., Kuster, W.C., Lang, P.M., Novelli, P.C., Santoni, G.W., Trainer, M., Wofsy, S.C., Parrish, D.D., 2013. Quantifying sources of methane using light alkanes in the Los Angeles basin, California. Journal of Geophysical Research: Atmospheres 118, 4974–4990. https://doi.org/10.1002/jgrd.50413

Petrescu, A.M.R., Peters, G.P., Janssens-Maenhout, G., Ciais, P., Tubiello, F.N., Grassi, G., Nabuurs, G.-J., Leip, A., Carmona-Garcia, G., Winiwarter, W., Höglund-Isaksson, L., Günther, D., Solazzo, E., Kiesow, A., Bastos, A., Pongratz, J., Nabel, J.E.M.S., Conchedda, G., Pilli, R., Andrew, R.M., Schelhaas, M.-J., Dolman, A.J., 2020. European anthropogenic AFOLU greenhouse gas emissions: a review and benchmark data. Earth System Science Data 12, 961–1001. https://doi.org/10.5194/essd-12-961-2020

Qu, Z., Jacob, D.J., Shen, L., Lu, X., Zhang, Y., Scarpelli, T.R., Nesser, H., Sulprizio, M.P., Maasakkers, J.D., Bloom, A.A., Worden, J.R., Parker, R.J., Delgado, A.L., 2021. Global distribution of methane emissions: a comparative inverse analysis of observations from the TROPOMI and GOSAT satellite instruments. Atmospheric Chemistry and Physics 21, 14159–14175. https://doi.org/10.5194/acp-21-14159-2021

Randerson, J.T., Chen, Y., van der Werf, G.R., Rogers, B.M., Morton, D.C., 2012. Global burned area and biomass burning emissions from small fires. Journal of Geophysical Research: Biogeosciences 117. https://doi.org/10.1029/2012JG002128

Roberts, G., Wooster, M.J., Perry, G.L.W., Drake, N., Rebelo, L.-M., Dipotso, F., 2005. Retrieval of biomass combustion rates and totals from fire radiative power observations: Application to southern Africa using geostationary SEVIRI imagery. Journal of Geophysical Research: Atmospheres 110. https://doi.org/10.1029/2005JD006018

Rodgers, C.D., 2000. Inverse Methods for Atmospheric Sounding: Theory and Practice, in: Series on Atmospheric, Oceanic and Planetary Physics. WORLD SCIENTIFIC. https://doi.org/10.1142/3171

Roelfsema, M., van Soest, H.L., Harmsen, M., van Vuuren, D.P., Bertram, C., den Elzen, M., Höhne, N., Iacobuta, G., Krey, V., Kriegler, E., Luderer, G., Riahi, K., Ueckerdt, F., Després, J., Drouet, L., Emmerling, J., Frank, S., Fricko, O., Gidden, M., Humpenöder, F., Huppmann, D., Fujimori, S., Fragkiadakis, K., Gi, K., Keramidas, K., Köberle, A.C., Aleluia Reis, L., Rochedo, P., Schaeffer, R., Oshiro, K., Vrontisi, Z., Chen, W., Iyer, G.C., Edmonds, J., Kannavou, M., Jiang, K., Mathur, R.,

Safonov, G., Vishwanathan, S.S., 2020. Taking stock of national climate policies to evaluate implementation of the Paris Agreement. Nat Commun 11, 2096. https://doi.org/10.1038/s41467-020-15414-6

Rosa, I.M.D., Pereira, J.M.C., Tarantola, S., 2011. Atmospheric emissions from vegetation fires in Portugal (1990–2008): estimates, uncertainty analysis, and sensitivity analysis. Atmospheric Chemistry and Physics 11, 2625–2640. https://doi.org/10.5194/acp-11-2625-2011

Samborska, V., Ritchie, H., Roser, M., 2024. Wildfires [https://ourworldindata.org/wildfires]. Our World in Data.

San-Miguel-Ayanz, J., Durrant, T., Boca, R., Maianti, P., Liberta', G., Artes, V.T., Jacome, F.O.D., Branco, A., De, R.D., Ferrari, D., Pfeiffer, H., Grecchi, R., Onida, M., Loffler, P., 2022. Forest Fires in Europe, Middle East and North Africa 2021 [WWW Document]. JRC Publications Repository. https://doi.org/10.2760/34094

Scarpa, C., Bacciu, V., Ascoli, D., Costa-Saura, J.M., Salis, M., Sirca, C., Marchetti, M., Spano, D., 2024. Estimating annual GHG and particulate matter emissions from rural and forest fires based on an integrated modelling approach. Science of The Total Environment 907, 167960. https://doi.org/10.1016/j.scitotenv.2023.167960

Schicker, I., Radanovics, S., Seibert, P., 2010. Origin and transport of Mediterranean moisture and air. Atmospheric Chemistry and Physics 10, 5089–5105. https://doi.org/10.5194/acp-10-5089-2010

Shaman, J., 2014. The Seasonal Effects of ENSO on European Precipitation: Observational Analysis. https://doi.org/10.1175/JCLI-D-14-00008.1

Shen, L., Jacob, D.J., Gautam, R., Omara, M., Scarpelli, T.R., Lorente, A., Zavala-Araiza, D., Lu, X., Chen, Z., Lin, J., 2023. National quantifications of methane emissions from fuel exploitation using high resolution inversions of satellite observations. Nat Commun 14, 4948. https://doi.org/10.1038/s41467-023-40671-6

Shi, Y., Gong, S., Zang, S., Zhao, Y., Wang, W., Lv, Z., Matsunaga, T., Yamaguchi, Y., Bai, Y., 2021. High-resolution and multi-year estimation of emissions from open biomass burning in Northeast China during 2001–2017. Journal of Cleaner Production 310, 127496. https://doi.org/10.1016/j.jclepro.2021.127496

Song, Y., Liu, B., Miao, W., Chang, D., Zhang, Y., 2009. Spatiotemporal variation in nonagricultural open fire emissions in China from 2000 to 2007. Global Biogeochemical Cycles 23. https://doi.org/10.1029/2008GB003344

Stohl, A., Seibert, P., Arduini, J., Eckhardt, S., Fraser, P., Greally, B.R., Lunder, C., Maione, M., Mühle, J., O'Doherty, S., Prinn, R.G., Reimann, S., Saito, T., Schmidbauer, N., Simmonds, P.G., Vollmer, M.K., Weiss, R.F., Yokouchi, Y., 2009. An analytical inversion method for determining regional and global emissions of greenhouse gases: Sensitivity studies and application to halocarbons. Atmospheric Chemistry and Physics 9, 1597–1620. https://doi.org/10.5194/acp-9-1597-2009

Turco, M., Herrera, S., Tourigny, E., Chuvieco, E., Provenzale, A., 2019. A comparison of remotely-sensed and inventory datasets for burned area in Mediterranean Europe. International Journal of Applied Earth Observation and Geoinformation 82, 101887. https://doi.org/10.1016/j.jag.2019.05.020

Turco, M., Llasat, M.C., von Hardenberg, J., Provenzale, A., 2013. Impact of climate variability on summer fires in a Mediterranean environment (northeastern Iberian Peninsula). Climatic Change 116, 665–678. https://doi.org/10.1007/s10584-012-0505-6

Turco, M., Rosa-Cánovas, J.J., Bedia, J., Jerez, S., Montávez, J.P., Llasat, M.C., Provenzale, A., 2018. Exacerbated fires in Mediterranean Europe due to anthropogenic warming projected with non-stationary climate-fire models. Nat Commun 9, 3821. https://doi.org/10.1038/s41467-018-06358-z

United Nations Environment Programme, 2024. Global Waste Management Outlook 2024: Beyond an age
of waste – Turning rubbish into a resource. Nairobi.
https://wedocs.unep.org/20.500.11822/44939

UNFCCC, 2015. Paris Agreement. https://climate.ec.europa.eu/eu-action/international-action-climate-
change/climate-negotiations/paris-agreement_en (accessed 26 June 2023).

van der Werf, G.R., Randerson, J.T., Giglio, L., van Leeuwen, T.T., Chen, Y., Rogers, B.M., Mu, M., van Marle,
M.J.E., Morton, D.C., Collatz, G.J., Yokelson, R.J., Kasibhatla, P.S., 2017. Global fire emissions
estimates during 1997–2016. Earth System Science Data 9, 697–720.
https://doi.org/10.5194/essd-9-697-2017

Volkova, L., Roxburgh, S.H., Surawski, N.C., Meyer, C.P. (Mick), Weston, C.J., 2019. Improving reporting of
national greenhouse gas emissions from forest fires for emission reduction benefits: An example
from Australia. Environmental Science & Policy 94, 49–62.
https://doi.org/10.1016/j.envsci.2018.12.023

Wilcox, R.R., 2010. Fundamentals of Modern Statistical Methods: Substantially Improving Power and
Accuracy. Springer, New York, NY. https://doi.org/10.1007/978-1-4419-5525-8

Yona, L., Cashore, B., Jackson, R.B., Ometto, J., Bradford, M.A., 2020. Refining national greenhouse gas
inventories. Ambio 49, 1581–1586. https://doi.org/10.1007/s13280-019-01312-9

Zhai, P., Yu, R., Guo, Y., Li, Q., Ren, X., Wang, Y., Xu, W., Liu, Y., Ding, Y., 2016. The strong El Niño of 2015/16
and its dominant impacts on global and China's climate. J Meteorol Res 30, 283–297.
https://doi.org/10.1007/s13351-016-6101-3

Zhang, W., Wang, Z., Stuecker, M.F., Turner, A.G., Jin, F.-F., Geng, X., 2019. Impact of ENSO longitudinal
position on teleconnections to the NAO. Clim Dyn 52, 257–274. https://doi.org/10.1007/s00382-
018-4135-1

Zhang, Y., Jacob, D.J., Lu, X., Maasakkers, J.D., Scarpelli, T.R., Sheng, J.-X., Shen, L., Qu, Z., Sulprizio, M.P.,
Chang, J., Bloom, A.A., Ma, S., Worden, J., Parker, R.J., Boesch, H., 2021. Attribution of the
accelerating increase in atmospheric methane during 2010–2018 by inverse analysis of GOSAT
observations. Atmospheric Chemistry and Physics 21, 3643–3666. https://doi.org/10.5194/acp-
21-3643-2021

World Meteorological Organization, 2023. WMO Greenhouse Gas Bulletin. No. 19 – 15 November 2023.
https://library.wmo.int/idurl/4/68532