Partial redundancy buffers deleterious effects of mutating *DNA methyltransferase 1-1* (*MET1-1*) in polyploid wheat

Samuel Burrows[1]*, Delfi Dorussen[1]*, Joseph Crudgington[1], Giorgia Di Santolo[1], James Simmonds[1], Marco Catoni[2], Philippa Borrill[1†]

[1] Department of Crop Genetics, John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, United Kingdom.

[2] School of Biosciences, University of Birmingham, Birmingham, B15 2TT, United Kingdom.

* These authors contributed equally

[†] Corresponding author, email: philippa.borrill@jic.ac.uk

Keywords: DNA methylation, DNA methyltransferases, Wheat (*Triticum aestivum*), polyploidy, epibreeding

Running title: Partial redundancy of *MET1-1* in wheat

The author responsible for distribution of materials integral to the findings presented in this article is: Philippa Borrill (philippa.borrill@jic.ac.uk).

Abstract

DNA methylation is conserved across biological kingdoms, playing important roles in gene expression, transposable element silencing and genome stability. Altering DNA methylation could generate additional phenotypic variation for crop breeding, however the lethality of epigenetic mutants in crop species has hindered its investigation. Here, we exploit partial redundancy between homoeologs in polyploid wheat to generate viable mutants in the DNA *methyltransferase 1-1* (*MET1-1*) gene with altered methylation profiles. In both *Triticum turgidum* (tetraploid wheat) and *Triticum aestivum* (hexaploid wheat) we identified clear segregation distortions of higher-order mutants (5/6 and 6/6 mutant *met1-1* copies in hexaploid and 3/4 and 4/4 copies in tetraploid) when genotyping segregating seeds and seedlings, which we attribute to reduced transmission of null mutant gametes. We found that the reduced transmission occurred from both the maternal and paternal gametes, however, we did not detect any deleterious effects on pollen development. The loss of four or more functional copies of *MET1-1* results in decreased CG methylation in hexaploid wheat.

Changes to gene expression increase stepwise with the number of mutant alleles suggesting a

dosage dependent effect. Finally, we identify heritable changes to flowering and awn

phenotypes which segregate independently of *MET1-1*. Together our results demonstrate that

polyploidy can be leveraged to generate quantitative changes to CG methylation without the

lethal consequences observed in other crops, opening the potential to exploit novel epialleles

in plant breeding.

Introduction

DNA methylation on cytosine bases is an epigenetic modification which is conserved across

biological kingdoms and plays important roles in gene expression, transposable element

silencing and genome stability. In plants, cytosine methylation can occur in any context (CG,

CHG and CHH, where H is A, T, or C) although CG methylation is the most prevalent

context in plant species examined to date (Feng *et al.*, 2010; Niederhuth *et al.*, 2016; Domb *et*

al., 2020; Li *et al.*, 2023). Methylation is established and maintained by multiple enzymes

which function in specific or overlapping contexts. CG methylation is maintained during cell

division by the methyltransferase 1 (MET1) which recognises hemi-methylated CG sites

following DNA replication and methylates the opposite strand (Kankel *et al.*, 2003; Zhang *et*

al., 2018).

Disruption of the *MET1* gene in Arabidopsis (*Arabidopsis thaliana*), tomato (*Solanum*

lycopersicum) and rice (*Oryza sativa*) has revealed major differences in the phenotypic

impact and severity of null mutations between species. Arabidopsis *met1* mutants, including

complete and partial loss-of-function alleles, exhibit dwarfism, delayed flowering, abnormal

embryo development and seed abortion (Kankel *et al.*, 2003; Saze *et al.*, 2003; Xiao *et al.*,

2006; Srikant *et al.*, 2022). Null *met1* alleles have severe consequences with segregation

distortion resulting in an underrepresentation of homozygous mutants, which are only

occasionally viable to maturity (Saze *et al.*, 2003; Srikant *et al.*, 2022). In rice, null

homozygous mutants were formed at the expected ratio but underwent necrotic death at

seedling stage (Hu *et al.*, 2014) and in tomato, homozygous null mutants could not be

recovered (Yang *et al.*, 2019). These species also vary in CG methylation loss in *met1*

mutants, with a stronger reduction in Arabidopsis than in rice (98.3% vs 75.7%) and only a

25% reduction in tomato CRISPR mutants, although this may be due to chimerism (Cokus *et*

al., 2008; Hu *et al.*, 2014; Yang *et al.*, 2019).

Differences in the segregation and viability of homozygous null mutants may be due to
variation in epigenetic backup systems: in Arabidopsis these backup systems operate through
changes to the RNA-directed DNA methylation (RdDM) process, changes to expression of
DNA demethylases and progressive H3K9 remethylation of heterochromatin (Mathieu *et al.*,
2007). Stochasticity in these processes explains the rarity of homozygous null mutants
(Mathieu *et al.*, 2007). In rice, a duplicate copy of *MET1* has been proposed to provide a
back-up mechanism for essential CG methylation, although the lethality of homozygous
mutants at seedling stage indicates incomplete compensation (Hu *et al.*, 2014; Yamauchi *et*
al., 2014).

Mutants in *met1* have not been examined within a polyploid species where duplicate gene
copies could provide a back-up mechanism. Alternatively, gene copies could be dosage-
sensitive with a mutation in just one copy of *met1* affecting DNA methylation. The gene
balance hypothesis proposes that dosage-sensitive genes are generally involved in
macromolecular complexes or are regulatory genes such as transcription factors, rather than
enzymes like *MET1* (Veitia & Birchler, 2010; Birchler & Veitia, 2012). Moreover, it is
generally expected that mutations in enzymes would be recessive, making it unlikely to find
phenotypic effects in a partial *met1* mutant in a polyploid (Kondrashov & Koonin, 2004).
However, some enzymes are dosage-sensitive in polyploids, such as the E3 ubiquitin ligase
gene *GW2* which has a dosage effect on grain width in wheat (Wang *et al.*, 2018). MET1 has
been reported to interact at the protein level with histone deacetylase HDA6 and MEDEA in
Arabidopsis (Liu *et al.*, 2012; Schmidt *et al.*, 2013), making a dosage effect more likely since
it could be part of a macromolecular complex. Opposing this view, heterozygous mutants in
tomato and rice maintained similar CG methylation to wild type plants (Hu *et al.*, 2014; Yang
et al., 2019), indicating only one allele was sufficient to compensate, and the same was found
for some Arabidopsis accessions, although CG methylation was reduced up to 64% in others
(Srikant *et al.*, 2022). Therefore, it is difficult to predict whether a partial mutant in *MET1* in
a polyploid species would induce changes to CG methylation.

If *MET1* is dosage-sensitive in polyploids, partial *met1* mutants could be used to alter
methylation status and produce stable epialleles for use in plant breeding. Epialleles affecting
diverse traits have been identified in a range of plant species, including alterations in floral
morphology, fruit ripening, plant height and climate adaptation (Cubas *et al.*, 1999; Manning
et al., 2006; Miura *et al.*, 2009; He *et al.*, 2018; Srikant & Tri Wibowo, 2021). The deliberate
induction of stable epialleles has been demonstrated through epiRIL populations generated

genetically using *met1* and *ddm1* (*Decreased DNA Methylation 1*) mutants in Arabidopsis

(Johannes *et al.*, 2009; Reinders *et al.*, 2009; Zhang *et al.*, 2013; Catoni & Cortijo, 2018). The

potential to exploit novel epialleles in plant breeding has been proposed, but the production

of epiRIL populations in crop species as a foundation to generate novel epialleles has been

limited by the lethality of mutants (Springer & Schmitz, 2017). For example, no null mutants

were identified in *MET1* orthologs in crop species with large genomes including in a barley

TILLING population (genome size of ~5Gb) (Schreiber *et al.*, 2019) and in maize TILLING

and transposon populations (genome size of ~2.4 Gb) (Li *et al.*, 2014). This may be due the

large numbers of transposable elements in large genomes causing strong genetic and

developmental defects in *met1* mutants through their remobilisation (Mirouze & Vitte, 2014).

Here we explore whether it is possible to use a polyploid crop species to generate quantitative

changes to CG methylation by knocking out differing numbers of *met1* alleles, without

causing lethal consequences. We use polyploid wheat as a model system due to the

availability of sequenced mutant populations which can be readily applied to breeding

worldwide without legislative complications surrounding gene editing (Krasileva *et al.*,

2017). We generate *met1* mutants in tetraploid (*Triticum turgidum*) and hexaploid (*Triticum*

aestivum) wheat with up to three null alleles (of four) in tetraploid wheat, and five null alleles

(of six) in hexaploid wheat. Contrary to the predictions of the gene dosage balance hypothesis

(Veitia & Birchler, 2010) we find changes to global DNA methylation, transcriptional

responses and effects on fertility and plant growth linked to mutant allele copy number. We

also identify stably inherited phenotypic alterations. Our findings provide insights into using

polyploidy to generate quantitative changes to CG methylation for epigenetic breeding in

crops.

Results

Null *met1-1* mutants cannot be recovered in polyploid wheat

METHYLTRANSFERASE 1 (*MET1*) genes were previously identified in hexaploid wheat

(*Triticum aestivum*) on the group 2, 5, and 7 chromosomes, with the group 5 and 7 *MET1*

homologs predicted to be pseudogenes (Figure 1A) (Thomas *et al.*, 2014). The group 2 genes

are orthologous to *MET1b* in rice, also known as *OsMET1-2,* which strongly affects CG

methylation (Figure 1A) (Hu *et al.*, 2014). The group 2 genes (*TraesCS2A02G235900,*

TraesCS2B02G260800, *TraesCS2D02G241800*) are expressed in numerous above-ground

tissues (Figure 1B), whereas expression of the group 5 and group 7 genes was virtually

undetectable in most of these tissues (Supplementary Figure 1). We therefore concentrated

our analysis on the group 2 genes which will be referred to as *MET1-A1*, *MET1-B1*, and

MET1-D1 in accordance with the gene nomenclature guidelines for wheat (Boden *et al.*,

2023) and a previous study by Li *et al.* (2021).

We produced independent segregating F_2 populations in hexaploid wheat from which, in

theory, all combinations of loss-of-function *met1-1* mutants could be recovered (Figure 1C,

Supplementary Figure 2). However, we were unable to recover null *met1-1* mutants with

homozygous loss-of-function mutations in all three homoeologs from either population,

despite screening 3,068 individuals from the C0465xC0884 population, and 328 individuals

from the C0451xC2028xC1292 population. Moreover, we only found one plant with five

non-functional *MET1-1* alleles which was capable of germination, although its growth was

severely stunted and the plant died before progressing to tillering (Zadoks growth stage 20)

(Figure 1D). The genotype ratios observed in both populations were significantly different

from expected Mendelian ratios as individuals with four or more non-functional *MET1-1*

alleles were underrepresented (C0465xC0884 Figure 1E, χ^2 p=1.63x10^{-103};

C0451xC2028xC2192 Supplementary Table 1, χ^2 p=5.05x10^{-12}). We concluded that higher-

order *met1-1* mutants have reduced viability in hexaploid wheat.

Figure 1. Null *met1-1* mutants cannot be produced by crossing TILLING lines with
disruptions in the three *MET1-1* homoeologs. A) Evolutionary relationships between *MET1*
homologs in Arabidopsis (*AtMET1*), rice (*OsMET1a* and *OsMET1b*), and wheat (homoeolog
groups *TaMET1-1*, *TaMET1-2*, and *TaMET1-3*). B) Expression of the three *MET1-1*
homoeologs (*TraesCS2A02G235900, TraesCS2B02G260800, TraesCS2D02G241800*) in
above-ground tissues across development in the Chinese Spring cultivar. SAM = shoot apical
meristem. C) Structure of the three *MET1-1* homoeologs. Filled rectangles represent exons,
empty rectangles represent untranslated regions, and lines represent introns. Triangles

indicate the positions of the premature termination codon mutations in the TILLING lines
used to produce the two hexaploid F_2 populations (C0465xC0884, orange, and
C0451xC2028xC1292, purple). Green box signifies Bromo adjacent homology domain
(BAH) and blue box signifies C-5 cytosine methyltransferase domain. D) Cadenza WT, WT
segregant (AABBDD), and Aabbdd individuals from the C0465xC0884 segregating F_2
population. E) Expected (grey) and observed (orange) number of individuals with each
possible number of mutant *met1-1* alleles in C0465xC0884 segregating F_2 population.
164

Reduced viability of null *met1-1* gametes hinders recovery of higher-order *met1-1* mutants

We hypothesized that the absence of functional *MET1-1* alleles within a gamete might cause

this segregation distortion. We developed a grain genotyping method using the endosperm-

containing half of the grain (Figure 2A) which contains maternal to paternal DNA at a 2:1

ratio (endosperm originates from two female central cell nuclei being fertilised by one male

sperm cell). This allowed us to determine which parent donated mutant alleles to the

offspring (grain) by separating the heterozygous genotypes into two clusters (Figure 2B). We

confirmed the accuracy of this method using reciprocal crosses for an independent mutation

in the semi-dwarfing gene *Rht13* (Borrill *et al.*, 2022) (Figure 2B). Grain genotypes were

consistent with seedlings grown from the embryo-containing halves of the grain

(Supplementary Table 2). We genotyped 1,710 grains from a segregating F_2 hexaploid wheat

population (C0465xC0884) and observed far fewer grains which inherited three mutant *met1-*

1 alleles from one parent than expected (generalised linear model with log link, $p<2x10^{-16}$;

Figure 2C). We detected fewer grains which inherited three mutant alleles from the maternal

side than from the paternal side (18 vs 32 grains). There was no reduction in the number of

grains which inherited one or two mutant alleles from the same parent (Figure 2C).

We confirmed these observations in an independent segregating F_4 tetraploid wheat

population (332 grains) (Supplementary Figure 3). We observed a significant segregation

distortion (χ^2 p= $3.39x10^{-30}$) and were unable to identify any homozygous null mutants

(aabb). Tetraploid wheat gametes with a complete knock-out of *MET1-1* (i.e. ab gametes)

were also strongly underrepresented (generalised linear model with log link, $p<2x10^{-16}$), with

reduced maternal and paternal transmission (Supplementary Table 3). In tetraploid wheat we

also detected fewer grains which inherited two mutant alleles from the maternal side than

from the paternal side (1 vs 18 grains; Supplementary Table 3).

In hexaploid wheat, we found the distribution of genotypes in grains was significantly

different from seedling leaf tissue (χ^2 p=0.00058), with a greater proportion of individuals

with five non-functional *MET1-1* alleles observed in the grain population (12 out of 1,710 in

the grain population [0.702%] compared to 1 out of 1,352 in the seedling population

[0.074%]). This suggests that higher-order *met1-1* mutants may be formed as grains, but not

be capable of germination. We hypothesised *met1-1* mutants inheriting three non-functional

alleles from the same parent may be associated with decreased grain size. We found that, for

grains with a total of three or four mutant *met1-1* alleles, those inheriting three mutant alleles

from one parent were smaller by ~1.5mm than those inheriting the mutant alleles separately

from both parents (ordered logistic regression model, p<0.05; Supplementary Figure 4).

To further understand why the recovery of *met1-1* mutants formed from at least one null

mutant gamete is reduced, we investigated whether pollen development was affected in the

met1-1 mutants. We measured the number of pollen grains per anther, pollen diameter, pollen

viability, and the proportion of pollen grains that were non- or mononucleated for plants from

the segregating F_2 population, including those that should be capable of producing gametes

with no functional copies of *MET1-1* (AaBbdd, AabbDd, and aaBbDd). However, no

significant differences (at α=0.05) were detected between those with an AaBbdd, AabbDd, or

aaBbDd genotype and the WT segregant (AABBDD) for any of these phenotypes

(Supplementary Table 4).

To confirm whether inheritance of three mutant *met1-1* alleles from the same parent reduces

viability, we performed reciprocal crosses between triple heterozygous (AaBbDd) *met1-1*

mutants and Cadenza WT. In total, 454 progeny were genotyped, 235 with AaBbDd as the

female parent and 219 with AaBbDd as the male parent. Again, we found that plants

inheriting three non-functional *MET1-1* alleles from one parent were significantly

underrepresented compared to the expected Mendelian ratio (χ^2 p<0.001 for both maternal

and paternal donation; Figure 2D). The proportion of abd gametes was lower from the

paternal side of the reciprocal crosses than the maternal side, opposite to the trend observed

in the selfed progeny of a triple heterozygous plant (Figure 2C).

220
Figure 2. Inheritance of three mutant alleles *met1-1* from the female or male parent (i.e. abd
gamete) is rare relative to inheritance of zero, one or two mutant alleles. A) Wheat grains
divided transversely to separate two halves of the grain, with one half containing the embryo
for future germination and the other half containing largely endosperm for DNA extraction
and genotyping. B) KASP genotyping for *Rht13* mutants (*Rht-B13b*) from reciprocal crosses
between mutant and wild type parents with known genotypes, showing homozygous wild
type (WT) (red cluster), homozygous mutant (blue cluster), heterozygous clusters with a
maternally inherited WT allele and paternally inherited mutant allele (brown), a paternally
inherited WT allele and maternally inherited mutant allele (grey) and no template control
(black). C) Expected (grey) and observed (orange) number of individuals inheriting zero, one,
two, or three mutant *met1-1* alleles from the maternal or paternal gamete from a self-
pollinated AaBbDd plant. D) Expected and observed proportion of individuals formed from
each type of gamete when inherited maternally (♀ donation) or paternally (♂ donation) from
an AaBbDd parent when crossed with AABBDD. Orange shades indicate proportions below
the expected proportion of 0.125.

Loss of four or more functional copies of *MET1-1* results in decreased CG methylation

We carried out whole genome bisulfite sequencing to determine whether a stepwise reduction in the number of functional *MET1-1* alleles was associated with a proportional change in DNA methylation. We observed partial functional redundancy between the *MET1-1* homoeologs; genome-wide CG methylation in the single mutants (aaBBDD, AAbbDD, and AABBdd) was between 87.0% and 87.1% which was very similar to the WT segregant (AABBDD, 87.3% CG methylation). Genome-wide CG methylation was reduced in the double mutants, ranging from 74.0% in the aabbDD mutant, to 79.3% in the aaBBdd mutant, and 81.2% in the AAbbdd mutant. The lowest level of CG methylation was observed in the Aabbdd mutant, with 72.4% (Figure 3A). However, the Aabbdd mutant had the highest CHG (57.4%) and CHH (4.1%) methylation of all the lines tested (Figure 3A).

To explore the impact of these global changes in methylation, we examined methylation across genes and transposable elements (TEs), using chromosome 5D as an example. Average CG methylation was decreased across genes and TEs relative to the WT segregant (AABBDD) in both the double mutant (AAbbdd) and the Aabbdd mutant (Fig 3B and C). For both sequence features, the greatest decrease was observed for the Aabbdd mutant, in line with the global changes in methylation. No obvious difference was observed in CHG methylation across genes or TEs, while CHH methylation was highest in the Aabbdd mutant across both genes and TEs (Supplementary Figure 5).

Spatial correlations between different methylation contexts were observed when looking at the distribution of cytosine methylation across each chromosome. Along chromosome 5D, CG methylation is reduced in the Aabbdd mutant compared to the WT segregant, single and double mutants in the R1, R2a, C, and R2b regions. In the R3 region, however, CG methylation in the Aabbdd mutant is comparable to the AAbbdd double mutant (Figure 3D). In this region, there is also an increase in CHG methylation in the Aabbdd mutant, such that it is higher than any of the other lines (Figure 3E), and an increase in CHH methylation (Figure 3F). There is a clear transition point between states in the Aabbdd mutant across all three methylation contexts (Supplementary Figure 6). The simultaneous increase in CG and CHG/CHH methylation is also observed in specific regions of another ten chromosomes, affecting all possible chromosome regions not only R3, and across the whole of

chromosomes 3A, 4D and 6B (Supplementary Figures 7-13). These region-specific increases

in CG, CHG and CHH methylation were only observed in the Aabbdd mutant, and not in any

of the single or double mutants, as shown for chromosome 5D (Supplementary Figure 14).

To further understand the spatial nature of methylation changes in the *met1-1* mutants, we

identified 100 bp differentially methylated regions (DMRs) in the CG context in each of the

mutants relative to the WT segregant. Substantially more DMRs were detected in the Aabbdd

mutant (1,156,551 DMRs), compared to between 98,626 and 282,113 DMRs in the double

mutants, and between 10,616 and 11,803 DMRs in the single mutants (Figure 3G). The vast

majority of DMRs were hypo-methylated in the *met1-1* mutants – only 0.27% of the DMRs

in the Aabbdd mutant were hyper-methylated (Figure 3G). We observed that, in the double

mutants, the distal regions R1 and R3 have the greatest density of DMRs, while the

centromeric region C has the lowest density (Figure 3H). However, in the Aabbdd mutant the

centromeric region has the highest density of DMRs (Figure 3H). In all *met1-1* mutants, the

density of hyper-methylated DMRs is lowest in the proximal and centromeric regions, and

highest in the distal regions (Supplementary Figure 15).

283
Figure 3: Reduced number of functional copies of *MET1-1* results in loss of CG methylation
and the formation of differentially methylated regions (DMRs). A) Genome-wide percentage
of cytosines methylated in the CG, CHG, and CHH contexts. B-C) Percentage CG
methylation across genes (B) and transposable elements (TEs) (C) on chromosome 5D,
including 2 kb up and downstream of the feature. The WT segregant (AABBDD, black),
AAbbDD single mutant (yellow), AAbbdd double mutant (blue), and Aabbdd mutant (red)
are shown. D-F) Percentage of methylated cytosines in the CG (D), CHG (E), and CHH (F)
contexts across chromosome 5D, calculated as an average in 1 Mb bins across the
chromosome. The WT segregant (AABBDD, black), AAbbDD single mutant (yellow),

AAbbdd double mutant (blue), and Aabbdd mutant (red) are shown. Partitions between the
chromosome regions are indicated by the dashed lines. G) Number of differentially
methylated regions (DMRs) in 1000s relative to AABBDD in each genotype. Both hypo-
methylated (dark grey) and hyper-methylated (light grey) regions are shown. H) Number of
hypo-methylated DMRs per kb in each of the chromosome regions – the distal regions R1
and R3 are shown in dark grey, the proximal regions R2a and R2b in medium grey, and the
centromeric region C in light grey.

Reduced function of *MET1-1* results in stepwise changes in gene expression

To determine the effect of these changes in methylation, we conducted RNA-sequencing and

differential expression analysis. The number of differentially expressed genes (DEGs)

compared to the WT segregant (FDR adjusted $p<0.01$, fold change ≥ 2) was calculated for

each mutant class (Figure 4A). The Aabbdd mutant had the largest number of DEGs (5,904),

while double mutants had on average 79 DEGs, and single mutants had on average 9 DEGs.

Generally, there were more up-regulated DEGs than down-regulated DEGs, except for the

Aabbdd mutant, with 3,900 genes down-regulated and 2,004 genes up-regulated (Figure 4A).

There was also a stronger change in transposable element (TE) expression in the double and

Aabbdd mutants compared to single mutants, in which no differentially expressed TEs

(DETs) were observed (on average 36 DETs in double mutants and 143 DETs in Aabbdd

mutant; FDR adjusted $p<0.01$, fold change ≥ 2, Figure 4B). These changes in gene and TE

expression are in line with the global changes in CG methylation, with mutants maintaining

the highest level of CG methylation relative to the WT segregant having the fewest DEGs,

and vice versa (Figures 3A, 4A and 4B).

To further examine the transcriptional response, we identified genes that were differentially

expressed in both the Aabbdd mutant and at least one of the double mutants. A total of 86

DEGs were identified in this way, 76 of which were up-regulated, and 10 of which were

down-regulated. Several of these DEGs have putative functions relating to DNA methylation

and other epigenetic marks (Figure 4C), suggesting that reduced *MET1-1* function initiates

active changes in transcriptional regulation. The gene *TraesCS6B02G315200* is up-regulated

in all double mutants and the Aabbdd mutant (adjusted $p<0.01$) (Figure 4C), and is annotated

in the Panther Classification System as RNA Polymerase IV and V Subunit 4

(PTHR21297:SF2). The homoeologous gene *TraesCS6D02G266900* is also up-regulated in

all double mutants and the Aabbdd mutant, while its other homoeolog *TraesCS6A02G286200*

and tandem duplicated gene copy *TraesCS6B02G15300*, are up-regulated only in the aabbDD

and Aabbdd mutants. A gene with the same Panther classification in Arabidopsis, *NRPD4*

(*AT4G15950*), is required for RNA-directed DNA methylation (RdDM) (He *et al.*, 2009). The

gene *TraesCS2D02G114600* was also up-regulated in all double *met1-1* mutants (Figure 4C).

This gene encodes a small RNA 2'-O-methyltransferase annotated as HEN1 in the Panther

Classification System (PTHR21404). In Arabidopsis, *HEN1* is also implicated in RdDM

(Erdmann & Picard, 2020). Furthermore, in both the aabbDD and Aabbdd mutants,

TraesCS3B02G023200, a DEMETER (DME)-like DNA glycosylase, is significantly down-

regulated (Figure 4C). In Arabidopsis, loss-of-function mutations in DME-like DNA

glycosylases *DML2* and *DML3* result in increased cytosine methylation in regions with low

to intermediate methylation levels in wild type (Ortega-Galisteo *et al.*, 2008). Overall, this

suggests that reduced *MET1-1* function is buffered by up-regulation of genes that facilitate

DNA methylation and down-regulation of genes that remove DNA methylation.

We also observed up-regulation of *TraesCS2B02G503200* in the Aabbdd, aabbDD and

AAbbdd mutants (adjusted $p<0.01$) and the aaBBdd mutants (adjusted $p<0.05$) (Figure 4C).

This gene is predicted to contain a histone-lysine N-methyltransferase

SUVR4/SUVR1/SUVR2 domain (IPR025776). Reduced function of *MET1-1* may therefore

promote changes in histone methylation that could further alter transcription.

344
Figure 4. Reduced number of functional copies of *MET1-1* results in transcriptional
reprogramming. A) Number of differentially expressed genes (DEGs) relative to AABBDD in
each genotype. Both down-regulated (light purple) and up-regulated (dark purple) genes are
shown. B) Number of differentially expressed transposable elements (DETs) relative to
AABBDD in each genotype (uniquely mapped reads only). Both down-regulated (light
purple) and up-regulated (dark purple) transposable elements are shown. C) Normalised
transcript counts in each genotype for genes putatively encoding an RNA Polymerase IV/V
subunit (*TraesCS6B02G315200*), a HEN1 small RNA 2'-O-methyltransferase

(*TraesCS2D02G114600*), a DEMETER (DME)-like DNA glycosylase
(*TraesCS3B02G023200*), and a histone-lysine methyltransferase (*TraesCS3B02G503200*).
Error bars represent the 95% confidence interval, estimated as 1.96 x standard error. Asterisks
show adjusted p-values for differential expression relative to AABBDD; * p<0.05, ** p<0.01,
*** p<0.001. D-E) The percentage of genes (D) or transposable elements (E) which intersect
at least one DMR in each of the *met1-1* mutants. Hyper-DMRs are shown in light grey and
hypo-DMRs are shown in dark grey. F) The percentage of DEGs or DETs associated with a
hypo-DMR which are down-regulated (light purple) or up-regulated (dark purple) in the
aabbDD and Aabbdd *met1-1* mutants.

362
363

Next, we investigated whether the CG-DMRs identified in the *met1-1* mutants (Figure 3)

overlapped genes and TEs. Up to 35.7% of high-confidence genes (IWGSC *et al.*, 2018)

overlapped with CG-DMRs in the *met1-1* mutants (either within the gene body, or within 1

kb up- or downstream) (Figure 4D). Although the Aabbdd mutant had 4.1-11.7x as many CG-

DMRs as the double mutants (Figure 3G), the percentage of genes intersected by at least one

CG-DMR (26.0%) was within the range observed in the double mutants (between 17.9% and

35.7%). In line with the overall distribution of DMRs, the majority of DMRs intersecting

genes were hypo-methylated in the *met1-1* mutants (Figure 4D).

In contrast to the DMRs intersecting genes, we observed a stark difference between the

percentage of TEs overlapped by at least one CG-DMR in the Aabbdd mutant and the double

mutants (Figure 4E). In the Aabbdd mutant, 20.1% of non-fragmented TEs (Wicker *et al.*,

2018) were intersected by at least one DMR, compared to between 0.95% and 4.7% in the

double mutants. This is in line with the greater density of hypo-DMRs in the transposon-

dense centromeric region in the Aabbdd mutant (Figure 3H). 87.0% of hypomethylated TEs

in the Aabbdd mutant were long terminal repeat retrotransposons (LTR-RTs), compared to

between 56.5% and 72.5% in the double mutants (Supplementary Table 5).

To examine whether changes in methylation status may be directly related to changes in gene

expression, we quantified the percentage of DEGs which overlapped at least one CG-DMR.

We focused on the double and Aabbdd mutants, as only a small number of DEGs were

identified in the single mutants (Figure 4A). Between 32.6% and 51.5% of DEGs were

associated with a DMR. For each mutant, the proportion of DEGs associated with a DMR is

greater than the proportion of genes associated with a DMR, suggesting that changes in DNA

methylation may be directly responsible for the change in expression of a subset of DEGs. In

the aabbDD double mutant, 80.6% of hypomethylated DEGs increased in expression relative

to the WT segregant (Figure 4F). However, in the Aabbdd mutant, only 26.9% of

hypomethylated DEGs were up-regulated (Figure 4F). This could reflect the overall greater

proportion of down-regulated genes in the Aabbdd mutant compared to the double mutants

(Figure 4A). The majority of DETs associated with a hypo-DMR were up-regulated in both

the aabbDD and Aabbdd mutants (77.8% in aabbDD and 80.4% in Aabbdd) (Figure 4F). The

same trend was observed for the other double mutants (Supplementary Figure 16).

Heritable phenotypic changes were identified in the *MET1-1* segregating F$_2$

population

When growing the C0451xC1292xC2028 F$_2$ population we identified one plant (AaBbdd

genotype) that flowered 23 days later than any other plant and had lax ears (thin ears

composed of many spikelets). Offspring from this late-flowering individual segregated into

normal flowering types with normal ears (<10 days relative to earliest flowering plant), and

late-flowering types with lax ears (≥10 days relative to earliest flowering plant) (Figure 5A-

C). The numbers of segregant plants in the late and normal flowering categories did not

significantly differ from a 3:1 ratio (χ^2 = 0.214, p-value = 0.644), suggesting that a single

dominant gene controls the phenotype. The late flowering phenotype was independent of

MET1-1 genotype (Figure 5D). The late flowering plants were also significantly taller than

normal flowering plants (FDR adjusted p-value = 8.2×10^{-15}; Figure 5A, E). There was no

significant difference in spike length between the late and normal flowering segregants (FDR

adjusted p-value = 0.22, Figure 5F), however late flowering segregants had a higher number

of fully formed spikelets per spike than normal flowering segregants (mean 25.3 vs 18.6;

FDR adjusted p-value = 5.5×10^{-9}, Figure 5B and G). These data show that the tall, late

flowering phenotype is heritable and is independent of *MET1-1* genotype, suggesting that it is

controlled by a single stable allele generated in the mutant background.

Figure 5. The late flowering phenotype in the C0451xC1292xC2028 *MET1-1* population was

independent of *MET1-1* genotype and was accompanied by increased plant height and

number of fully formed spikelets. A) Phenotypes of Cadenza WT and normal and late

flowering F$_3$ segregant plants. B) Normal and late flowering segregant plants showing the

increased number of fully formed spikelets in the late flowering plant. C) Histogram showing

the days to flowering (full emergence of primary spike) of the different flowering time

categories relative to the first plant. D) *MET1-1* genotypes of late and normal flowering

plants. Boxplots of plant height (E), primary spike length (F) and number of fully formed

spikelets (G) for each of the phenotype categories. For (E) to (G), the means were compared

by a t-test and the False Discovery Rate (FDR) adjusted p-value calculated is shown.

A second phenotype was observed in the same segregating F_2 population (C0451xC1292xC2028): one plant (AAbbdd genotype) exhibited an awned phenotype, despite all three parents being awnless. We generated a segregating population by backcrossing the awned plant with Cadenza WT and selfing the F_1 generation, which showed an intermediate (awnletted) phenotype. Amongst 215 F_2 plants the segregation ratio of awnless:awnletted:awned individuals did not significantly deviate from a single co-dominant gene 1:2:1 Mendelian ratio ($\chi^2 = 1.13$, p-value = 0.568) (Figure 6A and B). The awnletted individual had longer awns at the top of the ear, similar to the effect of the *Tipped1* (*B1*) awn suppressor gene (Huang *et al.*, 2020) and we hypothesised that this was the causal gene controlling the phenotype. We used published primer sequences to screen the *B1* region in the F_2 population (Figure 6C) (Huang *et al.*, 2020). PCR products could be obtained for awnless and awnletted plants, but not for awned plants either across the gene body or in the surrounding 28.5 Mbp (Figure 6C and D). Together, these results indicate that the presence of awns in our population is due to a heritable deletion >28.5Mbp which includes *B1*.

Figure 6. The observed awned phenotype in the C0451xC1292xC2028 *MET1-1* population followed a single gene semi dominant mendelian inheritance pattern, which was caused by a deletion encompassing *Tipped1* (*B1*). A) Phenotypes of the awnless Cadenza WT parent and spontaneous awned individual which were crossed together to generate the F_1 generation. The F_1 was self-pollinated to generate the segregating F_2 population which displayed awnless, awnletted and awned phenotypes. B) Expected (grey) and observed (purple) number of individuals which were awnless, awnletted and awned in the segregating F_2 population. C)

Diagram of the *B1* gene (black) and surrounding region (grey) with KASP marker binding
sites (black arrows) and PCR primer binding sites (maroon half arrows). Schematic
summarising successful (green tick) and unsuccessful (red cross) amplification of KASP
marker sets for each phenotype class of plant (awned, awnletted and awnless). D) Agarose gel
image showing PCR product of the ZF6F3 and ZF6R3 primer set (expected product ≈2kb)
from 3 individual F_2 segregant plants of each phenotype class and 1kb DNA ladder (1kb) and
no DNA (-C) controls.
454

Discussion

Complete loss of *MET1-1* is lethal in wheat, but single and double mutants can be recovered

457
458

Despite genotyping over 3,000 progeny from two segregating *MET1-1* populations, we were
unable to recover any null *met1-1* mutants in hexaploid wheat (Figure 1F, Supplementary
Table 1). Similarly, we were unable to recover any null *met1-1* mutants in tetraploid wheat
(Supplementary Table 3). This indicates that complete loss of *MET1-1* is lethal in wheat,
consistent with other monocotyledonous crop species with large genomes such as maize and
barley (Li *et al.*, 2014; Schreiber *et al.*, 2019).

However, unlike maize and barley, wheat is a polyploid, allowing intermediate mutants to be
recovered. We recovered all possible homozygous single and double *met1-1* mutants from the
segregating *MET1-1* populations suggesting that the function of *MET1-1* homoeologs in
wheat is mostly redundant. We also recovered a single Aabbdd mutant, in which five out of
six copies of *MET1-1* are non-functional (Figure 1D). We found that *MET1-1* has a threshold
for dosage-sensitivity: while the single mutants had no decrease in CG methylation compared
to the WT segregant, the double and Aabbdd mutants showed altered patterns of CG
methylation and gene expression (Figure 3 and 4). The double mutant classes showed
variability in the degree of CG methylation loss and changes to gene expression, with the
aabbDD mutants showing the greatest change compared to the AAbbdd and aaBBdd double
mutants (Figure 3 and 4). This suggests that *MET1-D1* is less effective at compensating for
the loss of the other two homoeologs than *MET1-A1* or *MET1-B1*. Furthermore, the double
mutants did not show any growth defects, despite the aabbDD mutants having a similar
degree of demethylation to the Aabbdd mutant (74.0% CG methylation and 72.4% CG
methylation respectively). Together these features suggests that double mutants with four

mutant alleles may allow the generation of epialleles which could be utilised in epigenetic

breeding (Figure 7).

Figure 7. Increasing the number of mutant *met1-1* alleles in hexaploid wheat has quantitative
yet variable effects on CG methylation, differentially expressed (DE) genes, differentially
methylated regions (DMRs) in genes and in transposable elements (TEs), viability and
phenotypic variation. Dashed regions indicate hypothetical data for a complete null mutant.
The region highlighted in yellow represents an optimum where CG methylation is
substantially reduced and viability is retained, made possible by partial redundancy between
homoeologs.

We observed novel, heritable phenotypic variation in one of the segregating *MET1-1*

populations (Figure 5 and 6). These phenotypes (delayed flowering and the development of

awns) were both identified in plants that had four out of six non-functional copies of *MET1-1*,

indicating that a null *met1-1* mutant is not necessary to generate phenotypic variation. Loss of

DNA methylation can generate novel variation in multiple ways. Firstly, epialleles may be

generated, resulting in altered patterns of gene expression. Secondly, hypomethylation of

transposable elements (TEs) is associated with their transcription and remobilisation, leading

to abnormal developmental phenotypes (Miura *et al.*, 2001; Mirouze *et al.*, 2009; Reinders *et*

al., 2009; Tsukahara *et al.*, 2009). We found that one fifth of TEs were associated with at least

one DMR in the Aabbdd mutant, and 45 of these TEs had up-regulated expression, potentially

resulting in their activation. This may explain the stunted growth of the Aabbdd mutant, and

the severely reduced viability of higher-order *met1-1* mutants. Genome instability caused by

loss of DNA methylation can also cause structural variation (Zhang *et al.*, 2023), which we

observed in the spontaneous awned *met1-1* mutant that had a substantial deletion (Figure 6).

A similar large deletion was found in wheat treated with the demethylation agent zebularine;
the deletion encompassed *FT-B1* and was associated with an increase in spikelet number and
a delay in flowering (Finnegan *et al.*, 2019).

Lethal effect of null *met1-1* mutants associated with mutant (abd) gametes
We observed a clear segregation distortion when genotyping selfed seed from a triple
heterozygous (AaBbDd) *MET1-1* parent, with underrepresentation of higher-level mutants
(5/6 and 6/6 non-functional copies). We attributed this distortion to a reduced transmission of
gametes with no functional copy of *MET1-1* (abd gametes) when donated maternally or
paternally. In reciprocal crosses, abd gametes transmitted paternally were rarer than abd
gametes transmitted maternally, whereas the opposite trend was observed using selfed
heterozygous (AaBbDd) mutants. This discrepancy may be explained by post-fertilisation
effects of *met1-1* being made more severe in a scenario where mutant alleles may be present
in both gametes, as for the selfed heterozygous plants. This is consistent with work in
Arabidopsis that found that selfed heterozygous *met1* mutants had higher rates of embryo
abortion than wild type x homozygous *met1* mutant reciprocal crosses (Xiao *et al.*, 2006).

In Arabidopsis, transmission of mutant *met1* alleles has also been shown to be distorted in the
gametes. However, in contrast to our results where both maternal and paternal transmission
was affected, only the paternal transmission of *met1* was reduced to 36.8% of the expected
rate in Arabidopsis while maternal transmission was not reduced (Liang *et al.*, 2022). This
single parental bias in *met1* transmission was, in part, due to impaired pollen development
resulting in abnormal numbers of nuclei in the pollen grains. In our work we found no change
to any pollen phenotype measured in the *met1-1* mutant plants compared to WT segregant
plants, suggesting that instead the barrier to the production of null mutants may occur during
fertilisation and/or embryogenesis. Post-zygotic effects are also likely important since we
found over 9x more offspring with five non-functional *MET1* alleles in grains than in leaves
of seedlings. Reduced germination may be due to abnormal endosperm development as
observed in Arabidopsis *met1* mutants (Jullien *et al.*, 2006). This would also be consistent
with the smaller seed sizes we observed in wheat mutants inheriting *met1-1* alleles through
abd gametes, although the effect was not as extreme as observed in rice (Hu *et al.*, 2014;
Yamauchi *et al.*, 2014), nor was it exclusively paternal as in Arabidopsis (Xiao *et al.*, 2006).

Reduced function of *MET1-1* is buffered by activation of other epigenetic pathways

Loss of five out of six functional copies of *MET1-1* resulted in a significant decrease in CG methylation, but an increase in CHG and CHH methylation (Figure 3), suggesting that other methylation pathways are activated upon reduction of *MET1-1* activity. A similar increase in CHG and CHH methylation was observed in tomato *met1* mutants (Yang *et al.*, 2019) and in rice *MET1/met1* heterozygotes, but not the homozygous *met1* mutant (Hu *et al.*, 2014). In Arabidopsis, CHG methylation was increased in *met1* mutants, while CHH methylation was decreased (Srikant *et al.*, 2022). In wheat, the increases in CHG and CHH methylation are found in extremely well-defined regions where CG methylation is maintained to at least the level of the AAbbdd double mutant (Figure 3, Supplementary Figures 5-12). Future studies are required to determine whether these regions feature only in species with large genomes, making them undetectable until now.

The compensatory increase in CHG and CHH methylation we observed may be promoted by transcriptional up-regulation of RNA-directed DNA methylation (RdDM). We found increased expression of genes involved in generating short interfering RNAs (siRNAs) which direct the RdDM machinery to loci to be methylated (He *et al.*, 2009; Erdmann & Picard, 2020); these include RNA Polymerase IV/V Subunit 4 and the HEN1 RNA methyltransferase in the double and Aabbdd *met1-1* mutants (Figure 4C). We also found down-regulation of a DME-like DNA glycosylase in the mutants with the largest reduction in CG methylation (aabbDD and Aabbdd) (Figure 4C). Down-regulation of DNA glycosylases has also been observed in *met1* mutants in Arabidopsis and rice (Mathieu *et al.*, 2007; Hu *et al.*, 2014). Transcriptional down-regulation of the Arabidopsis *ROS1* DNA glycosylase is caused by loss of methylation in its promoter at the methylation monitoring sequence (MEMS) (Lei *et al.*, 2015). We found that the DME-like DNA glycosylase down-regulated in the aabbDD and Aabbdd *met1-1* mutants overlapped a hypo-DMR, suggesting that a similar MEMS mechanism may regulate the expression of this DNA glycosylase, allowing the plant to sense changes in the DNA methylation status and compensate for them.

Compensation for reduced CG methylation can also occur through changes in histone modifications. We observed transcriptional up-regulation of a SUVR4/SUVR1/SUVR2 histone lysine methyltransferase in the double and Aabbdd *met1-1* mutants (Figure 4C). SUVR4/SUVR1/SUVR2 histone lysine methyltransferases have a preference for H3K9 methylation (Thorstensen *et al.*, 2006), suggesting that this epigenetic mark may also be

altered upon reduction of CG methylation in wheat, as has been reported in Arabidopsis *met1*
mutants (Tariq *et al.*, 2003; Mathieu *et al.*, 2007).

In summary, we leveraged partial redundancy between homoeologs in polyploid wheat to
recover *met1-1* mutants. Similar to other plants with large genomes, complete null mutants
were lethal. However, wheat plants with four null alleles (i.e. retaining two functional alleles)
develop normally, have substantially reduced CG methylation and occasionally exhibit novel
heritable phenotypes. Generating partial mutants may also be effective in other polyploid
species to manipulate CG methylation without lethal consequences, opening new avenues to
understand the role of CG methylation in polyploid species and generate epialleles for crop
breeding.

Methods

Plant materials and growth

Exome capture sequenced EMS mutagenized populations of hexaploid (*Triticum aestivum* cv.
Cadenza) and tetraploid wheat (*Triticum turgidum* cv. Kronos) were screened for premature
termination codon mutations in *MET1-1* (*TraesCS2A02G235900*, *TraesCS2B02G260800* and
TraesCS2D02G241800) predicted to eliminate C5 Cytosine methyltransferase domain
activity (Figure 1C) (Krasileva *et al.*, 2017).

Two independent Cadenza *MET1-1* populations were developed. The C0465xC0884
population was developed by crossing the double *met1-1* mutant Cadenza0465 with
Cadenza0884 to generate a triple heterozygous F_1, from which a segregating F_2 population
was produced (Supplementary Figure 2). The C0451xC2028xC1292 population was
developed by crossing Cadenza0451 with Cadenza2028 and Cadenza2028 with
Cadenza1292. The F_1 progeny were crossed to produce a triple heterozygous F_1 which was
self-pollinated to generate a segregating F_2 population (Supplementary Figure 2).

The Kronos *MET1-1* population was produced by crossing Kronos3085 with Kronos0809,
giving double heterozygous F_1 progeny, which were selfed to generate a segregating F_2
population. From the F_2 population, seeds from double heterozygous plants were sown to give
a segregating F_3 generation. F_4 seeds from double heterozygous F_3 plants were used for seed
genotyping (Supplementary Figure 3).

Seeds were germinated in petri dishes at 4°C for 48 hours, followed by room temperature for

48 hours before sowing into 96-cell trays containing John Innes F$_2$ starter + Grit (90% peat,

10% grit, 4kg m^{-3} dolomitic limestone, 1.2kg m^{-3} osmocote start). Selected plants were potted

on into 1L pots containing John Innes Cereal Mix (65% peat, 25% loam, 10% grit, 3kg m^{-3}

dolomitic limestone, 1.3kg m^{-3} pg mix, 3kg m^{-3} osmocote exact). Plants were grown in the

John Innes Centre glasshouses (Norwich, UK) with supplementary lighting and heating as

required for minimum 16h light with 16°C day and 14°C night.

Genotyping

Leaf

At the second leaf stage (Zadoks growth stage 12), DNA was extracted from 2cm leaf

samples following the protocol from www.wheat-training.com adapted from Pallotta *et al.*

(2003). Genotyping was carried out using KASP markers (Supplementary Table 6) and PACE

mix, according to the manufacturer's instructions. Kluster-Caller (v3.4.1.36) software was

used for data analysis. We developed a CAPS marker for C1292 (Supplementary Table 6),

which was amplified using Taq DNA Polymerase (New England Biolabs, 94°C for 2:00, 40

cycles of 94°C for 0:15, 50°C for 0:25, 72°C for 1:00 ending with 2:30 final extension at

72°C). The product was digested by restriction enzyme PstI-HF (New England Biolabs,

37°C, 1 hour). Homozygous *MET1-1* plants produced 3 digestion products (94bp, 157bp,

249bp) and homozygous *met1-1* mutants produced 2 digestion products (94bp and 406bp).

Grain

Grains were surface sterilised and cut transversely. The half of the grain containing the

embryo was germinated and the other half (containing endosperm) was used for genotyping.

Full details are available at dx.doi.org/10.17504/protocols.io.6qpvr84xplmk/v1. Parental

donation of the *met1-1* allele was determined using KASP genotyping, where the usual single

heterozygous cluster was split into two, corresponding to maternal or paternal donation of the

mutant *met1-1* allele. This method was validated using reciprocal crosses between *Rht-B13b*

(semi-dwarf) and wild type plants using KASP markers from (Borrill *et al.*, 2022).

Phenotyping

Grain size measurements

Grains were graded into size categories using 3D printed sieves with apertures from 1.0mm – 3.0mm increasing by 0.5mm increments The sieve design is available at https://www.printables.com/model/890045-grain-sieve. Following grain size grading, grain genotyping was carried out. The effect of total number of mutant *met1-1* alleles and inheritance of an abd gamete on grain size was tested using an ordered logistic regression model with the MASS package in R (v4.2.2).

Pollen Assays

Pollen grains per anther and pollen grain diameter were analysed using the Multisizer 4e Coulter Counter (Beckman Coulter, California, USA), based on the method described by (Alabdullah *et al.*, 2021). Pollen viability was analysed using the Ampha Z32 Pollen Analyzer (Amphasys, Lucerne, Switzerland). To determine the number of nuclei per pollen grain, pollen grains were stained with DAPI (4',6-diamidine-2-phenylindole) and imaged. Full method details can be found in the Supplementary Information.

Flowering Time

A late-flowering plant from the C0451xC2028xC1292 F_2 population was selfed. The progeny were genotyped and the date of complete emergence of the primary spike (Zadok's growth stage Z59) recorded. Plant height and the length of the primary spike were measured. The number of fully formed spikelets on the primary spike were counted.

Awn Development

An awned plant from the C0451xC2028xC1292 F_2 population was crossed to Cadenza WT. BC_1F_1 plants were grown and selfed.

215 segregating BC_1F_2 plants were grown and scored for the presence of awns as previously described (Huang *et al.*, 2020). Fully awned individuals had awns >1cm throughout the spike whereas awnletted plants had awns <1cm in the mid and basal portions of the spike while the length of awns at the apex reached up to 3cm. We scored any plants that had awns <1cm across the entire spike as awnless. We investigated the *Tipped1* (*B1*) gene (*TraesCS5A02G542800*), a characterised awn suppression gene, using published PCR and KASP markers (Huang *et al.*, 2020).

Reciprocal crossing

To test the transmissibility of *met1-1* mutant alleles in each of the parental gametes, we made reciprocal crosses between Cadenza WT and triple heterozygous plants (AaBbDd) from the C0465xC0884 population. The seeds of these crosses were harvested and grown. Once the seedlings reached the two-leaf stage (Zadoks growth stage Z12) leaf genotyping was carried out.

Whole Genome Bisulfite Sequencing

We collected leaf samples from the fourth leaf at Zadok's growth stage Z15.6/23, or from the sixth leaf at Z16 for the Aabbdd mutant. Samples were frozen in liquid nitrogen and stored at -70°C. DNA was extracted using the DNeasy Plant Mini Kit (Qiagen, Hilden, Germany) from three biological replicates per genotype, excluding the Aabbdd genotype, of which only one plant was recovered. DNA concentration was measured using an Invitrogen Qubit 4 fluorometer (Thermo Fisher Scientific, Massachusetts, USA) and the three replicates per genotype were pooled in equimolar ratios. Library preparation, bisulfite conversion, and sequencing by Illumina NovaSeq 6000 were performed by BMKgene (Münster, Germany), to obtain an average of 2.62 billion 150 bp paired-end reads per sample. Bisulfite conversion rates were calculated by BMKgene using the Bismark software (Krueger & Andrews, 2011), and rates between 99.52% and 99.70% were obtained. Fastp (Chen *et al.*, 2018) was used to remove adapter sequences, low quality reads, and polyG sequences. The Bismark Bisulfite Mapper (Krueger & Andrews, 2011) was used to process the sequencing data. Reads were aligned to a bisulfite-converted version of the IWGSC RefSeq v1.0 reference genome (IWGSC *et al.*, 2018). The average mapping efficiency was 40.49%, resulting in average coverage of 11X. Methylation calls in all sequence contexts (CpG, CHG, CHH) were extracted. Average cytosine methylation across genes and transposable elements was plotted using deepTools (Ramírez *et al.*, 2016) using GTF files generated using gffread (Pertea & Pertea, 2020) from previously published gff3 files (IWGSC *et al.*, 2018; Wicker *et al.*, 2018). Only cytosines with at least 4 reads were included in the analysis. Methylation profiles across chromosomes were calculated using computeMethylationProfile from DMRCaller (Catoni *et al.*, 2018), with bins of 1 Mb. Partitions between chromosome segments (R1, R2a, C, R2b, and R3) were assigned based on the segments defined in (IWGSC *et al.*, 2018), available at https://opendata.earlham.ac.uk/wheat/under_license/toronto/Ramirez-Gonzalez_etal_2018-06025-Transcriptome-Landscape/data/TablesForExploration/. Differentially methylated

regions (DMRs) in the CG context were called for each chromosome separately using computeDMRs from DMRCaller (Catoni *et al.*, 2018), specifying the method as 'bins', binSize as 100 bp, pValueThreshold as 0.01, minCytosinesCount as 10, minProportionDifference as 0.4, minGap as 0, and minReadsPerCytosine as 4. DMRs associated with genomic features were identified using bedtools intersect (Quinlan & Hall, 2010) to find overlapping genomic coordinates between previously annotated high-confidence genes (including 1 kb up- and downstream of the coding sequence) (IWGSC *et al.*, 2018) and transposable elements (Wicker *et al.*, 2018).

RNA-Sequencing

We collected leaf samples from the fourth leaf at Zadok's growth stage Z15.6/23, or from the sixth leaf at Z16 for the Aabbdd mutant. The samples were frozen in liquid nitrogen and stored at -70°C. RNA was extracted using trizol-chloroform and the RNA Clean & Concentrator Kit (Zymo Research). RNA was extracted from three biological replicates per genotype, excluding the Aabbdd genotype. Library preparation and sequencing by Illumina NovaSeq 6000 was performed by BMKgene (Münster, Germany), to obtain 150 bp paired-end reads. Fastp (Chen *et al.*, 2018) was used to remove adapter sequences, low quality reads, and polyG sequences. Reads were pseudoaligned to the IWGSC RefSeq v1.1 reference transcriptome (IWGSC *et al.*, 2018) and quantified using kallisto (Bray *et al.*, 2016). The output from kallisto was imported into R (v4.3.2) using tximport (Soneson *et al.*, 2015). Differentially expressed genes between genotypes were identified using DESeq2, using an adjusted p-value threshold of <0.01 and ≥ 2 fold change (Love *et al.*, 2014). For TE expression, reads were mapped to the IWGSC reference sequence using HISAT2 (Pertea *et al.*, 2016) and reads uniquely mapping to complete TE positions (Wicker *et al.*, 2018) were counted using HTSeq (Anders *et al.*, 2015), with default parameters. Differential expression analysis was carried out as for genes.

Gene expression data across tissues was downloaded as tpm values from https://bar.utoronto.ca/efp_wheat/cgi-bin/efpWeb.cgi (Choulet *et al.*, 2014; Borrill *et al.*, 2016; Ramírez-González *et al.*, 2018).

Data availability

Raw reads from whole genome bisulfite sequencing have been deposited in the European Nucletoide Archive under project PRJEB77426 and reads from RNA-sequencing have been deposited in project PRJEB77425.

All scripts used for whole genome bisulfite sequencing and RNA-sequencing analyses can be found on GitHub at https://github.com/Borrill-Lab/Wheat_MET1/ .

Funding

This work was supported by the UK Biotechnology and Biological Science Research Council (BBSRC) through grant BB/T013524/2 and the Institute Strategic Programmes Delivering Sustainable Wheat (DSW) (BB/X011003/1) and Building Robustness in Crops (BRiC) (BB/X01102X/1). DD was funded through a BBSRC Doctoral Training Partnership studentship (BB/T008717/1). GDS was funded by a scholarship from Sant'Anna School of Advanced Studies, Pisa.

Author contributions

MC and PB conceived the study. SB, DD and PB designed the research. JS and SB generated the mutant populations. SB and DD performed most experiments and statistical analysis. DD carried out the data analysis for the RNA-seq and whole genome bisulfite sequencing. JC assisted with genotyping and phenotyping of awn development. GDS helped with genotyping and carried out pollen phenotyping experiments. DD analysed the pollen phenotyping data. SB, DD and PB created the figures. SB, DD and PB wrote the manuscript with input from all authors. All authors have read and approved the manuscript.

Acknowledgements

We thank Abdulkader Alabdullah for advice on pollen phenotyping, Marek Glombik for advice on genomic analysis and James Brown for advice on statistical analysis of grain genotyping data. We thank Phil Robinson from Scientific Photography at JIC for taking high-quality images of plants and spikes. This research was supported by the NBI Research Computing group through HPC resources.

Declaration of interests

The current affiliation for Joseph Crudgington is the University of Edinburgh, United Kingdom and the current affiliation for Giorgia Di Santolo is Sant'Anna School of Advanced Studies, Pisa, Italy. All other authors declare no competing interests.

References

Alabdullah AK, Moore G, Martín AC. 2021. A duplicated copy of the meiotic gene *ZIP4* preserves up to 50% pollen viability and grain number in polyploid wheat. *Biology-Basel* **10**: 290.

Anders S, Pyl PT, Huber W. 2015. HTSeq-a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**: 166-169.

Birchler JA, Veitia RA. 2012. Gene balance hypothesis: Connecting issues of dosage sensitivity across biological disciplines. *Proceedings of the National Academy of Sciences* **109**: 14746-14753.

Boden SA, McIntosh RA, Uauy C, Krattinger SG, Dubcovsky J, Rogers WJ, Xia XC, Badaeva ED, Bentley AR, Brown-Guedira G, et al. 2023. Updated guidelines for gene nomenclature in wheat. *Theoretical and Applied Genetics* **136**: 72.

Borrill P, Mago R, Xu T, Ford B, Williams SJ, Derkx A, Bovill WD, Hyles J, Bhatt D, Xia X, et al. 2022. An autoactive *NB-LRR* gene causes *Rht13* dwarfism in wheat. *Proceedings of the National Academy of Sciences* **119**: e2209875119.

Borrill P, Ramirez-Gonzalez R, Uauy C. 2016. expVIP: a Customizable RNA-seq data analysis and visualization platform *Plant Physiology* **170**: 2172-2186.

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology* **34**: 525-527.

Catoni M, Cortijo S 2018. Chapter Four - EpiRILs: Lessons From Arabidopsis. In: Mirouze M, Bucher E, Gallusci P eds. *Advances in Botanical Research*: Academic Press, 87-116.

Catoni M, Tsang JM, Greco AP, Zabet NR. 2018. DMRcaller: a versatile R/Bioconductor package for detection and visualization of differentially methylated regions in CpG and non-CpG contexts. *Nucleic Acids Research* **46**: e114.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**: i884-i890.

Choulet F, Alberti A, Theil S, Glover N, Barbe V, Daron J, Pingault L, Sourdille P, Couloux A, Paux E, et al. 2014. Structural and functional partitioning of bread wheat chromosome 3B. *Science* **345**: 1249721.

Cokus SJ, Feng S, Zhang X, Chen Z, Merriman B, Haudenschild CD, Pradhan S, Nelson SF, Pellegrini M, Jacobsen SE. 2008. Shotgun bisulphite sequencing of the Arabidopsis genome reveals DNA methylation patterning. *Nature* **452**: 215-219.

Cubas P, Vincent C, Coen E. 1999. An epigenetic mutation responsible for natural variation in floral symmetry. *Nature* **401**: 157-161.

Domb K, Katz A, Harris KD, Yaari R, Kaisler E, Nguyen VH, Hong UVT, Griess O, Heskiau KG, Ohad N, et al. 2020. DNA methylation mutants in *Physcomitrella patens* elucidate individual roles of CG and non-CG methylation in genome regulation. *Proceedings of the National Academy of Sciences* **117**: 33700-33710.

Erdmann RM, Picard CL. 2020. RNA-directed DNA Methylation. *PLOS Genetics* **16**: e1009034.

Feng S, Cokus SJ, Zhang X, Chen PY, Bostick M, Goll MG, Hetzel J, Jain J, Strauss SH, Halpern ME, et al. 2010. Conservation and divergence of methylation patterning in plants and animals. *Proceedings of the National Academy of Sciences* **107**: 8689-8694.

Finnegan EJ, Ford B, Wallace X, Pettolino F, Griffin PT, Schmitz RJ, Zhang P, Barrero JM, Hayden MJ, Boden SA, et al. 2019. Zebularine treatment is associated with deletion of *FT-B1* leading to an increase in spikelet number in bread wheat *Plant Cell and Environment* **41**: 1346–1360.

He L, Wu W, Zinta G, Yang L, Wang D, Liu R, Zhang H, Zheng Z, Huang H, Zhang Q, et al.
2018. A naturally occurring epiallele associates with leaf senescence and local climate
adaptation in *Arabidopsis* accessions. *Nature Communications* **9**: 460.
He XJ, Hsu YF, Pontes O, Zhu JH, Lu J, Bressan RA, Pikaard C, Wang CS, Zhu JK. 2009.
NRPD4, a protein related to the RPB4 subunit of RNA polymerase II, is a component of RNA
polymerases IV and V and is required for RNA-directed DNA methylation. *Genes &*
Development **23**: 318-330.
Hu L, Li N, Xu C, Zhong S, Lin X, Yang J, Zhou T, Yuliang A, Wu Y, Chen Y-R, et al. 2014.
Mutation of a major CG methylase in rice causes genome-wide hypomethylation,
dysregulated genome expression, and seedling lethality. *Proceedings of the National Academy*
of Sciences **111**: 10642-10647.
Huang D, Zheng Q, Melchkart T, Bekkaoui Y, Konkin DJ, Kagale S, Martucci M, You FM,
Clarke M, Adamski NM. 2020. Dominant inhibition of awn development by a putative zinc-
finger transcriptional repressor expressed at the B1 locus in wheat. *New Phytologist* **225**: 340-
355.
IWGSC, Appels R, Eversole K, Stein N, Feuillet C, Keller B, Rogers J, Pozniak CJ, Choulet F,
Distelfeld A, et al. 2018. Shifting the limits in wheat research and breeding using a fully
annotated reference genome. *Science* **361**: eaar7191.
Johannes F, Porcher E, Teixeira FK, Saliba-Colombani V, Simon M, Agier N, Bulski A,
Albuisson J, Heredia F, Audigier P, et al. 2009. Assessing the impact of transgenerational
epigenetic variation on complex traits. *PLOS Genetics* **5**: e1000530.
Jullien PE, Kinoshita T, Ohad N, Berger F. 2006. Maintenance of DNA methylation during the
Arabidopsis life cycle is essential for parental imprinting. *The Plant Cell* **18**: 1360-1372.
Kankel MW, Ramsey DE, Stokes TL, Flowers SK, Haag JR, Jeddeloh JA, Riddle NC, Verbsky
ML, Richards EJ. 2003. Arabidopsis *MET1* cytosine methyltransferase mutants. *Genetics*
163: 1109-1122.
Kondrashov FA, Koonin EV. 2004. A common framework for understanding the origin of genetic
dominance and evolutionary fates of gene duplications. *Trends in Genetics* **20**: 287-290.
Krasileva KV, Vasquez-Gross HA, Howell T, Bailey P, Paraiso F, Clissold L, Simmonds J,
Ramirez-Gonzalez RH, Wang X, Borrill P, et al. 2017. Uncovering hidden variation in
polyploid wheat. *Proceedings of the National Academy of Sciences* **114**: E913-E921.
Krueger F, Andrews SR. 2011. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
applications. *Bioinformatics* **27**: 1571-1572.
Lei MG, Zhang HM, Julian R, Tang K, Xie SJ, Zhu JK. 2015. Regulatory link between DNA
methylation and active demethylation in. *Proceedings of the National Academy of Sciences of*
the United States of America **112**: 3553-3557.
Li D, Feng BE, Liu YJ, Gong J, Tang YM, Zhang LP, Pang BS, Sun RW, Zhang FT, Chen ZB, et
al. 2021. Genome-wide identification and transcriptional characterization of DNA
methyltransferases conferring temperature-sensitive male sterility in wheat. *BMC Genomics*
22: 310
Li J, Han F, Yuan T, Li W, Li Y, Wu HX, Wei H, Niu S. 2023. The methylation landscape of giga-
genome and the epigenetic timer of age in Chinese pine. *Nature Communications* **14**: 1947.
Li Q, Eichten SR, Hermanson PJ, Zaunbrecher VM, Song J, Wendt J, Rosenbaum H, Madzima
TF, Sloan AE, Huang J, et al. 2014. Genetic perturbation of the maize methylome. *The Plant*
Cell **26**: 4602-4616.
Liang W, Li J, Sun L, Liu Y, Lan Z, Qian W. 2022. Deciphering the synergistic and redundant roles
of CG and non-CG DNA methylation in plant development and transposable element
silencing. *New Phytologist* **233**: 722-737.
Liu X, Yu CW, Duan J, Luo M, Wang K, Tian G, Cui Y, Wu K. 2012. HDA6 directly interacts
with DNA methyltransferase MET1 and maintains transposable element silencing in
Arabidopsis. *Plant Physiol* **158**: 119-129.
Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biology* **15**: 550.

Manning K, Tör M, Poole M, Hong Y, Thompson AJ, King GJ, Giovannoni JJ, Seymour GB.
2006. A naturally occurring epigenetic mutation in a gene encoding an SBP-box transcription
factor inhibits tomato fruit ripening. *Nature Genetics* **38**: 948-952.
Mathieu O, Reinders J, Čaikovski M, Smathajitt C, Paszkowski J. 2007. Transgenerational
stability of the *Arabidopsis* epigenome is coordinated by CG methylation. *Cell* **130**: 851-862.
Mirouze M, Reinders J, Bucher E, Nishimura T, Schneeberger K, Ossowski S, Cao J, Weigel D,
Paszkowski J, Mathieu O. 2009. Selective epigenetic control of retrotransposition in. *Nature*
461: 427-U130.
Mirouze M, Vitte C. 2014. Transposable elements, a treasure trove to decipher epigenetic variation:
insights from Arabidopsis and crop epigenomes. *Journal of Experimental Botany* **65**: 2801-
2812.
Miura A, Yonebayashi S, Watanabe K, Toyama T, Shimada H, Kakutani T. 2001. Mobilization of
transposons by a mutation abolishing full DNA methylation in. *Nature* **411**: 212-214.
Miura K, Agetsuma M, Kitano H, Yoshimura A, Matsuoka M, Jacobsen SE, Ashikari M. 2009. A
metastable *DWARF1* epigenetic mutant affecting plant stature in rice. *Proceedings of the*
National Academy of Sciences **106**: 11218-11223.
Niederhuth CE, Bewick AJ, Ji L, Alabady MS, Kim KD, Li Q, Rohr NA, Rambani A, Burke
JM, Udall JA, et al. 2016. Widespread natural variation of DNA methylation within
angiosperms. *Genome Biology* **17**: 194.
Ortega-Galisteo AP, Morales-Ruiz T, Ariza RR, Roldán-Arjona T. 2008. Arabidopsis DEMETER-
LIKE proteins DML2 and DML3 are required for appropriate distribution of DNA
methylation marks. *Plant Molecular Biology* **67**: 671-681.
Pallotta M, Warner P, Fox R, Kuchel H, Jefferies S, Langridge P 2003. Marker assisted wheat
breeding in the southern region of Australia. *Proceedings of the Tenth International Wheat*
Genetics Symposium: Istituto Sperimentale per la Cerealicoltura Puglia, Italy. 789-791.
Pertea G, Pertea M. 2020. GFF Utilities: GffRead and GffCompare. *F1000Res* **9**.
Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. 2016. Transcript-level expression analysis of
RNA-seq experiments with HISAT, StringTie and Ballgown. *Nature Protocols* **11**: 1650-1667.
Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26**: 841-842.
Ramírez-González R, Borrill P, Lang D, Harrington S, Brinton J, Venturini L, Davey M, Jacobs
J, Van Ex F, Pasha A. 2018. The transcriptional landscape of polyploid wheat. *Science* **361**:
eaar6089.
Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke
T. 2016. deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic*
Acids Research **44**: W160-W165.
Reinders J, Wulff BB, Mirouze M, Marí-Ordóñez A, Dapp M, Rozhon W, Bucher E, Theiler G,
Paszkowski J. 2009. Compromised stability of DNA methylation and transposon
immobilization in mosaic *Arabidopsis* epigenomes. *Genes & Development* **23**: 939-950.
Saze H, Scheid OM, Paszkowski J. 2003. Maintenance of CpG methylation is essential for
epigenetic inheritance during plant gametogenesis. *Nature Genetics* **34**: 65-69.
Schmidt A, Wöhrmann HJP, Raissig MT, Arand J, Gheyselinck J, Gagliardini V, Heichinger C,
Walter J, Grossniklaus U. 2013. The polycomb group protein MEDEA and the DNA
methyltransferase MET1 interact to repress autonomous endosperm development in
Arabidopsis. *The Plant Journal* **73**: 776-787.
Schreiber M, Barakate A, Uzrek N, Macaulay M, Sourdille A, Morris J, Hedley PE, Ramsay L,
Waugh R. 2019. A highly mutagenised barley (cv. Golden Promise) TILLING population
coupled with strategies for screening-by-sequencing. *Plant Methods* **15**: 99.
Soneson C, Love MI, Robinson MD. 2015. Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. *F1000Res* **4**: 1521.
Springer NM, Schmitz RJ. 2017. Exploiting induced and natural epigenetic variation for crop
improvement. *Nature Reviews Genetics* **18**: 563-575.
Srikant T, Tri Wibowo A. 2021. The underlying nature of epigenetic variation: origin, establishment,
and regulatory function of plant epialleles. *International Journal of Molecular Sciences* **22**:
8618.

Srikant T, Yuan W, Berendzen KW, Contreras-Garrido A, Drost H-G, Schwab R, Weigel D.
2022. Canalization of genome-wide transcriptional activity in *Arabidopsis thaliana*
accessions by MET1-dependent CG methylation. *Genome Biology* 23: 263.

Tariq M, Saze H, Probst AV, Lichota J, Habu Y, Paszkowski J. 2003. Erasure of CpG methylation
in *Arabidopsis* alters patterns of histone H3 methylation in heterochromatin. *Proceedings of
the National Academy of Sciences of the United States of America* 100: 8823-8827.

Thomas M, Pingault L, Poulet A, Duarte J, Throude M, Faure S, Pichon JP, Paux E, Probst AV,
Tatout C. 2014. Evolutionary history of Methyltransferase 1 genes in hexaploid wheat. *BMC
Genomics* 15: 922.

Thorstensen T, Fischer A, Sandvik SV, Johnsen SS, Grini PE, Reuter G, Aalen RB. 2006. The
SUVR4 protein is a nucleolar histone methyltransferase with preference for monomethylated
H3K9. *Nucleic Acids Research* 34: 5461-5470.

Tsukahara S, Kobayashi A, Kawabe A, Mathieu O, Miura A, Kakutani T. 2009. Bursts of
retrotransposition reproduced in. *Nature* 461: 423-U125.

Veitia RA, Birchler JA. 2010. Dominance and gene dosage balance in health and disease: why levels
matter! *The Journal of Pathology* 220: 174-185.

Wang W, Simmonds J, Pan Q, Davidson D, He F, Battal A, Akhunova A, Trick HN, Uauy C,
Akhunov E. 2018. Gene editing and mutagenesis reveal inter-cultivar differences and
additivity in the contribution of TaGW2 homoeologues to grain size and weight in wheat.
Theoretical and Applied Genetics 131: 2463-2475.

Wicker T, Gundlach H, Spannagl M, Uauy C, Borrill P, Ramirez-Gonzalez RH, De Oliveira R,
International Wheat Genome Sequencing C, Mayer KFX, Paux E, et al. 2018. Impact of
transposable elements on genome structure and evolution in bread wheat. *Genome Biology*
19: 103.

Xiao W, Custard KD, Brown RC, Lemmon BE, Harada JJ, Goldberg RB, Fischer RL. 2006.
DNA methylation is critical for *Arabidopsis* embryogenesis and seed viability. *The Plant Cell*
18: 805-814.

Yamauchi T, Johzuka-Hisatomi Y, Terada R, Nakamura I, Iida S. 2014. The *MET1b* gene
encoding a maintenance DNA methyltransferase is indispensable for normal development in
rice. *Plant Molecular Biology* 85: 219-232.

Yang Y, Tang K, Datsenka TU, Liu W, Lv S, Lang Z, Wang X, Gao J, Wang W, Nie W, et al.
2019. Critical function of DNA methyltransferase 1 in tomato development and regulation of
the DNA methylome and transcriptome. *Journal of Integrative Plant Biology* 61: 1224-1242.

Zhang H, Lang Z, Zhu J-K. 2018. Dynamics and function of DNA methylation in plants. *Nature
Reviews Molecular Cell Biology* 19: 489-506.

Zhang P, Mbodj A, Soundiramourtty A, Llauro C, Ghesquière A, Ingouff M, Keith Slotkin R,
Pontvianne F, Catoni M, Mirouze M. 2023. Extrachromosomal circular DNA and structural
variants highlight genome instability in Arabidopsis epigenetic mutants. *Nature
Communications* 14: 5236.

Zhang Y-Y, Fischer M, Colot V, Bossdorf O. 2013. Epigenetic variation creates potential for
evolution of plant phenotypic plasticity. *New Phytologist* 197: 314-322.