Statistical learning beyond words in human neonates

Ana Fló[1,2*], Lucas Benjamin[1,3,4], Marie Palu[1], and Ghislaine Dehaene-Lambertz[1]

[1]Cognitive Neuroimaging Unit, CNRS ERL 9003, INSERM U992, CEA, Université Paris-Saclay, NeuroSpin center, Gif/Yvette, France.

[2]Department of Developmental Psychology and Socialisation and Department of Neuroscience, University of Padova, Padova, Italy

[3]Departement d'étude Cognitives, École Normale Supérieure, Paris, France

[4]Aix Marseille Univ, INSERM, INS, Inst Neurosci syst, Marseille, France

* Corresponding author

Email: ana.flo@unipd.it (AF)

ABSTRACT

Interest in statistical learning in developmental studies stems from the observation that 8-month-olds were able to extract words from a monotone speech stream solely using the transition probabilities (TP) between syllables (Saffran et al., 1996). A simple mechanism was thus part of the human infant's toolbox for discovering regularities in language. Since this seminal study, observations on statistical learning capabilities have multiplied across domains and species, challenging the hypothesis of a dedicated mechanism for language acquisition. Here, we leverage the two dimensions conveyed by speech –speaker identity and phonemes– to examine (1) whether neonates can compute TPs on one dimension despite irrelevant variation on the other and (2) whether the linguistic dimension enjoys an advantage over the voice dimension. In two experiments, we exposed neonates to artificial speech streams constructed by concatenating syllables while recording EEG. The sequence had a statistical structure based either on the phonetic content, while the voices varied randomly (Experiment 1) or on voices with random phonetic content (Experiment 2). After familiarisation, neonates heard isolated duplets adhering, or not, to the structure they were familiarised with. In both experiments, we observed neural entrainment at the frequency of the regularity and distinct Event-Related Potentials (ERP) to correct and incorrect duplets, highlighting the universality of statistical learning mechanisms and suggesting it operates on virtually any dimension the input is factorised. However, only linguistic duplets elicited a specific ERP component consistent with an N400, suggesting a lexical stage triggered by phonetic regularities already at birth. These results show that, from birth, multiple input regularities can be processed in parallel and feed different higher-order networks.

KEYWORDS

Neonates, statistical learning, neural entrainment, speech, phonemes, voices, ERP, language, steady-state, frequency tagging

HIGHLIGHTS

- Human neonates are sensitive to regularities in speech, encompassing both phonetic content and voice dimension.
- There is no observed advantage for linguistic content over voice content
- Both speech dimensions are processed in parallel by distinct networks.
- Only phonetic regularities evoked an N400, suggesting activations along the pathway to the lexicon.

INTRODUCTION

Since speech is a continuous signal, one of the infants' first challenges during language acquisition is to break it down into smaller units, notably to be able to extract words. Parsing has been shown to rely on prosodic cues (e.g., pitch and duration changes) but also on identifying regular patterns across perceptual units. Almost 20 years ago, Saffran, Newport, and Aslin (1996) demonstrated that infants are sensitive to local regularities between syllables. After hearing a stream of continuous and monotonous syllables constructed by concatenating four tri-syllabic pseudo-words, 8-month-old infants distinguished a list of these triplets from a list of triplets formed by the first part of one pseudo-word and the last part of another (called part-words). Indeed, for the correct triplets (called words), the TP between syllables was 1, whereas it drops to 1/3 for the transition encompassing two words present in the part-words. Since this seminal study, statistical learning has been regarded as an essential mechanism for language acquisition because it allows for the extraction of regular patterns without prior knowledge.

During the last two decades, many studies have extended this finding by demonstrating sensitivity to statistical regularities in sequences across domains and species. For example, segmentation capacities analogous to those observed for a syllable stream are observed throughout life in the auditory modality for tones (Kudo et al., 2011; Saffran et al., 1999) and in the visual domain for shapes (Bulf et al., 2011; Fiser and Aslin, 2002; Kirkham et al., 2002) and actions (Baldwin et al., 2008; Monroy et al., 2017). Non-human animals, such as cotton-top tamarins (Hauser et al., 2001), rats (Toro and Trobalón, 2005), dogs (Boros et al., 2021), and chicks (Santolin et al., 2016) are also sensitive to TPs. While the level of complexity that each species can track might differ, statistical learning between events appears as a general learning

mechanism for auditory and visual sequence processing (for a review of statistical learning

capacities across species, see (Santolin and Saffran, 2018)).

Using near-infra-red spectroscopy (NIRS) and electroencephalography (EEG), we have shown

that statistical learning is observed in sleeping neonates (Flo et al., 2022; Fló et al., 2019),

highlighting the automaticity of this mechanism. We also discovered that tracking statistical

probabilities might not lead to stream segmentation in the case of quadrisyllabic words in both

neonates and adults, revealing an unsuspected limitation of this mechanism (Benjamin et al.,

2022). Here, we aimed to further characterise the characteristics of this mechanism in order to

shed light on its role in the early stages of language acquisition. In particular, we wanted to

clarify whether, in human neonates, statistical learning is a general learning mechanism

applicable to any speech feature or whether there is a bias in favour of calculations on linguistic

content to extract words; and secondly, at what level infant newborns compute transitions

between syllables, at a low auditory level, i.e. between the presented events or later in the

processing chain, at the phonetic level after normalisation through an irrelevant dimension such

as voices.

We have, therefore, taken advantage of the fact that syllables convey two important pieces of

information for humans: what is being said and who is speaking, i.e. linguistic information and

information about the speaker's identity. While statistical learning can be helpful to word

extraction, a statistical relationship between successive voices is not of obvious use and could

even hinder word extraction if instances of a word uttered by a different speaker are considered

independently. However, as auditory processing is organised along several hierarchical and

parallel pathways integrating different spectro-temporal dimensions (Belin et al., 2000; DeWitt

and Rauschecker, 2012; Norman-Haignere et al., 2015; Zatorre and Belin, 2001), statistical

learning might be computed on one dimension independently of the variation of the other along

the linguistic and the voice pathways in parallel. Given the numerous behavioural and brain

imaging studies showing phonetic normalisation across speakers in infants (Dehaene-Lambertz

and Pena, 2001; Gennari et al., 2021; Kuhl and Miller, 1982), as also the statistical learning

studies in the second half of the first year using different voices (Estes and Lew-Williams, 2015)

and natural speech in which there is already variations of production from one instance to

another (Hay et al., 2011; Pelucchi et al., 2015), we were expecting that even if each syllable is

produced by a different speaker, TPs between syllables would be computed after a normalisation

process at the syllable or phonetic level, even in neonates. The predictions for learning the TPs

between different voices were more open. Either statistical learning is universal and can be

similarly computed over any dimension comprising voices, or listening to a speech stream

favours the processing of phonetic regularities over other non-linguistic dimensions of speech

and thus hinders the possibility of computing regularities over voices.

To study these possibilities, we constructed artificial streams based on the random concatenation

of three bi-syllabic pseudo-words (duplets). To build the duplets, we used six consonant-vowel

(CV) syllables produced by six voices (Table S1 and S2), resulting in 36 possible tokens. To

form the streams, tokens were combined either by considering their phonetic content

(Experiment 1: Structure over Phonemes) or their voice content (Experiment 2: Structure over

Voices), while the second dimension varied randomly (Figure 1). For example, in Experiment 1,

one duplet could be *petu* with *pe* and *tu* uttered by a random voice each time. In contrast, in

Experiment 2, one duplet could be the combination [*yellow* voice- *red* voice], each uttering

randomly any of the syllables.

If infants at birth compute regularities on the pure auditory signal, this implies computing the TPs over the 36 tokens. Thus, they should compute a 36×36 TPs matrix relating each acoustic signal, with TPs alternating between 1/6 within words and 1/12 between words. With this type of computation, we predict infants should fail the task in both experiments since previous studies showing successful segmentation in infants use high TP within words (usually 1) and much fewer elements (most studies 4 to 12) (Saffran and Kirkham, 2018) . If speech input is processed along the two studied dimensions in distinct pathways, it enables the calculation of two independent TP matrices of 6×6 between the six voices and six syllables. These computations would result in TPs alternating between 1 and 1/2 for the informative feature and uniform at 1/5 for the uninformative feature, leading to stream segmentation based on the informative dimension.

As in our previous experiments (Benjamin et al., 2022; Flo et al., 2022), we used high-density EEG (128 electrodes) to study speech segmentation abilities. Using artificial language with syllables with a fixed duration elicits Steady State Evoked Potentials (SSEP) at the syllable rate. Crucially, if the artificial language presents a regular structure between the syllables (i.e., regular drops in TPs marking word boundaries), if the structure is perceived, then the neural responses reflect the slower frequency of the word as well (Buiatti et al., 2009). In other words, the brain activity becomes phase-locked to the regular input, increasing the Inter Trial Coherence (ITC) and power at the input regularity frequencies. Under these circumstances, the analysis in the frequency domain is advantageous since fast and periodic responses can be easily investigated by looking at the target frequencies without considering their specific timing (Kabdebon et al., 2022). The phenomenon is also named frequency tagging or neural entrainment in the literature. Here, we will refer to it indistinctively as SSEP or neural entrainment since we do not aim to

make any hypothesis on the origin of the response (i.e., pure evoked or phase reset of endogenous oscillations (Giraud and Poeppel, 2012)). Our study used an orthogonal design across two groups of 1-4 day-old neonates. In Experiment 1 (34 infants), the regularities in the speech stream were based on the phonetic content, while the voices varied randomly (Phoneme group). Conversely, in Experiment 2 (33 infants), regularities were based on voices, while the phonemes changed randomly (Voice group). Both experiments started with a control stream in which both features varied randomly (i.e. Random stream, 120 s). Next, neonates were exposed to the Structured stream (120 s) with statistical structure over one or the other feature. The experiments ended with ten sets of 18 test duplets presented in isolation, preceded by short Structured streams (30 s) to maintain learning (Figure 1). Half of the test duplets corresponded to familiar regularities (Words, TP =1), and the other half were duplets present in the language but which straddled a drop in TPs (Partwords, TP = 0.5).

To investigate online learning, we quantified the ITC as a measure of neural entrainment at the syllable (4 Hz) and word rate (2 Hz) during the presentation of the continuous streams. For the recall process, we compared ERPs to Word and Part-Word duplets. We also tested 57 adult participants in a comparable behavioural experiment to investigate adults' segmentation capacities under the same conditions.

RESULTS

Neural entrainment during the familiarisation phase.

To measure neural entrainment, we quantified the ITC in non-overlapping epochs of 7.5 s. We compared the studied frequency (syllabic rate 4 Hz or duplet rate 2 Hz) with the 12 adjacent frequency bins following the same methodology as in our previous studies.

For the Random streams, we observed significant entrainment at syllable rate (4Hz) over a broad

set of electrodes in both experiments ($p < 0.05$, FDR corrected) and no enhanced activity at the

duplet rate for any electrode ($p > 0.05$, FDR corrected). Concerning the Structured streams, ITC

increased at both the syllable and duplet rate ($p < 0.05$, FDR corrected) in both experiments

(Figure 2A, 2B). The duplet effect was localised over occipital and central-left electrodes in the

Phoneme group and over occipital and temporal-right electrodes in the Voice group.

We also directly compared the ITC at both frequencies of interest between the Random and

Structured conditions (Figure 3C). We found electrodes with significantly higher ITC at the

duplet rate during Structured streams than Random streams in both experiments ($p < 0.05$, FDR

corrected). We also found electrodes with higher entrainment at syllable rate during the

Structured than Random streams in both experiments ($p < 0.05$, FDR corrected). This effect

might result from stronger or more phase-locked responses to syllables over those electrodes

when the input is structured. While the first harmonic of the duplet rate that coincides with the

syllable rate could also contribute to this effect, this contribution is unlikely since the electrodes

differ from the electrodes, showing enhanced word-rate activity at 2 Hz.

Finally, we looked for an interaction effect between groups and conditions (Structured vs.

Random streams) (Figure 2C). A few electrodes show differential responses between groups,

reflecting the topographical differences observed in the previous analysis, notably the trend for

stronger ITC at 2 Hz over the left central electrodes for the Phoneme group compared to the

Voice group, but none survive multiple comparison corrections.

Learning time-course: To investigate the time course of the learning, we computed neural

entrainment at the duplet rate in sliding time windows of 2 minutes with a 1 s step across both

random and structured streams (Figure 2D). Notice that because the integration window was two

minutes long, the entrainment during the first minute of the structure stream included data from

the random stream. To test whether ITC at 2 Hz increased during long Structured familiarisation

(120 s), we fitted a Linear Mixed Model (LMM) with a fixed effect of time and random slopes

and interceptions for individual subjects: $ITC \sim -1 + time + (1 + time|subject)$. In the

Phoneme group, we found a significant time effect (β=4.16×10^{-3}, 95% CI=[2.06×10^{-3}, 6.29×10^{-}

^{3}], SE=1.05×10^{-3}, p=4×10^{-4}), as well as in the Voice group (β=2.46×10^{-3}, 95% CI=[2.6×10^{-4},

4.66×10^{-3}], SE=1.09×10^{-3}, p=0.03). To test for differences in the time effect between groups, we

included all data in a single LMM: $ITC \sim -1 + time * group + (1 + time|subject)$. The

model showed a significant fixed effect of time for the Phoneme group consistent with the

previous results (β=4.22×10^{-3}, 95% CI=[1.07×10^{-3}, 7.37×10^{-3}], SE=1.58×10^{-3}, p=0.0096), while

the fixed effect estimating the difference between the Phoneme and Voice groups was not

significant (β=-1.84×10^{-3}, 95% CI=[-6.29×10^{-3}, 2.61×10^{-3}], SE=2.24×10^{-3}, p=0.4).

ERPs during the test phase.

To test the recall process, we also measured ERP to isolated duplets afterwards. The average

ERP to all conditions merged is shown in Figure S1. We investigated (1) the main effect of test

duplets (Word vs. Part-word) across both experiments, (2) the main effect of familiarisation

structure (Phoneme group vs. Voice group), and finally (3) the interaction between these two

factors. We used non-parametric cluster-based permutation analyses (i.e. without a priori ROIs)

(Oostenveld et al., 2011).

The difference between Word and Part-word consisted of a dipole with a median positivity and a

left temporal negativity ranging from 400 to 1500 ms, with a maximum around 800-900 ms (Fig

3). Cluster-based permutations recovered two significant clusters around 500-1500 ms: a frontal-

right positive cluster (p = 0.019) and a left temporal negative cluster (p = 0.0056). A difference

between groups was observed consisting of a dipole that started with a right temporal positivity

left temporo-occipital negativity around 300 ms and rotated anti-clockwise to bring the positivity

over the frontal electrodes and the negativity at the back of the head (500-800 ms) (Fig S2).

Cluster-based permutations on the Phoneme group vs. Voice group recovered a posterior cluster

(p = 0.018) around 500 ms; with no positive cluster reaching significance (p >0.10). A cluster-

based permutation analysis on the interaction effect, i.e., comparing Words - Part-Words

between both experiments, showed no significant clusters (p > 0.1).

As cluster-based statistics are not very sensitive, we also analysed the ERPs over seven ROIS

defined on the grand average ERP of all merged conditions (see Methods). Results replicated

what we observed with the cluster-based permutation analysis with similar differences between

Words and Part-words for the effect of familiarisation and no significant interactions. Results are

presented in SI. The temporal progression of voltage topographies for all ERPs is presented in

Figure S2. To verify that the effects were not driven by one group per duplet type condition, we

ran a mixed two-way ANOVA for the average activity in each ROI and significant time window,

with duplet type (Word/Part-word) as within-subjects factor and familiarisation as between-

subjects factor. We did not observe significant interactions in any case. Future studies should

consider a within-subject design to gain sensitivity to possible interaction effects.

Adult's behavioural performance in the same task.

Adult participants heard a Structure Learning stream lasting 120 s and then ten sets of 18 test

duplets preceded by Short Structure streams (30 s). For each test bigram, they had to rate their

familiarity with it on a scale from 1 to 6. For the group familiarised with the Phoneme structure, there was a significant difference between the score attributed to Words and Part-words ($t(26)=2.92$, $p = 0.007$, *Cohen's d =0.562*). The difference was marginally significant for the group familiarised with the Voice structure ($t(29)=2.0443$, $p = 0.050$, *Cohen's d=0.373*). A 2-way ANOVA with test-duplets and familiarisation as factors revealed a main effect of Word ($F(1,55)=12.52$, $p=0.0008$, $\eta_g^2=0.039$), no effect of familiarisation ($F(1,55) <1$), and a significant interaction Word × Familiarisation ($F(1,55) = 5.28$, $p= 0.025$, $\eta_g^2=0.017$).

DISCUSSION

Statistical learning is a general learning mechanism

In two experiments, we compared STATISTICAL LEARNING over a linguistic and a non-linguistic dimension in sleeping neonates. We took advantage of the possibility of constructing streams based on the same tokens, the only difference between the experiments being the arrangement of the tokens in the streams. We showed that neonates were sensitive to regularities based either on the phonetic or the voice dimensions of speech, even in the presence of a non-informative feature that must be disregarded.

Parsing based on statistical information was revealed by steady-state evoked potentials at the duplet rate observed around 2 min after the onset of the familiarisation stream and by different ERPs to Words and Part-words presented during the test in both experiments. Despite variations in the other dimension, statistical learning was possible, showing that this mechanism operates at a stage when these dimensions have already been separated along different processing pathways. Our results, thus, revealed that linguistic content and voice identity are calculated independently and in parallel. This result confirms that, even in newborns, the syllable is not a holistic unit

(Gennari et al., 2021) but that the rich temporo-frequential spectrum of speech is processed in

parallel along different networks, probably using different integration factors (Boemio et al.,

2005; Moerel et al., 2012; Zatorre and Belin, 2001) .

Second, we observed no obvious advantage for the linguistic dimension in neonates. These

results reveal the universality of statistical learning. While statistical learning has already been

described in many domains and species (Frost et al., 2015; Ren and Wang, 2023; Santolin and

Saffran, 2018), we add here that even sleeping neonates distinctly applied it to possibly all the

dimensions in which speech is factorised in the auditory cortex (Gennari et al., 2021; Gwilliams

et al., 2022). This mechanism gives them a powerful tool to create associations between

recurrent events.

Differences between statistical learning over voices and over phonemes

While the main pattern of results between experiments was comparable, we did observe some

differences. The word-rate steady-state response (2 Hz) for the group of infants exposed to

structure over phonemes was left lateralised over central electrodes, while the group of infants

hearing structure over voices showed mostly entrainment over right temporal electrodes. These

results are compatible with statistical learning in different lateralised neural networks for

processing speech's phonetic and voice content. Recent brain imaging studies on infants do

indeed show precursors of later networks with some hemispheric biases (Blasi et al., 2011;

Dehaene-Lambertz et al., 2010), even if specialisation increases during development (Shultz et

al., 2014; Sylvester et al., 2023). The hemispheric differences reported here should be considered

cautiously since the group comparison did not survive multiple comparison corrections. Future

work investigating the neural networks involved should implement a within-subject design to

gain statistical power.

The time course of the entrainment at the duplet rate revealed that entrainment emerged at a

similar time for both statistical structures. While this duplet rate response seemed more stable in

the Phoneme group (i.e., the ITC at the word rate was higher than zero in a sustained way only in

the Phoneme group, and the slope of the increase was steeper), no significant difference was

observed between groups. Since we did not observe group differences in the ERPs to Words and

Part-words during the test, it is unlikely that these differences during learning were due to a

worse computation of the statistical transitions for the voice stream relative to the phoneme

stream. An alternative explanation might be related to the nature of the duplet rate entrainment.

Entrainment might result either from a different response to low and high TPs or (and) from a

response to chunks in the stream (i.e., "Words"). In a previous study (Benjamin et al., 2022), we

showed that in some circumstances, neonates compute TPs, but entrainment does not emerge,

likely due to the absence of chunking. It is thus possible that chunking was less stable when the

regularity was over voices, consistent with the results of previous studies reporting challenges

with voice identification in infants as in adults (Johnson et al., 2011; Mahmoudzadeh et al.,

2016).

Phoneme regularities might trigger a lexical search

In the test part on isolated duplets, we also observed a significant difference between groups: A

dipole, consisting of a posterior negative pole and a frontal positivity, was observed around 500

ms following linguistic duplets but not voice duplets. Since the acoustic properties of the duplets

were the same in the two experiments, the difference can only be due to an endogenous process

that modulates stimuli processing. Given its topography, latency and the context in which it

appears, we hypothesise that this component is congruent with an N400, a component elicited by

lexico-semantic manipulations in adults (Kutas and Federmeier, 2011). As often in infants, the

component latency is delayed, and its topography is more posterior relative to later ages

(Friedrich and Friederici, 2005; Junge et al., 2021). A larger posterior negativity has been

reported in infants as young as five months when they hear their name compared to a stranger's

name (Parise et al., 2010), when a pseudo-word is consistent vs inconsistently associated with an

object (Friedrich and Friederici, 2011), and when they see unexpected vs expected actions (Reid

et al., 2009), suggesting that such negativity might be related to semantic.

There is also evidence that infants extract and store possible word-forms (Jusczyk and Hohne,

1997). A stronger fMRI activation for forward speech than backward speech in the left angular

gyrus in 3-mo-olds was assumed to be related to activations of possible word forms in a proto-

lexicon for native language sentences (Dehaene-Lambertz et al., 2002). As elegantly shown by

(Shukla et al., 2011), these chunks extracted from the speech stream are candidate words to

which meanings can be attached. These authors show that 6-month-olds spontaneously associate

a non-word extracted from natural sentences with a visual object. Although in their experiment,

infants used prosodic cues to extract the word, and here statistical cues, a spontaneous bias to use

possible word-forms as referring to a meaning (see also (Bergelson and Aslin, 2017) might

trigger activation along a lexicon pathway and explain the difference seen here between the two

groups. Only speech chunks based on phonetic regularities, and not voice regularities, would be

considered as possible candidate.

A similar interpretation of an N400 induced by possible words, even without a clear semantic,

explains the observation of an N400 in adult participants listening to artificial languages. Sanders

et al. (2002) observed an N400 in adults listening to an artificial language only when they were previously exposed to the isolated pseudo-words. Other studies reported larger N400 amplitudes when adult participants listened to a structured stream compared to a random sequence of syllables (Cunillera et al., 2009, 2006), tones (Abla et al., 2008), and shapes (Abla and Okanoya, 2009). Our results show an N400 for both Words and Part-words in the post-learning phase, possibly related to a top-down effect induced by the familiarisation stream. Since computing ERPs during the streams has inherent baseline issues that become critical with young infants' EEG recordings due to the large amplitude of the slow waves at this age (Eisermann et al., 2013), we cannot perform the ERP analysis during the stream as in previous adult studies preventing a direct comparison. However, the component we observed for duplets presented after the familiarisation streams might result from a related phenomenon. Learning the phonetic structure, but not the voice sequence, might induce a lexical search during the presentation of the isolated duplets, as it happens in the Structured stream vs. Random streams in the previously mentioned adult studies (Abla et al., 2008; Abla and Okanoya, 2009; Cunillera et al., 2009, 2006). A lexical entry might also explain the more sustained activity during the familiarisation stream in the phonemes group, as the chunk might be encoded as a putative word in this admittedly rudimentary but present lexical store, and the neural entrainment reflects not only the TP but also the recovery of the "lexical" item.

Finally, we would like to point out that it is not natural for a word not to be produced by the same speaker, nor for speakers to have statistical relationships of the kind we used here. Neonates, who have little experience and therefore no (or few) expectations or constraints, are probably better revealers of the possibilities opened by statistical learning than older participants. In fact, adults obtained better results for phoneme structure than for voice structure, perhaps

because of an effective auditory normalisation process or the use of a writing code for phonemes

but not for voices. It is also possible that the difference between neonates and adults is related to

the behavioural test being a more explicit measure of word recognition than the implicit task

allowed by EEG recordings. In any case, results show that even adults displayed some learning

on the voice duplets.

Altogether, our results show that statistical learning works similarly on different speech features

with no clear advantage for computing linguistically relevant regularities in speech. This

supports the idea that statistical learning is a general and evolutionarily ancient learning

mechanism, probably operating on common computational principles but within different neural

networks with a different chain of operations: phonetic regularities induce a supplementary

component - not seen in the case of voice regularities- that we related to a lexical N400.

Understanding how statistical learning computations over linguistically relevant dimensions,

such as the phonetic content of speech, are extracted and passed to other networks might be

fundamental to understanding what enables the infant brain to acquire language. Further work is

needed to understand how extracting regularities over different features articulates the language

network.

ACKNOWLEDGEMENTS

353 We want to thank all the families who participated in the study. This research has received

354 funding from the European Research Council (ERC) under the European Union's Horizon 2020

355 research and innovation program (grant agreement No. 695710).

AUTHOR CONTRIBUTIONS

357 A.F. and G.D.L. conceptualised the research; A.F., L.B. and M.P. performed the research; A.F.

358 analysed the data; and A.F., L.B., and G.D.L. wrote the paper

359 **DECLARATION OF INTERESTS**

360 The authors declare no competing interests.

Figure 1. Experimental protocol.

The experiments started with a Random stream (120 s) in which both syllables and voices changed randomly, followed by a long-Structured stream (120 s). Then, 10 short familiarisation streams (30 s), each followed by test blocks comprising 18 isolated duplets (SOA 2-2.3 s) were presented. Example streams are presented to illustrate the construction of the streams, with different colours representing different voices. In Experiment 1, the Structured stream had a statistical structure based on phonemes (TPs alternated between 1 and 0.5), while the voices were randomly changing (uniform TPs of 0.2). For example, the two syllables of the word "petu" were produced by different voices, which randomly changed at each presentation of the word. In Experiment 2, the statistical structure was based on voices (TPs alternated between 1 and 0.5), while the syllables changed randomly (uniform TPs of 0.2). For example, the "green" voice was always followed by the "red" voice, but they were randomly saying different syllables "boda" then "tupe" in our example. The test duplets were either Words (TP=1) or Partwords (TP=0.5). Words and Partwords were defined in terms of phonetic content for Experiment 1 and voice content for Experiment 2.

376
Figure 2. Neural entrainment during the random and structured streams.

(A) SNR for the ITC during the Random and Structured streams of Experiment 1 (structure on phonetic content). The topographies represent the entrainment in the electrode space at the syllabic (4 Hz) and duplet rates (2 Hz). Crosses indicate the electrodes showing enhanced neural entrainment (cross: p <

0.05, one-sided paired permutation test, FDR corrected by the number of electrodes; dot: p < 0.05, without FDR correction). Colour scale limits [-1.8, 1.8]. The entrainment for each electrode is shown in light grey. The thick orange line shows the mean over the electrodes with significant entrainment relative to the adjacent frequency bins at the syllabic rate (4 Hz) (p < 0.05 FDR corrected). The thick green line shows the mean over the electrodes showing significant entrainment relative to the adjacent frequency bins at the duplet rate (2 Hz) (p < 0.05 FDR corrected). The asterisks indicate frequency bins with entrainment significantly higher than on adjacent frequency bins for the average across electrodes (p < 0.05, one-sided permutation test, FDR corrected for the number of frequency bins). (B) Analog to A for Experiment 2 (structure on voice content). (C) The first two rows show the topographies for the difference in entrainment during the Structured and Random streams at 4 Hz and 2 Hz for both experiments. Crosses indicate the electrodes showing stronger entrainment during the Structured stream (cross: p < 0.05, one-sided paired permutation test, FDR corrected by the number of electrodes; dot: p < 0.05, without FDR correction). The bottom row shows the interaction effect by comparing the difference in entrainment during the Structured and Random streams between Experiments 1 and 2. Crosses indicate significant differences (cross: p < 0.05, two-sided unpaired permutation test, FDR corrected by the number of electrodes; dot: p < 0.05, without FDR correction). (D) Time course of the neural entrainment at 4 Hz for the average over electrodes showing significant entrainment during the Random stream and at 2 Hz for the average over electrodes showing significant entrainment during the Structured stream (Phoneme: green line, Voice blue line). The shaded area represents standard errors. The horizontal lines on the bottom indicate when the entrainment was larger than 0 (p < 0.05, one-sided t-test, corrected by FDR by the number of time points).

Figure 3. Cluster-based permutation analysis of ERPs to isolated duplets during recall

The topographies show the difference between the two conditions corresponding to each main effect. Results obtained from the cluster-based permutation analyses are shown at the bottom of each panel. Thick lines correspond to the grand averages for the two main tested conditions. Shaded areas correspond to the standard error across participants. Thin lines show the ERPs separated by duplet type and familiarisation type. The shaded areas between the thick lines show the time extension of the cluster. The topographies correspond to the difference between conditions during the time extension of the cluster. The electrodes belonging to the cluster are marked with a cross. Significant clusters are indicated with an asterisk. Color scale limits [-0.07, 0.07] a.u. (A) Main effect of Test-duplets (Words - Part-words) over a frontal-right positive cluster (p = 0.019) and a left temporal negative cluster (p = 0.0056). (B) Main effect of familiarisation (Phonemes - Voices) over a posterior negative cluster (p = 0.018). The frontal positive cluster did not reach significance (p = 0.12). Results are highly comparable to the ROIs-based analysis presented in SI (Fig S3).

419 **Figure 4. Adults' behavioural experiment.**

420 Each subject's average score attributed to the Words (blue) and Partwords (orange) is represented. On
421 the right, for the group familiarised with the Phoneme structure and on the left, for the group familiarised
422 with the Voice structure. The difference between test duplets was significant for the Phoneme group (p =
423 *0.007*) and only marginally significant for the Voice group (p = *0.050*).

METHODS

Participants

Participants were healthy-full-term neonates with normal pregnancy and birth (GA > 38 weeks, Apgar scores ≥ 7/8 at 1/5 minute, birthweight > 2.5 Kg, cranial perimeter ≥ 33.0 cm), tested at the Port Royal Maternity (AP-HP), in Paris, France. Parents provided informed consent. The regional ethical committee for biomedical research (Comité de Protection des Personnes Region Centre Ouest 1, EudraCT/ID RCB: 2017-A00513-50) approved the protocol, and the study was carried out according to relevant guidelines and regulations. 67 participants (34 in Experiment 1 and 33 in Experiment 2) who provided enough data without motion artefacts were included (Experiment 1: 19 females; 1 to 4 days old; mean GA: 39.3 weeks; mean weight: 3387 g; Experiment 2: 15 females; 1 to 4 days old; mean GA: 39.0 weeks; mean weight: 3363 g). 12 other infants were excluded from the analyses (11 due to fussiness; 1 due to bad data quality).

Stimuli

The stimuli were synthesised using the MBROLA diphone database (Dutoit et al., 1996). Syllables had a consonant-vowel structure and lasted 250 ms (consonants 90 ms, vowels 160 ms). Six different syllables (*ki*, *da*, *pe*, *tu*, *bo*, *gɛ*) and six different voices were used (*fr3*, *fr1*, *fr7*, *fr2*, *it4*, *fr4*), resulting in a total of 36 syllable-voice combinations, from now on, tokens. The voices could be female or male and have three different pitch levels (low, middle, and high) (Table S1). The 36 tokens were synthesised independently in MBROLA, their intensity was normalised, and the first and last 5 ms were ramped to zero to avoid "clicks." The streams were synthesised by concatenating the tokens' audio files, and they were ramped up and down during the first and last 5 s to avoid the start and end of the stream serving as perceptual anchors.

The Structured streams were created by concatenating the tokens in such a way that they resulted

in a semi-random concatenation of the duplets (i.e., pseudo-words) formed by one of the features

(syllable/voice) while the other feature (voice/syllable) vary semi-randomly. In other words, in

Experiment 1, the order of the tokens was such that Transitional Probabilities (TPs) between

syllables alternated between 1 (within duplets) and 0.5 (between duplets), while between voices,

TPs were uniformly 0.2. The design was orthogonal for the Structured streams of Experiment 2

(i.e., TPs between voices alternated between 1 and 0.5, while between syllables were evenly 0.2).

The random streams were created by semi-randomly concatenating the 36 tokens to achieve

uniform TPs equal to 0.2 over both features. The semi-random concatenation implied that the

same element could not appear twice in a row, and the same two elements could not repeatedly

alternate more than two times (i.e., the sequence $X_kX_jX_kX_j$, where X_k and X_j are two elements, was

forbidden). Notice that with an element, we refer to a duplet when it concerns the choice of the

structured feature and to the identity of the second feature when it involves the other feature. The

same statistical structures were used for both Experiments, only changing over which dimension

the structure was applied. The learning stream lasted 120 seconds, with each duplet appearing 80

times. The 10 short structured streams lasted 30 seconds each, each duplet appearing a total of

200 times (10×20). The same random stream was used for both Experiments, and it lasted 120

seconds.

In Experiment 1, the duplets were created to prevent specific phonetic features from facilitating

stream segmentation. In each experiment, two different structured streams (lists A and B) were

used by modifying how the syllables/voices were combined to form the duplets (Table S2).

Crucially, the Words/duplets of list A are the Part-words of list B and vice versa any difference

between those two conditions can thus not be caused by acoustical differences. Participants were

randomly assigned and balanced between lists and Experiments.

The test words were duplets formed by the concatenation of two tokens, such that they formed a

Word or a Part-word according to the structured feature.

Procedure and data acquisition

Scalp electrophysiological activity was recorded using a 128-electrode net (Electrical Geodesics,

Inc.) referred to the vertex with a sampling frequency of 250 Hz. Neonates were tested in a

soundproof booth while sleeping or during quiet rest. The study involved: (1) 120 s of a random

stream, (2) 120 s of a structured stream, (3) 10 series of 30 s of structured streams followed by

18 test sequences (SOA 2-2.3s).

Data pre-processing

Data were band-pass filtered 0.1-40 Hz and pre-processed using custom MATLAB scripts based

on the EEGLAB toolbox 2021.0 according to the APICE pre-processing pipeline to recover as

much free-artifacts data as possible (Fló et al., 2022).

Neural entrainment

The pre-processed data were further high-pass filtered at 0.2 Hz. Then, data was segmented from

the beginning of each phase into 0.5 s long segments (240 duplets for the Random, 240 duplets

for the long Structured, and 600 duplets for the short Structured). Segments containing samples

with artefacts defined as bad data in more than 30% of the channels were rejected, and the

remaining channels with artefacts were spatially interpolated.

<u>Neural entrainment per condition.</u> The 0.5 s epochs belonging to the same condition were

reshaped into non-overlapping epochs (Benjamin et al., 2021) of 7.5 s (15 duplets, 30 syllables),

retaining the chronological order; thus, the timing of the steady-state response. Subjects who did

not provide at least 50% artifact-free epochs for each condition (at least 8 long epochs during

Random and 28 during Structured) were excluded from the entrainment analysis (32 included

subjects in Experiment 1, and 32 included subjects in Experiment 2). The retained subjects for

Experiment 1, on average provided 13.59 epochs for the Random condition (SD 2.07, range [8,

16]) and 48.16 for the Structured conditions (SD 5.89, range [33, 55]). The retained subjects for

Experiment 2, on average provided 13.78 epochs for the Random condition (SD 1.93, range [8,

16]) and 46.88 for the Structured conditions (SD 5.62, range [36, 55]). After data rejection, data

were referenced to the average and normalized by dividing by the standard deviation within an

epoch across electrodes and time. Next, data were converted to the frequency domain using the

Fast Fourier Transform (FFT) algorithm, and the ITC was estimated for each electrode during

each condition (Random, Structured) as $ITC(f) = \frac{1}{N}\left|\sum_{i=1}^{N} e^{i\varphi(f,i)}\right|$, where N is the number of

trials and φ(f,i) is the phase at frequency f and trial i. The ITC ranges from 0 to 1 (i.e.,

completely desynchronized activity to perfectly phased locked activity). Since we aim to detect

an increase in signal synchronization at specific frequencies, the SNR was computed relative to

the twelve adjacent frequency bins (six of each side corresponding to 0.8 Hz) (Kabdebon et al.,

2022). This procedure also enables correcting differences in the ITC due to a different number of

trials. Specifically, the SNR was *SNR(f) = (ITC(f)-mean(ITC$_{noise}$(f)))/std(ITC$_{noise}$(f))*, where

ITC$_{noise}$(f) is the ITC over the adjacent frequency bins. For statistical analysis, we compared the

SNR at syllable rate (4 Hz) and duplet rate (2 Hz) against the average SNR over the 12 adjacent

frequency bins using a one-tail paired permutation test (5000 permutations). We also directly

compared the entrainment during the two conditions to individuate the electrodes showing a greater entrainment during the Structured than Random streams. We evaluated the interaction between Stream type (Random and Structured) and Familiarization type (Structured over Phonemes or Voices) by comparing the difference in entrainment between Structured and Random during the two experiments using a two sides unpaired permutation test (5000 permutations). All P-values were corrected across electrodes by FDR.

Neural entrainment time course. The 0.5 s epochs were concatenated chronologically (2 minutes of Random, 2 minutes of long Structured stream, and 5 minutes of short Structured blocks). The same analysis as above was performed in sliding time windows of 2 minutes with a 1 s step. A time window was considered valid if at least 8 out of the 16 epochs were free of motion artefacts. Missing values due to the presence of motion artifacts where linearly interpolated. Then, the entrainment time course at the syllable rate was computed as the average over the electrodes showing significant entrainment at 4 Hz during the Random condition, and at the duplet rate, as the average over the electrodes showing significant entrainment at 2 Hz during the Structured condition. Finally, data was smooth over a time window of 30 s.

To investigate the increase in the neural activity locked to the regularity during the long familiarisation, we fitted a LMM for each group of subjects. We included time as a fixed effect and random slopes and interceptions for individual subjects: $ITC \sim -1 + time + (1 + time|subject)$. We then compare the time effect between groups by including all data in a single LMM with time and group as fixed effects: $ITC \sim -1 + time * group + (1 + time|subject)$.

ERPs to test words

The pre-processed data were filtered between 0.2 and 20 Hz, and epoched between [-0.2, 2.0] s

from the onset of the duplets. Epochs containing samples identified as artifacts by APICE

procedure were rejected. Subjects who did not provide at least half of the trials (45 trials) per

condition were excluded (34 subjects kept for Experiment 1, and 33 for Experiment 2). None

subjects were excluded based on this criteria in the Phonemes groups, and one subject was

excluded in the Voice groups. For Experiment 1, we retained on average 77.47 trials (SD 9.98,

range [52, 89]) for the Word condition and 77.12 trials (SD 10.04, range [56, 89]) for the

Partword condition. For Experiment 2, we retained on average 73.73 trials (SD 10.57, range [47,

90]) for the Word condition and 74.18 trials (SD 11.15, range [46, 90]) for the Partword

condition. Data were reference averaged and normalised within each epoch by dividing by the

standard deviation across electrodes and time.

Since the gran average response across both groups and conditions returned to the pre-stimulus

level at around 1500 ms, we defined [0, 1500] ms as time windows of analysis. We first analysed

the data using non-parametric cluster-based permutation analysis (Oostenveld et al., 2011) in the

time window [0, 1500] ms (alpha threshold for clustering 0.10, neighbour distance $\leq$ 2.5 cm,

clusters minimum size 3 and 5,000 permutations).

We also analysed the data in seven ROIs to ensure that no other effects were present that were

not caught by the cluster-based permutation analysis. By inspecting the grand average ERP

across both Experiments and conditions, we identified three characteristic topographies: (a)

positivity over central electrodes, (b) positivity over frontal electrodes and negativity over

occipital electrodes, and (c) positivity over prefrontal electrodes and negativity over temporal

electrodes (Figure S1). Then, we defined 7 symmetric ROIs: Central, Frontal Left, Frontal Right,

Occipital, Prefrontal, Temporal Left, Temporal Right. We evaluated the main effect of Test-word

type by comparing the EPRs between Words and Partwords (paired t-test) and the main effect of Familiarization type by comparing ERPs between Experiment 1 (structured over Phonemes) and Experiment 2 (structure over Voices) (unpaired t-test). All p-values were FDR corrected by the number of time points (n = 376) and ROIs (n = 7). To test for possible interaction effects, we compared the difference between Words and Partwords between the two groups. To verify that the main effects were not driven by one condition or group, we computed the average on each of the time windows where a main effect was identified considering both the Test-word type and Familiarization type factors, and we ran a two ways-ANOVA (Test-word type x Familiarization type). Results are presented in SI (Figure S3).

Adults' behavioural experiment

57 French-speaking adults were tested in an online experiment analogous to the infant study through the Prolific platform. All participants provided informed consent and received monetary compensation for their participation. The study was approved by the Ethical Research Committee of Paris Saclay University under the reference CER-Paris-Saclay-2019-063. The same stimuli as in the infants' experiment were used. Participants first heard 2 min of familiarisation with the Structured stream. Then, they completed ten sessions of re-familiarisation and testing. Each re-familiarization lasted 30 s, and in each test session, all 18 test words were presented. The structure could be either over the phonetic or the voice content, and two lists were used (see Table S2). Participants were randomly assigned to one of the groups and to one list. The Phonemes group included 27 participants, and the Voices group 30 participants. Before starting the experiment, subjects were instructed to pay attention to an invented language because later, they would have to answer if different sequences adhered to the structure of the

580   language. During the test phase, subjects were asked to scale their familiarity with each test word

581   by clicking with a cursor on a scale from 1 to 6.

REFERENCES

Abla D, Katahira K, Okanoya K. 2008. On-line assessment of statistical learning by event-related potentials. *Journal of Cognitive Neuroscience* **20**:952–64. doi:10.1162/jocn.2008.20058

Abla D, Okanoya K. 2009. Visual statistical learning of shape sequences: An ERP study. *Neuroscience Research* **64**:185–190. doi:10.1016/j.neures.2009.02.013

Baldwin D, Andersson A, Saffran J, Meyer M. 2008. Segmenting dynamic human action via statistical structure. *Cognition* **106**:1382–1407. doi:10.1016/j.cognition.2007.07.005

Belin P, Zatorre RJ, Lafaille P, Ahad P, Pike B. 2000. Voice-selective areas in human auditory cortex **403**.

Benjamin L, Dehaene-Lambertz G, Fló A. 2021. Remarks on the analysis of steady-state responses: spurious artifacts introduced by overlapping epochs. *Cortex*. doi:10.1016/j.cortex.2021.05.023

Benjamin L, Fló A, Palu M, Naik S, Melloni L, Dehaene-Lambertz G. 2022. Tracking transitional probabilities and segmenting auditory sequences are dissociable processes in adults and neonates. *Developmental Science* **n/a**:e13300. doi:10.1111/desc.13300

Bergelson E, Aslin RN. 2017. Nature and origins of the lexicon in 6-mo-olds. *Proc Natl Acad Sci USA* **114**:12916–12921. doi:10.1073/pnas.1712966114

Blasi A, Mercure E, Lloyd-Fox S, Thomson A, Brammer M, Sauter D, Deeley Q, Barker GJ, Renvall V, Deoni S, Gasston D, Williams SCR, Johnson MH, Simmons A, Murphy DGM. 2011. Early Specialization for Voice and Emotion Processing in the Infant Brain. *Current Biology* **21**:1220–1224. doi:10.1016/j.cub.2011.06.009

Boemio A, Fromm S, Braun A, Poeppel D. 2005. Hierarchical and asymmetric temporal sensitivity in human auditory cortices. *Nat Neurosci* **8**:389–395. doi:10.1038/nn1409

Boros M, Magyari L, Török D, Bozsik A, Deme A, Andics A. 2021. Neural processes underlying statistical learning for speech segmentation in dogs. *Current Biology*. doi:10.1016/j.cub.2021.10.017

Buiatti M, Peña M, Dehaene-Lambertz G. 2009. Investigating the neural correlates of continuous speech computation with frequency-tagged neuroelectric responses. *NeuroImage* **44**:509–519. doi:10.1016/j.neuroimage.2008.09.015

Bulf H, Johnson SP, Valenza E. 2011. Visual statistical learning in the newborn infant. *Cognition* **121**:127–132. doi:10.1016/j.cognition.2011.06.010

Cunillera T, Càmara E, Toro JM, Marco-Pallares J, Sebastián-Galles N, Ortiz H, Pujol J, Rodríguez-Fornells A. 2009. Time course and functional neuroanatomy of speech segmentation in adults. *NeuroImage* **48**:541–553. doi:10.1016/j.neuroimage.2009.06.069

Cunillera T, Toro JM, Sebasti??n-Gall??s N, Rodr??guez-Fornells A. 2006. The effects of stress and statistical cues on continuous speech segmentation: An event-related brain potential study. *Brain Research* **1123**:168–178. doi:10.1016/j.brainres.2006.09.046

Dehaene-Lambertz G, Dehaene S, Hertz-Pannier L. 2002. Functional Neuroimaging of Speech Perception in Infants. *Science* **298**:2013–2015. doi:10.1126/science.1077066

Dehaene-Lambertz G, Montavont A, Jobert A, Allirol L, Dubois J, Hertz-Pannier L, Dehaene S. 2010. Language or music, mother or Mozart? Structural and environmental influences on infants' language networks. *Brain and Language* **114**:53–65. doi:10.1016/j.bandl.2009.09.003

Dehaene-Lambertz G, Pena M. 2001. Electrophysiological evidence for automatic phonetic processing in neonates: *Neuroreport* **12**:3155–3158. doi:10.1097/00001756-200110080-00034

Delorme A, Makeig S. 2004. EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods* **134**:9–21. doi:10.1016/j.jneumeth.2003.10.009

DeWitt I, Rauschecker JP. 2012. Phoneme and word recognition in the auditory ventral stream. *Proc Natl Acad Sci U S A* **109**:E505-514. doi:10.1073/pnas.1113427109

Dutoit T, Pagel V, Pierret N, Bataille F, van der Vrecken O. 1996. The MBROLA project: towards a set of high quality speech synthesizers free of use for non commercial purposesProceeding of Fourth International Conference on Spoken Language Processing. ICSLP '96. IEEE. pp. 1393–1396. doi:10.1109/ICSLP.1996.607874

Eisermann M, Kaminska A, Moutard M-L, Soufflet C, Plouin P. 2013. Normal EEG in childhood: From neonates to adolescents. *Neurophysiologie Clinique/Clinical Neurophysiology* **43**:35–65. doi:10.1016/j.neucli.2012.09.091

Estes KG, Lew-Williams C. 2015. Listening through voices: Infant statistical word segmentation across multiple speakers. *Dev Psychol* **51**:1517–1528. doi:10.1037/a0039725

Fiser J, Aslin RN. 2002. Statistical learning of new visual feature combinations by infants. *Proceedings of the National Academy of Sciences of the United States of America* **99**:15822–6. doi:10.1073/pnas.232472899

Flo A, Benjamin L, Palu M, Dehaene-Lambertz G. 2022. Sleeping neonates track transitional probabilities in speech but only retain the first syllable of words. *Sci Rep* **12**:4391. doi:10.1038/s41598-022-08411-w

Fló A, Brusini P, Macagno F, Nespor M, Mehler J, Ferry AL. 2019. Newborns are sensitive to multiple cues for word segmentation in continuous speech. *Dev Sci* **22**:e12802. doi:10.1111/desc.12802

Fló A, Gennari G, Benjamin L, Dehane-Lambertz G. 2022. Automated Pipeline for Infants Continuous EEG (APICE): A flexible pipeline for developmental cognitive studies | Elsevier Enhanced Reader. *Developmental Cognitive Neuroscience*. doi:10.1016/j.dcn.2022.101077

Friedrich M, Friederici AD. 2011. Word Learning in 6-Month-Olds: Fast Encoding–Weak Retention. *Journal of Cognitive Neuroscience* **23**:3228–3240. doi:10.1162/jocn_a_00002

Friedrich M, Friederici AD. 2005. Semantic sentence processing reflected in the event-related potentials of one- and two-year-old children. *NeuroReport* **16**:1801. doi:10.1097/01.wnr.0000185013.98821.62

Frost R, Armstrong BC, Siegelman N, Christiansen MH. 2015. Domain generality versus modality specificity: The paradox of statistical learning. *Trends in Cognitive Sciences* **19**:117–125. doi:10.1016/j.tics.2014.12.010

Gennari G, Marti S, Palu M, Flo A, Dehaene-Lambertz G. 2021. Orthogonal neural codes for speech in the infant brain. *Proc Natl Acad Sci USA* **118**:e2020410118. doi:10.1073/pnas.2020410118

Giraud A-L, Poeppel D. 2012. Cortical oscillations and speech processing: emerging computational principles and operations. *Nature neuroscience* **15**:511–7. doi:10.1038/nn.3063

Gwilliams L, King J-R, Marantz A, Poeppel D. 2022. Neural dynamics of phoneme sequences reveal position-invariant code for content and order. *Nat Commun* **13**:6606. doi:10.1038/s41467-022-34326-1

Hauser MD, Newport EL, Aslin RN. 2001. Segmentation of the speech stream in a non-human primate: Statistical learning in cotton-top tamarins. *Cognition* **78**:53–64. doi:10.1016/S0010-0277(00)00132-3

Hay JF, Pelucchi B, Estes KG, Saffran JR. 2011. Linking sounds to meanings: Infant statistical learning in a natural language. *Cognitive Psychology* **63**:93–106. doi:10.1016/j.cogpsych.2011.06.002

Johnson EK, Westrek E, Nazzi T, Cutler A. 2011. Infant ability to tell voices apart rests on language experience: Infant voice discernment. *Developmental Science* **14**:1002–1011. doi:10.1111/j.1467-7687.2011.01052.x

Junge C, Boumeester M, Mills DL, Paul M, Cosper SH. 2021. Development of the N400 for Word Learning in the First 2 Years of Life: A Systematic Review. *Frontiers in Psychology* **12**:2420. doi:10.3389/fpsyg.2021.689534

Jusczyk PW, Hohne EA. 1997. Infants' Memory for Spoken Words. *Science* **277**:1984–1986. doi:10.1126/science.277.5334.1984

Kabdebon C, Flo A, de Heering A, Aslin R. 2022. The power of rhythms: how steady-state evoked responses reveal early neurocognitive development. *NeuroImage* 119150. doi:10.1016/j.neuroimage.2022.119150

Kirkham NZ, Slemmer J a, Johnson SP. 2002. Visual statistical learning in infancy: Evidence for a domain-general learning mechanism. *Cognition* **83**:4–5.

Kudo N, Nonaka Y, Mizuno N, Mizuno K, Okanoya K. 2011. On-line statistical segmentation of a non-speech auditory stream in neonates as demonstrated by event-related brain potentials. *Developmental Science* **14**:1100–1106. doi:10.1111/j.1467-7687.2011.01056.x

Kuhl PK, Miller JD. 1982. Discrimination of auditory target dimensions in the presence or absence of variation in a second dimension by infants. *Perception & Psychophysics* **31**:279–292. doi:10.3758/BF03202536

Kutas M, Federmeier KD. 2011. Thirty years and counting: Finding meaning in the N400 component of the event related brain potential (ERP). *Annu Rev Psychol* **62**:621–647. doi:10.1146/annurev.psych.093008.131123

Mahmoudzadeh M, Wallois F, Kongolo G, Goudjil S, Dehaene-Lambertz G. 2016. Functional Maps at the Onset of Auditory Inputs in Very Early Preterm Human Neonates. *Cereb Cortex* **27**:2500–2512. doi:10.1093/cercor/bhw103

Moerel M, De Martino F, Formisano E. 2012. Processing of Natural Sounds in Human Auditory Cortex: Tonotopy, Spectral Tuning, and Relation to Voice Sensitivity. *Journal of Neuroscience* **32**:14205–14216. doi:10.1523/JNEUROSCI.1388-12.2012

Monroy CD, Gerson SA, Hunnius S. 2017. Toddlers' action prediction: Statistical learning of continuous action sequences. *Journal of Experimental Child Psychology* **157**:14–28. doi:10.1016/j.jecp.2016.12.004

Norman-Haignere S, Kanwisher NG, McDermott JH. 2015. Distinct Cortical Pathways for Music and Speech Revealed by Hypothesis-Free Voxel Decomposition. *Neuron* **88**:1281–1296. doi:10.1016/j.neuron.2015.11.035

Oostenveld R, Fries P, Maris E, Schoffelen J-M. 2011. FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational Intelligence and Neuroscience* **2011**:1–9. doi:10.1155/2011/156869

Parise E, Friederici AD, Striano T. 2010. "Did You Call Me?" 5-Month-Old Infants Own Name Guides Their Attention. *PLoS ONE* **5**:e14208. doi:10.1371/journal.pone.0014208

Pelucchi B, Hay JF, Saffran JR. 2015. Statistical learning in a natural language by 8-month-old infants. *Child development* **80**:674–685. doi:10.1111/j.1467-8624.2009.01290.x

Reid VM, Hoehl S, Grigutsch M, Groendahl A, Parise E, Striano T. 2009. The neural correlates of infant and adult goal prediction: Evidence for semantic processing systems. *Developmental Psychology* **45**:620–629. doi:10.1037/a0015209

Ren J, Wang M. 2023. Development of statistical learning ability across modalities, domains, and languages. *Journal of Experimental Child Psychology* **226**:105570. doi:10.1016/j.jecp.2022.105570

Saffran JR, Aslin RN, Newport EL. 1996. Statistical learning by 8-month-old infants. doi:10.1126/science.274.5294.1926

Saffran JR, Johnson EK, Aslin RN, Newport EL. 1999. Statistical learning of tone sequences by human infants and adults. *Cognition* **70**:27–52. doi:10.1016/S0010-0277(98)00075-4

Saffran JR, Kirkham NZ. 2018. Infant Statistical Learning. *Annual Review of Psychology* **69**:181–203. doi:10.1146/annurev-psych-122216-011805

Santolin C, Rosa-Salva O, Vallortigara G, Regolin L. 2016. Unsupervised statistical learning in newly hatched chicks. *Current Biology* **26**:R1218–R1220. doi:10.1016/j.cub.2016.10.011

Santolin C, Saffran JR. 2018. Constraints on Statistical Learning Across Species. *Trends in Cognitive Sciences* **22**:52–63. doi:10.1016/j.tics.2017.10.003

Shukla M, White KS, Aslin RN. 2011. Prosody guides the rapid mapping of auditory word forms onto visual objects in 6-mo-old infants. *Proc Natl Acad Sci USA* **108**:6038–6043. doi:10.1073/pnas.1017617108

Shultz S, Vouloumanos A, Bennett RH, Pelphrey K. 2014. Neural specialization for speech in the first months of life. *Developmental Science* **17**:766–774. doi:10.1111/desc.12151

Sylvester CM, Kaplan S, Myers MJ, Gordon EM, Schwarzlose RF, Alexopoulos D, Nielsen AN, Kenley JK, Meyer D, Yu Q, Graham AM, Fair DA, Warner BB, Barch DM, Rogers CE, Luby JL, Petersen SE, Smyser CD. 2023. Network-specific selectivity of functional connections in the neonatal brain. *Cerebral Cortex* **33**:2200–2214. doi:10.1093/cercor/bhac202

Toro JM, Trobalón JB. 2005. Statistical computations over a speech stream in a rodent. *Perception & psychophysics* **67**:867–875. doi:10.3758/BF03193539

Zatorre RJ, Belin P. 2001. Spectral and Temporal Processing in Human Auditory Cortex. *Cerebral Cortex*.