Regeneration following tissue necrosis is mediated by non-apoptotic caspase activity

Jacob W. Klemm, Chloe Van Hazel, Robin E. Harris

Arizona State University, 427 E Tyler Mall LSE 229, Tempe, AZ 85287-4501

Corresponding author: robin.harris@asu.edu

Running title: Dronc promotes necrosis-induced regeneration.

Abstract

Tissue necrosis is a devastating complication for many human diseases and injuries. Unfortunately, our understanding of necrosis and how it impacts surrounding healthy tissue – an essential consideration when developing effective methods to treat such injuries – has been limited by a lack of robust genetically tractable models. Our lab previously established a method to study necrosis-induced regeneration in the *Drosophila* wing imaginal disc, which revealed a unique phenomenon whereby cells at a distance from the injury upregulate caspase activity in a process called Necrosis-induced Apoptosis (NiA) that is vital for regeneration. Here we have further investigated this phenomenon, showing that NiA is predominantly associated with the highly regenerative pouch region of the disc, shaped by genetic factors present in the presumptive hinge. Furthermore, we find that a proportion of NiA fail to undergo apoptosis, instead surviving effector caspase activation to persist within the tissue and stimulate reparative proliferation late in regeneration. This proliferation relies on the initiator caspase Dronc, and occurs independent of JNK, ROS or mitogens associated with the previously characterized Apoptosis-induced Proliferation (AiP) mechanism. These data reveal a new means by which non-apoptotic Dronc signaling promotes regenerative proliferation in response to necrotic damage.

Introduction

Necrosis is the rapid, disordered death of cells characterized by the loss of membrane integrity and release of cytoplasmic contents into the surrounding tissue (Hajibabaie *et al.* 2023). This catastrophic type of cell death can occur in diverse tissues and is central to many human conditions, particularly those related to ischemic injuries. Such conditions include chronic illnesses like diabetes, joint disorders, sickle-cell anemia and other inherited and congenital diseases (Masi *et al.* 2007, Mulay *et al.* 2016, Karsch-Bluman *et al.* 2019, Tonnus *et al.* 2021, Li *et al.* 2023), as well as more acute medical events like strokes, heart attacks, bacterial infections and common traumatic injuries (Konstantinidis *et al.* 2012, Hakkarainen *et al.* 2014, Bonne and Kadri 2017, Wu *et al.* 2018). Even therapeutic interventions, in particular treatments for cancer, can result in this devastating form of damage (Robertson *et al.* 2017, Nakada *et al.* 2019, Yang *et al.* 2021). Unfortunately, current strategies to treat necrosis mainly focus on invasive procedures that are often met with limited success. With such substantial bearings on human health, it is crucial to better understand the effects of necrosis in disease and injury, particularly in the context of tissue repair and regeneration.

We currently have a limited understanding of how necrosis impacts surrounding healthy tissue during wound healing. Indeed, much of our understanding about how cell death influences tissue repair comes instead from models involving programmed cell death (PCD) like apoptosis (Hajibabaie *et al.* 2023). This highly regulated process can be triggered by intrinsic or extrinsic pathways, both of which lead to the activation of caspases that mediate the controlled disassembly of the cell (Ashkenazi and Salvesen 2014). Studies of PCD in a variety of species have shown that cells undergoing apoptosis can release signaling molecules that are interpreted by surrounding tissues to drive wound healing events, such as tissue remodeling, immune responses, survival and proliferation of surrounding cells (Tseng *et al.* 2007, Fan and Bergmann 2008, Chera *et al.* 2009, Bergmann and Steller 2010, Li *et al.* 2010, Pellettieri *et al.* 2010, Ryoo and Bergmann 2012, Vriz *et al.* 2014, Fuchs and Steller 2015, Perez-Garijo and Steller 2015, Fogarty and Bergmann 2017, Perez-Garijo 2018). For example, the signaling molecules Prostaglandin E2 and Hedgehog are produced by dying hepatocytes to induce regenerative proliferation in the vertebrate liver (Jung *et al.* 2010, Li *et al.* 2010), while apoptotic-deficient mice show both impaired liver regeneration and epidermal recovery after wounding (Li *et al.* 2010). Although mitogenic signaling by apoptotic cells is an established and conserved process, whether similar signaling events occur following necrotic cell death is less clear.

Evidence of apoptotic signaling first originated from studies of the larval wing primordia in *Drosophila* (Perez-Garijo *et al.* 2004, Ryoo *et al.* 2004). This epithelial tissue has been extensively characterized as a model for growth, development and regeneration, including the role that cell death plays in these processes (Beira and Paro 2016, Worley and Hariharan 2022). Ongoing studies of this model have identified an essential signaling network centered on the highly conserved JNK pathway. JNK activates several major signaling pathways including Hippo and JAK/STAT, which have conserved roles in promoting regeneration across

species (Worley *et al.* 2012, Hariharan and Serras 2017, Fox *et al.* 2020, Worley and Hariharan 2022), as well as activating JNK itself via overlapping positive feedback loops. One such feedback loop acts through the initiator caspase Dronc (*Drosophila* Caspase-9), which, independent of its role in apoptosis, is translocated to the cell membrane to activate the release of ROS from the NADPH oxidase Duox (Amcheslavsky *et al.* 2018). ROS attracts hemocytes to further activate JNK signaling in the disc through the release of the TNF ligand Eiger (Fogarty *et al.* 2016, Diwanji and Bergmann 2018). In a related pathway, JNK can also lead to the expression of the *Duox* maturation factor *moladietz* (*mol*), thus activating this feedback loop without Dronc (Khan *et al.* 2017, Pinal *et al.* 2018). An important advance in elucidating this network was the ability to generate "undead cells", using the baculovirus caspase inhibitor P35 to prevent apoptotic cells from dying (Hay *et al.* 1994). These cells therefore persist, releasing mitogenic factors including Wingless (Wg, Wnt1), Decapentaplegic (Dpp, BMP2/4), Spitz (Spi, EGF) or Hedgehog (Hh) (Huh *et al.* 2004, Perez-Garijo *et al.* 2004, Ryoo *et al.* 2004, Perez-Garijo *et al.* 2005, Fan and Bergmann 2008, Perez-Garijo *et al.* 2009, Morata *et al.* 2011, Fan *et al.* 2014). These signals subsequently promote proliferation of the surrounding cells in a phenomenon known as Apoptosis-induced Proliferation (AiP) (Fan and Bergmann 2008, Ryoo and Bergmann 2012, Fogarty and Bergmann 2017).

By contrast, the genetic events following necrosis are less well explored. Necrosis is characterized by swelling and loss of cellular membrane integrity, with the release of cellular contents into the intercellular space causing a significant inflammatory response (Festjens *et al.* 2006, D'Arcy 2019, Hajibabaie *et al.* 2023). Necrosis is highly variable, occurring as a regulated process, for example necroptosis, or as unregulated, caspase-independent lysis (Ashkenazi and Salvesen 2014, D'Arcy 2019). The factors released from necrotic cells are collectively termed Damage-Associated Molecular Patterns (DAMPs)(Venereau *et al.* 2015, Roh and Sohn 2018), which are thought to interact with pattern recognition receptors (PRRs) on nearby cells, mostly of the Toll-like receptor (TLR) family (Ming *et al.* 2014, Gong *et al.* 2020). DAMPs are understood to mainly consist of fundamental cellular components like histones, chromatin, and actin (Venereau *et al.* 2015, Gordon *et al.* 2018, Roh and Sohn 2018), although specific factors have also been described. For example, High-mobility group box 1 (HMBG1) has been characterized as a DAMP in models of spinal cord, cardiac and muscle injury where it promotes angiogenesis, attracts repair cells and induces proliferation (Venereau *et al.* 2015), as well as in *Drosophila* models of necrosis (Nishida *et al.* 2024). However, the overall role of DAMPs and how they influence healing and regeneration through interaction with healthy tissues has yet to be fully explored.

To better investigate necrosis-induced wound repair and regeneration, our lab developed a method to rapidly and reproducibly induce necrotic cell death within the developing *Drosophila* wing imaginal discs (Klemm *et al.* 2021). Using a genetic ablation system we previously established, named Duration and Location (DUAL) Control (Harris 2023), we can induce necrosis in the wing disc via expression of a leaky cation channel *GluR1^{LC}* (Liu *et al.* 2013, Yang *et al.* 2013). Using this system (*DC^{GluR1}*), we showed that

wing discs are capable of fully regenerating following necrotic injury at a rate comparable to that of damage

induced by apoptosis (Klemm *et al.* 2021). However, while apoptotic ablation leads to JNK signaling and

extensive caspase activity, we found that necrosis leads to only minor levels of JNK-mediated apoptosis,

which is confined to the wound edge, but unexpectedly generates significant caspase activity in cells distant

from the injury. We called this non-autonomous caspase activation Necrosis-induced Apoptosis (NiA)

(Klemm *et al.* 2021). Unlike normal apoptotic cells, NiA form entirely independent of JNK signaling, and

cannot be made undead using P35. We also demonstrated that NiA is essential for regeneration, although

the mechanism was unclear.

Here we have further characterized the NiA phenomenon, finding that only regeneration-competent areas of

the wing disc can produce NiA following damage, in part due to WNT and JAK/STAT signaling in the hinge

that limits NiA to the pouch. Building upon our finding that NiA is necessary for regeneration, we show that

NiA leads to localized proliferation significantly later in regeneration than previously observed. Using tools to

trace caspase activity and cell death, we demonstrate that this is possible because a proportion of NiA

survive effector caspase activation and persist late into regeneration where they promote proliferation.

Finally, we show that this proliferation relies on the initiator caspase Dronc, but surprisingly does not involve

established AiP mechanisms. Our data suggest a model in which necrotic injuries induce caspase activity in

cells at a distance from the injury, some of which undergo JNK-independent apoptosis (NiA), while others

survive and promote proliferation through a novel non-apoptotic function of Dronc, which is separate from its

role in AiP. We refer to these surviving NiA cells as Necrosis-induced Caspase-Positive (NiCP) cells. These

findings reveal an important genetic response to lytic cell death that could potentially be leveraged to

augment regeneration of necrotic wounds.

Results

Formation of NiA occurs primarily in the wing pouch.

Previously, we found that NiA occurs in the lateral pouch (LP) upon induction of necrosis in the distal pouch

with DC^{GluR1} (Figure 1A, D and E, yellow arrowheads in E) (Klemm *et al.* 2021). The wing disc itself

comprises different identities reflecting the adult structures they ultimately create, including the pouch, hinge

and notum, which are themselves divided into compartments; anterior/posterior and dorsal/ventral (Figure

1B). Since these various disc identities have distinct regenerative capacities stemming from their different

genetic responses to damage (Martin *et al.* 2017), to better understand the formation of NiA and the role it

plays in regeneration we tested whether necrosis occurring in different areas of the disc leads to NiA. To do

so, we utilized $GAL4/UAS/GAL80^{ts}$ to conditionally express $UAS\text{-}GluR1^{LC}$ (Liu *et al.* 2013, Yang *et al.* 2013)

in the pouch, hinge or notum tissues (Figure 1C). As an initial test, we attempted to recapitulate our original

observations made using DC^{GluR1} by employing an enhancer of the *spalt* gene driving *GAL4*

(*R85E08^{ts}>GluR1*) to cause necrosis in the distal pouch (Figure 1F and G). As anticipated, NiA are formed

in the lateral pouch following 20 hr of ablation (denoted as 0 hr, when larvae are downshifted to 18°C)

(Figure 1G). In this figure and others, NiA are recognized as cells positive for the cleaved caspase cDcp-1

and negative for GFP that labels the ablation domain (*UAS-GFP*) (Figure 1E and G, yellow arrowheads),

which indicates that these caspase-positive cells originate outside the area of ablation. This test confirms

that the NiA phenomenon occurs independent of the ablation system used. Notably, NiA are consistently

absent from the presumptive hinge region surrounding the pouch following both *R85E08^{ts}>GluR1* or *DC^{GluR1}*

ablation (Figure 1E and G). Indeed, outside of the pouch, cDcp-1 staining is only observed in a small area

of the posterior pleura (Figure 1E, red arrowhead) and at low levels stochastically across the disc due to

stress induced by temperature changes (Klemm *et al.* 2021). To further investigate the extent to which

necrotic pouch tissue can induce NiA we next ablated the entire pouch using *rotund-GAL4* (*rn^{ts}>GluR1*) or

nubbin-GAL4 (*nub^{ts}>GluR1*) (Figure 1H, I and J, figure 1- figure supplement 1A). Necrosis of the whole

pouch with either driver fails to result in NiA (Figure 1I and Figure 1- figure supplement 1A), even after 24 hr

of recovery (Figure 1J). Importantly, the majority of cDcp-1-positive cells overlap with expression of the JNK

target *Mmp1* (Figure 1I and J), and therefore resembles cells undergoing JNK-mediated apoptosis, such as

those seen at the wound edge (WE) following distal pouch ablation (Figure 1 – figure supplement 1B,

arrowhead) (Klemm *et al.* 2021)). These data suggest that cells outside of the pouch are generally unable to

respond to DAMPs released by pouch cells undergoing necrosis to generate NiA, or that such DAMPs are

spatially limited.

To test whether necrosis in areas outside of the pouch can induce NiA, we ablated the proximal notum using

pannier-GAL4 (*pnr^{ts}>GluR1*) (Figure 1K), using the absence of the notum Wg stripe to confirm loss of this

area (Figure 1L and M, open arrowheads). In this ablation we observed only minimal NiA, with sporadic

cDcp-1-positive, GFP-negative cells in the unablated areas of the notum (Figure 1L), which does not change

after 24 hr (Figure 1M). To test the hinge, we used a putative *zfh1* enhancer driving *GAL4*

(*R73G07^{ts}>GluR1*), which has hinge-specific expression (Figure 1N). Ablation of the hinge fails to generate

NiA in the notum, and surprisingly does not induce a response in the neighboring pouch cells (Figure 1O

and P), despite their demonstrated ability to form NiA (Figure 1G). Given that the regenerative capacity of

the wing disc is mainly constrained to the pouch (Martin *et al.* 2017, Martin and Morata 2018), it is possible

that the intrinsic differences in regenerative capacity across the disc accounts for the disparity in NiA/NiCP

occurrence in the pouch versus the notum. A difference in the availability of the relevant DAMPs or PRRs

may also be responsible for the pattern of NiA formation throughout the disc. Alternatively, there may be

inhibitory factors present in notum cells that protect against the induction of pro-apoptotic pathway elements.

Indeed, differences in caspase 3 activation within the notum and pouch have been shown (Bergantinos *et al.*

2010). Thus, these data suggest that either only the pouch releases DAMPs - and has the requisite PRRs to

respond to these DAMPs - that lead to NiA following necrosis, or that unknown genetic factors limit the
capacity of the notum and hinge to respond to damage.

As the efficacy of DAMPs might be limited by how far they can reach after being released from lysed cells,
we also induced necrosis in the entire posterior disc compartment with *hedgehog-GAL4* (*hh^{ts}>GluR1*)
(Figure 1Q), and in an anterior stripe along the anterior/posterior compartment boundary using *patched-*
GAL4 (*ptc^{ts}>GluR1*) (Figure 1 – figure supplement 1C). These experiments cause the simultaneous necrosis
of pouch, hinge and notum tissues, allowing us to determine the potential of these different tissue identities
to produce NiA. In both experiments, NiA cells are observed in the pouch, and to a lesser extent the notum,
but are still strikingly absent from the hinge (Figure 1R, R' and T, Figure 1 – figure supplement 1D, D' and
F). After 24 hr of recovery, there is an increase in the number of NiA within the pouch and notum, but not the
hinge (Figure 1S, S' and T, Figure 1 – figure supplement 1E, E' and F). Thus, it appears that NiA can occur
outside of the pouch when a large enough area, or an area that also includes the pouch, is ablated.
However, the hinge remains completely refractory to NiA formation.

To avoid any bias in the use of tissue-specific *GAL4* drivers, we also made RFP-labeled stochastic clones
that have the potential to undergo necrosis upon changing the growth temperature to 30°C (Figure 1 – figure
supplement 1G). After allowing these clones to develop, we triggered necrosis in early third larval instar and
examined the extent of active caspase in the different disc regions (Figure 1 – figure supplement 1H). As
expected, necrosis of clones in the pouch leads to active caspases both within and surrounding the ablated
area, including cells without the RFP clone label, suggesting that NiA has occurred (Figure 1 – figure
supplement 1H', arrowheads). We also found that necrosis in the notum leads to comparatively little
caspase labeling (Figure 1 – figure supplement 1H'') consistent with the notum being less able to generate
NiA. Necrotic clones in the hinge also produces caspase activity, but most of these cells also have RFP,
suggesting again that NiA does not occur in the hinge (Figure 1 – figure supplement 1H'). Using this
approach we also examined whether NiA formation occurs in other larval tissues as well, and found that
necrotic clones in both haltere and leg discs appear to result in an NiA response (Figure 1 – Figure
supplement 1J). Of note, the use of clones that naturally vary in size also demonstrates that the area of
ablation is related to the amount of NiA produced (Figure 1 – figure supplement 1I), which is also true of our
tissue-specific ablation experiments (Figure 1 – figure supplement 1I).

Together, these data infer three important conclusions: 1) all areas of the disc can be killed by necrosis and
therefore potentially can release DAMPs, 2) NiA is limited to the pouch when local necrosis occurs, but
when multiple (or large) areas of the disc are killed, limited NiA can also be induced in the notum, although
we cannot rule out that this is due to DAMPs from dying pouch cells, and 3) the hinge is refractory to NiA,
which is consistent with other findings that show its resistance to apoptosis in response to irradiation
(Verghese and Su 2016). Thus, the overall pattern of competence to undergo NiA appears to reflect the

uneven regenerative capacity of the wing disc, with NiA formation predominantly associated with the highly
regenerative wing pouch.

NiA is regulated by WNT and JAK/STAT signaling.

As NiA readily occurs in the pouch but is excluded from the nearby hinge, we used this contrasting response
to identify genetic factors that might regulate NiA formation. The wing hinge is specified by JAK/STAT
signaling during disc development, which can protect cells from irradiation-induced apoptosis potentially via
the expression of *Zn finger homeodomain* 2 (*zfh2*) (La Fortezza *et al.* 2016, Verghese and Su 2016,
Verghese and Su 2018). Alongside JAK/STAT, the presence of Wingless (Wg, *Drosophila* Wnt1), which
encircles the pouch, may also protect cells from death and permit regeneration of the pouch through the
repression of *reaper* (*rpr*) (Verghese and Su 2016). As such, we investigated both JAK/STAT and Wg to
determine whether they regulate NiA formation.

The activity of the JAK/STAT pathway can be visualized in the hinge of early third instar larval discs by a
10XSTAT-GFP reporter (Bach *et al.* 2007)(Figure 2A). Upon ablation with DC^{GluR1}, high levels of JAK/STAT
activity are observed in the pouch at the immediate WE (Figure 2B, arrowhead), similar to its upregulation
following irradiation or apoptotic ablation (Herrera and Bach 2019). As JNK signaling is induced at the WE
(Figure 1 – figure supplement 1B) (Klemm *et al.* 2021), and the *unpaired* ligands are targets of JNK signaling
(Katsuyama *et al.* 2015, Jaiswal *et al.* 2023), this JAK/STAT activity is likely to be JNK-mediated. By
contrast, low levels of JAK/STAT activity are observed in the areas of the pouch where NiA occurs (Figure
2B, open arrowheads), surrounded by the higher developmental JAK/STAT in the hinge (Figure 2B). To
determine if low JAK/STAT activity is important for NiA, we knocked down the receptor *domeless* (*UAS-*
dome^{RNAi}) in the pouch, which results in a significant increase in NiA (Figure 2C, D and E), but has no effect
when expressed without damage (Figure 2 – Figure supplement 1A), suggesting that JAK/STAT signaling
may negatively regulate the formation of NiA. To support this finding, we next ectopically activated
JAK/STAT (*UAS-hop48A*). However, even in the absence of damage, this expression results in high levels
of caspase positive cells (Figure 2 – figure supplement 1B and C), making it difficult to determine an effect
on NiA formation. Therefore, to further investigate if JAK/STAT regulates NiA formation, we asked whether
reducing developmental JAK/STAT in the hinge might lead to NiA spreading further into this region. We
generated a version of DUAL Control that expresses *GAL4* in the posterior compartment by replacing the
pouch-specific *DVE>>GAL4* with *hh-GAL4* (Figure 2F and G, Figure 2 – figure supplement 1D). To prevent
hh-GAL4 from being active throughout development, we included $GAL80^{ts}$ (hereafter $DC^{GluR1}hh^{ts}$) and used
temperature changes to limit GAL4 activity to the period just prior to ablation (Figure 2F). With this system,
we knocked down the expression of the JAK/STAT transcription factor *Stat92E* (*UAS-Stat92E^{RNAi}*) in the
posterior compartment (Figure 2 – Figure supplement 1E and F) and ablated the distal pouch, which again

shows an increase in caspase-positive cells in the pouch (Figure 2H, arrowhead, quantified in Figure 2L), but surprisingly NiA cells are still not observed in the hinge (Figure 2H, open arrowhead). To further test this result, we also targeted *zinc finger homeodomain 2* (*Zfh2*), a downstream target of the JAK/STAT pathway that potentially protects cells from apoptosis (La Fortezza *et al.* 2016, Verghese and Su 2018). The knockdown of *Zfh2* slightly increases the formation of NiA, although not significantly (Figure 2 – figure supplement 1I and J, quantified in Figure 2L). However, we noted that the level of Zfh2 protein did not appear strongly affected by this knockdown (Figure 2 – figure supplement 1I), and therefore we cannot confidently conclude that Zfh2 is involved in caspase activation in the context of necrotic damage. Expression of these RNAi lines targeting *Stat92E* or *Zfh2* under non-ablating conditions does not yield any increase in caspase signal (Figure 2 – figure supplement 1G and I). Thus, JAK/STAT signaling appears to limit NiA formation in the pouch, while the inability for NiA to expand into the hinge upon reducing JAK/STAT suggests that other hinge-specific factors may be involved.

Wg has also been shown to protect cells from apoptosis in the hinge (Verghese and Su 2016), and therefore could influence the formation of NiA. Unlike the stochastic temperature change-induced apoptosis (Figure 2I), we noted that NiA in the pouch frequently occurs in discrete populations that avoid Wg at the margin stripe and the inner Wg circle at the boundary of the pouch and hinge (Figure 2J), limiting formation of the NiA cells to regions that appear to overlap the *vestigial quadrant enhancer* (*vgQE-lacZ*, Figure 2 – figure supplement 1K and L) (Kim *et al.* 1996). By contrast, cDcp-1-positive cells at the WE do not avoid the Wg margin stripe (Figure 2J, arrowhead), suggesting that this behavior may be specific to NiA. To test this, we utilized $DC^{GluR1}hh^{ts}$ to knock down *wg* in the posterior compartment of the disc (*UAS-wg^{RNAi}*) and found that NiA now occurs in areas of the pouch where *wg* expression is lost, including the wing margin and inner hinge (Figure 2K, arrowhead, quantified in Figure 2L), unlike NiA in the anterior (Figure 2K, open arrowhead). The increase in cDcp-1-positive cells does not occur when *wg* signaling is similarly blocked without damage (Figure 2 – Figure supplement 1H). Notably, the converse experiment in which *wg* is ectopically expressed during ablation does not suppress NiA (Figure 2 – Figure supplement 1M and N), consistent with our hypothesis that other factors, such as targets downstream of JAK/STAT, might act alongside Wg to regulate NiA. Together, these data demonstrate that both WNT and JAK/STAT signaling act to limit NiA, thus potentially constraining it to the pouch following necrosis.

NiA promotes proliferation late in regeneration.

As NiA is spatially regulated by at least two major signaling pathways in the disc, we next focused on how the localization of NiA relates to its role in promoting regeneration. In our previous work, using an E2F reporter (*PCNA-GFP*) we found that the appearance of NiA coincides with localized proliferation in the distal pouch at 18 hr post-ablation close to the wound, which persists at 24 hr post ablation (Klemm *et al.* 2021).

However, investigation at subsequent time points of recovery shows that regenerative proliferation continues

to increase through 36 hr and 48 hr of regeneration, later than we initially assayed, with a significant

increase in E2F reporter expression occurring between 18 and 36 hr in the pouch relative to the whole disc,

(Figure 3A-A''', Figure 3 – Figure supplement 1A and B, quantified in C). To investigate this proliferative

response, we used EdU to assay the relative level of cell proliferation in discs throughout regeneration from

18 hr to 48 hr post-ablation with DC^{GluR1} (Figure 3B-B'''), using folds as landmarks to normalize EdU intensity

in the pouch relative to the disc (Figure 3 – figure supplement 1D-D'''). We also performed the same time

course with apoptotic ablation using DC^{hepCA} for comparison (Figure 3C-C'''). At 18 hr of regeneration

following ablation with either DC^{GluR1} or DC^{hepCA}, cells at the WE have already migrated distally to close the

injury, while EdU is absent from an area immediately adjacent to the wound (Figure 3B and C). This is

consistent with the recently described JNK-mediated pause in proliferation that occurs in regenerating wing

discs (Jaiswal *et al.* 2023). At 24 hr, this proliferation-devoid area continues to persist following apoptotic

injury (DC^{hepCA}) and the EdU signal becomes elevated broadly across the rest of the pouch, representing the

formation of a blastema (Figure 3C'). By contrast, at 24 hr after necrosis (DC^{GluRl}), the EdU label is

reestablished in cells around the wound showing that proliferation has restarted in these cells (Figure 3B').

Unlike DC^{hepCA}, the rest of the pouch does not appear to change its rate of proliferation (Figure 3B'). By 36

hr following necrosis, a higher EdU signal occurs broadly across the pouch relative to the rest of the disc,

which is significantly stronger compared to discs ablated by apoptosis (Figure 3B'', C'' and E). This

localization is maintained at 48 hr and remains consistently higher in DC^{GluR1} versus DC^{hepCA} ablated discs

(Figure 3B''', C''' and E). Thus, the timeline of recovery from necrosis appears to be distinct from that of

apoptotic injury, with the strongest changes in regenerative proliferation occurring at comparatively later

stages.

The localized proliferation at 36 hr occurs after the appearance of NiA. We previously showed that blocking

the apoptotic pathway by simultaneously knocking down DIAP1 inhibitors *rpr*, *hid* and *grim* (*UAS-mir(RHG),*

(Siegrist *et al.* 2010)) throughout the pouch limits the initial change in proliferation at early stages (18 – 24

hr) and inhibits regeneration (Klemm *et al.* 2021). However, it remains unclear whether this newly observed

localized proliferation at 36 hr and 48 hr also relies on a functional apoptotic pathway, and moreover, to what

extent this regenerative proliferation relies on the JNK-mediated apoptosis at the WE versus the JNK-

independent NiA in the LP. To answer these questions, we blocked apoptosis throughout the pouch and this

time examined proliferation in late regeneration using EdU (Figure 3D-D'''). We found that the localized

increase in EdU signaling at 36 hr is lost (Figure 3D'' and F), although by 48 hr this increase is mostly

restored (Figure 3D''' and F). Importantly, the expression of *mir(RHG)* does not influence EdU levels in the

absence of damage (Figure 3 – figure supplement 1E, F and G). Together, these data confirm that a

functional apoptotic pathway is necessary to induce a localized proliferative response late in regeneration

following necrosis. To understand the relative contribution of WE apoptosis or the NiA in the LP, we

designed experiments to block apoptosis in each disc area alone (*LP>mir(RHG),* Figure 3I, and

WE>mir(RHG), Figure 3J, see Materials and Methods for genotypes) relative to the whole pouch knock

down (Figure 3G and H). Strikingly, the high levels of EdU normally present at 36 hr are strongly reduced

when apoptosis is blocked in either population (Figure 3K-O), suggesting that both dying cells at the WE and

the NiA in the LP contribute to regenerative proliferation following necrosis. These data agree with our

previous findings that both populations are necessary for the overall ability to regenerate adult wings

(Klemm *et al.* 2021).

NiA does not promote proliferation through Apoptosis-induced Proliferation (AiP).

The question remains as to how NiA promotes regenerative proliferation. In *Drosophila*, cells undergoing

apoptosis secrete factors such as Wg and Dpp to induce the proliferation of neighboring cells as part of a

JNK-dependent Apoptosis-induced Proliferation (AiP) (Fogarty and Bergmann 2017). Although NiA occurs

independent of JNK, to determine whether NiA-induced proliferation relies on any of the same signaling

factors as AiP, we examined the damage-specific expression of these various secreted factors. To ensure

we could visualize such signals, we used *lacZ*-based reporters and generated undead cells by expressing

the baculoviral P35 (*UAS-P35*) in the whole pouch. This protein inhibits activity of the effector caspases

Drice and cDcp-1 to block cell death (Hawkins *et al.* 2000, Meier *et al.* 2000), thus allowing signals produced

by these cells to be readily detected. Following ablation with DC^{GluR1}, ectopic *wg* and *dpp* expression (*wg-*

lacZ and *dpp-lacZ*) is observed at the WE (Figure 4C and C' versus A-B', and Figure 4F and F' versus D-E',

arrowheads in C' and F') coinciding with JNK activity in this region (Klemm *et al.* 2021). However, *lacZ* is not

observed in the LP where NiA occurs (Figure 4C' and F', open arrowheads), indicating that these cells do

not activate these mitogens. Similarly, we did not see expression of the EGF ligand *spitz* (*spi-lacZ)* in

DC^{GluR1} ablated discs (Figure 4 – figure supplement 1A and B), which is observed during AiP in the eye (Fan

et al. 2014). These results suggest that NiA does not promote proliferation through the same signaling

factors as those seen during AiP.

We also tested whether other elements required for AiP are involved in NiA-induced proliferation. In addition

to mitogen production, AiP also involves the production of extracellular reactive oxygen species (ROS)

through a non-apoptotic function of Dronc that activates Duox. (Fogarty *et al.* 2016, Fogarty and Bergmann

2017, Amcheslavsky *et al.* 2018, Diwanji and Bergmann 2018). We first examined the extent of ROS

production using dihydroethidium (DHE). This assay showed high levels of ROS localized to the WE but not

in the LP (Figure 4G, arrowhead), suggesting that NiA does not produce ROS during regeneration.

Consistent with this finding, the removal of ROS through pouch-wide expression of either the ROS chelators

Catalase and *Superoxide dismutase 1* (*UAS-Cat; UAS-Sod1*) or knockdown of *Duox* (*UAS-Duox^{RNAi}*) has no

observable effect on the appearance of NiA (Figure 4H and I), although apoptosis at the WE is strongly

suppressed in both experiments (Figure 4H and I, open arrowhead). It has also been shown that the Duox

maturation factor *moladietz* (*mol*) is upregulated following injury to sustain the production of ROS (Khan *et*

al. 2017, Pinal *et al.* 2018). However, while a minor increase in lacZ expression occurs at the WE (Figure 4 –

figure supplement 1C and D, arrowhead), no change in the expression of a *mol* reporter (*mol-lacZ*) is

observed in response to necrosis in the LP. Finally, to functionally test whether AiP is required for the

proliferation associated with NiA, we examined EdU levels across the disc at 36 hr when P35 is expressed.

Normally, when undead cells are created via P35, ectopic mitogen production results increased proliferation

and tumorous overgrowth. However, when P35 is expressed solely in the LP we saw no change in EdU

labeling versus controls (Figure 4J, L and M), suggesting NiA do not form undead cells. When expressed in

the whole pouch, P35 has a small but non-significant effect on EdU (Figure 4J, K and M), consistent with

undead cells now being generated at the WE. Together, these data indicate that cells at the WE undergo

AiP to contribute to regenerative proliferation, while NiA promote proliferation through a different

mechanism.

A subset of cells undergoing NiA are both caspase-positive and have markers of DNA repair

and proliferation.

Robust populations of NiA appear in the wing pouch at around 18 hr of regeneration, (Klemm *et al.* 2021),

while the localized change in EdU labeling that encompasses much of the damaged pouch is detected later

in regeneration at 36 hr and 48 hr (Figure 3B-B'''', E and F). Since the loss of NiA abolishes this change in

proliferation (Figure 3F, L-M and O), we sought to understand how NiA might be influencing proliferation at

these later time points. To do so, we used a robust sensor for the activity of the effector caspases cDcp-1

and Drice called *Green Caspase-3 Activity indicator* (*UAS-GC3Ai*, (Schott *et al.* 2017) to label NiA

throughout regeneration (Figure 5A-E). GC3Ai consists of a cyclized GFP with the N and C termini linked by

a short peptide that contains a caspase-3 cleavage site (DEVD), which prevents fluorescence under non-

apoptotic conditions. Activated effector caspases can cleave the linker peptide, resulting in a fluorescent

GFP signal that reports caspase activity in real time. Schott et al. demonstrated that apoptotic cells labeled

by GC3Ai are extruded within 1 hr of labeling, consistent with previously published work on apoptotic cell

extrusion in imaginal discs (Monier *et al.* 2015). With this reporter we found that, unlike normal apoptotic

cells that are rapidly cleared from the wing disc, NiA persist and increase in abundance by 36 hr when

proliferation localizes to the distal pouch (Figure 5C, quantified in Figure 5F), as well as at 48 hr (Figure 5D,

quantified in Figure 5F), and even up to 64 hr (wandering stage), when regeneration is complete and pouch

tissue is restored (Figure 5E, arrowhead). These persistent *GC3Ai*-positive cells are also frequently

associated with adjacent cells actively undergoing mitosis by 36 hr, indicated by PH3 labelling (Figure 5G-

G'). Since the position of NiA cells change over time from 18 hr when they first appear to 36 hr when the

change in proliferation occurs, to confirm that these GC3Ai-positive cells originally derive from NiA in the LP

rather than the WE, we restricted *GC3Ai* expression to this part of the pouch (*LP>GC3Ai*, Figure 5H and I).

The *GC3Ai* construct is tagged with an HA epitope that can be used to show its expression even in the

absence of activation by caspases (Figure 5 – figure supplement 1A and B), which we used to confirm that

its expression is limited to the LP (Figure 5H and I, Figure 5 – figure supplement 1C). With this experimental

setup, we found that *GC3Ai*-positive cells are still present throughout the pouch at 36 hr, including the distal

salm domain (Figure 5I, arrowhead, versus 5H, open arrowhead, quantified in Figure 5 – Figure supplement

1D), confirming that the source of persistent caspase-positive cells is indeed NiA rather than the WE.

As NiA cells appear to be maintained in the disc for an extended period, we wondered how their behavior

and morphology compared to that of normal apoptotic cells. When apoptosis is induced in wing discs using

DC^{hepCA}, GC3Ai labeling shows apoptotic cells present in the pouch at 18 hr (Figure 5J) throughout the disc

proper (Figure 5J'). At 36 hr these cells appear pyknotic and are basally extruded by 64 hr (Figure 5K and

K'). By comparison, upon ablation with DC^{GluR1}, caspase-positive cells are seen to occupy different regions

of the disc at 18 hr, with WE apoptotic cells closer towards the basal surface and NiA derived from the LP

still within the disc proper (Figure 5L and L'). By 36 hr, the NiA form two distinct populations, some that are

rounded up and appear closer to the basal surface, similar to the WE cells (Figure 5M and M' red

arrowhead), and others that continue to exhibit a columnar appearance and contact both the apical and

basal surfaces of the disc (Figure 5M and M', yellow arrowheads). The appearance and position of these

cells suggest that a proportion of NiA cells may fail to complete apoptosis but instead persist into late stages

of regeneration, despite the presence of detectible caspase activity. This is supported by the observation

that only a minority of GC3Ai-positive cells have blebbing and pyknotic nuclei (Figure 5G', white arrowhead),

while the majority appear to have a consistent and undisturbed cytoplasmic fluorescent label (Figure 5G',

yellow arrowhead). To test our hypothesis that these cells are not undergoing apoptosis, we performed a

TUNEL assay to fluorescently label cells with double stranded DNA breaks in GC3Ai-expressing discs. We

found that most GC3Ai-labeled cells are co-labeled by TUNEL in early (18 hr) and late (36 hr) stages of

regeneration (Figure 5 – figure supplement 1E and F, arrowheads). However, while TUNEL is associated

with apoptosis, by itself it does not confirm that cells are dying (Grasl-Kraupp *et al.* 1995). Therefore, we

also examined levels of γH2Av, a histone variant associated with DNA repair and inhibition of apoptosis

following damaging stimuli such as irradiation (Madigan *et al.* 2002). Interestingly, we found that GC3Ai-

positive cells initially have high levels of γH2Av at 18 hr of regeneration, including those at the WE and the

NiA (Figure 5 – figure supplement 1G, arrowheads), which later is lost from the NiA at 36 hr (Figure 5 –

figure supplement 1H, open arrowheads). This indicates that NiA are undergoing active DNA repair rather

than apoptosis. Taken together, these data suggest that a majority of NiA may upregulate caspase activity,

but rather than undergoing apoptosis, they repair cellular damage and persist in the tissue into late stages of

regeneration where they promote proliferation. As such we are referring to this population of persistent and

potentially non-apoptotic NiA as Necrosis-induced Caspase Positive (NiCP) cells.

NiCP cells have initiator caspase activity but sublethal effector caspase activity.

While this characterization of caspase activation and discovery of NiCP cells do not change any of the
conclusions we have made up until this point, it does raise the question as to why some cells (NiA) undergo
apoptosis and are removed from the tissue in response to necrosis, while others (NiCP) seem to persist
despite caspase activity. The ability for cells to survive caspase activity is not surprising, as many non-
apoptotic roles for caspases have been documented (Su 2020), and includes promoting proliferation. As
mentioned, the initiator caspase Dronc functions in a non-apoptotic role to activate Duox and thus promote
proliferation during AiP in damaged wing discs (Fogarty *et al.* 2016). Therefore, we wondered whether the
difference between NiA and NiCP might arise from the level or activity of caspases within these cells. The
GC3Ai reporter indicates activity of Dcp-1 and Drice (Schott *et al.* 2017) while the anti-cleaved-Dcp-1
antibody is thought to also detect Drice (Li *et al.* 2019). Thus, these tools exclusively detect effector
caspases. As such, to gain a better understanding of caspase activity in NiCP we used two lineage tracing
tools to study the activity of initiator versus effector caspases. The initiator caspase-sensitive Drice-Based
Sensor *(DBS-GFP)* consists of a modified version of the effector caspase Drice that is 1) a membrane-
bound via mCD8, and 2) linked to a nuclear GFP (Baena-Lopez *et al.* 2018). Upon cleavage by Dronc the
GFP is released and translocates to the nucleus, providing a readout for Dronc activity. (Baena-Lopez *et al.*
2018). The second tool, CasExpress, is also membrane-bound via mCD8 and attached to GAL4 through a
short linker peptide that contains the Drice/Dcp-1 cleavage site DQVD, thus providing a readout for the
activity of both effector caspases Drice and Dcp-1 (Ding *et al.* 2016). This tool is highly sensitive and is
capable of detecting non-apoptotic caspase activity that occurs during development. However, since GAL4
activity can be modulated by temperature, the sensitivity of CasExpress can be altered by a combination of
GAL80^{ts} and temperature to suppress this background signal and optimize damage-induced caspase
labeling (Colon Plaza and Su 2024). Our results show a strong overlap of *DBS-GFP* with anti-cDcp-1 in the
LP of DC^{GluR1} ablated discs (Figure 6A and A'), indicating that these cells have robust Dronc activity, as do
the cells at the WE (Figure 6A and A'). However, when we used CasExpress to examine effector caspase
activity with a protocol that eliminates developmental caspase activity (Colon Plaza and Su 2024), we noted
that only cells at the WE were labelled (Figure 6B, B' and B'', arrowhead), most of which had pyknotic nuclei
showing they are actively undergoing apoptosis, while few cells in the LP were labelled (Figure 6B''). We
hypothesized that the level of effector caspase activity might be high enough to be detected by GC3Ai and
the cDcp-1 antibody, but not by CasExpress. To test this idea, we further attempted to detect NiCP with
CasExpress by combining it with G-TRACE (Evans *et al.* 2009), which should lineage trace cells that have
effector caspase activity, permanently labeling them with GFP at the start of ablation. Again, we found that
only cells of the WE are labelled during regeneration (Figure 6B''', arrowhead), with minimal labelling of cells
in the LP (Figure 6B''', open arrowhead), even after 36 hr of regeneration when the NiCP-induced uptick in

proliferation occurs (Figure 6 – figure supplement 1A). By comparison, performing these same experiments

using DC^{hepCA} to induce extensive apoptotic cell death leads to a significant proportion of cDcp-1-positive

cells being labelled by CasExpress under both normal and lineage tracing conditions (Figure 6C,

arrowhead), suggesting that in the context of necrosis there is not enough effector caspase activity to label

NiCP using these methods. Indeed, if the CasExpress experiment is performed in the absence of the

GAL80^{ts} that suppresses the background developmental caspase signal, the NiCP cells now become

labeled by GFP (Figure 6D), indicating that effector caspases are indeed present in these cells, but at low

enough levels to avoid death. Evidence for the existence of a cellular execution threshold of caspase activity

in cells of the wing disc, which must be reached to induce apoptosis, has previously been documented

(Florentin and Arama 2012). This is further supported by the observation that the GFP label becomes more

apparent later in regeneration (Figure 6E), confirming that cells with effector caspase activity persist rather

than die. Thus, it appears that in response to necrosis, the cells of the LP activate the initiator caspase

Dronc and the effector caspase(s) Drice/cDcp-1 (to an extent) but fail to undergo programmed cell death.

Instead, these cells persist in the disc late into regeneration where they stimulate regenerative proliferation.

Dronc in NiCP cells promote proliferation independent of AiP.

While NiCP cells have both initiator (Dronc) and effector caspase (Drice/cDcp-1) activity, it appears that the

level or function of effector caspases is insufficient to cause apoptosis, and is also inconsequential for

promoting regeneration – indeed, blocking Drice/cDcp-1 activity with P35 does not affect the localized

regenerative proliferation observed at 36 hr (Figure 4L and M), or the overall ability to regenerate (Klemm *et*

al. 2021). This is in contrast with the ability of effector caspases to drive proliferation in the eye disc (Fan

and Bergmann 2008). However, blocking the apoptotic pathway upstream of Dronc using *mir(RHG)*

eliminates NiA/NiCP (Figure 7A, C and K, (Klemm *et al.* 2021), blocks the increase in proliferation (Figure

7B, D and L), and limits regeneration (Klemm *et al.* 2021). These observations demonstrate that proliferation

induced by NiCP must depend on factors downstream of *rpr/hid/grim* and upstream of the effector caspases

Drice/cDcp-1, thus potentially pointing to a role for *Dronc*. To test this, we ablated discs with DC^{GluR1} while

reducing the activity of Dronc using a null allele (*Dronc^{I29}/+,* (Xu *et al.* 2005)). With this genetic background

there is a significant reduction in regenerative proliferation at 36 hr (Figure 7F and L) and regeneration is

limited, shown by both adult wing scores (Figure 7M) and size (Figure 7 – figure supplement 1A). These

data demonstrate an essential role for Dronc in NiCP to promote proliferation and subsequent regeneration

of the disc following necrosis. We noted that this mutant does not strongly affect the number of NiCP

assayed by cDcp-1 (Figure 7E and K), likely because this allele does not completely suppress apoptosis

unless homozygous (Xu *et al.* 2005), which is precluded by the genetics of this experiment. As an

alternative, we interfered with Dronc function by expressing a dominant negative form of *Dronc* that contains

only the caspase recruitment domain (CARD), here referred to as *Dronc^{DN}*, which blocks activation of Dcp-

1/Drice and apoptosis (Meier *et al.* 2000). *Dronc^DN* reduces NiCP number, but to a lesser degree than

mi(RHG) (Figure 7G and K) and does not affect proliferation (Figure 7H and L), suggesting that the CARD

domain is dispensable for NiCP-induced proliferation.

Finally, we wondered how this might relate to the previously documented role of Dronc in promoting

proliferation following apoptotic cell death during AiP (Fogarty and Bergmann 2017). AiP depends on both

JNK and ROS (Fogarty *et al.* 2016), which we have shown are only present at the WE and are not

associated with NiA/NiCP (Figure 1 – figure supplement 1B, 4G, H and I). Thus, it is possible that Dronc's

function in response to necrosis occurs via a distinct mechanism. Importantly, the activity of Dronc in both

apoptosis and in AiP is influenced by the Dronc's upstream regulator, DIAP1 (Meier *et al.* 2000). DIAP1

modifies Dronc's CARD domain through K78 mono-ubiquitylation, which blocks both apoptosis (Kamber

Kaya *et al.* 2017), and AiP (Fan and Bergmann 2008, Fogarty and Bergmann 2017). Thus, we expressed

DIAP1 (*UAS-DIAP1*) following ablation with *DC^GluRI* and found that it strongly suppresses the number of

cDcp-1-positive cells at the WE and the NiA/NiCP in the LP (Figure 7I and K), but strikingly has no effect on

regenerative proliferation at 36 hr (Figure 7J and L). These results demonstrate a key role for the initiator

caspase *Dronc* in promoting regenerative proliferation following necrosis, which is not affected by DIAP1,

and therefore is likely separate from its role in apoptosis and the AiP mechanism.

Taken together, our data suggest a model in which necrosis leads to the establishment of distinct cell

populations important for regeneration (Figure 7N and O). After injury, cells at the immediate WE undergo

JNK-mediated apoptosis and contribute to proliferation via the established AiP mechanism (Figure 7O, left),

while cells at a distance from the injury in the LP activate Dronc via an unknown DAMP-like signal(s) that

occurs independent of JNK. Some of these cells go on to activate effector caspases at levels high enough to

result in apoptosis (NiA, Figure 7O, right), while others activate these caspases at a low enough level to be

detectable but insufficient to induce death (NiCP, Figure 7O right). Instead, these cells persist in the tissue

late into the repair process, where they promote proliferation via a novel non-apoptotic and AiP-independent

function of the initiator caspase Dronc.

Discussion

An important and often overlooked factor when studying regeneration is the type of injury – and consequently the type of cell death it causes – that can significantly impact repair processes. The existence of conserved signaling events that promote recovery has been well established in the context of apoptosis (Fogarty and Bergmann 2017, Perez-Garijo 2018), while the importance of such events in necrosis are less well established. Here, we have investigated the genetic events that occur in the aftermath of necrosis and how they influence the ability of a tissue to recover and regenerate. We previously showed that wing discs can regenerate effectively in the face of necrotic cell death (Klemm *et al.* 2021), triggering local JNK-dependent AiP at the WE, which is likely in response to tissue disruption (La Marca and Richardson 2020), while also inducing caspase activity in cells at a distance from the wound. This induction is independent of JNK signaling and, alongside having effector caspase activity, these cells can be marked by TUNEL and can be blocked by inhibiting the apoptotic pathway (Klemm *et al.* 2021). As such, we determined that these cells are undergoing PCD and named this phenomenon Necrosis-induced Apoptosis (NiA). We also showed that both NiA and AiP at the wound contribute to regeneration. Our current work further characterizes the NiA phenomenon, and we have now shown that cells undergoing NiA actually comprise two populations with separate behaviors. Upon necrosis, cells of the LP appear to activate effector caspases, indicated by cDcp-1 antibody staining and the transgenic reporter GC3Ai. While a proportion of these cells develop apoptotic morphology, round up and are cleared over time as part of NiA, a large number of cells appear able to persist despite the presence of caspases, where they promote regenerative proliferation dependent on the initiator caspase Dronc (Figure 7N). To reflect these new findings, we have called these Necrosis-induced Caspase Positive (NiCP) cells. While NiCP appear to promote proliferation, it remains unclear whether they do so through cell autonomous or non-autonomous effects. We were able to determine that NiCP do not utilize AiP as a potential mechanism, as these events occur in the absence of JNK, and exhibit none of the established hallmarks of AiP, including the presence and requirement for ROS, the ability to be blocked by DIAP1, or the production of mitogens such as Wg and Dpp. However, it is possible that other signaling factors may be secreted by these NiCP cells. Alternatively, the observed non-apoptotic caspase activity in NiCP cells may interact with intracellular factors to promote cell-autonomous proliferation. In either case, our results have identified an important role for Dronc in promoting regeneration in response to necrotic cell death.

The signal from necrotic cells that leads to NiA/NiCP remains unknown.

Although we have shown that necrosis leads to NiCP cells and NiA at a distance from the site of injury, the signal that leads to these events upon wounding is still unknown. Cells undergoing necrosis release DAMPs, a category of molecules that includes both common cellular contents as well as specific proteins, both of which can produce downstream responses like inflammation and the activation of effectors that promote a healing response (Gordon *et al.* 2018). DAMPs of both categories have been demonstrated in *Drosophila*; in

apoptotic-deficient larvae, circulating DAMPs in the hemolymph can constitutively activate immune signaling

in the fat body (Nishida *et al.* 2024), while specific factors such as α-actinin (Gordon *et al.* 2018), and

HMGB1 (Nishida *et al.* 2024) are known to possess DAMP activity. Recent work describing an *in vivo* sensor

for HMGB1 demonstrates the release of this DAMP from wing disc cells in response to a necrotic stimulus

similar to that used here (Nishida *et al.* 2024), making this an important candidate to test for a potential role

in producing NiCP. However, it is equally possible that the causative DAMP(s) is/are one or more of the

common fundamental cellular components released upon cell lysis, which would be more challenging to

manipulate and test. Considering the potential diversity and variable nature of DAMPs, it may be more

feasible to instead identify the downstream PRRs that are required to interpret this unknown signal. Several

groups of genes can act as PRRs, most notably the Toll-like receptor (TLR) family, various members of

which can respond to DAMPs (Ming *et al.* 2014, Gong *et al.* 2020). Nine TLR genes have been identified in

Drosophila, which have assorted roles in development and innate immunity (Anthoney *et al.* 2018), and have

been implicated in DAMP sensing (Ming *et al.* 2014). Thus, it would be valuable to test the requirement for

TLRs, alongside other suspected PRRs such as scavenger receptors (Cao 2016, Roh and Sohn 2018, Gong

et al. 2020) for their requirement to cause NiCP.

An additional approach to elucidating how these cells are generated is by leveraging our finding that both

JAK/STAT and WNT signaling seem to block NiCP/NiA in the disc. JAK/STAT signaling promotes survival of

cells in response to stress by repressing JNK signaling, thus minimizing JNK-mediated apoptosis (La

Fortezza *et al.* 2016). Although NiCP appear to avoid damage-induced JAK/STAT upregulation in the pouch,

it is unlikely they are regulated by this mechanism, since NiCP cells occur independent of JNK activity

(Klemm *et al.* 2021). However, within the hinge, which is completely devoid of NiCP, JAK/STAT protects

cells from apoptosis potentially by upregulating *DIAP1* (Verghese and Su 2016) and *zfh2* (Verghese and Su

2018), while Wg represses the transcription of *rpr*, (Verghese and Su 2016). Both signaling pathways are

required autonomously in hinge cells for their ability to replace and regenerate ablated pouch tissue

(Verghese and Su 2016, Ledru *et al.* 2022). Our results show that DIAP1 can prevent cDcp-1 activation in

NiCP in the pouch, but not the corresponding proliferation that they produce, suggesting one reason that

NiCP is not seen in the hinge is possibly due to the high DIAP1 threshold that results from developmental

JAK/STAT activity, which blocks Dronc's ability to activate effector caspases in this region. Thus, it remains

to be seen whether lowering DIAP1 levels in the hinge, or manipulating Wg to allow *rpr* expression alongside

changes in JAK/STAT, could allow NiCP to form in the hinge, and whether this might lead to regenerative

proliferation in this region. Finally, the observation that both NiCP cells and their associated proliferative

effects appear to occur independent of JNK signaling, which is normally central in models of stress and

damage (Pinal *et al.* 2019), could help to identify the upstream signaling that leads to NiCP following

necrosis. Together, these approaches could provide essential information as to the underlying genetic and

cellular events that connect the lysis of cells during necrosis to the formation of NiCP cells required for
regeneration.

The persistence of NiCP cells versus their elimination via NiA.

Our work shows that cells in the LP can either persist as NiCP and contribute to regenerative proliferation, or
progress to apoptosis as part of NiA, but how this decision is made is unknown. During apoptosis in the wing
disc, a threshold level of effector caspases must be reached for the cell to complete PCD (Florentin and
Arama 2012). Thus, we hypothesize that DAMP signals from necrotic cells may result in inconstant levels of
effector caspase activity in cells of the LP – some with high caspases that advance to apoptosis (NiA),
recognized by the changes in morphology and position in the disc characteristic of PCD, which are ultimately
cleared, and others (the NiCP cells) that have low enough caspase activity levels to survive, but this activity
can still be detected by sensitized reagents. This is supported by our observations made using effector
caspase-based lineage tracing using CasExpress (Ding *et al.* 2016), in which caspases can only be
detected in NiCP cells by lowering the detection threshold. Alternatively, since CasExpress is a membrane-
based reporter (Ding *et al.* 2016), it is also possible that effector caspases within NiCP cells have a different
subcellular localization that reflects a non-apoptotic function, thus preventing reporter activation, rather than
a difference in expression level or activity. The idea that caspases have different functions, both apoptotic
and non-apoptotic, based on localization is well established, such as the targeting of specific cleavage
substrates in distinct subcellular compartments (Brown-Suedel and Bouchier-Hayes 2020), or the trafficking
of Dronc to the membrane to promote ROS (Amcheslavsky *et al.* 2018). Whatever the mechanism is that
distinguishes between elimination via NiA or persistence as NiCP, the proportion of LP cells that participate
in each remains difficult to assay due to the dynamic nature of cell death (Nano and Montell 2024) and
overall variability of the NiCP/NiA phenotype. Anecdotal evidence from our experiments examining GC3Ai-
labelled cells over time suggests the majority of cells in fact do not undergo apoptosis and are therefore
NiCP. Importantly, however, while our experiments demonstrate that GC3Ai labeled NiCP increase over
time, it is unclear whether this increase in signal abundance is due to cells with newly activated caspases
that cleave and activate GC3Ai, or if GC3Ai labeled NiCP are instead proliferating. The use of other tools
with potentially different sensitivities to effector caspases, such as Apoliner (Bardet *et al.* 2008), or CD8-
PARP-Venus (Florentin and Arama 2012), may shed further light on this issue. Nevertheless, it remains to
be seen how a cell becomes NiCP or undergoes NiA and dies, and whether distinct levels of caspases
(initiator or effector) their localization, or a different attribute is responsible.

How does NiCP contribute to regeneration?

One of the most important questions that future work must address is how the phenomena we have
identified lead to regeneration following necrosis. Our experiments suggest that different caspase-positive

populations of cells may contribute to regenerative proliferation: Firstly, the smaller number of apoptotic cells

at the WE generated by epithelial disruption likely contribute by the established process of JNK-dependent

AiP (Fogarty and Bergmann 2017). Secondly, the caspase-surviving NiCP cells in the LP that contribute via

the initiator caspase Dronc. Crucially, this non-apoptotic function of Dronc cannot be inhibited by the

expression of DIAP1, which has previously been shown to block both the apoptotic and non-apoptotic AiP

functions of Dronc (Kamber Kaya *et al.* 2017). Thus, our results suggest that Dronc acts via a different

mechanism to induce growth in response to necrosis. The current understanding of how Dronc functions at

the molecular level may provide valuable clues as to how. DIAP1 suppresses Dronc through its E3 ubiquitin

ligase activity, which mono-ubiquitylates the caspase recruitment (CARD) domain of Dronc to suppress both

apoptotic and non-apoptotic functions related to AiP (Kamber Kaya *et al.* 2017). Here, we find that the

ectopic expression of *DIAP1* or just the CARD-containing pro-domain of Dronc ($Dronc^{DN}$) cannot block

regenerative proliferation, while by contrast heterozygosity for the $Dronc^{I29}$ null allele that can inhibit AiP

(Kamber Kaya *et al.* 2017) is sufficient to block NiCP-mediated proliferation. Thus, regulation of Dronc

activity in the context of necrosis may not rely on the modification of its CARD domain. The documented

functions of Dronc have also shown a requirement for its catalytic domain; mutations in this domain can

block the activation of Drice and AiP-induced overgrowth (Fan *et al.* 2014). Thus, the inability of DIAP1 to

suppress NiCP-mediated proliferation may be due to the catalytic activity of Dronc. Importantly, although the

CARD and catalytic domains are important points of regulation for Dronc activity, it has also been shown

that damage-specific context cues are also vital. For example, the ectopic expression of $Dronc^{K78R}$, a mutant

that renders it irrepressible by DIAP1, might be expected to induce significant apoptotic cell death. However,

this is not the case unless its structural binding partner, the APAF1 ortholog encoded by *Dark*, is expressed

alongside (Shapiro *et al.* 2008), demonstrating the importance of stoichiometry between Dronc and Dark for

the apoptotic function of Dronc. As such, further investigation into Dronc's functional domains and context-

dependent interactions with respect to its CARD and catalytic domains as well as its binding partners,

including DIAP1 and Dark, will be necessary to understand how Dronc is involved in promoting regenerative

proliferation in response to necrotic injury.

NiCP as a general mechanism to promote regeneration

Although having initially been characterized for their central role in apoptosis, many non-apoptotic functions

of caspases have since been discovered, revealing them to be dynamic regulators of diverse process

including cell fate specification, cellular remodeling, tissue growth, development, metabolism and others

(Shinoda *et al.* 2019, Wang and Baker 2019, Su 2020). Studies of caspase signaling during regeneration

have revealed essential aspects of non-apoptotic caspase activity, such as initiator and effector-dependent

models of AiP that contribute to repair (Ryoo and Bergmann 2012, Fogarty and Bergmann 2017). The

contrasts in caspase functions that we have observed between apoptotic and necrotic damage, despite

ultimately resulting in comparable levels of regeneration in an organ (Klemm *et al.* 2021), underscores the

nuance that exists in damage signaling between different injury contexts. It is clear that caspase activity in

response to injury as a mechanism to promote regeneration is a highly conserved process that occurs in

many organisms, regardless of tissue identify or type of damage incurred (Bergmann and Steller 2010, Vriz

et al. 2014, Fuchs and Steller 2015, Perez-Garijo and Steller 2015, Fogarty and Bergmann 2017, Perez-

Garijo 2018). Our findings reinforce the idea that much remains to be understood about the role of caspases

in tissue repair, and that the nature of an injury – and therefore the type of cell death involved – is an

important aspect that must also be considered.

Materials and Methods

Drosophila stocks

Flies were cultured in conventional dextrose fly media at 25^{o}C with 12h light–dark cycles. The recipe for dextrose media contains 9.3 g agar, 32 g yeast, 61 g cornmeal, 129 g dextrose, and 14 g tegosept in 1 L distilled water. Genotypes for each figure panel are listed in the Supplementary Genotypes file. Fly lines used as ablation stocks are as follows: hs-FLP; hs-p65; salm-LexADBD, DVE>>GAL4 (DC^{NA}), hs-FLP; hs-p65; salm-LexADBD/ TM6C, sb (DC^{NA} no GAL4), hs-FLP; lexAop-GluR1^{LC}, hs-p65/ CyO; salm-LexADBD, DVE>>GAL4/ TM6B, Tb (DC^{GluR1}), hs-FLP; lexAop-GluR1^{LC}, hs-p65/CyO; salm-LexADBD/TM6C, sb (DC^{GluR1} no GAL4), hs-FLP;lexAOp-GluR1^{LC}/ CyO; salm-LexADBD, hh-GAL4/ TM6B, Tb (DC^{GluR1} hh-GAL4), hs-FLP;lexAop-hepCA, hs-p65/CyO; salm-DBD, DVE>>GAL4/ TM6B, Tb (DC^{hepCA}), UAS-GluR1; tubGAL80^{ts}, and tubGAL80^{ts}. The stock DR^{WNT}-GAL80 was used to limit *UAS-* transgenes to the lateral pouch (LP) where NiA occur (Klemm *et al.* 2021), while *R85E08-GAL4* was used to drive *UAS-* expression at the wound edge (WE). The following stocks were obtained from Bloomington Drosophila Stock Center: w^{1118} (BL#3605), UAS-y^{RNAi} (BL#64527), UAS-p35 (BL#5073), UAS-GC3Ai (II, BL#84346), UAS-GC3Ai (III, BL#84343), rn-GAL4 (BL#7405), hh-GAL4 (BL#600186), ptc-GAL4 (BL#2017), pnr-GAL4 (BL#), (BL#25758), nub-GAL4 (BL#25754), R73G07-GAL4(BL#39829), UAS-Zfh2^{RNAi} (BL#50643), UAS-wg^{RNAi} (BL#32994), UAS-Stat92E^{RNAi} (BL#35600), dpp-lacZ (BL#8412), wg-lacZ (BL#50763), spi-lacZ (BL#10462), UAS-p35 (II, BL#5072), UAS-p35 (III, BL#5073), AP-1-GFP (Chatterjee and Bohmann 2012), act>>GAL4, UAS-RFP (BL#30558), DBS-GFP (III, BL#83130), CasExpress (BL#65419), G-TRACE (III, BL#28281), tubGAL80^{ts} (II, BL#7019), tubGAL80^{ts} (III, BL#7017), 10xSTAT-GFP (BL#), UAS-dome^{RNAi} (BL#32860), UAS-hop48A (BL#), PCNA-GFP (BL#25749), UAS-Cat (BL#24621), UAS-Sod1 (BL#24754), UAS-Duox^{RNAi} (BL#32903), mol-lacZ (BL#12173), UAS-Dronc^{DN} (BL#58992), UAS-DIAP1 (BL#6657), and dronc^{I29}/TM3, Sb (BL#98453). UAS-mir(RHG) was gifted from the Hariharan lab at UC Berkeley. vgQE-lacZ was gifted from Tin Tin Su. UAS-GluR1^{LC} (Liu *et al.* 2013) was gifted from the Xie lab at Stowers Institute.

Ablation experiments

DUAL Control ablation with *DVE>>GAL4*. DUAL Control experiments were performed essentially as described in Harris et al. (2020). Briefly, experimental crosses were cultured at 25°C and density controlled at 50 larvae per vial. Larvae were heat shocked on day 3.5 of development (84 hr after egg deposition (AED)) by placing vials in a 37°C water bath for 45 min, followed by a return to 25°C. Larvae were allowed to recover for 18 hr before being dissected, fixed and immunolabeled, unless otherwise indicated. *UAS-y^{RNAi}* and *UAS*-GFP were used as control lines for RNAi-based experiments. w^{1118} was used as a control for *dronc^{I29}* experiments. The *DVE>>GAL4* driver drives expression in the wing pouch, allowing for the regenerating wound edge cells and NiA cells to be targeted for interrogation (Klemm *et al.* 2021). The

DR^{WNT}-GAL80 transgene (Klemm *et al.* 2021) was included as necessary to restrict *UAS-* expression to the

lateral pouch (LP), where NiA/NiCP occurs.

DUAL Control ablation without *DVE>>GAL4*. To restrict *UAS-* expression to wound edge apoptotic cells,

a version of DUAL Control lacking the DVE>>GAL4 (*DC^{GluR1} no GAL4*) was crossed to the *R85E08-GAL-4*

driver. *DC^{GluR1} no GAL4* experiments were performed along the same parameters as *DC^{GluR1}* experiments.

The *R85E08-GAL4* driver was used alongside *DC^{GluR1} no GAL4*, when necessary, to target *UAS-* expression

to the wound edge (WE), where the JNK-dependent wound edge apoptotic cells occur.

DUAL Control ablation with *hh-GAL4*. DUAL Control flies bearing *hh-GAL4* (*DC^{GluR1} hh^{ts}*) were cultured at

18°C and density controlled at 50 larvae per vial. *tubGAL80^{ts}* was included to conditionally express *UAS-*

based constructs after ablation. Larvae were heat-shocked on day 7 of development (168 hr AED) for 45

min at 37°C, followed by incubation at 30°C to inactivate *tubGAL80^{ts}* and permit *UAS*-based expression.

Larvae were held at 30°C for 18 hr before being dissected, fixed, and immunolabeled.

GAL4/UAS **ablation.** *GAL4/UAS*-based ablation experiments were performed essentially as described in

Smith-Bolton et al. (2009). Briefly, larvae bearing either *UAS-GluR1;tubGAL80^{ts}* were cultured at 18°C and

density controlled at 50 larvae per vial. Larvae upshifted on day 7 of development (168 hr AED) for 20 hr at

30°C and were either immediately dissected (denoted as 0 hr) or were allowed to recover for 24 hr before

being dissected, fixed, and imaged. *tubGAL80^{ts}* flies were used as a non-ablating control.

*FLP/*FRT **ablation experiments.** To generate clonal patches of *UAS-GluR1;UAS-RFP*-expressing cells,

flies of the genotype *hs-FLP; AP-1-GFP; act>>GAL4, UAS-RFP/ T(2:3)SM6A, TM6B, Tb* were crossed to

flies bearing *UAS-GluR1; tubGAL80^{ts}*. Larvae were cultured at 18°C and heat shocked in a 37°C water bath

for 10 min at 82 hr AEL, returned to 18°C, and upshifted to 30°C for 20 hr at 168 h AEL, followed by

dissection and immunostaining. *tubGAL80^{ts}* flies were used as a non-ablating control.

Regeneration scoring and wing measurements.

Wings of adult flies from heat shocked larvae were scored and measured after genotype blinding by another

researcher. Scoring was performed on anesthetized adults by binning into a regeneration scoring category

(Harris *et al.* 2020, Klemm *et al.* 2021). Wing measurements were performed by removing wings, mounting

in Permount solution (Fisher Scientific) and imaged using a Zeiss Discovery.V8 microscope. Wing area was

measured using the Fiji software. Male and female adults were measured separately to account for sex

differences in wing size using a reproducible measuring protocol that excludes the variable hinge region of

the wing (details of measuring protocol available on request). Statistics were performed using GraphPad
Prism 10.0.
716

Immunohistochemistry

Larvae were dissected in 1 x PBS followed by a 20 min fix in 4 % paraformaldehyde in PBS (PFA). After 3
washes in 0.1 % PBST (1 x PBS + 0.1 % Triton-X), larvae were washed in 0.3% PBST and then blocked in
0.1 % PBST with 5 % normal goat serum (NGS) for 30 min. Primary staining was done overnight at 4°C, and
secondary staining was done for 4 hr at room temperature. The following primary antibodies were obtained
from the Developmental Studies Hybridoma Bank: mouse anti-Nubbin (1:25), mouse anti-Wg (1:100),
mouse anti-Mmp1 C-terminus (1:100), mouse anti-Mmp1 catalytic domain (1:100), mouse anti-LacZ (1:100),
mouse anti-discs large (1:50), mouse anti-yH2Av (1:100), and rat anti-DE-cadherin (1:100). Rabbit anti-
cDcp-1 (1:1000), mouse anti-PH3 (1:500), and rabbit anti-HA (1:1000) were obtained from Cell Signaling
Technologies. Rat anti-Zfh-2 was generously gifted by Chris Doe. Anti-rabbit 647, anti-rat 647, anti-mouse
555, and anti-mouse 488 secondary antibodies were obtained from Invitrogen and used at a 1:500 dilution.
DAPI (1:1000) was used as a counterstain. Images were obtained on a Zeiss AxioImager.M2 with ApoTome
and a Leica TCS SP8 LCSM (NIH SIG award 1 S10 OD023691-01) housed in the Regenerative Medicine
Imaging Facility at Arizona State University. For each experiment at least 15 discs were analyzed prior to
choosing a representative image. Images were processed using Affinity Photo.
732

733 **EdU staining, TUNEL assay, and DHE staining**

734 **EdU.** The Click-It EdU Alexa Fluor 555 Imaging Kit (Invitrogen C10338) was used to assay cell proliferation.
735 Briefly, imaginal discs were dissected and labeled with 1 µl EdU in 1 ml PBS for 20 min, fixed in 4 % PFA for
736 20 min and immunolabeled (as necessary), followed by a 30 min Click-It reaction that was performed as
737 directed in the EdU manual.
738

TUNEL. The TUNEL assay was performed with the ApopTag Red In Situ Apoptosis Detection Kit (Millipore
S7165). Dissected larvae were fixed in 4 % paraformaldehyde, followed by a 10 min wash in 75 ml
equilibration buffer. Discs were then submerged in 55 µl working strength TdT enzyme for 3 hr at 37°C. The
reaction was stopped by adding 1ml stop/wash buffer and incubating for 10min at room temperature,
followed by three washes in PBS. Immunolabeling was performed by incubating the tissue preps with 65 µl
of anti-digoxigenin rhodamine overnight at room temperature.
745

746 **Dihydroethidium (DHE)**. DHE labeling was performed by incubating freshly dissected wing imaginal discs
747 in Schneider's Media with 1 µl of 10 mM DHE reconstituted in 1 ml DMSO (for a working concentration of 10
748 µm DHE) for 10 min, followed by three 5 min washes in PBS and immediately mounting and imaging.

Quantification and Statistical Analysis

Adult wings, mean fluorescence intensity, and cell counts were measured using Fiji. GraphPad Prism 10.0 was used for statistical analysis and graphical representation. Graphs depict the mean of each treatment, while error bars represent the standard deviation. The mean fluorescence intensity of EdU labeling was quantified in Fiji. In each experiment, the mean fluorescence intensity (MFI) of the wing pouch was normalized to the MFI of the entire disc. The mean and standard deviation for each normalized treatment was calculated and used for statistical analysis. The sample size and P values for all statistical analyses are indicated in the figure legends. Statistical significance was evaluated in Prism 10.0 using a Student's T-test or a one-way ANOVA with a multiple comparisons test.

Data and reagent availability

Stocks are available upon request and details of stocks and reagents used in this study are available in the materials and methods. The authors affirm that all data necessary for confirming the conclusions of the article are present within the article, figures, and tables.

Acknowledgements

The authors would like to thank Dr. Tin Tin Su of UC Boulder, Dr. Chris Doe of the University of Oregon, and Dr. Tian Xie of the Stowers Institute for their generous gift of stocks and reagents. We thank the current members of the Harris lab for useful input and feedback. We thank the Bloomington Drosophila Stock Center and Developmental Studies Hybridoma Bank for stocks and reagents.

Funding

This work was supported by a grant from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) R21HD102765-01 and the National Institute of General Medical Sciences (NIGMS) R01GM147615 to Robin Harris.

Author contributions

Jacob Klemm, Conceptualization, Investigation Data curation, Formal analysis, Validation, Writing – original draft, Writing – review and editing; Chloe Van Hazel, Resources; Robin Harris, Supervision, Funding acquisition, Investigation, Resources, Writing – review and editing.

Conflicts of interest

782   The authors declare that there is no conflict of interest.

788
789

References

References

1. Amcheslavsky, A., S. Wang, C. E. Fogarty, J. L. Lindblad, Y. Fan and A. Bergmann (2018). "Plasma Membrane
Localization of Apoptotic Caspases for Non-apoptotic Functions." *Dev Cell* **45**(4): 450-464 e453.

2. Anthoney, N., I. Foldi and A. Hidalgo (2018). "Toll and Toll-like receptor signalling in development."
Development **145**(9).

3. Ashkenazi, A. and G. Salvesen (2014). "Regulated cell death: signaling and mechanisms." *Annu Rev Cell Dev*
Biol **30**: 337-356.

4. Bach, E. A., L. A. Ekas, A. Ayala-Camargo, M. S. Flaherty, H. Lee, N. Perrimon and G. H. Baeg (2007). "GFP
reporters detect the activation of the Drosophila JAK/STAT pathway in vivo." *Gene Expr Patterns* **7**(3): 323-331.

5. Baena-Lopez, L. A., L. Arthurton, M. Bischoff, J. P. Vincent, C. Alexandre and R. McGregor (2018). "Novel
initiator caspase reporters uncover previously unknown features of caspase-activating cells." *Development*
145(23).

6. Bardet, P. L., G. Kolahgar, A. Mynett, I. Miguel-Aliaga, J. Briscoe, P. Meier and J. P. Vincent (2008). "A
fluorescent reporter of caspase activity for live imaging." *Proc Natl Acad Sci U S A* **105**(37): 13901-13905.

7. Beira, J. V. and R. Paro (2016). "The legacy of Drosophila imaginal discs." *Chromosoma* **125**(4): 573-592.

8. Bergantinos, C., M. Corominas and F. Serras (2010). "Cell death-induced regeneration in wing imaginal discs
requires JNK signalling." *Development* **137**(7): 1169-1179.

9. Bergmann, A. and H. Steller (2010). "Apoptosis, stem cells, and tissue regeneration." *Sci Signal* **3**(145): re8.

10. Bonne, S. L. and S. S. Kadri (2017). "Evaluation and Management of Necrotizing Soft Tissue Infections."
Infect Dis Clin North Am **31**(3): 497-511.

11. Brown-Suedel, A. N. and L. Bouchier-Hayes (2020). "Caspase-2 Substrates: To Apoptosis, Cell Cycle Control,
and Beyond." *Front Cell Dev Biol* **8**: 610022.

12. Cao, X. (2016). "Self-regulation and cross-regulation of pattern-recognition receptor signalling in health and
disease." *Nat Rev Immunol* **16**(1): 35-50.

13. Chatterjee, N. and D. Bohmann (2012). "A versatile PhiC31 based reporter system for measuring AP-1 and
Nrf2 signaling in Drosophila and in tissue culture." *PLoS One* **7**(4): e34063.

14. Chera, S., L. Ghila, K. Dobretz, Y. Wenger, C. Bauer, W. Buzgariu, J. C. Martinou and B. Galliot (2009).
"Apoptotic cells provide an unexpected source of Wnt3 signaling to drive hydra head regeneration." *Dev Cell*
17(2): 279-289.

15. Colon Plaza, S. and T. T. Su (2024). "Ionizing radiation induces cells with past caspase activity that contribute
to the adult organ in Drosophila and show reduced Loss of Heterozygosity." *Cell Death Discov* **10**(1): 6.

16. D'Arcy, M. S. (2019). "Cell death: a review of the major forms of apoptosis, necrosis and autophagy." *Cell Biol*
Int **43**(6): 582-592.

17. Ding, A. X., G. Sun, Y. G. Argaw, J. O. Wong, S. Easwaran and D. J. Montell (2016). "CasExpress reveals
widespread and diverse patterns of cell survival of caspase-3 activation during development in vivo." *Elife* **5**.

18. Diwanji, N. and A. Bergmann (2018). "An unexpected friend - ROS in apoptosis-induced compensatory
proliferation: Implications for regeneration and cancer." *Semin Cell Dev Biol* **80**: 74-82.

19. Evans, C. J., J. M. Olson, K. T. Ngo, E. Kim, N. E. Lee, E. Kuoy, A. N. Patananan, D. Sitz, P. Tran, M. T. Do,

K. Yackle, A. Cespedes, V. Hartenstein, G. B. Call and U. Banerjee (2009). "G-TRACE: rapid Gal4-based cell

lineage analysis in Drosophila." *Nat Methods* **6**(8): 603-605.

20. Fan, Y. and A. Bergmann (2008). "Apoptosis-induced compensatory proliferation. The Cell is dead. Long live

the Cell!" *Trends Cell Biol* **18**(10): 467-473.

21. Fan, Y. and A. Bergmann (2008). "Distinct mechanisms of apoptosis-induced compensatory proliferation in

proliferating and differentiating tissues in the Drosophila eye." *Dev Cell* **14**(3): 399-410.

22. Fan, Y., S. Wang, J. Hernandez, V. B. Yenigun, G. Hertlein, C. E. Fogarty, J. L. Lindblad and A. Bergmann

(2014). "Genetic models of apoptosis-induced proliferation decipher activation of JNK and identify a requirement

of EGFR signaling for tissue regenerative responses in Drosophila." *PLoS Genet* **10**(1): e1004131.

23. Festjens, N., T. Vanden Berghe and P. Vandenabeele (2006). "Necrosis, a well-orchestrated form of cell

demise: signalling cascades, important mediators and concomitant immune response." *Biochim Biophys Acta*

1757(9-10): 1371-1387.

24. Florentin, A. and E. Arama (2012). "Caspase levels and execution efficiencies determine the apoptotic

potential of the cell." *J Cell Biol* **196**(4): 513-527.

25. Fogarty, C. E. and A. Bergmann (2017). "Killers creating new life: caspases drive apoptosis-induced

proliferation in tissue repair and disease." *Cell Death Differ* **24**(8): 1390-1400.

26. Fogarty, C. E., N. Diwanji, J. L. Lindblad, M. Tare, A. Amcheslavsky, K. Makhijani, K. Bruckner, Y. Fan and A.

Bergmann (2016). "Extracellular Reactive Oxygen Species Drive Apoptosis-Induced Proliferation via Drosophila

Macrophages." *Curr Biol* **26**(5): 575-584.

27. Fox, D. T., E. Cohen and R. Smith-Bolton (2020). "Model systems for regeneration: Drosophila." *Development*

147(7).

28. Fuchs, Y. and H. Steller (2015). "Live to die another way: modes of programmed cell death and the signals

emanating from dying cells." *Nat Rev Mol Cell Biol* **16**(6): 329-344.

29. Gong, T., L. Liu, W. Jiang and R. Zhou (2020). "DAMP-sensing receptors in sterile inflammation and

inflammatory diseases." *Nat Rev Immunol* **20**(2): 95-112.

30. Gordon, O., C. M. Henry, N. Srinivasan, S. Ahrens, A. Franz, S. Deddouche, P. Chakravarty, D. Phillips, R.

George, S. Kjaer, D. Frith, A. P. Snijders, R. S. Valente, C. J. Simoes da Silva, L. Teixeira, B. Thompson, M. S.

Dionne, W. Wood and C. Reis e Sousa (2018). "alpha-actinin accounts for the bioactivity of actin preparations in

inducing STAT target genes in Drosophila melanogaster." *Elife* **7**.

31. Grasl-Kraupp, B., B. Ruttkay-Nedecky, H. Koudelka, K. Bukowska, W. Bursch and R. Schulte-Hermann

(1995). "In situ detection of fragmented DNA (TUNEL assay) fails to discriminate among apoptosis, necrosis, and

autolytic cell death: a cautionary note." *Hepatology* **21**(5): 1465-1468.

32. Hajibabaie, F., N. Abedpoor and P. Mohamadynejad (2023). "Types of Cell Death from a Molecular

Perspective." *Biology (Basel)* **12**(11).

33. Hakkarainen, T. W., N. M. Kopari, T. N. Pham and H. L. Evans (2014). "Necrotizing soft tissue infections:

review and current concepts in treatment, systems of care, and outcomes." *Curr Probl Surg* **51**(8): 344-362.

34. Hariharan, I. K. and F. Serras (2017). "Imaginal disc regeneration takes flight." *Curr Opin Cell Biol* **48**: 10-16.

35. Harris, R. E. (2023). "Investigating Tissue Regeneration Using the DUAL Control Genetic Ablation System."

Methods Mol Biol **2599**: 255-270.

36. Harris, R. E., M. J. Stinchfield, S. L. Nystrom, D. J. McKay and I. K. Hariharan (2020). "Damage-responsive,

maturity-silenced enhancers regulate multiple genes that direct regeneration in Drosophila." *Elife* **9**.

37. Hawkins, C. J., S. J. Yoo, E. P. Peterson, S. L. Wang, S. Y. Vernooy and B. A. Hay (2000). "The Drosophila

caspase DRONC cleaves following glutamate or aspartate and is regulated by DIAP1, HID, and GRIM." *J Biol*

Chem **275**(35): 27084-27093.

38. Hay, B. A., T. Wolff and G. M. Rubin (1994). "Expression of baculovirus P35 prevents cell death in

Drosophila." *Development* **120**(8): 2121-2129.

39. Herrera, S. C. and E. A. Bach (2019). "JAK/STAT signaling in stem cells and regeneration: from Drosophila to

vertebrates." *Development* **146**(2).

40. Huh, J. R., M. Guo and B. A. Hay (2004). "Compensatory proliferation induced by cell death in the Drosophila

wing disc requires activity of the apical cell death caspase Dronc in a nonapoptotic role." *Curr Biol* **14**(14): 1262-

1266.

41. Jaiswal, J., J. Egert, R. Engesser, A. A. Peyroton, L. Nogay, V. Weichselberger, C. Crucianelli, I. Grass, C.

Kreutz, J. Timmer and A. K. Classen (2023). "Mutual repression between JNK/AP-1 and JAK/STAT stratifies

senescent and proliferative cell behaviors during tissue regeneration." *PLoS Biol* **21**(5): e3001665.

42. Jung, Y., R. P. Witek, W. K. Syn, S. S. Choi, A. Omenetti, R. Premont, C. D. Guy and A. M. Diehl (2010).

"Signals from dying hepatocytes trigger growth of liver progenitors." *Gut* **59**(5): 655-665.

43. Kamber Kaya, H. E., M. Ditzel, P. Meier and A. Bergmann (2017). "An inhibitory mono-ubiquitylation of the

Drosophila initiator caspase Dronc functions in both apoptotic and non-apoptotic pathways." *PLoS Genet* **13**(2):

e1006438.

44. Karsch-Bluman, A., A. Feiglin, E. Arbib, T. Stern, H. Shoval, O. Schwob, M. Berger and O. Benny (2019).

"Tissue necrosis and its role in cancer progression." *Oncogene* **38**(11): 1920-1935.

45. Katsuyama, T., F. Comoglio, M. Seimiya, E. Cabuy and R. Paro (2015). "During Drosophila disc regeneration,

JAK/STAT coordinates cell proliferation with Dilp8-mediated developmental delay." *Proc Natl Acad Sci U S A*

112(18): E2327-2336.

46. Khan, S. J., S. N. F. Abidi, A. Skinner, Y. Tian and R. K. Smith-Bolton (2017). "The Drosophila Duox

maturation factor is a key component of a positive feedback loop that sustains regeneration signaling." *PLoS*

Genet **13**(7): e1006937.

47. Kim, J., A. Sebring, J. J. Esch, M. E. Kraus, K. Vorwerk, J. Magee and S. B. Carroll (1996). "Integration of

positional signals and regulation of wing formation and identity by Drosophila vestigial gene." *Nature* **382**(6587):

133-138.

48. Klemm, J., M. J. Stinchfield and R. E. Harris (2021). "Necrosis-induced apoptosis promotes regeneration in

Drosophila wing imaginal discs." *Genetics* **219**(3).

49. Konstantinidis, K., R. S. Whelan and R. N. Kitsis (2012). "Mechanisms of cell death in heart disease."
Arterioscler Thromb Vasc Biol **32**(7): 1552-1562.
50. La Fortezza, M., M. Schenk, A. Cosolo, A. Kolybaba, I. Grass and A. K. Classen (2016). "JAK/STAT signalling
mediates cell survival in response to tissue stress." *Development* **143**(16): 2907-2919.
51. La Marca, J. E. and H. E. Richardson (2020). "Two-Faced: Roles of JNK Signalling During Tumourigenesis in
the Drosophila Model." *Front Cell Dev Biol* **8**: 42.
52. Ledru, M., C. A. Clark, J. Brown, S. Verghese, S. Ferrara, A. Goodspeed and T. T. Su (2022). "Differential
gene expression analysis identified determinants of cell fate plasticity during radiation-induced regeneration in
Drosophila." *PLoS Genet* **18**(1): e1009989.
53. Li, F., Q. Huang, J. Chen, Y. Peng, D. R. Roop, J. S. Bedford and C. Y. Li (2010). "Apoptotic cells activate the
"phoenix rising" pathway to promote wound healing and tissue regeneration." *Sci Signal* **3**(110): ra13.
54. Li, M., S. Sun, J. Priest, X. Bi and Y. Fan (2019). "Characterization of TNF-induced cell death in Drosophila
reveals caspase- and JNK-dependent necrosis and its role in tumor suppression." *Cell Death Dis* **10**(8): 613.
55. Li, Z., W. Shao, X. Lv, B. Wang, L. Han, S. Gong, P. Wang and Y. Feng (2023). "Advances in experimental
models of osteonecrosis of the femoral head." *J Orthop Translat* **39**: 88-99.
56. Liu, K., Y. Li and L. Liu (2013). "Modeling calcium-overload mediated necrosis in Drosophila." *Methods Mol
Biol* **1004**: 203-213.
57. Madigan, J. P., H. L. Chotkowski and R. L. Glaser (2002). "DNA double-strand break-induced phosphorylation
of Drosophila histone variant H2Av helps prevent radiation-induced apoptosis." *Nucleic Acids Res* **30**(17): 3698-
3705.
58. Martin, R. and G. Morata (2018). "Regenerative response of different regions of Drosophila imaginal discs."
Int J Dev Biol **62**(6-7-8): 507-512.
59. Martin, R., N. Pinal and G. Morata (2017). "Distinct regenerative potential of trunk and appendages of
Drosophila mediated by JNK signalling." *Development* **144**(21): 3946-3956.
60. Masi, L., A. Falchetti and M. L. Brandi (2007). "Osteonecrosis in genetic disorders." *Clin Cases Miner Bone
Metab* **4**(1): 27-29.
61. Meier, P., J. Silke, S. J. Leevers and G. I. Evan (2000). "The Drosophila caspase DRONC is regulated by
DIAP1." *EMBO J* **19**(4): 598-611.
62. Ming, M., F. Obata, E. Kuranaga and M. Miura (2014). "Persephone/Spatzle pathogen sensors mediate the
activation of Toll receptor signaling in response to endogenous danger signals in apoptosis-deficient Drosophila."
J Biol Chem **289**(11): 7558-7568.
63. Monier, B., M. Gettings, G. Gay, T. Mangeat, S. Schott, A. Guarner and M. Suzanne (2015). "Apico-basal
forces exerted by apoptotic cells drive epithelium folding." *Nature* **518**(7538): 245-248.
64. Morata, G., E. Shlevkov and A. Perez-Garijo (2011). "Mitogenic signaling from apoptotic cells in Drosophila."
Dev Growth Differ **53**(2): 168-176.
65. Mulay, S. R., A. Linkermann and H. J. Anders (2016). "Necroinflammation in Kidney Disease." *J Am Soc
Nephrol* **27**(1): 27-39.

66. Nakada, H., M. Inoue, K. Furuya, H. Watanabe, K. Ikegame, Y. Nakayama, M. Ohmori and H. Nakagomi

(2019). "Fat necrosis after breast-conserving oncoplastic surgery." *Breast Cancer* **26**(1): 125-130.

67. Nano, M. and D. J. Montell (2024). "Apoptotic signaling: Beyond cell death." *Semin Cell Dev Biol* **156**: 22-34.

68. Nishida, H., A. B. Albero, K. Onoue, Y. Ikegawa, S. Sulekh, U. Sakizli, Y. Minami, S. Yonemura, Y. C. Wang

and S. K. Yoo (2024). "Necrosensor: a genetically encoded fluorescent sensor for visualizing necrosis in

Drosophila." *Biol Open* **13**(1).

69. Pellettieri, J., P. Fitzgerald, S. Watanabe, J. Mancuso, D. R. Green and A. Sanchez Alvarado (2010). "Cell

death and tissue remodeling in planarian regeneration." *Dev Biol* **338**(1): 76-85.

70. Perez-Garijo, A. (2018). "When dying is not the end: Apoptotic caspases as drivers of proliferation." *Semin*

Cell Dev Biol **82**: 86-95.

71. Perez-Garijo, A., F. A. Martin and G. Morata (2004). "Caspase inhibition during apoptosis causes abnormal

signalling and developmental aberrations in Drosophila." *Development* **131**(22): 5591-5598.

72. Perez-Garijo, A., F. A. Martin, G. Struhl and G. Morata (2005). "Dpp signaling and the induction of neoplastic

tumors by caspase-inhibited apoptotic cells in Drosophila." *Proc Natl Acad Sci U S A* **102**(49): 17664-17669.

73. Perez-Garijo, A., E. Shlevkov and G. Morata (2009). "The role of Dpp and Wg in compensatory proliferation

and in the formation of hyperplastic overgrowths caused by apoptotic cells in the Drosophila wing disc."

Development **136**(7): 1169-1177.

74. Perez-Garijo, A. and H. Steller (2015). "Spreading the word: non-autonomous effects of apoptosis during

development, regeneration and disease." *Development* **142**(19): 3253-3262.

75. Pinal, N., M. Calleja and G. Morata (2019). "Pro-apoptotic and pro-proliferation functions of the JNK pathway

of Drosophila: roles in cell competition, tumorigenesis and regeneration." *Open Biol* **9**(3): 180256.

76. Pinal, N., M. Martín, I. Medina and G. Morata (2018). "Short-term activation of the Jun N-terminal kinase

pathway in apoptosis-deficient cells of Drosophila induces tumorigenesis." *Nature Communications* **9**(1).

77. Robertson, S. A., J. A. Jeevaratnam, A. Agrawal and R. I. Cutress (2017). "Mastectomy skin flap necrosis:

challenges and solutions." *Breast Cancer (Dove Med Press)* **9**: 141-152.

78. Roh, J. S. and D. H. Sohn (2018). "Damage-Associated Molecular Patterns in Inflammatory Diseases."

Immune Netw **18**(4): e27.

79. Ryoo, H. D. and A. Bergmann (2012). "The role of apoptosis-induced proliferation for regeneration and

cancer." *Cold Spring Harb Perspect Biol* **4**(8): a008797.

80. Ryoo, H. D., T. Gorenc and H. Steller (2004). "Apoptotic cells can induce compensatory cell proliferation

through the JNK and the Wingless signaling pathways." *Dev Cell* **7**(4): 491-501.

81. Schott, S., A. Ambrosini, A. Barbaste, C. Benassayag, M. Gracia, A. Proag, M. Rayer, B. Monier and M.

Suzanne (2017). "A fluorescent toolkit for spatiotemporal tracking of apoptotic cells in living Drosophila tissues."

Development **144**(20): 3840-3846.

82. Shapiro, P. J., H. H. Hsu, H. Jung, E. S. Robbins and H. D. Ryoo (2008). "Regulation of the Drosophila

apoptosome through feedback inhibition." *Nat Cell Biol* **10**(12): 1440-1446.

83. Shinoda, N., N. Hanawa, T. Chihara, A. Koto and M. Miura (2019). "Dronc-independent basal executioner
caspase activity sustains Drosophila imaginal tissue growth." *Proc Natl Acad Sci U S A* **116**(41): 20539-20544.
84. Siegrist, S. E., N. S. Haque, C. H. Chen, B. A. Hay and I. K. Hariharan (2010). "Inactivation of both Foxo and
reaper promotes long-term adult neurogenesis in Drosophila." *Curr Biol* **20**(7): 643-648.
85. Su, T. T. (2020). "Non-apoptotic roles of apoptotic proteases: new tricks for an old dog." *Open Biol* **10**(8):
200130.
86. Tonnus, W., A. Belavgeni, F. Beuschlein, G. Eisenhofer, M. Fassnacht, M. Kroiss, N. P. Krone, M. Reincke, S.
R. Bornstein and A. Linkermann (2021). "The role of regulated necrosis in endocrine diseases." *Nat Rev*
Endocrinol **17**(8): 497-510.
87. Tseng, A. S., D. S. Adams, D. Qiu, P. Koustubhan and M. Levin (2007). "Apoptosis is required during early
stages of tail regeneration in Xenopus laevis." *Dev Biol* **301**(1): 62-69.
88. Venereau, E., C. Ceriotti and M. E. Bianchi (2015). "DAMPs from Cell Death to New Life." *Front Immunol* **6**:
422.
89. Verghese, S. and T. T. Su (2016). "Drosophila Wnt and STAT Define Apoptosis-Resistant Epithelial Cells for
Tissue Regeneration after Irradiation." *PLoS Biol* **14**(9): e1002536.
90. Verghese, S. and T. T. Su (2018). "Ionizing radiation induces stem cell-like properties in a caspase-dependent
manner in Drosophila." *PLoS Genet* **14**(11): e1007659.
91. Vriz, S., S. Reiter and B. Galliot (2014). "Cell death: a program to regenerate." *Curr Top Dev Biol* **108**: 121-
151.
92. Wang, L. H. and N. E. Baker (2019). "Salvador-Warts-Hippo pathway regulates sensory organ development
via caspase-dependent nonapoptotic signaling." *Cell Death Dis* **10**(9): 669.
93. Worley, M. I. and I. K. Hariharan (2022). "Imaginal Disc Regeneration: Something Old, Something New." *Cold*
Spring Harb Perspect Biol **14**(11).
94. Worley, M. I., L. Setiawan and I. K. Hariharan (2012). "Regeneration and transdetermination in Drosophila
imaginal discs." *Annu Rev Genet* **46**: 289-310.
95. Wu, M. Y., G. T. Yiang, W. T. Liao, A. P. Tsai, Y. L. Cheng, P. W. Cheng, C. Y. Li and C. J. Li (2018). "Current
Mechanistic Concepts in Ischemia and Reperfusion Injury." *Cell Physiol Biochem* **46**(4): 1650-1667.
96. Xu, D., Y. Li, M. Arcaro, M. Lackey and A. Bergmann (2005). "The CARD-carrying caspase Dronc is essential
for most, but not all,developmental cell death inDrosophila." *Development* **132**(9): 2125-2134.
97. Yang, X., H. Ren and J. Fu (2021). "Treatment of Radiation-Induced Brain Necrosis." *Oxid Med Cell Longev*
2021: 4793517.
98. Yang, Y., L. Hou, Y. Li, J. Ni and L. Liu (2013). "Neuronal necrosis and spreading death in a Drosophila
genetic model." *Cell Death Dis* **4**: e723.
1006

Article and author information

Jacob Klemm

School of Life Sciences, Arizona State University, Life Sciences E (LSE) 354, 427 Tyler Mall, Tempe, Arizona, 85287-4501

jwklemm@asu.edu

ORCID iD: 0000-0002-1414-5089

Chloe Van Hazel

School of Life Sciences, Arizona State University, Life Sciences E (LSE) 354, 427 Tyler Mall, Tempe, Arizona, 85287-4501

Robin Harris

School of Life Sciences, Arizona State University, Life Sciences E (LSE) 354, 427 Tyler Mall, Tempe, Arizona, 85287-4501

For correspondence: reharri4@asu.edu

ORCID iD: 0000-0001-6945-6741

Figure Legends

Figure 1. The NiA response is restricted to pouch cells following ablation. (**A**) A schematic of the DC^{GluR1} ablation scheme. This system utilizes a split LexA transcriptional activator that promotes the expression of *lexAop-GluR1* following a short heat shock (45 min. at 37°C). (**B**) A schematic of the different tissue domains and compartments targeted for *GAL4/UAS/GAL80^{ts}* (*GAL4^{ts}*) ablation. (**C**) A schematic of the temperature scheme used to induce necrosis with *GAL4^{ts}>GluR1* following the heat shift (20 hr at 30°C) induced inactivation of GAL80^{ts}. (**D**) A control disc for DUAL Control (*DC^{NA}*) bearing *lexAop-GFP*, which labels the domain targeted for ablation in DC^{GluR1} experiments. (**E**) A DC^{GluR1} ablated disc at 18 hr of regeneration bearing *lexAop-GFP*, yellow arrowheads indicate NiA cells, red arrowhead indicates potential NiA response in the posterior pleura. (**F**) A control disc for *R85E08-GAL4^{ts}* (*R85^{ts}>GFP*) ablation. (**G**) An *R85E08^{ts}>GluR1,GFP* ablated disc, arrowheads indicate the presence of NiA cells following ablation. (**H**) A control disc for whole pouch (*rn^{ts}>GFP*) ablation. (**I-J**) A *rn^{ts}>GluR1,GFP* ablated disc dissected immediately following the downshift (0 hr, **I**) and after 24 hr of regeneration (R24, **J**). (**K**) A control disc for notum ablation (*pnr^{ts}>GFP*). (**L-M**) *pnr^{ts}>GluR1,GFP* ablated discs at 0 hr (**L**) and 24 hr (**M**), open arrowheads highlight the absence of the Wg notum stripe following ablation. (**N**) A control disc for hinge ablation (*R73G07^{ts}>GFP*). (**O-P**) *R73G08^{ts}>GluR1,GFP* ablated discs at 0 hr (**O**) and 24 hr (**P**). Open arrowhead in (**O**) indicates the absence of NiA in the adjacent wing pouch. (**Q**) A control disc for posterior compartment ablation (*hh^{ts}>GFP*). (**R-S'**) *hh^{ts}>GluR1,GFP* ablated discs at 0 hr (**R-R'**) and 24 hr (**S-S'**). Dotted lines in (**R'**) and (**S'**) show an absence of NiA in the anterior hinge. (**T**) A quantification of the NiA response in the anterior pouch, hinge, and notum in control (n= 13), 0 hr (n= 14), and 24 hr (n= 12) discs in response to *hh* ablation. Error bars represent the standard deviation. NiA are defined as cDcp-1-positive (cDcp-1+ve), GFP-negative (GFP-ve) cells. See Supplementary Genotypes file for exact genotypes.

1049

1050

Figure 1 – figure supplement 1. (**A**) A *nub^{ts}>GluR1,GFP* ablated disc at 0 hr. (**B**) A *DC^{GluR1}* ablated disc
highlighting the overlap between wound edge apoptosis and the JNK target Mmp1 (arrowhead) and the
absence of JNK signaling in NiA cells. (**C**) A control disc for anterior compartment ablation (*ptc^{ts}>RFP*). (**D-**
E') *ptc^{ts}>GluR1,RFP* ablated discs at 0 hr (**D-D'**) and 24 hr (**E-E'**). Dotted lines in (**D'**) and (**E'**) highlight the
lack of NiA cells in the hinge. (**F**) A quantification of the NiA response to *ptc* ablation in the pouch, hinge, and
notum of control (n= 14), 0 hr (n= 15), and 24 hr (n= 14) discs, with NiA being defined as cDcp-1-positive
(+ve), RFP-negative (-ve) cells. Error bars represent the standard deviation. (**G**) A control disc for *FLP/FRT*
clonal ablation, where actin "flipout" GAL4 (*act^{ts}>>RFP*) drives the expression of *UAS-RFP* in clonal patches
of cells after a short heat shock (see materials and methods for details). (**H-H''**) A wing disc with clonal
patches of *UAS-RFP*, *UAS-GluR1* expressing cells. (**H'**) is a close-up of pouch and hinge clones with
arrowheads pointing to NiA cells, while (**H''**) is a close-up of notum clones. (**I**) A quantification of the number
of NiA cells induced following ablation with *act^{ts}>>GluR1* (n= 11), *R85E08^{ts}>GluR1* (n= 10), *ptc^{ts}>GluR1* (n=
15), and *hh^{ts}>GluR1* (n= 14), highlighting that NiA number increases with increasing area of ablation. (**J**) A
leg and a haltere disc with clonal patches of *UAS-RFP, UAS-GluR1* expressing cells. Small patches of
cDcp-1+ve, RFP-ve cells in both discs suggests that the NiA response may occur in other imaginal discs
and is not limited to the wing disc. See Supplementary Genotypes file for exact genotypes.

1067

Figure 2. NiA formation in pouch cells is regulated by the Wg and JAK/STAT pathways. (**A**). A control
disc (DC^{NA}) bearing the *10xSTAT-GFP* reporter, showing the hinge-specific Stat92E activity that is normally
absent from the wing pouch. (**B**) A DC^{GluR1} ablated disc bearing *10xSTAT-GFP*. Following damage, pouch-
specific reporter expression is observed at high levels at the wound edge (arrowhead) and low levels at the
NiA area of the pouch (open arrowheads). (**C-E**) DC^{GluR1}>>y^{RNAi} (**C**, n=14) versus DC^{GluR1}>>$dome^{RNAi}$ (**D**,
n=14) ablated discs, with the number of NiA quantified in (**E**), P**** <0.0001. Data were analyzed with a one-
way ANOVA followed by a multiple comparisons test. Error bars represent the standard deviation. (**F**) A
DC^{GluR1}*hh-GAL4* ablation schematic ($DC^{GluR1}hh^{ts}$). A short heat shock (45 min at 37°C) induces split
LexA/lexAop-GluR1 ablation, while a heat shift (20 hr at 30°C) following the heat shock will inactivate
GAL80^{ts} and permit *UAS-X* expression. (**G**) A $DC^{GluR1}hh^{ts}$>*GFP* ablated disc, where GFP highlights the area
of the disc being targeted for JAK/STAT and Wg knockdown in the following experiments. The posterior NiA
will be assayed in response to knockdown while anterior NiA serve as an internal control. (**H**) A
$DC^{GluR1}hh^{ts}$>$Stat92E^{RNAi}$ ablated disc, arrowhead highlights an increase in pouch NiA, open arrowhead
indicates the lack of cDcp-1 signal in the hinge. (**I**) A control DC^{NA} disc showing developmental Wg
expression. (**J**) A DC^{GluR1}>>y^{RNAi} ablated disc showing that NiA cells avoid the areas of the pouch with high
Wg expression, while wound edge apoptotic cells overlap the developmental and damage-specific Wg-
expressing cells (arrowhead). (**K**) A $DC^{GluR1}hh^{ts}$>wg^{RNAi} ablated disc with NiA cells observed at the posterior
margin (arrowhead) while anterior NiA avoid the Wg margin stripe (open arrowhead). (**L**) A quantification of
posterior NiA cells (cDcp-1+ve cells within the pouch) following a control knockdown (y^{RNAi}, n = 10),
$Stat92E^{RNAi}$ (n = 10), wg^{RNAi} (n = 10), and $Zfh-2^{RNAi}$ (n = 11). P*** = 0.0002, P**** < 0.0001. ns = not
significant. Data were analyzed with a one-way ANOVA followed by a multiple comparisons test. Error bars
represent the standard deviation. See Supplementary Genotypes file for exact genotypes.

1093

1094

Figure 2 – figure supplement 1. (A) DC^{NA}>>$dome^{RNAi}$ control disc with no apoptosis in the absence of
ablation. (**B**) A DC^{NA}>>$hop48A$ control disc with high levels of cDcp-1 observed throughout the pouch. (**C**) A
DC^{GluR1}>>$hop48A$ ablated disc with high levels of apoptosis throughout the pouch, which may be induced by
UAS-$hop48A$ expression as seen in (**B**). (**D**) A $DC^{GluR1}hh^{ts}$>GFP ablated disc with nubbin and DCAD
highlighting the ablated $salm$ domain and GFP labeling the area of UAS-X expression. (**E**) A control hh-
$GAL4$ disc showing the normal expression pattern of the 10xSTAT-GFP reporter. (**F**) A hh-$GAL4$>UAS-
$Stat92E^{RNAi}$ disc expressing 10xSTAT-GFP, demonstrating a reduction of JAK/STAT activity as revealed by
the 10xSTAT-GFP reporter (arrowhead in H versus open arrowhead in I). (**G**) A control
hh^{ts}>$GFP,Stat92E^{RNAi}$ disc showing no changes in cell death. (**H**) A control hh^{ts}>GFP,wg^{RNAi} disc showing
the absence of Wg in the posterior pouch and no change in posterior cell death. (I) A control disc (**I**) A
DC^{NA}>>$Zfh2^{RNAi}$ control disc showing no increased apoptosis in the absence of ablation. (**J**) A
$DC^{GluR1}hh^{ts}$>$Zfh2^{RNAi}$ ablated disc showing a slight increase in pouch NiA. (**K**) A control disc bearing the
$vgQE$-$lacZ$ reporter showing the area of the pouch with vestigial quadrant enhancer expression. (**L**) A
DC^{GluR1} ablated disc bearing $vgQE$-$lacZ$, showing the high degree of overlap between NiA cells and lacZ-
expressing cells. (**M**) A control hh^{ts}>Wg disc with no strong changes in cell death observed. (**N**) A
$DC^{GluR1}hh^{ts}$>Wg ablated disc showing an increase in apoptosis across the wing pouch and in the posterior
pleura. See Supplementary Genotypes file for exact genotypes.

1112

1113

1114

Figure 3. NiA promotes proliferation late in regeneration. (**A-A'''**) A time course of DC^{GluR1} ablated discs
bearing the *PCNA-GFP* reporter at 18 hr (**A**), 24 hr (**A'**), 36 hr (**A''**), and 48 hr (**A'''**). (**B-B'''**) A time course
of EdU in DC^{GluR1}>>y^{RNAi} ablated discs at 18 hr (**B**), 24 hr (**B'**), 36 hr (**B''**), and 48 hr (**B'''**). Over time, high
EdU signal localizes to the damaged pouch by 36 hr and remains elevated at 48 hr. (**C-C'''**) A time course of
DC^{hepCA}>>y^{RNAi} ablated discs at 18 hr (**C**), 24 hr (**C'**), 36 hr (**C''**), and 48 hr (**C'''**). (**D-D'''**), demonstrating that
EdU initially becomes elevated in the outer regions of the pouch by 24, but then localizes to the central
pouch region by 36 hr. A time course of DC^{GluR1}>>*mir(RHG)* ablated discs at 18 hr (**D**), 24 hr (**D'**), 36 hr
(**D''**), and 48 hr (**D'''**), demonstrating the loss of EdU localization. (**E**) A graph of the DC^{GluR1}, DC^{hepCA}, and
DC^{NA} EdU time courses highlight the pattern of EdU labeling in the wing pouch between each system. Error
bars represent the standard deviation (**F**) A quantification of EdU signal intensity between DC^{GluR1}>>y^{RNAi} at
18 hr (n= 10), 24 hr (n= 10), 36 hr (n= 10), 48 hr (n= 10), and DC^{GluR1}>>*mir(RHG)* at 18 hr (n= 10), 24 hr (n=
8), 36 hr (n= 11), and 48 hr (n= 10) time courses, ns, not significant; ***P = 0.0002; ****P <0.0001; data were
analyzed with a one-way ANOVA and multiple comparisons test. Error bars represent the standard
deviation. (**G-J**) DC^{GluR1} ablated discs with different populations of apoptotic cells suppressed by *mir(RHG)*.
(**G**) A control DC^{GluR1}>>y^{RNAi} ablated disc showing the typical pattern of NiA formation. (**H**) A
DC^{GluR1}>>*mir(RHG)* ablated disc suppressing both wound edge apoptosis and NiA cells. (**I**) A
DC^{GluR1}>>*miRHG;DR^{WNT}-GAL80* ablated disc, which targets the NiA area of the pouch for *UAS-mir(RHG)*
expression (hereafter Lateral Pouch, or *LP>mir(RHG)*). (**J**) A DC^{GluR1} x *R85E08>mir(RHG)* ablated disc,
which targets wound edge apoptotic cells for suppression (hereafter Wound Edge, or *WE>mir(RHG)*). The
dotted lines each panel highlight the area of *UAS-mir(RHG)* expression. (**K-N**) DC^{GluR1} ablated discs with
representative EdU labels at R36 in response to y^{RNAi} (**K**), whole-pouch *mir(RHG)* (**L**), *LP>mir(RHG)* (**M**),
and *WE>mir(RHG)* (**N**). (**O**) A quantification of the normalized EdU fluorescent intensity of DC^{GluR1}>>y^{RNAi}
R18 (n= 10), DC^{GluR1}>>y^{RNAi} R36 (n= 10), DC^{GluR1}>>*mir(RHG)* R36 (n= 11), *LP>mir(RHG)* (n= 10), and
WE>mir(RHG) (n= 9) ablated discs; *P = 0.0189, **P = 0.0044, ***P = 0.0008, ****P < 0.0001; data were
analyzed with a one-way ANOVA and multiple comparisons tests. Error bars represent the standard
deviation. LP = lateral pouch, WE = wound edge. See Supplementary Genotypes file for exact genotypes.

1141

1142

1143

Figure 3 – figure supplement 1. (**A**) A DC^{NA} control disc expressing *PCNA-GFP*. (**B**) A DC^{GluR1} ablated
disc expressing *PCNA-GFP*, with reporter expression localizing to the damaged pouch over time. (**C**) A
quantification of DC^{NA} (n = 10) and *DCGluR1* ablated discs at 18 hr (n = 10), 24 hr (n = 10), and 36 (n = 11),
with the pouch-specific PCNA-GFP signal normalized to entire discs in each time point. This time course
demonstrates that PCNA-GFP expression localizes to the distal pouch by 36 hr of regeneration, Data were
analyzed with a one-way ANOVA followed by a multiple comparisons test. P** = 0.0095. ns = not significant.
Error bars represent the standard deviation. (**D-D'''**) A DC^{NA}>>y^{RNAi} control time course at 18 hr (**F**), 24 hr
(**F'**), 36 hr (**F''**), and 48 hr (**F'''**). The folds of the wing pouch, visualized by DAPI, were used to outline the
EdU signal for quantification in all experiments. (**E-F**) A DC^{NA}>>y^{RNAi} control disc (**E**) with EdU labeling
relative to a DC^{NA}>>*mir(RHG)* control disc (**F**). (**G**) A quantification of discs in (**E**, n=10) and (**F**, n=10)
demonstrating that miRHG expression does not alter EdU levels, ns, not significant. Data were analyzed
with a Student's T-test. See Supplementary Genotypes file for exact genotypes.

1156

Figure 4 NiA do not secrete mitogens associated with Apoptosis-induced Proliferation (AiP). (**A-A'**) A

DC^{NA} control disc bearing the *wg-lacZ* reporter. (**B-B'**) A DC^{GluR1} ablated disc with *wg-lacZ* showing the

expression of the reporter during regeneration, note several *lacZ*-expressing cells at the wound edge. The

24 hr time point was chosen as an intermediate between NiA formation (18 hr) and blastema formation (36

hr), as any secreted factors involved in AiP are likely to be detected prior to 36 hr. (**C-C'**) A DC^{GluR1}>>*P35*

ablated disc bearing *wg-lacZ* with *lacZ* expressing cells observed at the wound edge (arrowhead) but not in

the lateral pouch where NiA occur (open arrowhead). (**D-D'**) A DC^{NA} control disc bearing the *dpp-lacZ*

reporter. (**E-E'**) A DC^{GluR1} ablated disc with *dpp-lacZ*; *lacZ*-expressing cells are again observed at the wound

edge. (**F-F'**) A DC^{GluR1}>>*P35* ablated disc bearing *dpp-lacZ*, with *lacZ*-expressing cells at the wound edge

(arrowhead) but not where NiA cells occur (open arrowhead), demonstrating that NiA do not secrete factors

(Dpp and Wg) involved in AiP. (**G**) A DC^{GluR1} ablated disc bearing the *AP-1-GFP* reporter and labeled with

dihydroethidium (DHE) to visualize reactive oxygen species (ROS). Low levels of DHE labeling are observed

at the wound edge overlapping *AP-1-GFP* expression (arrowhead), while no DHE labeling is observed in the

lateral pouch where NiA occur. (**H**) A DC^{GluR1}>>*Cat,Sod1* ablated disc showing reduced wound edge

apoptosis (open arrowhead) but no change in NiA formation. (**I**) A DC^{GluR1}>>*Duox^{RNAi}* ablated disc showing a

similar pattern to (**H**) with an observed loss of wound edge apoptosis (open arrowhead) but no change in

NiA formation. (**J-L**) DC^{GluR1} ablated discs bearing *yRNAi* (**J**), *P35* (**K**), and *LP*>*P35* (**L**) at 36 hr and labeled

with EdU. (**M**) A quantification of discs in (**J-L**) demonstrate no change in EdU labeling upon the expression

of P35 in the whole pouch (**K**) or in NiA cells (*LP*>*P35*, **L**), confirming that NiA promote proliferation

independent of AiP, (*y^{RNAi}*, n=8, *P35*, n=11, *LP*>*P35*, n=10), ns, not significant. Data were analyzed with a

one-way ANOVA followed by a multiple comparisons test. LP = lateral pouch, AiP = apoptosis-induced

proliferation. See Supplementary Genotypes file for exact genotypes.

1182

1183

1184 **Figure 4 – figure supplement 1.** (**A**) A DC^{NA} control disc bearing the *spi-lacZ* reporter, showing no lacZ in

1185 the wing disc. (**B**) A DC^{GluR1}>>*P35* ablated disc bearing *spi-lacZ* showing few *lacZ*-expressing cells at the

1186 wound edge. (**C**) A DC^{NA} control disc bearing *mol-lacZ* highlighting the developmental expression pattern of

1187 *mol* in the wing disc. (**D**) A DC^{GluR1} ablated disc bearing *mol-lacZ*. A slight increase in *lacZ* expression is

1188 observed at the wound edge (arrowhead). See Supplementary Genotypes file for exact genotypes.

1189

1190

1191

Figure 5. NiCP cells persist in the disc proper throughout regeneration. (**A-D**) A time course of DC^{GluR1}

ablated discs bearing the fluorescent caspase reporter *UAS-GC3Ai* at 18 hr (**A**), 24 hr (**B**), 36 hr (**C**), and 48

hr (**D**). (**E**) A DC^{GluR1}>>*GC3Ai* ablated disc at 64 hr, with GC3Ai-positive cells present at the distal pouch

after the nubbin marker is reestablished (arrowhead). (**F**) An area quantification of the GC3Ai signal in the

whole pouch at 18 hr (n = 10), 24 hr (n = 3), 36 hr (n = 7), and 48 hr (n = 7), demonstrating that GC3Ai

signal area increases over time. Data were analyzed with a one-way ANOVA followed by a multiple

comparisons test. P* = 0.0308, P*** = 0,0001, P**** < 0.0001. ns = not significant. Error bars represent the

standard deviation. (**G-G'**) A DC^{GluR1}>>*GC3Ai* ablated disc at 36 hr with GC3Ai-positive cells associated with

the mitotic PH3 marker. Yellow arrowhead in (**G'**) points to non-apoptotic GC3Ai-positive cells, while the

white arrowhead points to GC3Ai labeled cells with an apoptotic morphology. (**H-I**) A DC^{GluR1} *LP>GC3Ai*

ablated disc at 18 hr (**H**) and 36 hr (**I**). The open arrowhead in (**H**) points to the lack of GC3Ai labeling at the

WE. The arrowhead in (**I**) highlights the presence of GC3Ai-labeled NiA cells at the wound edge. (**J-J'**) A

DC^{hepCA}>>*GC3Ai* ablated disc at 18 hr with apoptotic cells present throughout the disc proper. (**K-K'**) A

DC^{hepCA}>>*GC3Ai* ablated disc at 36 hr (**K**) and 64 hr (**K'**); all apoptotic cells appear to be extruded towards

the basal surface of the epithelium. (**L-L'**) A DC^{GluR1}>>*GC3Ai* ablated discs at 18 hr with NiA cells present in

the disc proper. (**M-M'**) A DC^{GluR1} *LP>GC3Ai* ablated disc at 36 hr, with columnar-shaped GC3Ai-positive

cells in the disc proper (yellow arrowhead) alongside pyknotic, rounded GC3Ai-positive cells (red

arrowhead), showing that there are a mix of morphologically apoptotic NiA and non-apoptotic NiCP cells in

the disc proper. In each of the transverse sections (**J'**, **K'**, **L'**, **M'**), Ba indicates the basal surface of the disc,

while Ap indicates the apical surface of the disc. GC3Ai = green caspase-3 activity indicator, LP = lateral

pouch. See Supplementary Genotypes file for exact genotypes.

1213

1214

1215

Figure 5 – figure supplement 1. (**A**) A DC^{NA}>*GC3Ai* control disc driving *UAS-GC3Ai* in the pouch, HA
labels all *GC3Ai*-expressing cells, while the cells with activated GC3Ai also have green fluorescence. (**B**) A
DC^{GluR1}>>*GC3Ai* ablated disc showing GC3Ai-positive NiA cells and, to a lesser extent, GC3Ai-positive
wound edge apoptotic cells. (**C**) A DC^{GluR1} *LP*>*GC3Ai* ablated disc, with HA highlighting the restricted
expression of *GC3Ai* to the NiA area of the disc, thus excluding wound edge cells. (**D**) An area quantification
of the GC3Ai signal in the distal pouch at 18 hr (n = 10), 24 hr (n = 3), 36 hr (n = 7), and 48 hr (n = 7),
demonstrating that GC3Ai signal area increases over time. Data were analyzed with a one-way ANOVA
followed by a multiple comparisons test. P**** < 0.0001. ns = not significant. Error bars represent the
standard deviation. (**E-F**) DC^{GluR1} ablated discs at 18 hr (**E**) and 36 hr (**F**) with *GC3Ai/TUNEL* double-
positive NiA at both time points. (**G-H**) DC^{GluR1} ablated discs at 18 hr (**G**) and 36 hr (**H**) showing strong
overlap between NiA and yH2Av at 18 hr (arrowheads in **G**) that diminishes by 36 hr (open arrowheads in
(**H**). GC3Ai = green caspase-3 apoptosis indicator, LP = lateral pouch. See Supplementary Genotypes file
for exact genotypes.

1229

1230

1231

Figure 6 – NiCP have sublethal levels of activated executioner caspase. (**A-A'**) A DC^{GluR1} DBS-GFP

ablated disc at 18 hr showing strong overlap between cDcp-1 and DBS-GFP-positive NiA/NiCP. (**B-B'''**)

DC^{GluR1} $CasEx^{ts}$>G-$TRACE$ ablated discs at 18 hr with RFP labeled wound edge cells, (arrowhead), but no

RFP or lineage trace-positive labeling is observed in NiA/NiCP (open arrowheads). (**C**) A DC^{hepCA}

$CasEx^{ts}$>G-$TRACE$ ablated disc at 18 hr showing a high level of overlap between apoptotic cells and both

RFP and the lineage trace (arrowhead). (**D-E**) DC^{GluR1} $CasEx$>GFP ablated discs at 18 hr (**D**) and 36 hr (**E**)

showing that NiA/NiCP are labeled by $CasExpress$ in a sensitized background (when $tubGAL80^{ts}$ is omitted).

DBS = Drice-based sensor, CasEx = CasExpress. See Supplementary Genotypes file for exact genotypes.

1240

1241

1242

1243 **Figure 6 – figure supplement 1.** (**A**) A DC^{GluR1} $CasEx^{ts}$>G-TRACE ablated disc at 36 hr with no lineage-

1244 positive NiA/NiCP, despite the presence of clonal patches of cells traced by developmental caspase activity.

1245 See Supplementary Genotypes file for exact genotypes.

1246

Figure 7 – Dronc activity in NiCP promotes regeneration. (**A-J**) Representative DC^{GluR1} ablated discs at

18 hr (**A**,**C**,**E**,**G** and **I**) with cDcp-1 and Mmp1 labeling, and at 36 hr (**B**,**D**,**F**,**H** and **J**) with EdU labeling,

bearing UAS-y^{RNAi} (**A-B**), UAS-$mir(RHG)$ (**C-D**), $Dronc^{I29}/+$ (**E-F**), UAS-$Dronc^{DN}$ (**G-H**), and UAS-$DIAP1$ (**I-J**).

Open arrowhead in (**F**) indicates the strong reduction in EdU labeling. Open arrowheads in (**I**) show a strong

suppression of NiA/NiCP as detected by cDcp-1. (**K**) A quantification of NiA/NiCP number (cDcp-1+ve,

Mmp1-ve cells) in UAS-y^{RNAi} (n = 11), UAS-$mir(RHG)$ (n = 8), $Dronc^{I29}/+$ (n = 10), UAS- UAS-$Dronc^{DN}$ (n =

7), and UAS-$DIAP1$ (n = 8), ns, not significant, P**** <0.0001. Data were analyzed with a one-way ANOVA

followed by a multiple comparisons test. Error bars represent the standard deviation. (**L**) A quantification of

EdU fluorescence intensity with UAS-y^{RNAi} (n = 20), UAS-$mir(RHG)$ (n = 14), $Dronc^{I29}/+$ (n = 18) UAS- UAS-

$Dronc^{DN}$ (n = 17) and $DIAP1$ (n = 10), ns, not significant, P*** = 0.0004, P**** <0.0001. Data were analyzed

with a one-way ANOVA followed by a multiple comparisons test. Error bars represent the standard

deviation. (**M**) Regeneration scoring of DC^{GluR1} ablated flies in a +/+ versus a $Dronc^{I29}/+$ background. Note

the increase in frequency of full notched and ablated wings in $Dronc^{I29}/+$ relative to the +/+ treatment. (**N-O**)

Schematics depicting the apoptotic pathway in Drosophila (**N**) and the three different caspase-positive cell

types that occur following necrotic ablation (**O**). Following necrosis, JNK signaling at the wound edge (WE)

promotes Dronc activation, which mediates WE apoptosis and proliferation through the AiP feedback loop

(left). In the lateral pouch (LP), DAMP-like signal(s) leads to the formation of necrosis-induced caspase

positive (NiCP) cells that utilize Dronc to promote proliferation and subsequent regeneration independent of

both JNK and AiP signaling, while necrosis-induced apoptosis (NiA) undergo apoptotic death resulting from

high levels of dronc, cDcp-1 and Drice activity (right). Both WE apoptosis and NiCP act to promote

regeneration through separate mechanisms relying on Dronc-mediated regenerative proliferation. See

Supplementary Genotypes file for exact genotypes.

1272

1273

Figure 7 – figure supplement 1. (**A**) A quantification of regenerated wing size in an unablated control
(DC^{NA}) wildtype (+/+, males = 17, females = 11) and *Dronc^{I29}/+* (males = 11, females = 22) background, and
following *DC^{GluR1}* ablation in a +/+ (n = 29 males, n = 19 females) and *Dronc^{I29}/+* (n = 24 males, n = 28
females) background, ns = not significant, P*** = 0.0001, data were analyzed with a one-way ANOVA
followed by a multiple comparisons test. Error bars represent the standard deviation. Representative images
for regenerated +/+ (top) and *Dronc^{I29}/+* (bottom) wings are shown to the right. See Supplementary
Genotypes file for exact genotypes.

1281

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 1 - Figure supplement 1

Figure 2 - Figure supplement 1

Figure 3 - Figure supplement 1

Figure 4 - Figure supplement 1

Figure 5 - Figure supplement 1

Figure 6 - Figure supplement 1

Figure 7- Figure supplement 1

A