AutoMorFi: Automated Whole-image Morphometry in Fiji/ImageJ for Diverse Analyses and Discoveries

Ouzéna Bouadi[1], Chenkai Yao[2]*, Jason Zeng[1]*, Danielle Beason[2]*, Nyomi Inda[3]*, Zoe Malone[1]*, Jonathan Yoshihara[1], Amritha Vinayak Manjally[1], Clifton Johnson III[1], Jonathan Cherry[4], Chin-Yi Chen[5], Tzu-Chieh Huang[5], Bogdana Popovic[5], Maria Henley[5], Guangmei Liu[1], Hannah Aichelman[1], Sarah W. Davies[1], Yuan Tian[1], Hengye Man[1], Thomas Gilmore[1], Elif Ozsen[1], Kristen Harder[1], Peter Walentek[6,7], Elizabeth K. Kharitonova[8], Ella Zeldich[8], David Pitt[5], Tuan Leng Tay[1,8†]

Affiliations:

[1]Department of Biology, Boston University, Boston, MA 02215, USA

[2]Department of Biomedical Engineering, Boston University, Boston, MA 02215, USA

[3]Department of Electrical and Computer Engineering, Boston University, Boston, MA 02215, USA

[4]Department of Pathology and Laboratory Medicine, Boston University School of Medicine, Boston, MA 02118, USA

[5]Department of Neurology, Yale School of Medicine, Yale University, New Haven, CT 06510, USA

[6]Department of Internal Medicine IV, University Medical Center Freiburg, Freiburg 79106, Germany

[7]CIBSS - Center for Integrative Biological Signaling Studies, University of Freiburg, Freiburg, 79104, Germany

[8]Department of Anatomy & Neurobiology, Boston University School of Medicine, Boston, MA 02118, USA

*These authors contributed equally to this work.

†Correspondence:

Tuan Leng Tay, PhD <tltay@bu.edu>

ABSTRACT

Running on the highly popular and accessible ImageJ/Fiji platform for biological image analysis, we have established AutoMorFi as a streamlined interface for automated whole-image morphometric analysis that generates at least 47 measurements per cell or object in under 1 minute. We performed multiple validated cluster and principal component analyses on nonredundant morphometric parameters derived from AutoMorFi for various cell types, objects, and organisms. We used images of rodent macrophages, human postmortem brain tissues from multiple sclerosis (MS) and Alzheimer's disease (AD) patients, iPSC/animal models for Down's syndrome and autism spectrum disorder (ASD), and organisms such as sea anemone and corals. AutoMorFi's adaptability extends across diverse imaging modalities including brightfield, confocal, or widefield fluorescence microscopy as well as underwater photography. Due to its unlimited and unbiased sampling across any image and high potential for modification and customization, using AutoMorFi has led to the discovery of new distinguishing features in previously studied cell types and organisms as well as the development of rapid diagnostic approaches. AutoMorFi represents a transformative tool that will accelerate morphometric analysis and offer broad relevance in biological studies.

MAIN TEXT

Morphometry is a bedrock for understanding life sciences. As a key indicator of functional, behavioral, and physiological state, the study of cellular and organismic morphology has been achieved through numerous methods. Common approaches range from meticulously hand-drawn neurons and camera lucida traces of glia[1–3] to computational and machine-learning-based image analysis tools that classify cells and animal species by their diverse characteristics[4–6]. Current semi-automatic processes for common applications such as cell detection, quantification, and morphometry are, however, mostly out of reach for many researchers who are unfamiliar with coding languages that may pose a steep learning curve[7]. The day-to-day reality of image data analysis involves manual adjustments of brightness, contrast, and thresholds for each image as well as painstaking tracing and validation of individually segmented cells or objects. Accurate execution of these routine tasks requires varying in-depth subject knowledge and depends on the complexity and proximity of the objects of interest. Ideally, morphometric analysis tools should be user-accessible, generate accurate results that can be easily quality controlled, offer options for unbiased and batch sampling, and highly versatile and customizable to address multifaceted scientific questions.

For nearly 3 decades, ImageJ or Fiji[8,9] has been the go-to open-source platform for researchers

globally to process and analyze images as well as to reconstruct cells and objects acquired by

microscopy to obtain quantitative morphometric data. Taking advantage of this familiar resource,

we aimed to standardize and automate the acquisition of multiple morphological parameters of

complex and branched cells such as the highly ramified brain-resident microglia during

homeostasis, by looping a series of Fiji commands using a macro[10]. We invited 30 users ranging

from high school interns to principal investigators to test our first prototype, "AutoGliji", on a

standard fluorescence image of IBA-1-immunolabeled mouse microglial cells (**Fig. 1A**). Testers

were asked to perform manual measurements of the same 3 cells using Analyze Particles and

the Simple Neurite Tracer ImageJ plugin[11] for comparison (**Fig. 1A-C**). The time needed to

analyze all cells in the image by AutoGliji (5.5 ± 0.75 minutes) was drastically reduced when

compared to manual measurement of 3 cells (105 ± 19 minutes) (**Fig. 1D**). Furthermore, manual

measurements of over half of the parameters were highly inconsistent across testers due to

different levels of familiarity with microglial morphology, in stark contrast to the reproducible

results from AutoGliji which were generated by the same users running either Windows or MacOS

(**Fig. 1E-N**).

AutoMorFi is an optimized version of AutoGliji that was configured to generate 33 measurements

and 14 calculated parameters per cell or object with macro runtime reduced to under 1 minute

per image (**Table 1, Supplementary Notes 1–6**, **Supplementary Video 1**). Sampling limit within

any 2-dimensional 8-bit TIFF image that could be opened in Fiji was eliminated. To validate our

hypothesis that 47 measures provide sufficient dimensionality to distinguish and classify cells

based on morphology using standard clustering algorithms for high dimensional data, we

analyzed 393 microglia, 438 cardiac macrophages, 201 alveolar macrophages, 307 liver Kupffer

cells, 332 kidney macrophages, and 274 splenic macrophages from the same wildtype adult mice

that were immunolabeled with IBA-1[12–14] (**Fig. 2A-B**). As expected, 47 morphometric features

classified microglia and cardiac macrophages as most morphologically distinct and revealed that

macrophages of the kidney, liver, lung, and spleen share an intermediate phenotype (**Fig. 2C-F**).

Furthermore, UMAP visualization of the data has allowed us to determine the major morphological

characteristics that distinguish these macrophages of common lineage (**Fig. 2G-J)**.

To assess the potential of AutoMorFi for diagnostic and future large-scale applications, we tested

its ability to cluster microglia identified by a single IBA-1 marker in well-documented lesions of MS

from patients and controls (**Table 2**, **Fig. 3**). Microglia play different roles in all stages of
demyelination, oligodendrocyte death, and neuroaxonal injury in MS[15–18]. While microglia exhibit
degenerative and pro-inflammatory features as MS progresses, they also participate in
neuroprotection and repair. Thus, exploring microglial morphological diversity in different lesions
may unveil new strategies against microglia-driven neurodegeneration in MS. Cluster analysis
revealed a clear transition of microglial morphology from control white matter (WM) to the rim of
chronic active lesions (**Fig. 3C-F**) and 11 key morphological features that promoted optimal
classification of the pathological stages (**Fig. 3G-J**). AutoMorFi provides a cost-effective and time-
efficient alternative as a preliminary diagnostic tool with shorter turnaround time in contrast to
requiring multiple immunohistochemical assays on several disease markers. Furthermore,
evaluation of AutoMorFi against well-established open-source segmentation tools, including
CellProfiler, ilastik, Cellpose, and QuPath suggests that only AutoMorFi offers high confidence in
capturing intricate characteristics of complex and densely distributed cells with the operational
efficiency necessary for handling large datasets.

As further proof-of-principle that AutoMorFi can be easily applied for morphometry-based
discoveries, it was used to analyze diverse image types from widefield or confocal fluorescence
and brightfield microscopy to underwater photography (**Fig. 4**). These images represent diverse
applications such as the comparison of glial cells across model systems such as euploid human
induced pluripotent stem cells-derived microglia from Down's syndrome patients, aged human
microglia, *Drosophila* glia, and rodent microglia (**Fig. 4A**), characterization of *in vitro* neuronal
cultures (**Fig. 4B**), single cells (**Fig. 4B-C**), or whole organisms (**Fig. 4D**), genetic studies (**Fig.
4B-C**), and the impact of environmental stress on immune signaling and coral reef ecosystems
(**Fig. 4D-E**). AutoMorFi measurements unveiled additional morphological changes compared to
freehand tracing[19] of Ube3a 2x Tg neurons which model Ube3a overdosage and hyperactivity in
ASD[20–24] and confirmed the hypothesis of reduced dendritic arborization[25–27] (**Fig. 4B**). Motile cilia
are membrane-covered cell protrusions found in ciliated cells involved in facilitating movement or
transport of substances. Impaired ciliated cells can lead to chronic airway diseases, infertility, or
hydrocephalus. Ciliogenesis requires precise molecular regulation at the centriole to switch from
cell division to cilia formation employing molecules such as the centriolar protein Cp110[28]. Fast
readouts from multiplexing AutoMorFi and fluorescence intensity measurements from Centrin4
and acetylated α-tubulin markers of multi-ciliated cells (MCCs) validated previous findings of more
efficient rescue of *Cp110* knockdown by *Cp110-Δ3'UTR* DNA than *Cp110* DNA (**Fig. 4C**). This
approach revealed the heterogeneity of MCC phenotypes in the study and identified 30% of the

parameters that were significantly altered. Starvation of sea anemones was reported to reduce

NF-κB, leaving the animals vulnerable to infection[29]. NF-κB proteins are important for many

cellular processes including immunity, inflammation, development, cellular growth, and apoptosis.

They are differentially regulated in many diseases such as cancer, arthritis, chronic inflammation,

asthma, neurodegenerative disease, and heart disease. By reanalyzing the images of whole-

mount anemone polyps[29] using AutoMorFi as well as taking intensity and distribution (area

fraction) measurements of NF-κB, we newly discovered that starvation does not alter the gross

morphology of the anemones and validated the earlier findings of NF-κB reduction (**Fig. 4D**).

Coral reefs are critically threatened by climate change. Recent advancements in sequencing for

non-model systems have revealed previously unrecognized 'cryptic coral lineages' (*i.e.,*

genetically distinct but morphologically similar taxa) that have important implications for predicting

reef ecosystem futures and effectively restoring reefs[30]. Rapid and cost-effective methods to

identify these cryptic lineages, preferably alternative methods to DNA sequencing, are needed to

effectively characterize coral diversity present on reef ecosystems and ultimately improve lineage

assignment. AutoMorFi identified diagnostic skeletal phenotypes and replicated the same pattern

of cryptic lineage differentiation in much less time compared to previous analysis requiring ~25

minutes per image to measure all relevant traits (**Fig. 4E**).

In summary, we demonstrate the easy usage and versatility of AutoMorFi for rapid and

reproducible acquisition of large morphometric datasets across biological scales and research

objectives that have expanded the depth of information resulting in new discoveries. Using

AutoMorFi, statistical power is easily increased due to unlimited and unbiased sampling. To

promote broad usage and customization of the tool for novel applications, we provide a

troubleshooting guide (**Supplementary Note 3**) of common pitfalls identified from thorough

executions of AutoMorFi by multiple research teams. AutoMorFi could, for example, be adapted

to quantify and classify condensates of cellular proteins undergoing liquid-liquid phase separation

to introduce more quantitative descriptors of proteostasis in disease and aging to the field[31]. When

applied to multi-channel images (**Fig. 4C-D**) and combined with additional approaches such as

genetics and genomics as well as other classification algorithms (e.g., random forest,

bootstrapping[5]), morphometrics will serve as potent indicators and predictors of disease states

and environmental changes.

ACKNOWLEDGEMENTS

The authors thank AutoGliji testers E. Batchelor, S. Brayton, V. Chiplunkar, E. Choi, S. Guerra,
K. He, A. Hoo, M. Iordanov, S. Jo, T. Keefauver, J. Koo, E. Kraft, H. LeBlanc, A. Liu, E. Maloney,
C. Reddy, P. Ruiz, K. Sampat, and M. Zreik; I. Amin for assistance with microscopy; G. Lin (R
scripts) and A. Labadorf for data analysis and visualization advice. *In memoriam* M. Spelleken
and Scarlet. TLT was supported by a NARSAD Young Investigator Grant from the Brain &
Behavior Research Foundation, Spivack Neuroscience Pilot Award, and Patricia McLellan Leavitt
Research Fund Award.

AUTHOR CONTRIBUTIONS

ZM, DB, CY, and OB established the macro. CY wrote the coordinate-matching code. OB, JZ,
CY, NI, JY, AVM, CJ III, JC, C-YC, T-CH, BP, MH, GL, EKK, EZ, HA, SD, PW, YT, HM, TG, CHC,
B-BK, DP, and TLT generated data, provided/validated images, and performed analyses. EO and
KH ran pilot analysis. OB and TLT wrote the manuscript. TLT conceived and supervised the study.

COMPETING INTERESTS

The authors declare no competing interests.

MATERIAL AND METHODS

Sample preparation.

Mouse. Female C57Bl/6J (wild-type) mice were purchased from Charles River Laboratories and
raised in a specific-pathogen-free facility under 10h/14h dark/light cycle until 52 weeks old. Mice
were given food and water *ad libitum*. Animal experiments are approved by the Institutional Animal
Care and Use Committee (IPROTO202200000054). Mice were deeply anesthetized by
intraperitoneal administration of 100 mg ketamine and 5 mg xylazine per kg body weight and
transcardially perfused with phosphate-buffered saline (PBS). Brains, hearts, kidneys, livers,
lungs, and spleens were post-fixed overnight at 4°C in 4% paraformaldehyde (PFA) in PBS, rinsed
in PBS, and cryoprotected in 30% sucrose/PBS at 4°C until fully submerged. Samples were
embedded in Tissue-Tek O.C.T. compound (Sakura Finetek, USA) for frozen sectioning on a
cryostat (Cryostar NX50; Epredia Cryostat, USA).

Human. Human postmortem brain tissues from 2 MS and 2 control patients were obtained
according to an institutional review board-approved protocol (Yale HIC 1211011031) (**Table 2**).

Brain hemispheres were fixed with formalin for a minimum of 2 weeks. Brain slabs with lesions
were embedded with paraffin. A chronic active MS lesion was characterized using brightfield
staining. Images of human postmortem tissue from the dorsolateral frontal cortex were obtained
from the Boston University Alzheimer's Disease Research Center (BU ADRC): 5 cases from 3
females and 2 males aged 75–105 with either a non-AD or low NIA-Reagan score, and with no
other comorbid pathology were taken for the control group.
Human iPSC-derived microglia. A trisomic line (WC-24-02-DS-M) and an isogenic euploid
control cell line (WC-24-02-DS-B) between passages 18 and 27 were generated from a female
25-year-old with Down syndrome and used in this study. The cell lines were produced, validated,
and kindly provided by Dr. Anita Bhattacharyya from the University of Wisconsin-Madison. IPSCs
were routinely passaged and cultured on Matrigel® (cat. 354277 Corning®, Corning, NY, USA)
using mTeSR™ plus (cat. 85850, StemCell Technologies, Vancouver, BC, Canada). To generate
iPSC derived microglia, iPSCs were differentiated into CD-43 expressing hematopoietic
progenitor cells (HPCs) using the STEMdiff™ Hematopoietic Kit (cat. 05310, STEMCELL
Technologies) for 12 days. The expression of CD43, CD45, and CD34 was validated by flow
cytometry. Cells expressing >90% $CD43^+$ and >20% of $CD34^+$/$CD45^+$ on day 12 were confirmed
to be HPCs, cryoprotected, and used for the subsequent differentiation into microglia. The
microglial differentiation was performed on thawed HPCs (200,000 cells per well of a 6-well plate)
using the STEMdiff™ Microglia Culture System (cat. 100-0019 and 100-0020, STEMCELL
Technologies) for 24 days according to manufacturer's directions. On day 24, expression of >80%
$CD45^+$ and $CD11b^+$ microglial markers was confirmed by flow cytometry to ensure proper
differentiation. Cells were plated on poly-d-lysine coated glass coverslips (cat. A3890401,
ThermoFisher, Waltham, MA, USA) at 100,000 cells/well and matured until day 30. Cells were
fixed with 4% PFA for 30 min, washed twice with PBS, and stored at 4°C until use.
Primary neuronal culture. Primary neuronal cultures were prepared from postnatal day 0 (P0)
mouse pups. At P0, brain tissues from WT and Ube3a 2x Tg male pups were dissected and
digested, and then plated onto circular coverslips for neuron culture.
Drosophila. Fly strains *alrm-GAL4* (#67032) and *UAS-IVS-GFP* (#32201) were obtained from the
Bloomington *Drosophila* Stock Center, crossed at 25°C, and raised at 29°C on standard cornmeal
and molasses food. Flies aged to 2 days were dissected for immunohistochemistry. Only male
flies were included in the experiment.
Sea anemone. *Nematostella vectensis* (*Nv*) were obtained from Maryland by Mark Martindale
and Matt Gibson. The anemones were kept at ⅓ strength artificial seawater (ASW, ~12 parts per
1000) in a dark incubator at 19°C with weekly water changes. Clonal pairs of *Nv* were generated

by bisection and regeneration and then fed or starved for 30 days. Adult animals were bisected
perpendicularly to the oral-aboral axis. Halves were placed in wells of a 24-well plate. Anemones
regenerated for 30 days with feeding paused until tentacles formed from the aboral end. Adult *Nv*
were fed freshly hatched brine shrimp (*Artemia*; brineshrimpdirect.com) and young polyps were
fed ground *Artemia* in ⅓ ASW three times a week. Animals that were regenerated from aboral
and oral ends were included. 40-day-old polyps that were fed or starved for 30 days were used.
Xenopus. Mucociliary epithelial MCCs of *Xenopus* embryos were used to study cilia formation.
MCCs form >100 basal bodies and motile cilia that are important for generating directional fluid
flow to clear pathogens from the airways or the tadpole skin. *Centrin4-cfp* was injected into
embryos to identify targeted cells and basal bodies of MCCs. Knockdown of *cp110* in embryos
was achieved with morpholino oligonucleotide (MO) injection to study the role of Cp110 in MCC
ciliogenesis. To rescue cilia formation, *cp110* DNA was co-injected with *cp110* MO.
Corals. *Siderastrea siderea* (massive starlet coral) is one of the primary reef-building corals in
Bocas del Toro, Panamá[32,33]. Previous work identified three cryptic lineages of *S. siderea* that
occupy sites spanning the inshore reefs of Bocas del Toro[34].
242
Immunohistochemistry.
Mouse. Sagittal sections of brains, hearts, and kidneys and sections of livers, lungs and spleens
were captured at 16-μm thickness on Superfrost™ slides and stored at -20°C until use. Tissue
sections were rehydrated using PBS and permeabilized in blocking solution containing 0.5%
Triton-X-100 (Sigma Aldrich, USA) and 5% bovine serum albumin (BSA) (Sigma Aldrich, USA) in
PBS for 1 h at room temperature (RT). Tissues were incubated overnight at 4°C with rabbit anti-
IBA1 antibody (1:500; Wako 019-19741) diluted in PBS containing 0.1% Triton-X-100 (Sigma
Aldrich, USA) and 5% BSA (Sigma Aldrich, USA). After 5 10-min rinses in PBST, tissues were
incubated with Alexa Fluor 647-conjugated donkey anti-rabbit secondary antibody (1:1000;
Thermofisher Scientific A-31573) at RT for 2 h. After 5 10-min rinses in PBST, tissues were
coverslipped with Prolong Gold Antifade Mountant containing 4′,6-diamidino-2-phenylindole
(DAPI) (Cell Signaling, 8961).
Human. PLP was used to identify the areas of MS lesion and CD68 was used to identify the active
rim of microglial cells (**Table 2**). For fluorescent staining, the formalin-fixed paraffin embedded
(FFPE) block was sectioned at 7-μm thickness onto a slide, deparaffinized in xylene and
rehydrated via an ethanol series. The section was subjected to antigen retrieval using citrate
buffer at pH 6.0 followed by permeabilization using 0.5% Triton X-100 in Tris-buffered saline
(TBS). After washing, the section was quenched in 0.3% H_2O_2 in TBS and washed again. Tissue

was blocked for 1 h using 10% normal goat serum (NGS) and 10% Fc receptor inhibitor

(Invitrogen, 14-9161-73) in TBS. Primary antibody for IBA-1 (Synaptic Systems, #234004) was

applied with 10% NGS and incubated at 4°C overnight. After several washes in TBST, Alexa-

Fluor 488-conjugated secondary antibody (Invitrogen, #A1073) was applied at RT for 3 h. After

several washes, Sudan Black (MP Biomedicals, 152088) was applied to reduce autofluorescence,

followed by washes and Hoechst Counterstain (Life Technologies, H357). Slices were

coverslipped with 50% glycerol in TBS. Paraffin-embedded brain tissues from the BU ADRC were

cut at 20-µm thickness. DAB (3, 3'-diaminobenzidine) immunohistochemical staining for IBA-1

(1:500, Wako, 019-19741) and nuclear counterstaining with hematoxylin was performed.

Human iPSC-derived microglia. Glass coverslips were treated with blocking solution containing

5% normal donkey serum in PBST (0.1% Triton X-100) at RT for 1 h. Cells on coverslips were

incubated overnight at 4°C in rabbit anti-IBA1 (1:500; Wako, 019-19741) diluted in blocking

solution. Coverslips were washed 3X for 10 min in PBST and incubated at RT for 1 h with Alexa-

Fluor 546-conjugated donkey anti-rabbit antibody (1:1000; Invitrogen, A10040) in blocking

solution. Coverslips were washed 3X 10 min in PBST. ProLong Gold Antifade Mountant with DAPI

(Invitrogen, P36931) was applied to adhere coverslips to slides. Slides were stored at 4°C and

light-protected until imaging.

Primary neuronal culture. Neurons plated on coverslips were fixed at day *in vitro* 8 (DIV8) in 4%

PFA for 8 min and washed twice in 1X PBS. Cells were permeabilized for 10 min in 0.3% Triton

X-100 (Sigma) in 1X PBS, followed by 2 washes in 1X PBS. Cells were blocked for 1 h in 10%

NGS in 1X PBS. To label dendritic structures, cells were incubated overnight at 4°C in 1X PBS

containing MAP2 monoclonal antibody (1:300; ThermoFisher, #13-1500). The next day, cells

were washed 3X with cold 1X PBS, followed by incubation in 1X PBS containing Alexa Fluor 488

or 555-conjugated mouse IgG secondary antibody (1:500; Invitrogen, #A-21422. Cells were

counterstained with Hoechst (1:10,000 in 1X PBS; ThermoFisher, H3570) for 30 sec followed by

2 washes in 1X PBS. Coverslips were mounted onto slides with Prolong Gold (ThermoFisher,

P36930) and left to dry in the dark overnight before imaging.

Drosophila. Male flies were fixed in 4% PFA diluted in 1X PBS and 0.5% Triton X-100 (PBT) at

RT for 3 h. Flies were washed 4X in PBT for 15 min each with rotation and dissected in PBS.

Whole brains were blocked at RT for 1 h in PBT containing 0.5% BSA and 5% NGS (PBANG)

and then incubated in PBANG containing rabbit anti-GFP (1:100; Torry Pines Biolabs, TP401) at

4°C for 2 nights. Brains were washed 4X in PBT at RT and incubated in PBANG containing goat

anti-rabbit Cy3 (1:200; Jackson ImmunoResearch, 111-165-003) for 2 nights at 4°C. Brains were

washed 3X in PBT. Samples were placed in VectaShield with DAPI (Vector Labs) overnight at
4°C and subsequently mounted on slides.

Sea anemone. Whole-mount immunofluorescence staining of NF-κB was performed using a
custom antibody and Texas Red-labeled secondary antibody to visualize NF-κB levels in *Nv*. For
immunofluorescence, *Nv* polyps were fixed in 4% formaldehyde in ⅓ ASW overnight at 4°C and
then washed 3X with 0.1% Triton X-100 in PBS (PTx). Polyps were then microwaved in warm
urea (5% w/v) at the lowest setting for 5 min, cooled at RT for 20 min, then washed 3X with PTx.
Polyps were blocked overnight in PTx containing 5% NGS and 1% BSA at 4°C on a nutator. Anti-
Nv-NF-κB antibody diluted 1:100 in blocking buffer was used overnight at 4°C. Polyps were
washed 4X with PTx and incubated in Texas-red-conjugated anti-rabbit secondary antibody
(1:160; Invitrogen, #T-2767). DAPI was added to a final concentration of 5 mg/ml to visualize
nuclei.

Xenopus. Whole embryos were fixed at embryonic stages 30-32 overnight at 4°C, washed 3X 15
min with PBS, 2X 30 min in PBST (0.1% Triton X-100 in PBS), and blocked at RT in PBST-CAS
(CAS blocking; ThermoFisher #00-8120) for 1 h. To visualize cilia, embryos were incubated
overnight at 4°C in mouse monoclonal anti-acetylated-α-tubulin antibody (1:700; Sigma, #T6793)
and Alexa Fluor 555-labeled goat anti-mouse antibody (1:250; Molecular Probes, #A21422) in
CAS Blocking.

Image acquisition and processing.

Mouse. Z stacks of whole organ sections were acquired on a ZEISS AxioScan 7 slide scanner
using a 20X 1 NA objective lens at 1X zoom and 1.1 binning. Images were sampled with a voxel
size of 0.325-μm X 0.325-μm X 1-μm for the DAPI (excitation wavelength 353-nm; emission
wavelength 465-nm) and Alexa Fluor 647 (excitation wavelength 650-nm; emission wavelength
665-nm) channels. Image tiles were stitched, processed for maximum intensity projections
(MIPs), cropped to obtain >200 cells per image, and exported as TIFF images using the ZEISS
ZEN pro software (ZEN 3.2 blue edition). At least 4 images were analyzed per organ/mouse.

Human. Tissue sections from MS patients were scanned on a Leica Microsystems DM6B
microscope a 40X objective lens at 0.329-μm per pixel. Each tile was focused using the "Auto-
focus" function and subsequently captured in multiple z-stack images of 6-μm thickness in total.
Z-interval was set depending on tissue quality. Tiled images were merged with the "Mosaic
merge" function to include a 20% overlay area. The "Projection" function was used to obtain MIPs
that contained the best signal intensity for each pixel. Images were cleaned up using "Instant
Computational Cleaning" under the "Thunder" function before they were exported in TIFF for

image processing in ImageJ. Images containing MS lesions were characterized by PLP, CD68,

and IBA1 expression and precisely cropped for morphometric analysis. Tissue sections from BU

ADRC were imaged using an Aperio GT450 slide scanner at a 40X magnification. Images were

processed using the "extract region tool" in the Aperio ImageScope Pathology Slide Viewing

Software to generate 10–12 cropped images measuring 1000-µm X 1000-µm each from the white

and gray matter. To enhance the DAB signal and minimize the noise contributed by the nuclear

counterstain, each duplicated MIP was thresholded to exclude hematoxylin and subtracted from

the original MIP using "Image Calculator" to generate a new 8-bit TIFF image for AutoMorFi

analysis.

Human iPSC-derived microglia. Images were acquired on an Olympus Fluoview FV300 using

a 40X oil immersion objective lens and the FV31S-SW Viewer software. Images were captured in

DAPI (excitation wavelength 353-nm; emission wavelength 465-nm) and Alexa-Fluor 546

(excitation wavelength 556-nm; emission wavelength 573-nm) channels.

Primary neuronal culture. Neuronal images were captured using a Carl Zeiss Axiovert inverted

fluorescence microscope with a 40X 1.3 NA oil immersion objective lens and the AxioVision

Release 4.5 software.

Drosophila. Images of the central brain were captured on a Nikon C2+Si confocal microscope

using a 40X 1.25 NA water immersion objective lens and NIS-Elements software. Fifty planes 1-

µm apart were captured for the Z stacks of each brain with a resolution of 1024 X 1024 pixels

starting from the surface of the central brain at the first point where the antennal lobe becomes

visible. MIPs were generated in ImageJ at serial steps of 10 (e.g., 1-10, 11-20, …, and 41-50) to

distinguish individual glial cells in each brain.

Sea anemone. Images were acquired on a Nikon C2 Si confocal microscope and processed

using ImageJ.

Xenopus. Images were acquired using a ZEISS LSM700 confocal microscope. ImageJ was used

for Z-stack analysis and processing.

Corals. Photos of representative coral colonies of each of the three cryptic lineages were taken

from three sites across Bocas del Toro (Hospital Point, Cristobal Island, and STRI Point) using

an Olympus Tough TG-6 camera on the underwater macro setting. For each individual coral (N =

52), macro photos focused on individual corallites were taken in triplicate, with each of the three

photos taken on a different location on the coral surface. Photographs were converted to 8-bit

binary images and scaled using the 2-mm size standard with the Set Scale function in ImageJ/Fiji.

Images were individually adjusted by 0.35% contrast enhancement, smoothing, thresholding, and

despeckling.

AutoMorFi morphometric analysis of single cells. Images were blinded before the individuals who did not generate the primary data ran the AutoMorFi macro except for iPSC-derived microglial cells and *Drosophila* astrocyte-like glia which were known to the persons who performed image analysis. A different set of individuals performed quality control to validate the AutoMorFi output. They compared the outlines of the detected "soma" and "whole cell" with cells in the original images to assess for false positive or false negative detections and inaccurate object segmentation in consultation with authors of the original experiments. Incorrectly detected cells or objects were removed from the .csv output files. Output images with missing cells were reanalyzed using improved diameter measurements. Procedures for running the AutoMorFi macro in ImageJ/Fiji are detailed in **Supplementary Notes 1–4**. Each single-cell measurement of "soma", "whole cell", and "skeletonized" structure (generated by ImageJ/Fiji from the whole-cell outline) derived from AutoMorFi were matched by the coordinates of the objects using the Python packages numpy, pandas, and sklearn.neighbors, and collated in a single .csv data file (**Supplementary Note 6**). AutoMorFi generated 33 morphometric measurements that were in part used in arithmetic functions that produced additional parameters resulting in a total of 47 quantitative characteristics per cell (**Table 1**).

AutoMorFi morphometric analysis of single objects. Image blinding and data validation were carried out as described for AutoMorFi analysis of single cells. To customize AutoMorFi for measuring non-cellular objects, the use of the pipeline was modified as follows.

Sea anemone. The diameters of several whole polyps were measured from the DNA channel to obtain a range of representative values. While executing AutoMorFi, the Analyze Particle plugin was set to "Overlay" instead of "Mask" to save the polyp outline as an overlay that was now considered the "whole cell". The overlay was renamed and manually saved as a TIFF image in the output folder before continuing with the AutoMorFi macro. A region of interest (ROI) was marked in the background of the image and added to the ROI manager to generate a second pseudo-object to allow the acquisition of all standard parameters of the whole single polyp (see Error 7 in **Supplementary Note 3**). The macro was continued until the end. Measurements from the marked ROI and "soma" were excluded from the data analysis. "Whole cell" and "skeletonized" objects based on whole-polyp outlines contributing to 21 measured parameters were matched by their coordinates and collated in a single .csv data file. To acquire the total fluorescence intensity measurement of DNA and NF-κB signals, the overlay outline or "whole cell outline" that was saved earlier was added to the ROI manager in ImageJ/Fiji for each individual channel per organism.

The measured total fluorescence intensity values were normalized to the polyp area within the

whole cell outline. The NF-κB area fraction represents the percentage of non-zero pixels within

the polyp outline. An additional 12 calculated parameters (**Table 1**: 34, 35, 38–42, 46, 47; total

DNA fluorescence intensity, total NF-κB fluorescence intensity, and NF-κB area fraction) resulted

in a total of 33 quantitative characteristics per polyp. Area fraction (%Area) is acquired from the

table of results generated from ImageJ/Fiji (Analyze > Measure).

Xenopus. Focusing only on the acetylated-α-tubulin channel of the imaged MCCs, diameters of

several representative objects indicated by acetylated-α-tubulin staining were measured across

images to obtain a range of values for running the AutoMorFi macro. Measurements of "soma"

were excluded. "Whole cell" and "skeletonized" outlines were matched by their coordinates. Total

acetylated-α-tubulin fluorescence intensity was measured for each MCC. A total of 43 quantitative

characteristics was obtained per MCC by including the calculated parameters (**Table 1**: 34, 35,

38, 39, 40, 41, 42, 46, 47), total acetylated-α-tubulin fluorescence intensity, and morphometric

measurements of manually segmented MCCs based on *centrin4-cfp* signals (**Table 1**: 1 to 12).

Corals. Representative diameters of several corallites were acquired across coral lineages.

AutoMorFi adapted for color photography (**Supplementary Note 5**) was used. Measurements of

incomplete corallites, corallites covered by sand, and "somas" were excluded. "Whole cell" and

"skeletonized" parameters contributed to 21 morphometric measurements. Additional calculated

parameters (**Table 1**: 34, 35, 38, 39, 40, 41, 42, 46, and 47) resulted in a total of 30 quantitative

characteristics per corallite.

Data visualization. Unpaired t-tests, pairwise comparisons of morphometric parameters, and

principal component analysis (PCA) were performed between or across groups to determine the

need for dimensionality reduction[5,35] using tidyverse, ggplot2, and heatmap R packages.

Redundant morphometric features with high correlation coefficients between 0.5 and 1 were

removed to mitigate bias in subsequent cluster analysis. Unsupervised K-means clustering

analysis was performed on the reduced datasets using the R factoextra package. For data

visualization, UMAPs were generated using tidyverse, dplyr, ggplot2, and umap R packages, and

heatmaps were generated using dplyr, ggplot2, and heatmap R packages in R.4.3.2.

Statistical analysis. Paired and unpaired t-tests were performed using GraphPad Prism10. Data

are presented as mean ± s.e.m. and considered statistically significant at $p < 0.05$.

REFERENCES

1. Cajal, S. R. y. *Degeneration & Regeneration of the Nervous System*. (Oxford University Press, Humphrey Milford, 1928).

2. Lawson, L. J., Perry, V. H., Dri, P. & Gordon, S. Heterogeneity in the distribution and morphology of microglia in the normal adult mouse brain. *Neuroscience* **39**, 151–170 (1990).

3. Sierra, A. *et al.* The "Big-Bang" for modern glial biology: Translation and comments on Pío del Río-Hortega 1919 series of papers on microglia: 1919 Río-Hortega Papers on Microglia. *Glia* **64**, 1801–1840 (2016).

4. Falk, T. *et al.* U-Net: deep learning for cell counting, detection, and morphometry. *Nat. Methods* **16**, 67–70 (2019).

5. Colombo, G. *et al.* A tool for mapping microglial morphology, morphOMICs, reveals brain-region and sex-dependent phenotypes. *Nat. Neurosci.* **25**, 1379–1393 (2022).

6. Karbstein, K. *et al.* Species delimitation 4.0: integrative taxonomy meets artificial intelligence. *Trends Ecol. Evol.* (2024) doi:10.1016/j.tree.2023.11.002.

7. Reddaway, J., Richardson, P. E., Bevan, R. J., Stoneman, J. & Palombo, M. Microglial morphometric analysis: so many options, so little consistency. *Front. Neuroinformatics* **17**, (2023).

8. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).

9. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* **9**, 671–675 (2012).

10. Boulan, B. *et al.* AutoNeuriteJ: An ImageJ plugin for measurement and classification of neuritic extensions. *PLoS ONE* **15**, e0234529 (2020).

11. Longair, M. H., Baker, D. A. & Armstrong, J. D. Simple Neurite Tracer: open source software for reconstruction, visualization and analysis of neuronal processes. *Bioinformatics* **27**, 2453–2454 (2011).

12. Donadon, M. *et al.* Macrophage morphology correlates with single-cell diversity and prognosis

in colorectal liver metastasis. *J. Exp. Med.* **217**, e20191847 (2020).

13. Donovan, K. M. *et al.* Allograft Inflammatory Factor 1 as an Immunohistochemical Marker for

Macrophages in Multiple Tissues and Laboratory Animal Species. *Comp. Med.* **68**, 341–348

(2018).

14. Hagemeyer, N. *et al.* Transcriptome-based profiling of yolk sac-derived macrophages reveals

a role for Irf8 in macrophage maturation. *EMBO J.* **35**, 1730–1744 (2016).

15. Voet, S., Prinz, M. & van Loo, G. Microglia in Central Nervous System Inflammation and

Multiple Sclerosis Pathology. *Trends Mol. Med.* **25**, 112–123 (2019).

16. Yong, V. W. Microglia in multiple sclerosis: Protectors turn destroyers. *Neuron* **110**, 3534–

3548 (2022).

17. Zia, S. *et al.* Microglia Diversity in Health and Multiple Sclerosis. *Front. Immunol.* **11**, 588021

(2020).

18. Prineas, J. W. & Lee, S. Microglia subtypes in acute, subacute, and chronic multiple sclerosis.

J. Neuropathol. Exp. Neurol. **82**, 674–694 (2023).

19. Meijering, E. *et al.* Design and validation of a tool for neurite tracing and analysis in

fluorescence microscopy images. *Cytometry A* **58A**, 167–176 (2004).

20. Hogart, A., Wu, D., LaSalle, J. M. & Schanen, N. C. The comorbidity of autism with the

genomic disorders of chromosome 15q11.2-q13. *Neurobiol. Dis.* **38**, 181–191 (2010).

21. Khatri, N. & Man, H.-Y. The Autism and Angelman Syndrome Protein Ube3A/E6AP: The

Gene, E3 Ligase Ubiquitination Targets and Neurobiological Functions. *Front. Mol. Neurosci.*

12, 109 (2019).

22. Yi, J. J. *et al.* An Autism-Linked Mutation Disables Phosphorylation Control of UBE3A. *Cell*

162, 795–807 (2015).

23. Iossifov, I. *et al.* The contribution of de novo coding mutations to autism spectrum disorder.

Nature **515**, 216–221 (2014).

24. Weston, K. P. *et al.* Identification of disease-linked hyperactivating mutations in UBE3A
through large-scale functional variant analysis. *Nat. Commun.* **12**, 6809 (2021).

25. Khatri, N. *et al.* The Autism Protein Ube3A/E6AP Remodels Neuronal Dendritic Arborization
via Caspase-Dependent Microtubule Destabilization. *J. Neurosci. Off. J. Soc. Neurosci.* **38**,
363–378 (2018).

26. Smith, S. E. P. *et al.* Increased gene dosage of Ube3a results in autism traits and decreased
glutamate synaptic transmission in mice. *Sci. Transl. Med.* **3**, 103ra97 (2011).

27. Krishnan, V. *et al.* Autism gene Ube3a and seizures impair sociability by repressing VTA
Cbln1. *Nature* **543**, 507–512 (2017).

28. Walentek, P. *et al.* Ciliary transcription factors and miRNAs precisely regulate Cp110 levels
required for ciliary adhesions and ciliogenesis. *eLife* **5**, e17557 (2016).

29. Carrión, P. J. A. *et al.* Starvation decreases immunity and immune regulatory factor NF-κB in
the starlet sea anemone Nematostella vectensis. *Commun. Biol.* **6**, 698 (2023).

30. Grupstra, C. G. B. *et al.* Integrating cryptic diversity into coral evolution, symbiosis and
conservation. *Nat. Ecol. Evol.* **8**, 622–636 (2024).

31. Scott-Hewitt, N. *et al.* Microglial-derived C1q integrates into neuronal ribonucleoprotein
complexes and impacts protein homeostasis in the aging brain. *Cell* **187**, 1–20 (2024).

32. Seemann, J. *et al.* Assessing the ecological effects of human impacts on coral reefs in Bocas
del Toro, Panama. *Environ. Monit. Assess.* **186**, 1747–1763 (2014).

33. Guzmán, H. M. & Guevara, C. A. [Coral reefs of Bocas del Toro, Panamá: IV. Distribution,
structure and conservation state of continental reefs of Peninsula Valiente]. *Rev. Biol. Trop.*
49, 53–66 (2001).

34. Aichelman, H. E. Exploring coral symbiosis under climate change stress across spatial and
temporal scales. (2023).

35. van Weering, H. R. J., Nijboer, T. W., Brummer, M. L., Boddeke, E. W. G. M. & Eggen, B. J.
L. Microglia morphotyping in the adult mouse CNS using hierarchical clustering on principal

components reveals regional heterogeneity but no sexual dimorphism. *Glia* **71**, 2356–2371

(2023).

Figure 1: Acquisition and validation of whole-image cell morphometric data using the AutoMorFi prototype, AutoGliji. (A) Widefield fluorescence image of IBA-1-immunolabeled mouse cortical microglia (white) used for manual and automated morphometric analysis of 3 selected cells by 30 testers. Inset shows cell #3 corresponding to data in **B-N**. **(B)** Representative manual and automated cell thresholding and detection by 1 tester. **(C)** Representative manual trace using Simple Neurite Tracer and automated cell traces using AutoGliji by 2 testers. **(D)** Total time taken per tester for manual and automated data acquisition for cells #1–3. **(E-N)** Examples of manual and automated morphometric measurements of cell #3 for **(E)** whole-cell area, **(F)** whole-cell perimeter, **(G)** whole-cell roundness, **(H)** whole-cell circularity, **(I)** total branch length, **(J)** branch area, **(K)** branching to area ratio, **(L)** perimeter to area ratio, **(M)** whole cell convex area, and **(N)** cell spread. **(D-N)** Data are represented as mean ± s.e.m. Paired t-test. *N* = 30. Scale bars, 100 µm (overview in **A**); 10 µm (inset in **A**, **B**, and **C**).

Figure 2: Morphometric analysis of tissue-resident macrophages. (A) Representative widefield fluorescence images of IBA-1+ brain-resident microglia and liver Kupffer cells and corresponding elements generated by AutoMorFi including soma outline, whole-cell outline, and skeletonized structure of the whole-cell outline. **(B)** Representative images of homeostatic mouse IBA-1+ tissue-resident macrophages and their corresponding whole-cell outlines and skeletonized traces. **(C)** Correlation matrix of the 47 parameters from AutoMorFi across 1,945 tissue-resident macrophages. **(D)** Principal component analysis (PCA) of 1,945 macrophages based on 47 parameters from AutoMorFi. **(E)** Uniform manifold approximation and projection (UMAP) of 1,945 macrophages in 2 clusters based on 24 morphometric parameters after dimensionality reduction and unsupervised kmeans clustering. **(F)** UMAP of macrophages according to tissue origin. **(G-J)** UMAP of selected parameters, including **(G)** soma perimeter, **(H)** whole-cell perimeter, **(I)** whole-cell solidity, and **(J)** whole-cell convex area. Scale bars, 10 µm.

Figure 3: Microglial morphometric analysis in postmortem human MS and control brain. (A) Representative widefield fluorescence images of IBA-1[+] microglia (white) and their corresponding whole-cell outline and skeletonized trace. **(B)** Microglial cells analyzed from the 4 equally sampled brain areas: control white matter (WM), MS normal-appearing WM (NAWM), MS rim, and MS center. **(C-J)** UMAP of 450 microglia based on 27 morphometric parameters after dimensionality reduction from **(C)** control WM, **(D)** MS NAWM, **(E)** MS rim, and **(F)** MS center and of selected parameters including **(G)** total number of segments, **(H)** whole cell area, **(I)** soma area, and **(J)** whole cell roundness. Scale bars, 10 μm.

A **Glial cells from different models**

B **Primary neuronal cultures**

Morphometric parameter significantly different across groups

Genotypes
Wild type (14)
Ube3a 2x (14)

Morphometric parameters
1. Whole cell perimeter
2. Whole cell height
3. Whole cell minimum feret
4. Whole cell convex area
5. Perimeter to area ratio
6. Solidity

C *Xenopus* mucociliary epithelial multi-ciliated cells

*Cp110*MO (14)
*Cp110*MO-GFP-Cp110-D3'UTR-DNA (14)
*Cp110*MO-GFP-Cp110DNA (15)
Control (13)

D **Sea anemone NF-κB expression**

E **Underwater coral photography**

Figure 4: Diverse applications of AutoMorFi. (A) Comparative morphometric analysis of glial cells from different models. Representative widefield fluorescence and brightfield images of glial cells from experimental models (*Drosophila* astrocyte-like glia, hiPSCs-derived microglia, human microglia, and mouse microglia) and their corresponding whole-cell outlines. UMAPs of 1,100 glial cells clustered based on 47 morphometric parameters or model system. **(B) Single object morphometric analysis.** Representative widefield fluorescence images of MAP2-immunolabeled primary neurons (white) from Ube3a 2x transgenic and wildtype mice and their corresponding whole-cell and skeletonized traces with an example of an added pseudo-object (white arrowhead). Six of 33 morphometric parameters were significantly different between genotypes in unpaired t-tests: whole cell perimeter (p=0.0138), whole cell height (p=0.0270), whole cell minimum feret (p=0.0116), whole cell convex area (p=0.0177), perimeter to area ratio, and solidity (p=0.0351). *N* = 14 per group. **(C) Acetylated-α-tubulin marker-specific morphometric and fluorescence intensity measurements.** Representative confocal images of targeted *Xenopus* MCCs indicated by centrin4-cfp (blue) with varying ciliation indicated by acetylated-α-tubulin (red) and whole-cell outline of the marker (cyan). PCA based on 43 measurements of targeted MCCs averaged per individual embryo (N) in *Cp110* knockdown (red), rescue (green, turquoise), and control (blue) groups. Representative histograms (mean ± s.e.m.) of features with significant changes in ciliation analyzed by Kruskal-Wallis and Dunn's posthoc tests. **(D) Whole organism morphometric and NF-κB marker analyses.** Representative widefield fluorescence images of control and starved sea anemone polyps. DNA (blue), NF-κB (red), and AutoMorFi generated whole-object outline (yellow). Selected measurements (mean ± s.e.m.) are shown. Unpaired t-tests. *N* = 6 per group. **(E) Whole-image 2D morphometric analysis of photographed objects.** Representative image of corallites photographed under water and a corresponding whole-object outline and skeletonized trace. UMAPs of 628 corallites based on 21 morphometric parameters after dimensionality reduction distributed into clusters and lineages A–C. Histogram illustrates the proportion of the 3 clusters across lineages. Scale bars, 10 μm **(A-C)**; 100 μm **(D)**; 2 mm **(E)**.

Table 1 | List of 47 morphometric parameters.

Category	Parameter	Definition
"Soma"	1. Soma area	Area of the object (square micrometers)
	2. Soma perimeter	Length of the outside boundary of the object (micrometers)
	3. Soma width	Width of the smallest rectangle enclosing the object (micrometers)
	4. Soma height	Height of the smallest rectangle enclosing the object (micrometers)
	5. Soma major	Primary axis of the best fitting ellipse in the object (micrometers)
	6. Soma minor	Secondary axis of the best fitting ellipse in the object (micrometers)
	7. Soma circularity	Value of 1.0 indicates a perfect circle, $4\pi*area/perimeter^2$)
	8. Soma feret	Maximum caliper, longest distance between any two points in the object
	9. Soma minimum feret	Minimum caliper diameter
	10. Soma aspect ratio	Major axis/minor axis
	11. Soma roundness	Inverse of the aspect ratio, $4*area/(\pi*major\ axis^2)$
	12. Soma solidity	Area / convex area
Whole-cell	13. Whole cell area	Area of the object (square micrometers)
	14. Whole cell perimeter	Length of the outside boundary of the object (micrometers)
	15. Whole cell width	Width of the smallest rectangle enclosing the object (micrometers)
	16. Whole cell height	Height of the smallest rectangle enclosing the object (micrometers)
	17. Whole cell major (X)	Primary axis of the best fitting ellipse in the object (micrometers)
	18. Whole cell minor (Y)	Secondary axis of the best fitting ellipse in the object (micrometers)
	19. Whole cell circularity	Value of 1.0 indicates a perfect circle, $4\pi*area/perimeter^2$)
	20. Whole cell feret	Maximum caliper, longest distance between any two points in the object
	21. Whole cell minimum feret	Minimum caliper diameter
	22. Whole cell aspect ratio	Major axis/minor axis
	23. Whole cell roundness	Inverse of the aspect ratio, $4*area/(\pi*major\ axis^2)$
	24. Whole cell solidity	Area / convex area
"Skeletonized"	25. Total number of segments	Total number of slab segments
	26. Number of junctions	Number of actual segment junctions
	27. Number of end point voxels	Voxel with less than 2 neighbors
	28. Number of junction voxels	Voxel with more than 2 neighbors
	29. Number of slab voxels	Voxel with exactly 2 neighbors
	30. Average branch length	Average length of branches (micrometers)
	31. Number of triple points	Number of junctions with exactly 3 branches
	32. Number of quadruple points	Number of junctions with exactly 4 branches
	33. Maximum branch length	Maximum length of branches (micrometers)
Additional parameters	34. Total branch length	Maximum branch length / Number of junction voxels
	35. Cell spread	Whole cell aspect ratio / Number of end point voxels
	36. Branch area ratio	Whole cell area / Soma area
	37. Branch perimeter ratio	Whole cell perimeter / Soma perimeter
	38. Whole cell convex area	Whole cell solidity / Whole cell area
	39. Nodes	Number of end point voxels + Junction voxels + Slab voxels
	40. Ramification index	Whole cell perimeter / Whole cell area)/2 x [$\sqrt{(\pi Whole\ cell\ area\ x\ XY)}$]
	41. Perimeter to area ratio	Whole cell perimeter^2/ Whole cell area
	42. Density	Whole cell area / (Whole cell major x Whole cell minor)
	43. Process soma area ratio	Branch area / Soma area
	44. Process whole cell area ratio	Branch area / Whole cell area
	45. Skeleton to process ratio	Total branch length^2/ Branch area ratio
	46. Branch points endpoints ratio	Number of slab voxels / Number of end point voxels
	47. Whole cell solidity	Whole cell area/ Whole cell convex area

Table 2 | Morphometric analysis of IBA-1-immunostained microglia in postmortem human brain tissue.

Group	Patient ID	Area analyzed (μm^2)	Mean NND ± s.e.m. (μm)
Control WM	Control_56BT1, Control_56B9F, Control_31B1	320,000	41.69 ± 2.48
MS NAWM	MS_590939, MS_560L2	320,000	36.41 ± 1.93
MS Center	MS_590939, MS_560L2, MS_590937	320,000	38.62 ± 4.37
MS Rim	MS_590939, MS_560L2, MS_590937	320,000	27.42 ± 1.87

WM, white matter. MS, multiple sclerosis. NAWM, normal appearing WM.

SUPPLEMENTARY NOTE 1

Prerequisites to running AutoMorFi.
Step I: Download and add the following plugins to the "plugins" folder of your ImageJ:
- "Excel.macro.extensions-3.3.2-jar-with-dependencies.jar"
- "script-editor-0.7.4-SNAPSHOT.jar"

To find the "plugins" folder on Mac:
- Go to Finder > Desktop > Search: "Fiji"
- Right-click on the Fiji logo > "Show package contents"
- Open the "plugins" folder
- Copy-and-paste the downloaded plugins

Windows

 Mac

Step II: Update ImageJ/Fiji
- Start ImageJ/Fiji and select Help > Update… > Manage update site
- Select: ✅Excel Functions ✅Neuroanatomy
- Unselect: ResultToExcel
- Close the "Manage update sites" window
- Close the "ImageJ Updater" window
- Restart ImageJ/Fiji

Step III: Range of measurements
1. Ensure your version of Fiji/ImageJ is updated before each analysis.
2. Set your Fiji/ImageJ as follows:
   - Analyze > Set measurements
   - Select the top 18 criteria
   - Click "OK"

Window   Mac

3. AutoMorFi requires the user to enter the size range of the cell soma and whole cell areas. To get these values:
   - Drag and drop the image you want to analyze onto Fiji/ImageJ
   - Make sure to work with a single-channel .tif image that is 8-bit and scaled (i.e., unit per pixel).
   - Use the 'straight' tool to measure the diameter (in microns) of 5–10 cell somas and whole cells representing the average, smallest, and largest objects in the image.
   - With these diameter measurements, estimate the size ranges needed in the Analyze Particles wizard embedded within AutoMorFi. For example, if the soma diameters measured are 6.74, 7.56, 8.43, 13.74, and 12.33, the lower threshold value would be obtained as $6.74^2 = 45.43$ and the upper threshold value would be obtained as $13.74^2 = 188.79$. The range entered in the wizard could be 42-200.
   - Optimization tip: the ranges of soma and whole cell areas depend on the type of image and objects you are working with (e.g., cell type, cell density, method of image acquisition, etc.). After the first run, evaluate the accuracy of your soma and whole cell ranges by looking at the soma (SomaOutlines.tiff) and whole cell (WholeCellOutlines.tiff) outlines in your output folder.
     - too many nonspecific objects identified –> increase lower threshold value
     - small objects not captured –> decrease the lower threshold value
     - big objects not captured –> increase the upper threshold value

Windows

Mac

SUPPLEMENTARY NOTE 2

Running the AutoMorFi macro
AutoMorFi analyzes **one scaled single-channel 8-bit TIFF image** per cycle. Make sure the folder path leading to your image contains **no space, slash, or other special characters**. Use only letters, numerals, underscore, and dash.
Close any window that is open on Fiji/ImageJ. To start the AutoMorFi macro (AutoMorFi_3.ijm):
 1. drag and drop the macro file in Fiji/ImageJ > press [Run]
 2. select your image ("xxx.tif") > Click "Open"

Step 1: object identification by thresholding
On your screen, the original image (named "Temp") and the "Skeleton" image are displayed.
In the "Threshold" window, adjust the threshold of the "Skeleton" image:
 1. select the "Skeleton" image (an error message will result if a different image is selected)
 2. adjust the auto threshold method (ex: Default)
    Tip: set the display method to "Red" (instead of B&W) in the "Threshold" window
 3. adjust the threshold using the first sliding bar
 4. **click "Apply"** in the "Threshold" window
 5. click "OK" to continue the analysis

Windows

Mac

Step 2: connect disrupted processes using dilation (for objects with branches)

- Dilate Option window: if your cells are not branched, select "No" > click OK
- Dilate Option window: to connect disrupted processes, select "Yes" > click "OK"
- to **finish** the dilation: select "Finished" > "OK"
- to **revert** the dilation: select "Revert" > "OK" (only one dilation can be reversed)
- to **continue** dilating: select "Continue" > "OK" > "Yes" > "OK"

Windows

Mac

Step 3: object detection by diameter ranges

1. The **first** Analyze Particles window will analyze your cell **soma**
   - In "Size (micron)^2", enter the calculated soma size <u>range</u>
   - Show: <u>Masks</u>, ☑ Display results, ☑ Add to manager, ☑ Overlay > "OK"
2. The **second** Analyze particles window will analyze your **whole cell**
   - In "Size (micron^2)", enter the calculated whole cell size <u>range</u>
   - Show: <u>Masks</u>, ☑ Display results, ☑ Add to manager, ☑ Overlay > "OK"

Windows   Mac

Step 4: deletion of false positive objects in the ROI manager

1. Select the objects by their number in the ROI manager (click on "Labels" if the numbers are not visible)
2. Click "Delete"
3. When you are done, click "Update" > "Deselect"
4. Click "OK" in the "ROI Deletion" window

Let the macro run until the "Finish" window appears.

SUPPLEMENTARY NOTE 3

Troubleshooting for AutoMorFi

Error 1. Macro does not start
The image format, name, or image path is incorrect.
Solution 1: AutoMorFi only works on one-channel 8-bit TIFF images.
Solution 2: Make sure the folder path leading to your image contains no space, slash or special characters. Use only underscore or dash.

Error 2. Macro does not continue after step 1
A different image than "Skeleton" was selected for thresholding during step 2. The user did not click "Apply" in the Threshold window before clicking "OK" in the Action Required window.
Solution: Make sure to adjust the threshold of the "Skeleton" image in the Threshold window and click "Apply" in the Threshold wizard before "OK" in the Action Required window.

Error 3: Macro does not continue after step 3
No particle was detected based on the particle range set by the user in step 3 of AutoMorFi. The macro does not proceed without the "Results" window.
Solution 1: Check that the calculated size range provided for particle detection is correct (see **Step III** in **Supplementary Note 1**).

Solution 2: Locate the .tif image "Binary" in the output folder of the aborted AutoMorFi run. If the objects of interest are incorrectly represented in "Binary", adjust the extend of thresholding in step 1 (**Supplementary Note 2**).

Error 4. Only one row of data is shown in the BranchInformation.csv and SkeletonResult.csv output files
The user did not click "Update" followed by "Deselect" in the ROI Manager window before clicking "OK" during **step 4 (deletion of false positive objects in the ROI manager).**

Error 5. Unresponsive macro or to interrupt the current run
Solution: Click "Kill", at the bottom of the macro window.

Error 6. "Default" thresholding incorrectly detects "soma" of the objects

AutoMorFi has a built-in automatic thresholding function for "soma" identification instead of the manual thresholding for whole-cell object identification in step 1 (i.e., object identification by thresholding). This can lead to incorrect "soma" identification in the SomaOutline.tif output.

Solution: Choose the appropriate AutoThreshold method to use before running AutoMorFi:
- Open your image on ImageJ/Fiji to identify the optimal AutoThreshold method
- Select: Image > Adjust > AutoThreshold (https://imagej.net/plugins/auto-threshold)
- In the AutoMorFi macro window, replace "Default" with the optimal AutoThreshold method (case-sensitive) in line 68 of the standard AutoMorFi code in **Supplementary Note 4**
- Run AutoMorFi as usual

```
68    setAutoThreshold("Default");
```

Error 7. Macro does not generate output from image containing only one object

AutoMorFi requires a minimum of 2 objects per image to run.

Solution: Create a second object in the ROI manager after the primary object during step 4:

- In the ROI manager, click "More" > "Draw"
- Draw a rectangle in the background of the image (i.e., not on the object of interest)
- Adding the second object in the bottom right corner of the image is recommended for coordinate-based synchronization of "soma", "whole cell", and "skeleton" outputs using the cell matching app in **Supplementary Note 6**
- When completed, click "Update" > "Deselect"
- Click "OK" in the "ROI Deletion" window

SUPPLEMENTARY NOTE 4

<u>CODE FOR AUTOMORFI MACRO (IMAGEJ/FIJI-COMPATIBLE)</u>

```
//AutoMorFi_3 macro

// clean up: makes sure no data is stored from previous macro use
// close all images
close("*");
// empty the ROI manager
roiManager("reset");
// empty the results table
run("Clear Results");
// configure that binary image are black in background, objects are white
setOption("BlackBackground", true);

//Obtain Image Path, Folder Path
path = File.openDialog("Choose a File");
basename = File.nameWithoutExtension();
dirname = File.getParent(path);
folder = dirname + File.separator + basename;

//Create Result Folders
if (File.exists(folder)) {
    print("Directory already exists, override the existing directory");
    list = getFileList(folder);
    for (i = 0; i < list.length; i++) {
        File.delete(folder + File.separator + list[i]); // Delete each file
        }
}
else {
    File.makeDirectory(folder);
}

//Open Image thru Bio-Format Importer
run("Bio-Formats Importer", "open="+path);
imageOriginPath = folder + File.separator + basename;
saveAs("tiff", imageOriginPath);

//Get Image size, Set scales
```

```
getPixelSize(unit, pixelWidth, pixelHeight);
scale = 1 / pixelWidth;
nameOrigin = getTitle();
microWidth = getWidth() / scale;
microHeight = getHeight() / scale;

//All Output Directories
binaryPath = folder + File.separator + "Binary";
somaResultsPath = folder + File.separator + "SomaResults.csv";
somaOutlinesPath = folder + File.separator + "SomaOutlines";
wholeCellResultsPath = folder + File.separator + "WholeCellResults.csv";
wholeCellOutlinesPath = folder + File.separator + "WholeCellOutlines";
skeletonPath = folder + File.separator + "Skeleton";
skeletonResultsPath = folder + File.separator + "SkeletonResults.csv";
branchInformationPath = folder + File.separator + "BranchInformation.csv";

//Duplicate original input twice. One for skeleton, the other for threshold reference.
run("Duplicate...", " ");
rename("Skeleton");
run("Duplicate...", " ");
rename("Temp");

//Process Binary Image
selectWindow(basename+".tif");
run("Enhance Contrast", "saturated=0.35");
run("Apply LUT");
run("8-bit");
setAutoThreshold("Default");
run("Convert to Mask");
run("Invert LUT");
run("Make Binary");
run("Despeckle");
rename("Binary");

//Process Skeleton, Manually set threshold
selectWindow("Skeleton");
run("Threshold...");//https://imagej.net/plugins/auto-threshold#available-methods
waitForUser;   //User adjust threshold here
close("Threshold");
close("Temp");
```

```
//Dilation Dialogs
selectWindow("Skeleton");
stack_count = 0; //Keep track of the number of dilations
rename("Mask_" + stack_count);
while (true) {
    //Creating dialog with dilate option "Yes"/"No"
    Dialog.create("Dilate Option");
    Dialog.addChoice("Do you want to dilate the image?", newArray("Yes", "No"));
    Dialog.show();
    choice = Dialog.getChoice();

    //When choosing "Yes", dilate the image from the previous stack
    if (choice == "Yes") {
        selectWindow("Mask_" + stack_count);
        run("Duplicate...", "title=Temp");
        run("Dilate");
        stack_count++;
        selectWindow("Temp");
        rename("Mask_" + stack_count);

        //Creating dialog with options "Continue", "Revert", "Exit"
        Dialog.create("Continue or revert?");
        Dialog.addChoice("Continue dilation?", newArray("Continue", "Revert", "Finished"));
        Dialog.show();
        secondChoice = Dialog.getChoice();
        //When Continue, enter the next iteration
        //When Revert, return to the previous iteration
        //When Exit, save the current iteration and process to skeletonization
        if (secondChoice == "Continue") {
                        close("Temp");
        } else if (secondChoice == "Revert") {
            close("Temp");
            close("Mask_" + stack_count);
            stack_count--;
        } else {
            close("Temp");
            break;
        }
    } else {
        break;
    }
}
for (i = 0; i < stack_count; i++) {
        close("Mask_" + i);
```

```
    }
selectWindow("Mask_" + stack_count);
//Dilation Dialog ends

//Create skeleton, overlay with Binary using imageCalculator
rename("Skeleton");
run("Skeletonize");
imageCalculator("add create", "Binary", "Skeleton");
close("Skeleton");
close("Binary");
selectWindow("Result of Binary");
rename("newBinary");
saveAs("tiff", binaryPath);
rename("newBinary");

//Soma Analysis
selectWindow("newBinary");
run("Duplicate...", " ");
rename("newBinary2");
run("Watershed");
run("Clear Results");
run("Analyze Particles...");
n = roiManager("count");
somaArray = newArray(n);
for (var i = 0; i < n; i++) {
    somaArray[i] = i;
}
selectWindow("Results");
saveAs("Measurements", somaResultsPath);
run("Clear Results");
selectWindow("newBinary2");
open(imageOriginPath+".tif");
run("Enhance Contrast", "saturated=0.35");
run("Apply LUT");
roiManager("Select", somaArray);
selectWindow(basename+".tif");
setOption("Show All", true);
run("Flatten");
saveAs("tiff",somaOutlinesPath);
run("Close");
close("newBinary2");
roiManager("reset");
```

```
//Cell Analysis
selectWindow("newBinary");
run("Analyze Particles...");
n1 = roiManager("count");
// Can delete ROIs here
waitForUser("ROI Deletion", "In the ROI manager list, delete ROIs (using [Delete]). Click 'OK' to
continue.");
n2 = roiManager("count");
if (n2 < n1) {
        run("Clear Results");
        roiManager("multi-measure append");
}
cellArray = newArray(n2);

for (var i = 0; i < n2; i++) {
    cellArray[i] = i;
}
open(imageOriginPath+".tif");
run("Enhance Contrast", "saturated=0.35");
run("Apply LUT");
roiManager("Select", cellArray);
selectWindow(basename+".tif");
setOption("Show All", true);
run("Flatten");
saveAs("tiff",wholeCellOutlinesPath);
roiManager("XOR");
run("Create Mask");
selectWindow("Results");
saveAs("Measurements", wholeCellResultsPath);
close("newBinary");
run("Clear Results");
close("newBinary");

//Skeletonize and Analyze Skeletons
selectWindow("Mask");
rename("Skeleton");
run("Skeletonize");
saveAs("tiff",skeletonPath);
rename("Skeleton");
run("Analyze Skeleton (2D/3D)","prune=[lowest intensity branch] show display");
selectWindow("Results");
saveAs("Measurements", skeletonResultsPath);
```

```
selectWindow("Branch information");
saveAs("Measurements", branchInformationPath);

//Close all windows, end macro.
close("*");
selectWindow("ROI Manager");
run("Close");
selectWindow("Results");
run("Close");
if (isOpen("BranchInformation.csv")) {
        selectWindow("BranchInformation.csv");
        run("Close");
}

waitForUser("Finish", "Results in the output folder. Click 'OK' to exit.");
```

SUPPLEMENTARY NOTE 5

AUTOMORFI MACRO FOR COLOR PHOTOGRAPHY (IMAGEJ/FIJI-COMPATIBLE)

```
//AutoMorFi_3 macro adapted for color photography

close("*");
roiManager("reset");
run("Clear Results");
setOption("BlackBackground", false);

path = File.openDialog("Choose a File");
basename = File.nameWithoutExtension();
dirname = File.getParent(path);
folder = dirname + File.separator + basename;
if (File.exists(folder)) {
    print("Directory already exists, override the existing directory");
    list = getFileList(folder);
    for (i = 0; i < list.length; i++) {
        File.delete(folder + File.separator + list[i]); // Delete each file
        }
}
else {
    File.makeDirectory(folder);
}

binary = folder + File.separator + "Binary";
objectOutline = folder + File.separator + "ObjectOutline";
objectMeasure = folder + File.separator + "ObjectMeasurement.csv";
somaResultsPath = folder + File.separator + "SomaResults.csv";
somaOutlinesPath = folder + File.separator + "SomaOutlines";
skeletonPath = folder + File.separator + "Skeleton";
skeletonResultsPath = folder + File.separator + "SkeletonResults.csv";
branchInformationPath = folder + File.separator + "BranchInformation.csv";

run("Bio-Formats Importer", "open="+path);
imageOriginPath = folder + File.separator + basename;
saveAs("tiff", imageOriginPath);

getPixelSize(unit, pixelWidth, pixelHeight);
```

```
scale = 1 / pixelWidth;
nameOrigin = getTitle();
microWidth = getWidth() / scale;
microHeight = getHeight() / scale;

selectWindow(basename+".tif");
run("Duplicate...", " ");
rename("Binary");
run("8-bit");
run("Enhance Contrast", "saturated=0.35");
run("Smooth");
saveAs("tiff",binary);
run("Threshold...");
waitForUser;
close("Threshold");
run("Despeckle");
run("Invert LUT");

stack_count = 0; //Keep track of the number of dilations
rename("Mask_" + stack_count);
while (true) {
    //Creating dialog with dilate option "Yes"/"No"
    Dialog.create("Erode Option");
    Dialog.addChoice("Do you want to erode the image?", newArray("Yes", "No"));
    Dialog.show();
    choice = Dialog.getChoice();

    //When choosing "Yes", dilate the image from the previous stack
    if (choice == "Yes") {
        selectWindow("Mask_" + stack_count);
        run("Duplicate...", "title=Temp");
        run("Erode");
        stack_count++;
        selectWindow("Temp");
        rename("Mask_" + stack_count);

        //Creating dialog with options "Continue", "Revert", "Exit"
        Dialog.create("Continue or revert?");
        Dialog.addChoice("Continue erosion?", newArray("Continue", "Revert", "Finished"));
        Dialog.show();
        secondChoice = Dialog.getChoice();
        //When Continue, enter the next iteration
        //When Revert, return to the previous iteration
```

```
        //When Exit, save the current iteration and process to skeletonization
        if (secondChoice == "Continue") {
                        close("Temp");
        } else if (secondChoice == "Revert") {
           close("Temp");
           close("Mask_" + stack_count);
           stack_count--;
        } else {
           close("Temp");
           break;
        }
    } else {
        break;
    }
}
for (i = 0; i < stack_count; i++) {
        close("Mask_" + i);
}
selectWindow("Mask_" + stack_count);
rename("Binary");

run("Duplicate...", " ");
rename("Soma");
run("Duplicate...", " ");
rename("Skeleton");

selectWindow("Binary");
run("Analyze Particles...");
n1 = roiManager("count");
// Can delete ROIs here
waitForUser("ROI Deletion", "In the ROI manager list, delete ROIs (using [Delete]). Click 'OK' to
continue.");
n2 = roiManager("count");
if (n2 < n1) {
        run("Clear Results");
        roiManager("multi-measure append");
}
objectArray = newArray(n2);

for (var i = 0; i < n2; i++) {
    objectArray[i] = i;
}
```

```
selectWindow("Results");
saveAs("Measurements", objectMeasure);
close("Results");
roiManager("Select", objectArray);
selectWindow(basename+".tif");
setOption("Show All", true);
run("Flatten");
saveAs("tiff",objectOutline);
run("Close");
close("Binary");
roiManager("reset");
run("Clear Results");

selectWindow("Soma");
run("Watershed");
run("Analyze Particles...");
n = roiManager("count");
somaArray = newArray(n);
for (var i = 0; i < n; i++) {
    somaArray[i] = i;
}
selectWindow("Results");
saveAs("Measurements", somaResultsPath);
run("Clear Results");
selectWindow("Soma");
open(imageOriginPath+".tif");
run("Enhance Contrast", "saturated=0.35");
roiManager("Select", somaArray);
selectWindow(basename+".tif");
setOption("Show All", true);
run("Flatten");
saveAs("tiff",somaOutlinesPath);
run("Close");
close("Soma");
run("Clear Results");

selectWindow("Skeleton");
run("Skeletonize");
saveAs("tiff",skeletonPath);
rename("Skeleton");
run("Analyze Skeleton (2D/3D)","prune=[lowest intensity branch] show display");
selectWindow("Results");
saveAs("Measurements", skeletonResultsPath);
```

```
selectWindow("Branch information");
saveAs("Measurements", branchInformationPath);

close("*");
selectWindow("ROI Manager");
run("Close");
selectWindow("Results");
run("Close");
```

SUPPLEMENTARY NOTE 6

<u>CELL MATCHING APP ("ANALYZER")</u>

AutoMorFi generates "soma" and "whole cell" measurements in separate .csv output files. The cell matching algorithm combines the results by correlating every soma to its whole cell by using their common xy-coordinate system to find the soma with the shortest Euclidean distance to its cell. The algorithm runs on Python using the K-Nearest-Neighbours (KNN) model, which finds the k value of nearest neighbors to a given data point using the Euclidean distance between two points. Notably, in rare instances, a detected soma may be deemed have the shortest distance to a target cell to which the soma does not belong. To ensure an accurate match, an arbitrary default threshold is used to limit matched soma-cell pairs to within 12 µm in horizontal, vertical, and Euclidean distances.

Tutorial for Cell Matching App
1. Open **Command Prompt**
2. Connect to the folder containing the scripts (Supplementary Software 1)
   cd path/to/your/folder
3. Create the python virtual environment by **python -m venv "name"**
   Make sure Python is installed. Check Python version by typing **python -V**
4. Activate virtual environment by **.\name\Scripts\activate**

5. When the virtual environment is activated, your command prompt switches to a new window.

6. Use pip or pip3 to install all required libraries including numpy, scipy, scikit-learn, pandas, openpyxl, tk, and tkinterdnd2-universal. Check for all required libraries in **install.sh**. The installation statement is in the first line.

7. When all the libraries are installed, run **python main.py**

8. For future use, double click **main.py** to open the cell matching app without using the command prompt.

Troubleshooting for Cell Matching App

1. Use Python versions 3.7, 3.8, or 3.9.
2. Try different installing commands: pip, pip3, or pip3.x according to your version of Python.
3. Install tkinterdnd2-universal instead of tkinterdnd2 because the latter is incompatible with GitHub branches.
4. Run the code inside a virtual environment.
5. Run the code as the PC administrator.