Evidence of centromeric histone 3 chaperone involved in DNA damage repair

pathway

Authors: Prakhar Agarwal and Santanu Kumar Ghosh[*]

Affiliation: Department of Biosciences and Bioengineering, Indian Institute of Technology,
Bombay, Powai, Mumbai-400076, India.

*Corresponding author

Email: santanughosh@iitb.ac.in

Tel. +(91-22) 2576 7766

Fax. (+91)22 2572 3480

Key words: Scm3, HJURP, DNA damage, centromere, chromosome, budding yeast

Abstract

The centromeric protein-A (CENP-A) is an evolutionary conserved histone H3 variant that

marks the identity of the centromeres. Several mechanisms regulate the centromeric

deposition of CENP-A as its mislocalization causes erroneous chromosome segregation,

leading to aneuploidy-based diseases, including cancers. The most crucial deposition factor

is a CENP-A specific chaperone, HJURP (Scm3 in budding yeast), which specifically binds to

CENP-A. However, the discovery of HJURP as a DDR (DNA damage repair) protein and

evidence of its binding to Holliday junctions in vitro indicate a CENP-A-deposition-independent

role of these chaperones. In this study, using budding yeast, we demonstrate that Scm3 is

crucial for the DDR pathway as *scm3* cells are sensitive to DNA damage. We further observe

that the *scm3* mutant interacts with the *rad52* DDR mutant and is compromised in activating

DDR-mediated arrest. We demonstrate that Scm3 associates with the DNA damage sites and

undergoes posttranslational modifications upon DNA damage. Overall, from this report and

earlier studies on HJURP, we conclude that DDR functions of CENP-A chaperones are

conserved across eukaryotes. Thus, the revelation that these chaperones confer genome

stability in more than one pathway has clinical significance.

Introduction

The faithful replication and partitioning of the replicated chromatin as chromosomal units from mother to daughter cell is important for cell survival. New nucleosomes, the units of chromatin, are assembled on DNA during replication that requires deposition of the histones within the nucleosomes in a highly regulated step-by-step process that involves the deposition of $(H3-H4)_2$ tetramer followed by two H2A-H2B dimers (Hammond et al., 2017; Ransom et al., 2010). The deposition of the histones to their specific positions within the nucleosome occurs in a temporal manner assisted by a group of proteins called histone chaperones. For this, the chaperones physically interact with the histones in a pre-nucleosomal complex and thus prevent premature histone-DNA interaction or nucleosome assembly (Burgess and Zhang, 2013; Ransom et al., 2010). The deposition of the canonical histones at the gross chromatin takes place mostly during DNA replication via a replication-coupled (RC) pathway using the chaperones dedicated to the RC pathway (Serra-Cardona & Zhang, 2018). For instance, Asf1 and CAF-1 deposit $(H3-H4)_2$ tetramer (Burgess and Zhang, 2013; De Koning et al., 2007) and Nap1 facilitates $(H2A-H2B)_2$ assembly (De Koning et al., 2007; Eitoku et al., 2008) into the nucleosomes. On the other hand, the variant histones that earmark specialized chromatin domains committed to do specific biological functions are deposited at specific sites throughout the cell cycle using replication-independent (RI) pathway using variant-specific chaperones dedicated to RI pathway (Smith, 2002). For Instance, deposition of the variant histones H2A.Z, centromeric H3 (CenH3), MacroH2A signifying transcriptional activation (Giaimo et al., 2019), kinetochore formation (Earnshaw and Rothfield, 1985; Earnshaw and Tomkiel, 1992; McKinley and Cheeseman, 2016) and X chromosome inactivation (Sun and Bernstein, 2019) occurs via chromatin remodeler (Fan et al., 2022), HJURP chaperone (Dunleavy et al., 2009; Foltz et al., 2009) and acidic nuclear phosphoprotein 32B (ANP32B) (Mandemaker et al., 2023), respectively.

Almost all the eukaryotes harbor CenH3 at the centromeres, which promotes the formation of kinetochore at those loci (Blower and Karpen, 2001; Buchwitz et al., 1999; Howman et al., 2000; Stoler et al., 1995; Van Hooser et al., 2001). Since the kinetochores bind to the microtubules and drive faithful chromosome segregation, deposition of CenH3 at the centromeres is essential for cell survival. In metazoans, CENP-A (Centromeric protein- A) is the CenH3 (Earnshaw and Rothfield, 1985), and HJURP (Holliday junction recognition protein) is the cognate chaperone (Dunleavy et al., 2009; Foltz et al., 2006). The CENP-A binding domain within HJURP present at its N-terminal is highly conserved (Aravind et al., 2007; Sanchez-Pulido et al., 2009; Shuaib et al., 2010). Despite the well-established function of HJURP in CENP-A deposition at the centromeres, HJURP was initially identified as a

protein involved in DNA damage response (DDR) due to the following reasons (Kato et al.,
2007). HJURP was so named because of its ability to bind in vitro to Holliday junction DNA
substrates that are in vivo generated during DNA damage repair through homologous
recombination. In addition, HJURP was found to interact physically with DNA damage sensing
complex MRN (Mre11-Rad50-Nbs1), and the interaction was found to increase significantly
upon induction of double-strand breaks (DSBs). As a corollary, the level and subcellular
localization of HJURP change in the cells treated with DNA-damaging agents such as Infrared
Radiation (IR), Hydroxyurea (HU), cisplatin, and camptothecin (CPT). Finally, the expression
of HJURP was found to be regulated by ATM, a DDR kinase, as no expression was observed
in ATM-deficient cancerous cell lines. Perhaps due to genome wide function on DDR, HJURP
has been visualized to be diffusely present throughout the nucleus (Dunleavy et al., 2009;
Kato et al., 2007).

Given the functions of HJURP in CENP-A deposition and DDR pathway, it is not
surprising that misregulation of HJURP is involved in disease states. HJURP is considered an
oncogene as its overexpression has been associated with several cancers (Chen et al., 2018;
Kang et al., 2020) including breast cancer (Coates et al., 2010; Hu et al., 2010; Montes de
Oca et al., 2015), lung cancer (Kato et al., 2007), ovarian cancer (Dou et al., 2022),
glioblastoma (Valente et al., 2013, 2009), hepatocellular carcinoma (Chen et al., 2018; Hu et
al., 2017; Luo et al., 2022), colorectal cancer (Kang et al., 2020) and in pancreatic cancer
(Wang et al., 2020). Consequently, the inhibition of HJURP attenuates the properties of the
cancer cells and increases the survival rate of the patients, and thus, HJURP acts as a
prognostic marker for several cancers (Kang et al., 2020; Lai et al., 2021).

In budding yeast, Cse4 and Scm3 are the homologs of CENP-A and HJURP,
respectively. Scm3 deposits Cse4 at the centromeres during the S phase through its physical
interaction with Ndc10 that binds to the centromeric DNA as a part of the CBF3 complex
(Camahort et al., 2007; Cho and Harrison, 2012, 2011). It has been shown that following the
deposition of Cse4, Scm3 remains associated with the centromeres throughout the cell cycle
(Wisniewski et al., 2014; Xiao et al., 2011). Like HJURP, Scm3 has been found on the gross
chromatin besides its localization at the centromeres (Mizuguchi et al., 2007). Furthermore, a
comparison of the domain organization between HJURP and Scm3 suggests that besides the
highly conserved CENP-A binding domain, both HJURP and Scm3 also harbour homologous
DNA binding domain (Figure S1), which is predicted to be evolutionarily conserved in all
vertebrates and fission yeast (Müller et al., 2014). Notably, overexpressed Scm3, like HJURP,
shows chromosome instability, and it is believed that Scm3, when excess, binds to the
centromeric DNA independently and leads to chromosome missegregation (Choy et al., 2012;

Mishra et al., 2011). Both the proteins dimerizes and in *in-vitro* assay, they can bind even to

non-*CEN* DNA (Kato et al., 2007; Xiao et al., 2011). While there is no mammalian homolog of

budding yeast Ndc10 and no yeast homolog of mammalian Mis18, both these proteins are

critical for CENP-A deposition as they bind to their respective CENP-A chaperones through

regions independent of CENP-A or DNA binding domain. Therefore, despite the large

difference in the polypeptide length between HJURP and Scm3 (748 vs. 229), most of the

functions are highly conserved from humans to yeast. Given the physical interaction of HJURP

with the MRN complex, it is highly plausible that Scm3 might also interact with DDR proteins

in yeast. Altogether, from the above observations, we hypothesize that Scm3, like HJURP,

may also be involved in the DDR pathway in budding yeast. In this work, we report that the

cells lacking Scm3 are sensitive to the DNA damaging agents and are compromised both in

DNA damage repair. In addition to the known functions of HJURP in DDR pathway, in this

report we show, using Scm3, that these chaperones are required for mounting DDR-

dependent checkpoint activation. Importantly, we demonstrate for the first time that they

indeed associate with the DNA damage sites upon induction of DNA damage and persists

there perhaps to facilitate damage repair. Our results suggest that the phosphorylation of

Scm3 may be attributed to the DDR functions of Scm3. Taken together, this study reveals an

evolutionary kinship amongst CenH3 chaperones in conferring genome stability through

efficient DNA damage repair, besides their canonical role in CenH3 deposition.

Results

Scm3 depleted cells are sensitive to MMS mediated DNA damage.

The yeast cells compromised in DNA damage repair (DDR) function show sensitivity to DNA

damaging agents such as methyl methane sulfonate (MMS), HU and CPT. To investigate the

role of Scm3 in DDR, we constructed an auxin-inducible (Nishimura et al., 2009) Scm3

conditional mutant since the protein is essential for growth (Camahort et al., 2007; Mizuguchi

et al., 2007; Stoler et al., 2007), and assayed the behaviour of the mutant in the presence of

the DNA damaging agents. Since the cells devoid of the homologous recombination protein,

Rad52, are known to show extreme sensitivity to such drugs, *rad52Δ* cells were taken as a

positive control in our assays (Hallwyl et al., 2008; Kaytor and Livingston, 1994). Mid-log grown

cells were harvested, serially diluted, and spotted on plates supplemented with auxin and DNA

damaging agents (MMS, HU, CPT) in different combinations and found similar results for all

the drugs (Figure 1A, Figure S2A). We observed that *scm3* mutants (three independent

transformants) were more sensitive in the presence of MMS (plate supplemented with both

auxin and MMS) as compared to those without MMS (plate supplemented with only auxin),

whereas *rad52Δ* cells, as expected, were sensitive to MMS but not to auxin (Figure 1A). To

quantify the levels of sensitivity towards MMS, we performed a CFU count assay where *SCM3-*
AID cells were pregrown in rich media without auxin till the mid-log phase before an equal
number of cells were plated on media supplemented with auxin and MMS in different
combinations as indicated in Figure 1A. The number of colonies formed after 36 hrs were
counted and are quantified as follows. The number of colonies that appeared on the plates
free of auxin and MMS were taken as 100% cell viability; with respect to this, the number of
colonies that appeared on other plates (conditions) was used to calculate the viability
percentage for those conditions. We observed a statistically significant drop in cell viability in
the plate supplemented with both auxin and MMS compared to auxin alone (Figure 1B).

Since the presence of auxin in the plate can hinder the growth of the *SCM3-AID* cells,
in another approach to examine the sensitivity of *scm3* mutant to MMS, we pre-treated the
cells with 0.75 mM auxin for 2 hrs to deplete Scm3 before they were washed and spotted on
plates containing different concentrations of MMS, but without auxin. The depletion of Scm3
was verified by observing higher percentage of G2/M arrested cells (Figure S2B). Here also
we found that Scm3-depleted cells were more sensitive to MMS as compared to the wild-type
cells (Figure S2C). To quantify the sensitivity observed by this approach, we performed a CFU
count assay as above. Following auxin pretreatment, an equal number of wild-type, *SCM3-*
AID, and *rad52Δ* cells were spread on the YPD plate or YPD plate containing 0.01% MMS. By
calculating the viability percentage as described above, we again found a statistically
significant drop in cell viability on the plate where Scm3 was predepleted (Figure S2D). The
rad52Δ cells, as the positive control, showed high sensitivity, as expected.

The increased sensitivity of *scm3*-depleted cells to DNA-damaging agents could be
due to weakening of the kinetochores as Scm3-mediated deposition of Cse4 promotes
kinetochore assembly (Camahort et al., 2007; Cho and Harrison, 2011). If this holds true,
perturbation of the kinetochore by degradation of other kinetochore proteins must show a
similar sensitivity to MMS. In budding yeast, Ndc10 is recruited to the centromeres upstream
of Scm3 (Lang et al., 2018), whereas the centromeric localization of Mif2, another essential
inner kinetochore protein, depends on Scm3 and Cse4 (Xiao et al., 2017). We constructed
NDC10-AID and *MIF2-AID* strains and used them for our assay to represent the proteins
independent or dependent on Scm3 for centromeric localization, respectively. We also
included one non-essential kinetochore protein, Ctf19, a protein of the COMA complex, to
remove any possible mis-judgement in distinguishing cell-growth-arrest phenotype occurring
due to drug-sensitivity vs. auxin-mediated degradation of essential proteins. The COMA
complex is directly recruited to the centromeres through interaction with the N terminal tail of
Cse4, hence dependent on Scm3 (Chen et al., 2000; Fischböck-Halwachs et al., 2019). Mid-
log grown cells were harvested and spotted on the indicated plates, however, we did not
observe any increased sensitivity of such cells to MMS (Figure S3). In fact, Mif2 depleted cells

showed a slight resistance to MMS due to unknown reasons. Therefore, the increased
sensitivity to MMS in *scm3* mutant but not in other kinetochore mutant cells indicates that
Scm3 possesses an additional function in genome stability besides its role in kinetochore
assembly.

Since HJURP is involved in DDR through the HR pathway (Kato et al., 2007) and If
Scm3 mimics HJURP to repair damaged DNA, then the *scm3* mutant is expected to interact
genetically with the HR mutants. We tested that possibility by analyzing the genetic interaction
between *SCM3* and *RAD52,* as Rad52 is essential for efficient homologous recombination in
S. cerevisiae (Barlow & Rothstein, 2010; Lee et al., 2003). We deleted *RAD52* in the *SCM3-*
AID strain, and the mid-log grown cells were spotted on plates supplemented with increasing
concentrations of auxin (Figure 1C). Interestingly, the double mutant grew slowly as compared
to the single mutants and the wild-type strain. To further validate the genetic interaction
between the two genes, the mid-log grown cells were pre-treated with a 0.75 mM auxin to
deplete Scm3 for the indicated duration before the cells were spotted on YPD plates. Again,
we observed that the double mutant grew slower than the wild-type and single mutants (Figure
1D). Next, we wished to investigate how the double mutant behaves when challenged with
DNA damaging agent. As expected, when the double mutant was spotted in the presence of
0.005% MMS, all the single mutants grew better than the double mutant (Figure S2E). These
results indicate a genetic interaction between *SCM3* and *RAD52,* which becomes pronounced
in the presence of DNA-damaging agents*.*

Further, we wanted to identify the regions in Scm3 whose loss confers sensitivity to
DNA damage. The domain organization of Scm3 reveals two essential conserved regions,
namely NES (nuclear export signal) and HR (heptad repeat), having functions in DNA and
Cse4 binding, respectively (Figure S1A). Scm3 has a non-essential D/E-rich region, which is
important for protein stability but not involved in Cse4 binding (Stoler et al., 2007). Scm3 also
possesses two BR (bromodomain) regions, also referred to as the NLS (nuclear localization
signal). The deletion of the C-terminal 25 amino acids or the bromodomain regions does not
cause cell lethality (Shivaraju et al., 2011; Stoler et al., 2007). We used the versions of Scm3
mutants used elsewhere (Shivaraju et al., 2011) harbouring deletion in one or the other non-
essential region as the sole source of Scm3 and challenged these mutants (two independent
transformants) to MMS and CPT along with wild-type Scm3 as a control. We found no growth
defect in response to MMS or CPT (Figure S1B) suggesting that at least the nonessential
regions are not important for the DDR function of Scm3. Although this indicates that the
essential NES and/or HR might be involved but that is technically difficult to test using growth
based assays.

From these results, we conclude that the loss of Scm3 confers sensitivity to DNA
damaging agents, indicating that Scm3, much like its mammalian homolog HJURP, may play
a role in the DDR pathway.

The loss of Scm3 generates more Rad52 foci

Having shown that the *scm3* mutant is sensitive to DNA damage, we hypothesized that the
mutant cells might be deficient in repairing intrinsic and extrinsic DNA damages. To examine
this, we wished to visualize the Rad52-GFP foci in these cells with or without MMS-induced
DNA damage. Since in budding yeast, the primary pathway to repair DNA damage is through
homologous recombination (HR) mediated by Rad52, deficiency in repair will cause
accumulation of DNA lesions and hence Rad52 (Dornfeld and Livingston, 1991; Lisby et al.,
2003, 2001). The mid-log grown *RAD52-GFP SCM3-AID* cells were split into auxin- and
DMSO-treated halves. Following 2 hrs of treatment, the cells were again split and were mock-
treated or treated with 0.02% MMS for 90 mins (Figure 2A). We visualized the formation of
Rad52-GFP foci in these cells (Figure 2B). Interestingly, we observed that 18% of the Scm3
depleted (+auxin) cells showed Rad52-GFP foci in contrast to only 5% in the wild type (Figure
2C, untreated). In accord with this, when these cells were challenged with MMS, a significant
increase of cells with Rad52-GFP foci was observed in the mutant (70%) compared to the wild
type (45%) (Figure 2C, 0.02% MMS). Moreover, while the mutant cells harbored an average
of ~ 3 foci, the wild type showed ~2 foci per cell (Figure 2D) in response to MMS treatment.
These results suggest that both intrinsically and extrinsically generated DNA lesions remain
unrepaired in the absence of Scm3.

To understand the dynamics of Rad52 foci formation in *scm3* cells, we used a strategy
as shown in Figure S4A. Briefly, the *bar1Δ RAD52-GFP SCM3-AID* cells were arrested in the
G1 phase of the cell cycle by adding alpha factor. 1 hour before release from the G1 phase,
cells were treated with auxin to deplete Scm3 or with DMSO control. Subsequently, the cells
were washed several times and released into alpha factor free media supplemented with or
without MMS but in the presence of nocodazole in order to have all the cells eventually
arrested at metaphase. Following release, the cells were harvested every 30 mins and the
percentage of cells containing Rad52-GFP foci were visualized (Figure S4B). Scm3 depleted
cells, but not the wild type, with or without MMS, showed a peak of Rad52-GFP accumulation
at around 60 mins from G1 release (Figure S4C, D). Notably, this time point possibly coincides
with the timing of DNA replication (judged by bud morphology) when DNA is most susceptible
to intrinsic and extrinsic damages. Taken together, Scm3-depleted cells exhibit more Rad52
foci, indicating a compromised DDR pathway in these cells.

If the DNA lesions remain unrepaired, the cells might undergo an apoptotic pathway
leading to cell death. To determine the percentage of cell death in the *scm3* mutant culture,
the cells harvested at 90 mins of MMS or mock treatment (after auxin treatment), as shown in
Figure 2A were stained with propidium iodide (PI), which does not permeate the live cells and
stains only the dead cells (Figure S4E, arrow). We observed that while ~15% of the Scm3
depleted (+auxin, -MMS) cells were PI positive, it was only ~4% for the wild type (DMSO, -
MMS) (Figure S4E, untreated). The PI positive cell population increased after MMS treatment
in both the cultures with significant difference between them (35% vs 27%) (Figure S4E, 0.02%
MMS). These results suggest that unrepaired intrinsic and extrinsic DNA lesions in the
absence of *scm3* lead to compromised cell viability.

Scm3 associates with non-centromeric sites in response to DNA damage
256

Having observed that Scm3 depleted cells are compromised in DNA damage repair and *scm3*
interacts genetically with *rad52Δ*, it is reasonable to argue that Scm3 goes to the DNA damage
sites to promote repair. To examine this, we first wanted to see if Scm3 alters its localization
pattern in response to MMS-mediated DNA damage. Since Scm3 is also expected to localize
at the kinetochores (Mizuguchi et al., 2007), we visualized Scm3 along with the kinetochore
marker, Ndc10, on chromatin spreads. The cells harboring *NDC10-6HA SCM3-13MYC* were
mock-treated or treated with 0.02% MMS for 90 mins. Ndc10 showed bright foci representing
the kinetochore cluster, whereas Scm3 was localized to multiple locations across chromatin,
including its colocalization with Ndc10 (Figure 3A, arrow). Similar localization of Scm3 has
been observed earlier in live cells where Scm3-GFP showed a wider nuclear localization,
which might argue for its genome-wide functions (Luconi et al., 2011; Wisniewski et al., 2014).
We also observed a significant enrichment of Scm3 at the rDNA loops irrespective of the
extrinsic DNA damage (Figure 3A, arrowhead). The enrichment of Scm3 at the rDNA loops is
perhaps to maintain the highly repetitive organization of rDNA. HJURP has also been shown
to localize rDNA arrays in mammalian systems (Kato et al., 2007). Additionally, upon
quantifying the total intensity of Scm3-13Myc and Ndc10-6HA and normalizing that with the
background intensity, we observed a statistically significant increase in the Scm3-13Myc, but
not Ndc10-6HA intensity in response to MMS treatment (Figure 3B). No change in the intensity
of Ndc10-6HA post-MMS treatment justifies the absence of increased sensitivity of Ndc10
depleted cells in response to MMS (Figure S3A).
Having observed an increased chromatin localization of Scm3 in response to MMS on
chromatin spreads, we wished to verify this by determining genome wide association of Scm3
using ChIP-Seq. To negate any effect on Scm3's binding due to cell cycle progression we first
arrested the cells harboring *SCM3-13MYC* at metaphase by nocodazole. The arrested cells

were further mock treated or treated with 0.05% MMS for an additional 90 mins before they
were harvested for ChIP. The Scm3 ChIP-seq signal was normalized with the input signal,
and genome-wide plots were generated to see Scm3 binding along the entire chromosomes.
We also performed ChIP-seq from a no-tag control strain that provided the background signal
due to the antibody or the beads treatment. We could observe an enrichment of Scm3 in both
MMS treated and untreated cells at the centromeres, while no enrichment in the no-tag sample
was observed (Figures 3C, 3D, S5). In the untreated sample, the peaks called by MACS2 after
input normalization showed enrichment of Scm3 at a few other genomic locations as well
besides the centromeres (Figures 3C, 3D, S5, green arrows), which was also observed in our
chromatin spread data (Figures 3A, 3B). Interestingly, in MMS treated cells, following same
methodology, we could observe a greater number of peaks of Scm3 binding to the non-
centromeric locations (Figures 3C, 3D, S5, black arrows). While in the untreated sample,
Scm3 localized to 1-2 non-centromeric locations, the same increased drastically in the treated
sample. Such non-centromeric chromatin enrichment of Scm3 was observed for several
chromosomes (S5, black arrows) and corresponds to the increased chromatin localization of
Scm3 in response to MMS, as visualized on chromatin spreads (Figure 3B). Further, we could
observe the localization of Scm3 at the 9.1 kb rDNA region on ChrXII, validating our chromatin
spread data (Figure 3E).
299
Scm3 partially co-localizes with Rad52, perhaps at the DNA damage sites
301
Given the increased chromatin association of Scm3 in response to DNA damage we wished
to investigate if this occurs due to its binding to the DNA damage sites. For this, we examined
the colocalization of Scm3 with Rad52 upon DNA damage, as the latter is known to be targeted
to the damage sites (Lisby et al., 2003; Miyazaki et al., 2004). To visualize both Rad52 and
Scm3 simultaneously on chromatin spreads, cells harboring *RAD52-6HA* and *SCM3-13MYC*
were mock-treated or treated with 0.02% MMS for 90 mins. We did not observe any significant
co-localization between Scm3-13Myc and Rad52-6HA in untreated cells as judged by low
Pearson's Correlation Coefficient value (PCC < 0.4) (Figure 4A-B). However, in the cells
treated with MMS, we could observe a significant increase in the PCC value (PCC > 0.5)
(Figure 4B). This indicates a partial co-localization between Scm3 and Rad52 in the presence
of the DNA damage and implicates that Scm3 may localize to the damage sites in response
to DNA damage.
Next, we wished to examine the behavior of Scm3 with respect to another DNA
damage marker. In mammals, the DSB sites are marked by γ-H2Ax due to phosphorylation of
the H2A variant, H2Ax, by ATM/ATR kinases (Burma et al., 2001; Lee et al., 2014) In yeast
that lacks H2Ax, the DSBs are marked by γ-H2A due to phosphorylation of histone H2A by

Mec1 and Tel1 kinases (Lee et al., 2014). To be noted, γ-H2A is also found at the

heterochromatin region and co-localizes with yeast heterochromatin protein Sir3 (Javaheri et

al., 2006; Kirkland et al., 2015; Kitada et al., 2011). We performed a chromatin spread assay

using antibodies specific to γ-H2A and observed a drastic increase in the intensity of γ-H2A

when the cells were subjected to DNA damage, which is expected (Figure S6 A, B). To verify

that the phenomenon is DNA damage associated, we observed a significant increase in the

colocalization between Rad52 and γ-H2A in response to DNA damage (Figure S6 C-D). To

further correlate the localization of Scm3 with DNA damage, we then visualized both Scm3-

13Myc and γ-H2A simultaneously on chromatin spreads from *SCM3-13MYC* cells mock-

treated or treated with 0.02% MMS for 90 mins before they were harvested for the assay. We

did not observe any significant co-localization between Scm3-13Myc and γ-H2A in untreated

cells as judged by low Pearson's Correlation Coefficient value (PCC < 0.3) but found a

significant increase in the value (PCC > 0.6) in MMS treated cells (Figure 4C-D). Taken

together, we conclude that the association of Scm3 with chromatin and its co-localization with

DDR proteins increases with DNA damage.

Scm3 is recruited to a site-specific DSB

The overall increased association of Scm3 with the chromatin and its colocalization with

Rad52 and γ-H2A in response to MMS-mediated DNA damage prompted us to examine if

Scm3 is recruited specifically to DNA damage sites. For this, we utilized the chromatin

immunoprecipitation (ChIP) assay to visualize the dynamics of Scm3 recruitment at an

induced DSB site. We used a yeast strain (NA14) harboring an HO cut site integrated at the

mutant *ura3* locus, which is cleavable by HO endonuclease upon galactose induction (Agmon

et al., 2009; Fangaria et al., 2022). The strain also contains a distant *URA3* locus, which is

utilized to repair the cut *ura3* locus post-DSB induction. The cleavage by HO endonuclease

was first verified by PCR-based analysis using primers complementary to the DNA sequence

flanking the HO cut site (Figure 5A). The DNA damage was maximal at 2 hrs of galactose

induction, and the damage was majorly repaired by 4 hrs of galactose induction (Figure S7 A,

B).

We monitored the recruitment of Scm3 near DSB and at sites distal to that up to -3 kb

(Figure 5A) by performing ChIP of Scm3-6HA from the samples collected at every hr up to 4

hrs following galactose induction. As a positive control for Scm3 ChIP, we found significant

enrichment of Scm3 at the centromere (*CEN3)* (Figure S7C). We observed a significant

increase in the Scm3 binding near DSB at 2 hrs of galactose induction and a decrease in the

enrichment at 4 hrs, possibly due to the gradual repair of the damage (Figure 5B, C). We also

observed an increase in the enrichment of Scm3 up to -3 kb sites distal to the cut site, which

decreased gradually with damage repair (Figure 5C) and no significant enrichment at the
TUB2 locus, used as negative control (Figure 5C) or in the no antibody control (Figure 5D).
This suggests that Scm3 is maximally recruited to the cleavage sites at the time of DNA
damage and gradually dissociates as and when the DNA is repaired. To verify that HO-
induced DSB is able to recruit the DNA repair proteins, we also investigated the enrichment
of such a protein, Rad51, that is known to bind the DSB sites (Fangaria et al., 2022) and found
it is maximally enriched at these sites during 2-3 hrs of post-galactose induction (Figure S7D).
We conclude that upon DNA damage Scm3 associates with the damage sites and may act in
cis to facilitate damage repair.
363
364   **Activation of the DDR checkpoint is perturbed in the *scm3* mutant.**
365
Having shown the recruitment of Scm3 at the DNA damage sites, we wished to address the
role of Scm3 at these sites in case of DNA damage. Since we found that Scm3 depleted cells
are compromised in repairing DNA damage, it is possible that being at the damage sites Scm3
activates the DDR checkpoint, which is required for the damage repair. In the DDR pathway,
Rad53 is an effector kinase that becomes phosphorylated in a Mec1 sensor kinase-dependent
manner in response to DNA damage (Ciccia and Elledge, 2010; Melo and Toczyski, 2002;
Zhou and Elledge, 2000). Phosphorylation of Rad53 is essential to halt the cell cycle and
provide cells time to repair the damaged DNA. To examine whether Scm3 depleted cells have
defects in the activation of DDR checkpoint and in halting the cell cycle, we measured Rad53
phosphorylation status in the wild type and the Scm3 depleted cells after treating them with
0.02% MMS for 90 mins (Figure 6A). Anti-Rad53 antibodies that recognize both
phosphorylated and unphosphorylated forms of Rad53 were used. We observed lesser
phosphorylation of Rad53 in the *scm3* than in the wild-type cells treated with MMS, but no
difference was found in the untreated cells (Figure 6B).
Since Scm3-depleted cells, unlike the wild type, arrest at the G2/M stage due to
activation of spindle assembly checkpoint (SAC) (Camahort et al., 2007; Stoler et al., 2007),
the observed difference in Rad53 phosphorylation could be due to the difference in cell cycle
stages. To circumvent this issue, we grew *CDC20-AID* and *SCM3-AID* cells for 3 hrs before
treating them with 0.75 mM auxin for 2 hrs to arrest them first at G2/M by depleting Cdc20 and
Scm3, respectively (Figure 6C). The cells were then treated with 0.02% MMS and harvested
at indicated time points for total protein extraction. The *CDC20-AID* and *SCM3-AID* cultures
showed around ~80% metaphase arrest cells, in 2 hrs of auxin treatment (0 mins in Figure
S8A). Notably, while the Cdc20-depleted cells maintained the arrest even with a longer
incubation in auxin, the Scm3-depleted cells tend to bypass the arrest after a certain time and
showed accumulation of multibudded cells with fragmented DAPI, an arrest-release

phenotype (Figure S8B, Figure 6E) Consistent with our previous results (Figure 6C), we

observed that in Scm3 depleted cells, Rad53 is less phosphorylated than in Cdc20-depleted

cells (Figure 6D), in particular, while comparing the bands at 120 mins when the percentage

of metaphase cells was found similar in two cultures (Figure S8B). These results indicate that

in the absence of Scm3, when there is DNA damage, the activation of the DDR checkpoint is

improper, and consequently, the phosphorylation of Rad53 is compromised.

If the loss of Scm3 fails to activate the damage checkpoint properly, then it is expected

that the cell cycle arrest phenotype in response to DNA damage would also be defective and

would produce multibudded cells. We wished to examine this possibility. However, Scm3-

depleted cells arrest by spindle assembly checkpoint (SAC) at G2/M due to improper

kinetochore-microtubule attachment because of failure in Cse4 deposition (Camahort et al.,

2007; Stoler et al., 2007). To exclude the effect of SAC-mediated arrest, we deleted the SAC

component *MAD2* in *SCM3-AID* cells. These cells were first depleted of Scm3 by treatment

with auxin and then, along with *mad2Δ* control cells, were challenged with or without MMS for

90 mins before they were analyzed for percentage of multi budded cells (Figure 6E). While

the *mad2Δ* cells in the presence of MMS showed nominal multi budded cells (<5%), the same

cells upon depletion of Scm3 showed a steep rise in the phenotype (>40%), suggesting that

Scm3 depleted cells fail to activate DNA damage checkpoint and to arrest the cell cycle in the

presence of DNA damage.

Since Scm3 depleted cells showed reduced Rad53 phosphorylation, we wished to

examine if the lack of Scm3 perturbs Mec1 kinase function that phosphorylates Rad53. Mec1

phosphorylates histone H2A at S129 position upon DNA damage (Lee et al., 2014). However,

we failed to observe any significant difference in the phosphorylation status of H2AS129

between Scm3, or Cdc20 depleted cells (Figure S8C), indicating that the Mec1 function is not

perturbed in the absence of Scm3.

Phosphorylation of Scm3 in response to DNA damage

The function of Scm3 in DDR could be mediated by a damage-induced post-translational

modification of this protein. To examine if Scm3 becomes modified upon DNA damage, we

treated *SCM3-6HA* cells with increasing concentrations of MMS or with 0.02% MMS with

increasing duration. We observed slower migrating bands of Scm3 (Scm3*) that became

brighter with increasing concentrations of MMS and present very faintly even in the untreated

cells (Figure 7A, left panel). We observed an increase in the normalized intensity of Scm3*

and Scm3*/Scm3 ratio with increasing MMS concentration (Figure 7A, middle, and right

panels). A similar observation was also made when the cells were treated with 0.02% MMS

for increasing time (Figure S9A). Since we could detect Scm3* bands in the untreated cells as

well, its enrichment in response to MMS could be due to a cell cycle arrest caused by MMS.
To understand the dynamics of the appearance of Scm3* bands over different cell cycle stages
and in response to MMS, we released the *bar1Δ SCM3-6HA* cells from alpha-factor mediated
G1 arrest into the medium in the absence or presence of MMS, as shown in Figure 7B. We
harvested the cells at indicated time points and evaluated the cell cycle synchrony using bud
morphology and DAPI staining (Figure 7C, Figure S9 B, C), which depicts that the majority of
the cells at 0, 30, 60, 90, and at 120 mins are in G1, S, metaphase, metaphase/anaphase and
anaphase, respectively. We could faintly observe the Scm3* bands after release from the G1
phase in MMS-free (untreated) media, and the band intensity did not change over the time
period. However, when the cells were released in the presence of MMS, we observed an
increase in the intensity of the Scm3* bands over time (Figure 7D). These results suggest that
the accumulation of Scm3* in the MMS treated cells is due to DNA damage rather than an
effect of the pace of the cell cycle. It is possible that Scm3* bands are phosphorylated forms
of Scm3 since the protein has several serine/threonine residues. To test this, we challenged
the Scm3-6HA cells with MMS, as in Figure 7A, and resolved the proteins on SDS-PAGE with
or without Phos-Tag reagent. The presence of Phos-Tag in the gel enables slow mobility of
the phosphorylated proteins and thus further differentiates them from their unphosphorylated
forms. Figures S9D and E show the migration of the proteins isolated from *SCM3-6HA* cells
challenged with different concentrations of MMS in the absence or presence of Phos-Tag
reagent, respectively. The mobility of the Scm3* bands was further reduced with respect to
the Scm3 band in Phos-Tag gel compared to the normal gel, suggesting that the modification
is likely due to phosphorylation. Notably, in several high throughput studies, Scm3 has been
shown to be phosphorylated at S39, S64, and S165 residues (Lanz et al., 2021; MacGilvray
et al., 2020; Zhou et al., 2021). The S64 phosphorylation of Scm3 was found to be enriched
in cells after MMS treatment for 2 hrs as compared to G1 arrested cells, although the authors
could find S64 phosphorylation in only 2 out of 4 replicates used in their study (Lanz et al
2021). However, in the absence of further validation in those studies, our observation of a
gradual increase in the Scm3* band intensity with increasing dose of MMS convincingly
demonstrate DNA damage dependent phosphorylation of Scm3.
Since, Mec1 kinase majorly contributes to the activation of DDR through protein
phosphorylation, we tested if Scm3 undergoes a Mec1-dependent phosphorylation in
response to MMS. We constructed the *sml1Δ mec1Δ SCM3-6HA* strain as *mec1Δ* cells are
viable in the absence of Sml1 (Zhao et al., 2001, 1998) and challenged this and other control
strains with MMS. Briefly, wild type, *SCM3-6HA*, *sml1ΔSCM3-6HA*, and *sml1Δ mec1Δ SCM3-*
6HA cells were grown till mid-log and spotted on the plates containing MMS (Figure S9F). As
expected, the *sml1Δ mec1Δ SCM3-6HA* cells showed extreme sensitivity to MMS. However,
we could not observe any loss of Scm3 phosphorylation in the *sml1Δ mec1Δ* background in

the presence of MMS, suggesting that Scm3 phosphorylation in response to DNA damage is

not dependent on Mec1 kinase activity (Figure S9G).

Discussion

The bonafide function of the Cse4 chaperone, Scm3, is to deposit Cse4 at the centromeres

during the S phase in budding yeast to facilitate kinetochore formation. The observation that

Scm3 persists not only at the centromeres throughout the cell cycle but also associates with

the gross chromatin argues for additional kinetochore-independent functions of Scm3 in

maintaining genome stability. While in metazoan, Scm3 homolog HJURP was initially

identified as a protein involved in DNA damage response (DDR), how HJURP regulates the

DDR pathway besides its role in CENP-A deposition is not well understood. The DDR role of

Scm3 has never been reported in budding yeast. Whether Scm3 also contributes to DDR

pathway in budding yeast is important to address since this organism is distinct from

metazoans in terms of harboring small 125 bp point centromeres contrasting megabase pair

large regional centromeres in metazoans and having lesser epigenetic control on biological

functions, including DNA damage repair. Here, for the first time, we report that Scm3, perhaps

upon phosphorylation, promotes DNA damage repair by contributing to the activation of the

DNA damage checkpoint, and in the process, it associates with the DDR proteins at the DNA

damage sites (Figure 8). Our findings have a significant impact on advancing knowledge of

how CENP-A chaperones can be involved in the DDR pathway.

Lack of Scm3 increases DNA damage sensitivity

It is technically challenging to perform a growth-based sensitivity assay using cells depleted

for an essential gene. To circumvent this issue with Scm3, which is essential for cell survival,

we used partial depletion of Scm3 to assess the sensitivity of the partially depleted cells to the

genotoxic agents, including MMS, HU, and CPT and observed that such cells were sensitive

to those agents, (Figure 1, Figure S2A). However, since the primary function of Scm3 is to

promote kinetochore formation by depositing Cse4 at the centromeres, it is important to

address if the observed sensitivity is due to perturbation in kinetochores. Therefore, we

similarly partially depleted two essential kinetochore proteins, Ndc10 and Mif2, and deleted

one non-essential kinetochore protein, Ctf19, in separate cells but failed to detect any

increased sensitivity to DNA damage stress (Figure S3). These results indicate that the drug

sensitivity phenotype of Scm3 depleted cells is not due to weakly formed kinetochores. The

arguments favoring the possible centromere-independent DDR functions of Scm3 can be

gleaned as follows. From comparing the domain organization of Scm3 with HJURP (Figure

S1A), we noticed that besides the CENP-A/Cse4 binding domain, both HJURP and Scm3

have a common DNA binding domain, which is dispensable for CENP-A/Cse4 binding and

also for Ndc10 binding in budding yeast. Both in vitro and in vivo evidence are there demonstrating binding of the DNA binding domain to the DNA (Kato et al., 2007; Mizuguchi et al., 2007), suggesting this domain can indeed recruit Scm3 also at the non-centromeric DNA damage sites. In support of this, Scm3 has recently been shown to be phosphorylated in an MMS-dependent manner at S64 (Lanz et al., 2021), which, interestingly, lies within the DNA binding domain. Among other non-essential regions of Scm3, neither the NLS region nor the C-terminal region is involved in DDR (Figure S1B). On the other hand, the D/E-rich region, which is important for Scm3 stability (Stoler et al., 2007), forms a disordered domain as per alpha-fold prediction and hence is believed to be capable of interacting with multiple proteins or can act as a scaffold for engaging several protein-protein interactions which is required in DDR.

MMS halts replication forks due to the formation of DNA adducts, which eventually generate recombinogenic lesions (Conde and San-Segundo, 2008). In yeast, these lesions are repaired primarily through homologous recombination (HR), facilitated by the Rad52 epistasis group of proteins (Pizzul et al., 2022; Sirbu and Cortez, 2013). Rad52, the central component of the HR pathway, deposits Rad51 onto the broken DNA strand, which then searches for the homologous template and undergoes strand exchange to repair the broken DNA (Haber, 2018). In the presence of any DNA damage, Rad52 relocalizes to form bright foci, which mark the repair centres. Strikingly, we observed that the Rad52 foci are increased drastically in the absence of Scm3, even without any external DNA damage (Figure 2C, Left). The maximal accumulation of Rad52-GFP foci was noticed during the 60 mins post-G1 release, which coincides with the timing of DNA replication (Figure S4C), indicating that Scm3 might be required to avoid generating DNA lesions during replication, which itself is genotoxic. In support of this, previously, it was observed that a significant cell population arrests during the S phase in the absence of Scm3 (Camahort et al., 2007). As expected, the accumulation of Rad52-GFP foci was significantly higher in the *scm3* mutant as compared to the wild-type cells when they were treated with MMS, suggesting that cells become more vulnerable to DNA damage in the absence of Scm3 (Figure 2C, Right).

Association of Scm3 with non-centromeric loci

In addition to the localization of Scm3 at the kinetochores, we also found it at other loci, including at the ribosomal DNA (rDNA) loops (Figure 3A, arrowhead). In budding yeast, chromosome XII, harbors rDNA with around 150 repetitive loci (Dammann et al., 1993; Smith, 2022). Recombination among these repeats excise rDNA circles, and therefore, efficient DNA damage repair through HR at these loci is crucial for preserving genomic stability. Notably, HJURP has been identified in mammals to bind to the rDNA region and maintain its stability

(Kato et al., 2007). The degradation of HJURP using siRNA resulted in increased rDNA
recombination events and chromosome instability (Kato et al., 2007). The presence of Scm3
at the rDNA loops suggests that Scm3, like HJURP, is also important in regulating the
recombination and repair processes at those loci (Figure 3D).

Both Rad51 and Rad52 have been shown to mark the site of recombination and repair
(Lee et al., 2003; Lisby et al., 2001; Miyazaki et al., 2004). In our assay, we observed a
statistically significant co-localization of Scm3 with Rad52 in response to DNA damage (Figure
4A). Since during DSB repair, multiple lesions are recruited to a few repair centers, which are
marked by Rad51 and Rad52, the colocalization of these proteins with Scm3 implicates the
recruitment of Scm3 at the repair centers. Moreover, an overall increase in the chromatin
association of Scm3 in response to MMS also suggests that Scm3 is recruited to several repair
centers (Figure 3B, C, E, S5). However, as expected, no such increase in the chromatin
association was observed for an inner kinetochore protein, Ndc10. It is possible that Scm3
might physically interact with Rad51/Rad52 through the HR or DNA binding domain or the D/E
rich region present in the protein (Figure S1A). However, we failed to observe any interaction
of Scm3 with Rad52 using co-IP assay.

DNA damage sites are epigenetically marked by phosphorylation of H2AS129 (γ-H2A)
through the ATM/ATR kinase pathway (Burma et al., 2001; Lee et al., 2014). However, in
budding yeast, γ-H2A is also recruited at the heterochromatin regions (Kitada et al., 2011). In
unperturbed cells, we could visualize γ-H2A foci (Figure 4C), possibly due to its presence at
the stalled replication forks, fragile sites, telomeric regions, or at the heterochromatin regions
(Kirkland et al., 2015; Kitada et al., 2011; Rg Bewersdorf et al., 2006). The increase in intensity
of γ-H2A upon MMS treatment (Figure S6A, B) indicates the presence of γ-H2A at multiple
DNA damage sites. The significantly high co-localization of Scm3 (PCC > 0.6) with γ-H2A in
the MMS treated cells, therefore, indicates that Scm3 is indeed recruited to the DNA damage
sites (Figure 4D). This was further verified by using an ectopic DSB system, where we showed
the recruitment of Scm3 to the induced DSB sites (Figure 5B). Notably, Scm3's presence was
found not only limited to the regions very close to the cut site but also up to 3 kb distal from
there. This may be the reason for observing a high co-localization of Scm3 with γ-H2A in
response to MMS, as the latter also spreads distal to the cut site (Lee et al., 2014). Although
Scm3 binding at the cut site peaked at 2 hrs following HO induction, the time of maximal DNA
cleavage, it persisted for at least 3-4 hrs post induction even when ~60% of the cut DNA was
repaired. This suggests that Scm3 is recruited to the sites of DNA damage and persists there
during the repair process.

Scm3 regulates DNA damage checkpoint

We wanted to address the mechanism by which Scm3 might be involved in DDR. The DNA

damage checkpoint arrests the cell cycle in response to DNA damage, allowing cells to repair

the damage and maintain cell viability. The serine/threonine kinases Mec1/Tel1 in yeast are

the sensor kinases that are activated and recruited to the sites of DNA damage. These kinases

relay the signal downstream by phosphorylating and activating the effector kinases, including

Chk1 and Rad53 (Pizzul et al., 2022; Zhou and Elledge, 2000). The activation of the effector

kinases is critical for activating the further downstream substrates that eventually halts the cell

cycle. We observed a defect in the Rad53 phosphorylation in response to MMS-mediated DNA

damage in the absence of Scm3 (Figure 6B). Failure to activate Rad53 in the presence of

damage leads to improper G2/M cell cycle arrest, which can produce the multi-bud phenotype.

In the Scm3 depleted cells deleted for Mad2 to remove the ability of SAC mediated arrest (due

to kinetochore perturbation), we observed a significant population of multibudded cells (Figure

6E) when they were challenged with MMS. Only $mad2\Delta$ cells challenged with MMS did not

show such multi-bud phenotype, suggesting that Scm3 is important to arrest the cells in

response to DNA damage. Next, we wished to know why the Scm3 depleted cells are

compromised to phosphorylate Rad53. It is known that Rad9 acts as an adaptor upon

phosphorylation by upstream Mec1 kinase to phosphorylate Rad53 (Gilbert et al., 2001;

Pellicioli and Foiani, 2005). To test if lack of Scm3 attenuates Mec1 activity, we tested the

level of γ-H2A in the Scm3 depleted cells as Mec1 activity is also responsible to generate γ-

H2A (Burma et al., 2001; Lowndes and Toh, 2005). However, the level of γ-H2A was not

perturbed in the *scm3* mutant, suggesting that Mec1 activity is intact (Figure S8C). Therefore,

currently, we are uncertain about the cause of improper Rad53 phosphorylation; a defect in

its auto-activation property or in the action of some other mediator kinases, e.g., Tel1, may be

responsible, which can be tested. Nevertheless, given the increased phosphorylation of Scm3

in response to DNA damage, it is possible that the p-Scm3 is required to activate Rad53

(Figure 7A). Although we found that Scm3 phosphorylation is Mec1 independent (Figure S9G),

it is possible that other kinases are involved. While the involvement of CDKs in the DNA

damage checkpoint is not well understood, it is known that CDKs contribute to the

phosphorylation of both Rad52 and Rad51 to promote repair (Lim et al., 2020). Given that the

centromeric association of Scm3 varies with the cell cycle stage (Mishra et al., 2011), probably

due to oscillating levels of CDK activity, the DNA damage-dependent phosphorylation of Scm3

might also be regulated by CDKs. However, this does not rule out the involvement of other

kinases including Rad53 and Tel1, which are closely associated with the DNA damage

checkpoint pathway. Further investigation is required to precisely determine the kinase

involved in the Scm3 phosphorylation.

In summary, we report here that, like HJURP, Scm3 in budding yeast also possesses

functions in DDR, indicating that the centromere independent functions of the CenH3

chaperone are evolutionarily conserved. Although the chromatin landscape of budding yeast

is far more 'open' with lesser involvement of epigenetic determinants than in metazoans, the

inherent similarity in the DNA damage repair pathway may necessitate the utilization of these

chaperones in a similar fashion in both cell types. Not much has been studied about how

HJURP is involved in the DDR pathway, and as we reveal that Scm3 is involved in the proper

activation of the DNA damage checkpoint in the presence of DNA lesions (Figure 8), the same

can be tested for HJURP in metazoans as abrogation of such surveillance function in humans

can lead to aneuploidy and disease states. Our findings on Scm3 will pave the way for future

work using yeast as a model to decipher more genome wide functions of the CenH3

chaperone, which is crucial given its genome wide localization and evidence of its modulation

in several patients.

Acknowledgement:

We acknowledge Prof. Martin Kupeic from Tel Aviv University, Israel, for providing us with the

NA14 strain. SKG is supported by a grant (BT/PR43050/BRB/10/1992/2021) from the

Department of Biotechnology (DBT), Govt of India. PA is supported by CSIR fellowship

(09/087(0972)/2019- EMR-I).

Author contributions: PA performed experiments. PA and SKG designed experiments,

analyzed and interpreted the data. PA and SKG wrote the manuscript.

Materials and Methods:
630
631   **Yeast strains**
632
All *Saccharomyces cerevisiae* strains used in this study are derived from W303 genetic

background and are mentioned in Table S1. Standard growth conditions were used to grow

and harvest yeast for all the experiments. C-terminal tagging (AID, HA, Myc, GFP) or gene

deletions were performed using PCR-based integration at endogenous locations as described

before (Janke et al., 2004; Longtine et al., 1998; Wach, 1996). Standard lithium acetate-based

yeast transformation methods were employed for all genetic manipulations (Gietz and Woods,

2002).

Media, reagents and growth conditions

Yeast strains were grown using standard growth conditions (Sherman, 2002). For plates

containing MMS, HU and CPT (Sigma Chemicals, Switzerland), the drugs were top-plated on

YPD plates at appropriate concentrations. To arrest the cells at the G1 or metaphase stage of the cell cycle, alpha factor (Sigma Chemicals, Switzerland) or nocodazole (Sigma Chemicals, Switzerland) was added to the cells at indicated concentrations. For HO endonuclease induction, strains were grown on a rich medium containing 3% glycerol prior to the addition of 3% galactose. For western blotting, immunoprecipitation, or immunofluorescence studies the primary antibodies used were rat anti-HA (3F10, Roche), mouse anti-HA (12CA5, Roche), mouse anti-Myc (9E10, Roche), rat anti-tubulin (MCA78G, Serotec), rabbit anti-Rad53 (ab104232, Abcam), rabbit anti-Rad51 (PA5-34905, Invitrogen) and rabbit anti-Histone H2A (phospho-S129) (ab15083, Abcam). The secondary antibodies used were Rhodamine (TRITC)-conjugated goat anti-Rat (Jackson), Alexa Fluor 488 conjugated goat anti-Mouse IgG (Jackson), Rhodamine (TRITC)-conjugated goat anti-Rabbit, HRP-conjugated goat anti-mouse IgG (Jackson), HRP-conjugated goat anti-rat (Jackson), HRP- conjugated goat anti-rabbit (Jackson) at appropriate dilutions.

Cell spotting and viability assay

The cells were grown in standard media conditions till the mid-log phase. The cultures were then 10-fold serially diluted and spotted on the plates containing auxin (dissolved in DMSO) or other DNA-damaging agents. All the reagents were top-plated onto YPD or specific dropout plates. After spotting, the plates were incubated at 30°C for 2-3 days before they were photographed, depending on the growth rate.

For treatment of the cells in liquid culture, mid-log grown cells were treated with auxin (dissolved in DMSO) or MMS for the indicated time points before they were harvested and spotted on YPD plates or plates containing DNA damaging agents. To measure cell viability, an equal number of cells were plated on drug free or drug containing YPD plates. The cells were grown at 30°C for 2 days, and the number of viable colonies was counted. The colonies formed on the drug free plate were taken as 100% viability, and accordingly the viability percentage was calculated and plotted for drug containing plates.

G1 arrest and release

Cells were arrested in the G1 phase by adding 0.5 µg/mL α-factor to early log phase cells in liquid culture (Amon, 2002). The cells were incubated for 2-3 hours or until at least more than 90% of cells showed shmoo morphology. For the release of the cells from α-factor arrest, the cells were harvested and washed twice with pre-warmed water at 30°C which was followed by washing twice with pre-warmed α-factor free media at 30°C with at least 10 times the

680   volume of the initial culture. The cells were then released in α-factor free media, and samples
681   were collected at different time points.

683
684   **Fluorescence imaging**
685

For live cell imaging, the samples were harvested at indicated time points and washed twice
in 0.1 M phosphate buffer before imaging. The cells were processed for DAPI staining, as
mentioned elsewhere (Mittal et al., 2020; Shah et al., 2023). Typically, the cells were fixed with
4% formaldehyde at RT for 5 mins. The fixed cells were washed twice with 0.1M phosphate
buffer (pH 7.5), vortexed in freshly prepared 50% ethanol for 30 seconds, and rewashed with
0.1M phosphate buffer. Before imaging, the cells were resuspended in freshly prepared DAPI
(1 µg/ml, Invitrogen) solution for 20 mins. The images were acquired through the Zeiss Axio
Observer Z1 fluorescence microscope (63× or 100×) in z-stack mode (0.2-0.5 µm spacing).

695
Chromatin spread

Chromatin spread was performed, as mentioned previously (Ma et al., 2023; Mehta et al.,
2014; Shah et al., 2023). Briefly, mid-log (O.D.$_{600}$ = ~1.0) grown cells were harvested and
washed once with spheroplasting buffer (1.2 M sorbitol in 0.1 M phosphate buffer). The cells
were resuspended in 0.5 ml of the same buffer supplemented with 5 µl of β-mercaptoethanol
and 12.5 µl of zymolyase 20T (10 mg/ml, MP Biomedicals, USA) and incubated at 30ºC for 1
h. Once 80-90% of the cells were spheroplasted (by observing under the microscope), the
spheroplasting was stopped by adding 1 ml of ice-cold stop solution (0.1 M MES- pH 6.4, 1
mM EDTA, 0.5 mM MgCl$_2$, 1 M Sorbitol). The spheroplasts were centrifuged at 2000 rpm for
2 mins and resuspended in 120 µl of ice-cold stop solution. About 60 µl of spheroplasts were
mounted on an acid-washed clean glass slide. The cells were treated with freshly made
paraformaldehyde solution (4% paraformaldehyde, 3.4% sucrose), followed by 1% Lipsol (LIP
Equipment and services) solution for cell lysis. The lysed cells were homogenously spread
over the slide and air-dried at RT. After overnight drying, the slides were treated with 0.4%
Kodak Photoflow-200 to avoid photobleaching, followed by 5% skim milk as a blocking
solution, which was covered with coverslips. The coverslips were removed, and the slides
were incubated with primary and secondary antibodies (1:200) for 60 mins with three times
PBS washing between antibody treatments. The slides were then incubated with 100 µl of
DAPI (1 µg/ml, Invitrogen) for 30 mins. After PBS wash for 5 mins, 100 µl of mounting solution
(90% glycerol supplemented with 1 mg/ml p-phenylenediamine) was added over the slides. A
clean coverslip was placed over a slide and was sealed with transparent nail paint.

Microscopic image analysis

The images acquired through the Zeiss Axio (Observer Z1) fluorescence microscope were processed using Zeiss Zen 3.1 software. The z-stacks with the best signal for a particular channel were extracted and merged separately. For intensity calculation, a Region Of Interest (ROI) was drawn around the signal of interest, and the integrated density was determined to get the 'signal intensity' for both channels. An ROI of the same size was put elsewhere in the background, and integrated density was determined to get the 'background intensity.' Each signal intensity value was normalized with the background intensity values to get the 'normalized signal intensity' which was used to plot the graphs (Shah et al., 2023).

For Pearson's correlation coefficient (PCC) values to determine the extent of co-localization, the merged z-stacks were extracted, and the 'Coloc' tool from the 'Imaris 8.0.2' software was used. The 'automatic thresholding option' in the 'Coloc' tool was used to calculate the threshold for each fluorescence emission channel. The PCC values were recorded separately for each individual spread. The colocalization was described as no (PCC <0.1), partial (PCC = 0.3-0.5) and complete (PCC>0.5) (Ma et al., 2023; Prajapati et al., 2018; Zinchuk and Grossenbacher-Zinchuk, 2014).

Induction of a site-specific DSB

NA14 (Fangaria et al., 2022) cells were grown till 0.3 OD in the presence of 3% glycerol. 5 OD of the cells were harvested (0 h), and the remaining cells were treated with 3% galactose for 4 hrs. 5 OD cells were harvested every hour till 4 hrs. The cells were lysed using glass beads in lysis buffer (2% Triton X-100, 1% SDS, 100 mM NaCl, 10 mM Tris pH 8, and 1 mM EDTA pH 8.0), and total DNA was isolated as described before (Hoffman, 1997) and was solvent extracted using PCI. The genomic DNA was precipitated and finally resuspended in 30 µl 1X TE. The concentration of DNA was measured using a nano spectrophotometer. About 30 ng of DNA was used for PCR to detect the DNA break using primers OSB289 and kanB1 (Fangaria et al., 2022). Primers against *TUB2* ORF were taken as a control.

ChIP assay and qPCR quantification

ChIP assay was performed as mentioned previously (Fangaria et al., 2022; Kumar et al., 2021; Makrantoni et al., 2019; Mehta et al., 2014). First, cells were grown in the presence of 3% glycerol till 0.3 OD. A total of 60 OD cells were harvested (0 h), and the remaining cells were treated with 3% galactose. A total of 60 OD of cells were harvested every hr till 4 hrs. The cells were then fixed with 1% formaldehyde for 30 mins at 25°C at 100 rpm, followed by quenching

by the addition of glycine at a final concentration of 125 mM. The cells were then harvested and washed twice in ice-cold TBS buffer (20 mM Tris–HCl pH 7.5, 150 mM NaCl) and once with 10 ml of FA lysis buffer (50 mM Hepes–KOH pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% Na-deoxycholate) supplemented with 0.1 % SDS. The pellet was resuspended in 500 µl of 1X FA lysis buffer supplemented with 0.5% SDS, 1 mM PMSF, 1X PIC. The cells were lysed using 0.5 mm glass beads using a mini-bead beater (BIOSPEC, India) for 7 cycles (1 min ON, 2 mins OFF on ice). The lysate was collected in a pre-chilled 1.5 ml eppendorf tube and centrifuged twice at 16,000 x g for 15 mins at 4°C. The pellet was resuspended in 300 µl ice-cold 1X FA lysis buffer supplemented with 0.1% SDS, 1 mM PMSF, and 1X PIC. The chromatin was then sheared to 200-600 bps using a water bath sonicator (Diagenode SA, Picoruptor, BC 100,27 LAUDA Germany) for 30 cycles of 30 s ON / 30 s OFF at 4°C. After clarification of the lysate by centrifugation at 16,000 x g for 10 mins at 4°C, 3-5 µg of appropriate antibodies (were added and incubated at 4°C overnight with gentle rotation. Subsequently, 50 µl Protein-A conjugated Sepharose beads were added and incubated for 2 h at 4°C with rotation. The beads were washed twice with IP wash buffer I (1X FA lysis buffer, 0.1%SDS, 275 mM NaCl), followed by one wash in IP wash buffer II (1X FA lysis buffer, 0.1%SDS, 500 mM NaCl, III (10 mM Tris pH 8.0, 250 mM LiCl, 0.5% NP-40, 0.5% sodium deoxycholate, 1 mM EDTA), and 1X TE at room temperature The beads were resuspended in elution buffer and the chromatin was eluted by boiling at 65°C. The eluate was then decrosslinked overnight at 65°C, followed by Proteinase K (SRL, India) treatment for 2 hrs at 45°C. The DNA was purified by PCI-based purification and precipitated overnight in chilled ethanol at -80°C. The enrichment of obtained chromatin fragments was estimated using qPCR (BioRad CFX96, USA) using specific primers targeting specific and non-specific (negative control) chromatin loci, listed in Table S2. As mentioned before (Kumar et al., 2021; Shah et al., 2023) the following equation was used to estimate the percentage of chromatin enrichment. ChIP efficiency = Enrichment/Input X 100; Enrichment/Input = E^-ΔCT; ΔCT = CT(ChIP) − [CT(Input) − LogEX(D)]; E = primer efficiency value, CT = Threshold values obtained from qPCR, D = Input dilution factor. E was estimated as {[10^(-1/slope)]– 1} from standardization graphs of CT values against dilutions of the input DNA.

Chromatin immunoprecipitation (ChIP) and sequencing

ChIP assay was performed as mentioned above with certain modifications. The cells were grown till mid-log and treated with nocodazole (20 ug/ml) for 90 mins to arrest cells at metaphase and were subsequently treated with 0.05% MMS for additional 90 mins. The cells were harvested and processed for immunoprecipitation, as mentioned above except Protein-A conjugated dyna beads were used instead of sepharose beads. 10 ng of DNA eluate was

taken to prepare the NGS library using a TruSeq ChIP sample preparation kit from Illumina.
The library was prepared as per the manufacturer's protocol. Library QC was performed by
measuring the concentration and size of the libraries by using the 4150 Tapestation system
from Agilent and Qubit dsDNA HS assay kit (Thermo Fisher Scientific). The libraries were
sequenced using the Illumina platform in paired-end mode with a read length of 151 bp by
Macrogen Asia Pacific Pte. Ltd, Singapore.

Bioinformatics analysis
The ChIP-Seq analysis was performed by Genotypic (Bengaluru,India). The raw reads were
processed using FastQC (Andrews S., 2010) for quality assessment and pre-processing,
which includes removing the adapter sequences and low quality bases (<q30) using
TrimGalore (Krueger F., 2015). The pre-processed high quality read data was aligned to
reference *Saccharomyces cerevisiae* genome data base using Bowtie, which is an ultrafast,
memory-efficient short read aligner geared towards quickly aligning large sets of short DNA
sequences (reads) to large genomes. The aligned files (BAM files) were used for peak calling
using MACS2 software with input sample as a control. The bedgraph files generated from
MACS2 software were used to plot the graphs using the IGV browser (Robinson et al., 2011).

Protein extraction, gel running and western blotting

Yeast cells harvested at the mentioned time points were washed once with ice-cold water and
once with ice-cold 20% TCA. To preserve the phosphorylation moieties, the cells were lysed
in the presence of 1X PhosSTOP (Roche). The pellet was resuspended in 200 µl of 20% TCA,
and an equal volume of glass beads was added, and cells were lysed using a mini bead beater
(BIOSPEC, India) for 7 cycles (1 min ON, 2 mins OFF on ice). The supernatant was collected
after puncturing the O-ring tube and centrifugating at 2800 rpm for 2 mins at 4°C. 400 µl of 5%
TCA was added to the beads and spun again to collect the supernatant. The total supernatant
was spun at maximum speed for 10 mins at 4°C. The precipitate was kept for air-drying for
10-15 mins before resuspending in 2X SDS sample buffer and 1X Tris (pH 8.0) for
neutralization. Before loading, the protein was boiled for 5 mins at 95°C and spun at maximum
speed for 30 seconds. Standard procedure was followed for SDS-PAGE and western blot
transfer.
For the gel containing the PhosTag reagent, 10 mM $MnCl_2$ was added to the resolving
gel along with 50 µM of the PhosTag reagent (AAL-107, FUJIFILM, JAPAN). Before western
blotting, the gel was washed three times with 1 mM EDTA for 10 mins, followed by 2 washes
in transfer buffer. The blot was further incubated with primary antibodies at a dilution of 1:2500,
while the secondary antibodies were used at 1:5000 dilution.

Statistical analyses

The data presented in the main figures and the supplementary figures were obtained from two

to three independent experiments. The error bars in the individual bar graphs represent the

standard error (SE) derived from the standard deviation (SD). The statistical significance (p)

was determined by two-tailed Student's t-test, or one-way ANOVA test, as appropriate. When

comparing more than one group, a one-way ANOVA test was used; otherwise, Student's t-

test was used. The 'N' values denote the total number of cells analyzed from the combined 'n'

number of replicates of the individual assays. The p values less than or equal to 0.05 are

categorized as significant differences. The SD, SE, and statistical significance (p) values were

calculated using automated modules of GraphPad Prism 9.0 (Version 9.4.1) software.

840
Figures

Figure 1. Loss of Scm3 causes sensitivity to MMS-mediated DNA damage. (A) Wild-type

(*SCM3*), *SCM3-AID* (three independent transformants)*, and rad52Δ* cells grown till mid-log

were serially diluted and spotted on the indicated plates. The plates were incubated at 30°C

for 24-48 hrs before imaging. **(B)** An equal number of mid-log grown *SCM3-AID* cells were

spread on plates containing indicated supplements. From the number of colonies that

appeared after incubation at 30°C for 2 days, the cell viability was estimated (see materials

and methods) and plotted. Colonies formed on the plate without auxin and MMS were taken

as 100% viability, and accordingly, viability percentage was calculated for other plates. The *p*

values were estimated by one-way ANOVA test from three independent experiments. **(C)** Wild-

type (*SCM3*)*, SCM3-AID, SCM3-AID rad52Δ,* and *rad52Δ* cells grown till mid-log were serially

diluted and spotted in the presence of indicated auxin concentrations. The plates were
incubated at 30°C for 24-48 hrs before imaging. **(D)** The cells of the strains used in (C) were
grown till mid-log and treated with 0.75 mM auxin for the indicated duration before they were
washed, serially diluted, and spotted on YPD plates that were incubated at 30°C for 24-48 hrs
before imaging.

Figure 2. Loss of Scm3 causes increased Rad52 foci in the presence or absence of
MMS. (A) The schematic of the experimental strategy that was followed for visualization of
Rad52-GFP foci in the *RAD52-GFP SCM3-AID* cells either in the presence of DMSO (wild-
type) or auxin (Scm3 depleted) and each either treated with MMS or left untreated. **(B)**
Representative images showing different patterns of Rad52-GFP foci. Rad52-GFP produces
a nuclear signal (row 1) in the absence of any DNA damage, but with the damage, it coalesces
to form bright foci (rows 2-5) representing the DNA repair sites. **(C)** The percentage of
indicated cells containing Rad52-GFP foci with or without MMS treatment for 90 mins. **(D)** The
average number of Rad52-GFP foci per MMS-treated cell from the experiment described in
(C). At least 300 cells were analyzed from three independent experiments for each set in **(C)**

and **(D)**. The *p* values were estimated by unpaired two-tailed student's t-test. Scale bar = 2
μm.
873

874
875
Figure 3. The localization of Scm3 in response to DNA damage. (A) Representative
images showing the localizations of Ndc10-6HA and Scm3-13Myc using chromatin spread.
The mid-log grown *NDC10-6HA SCM3-13MYC* cells were treated with 0.02% MMS or mock-
treated for 90 mins for the assay. The arrow shows the co-localization of Ndc10-6HA and
Scm3-13Myc at the kinetochore cluster, and the arrowhead shows the rDNA loop. **(B)** The
intensities of Scm3-13Myc and Ndc10-6HA normalized with the background intensity were

quantified using ImageJ software and were plotted. At least 30 spreads from 2 independent
experiments were analyzed for each set in **(B).** The *p* values were estimated by unpaired two-
tailed student's t-test. Scale bar = 2 μm. **(C, D)** The ChIP-seq signal of Scm3 (Scm3-13Myc)
and input from cells either untreated or treated with 0.05% MMS for the entire chromosome I;
the no-tag strain was used as a control. The plot shows the binding of Scm3 across (C)
chromosome I and (D) chromosome IV. The green and black arrows represent the non-
centromeric localization of Scm3 in untreated, and MMS treated samples, respectively. **(E)**
Same as (C); only a portion of chromosome XII flanking 9.1 kb rDNA region is shown. The
scale is 7–50 for the input samples and 7-25 for IP samples.

891
892
893

894
895
Figure 4. Scm3 colocalizes with DDR proteins in response to DNA damage. (A)
Representative images showing the localizations of Rad52-6HA and Scm3-13Myc using
chromatin spread. The mid-log grown *RAD52-6HA SCM3-13MYC* cells were treated with
0.02% MMS or mock-treated for 90 mins for the assay. **(B)** The quantification of the
colocalization between Rad52-6HA and Scm3-13Myc was calculated using Pearson's
correlation coefficient (PCC). **(C)** Representative images showing the localizations of γ-H2A
and Scm3-13Myc using chromatin spread. The mid-log grown *SCM3-MYC* cells were treated
with 0.02% MMS or mock-treated for 90 mins for the assay. **(D)** The quantification of the
colocalization between γ-H2A and Scm3-13Myc was calculated using PCC. Anti-HA (12CA5,
Roche), anti-Myc (9E10, Roche), and anti-H2A (ab15083, Abcam) antibodies were used to
detect HA, Myc, and H2A-S129, respectively. At least 30 spreads from 2 independent

experiments were analyzed for each set in **(B),** and **(D)**. The *p* values were estimated by
unpaired two-tailed student's t-test. Scale bar = 2 µm.
909

Figure 5

Figure 5. Scm3 is recruited to a site-specific double-stranded break (DSB). (A)
Schematic representation of a cut site of the *HO* endonuclease (*HOcs*) integrated at the *URA3*
locus in the NA14 strain. Upon galactose induction, the *HO* endonuclease cleaves the DNA at
the specific site generating DSB. The positions of the primers (OSB289 and kanB1) are shown
that were used to detect the DSB, and the binding of Scm3 near and distal (-1 to -3 kb) to the
DSB **(B)** ChIP analyses were performed to quantify the association of Scm3-6HA in NA14
strain harboring *SCM3-6HA* with the indicated loci at the indicated time points. The PCR
amplified DNA bands following ChIP were run on agarose gel and stained with ethidium
bromide. **(C)** ChIP analyses for measuring the association of Scm3-6HA with the indicated loci
were performed at the indicated time points of HO induction using anti-HA (12CA5, Roche)
antibodies. The ChIP efficiency as percentage enrichment per input obtained by qPCR
analysis of the immunoprecipitated DNA was plotted. **(D)** Same analysis as in (C) using no
antibody (-Ab) control. The *p* values in **(C)** and **(D)** were estimated from two independent
experiments.
925
926

Figure 6. Evidence of improper activation of DNA damage checkpoint in absence of Scm3. (A) The schematic of the experimental strategy that was followed for detecting Rad53 in *SCM3-AID* cells in the presence of DMSO (wild-type) or auxin (Scm3 depleted) and each either treated with MMS or left untreated. **(B)** Left**,** western blots showing p-Rad53/Rad53 and tubulin (as loading control) using anti-Rad53 (ab104232, Abcam) and anti-tubulin (MCA78G, Serotec) antibodies, respectively, from indicated cells obtained from (A). Right, the intensities of the p-Rad53 measured by ImageJ were plotted after normalizing with the corresponding loading control. **(C)** The schematic of the experimental strategy to detect Rad53 in Scm3 or

Cdc20 depleted cells treated with MMS for different time points. **(D)** Western blots developed
as in **(B)** from indicated cells obtained from **(C)**. **(E)** *mad2Δ, SCM3-AID,* and *SCM3-AID mad2Δ*
cells were first treated with 0.75 mM auxin for 2 hrs to deplete Scm3 and then either treated
with 0.02% MMS for 90 mins or left untreated. The cells were harvested and were scored for
the presence of multibudded cells (shown on the left). At least 150 cells from three
independent experiments were analyzed for each set. The *p* values were estimated by one-
way ANOVA test. Scale bar = 2 μm.

Figure 7. Post-translational modification of Scm3 in response to DNA damage (A) Left,
western blots show Scm3-6HA and its modified form (Scm3*) along with tubulin as loading
control from the cells untreated or treated with indicated concentrations of MMS for 90 mins.
The cell lysates were probed using anti-HA (12CA5, Roche) or anti-tubulin (MCA78G, Serotec)
antibodies. Middle, the graph shows the intensity of Scm3* with increasing MMS
concentrations, normalized with respect to tubulin using ImageJ software. Right, the graph
represents the ratio of the modified to unmodified forms of Scm3. **(B)** The schematic of the
experimental strategy that was followed for detecting Scm3-6HA in the G1 arrested cells and
following their release from the arrest for the indicated time in the absence or presence of
MMS. **(C)** Representative images showing the cell cycle stages by DAPI staining and bud
morphology at indicated time points since their release from G1 arrest. **(D)** Western blot
showing Scm3-6HA along with tubulin as a loading control from the cells harvested, as shown

in (B). The cell lysates were probed using anti-HA and anti-tubulin antibodies to detect Scm3-
6HA and tubulin, respectively. Error bars represent the standard deviation from mean values
obtained from two independent experiments. Scale bar = 2 μm.

962
963

964
Figure 8: A working model summarizing the role of Scm3 in DNA damage response
(DDR). Left, in wild type cells, a subset of the Scm3 molecules present as a nuclear pool
(green) are phosphorylated upon DNA damage. The phosphorylated Scm3 (p-Scm3)
associates with DSB sites, colocalizes with yH2A, and spreads to the chromatin, flanking the
DSB site. Possibly, the p-Scm3 facilitates proper activation of Rad53 through its
phosphorylation, leading to cell cycle arrest. Additionally, p-Scm3 co-localizes with Rad52 and
assists in DSB repair. Right, in *scm3* mutant cells, there is also an increase in intrinsic DNA
damage, corresponding to more Rad52 foci formation, and due to the inability to repair DSBs,
the cell viability drops. In parallel, reduced Rad53 activation causes improper DDR-mediated
cell cycle arrest generating multibudded cells and aneuploids which causes cell death. The
phenotypes associated with the loss of Scm3 are shown in red.

976
977

References

1. Agmon N, Pur S, Liefshitz B, Kupiec M. 2009. Analysis of repair mechanism choice during
homologous recombination. *Nucleic Acids Res* **37**:5081–5092. doi:10.1093/NAR/GKP495
2. Amon A. 2002. Synchronization procedures. pp. 457–467. doi:10.1016/S0076-
6879(02)51864-4
3. Andrews, S., 2010. FastQC: a quality control tool for high throughput sequence data.
4. Aravind L, Iyer LM, Wu C. 2007. Domain architectures of the Scm3p protein provide
insights into centromere function and evolution. *Cell Cycle* **6**:2511–2515.
doi:10.4161/CC.6.20.4793
5. Barlow JH, Rothstein R. 2010. Timing is everything: cell cycle control of Rad52. *Cell Div*
5:7. doi:10.1186/1747-1028-5-7

6. Blower MD, Karpen GH. 2001. The role of Drosophila CID in kinetochore formation, cell-cycle progression and heterochromatin interactions. *Nat Cell Biol* **3**:730–739. doi:10.1038/35087045

7. Buchwitz BJ, Ahmad K, Moore LL, Roth MB, Henikoff S. 1999. A histone-H3-like protein in C. elegans. *Nature* **401**:547–548. doi:10.1038/44062

8. Burgess RJ, Zhang Z. 2013. Histone chaperones in nucleosome assembly and human disease. *Nat Struct Mol Biol* **20**:14. doi:10.1038/NSMB.2461

9. Burma S, Chen BP, Murphy M, Kurimasa A, Chen DJ. 2001. ATM Phosphorylates Histone H2AX in Response to DNA Double-strand Breaks. *Journal of Biological Chemistry* **276**:42462–42467. doi:10.1074/jbc.C100466200

10. Camahort R, Li B, Florens L, Swanson SK, Washburn MP, Gerton JL. 2007. Scm3 Is Essential to Recruit the Histone H3 Variant Cse4 to Centromeres and to Maintain a Functional Kinetochore. *Mol Cell* **26**:853–865. doi:10.1016/J.MOLCEL.2007.05.013

11. Chen T, Huang H, Zhou Y, Geng L, Shen T, Yin S, Zhou L, Zheng S. 2018. HJURP promotes hepatocellular carcinoma proliferation by destabilizing p21 via the MAPK/ERK1/2 and AKT/GSK3β signaling pathways. *Journal of Experimental and Clinical Cancer Research* **37**. doi:10.1186/s13046-018-0866-4

12. Chen Y, Baker RE, Keith KC, Harris K, Stoler S, Fitzgerald-Hayes M. 2000. The N Terminus of the Centromere H3-Like Protein Cse4p Performs an Essential Function Distinct from That of the Histone Fold Domain. *Mol Cell Biol* **20**:7037. doi:10.1128/MCB.20.18.7037-7048.2000

13. Cho US, Harrison SC. 2012. Ndc10 is a platform for inner kinetochore assembly in budding yeast. *Nat Struct Mol Biol* **19**:48–56. doi:10.1038/NSMB.2178

14. Cho US, Harrison SC. 2011. Recognition of the centromere-specific histone Cse4 by the chaperone Scm3. *Proc Natl Acad Sci U S A* **108**:9367–9371. doi:10.1073/PNAS.1106389108

15. Choy JS, Mishra PK, Au WC, Basrai MA. 2012. Insights into assembly and regulation of centromeric chromatin in Saccharomyces cerevisiae. *Biochim Biophys Acta Gene Regul Mech* **1819**:776–783. doi:10.1016/J.BBAGRM.2012.02.008

16. Ciccia A, Elledge SJ. 2010. The DNA Damage Response: Making It Safe to Play with Knives. *Mol Cell*. doi:10.1016/j.molcel.2010.09.019

17. Coates P, Dewar J, Thompson AM. 2010. At last, a predictive and prognostic marker for radiotherapy? *Breast Cancer Res* **12**:106. doi:10.1186/BCR2567

18. Conde F, San-Segundo PA. 2008. Role of Dot1 in the Response to Alkylating DNA Damage in Saccharomyces cerevisiae: Regulation of DNA Damage Tolerance by the Error-Prone Polymerases Polζ/Rev1. *Genetics* **179**:1197. doi:10.1534/GENETICS.108.089003

19. Dammann R, Lucchini R, Koller T, Sogo JM. 1993. Chromatin structures and transcription of rDNA in yeast Saccharomyces cerevisiae. *Nucleic Acids Res* **21**:2331. doi:10.1093/NAR/21.10.2331

20. De Koning L, Corpet A, Haber JE, Almouzni G. 2007. Histone chaperones: an escort network regulating histone traffic. *Nat Struct Mol Biol* **14**:997–1007. doi:10.1038/nsmb1318

21. Dornfeld KJ, Livingston DM. 1991. Effects of controlled RAD52 expression on repair and recombination in Saccharomyces cerevisiae. *Mol Cell Biol* **11**:2013–2017. doi:10.1128/MCB.11.4.2013-2017.1991

22. Dou Z, Qiu C, Zhang X, Yao S, Zhao C, Wang Z, Chu R, Chen J, Chen Z, Li R, Wang K, Liu P, Liu C, Song K, Kong B. 2022. HJURP Promotes Malignant Progression and Mediates Sensitivity to Cisplatin and WEE1-inhibitor in Serous Ovarian Cancer. *Int J Biol Sci* **18**:1188–1210. doi:10.7150/ijbs.65589

23. Dunleavy EM, Roche D, Tagami H, Lacoste N, Ray-Gallet D, Nakamura Y, Daigo Y, Nakatani Y, Almouzni-Pettinotti G. 2009. HJURP is a cell-cycle-dependent maintenance and deposition factor of CENP-A at centromeres. *Cell* **137**:485–497. doi:10.1016/J.CELL.2009.02.040

24. Earnshaw WC, Rothfield N. 1985. Identification of a family of human centromere proteins using autoimmune sera from patients with scleroderma. *Chromosoma 1985 91:3* **91**:313–321. doi:10.1007/BF00328227

25. Earnshaw WC, Tomkiel JE. 1992. Centromere and kinetochore structure. *Curr Opin Cell Biol* **4**:86–93. doi:10.1016/0955-0674(92)90063-I

26. Eitoku M, Sato L, Senda T, Horikoshi M. 2008. Histone chaperones: 30 years from isolation to elucidation of the mechanisms of nucleosome assembly and disassembly. *Cellular and Molecular Life Sciences* **65**:414–444. doi:10.1007/s00018-007-7305-6

27. Fan J, Moreno AT, Baier AS, Loparo JJ, Peterson CL. 2022. H2A.Z deposition by SWR1C involves multiple ATP-dependent steps. *Nat Commun* **13**:7052. doi:10.1038/s41467-022-34861-x

28. Fangaria N, Rani K, Singh P, Dey S, Kumar KA, Bhattacharyya S. 2022. DNA damage-induced nuclear import of HSP90α is promoted by Aha1. *Mol Biol Cell* **33**. doi:10.1091/MBC.E21-11-0554

29. Fischböck-Halwachs J, Singh S, Potocnjak M, Hagemann G, Solis-Mezarino V, Woike S, Ghodgaonkar-Steger M, Weissmann F, Gallego LD, Rojas J, Andreani J, Köhler A, Herzog F. 2019. The COMA complex interacts with Cse4 and positions Sli15/ipl1 at the budding yeast inner kinetochore. *Elife* **8**. doi:10.7554/ELIFE.42879

30. Foltz DR, Jansen LET, Bailey AO, Yates JR, Bassett EA, Wood S, Black BE, Cleveland DW. 2009. Centromere-Specific Assembly of CENP-A Nucleosomes Is Mediated by HJURP. *Cell* **137**:472–484. doi:10.1016/j.cell.2009.02.039

31. Foltz DR, Jansen LET, Black BE, Bailey AO, Yates JR, Cleveland DW. 2006. The human CENP-A centromeric nucleosome-associated complex. *Nat Cell Biol* **8**:458–469. doi:10.1038/NCB1397

32. Giaimo BD, Ferrante F, Herchenröther A, Hake SB, Borggrefe T. 2019. The histone variant H2A.Z in gene regulation. *Epigenetics Chromatin* **12**:37. doi:10.1186/s13072-019-0274-9

33. Daniel Gietz, R., & Woods, R. A. (2002). Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method (pp. 87–96). https://doi.org/10.1016/S0076-6879(02)50957-5

34. Gilbert CS, Green CM, Lowndes NF. 2001. Budding Yeast Rad9 Is an ATP-Dependent Rad53 Activating Machine. *Mol Cell* **8**:129–136. doi:10.1016/S1097-2765(01)00267-2

35. Haber JE. 2018. DNA repair: the search for homology. *Bioessays* **40**:e1700229. doi:10.1002/BIES.201700229

36. Hallwyl SCL, Shi I, Krejci L, Mü Ller ‡ C, Albertsen L, Sung P, Mortensen UH. 2008. Interaction with RPA Is Necessary for Rad52 Repair Center Formation and for Its Mediator Activity * □ S Iben Plate ‡1. doi:10.1074/jbc.M804881200

37. Hammond CM, Strømme CB, Huang H, Patel DJ, Groth A. 2017. Histone chaperone networks shaping chromatin function. *Nature Reviews Molecular Cell Biology 2017 18:3* **18**:141–158. doi:10.1038/nrm.2016.159

38. Hoffman CS. 1997. Preparation of Yeast DNA. *Curr Protoc Mol Biol* **39**. doi:10.1002/0471142727.mb1311s39

39. Howman E V., Fowler KJ, Newson AJ, Redward S, MacDonald AC, Kalitsis P, Choo KHA. 2000. Early disruption of centromeric chromatin organization in centromere protein A (*Cenpa*) null mice. *Proceedings of the National Academy of Sciences* **97**:1148–1153. doi:10.1073/pnas.97.3.1148

40. Hu B, Wang Q, Wang Y, Chen J, Li P, Han M. 2017. Holliday junction–recognizing protein promotes cell proliferation and correlates with unfavorable clinical outcome of hepatocellular carcinoma. *Onco Targets Ther* **10**:2601–2607. doi:10.2147/OTT.S127738

41. Hu Z, Huang G, Sadanandam A, Gu S, Lenburg ME, Pai M, Bayani N, Blakely EA, Gray JW, Mao JH. 2010. The expression level of HJURP has an independent prognostic impact and predicts the sensitivity to radiotherapy in breast cancer. *Breast Cancer Research* **12**. doi:10.1186/BCR2487

42. Janke C, Magiera MM, Rathfelder N, Taxis C, Reber S, Maekawa H, Moreno-Borchart A, Doenges G, Schwob E, Schiebel E, Knop M. 2004. A versatile toolbox for PCR-based

tagging of yeast genes: new fluorescent proteins, more markers and promoter substitution
cassettes. *Yeast* **21**:947–962. doi:10.1002/yea.1142

43. Javaheri A, Wysocki R, Jobin-Robitaille O, Altaf M, Cô Té § J, Kron SJ. 2006. Yeast G1
DNA damage checkpoint regulation by H2A phosphorylation is independent of chromatin
remodeling.

44. Kang DH, Woo J, Kim H, Kim SY, Ji S, Jaygal G, Ahn TS, Kim HJ, Kwak HJ, Kim CJ, Baek
MJ, Jeong D. 2020. Prognostic relevance of hjurp expression in patients with surgically
resected colorectal cancer. *Int J Mol Sci* **21**:1–12. doi:10.3390/ijms21217928

45. Kato T, Sato N, Hayama S, Yamabuki T, Ito T, Miyamoto M, Kondo S, Nakamura Y, Daigo
Y. 2007. Activation of Holliday junction-recognizing protein involved in the chromosomal
stability and immortality of cancer cells. *Cancer Res* **67**:8544–8553. doi:10.1158/0008-
5472.

46. Kaytor MD, Livingston DM. 1994. Saccharomyces cerevisiae RAD52 Alleles Temperature-
Sensitive for the Repair of DNA Double-Strand Breaks.

47. Kirkland JG, Peterson MR, Still Ii CD, Brueggeman L, Dhillon N, Kamakaka RT. 2015.
Heterochromatin formation via recruitment of DNA repair proteins **26**.
doi:10.1091/mbc.E14-09-1413

48. Kitada T, Schleker T, Sperling AS, Xie W, Gasser SM, Grunstein M. 2011. Cell Cycle γH2A
is a component of yeast heterochromatin required for telomere elongation View
supplementary material. *www.landesbioscience.com Cell Cycle* **293**:293–300.
doi:10.4161/cc.10.2.14536

49. Kumar D, Prajapati HK, Mahilkar A, Ma C-H, Mittal P, Jayaram M, Ghosh SK. 2021. The
selfish yeast plasmid utilizes the condensin complex and condensed chromatin for faithful
partitioning. *PLoS Genet* **17**:e1009660. doi:10.1371/journal.pgen.1009660

50. Krueger F. 2015. Trim Galore: a wrapper tool around Cutadapt and FastQC to consistently
apply quality and adapter trimming to FastQ
files. http://www.bioinformatics.babraham.ac.uk/projects/trim_galore.

51. Lai W, Zhu W, Xiao C, Li X, Wang Y, Han Y, Zheng J, Li Y, Li M, Wen X. 2021. HJURP
promotes proliferation in prostate cancer cells through increasing CDKN1A degradation
via the GSK3β/JNK signaling pathway. *Cell Death Dis* **12**. doi:10.1038/s41419-021-03870-
x

52. Lang J, Barber A, Biggins S. 2018. An assay for de novo kinetochore assembly reveals a
key role for the CENP-T pathway in budding yeast. *Elife* **7**. doi:10.7554/eLife.37819

53. Lanz MC, Yugandhar K, Gupta S, Sanford EJ, Faça VM, Vega S, Joiner AMN, Fromme
JC, Yu H, Smolka MB. 2021. In-depth and 3-dimensional exploration of the budding yeast
phosphoproteome. *EMBO Rep* **22**. doi:10.15252/EMBR.202051121

54. Lee C-S, Lee K, Legube G, Haber JE. 2014. Dynamics of yeast histone H2A and H2B
phosphorylation in response to a double-strand break. *Nat Struct Mol Biol* **21**:103–109.
doi:10.1038/nsmb.2737

55. Lee SE, Pellicioli A, Vaze MB, Sugawara N, Malkova A, Foiani M, Haber JE. 2003. Yeast
Rad52 and Rad51 Recombination Proteins Define a Second Pathway of DNA Damage
Assessment in Response to a Single Double-Strand Break. *Mol Cell Biol* **23**:8913–8923.
doi:10.1128/mcb.23.23.8913-8923.2003

56. Lim G, Chang Y, Huh W-K. 2020. Phosphoregulation of Rad51/Rad52 by CDK1 functions
as a molecular switch for cell cycle-specific activation of homologous recombination. *Sci*
Adv **6**.

57. Lisby M, Mortensen UH, Rothstein R. 2003. Colocalization of multiple DNA double-strand
breaks at a single Rad52 repair centre, NATURE CELL BIOLOGY.

58. Lisby M, Rothstein R, Mortensen UH. 2001. Rad52 forms DNA repair and recombination
centers during S phase. *Proc Natl Acad Sci U S A* **98**:8276–8282.
doi:10.1073/PNAS.121006298/ASSET/55754FA8-24FB-44C3-8D0D-
05A332BD04A3/ASSETS/GRAPHIC/PQ1210062005.JPEG

59. Longtine MS, Mckenzie III A, Demarini DJ, Shah NG, Wach A, Brachat A, Philippsen P,
Pringle JR. 1998. Additional modules for versatile and economical PCR-based gene

deletion and modification in Saccharomyces cerevisiae. *Yeast* **14**:953–961.
doi:10.1002/(SICI)1097-0061(199807)14:10<953::AID-YEA293>3.0.CO;2-U

60. Lowndes NF, Toh GWL. 2005. DNA repair: The importance of phosphorylating histone
    H2AX. *Current Biology* **15**:R99–R102. doi:10.1016/j.cub.2005.01.029

61. Luconi L, Araki Y, Erlemann S, Schiebel E. 2011. The CENP-A chaperone Scm3 becomes
    enriched at kinetochores in anaphase independently of CENP-A incorporation. *Cell Cycle*
    10:3369–3378. doi:10.4161/CC.10.19.17663

62. Luo D, Liao S, Liu Y, Lin Y, Li Y, Liao X. 2022. Holliday Cross-Recognition Protein HJURP:
    Association With the Tumor Microenvironment in Hepatocellular Carcinoma and With
    Patient Prognosis. *Pathology and Oncology Research* **28**.
    doi:10.3389/PORE.2022.1610506

63. Ma C-H, Kumar D, Jayaram M, Ghosh SK, Iyer VR. 2023. The selfish yeast plasmid
    exploits a SWI/SNF-type chromatin remodeling complex for hitchhiking on chromosomes
    and ensuring high-fidelity propagation. *PLoS Genet* **19**:e1010986.
    doi:10.1371/journal.pgen.1010986

64. MacGilvray ME, Shishkova E, Place M, Wagner ER, Coon JJ, Gasch AP. 2020. The
    phosphoproteome response to dithiothreitol reveals unique versus shared features of
    Saccharomyces cerevisiae stress responses. *J Proteome Res* **19**:3405.
    doi:10.1021/ACS.JPROTEOME.0C00253

65. Makrantoni V, Robertson D, Marston AL. 2019. Analysis of the Chromosomal Localization
    of Yeast SMC Complexes by Chromatin Immunoprecipitation. pp. 119–138.
    doi:10.1007/978-1-4939-9520-2_10

66. Mandemaker IK, Fessler E, Corujo D, Kotthoff C, Wegerer A, Rouillon C, Buschbeck M,
    Jae LT, Mattiroli F, Ladurner AG. 2023. The histone chaperone ANP32B regulates
    chromatin incorporation of the atypical human histone variant macroH2A. *Cell Rep*
    42:113300. doi:10.1016/j.celrep.2023.113300

67. McKinley KL, Cheeseman IM. 2016. The molecular basis for centromere identity and
    function. *Nat Rev Mol Cell Biol* **17**:16–29. doi:10.1038/NRM.2015.5

68. Mehta, G. D., Agarwal, M., & Ghosh, S. K. (2014). Functional characterization of
    kinetochore protein, Ctf19 in meiosis I: an implication of differential impact of Ctf19 on the
    assembly of mitotic and meiotic kinetochores in Saccharomyces cerevisiae. *Molecular
    Microbiology*, *91*(6), 1179–1199. https://doi.org/10.1111/mmi.12527

69. Melo J, Toczyski D. 2002. A unified view of the DNA-damage checkpoint. *Curr Opin Cell
    Biol*. doi:10.1016/S0955-0674(02)00312-5

70. Mishra PK, Au WC, Choy JS, Kuich PH, Baker RE, Foltz DR, Basrai MA. 2011.
    Misregulation of Scm3p/HJURP Causes Chromosome Instability in Saccharomyces
    cerevisiae and Human Cells. *PLoS Genet* **7**:1002303.
    doi:10.1371/JOURNAL.PGEN.1002303

71. Mittal P, Ghule K, Trakroo D, Prajapati HK, Ghosh SK. 2020. Meiosis-Specific Functions
    of Kinesin Motors in Cohesin Removal and Maintenance of Chromosome Integrity in
    Budding Yeast. *Mol Cell Biol* **40**. doi:10.1128/MCB.00386-19

72. Miyazaki T, Bressan DA, Shinohara M, Haber JE, Shinohara A. 2004. In vivo assembly
    and disassembly of Rad51 and Rad52 complexes during double-strand break repair.
    EMBO J **23**:939. doi:10.1038/SJ.EMBOJ.7600091

73. Mizuguchi G, Xiao H, Wisniewski J, Smith MM, Wu C. 2007. Nonhistone Scm3 and
    histones CenH3-H4 assemble the core of centromere-specific nucleosomes. *Cell*
    129:1153–1164. doi:10.1016/J.CELL.2007.04.026

74. Montes de Oca R, Gurard-Levin ZA, Berger F, Rehman H, Martel E, Corpet A, de Koning
    L, Vassias I, Wilson LOW, Meseure D, Reyal F, Savignoni A, Asselain B, Sastre-Garau X,
    Almouzni G. 2015. The histone chaperone HJURP is a new independent prognostic
    marker for luminal A breast carcinoma. *Mol Oncol* **9**:657–674.
    doi:10.1016/J.MOLONC.2014.11.002

75. Müller S, MontesdeOca R, Lacoste N, Dingli F, Loew D, Almouzni G. 2014.
    Phosphorylation and DNA binding of HJURP determine its centromeric recruitment and

function in CenH3CENP-A loading. *Cell Rep* **8**:190–203. doi:10.1016/J.CELREP.2014.06.002

76. Nishimura, K., Fukagawa, T., Takisawa, H., Kakimoto, T., & Kanemaki, M. (2009). An auxin-based degron system for the rapid depletion of proteins in nonplant cells. Nature Methods 2009 6:12, 6(12), 917–922. https://doi.org/10.1038/nmeth.1401

77. Pellicioli A, Foiani M. 2005. Signal Transduction: How Rad53 Kinase Is Activated. *Current Biology* **15**:R769–R771. doi:10.1016/j.cub.2005.08.057

78. Pizzul P, Casari E, Gnugnoli M, Rinaldi C, Corallo F, Longhese MP. 2022. The DNA damage checkpoint: A tale from budding yeast. *Front Genet* **13**:995163. doi:10.3389/FGENE.2022.995163/BIBTEX

79. Prajapati HK, Agarwal M, Mittal P, Ghosh SK. 2018. Evidence of *Zip1* Promoting Sister Kinetochore Mono-orientation During Meiosis in Budding Yeast. *G3 Genes|Genomes|Genetics* **8**:3691–3701. doi:10.1534/g3.118.200469

80. Ransom M, Dennehey BK, Tyler JK. 2010. Chaperoning histones during DNA replication and repair. *Cell* **140**:183. doi:10.1016/J.CELL.2010.01.004

81. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011. Integrative genomics viewer. Nature Biotechnology 2011 29:1 29:24–26. doi:10.1038/nbt.1754

82. Rg Bewersdorf J, Bennett BT, Knight KL. 2006. H2AX chromatin structures and their response to DNA damage revealed by 4Pi microscopy.

83. Sanchez-Pulido L, Pidoux AL, Ponting CP, Allshire RC. 2009. Common ancestry of the CENP-A chaperones Scm3 and HJURP. *Cell* **137**:1173–1174. doi:10.1016/J.CELL.2009.06.010

84. Serra-Cardona A, Zhang Z. 2018. Replication-Coupled Nucleosome Assembly in the Passage of Epigenetic Information and Cell Identity. *Trends Biochem Sci* **43**:136–148. doi:10.1016/j.tibs.2017.12.003

85. Shah S, Mittal P, Kumar D, Mittal A, Ghosh SK. 2023. Evidence of kinesin motors involved in stable kinetochore assembly during early meiosis. *Mol Biol Cell* **34**. doi:10.1091/mbc.E22-12-0569

86. Sherman F. 2002. Getting started with yeast. pp. 3–41. doi:10.1016/S0076-6879(02)50954-X

87. Shivaraju M, Camahort R, Mattingly M, Gerton JL. 2011. Scm3 is a centromeric nucleosome assembly factor. *J Biol Chem* **286**:12016–23. doi:10.1074/jbc.M110.183640

88. Shuaib M, Ouararhni K, Dimitrov S, Hamiche A. 2010. HJURP binds CENP-A via a highly conserved N-terminal domain and mediates its deposition at centromeres. *Proc Natl Acad Sci U S A* **107**:1349–1354. doi:10.1073/PNAS.0913709107

89. Sirbu BM, Cortez D. 2013. DNA Damage Response: Three Levels of DNA Repair Regulation. *Cold Spring Harb Perspect Biol* **5**. doi:10.1101/CSHPERSPECT.A012724

90. Smith JS. 2022. The long and short of rDNA and yeast replicative aging. doi:10.1073/pnas.2119593119

91. Smith MM. 2002. Centromeres and variant histones: what, where, when and why? *Curr Opin Cell Biol* **14**:279–285. doi:10.1016/S0955-0674(02)00331-9

92. Stoler S, Keith KC, Curnick KE, Fitzgerald-Hayes M. 1995. A mutation in CSE4, an essential gene encoding a novel chromatin-associated protein in yeast, causes chromosome nondisjunction and cell cycle arrest at mitosis. *Genes Dev* **9**:573–586. doi:10.1101/gad.9.5.573

93. Stoler S, Rogers K, Weitze S, Morey L, Fitzgerald-Hayes M, Baker RE. 2007. Scm3, an essential Saccharomyces cerevisiae centromere protein required for G2/M progression and Cse4 localization. *Proc Natl Acad Sci U S A* **104**:10571–10576. doi:10.1073/PNAS.0703178104

94. Sun Z, Bernstein E. 2019. Histone variant macroH2A: from chromatin deposition to molecular function. *Essays Biochem* **63**:59–74. doi:10.1042/EBC20180062

95. Valente V, Serafim RB, de Oliveira LC, Adorni FS, Torrieri R, da Cunha Tirapelli DP, Espreafico EM, Oba-Shinjo SM, Marie SKN, Paçó-Larson ML, Carlotti CG. 2013.

Modulation of HJURP (Holliday Junction-Recognizing Protein) levels is correlated with glioblastoma cells survival. *PLoS One* **8**. doi:10.1371/JOURNAL.PONE.0062200

96. Valente V, Teixeira SA, Neder L, Okamoto OK, Oba-Shinjo SM, Marie SKN, Scrideli CA, Paçó-Larson ML, Carlotti CG. 2009. Selection of suitable housekeeping genes for expression analysis in glioblastoma using quantitative RT-PCR. *BMC Mol Biol* **10**:1–11. doi:10.1186/1471-2199-10-17/TABLES/5

97. Van Hooser AA, Ouspenski II, Gregson HC, Starr DA, Yen TJ, Goldberg ML, Yokomori K, Earnshaw WC, Sullivan KF, Brinkley BR. 2001. Specification of kinetochore-forming chromatin by the histone H3 variant CENP-A. *J Cell Sci* **114**:3529–3542. doi:10.1242/jcs.114.19.3529

98. Wach A. 1996. PCR-synthesis of marker cassettes with long flanking homology regions for gene disruptions in S. cerevisiae. *Yeast* **12**:259–265. doi:10.1002/(SICI)1097-0061(19960315)12:3<259::AID-YEA901>3.0.CO;2-C

99. Wang C jing, Li X, Shi P, Ding H yan, Liu Y ping, Li T, Lin P ping, Wang Y shan, Zhang G qing, Cao Y. 2020. Holliday junction recognition protein promotes pancreatic cancer growth and metastasis via modulation of the MDM2/p53 signaling. *Cell Death Dis* **11**. doi:10.1038/s41419-020-2595-9

100. Wisniewski J, Hajj B, Chen J, Mizuguchi G, Xiao H, Wei D, Dahan M, Wu C. 2014. Imaging the fate of histone Cse4 reveals de novo replacement in S phase and subsequent stable residence at centromeres. *Elife* **2014**. doi:10.7554/ELIFE.02203\

101. Xiao H, Mizuguchi G, Wisniewski J, Huang Y, Wei D, Wu C. 2011. Nonhistone Scm3 Binds to AT-Rich DNA to Organize Atypical Centromeric Nucleosome of Budding Yeast. *Mol Cell* **43**:369–380. doi:10.1016/j.molcel.2011.07.009

102. Xiao H, Wang F, Wisniewski J, Shaytan AK, Ghirlando R, Fitzgerald PC, Huang Y, Wei D, Li S, Landsman D, Panchenko AR, Wu C. 2017. Molecular basis of CENP-C association with the CENP-A nucleosome at yeast centromeres. *Genes Dev* **31**:1958–1972. doi:10.1101/GAD.304782.117/-/DC1

103. Zhao X, Chabes A, Domkin V, Thelander L, Rothstein R. 2001. The ribonucleotide reductase inhibitor Sml1 is a new target of the Mec1/Rad53 kinase cascade during growth and in response to DNA damage. *EMBO J* **20**:3544. doi:10.1093/EMBOJ/20.13.3544

104. Zhao X, Muller EGD, Rothstein R. 1998. A Suppressor of Two Essential Checkpoint Genes Identifies a Novel Protein that Negatively Affects dNTP Pools. *Mol Cell* **2**:329–340. doi:10.1016/S1097-2765(00)80277-4

105. Zhou BBS, Elledge SJ. 2000. The DNA damage response: putting checkpoints in perspective. *Nature* **408**:433–439. doi:10.1038/35044005

106. Zhou X, Li W, Liu Y, Amon A. 2021. Cross-compartment signal propagation in the mitotic exit network. *Elife* **10**. doi:10.7554/eLife.63645

107. Zinchuk V, Grossenbacher-Zinchuk O. 2014. Quantitative Colocalization Analysis of Fluorescence Microscopy Images. *Curr Protoc Cell Biol* **62**. doi:10.1002/0471143030.cb0419s62