Habitat use of rewilded horses and cattle as related to the functional and structural composition of plant communities in a European restored wetland ecosystem

Lilla Lovász[1,2], Fränzi Korner-Nievergelt[2,3] and Valentin Amrhein[1,2]

[1] *Research Station Petite Camargue Alsacienne, 1 Rue de la Pisciculture, 68300 Saint Louis, France*
[2] *Department of Environmental Sciences, Universität Basel, Vesaglasse 1, 4051 Basel, Switzerland*
[3] *Oikostat GmbH, Rothmättli 16, 6218 Ettiswil, Switzerland*

Abstract

1. Rewilding initiatives in European open and semi-open lowlands increasingly involve cattle and horses for ecological restoration, especially in wetland areas of high conservation value. These large herbivores contribute to spatial heterogeneity and enhance biodiversity by shaping ecosystems through movement, grazing, and resting behaviours. However, the effect of their site-specific habitat use patterns on plant communities remains unclear.

2. In this study, we investigated the fine-scale spatiotemporal distribution of rewilded cattle and horses in a recently restored alluvial grassland in a French nature reserve. We explored differences in habitat use between the two species during summer and winter on a macrohabitat scale and examined structural and functional changes in vegetation traits over four years, focusing on plant height, patch cover, species richness, and light preference, nutrient-tolerance, and mowing/grazing tolerance of plants. The study site, a former agricultural area converted into a restored alluvial nature conservation site, allowed observation of ecological processes from a "zero state".

3. Our results suggest that cattle and horses exhibit similar habitat use with seasonal variations, potentially indicating partially shared feeding niches. The mixed-species grazing prevented vegetation overgrowth by keeping plant cover and vegetation height under control, yet without causing destructive impacts. The two herbivore species induced a clear increase in grazing-tolerant plants and slight changes in the abundance of light-preferring and nutrient-tolerant species.

4. *Synthesis*: Overall, we found that the varying spatiotemporal distribution of rewilded horses and cattle likely induces changes in plant community on the patch scale but results in vegetation stability on the landscape scale, which is known to facilitate ecosystem functioning. Our study therefore informs managers of conservation initiatives, proposing rewilding with year-round grazing horses and cattle a promising strategy for ecological restoration and natural habitat maintenance in wetland areas.

Introduction

Rewilding with domestic cattle and horse breeds is becoming increasingly widespread in ecological restoration initiatives in European open and semi-open lowlands, especially in wetland areas of high conservation value (Gordon et al., 1990; Wallis DeVries, 1998; Carver et al., 2021). Much evidence suggests that these large herbivores are able to substitute their ancestors – wild horses and aurochs – by creating and maintaining spatial heterogeneity (Trepel et al., 2024) and thus contributing to species richness and diversity (Olff and Ritchie, 1998; Stein et al., 2014; Lorimer et al., 2015; Pereira and Navarro, 2015; Gordon et al., 2021; Svenning et al., 2024). Large herbivores shape ecosystems through their movement-, grazing- and resting- behaviours (Liu et al., 2015) and by their habitat use (Martin et al., 2009). This habitat choice is influenced by resource availability (Aarts et al., 2013) and season (Zweifel-Schielly et al., 2009; Zielke et al., 2019), but it likely primarily depends on the herbivore species and their interspecific interaction, as bovids and equids differ in nutritional and behavioural needs (Olff and Ritchie, 1998; Esmaeili et al., 2021) but show a strong potential for competition due to shared habitat (Menard et al., 2002).

Only few studies of habitat selection of co-existing cattle and horses have been conducted to date (Menard et al., 2002; Lamoot et al., 2005; Thomassen et al., 2023), and these seem to reveal conflicting patterns. For example, Cromsigt et al. (2018) and (Lamoot et al., 2005) showed that horses almost exclusively feed on grasses and thus select grassland habitats, while cattle include a solid amount of woody vegetation in their diet and therefore select wood- and shrubland more often than horses. Thomassen et al. (2023) found high diet richness with low seasonal variability in horses, and less diverse diet yet with increasing summer-time variability in cattle. While these studies reflect considerable differences in habitat use of the two species, others, e.g., Nolte et al. (2014) and Menard et al. (2002), suggest strong competition for grasses between the two species especially in winter, meaning an overlapping habitat choice.

In the first part of our study, we therefore assessed the seasonal variation of space use intensity of semi-wild cattle and horses in a French nature reserve by following the spatiotemporal distribution of the grazers, and contrasted this space use pattern with the habitats on the study site to describe the habitat use of the grazers. Our first study question was: how do the habitat use patterns differ between horses and cattle, and how does such a difference vary between summer and winter?

Not only the habitat use patterns of co-existing cattle and horses are still unclear, but also the effect of this habitat use on plants is not well understood.
In a comprehensive review, Bakker et al. (2006) found the effects of large herbivores on plant diversity conflicting, with effects ranging from positive (Belsky, 1992; Collins et al., 1998) through neutral (Stohlgren et al., 1999; Adler et al., 2005) to even negative (Milchunas et al., 1998). This may be because grazers can have either compensatory or additive effects on vegetation through selective feeding on different competing plants, or through consumption of the same plants by multiple herbivore species (Ritchie and Olff, 1999). For example, Schmitz and Isselstein (2020) showed that cattle defoliate grasses more evenly, creating a homogenous sward of grazing-tolerant plants, while horses graze heterogeneously, avoiding foraging in latrine areas and strongly defoliating in other areas. As an outcome, responses of the plant community will result in changes in vegetation structure and diversity (Olff and Ritchie, 1998; Liu et al., 2015).

The response of plant community structure to grazers is believed to be principally
determined by functional traits of plants (Díaz et al., 2007). Investigating the response of
such functional traits are a "common currency" (McGill et al., 2006) to compare species
independently of species identity and across taxa (Reese et al., 2016). Changes in plant
functional traits as a function of grazer presence therefore offer generalizable predictions
about the effect of rewilding with large herbivores as a means for ecosystem restoration
(Atkinson et al., 2024). Yet, as Adler et al. (2005) points out, the challenge is identifying key
functional traits and then understanding how differences in their means and variances
translate into differences in response to grazing.
Such key traits may include tolerance for nutrient-rich soil, preference for exposure
to light, and tolerance of disturbance as predictors of changes in plant community as a
function of grazer presence (Milchunas et al., 1988; Augustine and McNaughton, 1998).
Further, simple vegetative traits such as plant height and patch cover may describe a
general response of herbaceous plant community to large herbivores (Díaz et al., 2001).
Results about how these traits predict plant responses are however often
contradicting. Herbivores seem to show opposing effect on plant species with different
nutrient-tolerance levels, as they alter rates of nutrient cycling and redistribute nutrient
availability (Trlica and Rittenhouse, 1993; Bardgett, 2005). For example, frequently grazed
patches may experience "nutrient export", resulting in a shift towards plant species with
lower nutrient demands (Nolte et al., 2014; Schmitz and Isselstein, 2020); but urine and
dung input by herbivores may also increase organic matter and nitrogen levels of the soil on
intensively visited patches (Harrison and Bardgett, 2008; Peco et al., 2017), leading to an
opposite effect: a shift towards nutrient-tolerant species.
Light-preference may similarly result in vegetation composition differences as a
function of grazers' space use, since heterogeneous grazing patterns can lead to spatial
heterogeneity of light availability (Bakker et al., 2003). Intense grazer presence at certain
areas may reduce competition for light by creating gaps and opening colonization windows
in the vegetation, and thus may favour light-demanding plants (Collins et al., 1998). Other
areas without frequent grazer visits may however increase competition for light, hosting
mainly the few most competitive light-preferent species and an array of shade-tolerant
plants as an understory (reviewed in: Marion et al., 2010). Yet, due to the heterogeneous
nature of grazing, such patterns are highly variable, and general predictions are hard to
draw.
Tolerance to disturbance, such as grazing and trampling (often termed as herbivory)
is regarded to be a basic predictor for changes in plant community composition (Augustine
and McNaughton, 1998; Díaz and Cabido, 2001; Evju et al., 2009). Herbivory affects plant
physiology, morphology, and genetics, thus plants have evolved to avoid or tolerate such
disturbances (Trlica and Rittenhouse, 1993). However, the several forms of herbivory
impacts and the variation in grazing pressure make predictions about the effects of grazing
on different plant species difficult (Reese et al., 2016). Furthermore, the tolerance of plants
to grazing depends also on how they interact with neighbouring plants (Hendrickson and
Olson, 2006).
As Herrero-Jáuregui and Oesterheld (2018) emphasised, most of our knowledge of
the effect of grazing animals on vegetation structure is based on grazed–non-grazed
contrasts, but the effects of a scenario where grazing intensity varies from low to high due
to the movement patterns of free-roaming animals are largely unknown. Also,

comparatively little attention has been paid to investigating the joint effect of co-existing
free-roaming cattle and horses (Liu et al., 2015), especially in nature conservation initiatives.
In the second part of our study, we thus carried out a large-scale macrohabitat
survey, and additionally we recorded vegetation characteristics on a microhabitat scale to
assess the changes that large herbivores induce in the plant community composition and
the physical vegetation structure.
Following Díaz et al. (2007), we assessed the structural responses of the vegetation
to answer the question: how does plant height, patch cover and species richness change as
a function of the spatiotemporal distribution of the grazers? We were also interested in how
plant functional traits of light preference, grazing tolerance and soil-nutrient tolerance
changed over four years, including one year before and three years after the introduction of
the cattle and horses on our study site.
The distinctive feature of our study lies in the fact that we were able to follow the
ecological processes from a "zero state" on an area of a recent ecological restoration
initiative. The study site used to be an agricultural area, where, after eliminating the
remains of pesticides, the colonisation by vegetation started from bare soil, and where wild-
living, free-roaming rewilded cattle and horses were then introduced to fulfil the functional
roles of their wild ancestors in regulating natural succession and in contributing to
ecosystem processes (Vera, 2000; Lovász, 2022; Schmitz et al., 2023).
Our overall aim was to contribute to a more comprehensive understanding of the
seasonal and interspecific variability of habitat use of large herbivores, and of the
vegetation functional trait-dynamics as an outcome of this habitat use in a rewilding-based
ecosystem restoration area. Our study thus helps to inform management strategies when
applying free-roaming horses and cattle in ecosystem restoration, and especially in
rewilding initiatives.
119
120   **Methods**
121
122   **Study area**
123
Our study area is situated on an island of the Rhine river and the Grand Canal of
Alsace in France, and is part of the national nature reserve Petite Camargue Alsacienne. A
total of 100 hectares of the island has been dedicated to an ecosystem restoration project
since 2014. Our study site is a 32-hectare area, part of the restored 100-ha area.
In the course of the restoration initiative, former crop fields in the area were
transformed into an alluvial environment. The reconstructed landscape comprises open
grasslands with scattered shrubs and tree groups, a few gravel sites without vegetation
cover, groundwater ponds, and a small creek allowing part of the Rhine water to flow
through the island. Remnants of old alluvial forest patches that surround the area remained
untouched during the ecosystem restoration. Natural succession from bare ground with
gravel areas started with an initial prompt to vegetation growth by seeding and planting
indigenous plant species in 2015 (Lachat, 2012). In addition, free-roaming Konik horses and
Highland cattle were introduced following the concept of rewilding, i.e., the passive
management of ecological succession to restore natural ecosystem processes and reduce
the human control of landscapes (Pereira and Navarro, 2015). The cattle and horses arrived
between September 2018 and March 2019 and their expected role was to act as a natural
disturbance regime (Mackey and Currie, 2000; Vera, 2000) on the restored alluvial

environment by creating spatial heterogeneity and thus enhancing biodiversity (Pereira and
Navarro, 2015; Carver et al., 2021). The animals have thus been grazing year-round in low
grazing intensity (0.3-0.5 animals per ha); human intervention such as supplementary
feeding has been restricted so that animals have a natural impact on their environment,
especially on the vegetation.
146
147 The national nature reserve Petite Camargue Alsacienne granted us permission to
148 conduct our study on this site.
149
150 **Grazer spatial distribution**
151
Upon their arrival to the Rhine island, the horses and the cattle were equipped with
GPS-collars (Followit, type Pellego) for management purposes of the nature reserve. The
collars recorded the position of the grazers every hour, and we used the data of summers
and winters from 2019 to 2021 (see Supplementary Material Table 1). We described the
space use of the grazers based on these hourly GPS positions. Parts of those data were used
in two other studies (Lovász et al, 2021, and Lovász et al 2024), but for better
understanding, we give a detailed explanation of the corresponding methods here as well.
During this three-year period, the number of grazers increased from the initial 5
horses and 5 cattle to 7 horses and 7 cattle.
The GPS fixes may show some imprecision because GPS accuracy can be affected by
satellite or receiver errors (Hurn 1993), satellite geometry (Dussault et al. 2001),
atmospheric conditions, topography, or overhead canopies (Di Orio et al. 2003; Moen et al.
1996). The GPS collars did not record DOP (Dilution of Precision) data and we therefore
could not correct for the inaccuracy of the fixes. However, since in an earlier study using a
part of the same GPS-collar data (Lovász et al., 2021), only 3.3% of all grazer positions fell
outside the fenced area, and since also earlier studies about GPS-error showed marginal
average inaccuracy (e.g., Ganskopp and Johnson, 2007), we assumed that inaccuracy would
not strongly influence our results; we thus excluded GPS positions erroneously falling
outside of the enclosure in our analysis.
The data were downloaded via satellite processing and through the interface of the
GPS collar provider (Followit, Lindesberg, Sweden) remotely, without contact to the grazers.
The use of GPS collars on both cattle and horses comply with animal welfare requirements
and reportedly does not cause disturbance to the animals (Ungar et al., 2005; Collins et al.,
2014).
176
177 **Macrohabitat data**
178
We carried out visual landcover surveys on the 32-ha study site in the median year of
our sampling period (in 2020). We divided the study area into 50x50-meter grid cells, and
with the support of a printed map with the projected grid cells, the observer (L.L.) surveyed
each grid cell by standing in the midpoint of the cell and visually estimating the percentages
of landcover types in the respective grid cell. The data from these surveys were also used in
(Lovász et al., 2024).
We distinguished the following landcover categories: trees (trees of ca. ≥ 3 years
old), saplings (all growth stages of young trees of ca. < 3 years old), shrubs, meadow, bare

ground, and surface of water in the respective grid cell. These macrohabitat-scale habitat
characteristics remained relatively stable throughout the 3-year study period.
189
190
191 **Microhabitat data**
192
We sampled herbaceous vegetation yearly between 2018 and 2021 on 80
permanent quadratic plots of 1 m^2 each. The locations of the plots were defined by the mid-
points of the 50-meter UTM-grid cells that we used also for the macrohabitat data. The
coordinates of the sample points were determined by using QGIS (version 3.4.4-Madeira).
Two months before the first data collection, we marked the sampling plots by digging
coloured flat concrete tiles into the ground, to facilitate finding the plots visually in high
vegetation and to avoid that the grazers stumble on them. To avoid potential effects of the
marking tiles on the vegetation, the closest corner of a sampling plot was placed at a
distance of 50 cm from the north-western corner of a tile.
Vegetation sampling (performed by L.L.) took place during the peak growing season,
between mid-May and mid-July. For the measurements, we used a modified Daubenmire
method (Daubenmire, 1959), by laying a 1x1-meter sized frame onto a sampling plot. Within
each plot, the following attributes were measured: plant height (cm); total vegetation cover
(%); species (name) of the three most abundant plant species. The variable 'plant height'
was the average of 20 measurements per plot that were taken along the two diagonals of
the 1-m^2 frame every 10 cm. We measured the total vegetation cover in the plot by visually
estimating the percentage of the plot covered by plants (on a continuous scale). The three
most abundant plant species in each plot were identified according to visual estimation of
their abundance; when identification to the species level was not possible due to missing
identification criteria (e.g., in graminoids, the lack of flowers due to grazing), we identified
the plant to the genus level.
Using the dataset of Flora Indicativa (Landolt et al., 2010), the following functional-
trait attributes were assigned to each species identified in our study area: light preference,
soil-nutrient tolerance and mowing/grazing tolerance (see Table 1 for the scales used to
describe the plant species).
218

Scale	Light (L)	Nutrients in the soil (N)	Mowing/grazing tolerance (MT)
1	full shade	very infertile	bad
2	shade	infertile	little
3	semi-shade	medium fertile	moderate
4	light	fertile	good
5	full light	over-rich	very good

Table 1. Plant functional trait attributes according to Landolt et al. (2010).

We estimated how the means of these functional traits of the most abundant
species in the 80 sampling plots changed as a function of horse and cattle location density.
Because vegetation sampling took place during the summers of the study period, we
assessed the effect of average grazer location density on the functional and structural traits
of plants in summer only. For each trait, we assessed the response to both horse and cattle
location density, and we also estimated average yearly changes.

In addition to the mean light preference of the three most abundant plant species in
a plot, we also assessed variation between plots by calculating the standard deviation of
light preference among the species.
We also calculated the overall species richness, species (alpha) diversity and relative
evenness of the plant community for each year of the study, using the species list arising
from taking the three most abundant species identified from all sampling plots. Species
richness refers to the sum of all species recorded on the study site in a given year. For
calculating species diversity, we used the Shannon-index H' = - $\sum p_i \ln(p_i)$, where p is the
proportion of individuals of the i species divided by the total number of species. We
computed the relative evenness using the Pielou's index J = H'/ln(N), where H' is the
Shannon-index and N is the total number of species in our study site. Both indices were
measured taking the number of occupied cells per species as if they were the numbers of
individuals.
239
240   **Statistical analysis**
241
For the macrohabitat analysis, we used the 50x50-m grid cells of the UTM grid over
the study area. We calculated the average number of locations in a grid cell across days and
individuals within each of the six seasons, i.e., summer and winter in the years 2019 to
2021. We averaged across days and individuals because seasons differed in duration and a
different number of animals were tracked in the different seasons (Suppl. Mat. Table 1). The
average number of locations in a grid cell was calculated separately for horses, cattle and
the sum of the two, i.e. "grazers". We increased the three variables by half of their minimal
non-zero value to get rid of the zero-values before we applied the logarithm transformation.
250
251
We analysed the logarithm transformed average number of locations per grid cell
using a liner mixed model with season (summer vs. winter), landcover variables and the
two-way interactions between season and each of the landcover variable as predictors. The
grid cell ID was used as a random factor to account for repeated measures of the same grid
cells over the six seasons. We included only five of the six landcover variables because the
six sum up to 100% which would cause redundancy in the model if all were included. We
decided to drop meadow because that was the most abundant landcover and leaving this
variable out produces a set of predictors with the lowest possible correlation. Of all
landcover variables, both linear and quadratic terms were included.
We checked the model assumptions (normal distribution, linear relationship,
homogeneity of variance) by standard diagnostic residual plots. We further checked for
spatial correlation by plotting the residuals in space (bubble plot) and by the semi-
variogram.
To analyse how grazer location density influences vegetation structure, we summed
grazer locations within each grid cell from 1 January to 1 June of each of four years. During
the first year 2018 grazer density was zero. Because not all animals were tracked and the
number of tracked animals varied over the five-month period, we corrected that number by
dividing it by the number of "animal-days" of tracked animals and multiplied it by the
number of "animal-days" of the total number of animals. The total number of horses varied
between 5 and 9 individuals (of which 3 to 5 were tracked) and the total number of cattle
varied between 3 and 8 individuals (of which 2 to 5 were tracked). In that way, we got for

every 50x50m grid cell an expected grazer density that was obtained from the space use of
the grazers and the total grazer abundance. We square-root transformed the horse and
cattle grazer density variables before we used them as predictors in the analyses of
vegetation structure.
We used a normal linear mixed model for analysing each of the six vegetation trait
variables: average light preference value, standard deviation of light preference value,
average nutrient-tolerance value, average mowing/grazing tolerance value, total cover,
average height. We used horse and cattle location density and year as numeric predictors,
and the cell ID was used as a random factor to account for the repeated measure of the
same cells over the four years. Standard diagnostic residual plots were used to assess
normal distribution of the residuals, linearity assumption and homogeneity of variance. We
further used semi-variograms to check for spatial correlation and auto-correlation plots to
check for temporal correlation.
Data preparation and model fitting was done in R 4.2.2 (RCoreTeam, 2022). For fitting the
normal linear mixed models, we used the function lmer of the package lme4 (Bates et al.,
2015). We used flat prior distributions for all model parameters to obtain posterior
distributions from which we report the 2.5% and 97.5% quantiles as lower and upper limits
of compatibility intervals of parameters and fitted values. The function sim from the
package arm (Gelman and Hill, 2007) was used to obtain the posterior distribution.
292
293

294 # Results

295
296 **Landcover characterization**
297
From the six landcover types (see Fig. 1), meadow cover was the most abundant
(42.3%). The relatively high percentage of tree cover (32.6%) was mostly due to an old oak
forest in the south-eastern corner of the grazed area, while at other parts of the study site,
trees were found only in small patches. The percentage of sapling cover (11.9%) indicated a
fast succession of mainly poplar trees. Open bare ground and shrub cover were present in
small proportions (4.7% and 2.4%). Water bodies represented 6.2% of all cover types and
were all groundwater ponds with stable water levels throughout the year.

Figure 1. Macrohabitat composition on the Rhine Island. Violin plots show the percentages for each land cover type, with mean percentage (white dot, and value in %) and standard deviation (white error bars) over all grid cells sampled

305
306
307

Macrohabitat use of horses and cattle

308
309
In our study site, Highland cattle and Konik horses used the different habitats in a
similar manner, but we found differences in seasonal patterns and the magnitude of the
patterns (Fig. 2 and 3, and Suppl. Mat. Fig. 1).
Both grazer species were similarly less frequently present in areas with high than
with low tree cover in summer, but in winter cattle densities outnumbered horse densities
in forested grid cells (Fig. 2 and 3).
Areas with higher sapling cover were increasingly used in winter by horses but not so
much by cattle. In summer, sapling-covered areas seemed not to be very important habitats
for either of the grazer species, as regression lines show a likely negative relationship (Fig. 2
and 3, but note that the compatibility intervals (Amrhein and Greenland, 2022) would allow
a slight positive relationship too).
Patterns were clearer regarding the use of grid cells with shrubs both in case of
cattle and horses (Fig. 2 and 3), with a preference for such shrub-covered habitats in
summer. However, note that shrub cover was rather patchy in our study site, with a
maximum of 20% cover by grid cell, and an average of 4.7% of the total area (Fig. 1),
therefore locations of animals in these habitats may be a consequence of selecting
neighbouring habitats (which were often meadows, since meadow was the most abundant
cover variable in our dataset).
328
329
330
331

332   .

Horse density   **Cattle density**

Meadows – i.e., habitats of extensive grass cover – were used rather evenly by both
grazer species, except in winter, when cattle densities decreased as meadow-cover
increased. This refers to a preference towards habitats other than only meadow-covered
areas in winter. In summer, the location densities of cattle were quite similar in all scales of
grass-cover, possibly meaning a preference for meadows compared to other cover variables,
yet without a strong trend. Horses on the other hand were aggregating on grid cells with the
highest meadow-cover percentages in summer.

Grid cells with average percentages of waterbodies were used by cattle more often
in summer than in winter, while we observed horses in similar densities in both seasons in
areas with water presence.

Unsurprisingly, neither grazer species selected areas with extensive bare-ground
percentages, both cattle and horses showed a clear negative relationship to bare-ground.

Grazer spatiotemporal densities and plant communities on a microhabitat scale

In the 80 sampling plots, the mean number of light-preferring plant species (Light
mean; Fig 4, Table 2) did not seem to change at areas with different intensity of occurrence
(i.e., GPS-location density) of horse and cattle (effect size: 0.00, CrI: 0.00 – 0.01 for both
horses and cattle). At the same time, throughout the 4 years of the study, there seemed to
be a slight decrease in species with higher light preference, but the effect was not strong
(effect size: -0.03, CrI: -0.07 – 0.01).

The variation between species of different light preference values (Light SD; Fig 4,
Table 2) showed a relatively flat relationship to horse density, but at higher cattle densities,

the within-plot variation in light preference among plant species was higher. The long-term
general trend was a rather slight decrease in the variation of species according to light
preference over the 4 years (effect size: -0.02, CrI: -0.06 – 0.02).
The mean number of plants with nutrient-rich soil preference (N mean; Fig 4, Table
2) did not seem to differ in areas with different intensity of horse presence (effect size:
0.00, CrI: -0.01 – 0.01), and their numbers were only slightly higher at higher cattle densities
(effect size: 0.01, CrI: -0.01 – 0.02). We found a slight yearly decrease in nutrient-preferring
plants (effect size: -0.07, CrI: -0.13 – 0.00).
Grazing-tolerant plant species (MV mean; Fig 4, Table 2) increased with increasing
horse densities but decreased with cattle densities. In the course of the four study years,
there was a clear increase in grazing-tolerant plant species (effect size: 0.23, CrI: 0.15–0.31).
Mean vegetation height (Fig 4, Table 2) was similar at low and high densities of both
horses and cattle. Yet, throughout the years, the mean vegetation height clearly decreased
(effect size (log): -0.25, CrI: -0,34 – 0.17).
The average of total vegetation cover seemed to increase in areas where horse
density was higher but the direction of the effect in case of cattle location densities
indicated a potentially negative relationship (although compatibility intervals might also
allow for the opposite direction of the effects, Fig. 4.). However, throughout the four years
of the study period – including the first year without grazer presence – the average total
cover seemed to stay rather stable (effect size: -0.1, CrI: -2,62 – 2.37). This may suggest that
the effects of horses and cattle potentially balance each other out on the overall vegetation
cover.
382
The total number of species decreased after grazers were introduced (2019), and
this trend continued until the third year of grazing (2021), when species richness slightly
increased again. Similarly, the Shannon-index showed a small decrease in diversity until
2021, when diversity increased again. The Pielou's index of evenness however almost did
not decrease from 2018 to 2019, and after a small decline, it slightly increased in the last
year of the study (Table 3.)
389
390

391
392
393

Figure 4. Relationship between horse and cattle location densities and plant functional (mean of light-preference values [Light mean]; standard deviation of light preference values [Light SD]; mean nutrient-tolerance [N mean]; mean of mowing/grazing tolerance values [MV mean]) and structural traits (mean height [cm] and total cover [%]), and the changes of the plant community along these traits throughout the 4 years of the study.

	Interc.	Int.lwr	Int.upr	horse	horse. lwr	horse. upr	cattle	cattle. lwr	cattle. upr	year.z	year. lwr	year. upr
Light.mean	3.38	3.30	3.46	0.00	0.00	0.01	0.00	0.00	0.01	-0.03	-0.07	0.01
Light.SD	0.40	0.33	0.47	0.00	-0.01	0.01	0.01	0.00	0.02	-0.02	-0.06	0.02
N.mean	3.02	2.90	3.15	0.00	-0.01	0.01	0.01	-0.01	0.02	-0.07	-0.13	0.00
MV.mean	2.88	2.73	3.03	0.01	0.00	0.03	-0.02	-0.04	0.00	0.23	0.15	0.31
log(height.m)	2.45	2.29	2.61	-0.01	-0.02	0.01	-0.01	-0.03	0.01	-0.25	-0.34	-0.17
total cover	70.66	64.76	76.70	0.67	0.15	1.20	-0.29	-0.82	0.25	-0.10	-2.62	2.37

Table 2. Response of plant functional and structural traits to grazing by horses and cattle. Effect sizes and compatibility intervals

Year	Nr of species	Shannon index	Pielou evenness
2018	51	3.352321	0.8526119
2019	46	3.263625	0.8524238
2020	42	3.099439	0.8292438
2021	43	3.138654	0.8344821

Table 3. Diversity of vegetation on the study site in the different years: Species richness (Nr of species), Shannon diversity index and Pielou evenness.

Discussion

Macrohabitat use of horses and cattle

Our investigation of the habitat use of co-existing free-roaming rewilded horses and cattle revealed overall similar habitat use, but with seasonal and, to some degree, directional differences. In light of our results, we assume that these two species share feeding niches to a certain level and may compete for certain resources, as for instance the findings of Scasta et al. (2016) suggest in western North American rangeland habitats, or as Menard et al. (2002) found in European wetlands. However, since overall grazing pressure remained low in our study site –0.3-0.4 animal per hectare –, resource availability stayed high enough throughout all seasons, therefore strong competition did likely not occur.

During summer, both grazer species exhibited low location densities in areas with high tree cover, but cattle were found in forested areas more frequently than horses. Especially in winter, cattle sought out grid cells in densely forested areas, while horses stayed away from high-percentage tree cover, similarly as for example Menard et al. (2002) and Cromsigt et al. (2018) found. The use of forested habitats by cattle in winter on our study site is in contrast with some earlier findings that showed the lowest probability of woodland selection by cattle in autumn and winter and the highest in spring and summer (Lamoot et al., 2005). On the other hand, the work by Pratt et al. (1986) detected a pattern similar to what we found, and assumed the shelter-seeking behaviour of cattle as a reason for searching woodlands in winter. This may have also played a role in our study area. However, on our study site, the reason why Highland cattle might have searched for forested areas in winter could have been their capacity to reach higher branches of trees with their horns and by bending them down, they could eat the leftover leaves and bark of the branch (L.L. personal observation). Previous studies (Putman et al., 1987; Bokdam et al., 2003) investigated the habitat use of horn-less cattle breeds, which may have been the cause of finding overwhelmingly higher grassland than woodland use by cattle. Our results may thus indicate that conservation grazing involving cattle breeds with long horns, such as Highland cattle, may facilitate the suppression of woody encroachment.

Horses, on the other hand, used areas with higher sapling cover during winter instead of old woodlands. Such habitat preference in free-roaming horses was, to our knowledge, so far not described. An earlier work found increased tree or shrub consumption of horses in winter (Thomassen et al., 2023), but in their study site, sapling-covered areas were not present in similarly high percentage as in our site. It seems therefore that in winter, upon availability, horses rather choose to feed on young trees than on older trees, by eating twigs and debarking thicker stands (L.L. personal observation). Since cattle only showed a relatively neutral relationship to sapling cover in winter, our results may refer to a possible niche separation by the two species.

Water-bodies on the Rhine Island did not seem to have a strong effect on horse location densities, but cattle seemed to search for specific grid cells with water-cover in them. This may be because – according to our personal observations (L.L.) – Highland cattle often cool themselves in warm weather in ponds, and likely also spend much time standing in water to avoid insect harassment. Such behaviour is a natural response to environmental conditions and was not observed in studies about domestic cattle (Senft et al., 1985). Our

475   result may therefore indicate that, to fulfil the natural needs of grazers in nature
476   conservation areas, the presence of large ponds is desirable over drinking wells.
477   As for the use of meadow habitats, horses seemed to be present constantly
478   throughout winter regardless of the percentage cover of meadows by grid cell, while in
479   summer, horse location density was larger in grid cells with very high (80-100%) meadow
480   cover. Our results are in line with other studies showing that horses prefer grassland areas
481   year-round (e.g., Girard et al., 2013; Scasta et al., 2016).
482   Cattle, on the other hand, were not found more often in habitats with high
percentage of grass cover in summer, even though the availability of meadows was high in
our study site (42,3% in total), and even though this species is also considered to be a grazer
rather than browser and to select for grasslands more often than woody areas (Pratt et al.,
1986). However, as it is theoretically assumed and also recently shown by Thomassen et al.
(2023), in co-existing herbivore communities, niche separation between cattle and horses
may lead to extended consumption of woody vegetation by cattle. Our results are in line
with this theory. Such patterns likely make it possible that the two grazer species may co-
exist in natural circumstances, and at the same time have a substantial effect on the
vegetation composition (Vermeulen, 2015), facilitating the aims of rewilding initiatives.
492
493
494   **Grazer spatiotemporal densities and plant communities on a microhabitat scale**
495
496   Our analysis of vegetation functional and structural traits in relation to grazer location
497   density showed different effects of horses and cattle, which seemed to either cancel out or
498   lead to different directional changes when looking at the long-term effects throughout the
499   years.
500   In the apparent response of the functional traits of plants to grazers, we did not
observe strong differences between herbivores and between years, except in the trait of
grazing tolerance. Only horse location densities seemed to be related to a higher number of
grazing-tolerant plants, while lower numbers of these plants were found at higher cattle
densities. This may be because free-roaming cattle are known to spend only about 30% of
their daily budget with grazing (Van Rees and Hutson, 1983; Tofastrud et al., 2018),
therefore the areas where their location densities accumulated in our study site were likely
not at feeding but rather at resting/ruminating areas. Resting and ruminating do not involve
defoliating disturbance on the vegetation, which can explain why cattle-presence was
negatively correlated with grazing-tolerant plant abundance. In contrast, since horses graze
about 50-70% of their daily time budget (Gudmundsson and Dyrmundsson, 1994; Houpt,
2005), high horse densities in our dataset likely indicated feeding activity (rather than only
trampling) by horses and thus defoliation disturbance on the vegetation. This may also
mean that in our dataset, the classification of plants by Landolt et al. (2010) as
grazing/mowing tolerant tended to include tolerance to defoliation rather than tolerance to
trampling. In view of this, it is not surprising that the average number of grazing/mowing
tolerant plants increased throughout the years (Pakeman, 2004; Evju et al., 2009), also
explaining why horse-density had a stronger influence on such plants than cattle-density.
518   Less clear were the effects of the herbivores on light- and nutrient-preferent plants.
519   The mean number and also the within-plot variation of light-preferent species
seemed to be mainly independent from densities of both horses and cattle.. The relatively
flat regression line of the correlations of both the mean and the standard deviation of light

preference and grazer densities throughout the four years of the study may indicate a
landscape-scale stability over years (Briske et al., 2003).Yet, the slight increase in the within-
plot variation of light-preferring plants at higher cattle densities may refer to heterogeneity
on the patch-scale and on the short term (Oñatibia et al., 2018; Li et al., 2021).
A similar trend seems to be apparent in the nutrient-preference trait of plants. The
nutrient distribution capacity of large herbivores (Doughty et al., 2016) may have resulted in
altered nutrient availability in the soil which could lead to differences in nutrient-tolerant
species abundance along the gradient of grazer densities (Harrison and Bardgett, 2008),
especially in cattle-frequented areas. Yet, the changes across years did not indicate a clear
difference and thus may suggest an overall stability of the nutrient preferring plant
community.
533
Structural changes, such as plant height and cover, seemed to be not considerable
on the long term. The mean total cover in the 80 sampling plots remained constant over the
four years of the study, probably because the percentage of vegetation cover increased with
high horse densities, and decreased with high cattle densities. Average height remained
relatively constant, with a small mean decline of approximately 10 cm-s between the first
and the last year of the study. The mixed-species grazing by horses and cattle therefore
seemed to have a conserving effect on plant cover and vegetation height, indicating that
such grazing effectively prevents vegetation overgrowth, but at the same time, does not
destroy plant physical structure. Our results indicate that, contrary to some earlier findings
(e.g., Eldridge et al., 2019) and the concerns of conservation managers, multi-species grazer
assemblages in close-to-natural (i.e. 0.3-0.4 animals per ha) densities do not have a
destructive impact on herbaceous plants, but are rather an effective tool in conservation
management, in line with recent findings (Bonavent et al., 2023).
Our estimation of species richness, diversity and evenness indices suggest slight
vegetation compositional changes, likely as a response to grazing. The limitation of our
estimation may be that we used only the three most abundant species rather than the
entire species pool in each of the 80 sampling plots as a proxy to calculate the diversity
indices, the results nevertheless seem to reflect a realistic trend. We detected a small
decrease in all indices at the beginning of the grazer introductions, which may be a result of
the sensitivity of some of the species (Olff and Ritchie, 1998) that were sown by the
management (L. Merckling, 2022, pers. com.) to establish the local species pool on a 'zero-
state' environment. Further, after introducing the herbivores, a visible decrease of invasive
plant species (e.g. , *Solidago gigantea*) was observed (L. Merckling, 2022, pers. com., and
L.L. personal observation) on the grazed area, which may be due to the effectiveness of
grazers in reducing certain alien species to the advantage of local species, similarly as Firn et
al. (2013) and Marchetto et al. (2021) found. These two factors therefore might have
decreased the total number and diversity of species shortly after the grazer introduction,
until it again started slightly increasing in the third year of the grazing management, possibly
by species that are more resistant to local conditions, as proposed by Olff and Ritchie
(1998). An opposite pattern was shown by Kimball and Schiffman (2003) and McIntyre et al.
(2003) but only for cattle grazing and in American and Australian environments, where
conditions substantially differ from the European context. Further research could thus
investigate the background of such effects in European mixed-grazer assemblages.
567
568

Conclusion

This study fills a gap in previous knowledge by linking the habitat use of rewilded cattle and horses with their effect on functional traits and structure of vegetation, since studies so far mainly investigated one of these subjects but not each of them. The results extend our current understanding of how these animals influence the plant community, by showing complementary habitat use and conserving effects on plant functional and structural traits. We conclude that the varying spatiotemporal distribution of these grazers likely involves changes on the patch scale but vegetation stability on the landscape scale, which is known to facilitate ecosystem functions. Our work may therefore serve as an important source for conservation managers: the results support nature conservation involving mixed species of large herbivores along rewilding principles.

Acknowledgements and funding

We thank the team of the Réserve Naturelle Petite Camargue Alsacienne for making it possible to conduct our research in the nature reserve. We are grateful for Heiner Lenzin for his support in plant species determination in the initial phase of the study.
This work was supported by the Fondation de bienfaisance Jeanne Lovioz, the Foundation Emilia Guggenheim-Schnurr, the Ornithologische Gesellschaft Basel, the Swiss Association Pro Petite Camargue Alsacienne, the Foundation Wolfermann-Nägeli, the Foundation Frey-Clavel, and the MAVA Foundation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

Lilla Lovász conceived, designed and performed the experiments, analyzed the data, prepared figures and/or tables, authored and/or reviewed drafts of the article, and approved the final draft.
Fränzi Korner-Nievergelt conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored and/or reviewed drafts of the article, and approved the final draft.
Valentin Amrhein conceived and designed the experiments, authored and/or reviewed drafts of the article, and approved the final draft.

References

Aarts, G., Fieberg, J., Brasseur, S., Matthiopoulos, J., 2013. Quantifying the effect of habitat availability on species distributions. J. Anim. Ecol. 82, 1135-1145.

Adler, P.B., Milchunas, D.G., Sala, O.E., Burke, I.C., Lauenroth, W.K., 2005. Plant traits and ecosystem grazing effects: comparison of US sagebrush steppe and Patagonian steppe. Ecol. Appl. 15, 774-792.

Amrhein, V., Greenland, S., 2022. Discuss practical importance of results based on interval estimates and p-value functions, not only on point estimates and null p-values. Journal of information technology 37, 316-320.

Atkinson, J., Gallagher, R., Czyżewski, S., Kerr, M., Trepel, J., Buitenwerf, R., Svenning, J.C., 2024. Integrating functional traits into trophic rewilding science. Journal of Ecology 112, 936-953.

Augustine, D.J., McNaughton, S.J., 1998. Ungulate effects on the functional species composition of plant communities: herbivore selectivity and plant tolerance. The Journal of wildlife management, 1165-1183.

Bakker, C., Blair, J., Knapp, A., 2003. Does resource availability, resource heterogeneity or species turnover mediate changes in plant species richness in grazed grasslands? Oecologia 137, 385-391.

Bakker, E.S., Ritchie, M.E., Olff, H., Milchunas, D.G., Knops, J.M., 2006. Herbivore impact on grassland plant diversity depends on habitat productivity and herbivore size. Ecol. Lett. 9, 780-788.

Bardgett, R., 2005. The biology of soil: a community and ecosystem approach. Oxford university press.

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software 67, 1 - 48.

Belsky, A.J., 1992. Effects of grazing, competition, disturbance and fire on species composition and diversity in grassland communities. Journal of vegetation science 3, 187-200.

Bokdam, J., Gleichman, J.M., van Wieren, S.E., Wallis De Vries, M.F., 2003. Free-ranging cattle as opportunistic exploiters of a successional woodland-grassland-heathland mosaic. Nature conservation and grazing management, 87-116.

Bonavent, C., Olsen, K., Ejrnæs, R., Fløjgaard, C., Hansen, M.D., Normand, S., Svenning, J.C., Bruun, H.H., 2023. Grazing by semi-feral cattle and horses supports plant species richness and uniqueness in grasslands. Appl. Veg. Sci. 26, e12718.

Briske, D.D., Fuhlendorf, S.D., Smeins, F.E., 2003. Vegetation dynamics on rangelands: a critique of the current paradigms. J. Appl. Ecol., 601-614.

Carver, S., Convery, I., Hawkins, S., Beyers, R., Eagle, A., Kun, Z., Van Maanen, E., Cao, Y., Fisher, M., Edwards, S.R., 2021. Guiding principles for rewilding. Conserv. Biol. 35, 1882-1893.

Collins, G.H., Petersen, S.L., Carr, C.A., Pielstick, L., 2014. Testing VHF/GPS collar design and safety in the study of free-roaming horses. PLoS One 9.

Collins, S.L., Knapp, A.K., Briggs, J.M., Blair, J.M., Steinauer, E.M., 1998. Modulation of diversity by grazing and mowing in native tallgrass prairie. Science 280, 745-747.

Cromsigt, J.P., Kemp, Y.J., Rodriguez, E., Kivit, H., 2018. Rewilding Europe's large grazer community: how functionally diverse are the diets of European bison, cattle, and horses? Restoration Ecology 26, 891-899.

Daubenmire, R.F., 1959. Canopy coverage method of vegetation analysis. Northwest Sci 33,
39-64.
Díaz, S., Cabido, M., 2001. Vive la différence: plant functional diversity matters to ecosystem
processes. Trends Ecol. Evol. 16, 646-655.
Díaz, S., Lavorel, S., McIntyre, S., Falczuk, V., Casanoves, F., Milchunas, D.G., Skarpe, C.,
Rusch, G., Sternberg, M., Noy-Meir, I., 2007. Plant trait responses to grazing–a global
synthesis. Global change biology 13, 313-341.
Díaz, S., Noy-Meir, I., Cabido, M., 2001. Can grazing response of herbaceous plants be
predicted from simple vegetative traits? J. Appl. Ecol. 38, 497-508.
Doughty, C.E., Roman, J., Faurby, S., Wolf, A., Haque, A., Bakker, E.S., Malhi, Y., Dunning,
J.B., Svenning, J.-C., 2016. Global nutrient transport in a world of giants. Proceedings of the
National Academy of Sciences 113, 868-873.
Eldridge, D.J., Travers, S.K., Val, J., Zaja, A., Veblen, K.E., 2019. Horse activity is associated
with degraded subalpine grassland structure and reduced habitat for a threatened rodent.
Rangel. Ecol. Manag. 72, 467-473.
Esmaeili, S., Jesmer, B.R., Albeke, S.E., Aikens, E.O., Schoenecker, K.A., King, S.R., Abrahms,
B., Buuveibaatar, B., Beck, J.L., Boone, R.B., 2021. Body size and digestive system shape
resource selection by ungulates: A cross-taxa test of the forage maturation hypothesis. Ecol.
Lett. 24, 2178-2191.
Evju, M., Austrheim, G., Halvorsen, R., Mysterud, A., 2009. Grazing responses in herbs in
relation to herbivore selectivity and plant traits in an alpine ecosystem. Oecologia 161, 77-
85.
Firn, J., Price, J.N., Whalley, R.D., 2013. Using strategically applied grazing to manage
invasive alien plants in novel grasslands. Ecological Processes 2, 1-11.
Ganskopp, D.C., Johnson, D.D., 2007. GPS error in studies addressing animal movements and
activities. Rangel. Ecol. Manag. 60, 350-358.
Gelman, A., Hill, J., 2007. Data Analysis Using Regression and Multilevel. Hierarchical
Models.
Girard, T.L., Bork, E.W., Nielsen, S.E., Alexander, M.J., 2013. Seasonal variation in habitat
selection by free-ranging feral horses within Alberta's forest reserve. Rangel. Ecol. Manag.
66, 428-437.
Gordon, I., Duncan, P., Grillas, P., Lecomte, T., 1990. The use of domestic herbivores in the
conservation of the biological richness of European wetlands. Bulletin d'écologie 21, 49-60.
Gordon, I.J., Pérez-Barbería, F.J., Manning, A.D., 2021. Rewilding lite: Using traditional
domestic livestock to achieve rewilding outcomes. Sustainability 13, 3347.
Gudmundsson, O., Dyrmundsson, O., 1994. Horse grazing under cold and wet conditions: a
review. Livestock Production Science 40, 57-63.
Harrison, K.A., Bardgett, R.D., 2008. Impacts of grazing and browsing by large herbivores on
soils and soil biological properties. The ecology of browsing and grazing, 201-216.
Hendrickson, J., Olson, B., 2006. Understanding plant response to grazing. Targeted Grazing:
A Natural Approach to Vegetation Management and Landscape Enhancement, ed.
Launchbaugh K, 32-39.
Herrero-Jáuregui, C., Oesterheld, M., 2018. Effects of grazing intensity on plant richness and
diversity: A meta-analysis. Oikos 127, 757-766.
Houpt, K.A., 2005. Maintenance behaviours, in: Mills, D.S., McDonnell, Sue M. (Ed.), The
domestic horse: the origins, development and management of its behaviour., Cambridge
University Press, pp. 94-108.

Kimball, S., Schiffman, P.M., 2003. Differing effects of cattle grazing on native and alien
plants. Conserv. Biol. 17, 1681-1693.
Lachat, B., 2012. Projet de renaturation d'un ancien bras sur l'Ile du Rhin, In: EDF (Ed.), EDF-
CIH.
Lamoot, I., Meert, C., Hoffmann, M., 2005. Habitat use of ponies and cattle foraging
together in a coastal dune area. Biol. Conserv. 122, 523-536.
Landolt, E., Bäumler, B., Erhardt, A., Hegg, O., Klötzli, F., Lämmler, W., Nobis, M., Rudmann-
Maurer, K., Schweingruber, F., Theurillat, J.-P., 2010. Flora indicativa= Ecological inicator
values and biological attributes of the flora of Switzerland and the Alps: ökologische
Zeigerwerte und biologische Kennzeichen zur Flora der Schweiz und der Alpen, Haupt
Verlag.
Li, W., Hooper, D.U., Wu, L., Bakker, J.D., Gianuca, A.T., Wu, X.B., Taube, F., Wang, C., Bai, Y.,
2021. Grazing regime alters plant community structure via patch-scale diversity in semiarid
grasslands. Ecosphere 12, e03547.
Liu, J., Feng, C., Wang, D., Wang, L., Wilsey, B.J., Zhong, Z., 2015. Impacts of grazing by
different large herbivores in grassland depend on plant species diversity. J. Appl. Ecol. 52,
1053-1062.
Lorimer, J., Sandom, C., Jepson, P., Doughty, C., Barua, M., Kirby, K.J., 2015. Rewilding:
science, practice, and politics. Annu. Rev. Environ. Resour. 40, 39-62.
Lovász, L., 2022. Nature conservation with large herbivores in a restored riparian
environment, University_of_Basel.
Lovász, L., Korner-Nievergelt, F., Amrhein, V., 2021. Grazer density and songbird counts in a
restored conservation area. PeerJ 9, e10657.
Lovász, L., Korner-Nievergelt, F., Amrhein, V., 2024. Effects of grazer density, season and
landcover on bird guilds in a restored conservation area. ResearchSquare.
Mackey, R.L., Currie, D.J., 2000. A re-examination of the expected effects of disturbance on
diversity. Oikos 88, 483-493.
Marchetto, K.M., Wolf, T.M., Larkin, D.J., 2021. The effectiveness of using targeted grazing
for vegetation management: a meta-analysis. Restoration Ecology 29, e13422.
Marion, B., Bonis, A., Bouzillé, J.-B., 2010. How much does grazing-induced heterogeneity
impact plant diversity in wet grasslands? Ecoscience 17, 229-239.
Martin, J., Tolon, V., Van Moorter, B., Basille, M., Calenge, C., 2009. On the use of telemetry
in habitat selection studies. Telemetry: Research, technology and applications. Hauppauge
(NY): Nova Science Publishers, 37-55.
McGill, B.J., Enquist, B.J., Weiher, E., Westoby, M., 2006. Rebuilding community ecology
from functional traits. Trends Ecol. Evol. 21, 178-185.
McIntyre, S., Heard, K., Martin, T., 2003. The relative importance of cattle grazing in
subtropical grasslands: does it reduce or enhance plant biodiversity? J. Appl. Ecol. 40, 445-
457.
Menard, C., Duncan, P., Fleurance, G., Georges, J.Y., Lila, M., 2002. Comparative foraging
and nutrition of horses and cattle in European wetlands. J. Appl. Ecol. 39, 120-133.
Milchunas, D., Sala, O., Lauenroth, W.K., 1988. A generalized model of the effects of grazing
by large herbivores on grassland community structure. The American Naturalist 132, 87-106.
Milchunas, D.G., Lauenroth, W.K., Burke, I.C., 1998. Livestock grazing: animal and plant
biodiversity of shortgrass steppe and the relationship to ecosystem function. Oikos 83, 65-
74.

Nolte, S., Esselink, P., Smit, C., Bakker, J.P., 2014. Herbivore species and density affect
vegetation-structure patchiness in salt marshes. Agriculture, Ecosystems & Environment
185, 41-47.
Olff, H., Ritchie, M.E., 1998. Effects of herbivores on grassland plant diversity. Trends Ecol.
Evol. 13, 261-265.
Oñatibia, G.R., Boyero, L., Aguiar, M.R., 2018. Regional productivity mediates the effects of
grazing disturbance on plant cover and patch-size distribution in arid and semi-arid
communities. Oikos 127, 1205-1215.
Pakeman, R.J., 2004. Consistency of plant species and trait responses to grazing along a
productivity gradient: a multi-site analysis. Journal of Ecology 92, 893-905.
Peco, B., Navarro, E., Carmona, C., Medina, N., Marques, M., 2017. Effects of grazing
abandonment on soil multifunctionality: The role of plant functional traits. Agriculture,
Ecosystems & Environment 249, 215-225.
Pereira, H.M., Navarro, L.M., 2015. Rewilding european landscapes. Springer Nature.
Pratt, R., Putman, R., Ekins, J., Edwards, P., 1986. Use of habitat by free-ranging cattle and
ponies in the New Forest, Southern England. J. Appl. Ecol., 539-557.
Putman, R., Pratt, R., Ekins, J., Edwards, P., 1987. Food and feeding behaviour of cattle and
ponies in the New Forest, Hampshire. J. Appl. Ecol., 369-380.
RCoreTeam, 2022. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. 2012.
Reese, A.T., Ames, G.M., Wright, J.P., 2016. Variation in plant response to herbivory
underscored by functional traits. PLoS One 11, e0166714.
Ritchie, M., Olff, H., 1999. Herbivore diversity and plant dynamics: compensatory and
additive effects. Herbivores: between plants and predators, 175-204.
Scasta, J.D., Beck, J.L., Angwin, C.J., 2016. Meta-analysis of diet composition and potential
conflict of wild horses with livestock and wild ungulates on western rangelands of North
America. Rangel. Ecol. Manag. 69, 310-318.
Schmitz, A., Isselstein, J., 2020. Effect of grazing system on grassland plant species richness
and vegetation characteristics: Comparing horse and cattle grazing. Sustainability 12, 3300.
Schmitz, O.J., Sylvén, M., Atwood, T.B., Bakker, E.S., Berzaghi, F., Brodie, J.F., Cromsigt, J.P.,
Davies, A.B., Leroux, S.J., Schepers, F.J., 2023. Trophic rewilding can expand natural climate
solutions. Nature Climate Change, 1-10.
Senft, R., Rittenhouse, L., Woodmansee, R., 1985. Factors influencing patterns of cattle
grazing behavior on shortgrass steepe. Rangeland Ecology & Management/Journal of Range
Management Archives 38, 82-87.
Stein, A., Gerstner, K., Kreft, H., 2014. Environmental heterogeneity as a universal driver of
species richness across taxa, biomes and spatial scales. Ecol. Lett. 17, 866-880.
Stohlgren, T.J., Schell, L.D., Vanden Heuvel, B., 1999. How grazing and soil quality affect
native and exotic plant diversity in Rocky Mountain grasslands. Ecol. Appl. 9, 45-64.
Svenning, J.-C., Buitenwerf, R., Le Roux, E., 2024. Trophic rewilding as a restoration
approach under emerging novel biosphere conditions. Curr. Biol. 34, R435-R451.
Thomassen, E.E., Sigsgaard, E.E., Jensen, M.R., Olsen, K., Hansen, M.D., Svenning, J.C.,
Thomsen, P.F., 2023. Contrasting seasonal patterns in diet and dung-associated
invertebrates of feral cattle and horses in a rewilding area. Mol. Ecol.
Tofastrud, M., Hegnes, H., Devineau, O., Zimmermann, B., 2018. Activity patterns of free-
ranging beef cattle in Norway. Acta Agriculturae Scandinavica, Section A—Animal Science
68, 39-47.

Trepel, J., le Roux, E., Abraham, A.J., Buitenwerf, R., Kamp, J., Kristensen, J.A., Tietje, M.,
Lundgren, E.J., Svenning, J.-C., 2024. Meta-analysis shows that wild large herbivores shape
ecosystem properties and promote spatial heterogeneity. Nature Ecology & Evolution 8,
705-716.
Trlica, M., Rittenhouse, L., 1993. Grazing and plant performance. Ecol. Appl. 3, 21-23.
Ungar, E.D., Henkin, Z., Gutman, M., Dolev, A., Genizi, A., Ganskopp, D., 2005. Inference of
animal activity from GPS collar data on free-ranging cattle. Rangel. Ecol. Manag. 58, 256-
266.
Van Rees, H., Hutson, G., 1983. The behaviour of free-ranging cattle on an alpine range in
Australia. Rangeland Ecology & Management/Journal of Range Management Archives 36,
740-743.
Vera, F.W.M., 2000. Grazing ecology and forest history. CABI publishing.
Vermeulen, R., 2015. Natural grazing: Practices in the rewilding of cattle and horses.
Rewilding Europe.
Wallis DeVries, M.F., 1998. Large herbivores as key factors for nature conservation, in:
WallisDeVries, M.F., Bakker, J.P., van Wieren, S.E. (Eds.) (Ed.), Grazing and Conservation
Management, Conservation Biology Series 11., Kluwer, Dordrecht, pp. pp1-17.
Zielke, L., Wrage-Mönnig, N., Müller, J., Neumann, C., 2019. Implications of spatial habitat
diversity on diet selection of European Bison and Przewalski's Horses in a Rewilding Area.
Diversity 11, 63.
Zweifel-Schielly, B., Kreuzer, M., Ewald, K.C., Suter, W., 2009. Habitat selection by an Alpine
ungulate: the significance of forage characteristics varies with scale and season. Ecography
32, 103-113.
815
816
817
818