A novel quinone biosynthetic pathway illuminates the evolution of aerobic metabolism

Felix J. Elling[1,2,*], Fabien Pierrel[3], Sophie-Carole Chobert[3], Sophie S. Abby[3], Thomas W. Evans[4,5,#],

Arthur Reveillard[3], Ludovic Pelosi[3], Juliette Schnoebelen[3], Jordon D. Hemingway[6], Ahcène

Boumendjel[7], Kevin W. Becker[8], Pieter Blom[9], Julia Cordes[5], Vinitra Nathan[1,§], Frauke Baymann[10],

Sebastian Lücker[9], Eva Spieck[11], Jared R. Leadbetter[12,13], Kai-Uwe Hinrichs[5], Roger E. Summons[4], Ann

Pearson[1]

[1] Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA 02138, USA

[2] Leibniz-Laboratory for Radiometric Dating and Isotope Research, Christian-Albrecht University of

Kiel, 24118 Kiel, Germany

[3] Univ. Grenoble Alpes, CNRS, UMR 5525, VetAgro Sup, Grenoble INP, TIMC, 38000 Grenoble,

France

[4] Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology,

Cambridge, MA 02139, USA

[5] MARUM – Center for Marine Environmental Sciences and Department of Geosciences, University

of Bremen, 28359 Bremen, Germany

[6] Geological Institute, Department of Earth Sciences, ETH Zurich, 8092 Zurich, Switzerland

[7] Univ. Grenoble Alpes, INSERM, LRB, Grenoble, 38000, France

[8] GEOMAR Helmholtz Centre for Ocean Research Kiel, 24148 Kiel, Germany

[9] Department of Microbiology, Radboud Institute for Biological and Environmental Sciences, Radboud

University, 6525 AJ Nijmegen, The Netherlands

[10] Laboratoire de Bioénergétique et Ingénierie des Protéines UMR 7281 CNRS/AMU, FR3479, F-

13402 Marseille Cedex 20, France

[11] Department of Microbiology and Biotechnology, University of Hamburg, 22609 Hamburg, Germany.

[12] Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA

91125, USA.

[13] Division of Engineering and Applied Science, California Institute of Technology, Pasadena, CA

91125, USA.

[*]Corresponding author: felling@leibniz.uni-kiel.de

[#]Present address: Shell Global Solutions International BV, 1031 CM Amsterdam, The Netherlands

[§]Present address: Department of Biology, Boston University, Boston, MA 02215, USA

Abstract

The dominant organisms in modern oxic ecosystems rely on respiratory quinones with high redox

potential (HPQs) for electron transport in aerobic respiration and photosynthesis. The diversification of

quinones, from low redox potential in anaerobes to HPQs in aerobes, is assumed to have followed

Earth's surface oxygenation ~2.3 billion years ago. However, the evolutionary origins of HPQs remain

unresolved. Here, we reconstruct the biosynthetic pathway of a novel HPQ, methyl-plastoquinone, that

is unique to bacteria of the phylum *Nitrospirota*. We demonstrate that the three extant HPQ biosynthetic

pathways, in *Nitrospirota*, *Cyanobacteriota*, and *Pseudomonadota*, share a common origin that predates

the emergence of these phyla. We show that aerobic metabolism using HPQs is ancestral to

Cyanobacteriota and *Pseudomonadota* and significantly predates Earth's surface oxygenation.

Introduction

The oxygenation of Earth's surface environments following the emergence of oxygenic photosynthesis in ancestors of *Cyanobacteriota* enabled the metabolic and genetic diversification of life (*1–4*). The use of oxygen as a terminal electron acceptor, i.e., aerobic respiration, enabled a higher energy yield compared to anaerobic metabolisms and was a prerequisite for the emergence of eukaryotes (*5, 6*). However, it remains poorly resolved how and when the electron transport chain (ETC) used for aerobic respiration evolved. While geochemical evidence indicates iron oxidation by acidophilic bacteria must have evolved by the time oxygen accumulated in the atmosphere during the great oxygenation event (GOE; ~2.4-2.3 Ga) (*3, 7–9*), there is now considerable evidence for an ancient origin of dioxygen-utilizing and detoxifying enzymes as early as 3.1 Ga (*10–12*). Though these enzymes may not have participated in aerobic respiration (*11, 13, 14*), their widespread occurrence in bacteria suggests the availability of oxygen in physiologically significant quantities, at least in some niches, before the GOE. Studying the evolution of ETC components, such as oxygen reductases that use electrons derived from the ETC (*15–20*), can help elucidate the origins of aerobic metabolisms. However, the interpretation of oxygen reductase evolution has remained contentious (*15–19, 21*), and alternative roles of ancestral oxygen reductases in oxygen detoxification and nitric oxide reduction rather than aerobic respiration have been proposed (*16*). Exploring the evolution of other ETC components, such as respiratory quinones, may yield new insights into the evolution of ETCs and aerobic respiration.

Strict anaerobes use ETCs and quinones with low redox potential (LPQs), while aerobes and facultative aerobes generally use high-potential quinones (HPQs) (*22–24*). HPQs require all parts of the ETC to operate at high redox potential (*24–26*) and confer no known benefit over LPQs under anaerobic conditions. However, under aerobic conditions HPQs are advantageous due to their decreased electron leakage to oxygen, thus reducing oxidative stress and minimizing free energy losses (*25, 27*). The occurrence of HPQs may represent a marker for high-potential ETCs and their evolution may be tied to the history of oxygenic photosynthesis and aerobic respiration. Within bacteria, HPQs have been found only in two phyla, oxygenic *Cyanobacteriota* (here used *sensu stricto*, including only *Cyanophyceae*)

and *Pseudomonadota* (formerly Proteobacteria, now comprising the classes Alpha-, Beta-, Gamma-,

and Zetaproteobacteria, Acidithiobacillia, and Hydrogenophilalia) (*28*). The *Cyanobacteriota* and

Pseudomonadota produce two distinct types of HPQs, plastoquinone (PQ) and ubiquinone (UQ),

respectively (*22, 23*), which became the quinones of plastids (PQ) and mitochondria (UQ) through

endosymbiosis during the early evolution of eukaryotes (*23, 29*). Yet, despite the dominance of HPQ-

utilizing organisms in Earth's oxic environments today (*30–35*), the co-evolution of HPQs and Earth

surface oxygenation remains largely unresolved (*36*).

Recent progress in metagenomic coverage of uncultivated bacteria and isolation of novel lineages

may help elucidate HPQ evolution through the discovery of new quinone structures and biosynthetic

pathways in unstudied lineages of aerobic bacteria. Here, we describe the discovery of a third, novel

type of HPQ, methyl-plastoquinone (mPQ). mPQ occurs only in aerobic members of the phylum

Nitrospirota (formerly Nitrospirae), a metabolically diverse group of bacteria that perform essential

transformations in the biogeochemical cycles of iron, nitrogen, and manganese. We characterize the

biosynthetic pathway of mPQ using bioinformatic, genetic, and biochemical techniques and use these

data to infer the evolutionary history of HPQs. Our study sheds new light on the evolutionary history

of ETCs by revealing a single origin of the three HPQ biosynthetic pathways prior to the radiation of

crown-group *Cyanobacteriota*, *Nitrospirota*, and *Pseudomonadota*, which evidently preceded the GOE.

Results & Discussion

Novel respiratory quinones in Nitrospirota

Despite their widespread distribution and the important roles of *Nitrospirota* in biogeochemical

cycles of iron, manganese, and nitrogen (*37–41*), many aspects of their chemotaxonomy and

bioenergetics remain understudied. Genome-based bioenergetic models implicate the presence of ETCs

in aerobic and anaerobic *Nitrospirota* (*41–45*), yet their corresponding respiratory quinones have not

been studied. During screening of *Nitrospirota* genomes for lipid biosynthetic pathways (*46*), we

observed that the genomes of aerobic *Nitrospirota* did not contain any of the characterized quinone

biosynthesis pathways (*23, 36*). In contrast, genomes of anaerobic sulfur-reducing *Nitrospirota*, i.e.,

Thermodesulfovibrio species and some *Nitrospirota* metagenome-assembled genomes from anoxic

environments, contained the futalosine pathway (MK$_{mqn}$, composed of *mqn* genes) for biosynthesis of

the LPQ, menaquinone (MK; Supplementary Datafile S1).

To evaluate the presence of respiratory quinones, we analyzed lipid extracts of one anaerobic and

eight aerobic species of *Nitrospirota*, covering all formally described genera (*Thermodesulfovibrio*,

Leptospirillum, *Nitrospira*, and *Manganitrophus*), using high-performance liquid chromatography

coupled to high-resolution tandem mass spectrometry. MKs were detected only in the anaerobic species,

Thermodesulfovibrio islandicus (Fig. 1a), and we did not find any of the previously known respiratory

quinone types in the aerobic *Nitrospirota*. Instead, all eight studied aerobic *Nitrospirota* contained a

novel type of quinone, identified as methyl-plastoquinone (mPQ). The polyprenyl chain of mPQ varied

in length and saturation depending on species (Fig. 1, Fig. S1, SI results). Mass spectrometric

characterization of mPQ revealed fragmentation spectra analogous to PQ but with a dominant ion at

m/z 165 instead of 151, indicative of a distinctive trimethyl-benzoquinone headgroup connected to the

isoprenoid tail (Fig. 1, Fig. S1, Table S1). Stable isotope labeling experiments and nuclear magnetic

resonance spectroscopy of mPQ confirmed the structural assignment of the headgroup (SI Results &

Discussion; Fig. S2-4). Specifically, [1]H-NMR spectra showed the absence of any proton linked to the

C2 of the quinone moiety and [1]H-NMR and [13]C-NMR confirmed the presence of a third methyl group

(Fig. S4; see SI Results & Discussion). mPQ is thus structurally related to both UQ (methylated at C2

of the benzoquinone) and PQ (methylated at C5 and C6).

Fig. 1. Novel quinones detected in aerobic *Nitrospirota*. a-d, Chromatograms showing presence of a distinct quinone type (methyl-plastoquinone, mPQ) in aerobic *Nitrospirota* (*Nitrospira marina*, *Leptospirillum ferrooxidans*, *Ca.* Manganitrophus noduliformans) and canonical menaquinones (MK) in the anaerobic *Nitrospirota* species *Thermodesulfovibrio islandicus*. Ubiquinone (UQ$_{8:8}$) in the *Ca.* Manganitrophus-*Ramlibacter* co-culture derives from *Ramlibacter* (see Fig. S1). **e-g**, High resolution mass spectrometric characterization of mPQ$_{9:9}$ and PQ$_{9:9}$ showing similar fragmentation patterns but suggesting the presence of a trimethyl-benzoquinone moiety in mPQ$_{9:9}$ (see Fig. S1); structure and fragmentation pattern of UQ$_{8:8}$ from *Ramlibacter* shown in **g** for reference.

Characterization of the biosynthetic pathway of mPQ

Based on the structure of mPQ, we hypothesized that its biosynthesis pathway might share characteristics with the UQ and PQ biosynthesis pathways. The PQ biosynthesis pathway of *Cyanobacteriota* has been partially resolved (*47, 48*) and contains several enzymes that are homologous to enzymes involved in the well-characterized bacterial UQ pathway (*36, 49*). In both pathways, the conversion of chorismate into 4-hydroxybenzoate (4-HBA) is mediated by a UbiC homolog and the

subsequent prenylation, decarboxylation, and hydroxylation of 4-HBA involve the UbiA, UbiD/X, and UbiH homologs, respectively (Fig. 2a) (*36*). Specific to the UQ pathway, methylation at C2 is mediated by UbiE (*50*). In the cyanobacteriotal PQ pathway, methylation at C5 and C6 has been proposed to be mediated by Sll0418 (PlqQ) (*49*, *51*).

Isotope labeling experiments further point to biochemical similarities between the HPQ biosynthesis pathways. Supplementation of cultures with ring-$^{13}C_6$-labeled substrates demonstrates that 4-HBA is the ring precursor in *Nitrospirota*, similar to *Cyanobacteriota* and *Pseudomonadota* (Fig. S5, SI Discussion). Further, experiments with methyl-$^2H_3$ methionine indicate that all three methyl groups of mPQ (at C2, C5, and C6) are derived from methionine via SAM-dependent methyltransferases (Fig. S2). We used this information to find multiple, homologous candidate genes for the biosynthetic pathway of mPQ in genomes of *Nitrospirota*. We suggest a gene nomenclature for the mPQ pathway (*mpq*) analogous to that of the UQ pathway and extend this to the PQ pathway (*plq*; Fig. 2). We identified a four-gene cluster in *Leptospirillum* spp. (Fig. S6), encoding a *ubiA* family prenyltransferase (*mpqA*; LFE_2122), *ubiC*-like chorismate pyruvate lyase (*mpqC*; LFE_2123), a cobalamin-binding radical *S*-adenosyl methionine (SAM) methyltransferase (LFE_2124), and a *ubiE*-like methyltransferase (*mpqE*; LFE_2125). The genes are not co-localized in other *Nitrospirota*, but *mpqA* and *mpqE* homologs are found in all aerobic *Nitrospirota*, in addition to a *ubiB*-like kinase (*mpqB*). By contrast, homologs of *mpqC*, *ubiD/X* (*mpqD/X*) and *plqQ* (*mpqQ*) occur only in a subset of aerobic *Nitrospirota* (Table S2). No clear *ubiH* homologs were identified. Consequently, aerobic *Nitrospirota* contain a mosaic pathway for mPQ biosynthesis composed of well conserved genes (*mpqA*, *mpqB*, *mpqE*) and alternative genes (*mpqC*, *mpqD/X*, *mpqQ*).

Fig. 2. Characterization of the mPQ biosynthetic pathway. a, Biosynthetic pathways of quinones showing homology of pathways for mPQ_9 in *Nitrospirota* (purple), PQ_9 in the cyanobacterium *Synechocystis sp.* PCC6803 (green) and UQ_8 in the gammaproteobacterium *Escherichia coli* (blue). Biosynthetic steps are numbered, and homologous steps are connected by colored lines. **b-d**, Heterologous complementation experiments using mPQ biosynthesis gene candidates to restore UQ_8 production in *E. coli* mutants lacking key genes for ubiquinone biosynthesis ($\Delta ubiC+mpqC$, $\Delta ubiA+mpqA$, $\Delta ubiX+mpqX$, $\Delta ubiD+mpqD$). **e**, PQ production in *E. coli* $\Delta ubiIFE$ mutants complemented with *mpqQ* from *N. inopinata* as well as PQ and mPQ in *E. coli* $\Delta ubiIFE$ mutants complemented with *mpqQ from N. inopinata* and *mpqE* from other *Nitrospirota*. WT=wild type; vec=empty vector; thick bars represent means and error bars represent standard deviations of the means, n=3-5; AU=arbitrary units. Abbreviations: *Ca.* N. nitrificans (Nnit), *N. moscoviensis* (Nmos), *N. inopinata* (Nino), *L. ferrooxidans* (Lfer), *Ca.* M. noduliformans (Mnod). The numbering of the carbon atoms on the 4-HBA precursor (panel **a**, light grey) defines the nomenclature for all intermediates described in the text. The octaprenyl and nonaprenyl chains are abbreviated with R_8 and R_9, respectively. See Fig. S7-S9 for details on compound identification and quantification. Stars indicate $p < 0.01$ (**), $p < 0.001$ (***), and $p < 0.0001$ (****) for unpaired Student's *t* tests relative to the empty vector.

We verified the mPQ candidate genes by assessing their functions in heterologous complementation assays using *Escherichia coli* mutants deficient in defined steps of UQ biosynthesis. When expressed in the *E. coli* $\Delta ubiC$ mutant, the *mpqC* from *L. ferrooxidans*, *N. nitrificans*, and *M. noduliformans* restored UQ biosynthesis up to wild-type levels (Fig. 2b). Likewise, the *mpqA* homologs from *L. ferrooxidans*, *N. moscoviensis*, and *M. noduliformans* restored UQ biosynthesis in an *E. coli* $\Delta ubiA$ mutant (Fig. 2, S7). Similarly, we observed recovery of UQ levels in *E. coli* $\Delta ubiD$ and $\Delta ubiX$ mutants upon expression of *mpqD/X* from *M. noduliformans* (*mpqD/X* do not occur in *Nitrospira* and *Leptospirillum* spp.; Fig. 2). Expression of the *plqQ* homolog (*mpqQ*) from *N. inopinata* in an *E. coli* $\Delta ubiIF$ mutant yielded PQ_8 and mPQ_8 (Fig. S8b-g). The $\Delta ubiIFE$ strain, in which the *E. coli ubiE* gene was additionally deleted, showed a strong increase in the amount of PQ_8 and the disappearance of mPQ_8

(Fig. 2e, S8b-g). Finally, expression of *mpqE* from *L. ferrooxidans*, *N. moscoviensis*, and *Ca.* M.

noduliformans in the Δ*ubiIFE* strain led to the accumulation of mPQ$_8$ (Fig. 2e, S8c).

Based on these heterologous expression and isotope labeling experiments, we reconstructed a

tentative mPQ biosynthetic pathway that shares homology with the UQ and PQ pathways (Fig. 2). The

ring precursor 4-HBA is generated from chorismate by MpqC and alternative enzymes, followed by

prenylation of 4-HBA by MpqA and decarboxylation by MpqD/X. The following hydroxylation step at

C1 is unresolved, but observations from *Pseudomonadota* indicate that a large diversity of

benzoquinone C1 hydroxylases exist in nature (*36*, *52–54*). Finally, methylations are introduced at C5

and C6 by MpqQ and at C2 by MpqE.

Distribution and function of mPQ

Analysis of mPQ biosynthesis proteins in a representative selection of high- and medium-quality

genomes and metagenome-assembled genomes revealed that mPQ is present in all aerobic lineages of

Nitrospirota (*n*=85), but not found outside this phylum (*n*=482). A few early-branching lineages of

Nitrospirota, which are anaerobes using the MK$_{mqn}$ pathway, are devoid of mPQ biosynthesis proteins

(Fig. 3). Since mPQ is the only respiratory quinone found in aerobic *Nitrospirota*, it is likely involved

in the ETC used for aerobic respiration (*41–43*), and the structural similarity between mPQ and UQ/PQ

suggests that mPQ has a high redox potential. Since *Nitrospirota* grow slowly and to low cell densities,

mPQ could not be isolated in quantities required for redox potential measurements. We therefore

calculated the redox potential of mPQ, UQ, and PQ using density functional theory (*55*). For a given

biologically relevant prenyl chain length, the calculated redox potential of mPQ (E^{0}Q/H2Q = 517±8

mV) is lower than that of PQ (551±8 mV) but higher than that of UQ (480±8 mV; Table S3).

Furthermore, all HPQs are described by significantly higher calculated redox potentials than the LPQ

MK (364±8 mV), confirming the validity of our computational approach. Calculations for simple 1,4-

benzoquinones indicate that redox potentials decrease by ~50 mV per methyl or methoxy group, with

methoxy additions having the larger effect, which explains the higher potential of mPQ (trimethyl)

relative to UQ (dimethoxy, methyl). These functional group combinations may reflect redox tuning of

HPQs to specific components of the ETC in *Cyanobacteriota*, *Nitrospirota*, and *Pseudomonadota*. Due

to the tight coupling of redox potentials of quinones to other ETC components (e.g., iron sulfur clusters

of Rieske proteins and hemes) (*26*), we infer that aerobic *Nitrospirota* have high potential ETCs. Indeed,

we find that Rieske proteins of aerobic *Nitrospirota* contain the 'SY' motif (Table S4) characteristic for

Rieske/cytochrome *b* complexes adapted to interact with HPQs in high potential ETCs (*56*). High

potential ETCs would be advantageous for minimizing ROS generation and maximizing proton motive

force (*26*) in the low energy-yielding chemoautotrophic metabolisms of aerobic *Nitrospirota*. Since the

use of HPQs requires adaptation of the entire ETC to higher redox potential (*26*), such a decisive step

may have been linked to a major event, such as Earth's surface oxygenation (*24*, *26*, *57*).

215
Fig. 3. Phylogenetic tree of bacteria showing the occurrence of respiratory quinones. Quinones with high
redox potential (UQ, PQ, mPQ) occur only in aerobic *Nitrospirota*, *Pseudomonadota*, and *Cyanobacteriota*. Low
potential quinones occur in anaerobic *Nitrospirota* (MK), some *Pseudomonadota* (MK), and all *Cyanobacteriota*
(PhQ). Asterisks indicate strains in which presence of mPQ has been verified experimentally. See Fig. S11-13 for
detailed trees. The maximum-likelihood phylogenetic tree was constructed from 120 concatenated single copy
marker proteins (*58*) of 547 isolate genomes and metagenome-assembled genomes, covering all bacterial phyla,
and rooted using the DST group to approximate the bacterial root (*59*). Quinone occurrences were derived from
instrumental analysis of isolates or inferred from the presence of key biosynthesis genes (SI results;

Ancient origin of high-potential quinones

The biochemical similarity of the HPQ biosynthesis pathways suggests a common ancestry. The

mPQ biosynthesis proteins of Nitrospirota are most closely related to homologs from the UQ and PQ

pathways of *Cyanobacteriota* and *Pseudomonadota*, which is unexpected given the phylogenetic

distance between these phyla (Fig. 3). Specifically, the phylogenies of the key prenyltransferases and

decarboxylases exhibit a consistent tree topology, with HPQ proteins being monophyletic relative to

homologous proteins of the LPQ biosynthesis pathways (Fig. 4; see SI for expanded discussion). HPQ

homologs from *Pseudomonadota* and *Nitrospirota* branch as sister lineages with respect to

Cyanobacteriota. Other proteins of the HPQ pathways (chorismate-pyruvate lyase, decarboxylase co-

factor) generally support this topology, although with lower branch support (Fig. S19-20). These

patterns suggest a single, shared origin of the universal core of HPQ biosynthesis in bacteria.

The distribution of LPQs, HPQs and their associated biosynthetic genes in Bacteria suggests that

HPQ biosynthesis is conserved in all known lineages of *Cyanobacteriota*, *Pseudomonadota*, and

aerobic *Nitrospirota* (Fig. 3, S11-13; see SI for expanded discussion). Conversely, HPQs are not found

in anaerobic *Nitrospirota*, nor in the sister phyla of *Cyanobacteriota*, *Nitrospirota*, or *Pseudomonadota*,

all of which produce LPQs via the MK_{mqn} pathway (Fig. 3; Table S5). Given that *Cyanobacteriota*,

Nitrospirota, and *Pseudomonadota* are paraphyletic, vertical inheritance of HPQ pathways from a

common ancestor is unlikely. Instead, HPQ occurrence and protein phylogenies indicate that an

ancestral HPQ pathway was laterally acquired by stem-group *Cyanobacteriota*, *Pseudomonadota*, and

aerobic *Nitrospirota* from an unknown or extinct donor lineage.

The ancestral HPQ pathway later diversified through changes to the C2, C5, and C6 substituents.

Specifically, C2 methyltransferases are present in all LPQ and HPQ biosynthesis pathways except PQ.

LPQ and HPQ C2 methyltransferases form sister clades (Fig. 4b) and C2 methyltransferases may thus

be as old as the divergence between LPQ and HPQ pathways. Consequently, it is likely that the ancestral

HPQ pathway contained a C2 methyltransferase that was lost prior to the radiation of crown group

Cyanobacteriota (Fig. 4c). Lack of C2 methylation increases the redox potential of PQ (Table S3) and

is essential for the functioning of the oxygen-evolving photosystem II (*60*). Loss of C2-methylation

was likely linked to the evolution of oxygenic photosynthesis and therefore did not occur in

Pseudomonadota and *Nitrospirota*. The evolution of the C5/C6 functional groups is less constrained.

The C5/C6 methyltransferases of the PQ/mPQ pathways are poorly conserved in *Nitrospirota* and

Cyanobacteriota but at least one subgroup of *Nitrospirota* laterally acquired a C5/C6 methyltransferase

from *Cyanobacteriota* (Fig. 4b; see SI discussion). C5/C6 methylation requires a single enzyme,

whereas methoxylation to yield UQ requires at least two enzymes that are specific to *Pseudomonadota*

(*36*, *54*). Thus, the most parsimonious explanation is that the ancestral HPQ was methylated at C5/C6

in addition to C2, i.e., identical to mPQ, and that methoxylation evolved later (Fig. 4c).

Fig. 4. High-potential quinones (HPQ) share a single origin predating the great oxygenation event. a, Phylogenetic trees of HPQ biosynthesis proteins demonstrating that prenyltransferases and decarboxylases of the ubiquinone (UQ, UbiAD), plastoquinone (PQ, PlqAD), and methylplastoquinone (mPQ, MpqAD) pathways form sister clades of the archaeal and bacterial futalosine pathway for biosynthesis of menaquinone (MK, MqnPL). **b,** Phylogenetic trees of quinone C5/C6 (PlqQ, MpqQ) and C2 methyltransferases (UbiE, MpqE), showing a nested topology of C5/C6 methyltransferases and that C2 methyltransferases form a sister lineage of menaquinone-associated methyltransferases (MqnK). Outgroups used for rooting the trees are discussed in the Supplementary Information. Scale bars indicate 0.5 substitutions per site. Open circles indicate ultra-fast bootstrap support ≥0.9. **c**, Conceptual sketch of HPQ evolution and resulting redox potentials. **d**, Timescale of LPQ and HPQ evolution (colors as in panel c) in relation to geochemical changes (evidence for localized O_2 oases (*61–63*), the great oxygenation event, GOE(*7*)) and biological innovations (Archean rapid genetic expansion (*10*), evolution of enzymes protecting against reactive oxygen species (ROS) (*12*), expansion of O_2 reductase diversity (*11*)). Shaded hexagons indicate minimum and maximum estimates of HPQ evolution timescale. Open symbols indicate median ages (colored bars: uncertainty range; quinone symbols: upper/lower estimate) of relevant clades estimated by previous molecular clock analyses (Boden et al. (*12*); Davín et al. (*64*); Fournier et al. (*65*); Oliver et al. (*66*); Magnabosco et al. (*67*); Shih et al. (*68*); Ward et al. (*69*)).

It has been proposed that LPQs were present in the last universal common ancestor or evolved shortly thereafter, given their nearly universal presence in archaea and bacteria (*70*). Of the two LPQ biosynthetic pathways, the MK_{mqn} pathway is considered ancestral to basal archaea and bacteria, whereas the MK_{men} pathway was laterally transferred from bacteria to a subset of archaea (*70*). Homologous proteins suggest that the LPQ and HPQ biosynthetic pathways are evolutionarily related. The HPQ pathways share five homologs with the MK_{mqn} pathway (prenyltransferase, two-component decarboxylase, C2 methyltransferase, kinase) and two with the alternative MK_{men} pathway (prenyltransferase, C2 methyltransferase) (*36*). Our analysis shows that the MK_{mqn} homologs from archaea and bacteria form sister groups to the HPQ proteins, whereas the two homologs of the MK_{men} pathway are more distantly related to both HPQ and MK_{mqn} proteins (Fig. 4, S14-16, and SI discussion). This topology suggests that contrary to previous conclusions (*71*), the HPQ pathways did not descend directly from extant MK pathways.

Instead, the HPQ and MK pathways likely evolved from an ancestral quinone biosynthesis pathway that, like all extant pathways, used a chorismate derivative as precursor. In the case of HPQ, this precursor is prenylated in the second step, whereas prenylation is a late step in MK biosynthesis. The specificity of prenyltransferase for its quinone substrate (*72*) combined with early prenylation in the HPQ pathways may have facilitated evolutionary divergence of the HPQ and MK pathways. Existing machinery from the ancestral quinone biosynthesis pathway such as decarboxylase, C2 methyltransferase, and kinase were then co-opted by these new pathways. The deep phylogenetic divergence between HPQ and LPQ proteins (Fig. S14-16) suggests that the ancestral HPQ pathway

could have emerged before the radiation of bacteria and archaea (*70*) 4.1-3.4 Ga ago in an extinct lineage

coeval to the evolution of the extant LPQ pathways (Fig. 4c-d). Such an early origin of HPQs is not

necessarily linked to aerobic respiration or oxygenic photosynthesis using high-potential ETCs. Instead,

ancestral HPQs could have been involved in different functions, such as oxygen detoxification or a

primordial form of high-potential photosynthesis (*73*, *74*), and only later adopted into high-potential

ETCs used for oxygenic photosynthesis and respiration using oxygen or other high-potential electron

acceptors, such as nitric oxide (*16*, *75*).

Early evolution of aerobic metabolism

The association of HPQ biosynthesis with oxygenic photosynthesis and aerobic respiration in

extant bacteria suggests that these traits became inseparably linked during evolution. The phylogeny of

HPQ biosynthesis proteins therefore allows dating the origin of aerobic metabolisms using HPQs

relative to Earth's oxygenation. Oxygen first accumulated permanently in the atmosphere during the

GOE(*7*) but geochemical tracers suggest oxygen was locally present during the late Archean (*61*, *62*,

76). The likely presence of oxygen during the Archean aligns with the diversification of electron

transport pathways, oxygenases, oxidoreductases, and antioxidant enzymes around 3.3-2.9 Ga (*10–12*,

77), i.e., long before the GOE. Alternative proposals place the emergence of crown group

Cyanobacteriota, oxygenic photosynthesis, and aerobic respiration coeval to, or after, the GOE (*4*, *68*,

78). Regardless of whether oxygenic photosynthesis emerged during the Archean or was coeval to the

GOE, the phylogenetic split between HPQ and LPQ proteins and the presence of the MK_{mqn} pathway

in the non-photosynthetic sister lineages (*Vampirovibrionophyceae*, "*Candidatus* Margulisbacteria",

"*Candidatus* Sericytochromatia"; Fig. 3, 4; Table S5; SI discussion) together indicate that the HPQ

pathway in *Cyanobacteriota* originated from lateral transfer after their divergence from these sister

lineages. Because PQ is central to the functioning of photosystem II in all extant oxygenic

photosynthesizers (*23*, *79*), emergence of PQ biosynthesis was likely tied to the evolution of oxygenic

photosynthesis and thus may have existed before the radiation of crown group *Cyanobacteriota*. This

supports earlier proposals that the extant oxygenic photosynthetic machinery originated in a lineage that

diverged from the non-photosynthetic sister lineages (*78*, *80*), but pre-dated the radiation of crown

group *Cyanobacteriota* (*74*). Collectively, these constraints indicate that HPQs are at least as old as oxygenic photosynthesis by *Cyanobacteriota* and therefore predate the GOE.

Aerobic metabolism preceding the GOE is supported by the near-universal occurrence of aerobic respiration in crown group *Cyanobacteriota* and *Pseudomonadota*. All basal clades of *Cyanobacteriota* and *Pseudomonadota* possess HPQs and are capable of aerobic respiration, with only few late-branching *Pseudomonadota* being obligate anaerobes (Fig. 3, S10, S12-13). Molecular clocks calibrated using cyanobacteriotal fossils place the last common ancestor of crown group *Cyanobacteriota* and the emergence of basal, aerobic *Pseudomonadota* (Magnetococci) around 2.5-3.2 Ga (*10*, *12*, *64–66*, *81*), whereas aerobic *Nitrospirota* may have emerged shortly after the GOE (*64*, *69*). HPQs were thus likely used for aerobic respiration by the time of the radiation of extant *Cyanobacteriota* and *Pseudomonadota*. Given the constraint that HPQs must have been present in stem-group *Cyanobacteriota* and *Pseudomonadota*, the minimum age of extant HPQs is between 2.5-3.2 Ga, whereas the ancestral HPQ pathway may be as old as crown group bacteria (3.4-4.1 Ga; Fig. 4d). We therefore suggest that aerobic respiration with high potential ETCs may have originated up to 800 Ma before oxygen permanently accumulated in the atmosphere during the GOE. Microbial mats could have provided a niche for chemoautotrophs and heterotrophs consuming oxygen provided by *Cyanobacteriota* directly (*31*), preventing escape to the atmosphere.

In modern ecosystems, some aerobic bacteria continue to use LPQ- rather than HPQ-dependent ETCs for aerobic respiration (*22*). In the presence of O_2, reduced HPQs are relatively stable but reduced LPQs are rapidly autoxidized, resulting in the loss of reducing equivalents to O_2 (*25*, *82*). Further, aerobic respiration with LPQs leads to increased formation of deleterious reactive oxygen species (*27*, *83*), requiring energy to be expended on the mitigation of cellular damage, thereby decreasing growth rates (*27*, *84*). Finally, the use of LPQs instead of HPQs for proton pumping by complex I is less efficient (*85*, *86*). Given these drawbacks, one might question why all aerobes have not switched from LPQs to HPQs to use oxygen without these disadvantages. The evolution of HPQs was a complex process closely tied to the evolution of the ETC itself, demanding not only the acquisition of a dedicated pathway for quinone biosynthesis, but also an upshift in redox potential of all other ETC components,

including hemes and iron-sulfur clusters (*26*). These complex requirements may explain why the

evolution of HPQ was successful only once and lateral transfers were rare. Distinct redox-tuning then

led to the extant diversity of HPQ structures and biosynthetic pathways in *Cyanobacteriota*,

Pseudomonadota, and *Nitrospirota*. Two of the three HPQ pathways were later obtained by eukaryotes,

UQ via incorporation of an alphaproteobacterium as the mitochondrion and PQ via incorporation of a

cyanobacterium as the chloroplast (*57*, *87*), while mPQ remained exclusively bacterial. Through their

high potential ETCs, these lineages were able to rise to dominance in modern oxic ecosystems (*31–35*),

as evidenced by the prevalence of HPQs in modern oxic environments (*30*, *88*).

Acknowledgements

We thank Susan Carter for laboratory assistance and Colleen Hansel, Gregory Fournier, and C.S.

Raman for discussions. Linda L. Jahnke and Mary N. Parenteau are thanked for providing biomass of

G. violaceus. Fay-Wey Li is thanked for providing the *A. panamensis* culture and Anja Engel for access

to UPLC-Orbitrap-MS/MS. This work was funded through National Science Foundation grants

1702262 and 1843285 (to A.P.), the Gordon and Betty Moore Foundation (to A.P.), and the Deutsche

Forschungsgemeinschaft grant 441217575 (to F.J.E). This work was also supported by the French

National Research Agency through the grant ANR-21-CE02-0018 (to S.S.A.), and the "Investissements

d'Avenir" program (ANR-15-IDEX-02) through the "Origin of Life" project and the IDEX-IRS 2020

call of the Grenoble-Alpes University (to S.S.A.). Research at MARUM was supported by the Deutsche

Forschungsgemeinschaft via Germany's Excellence Strategy, no. EXC-2077-390741603. T.W.E.

acknowledges funding by the Alexander-von-Humboldt-Stiftung through the Feodor-Lynen-

Fellowship. S.L. acknowledges funding by the Netherlands Organisation for Scientific Research

(016.Vidi.189.050). Research at MIT was otherwise supported by the NASA Astrobiology Institute

Exobiology Program grant number 18-EXXO18-0039. This result is part of a project that has received

funding from the European Research Council (ERC) under the European Union's Horizon 2020

research and innovation program (Grant agreement No. 946150, to J.D.H.). Research at Caltech was

supported by the NASA Astrobiology Institute Exobiology grant no. 80NSSC19K0480. This research

was supported through high-performance computing resources available at the Kiel University

Computing Centre. Redox potential calculations were performed on the Euler cluster operated by the

High-Performance Computing group at ETH Zurich. The authors thank Coline Fernandes from

University Grenoble Alpes for running the ^{1}H and ^{13}NMR experiments.

Author Contributions

F.J.E., F.P., S.S.A., and A.P. designed the research. F.P., L.P., and J.S. performed heterologous

expression experiments. F.J.E., P.B., S.L., E.S., and J.R.L. grew cultures and provided biomass samples.

F.J.E. and V.N. performed isotope labeling experiments. A.P., F.P., S.S.A., K.-U.H., and R.E.S

provided oversight and access to instruments and facilities. F.J.E., T.W.E, J.C., K.W.B., and F.P.

performed chromatographic and mass spectrometric analyses. F.P., A.R., and A.B. performed mPQ

purification and NMR characterization. J.D.H. performed redox potential calculations. F.J.E., S-C.C,

and S.S.A. performed phylogenetic analyses. F.B. performed sequence analyses of Rieske/cytb

complexes. F.J.E. wrote the manuscript with contributions from all authors.

Data Availability

All data are available in the supplementary materials and in the Dryad data repository:

10.5061/dryad.2ngf1vhz1

Code Availability

All code for the calculation of redox potentials is available in the Dryad data repository:

10.5061/dryad.2ngf1vhz1

Competing Interests

The authors declare no competing interests.

References

1.   J. Raymond, The Effect of Oxygen on Biochemical Networks and the Evolution of Complex
Life. *Science* **311**, 1764–1767 (2006).

2.   A. L. Zerkle, S. W. Poulton, R. J. Newton, C. Mettam, M. W. Claire, A. Bekker, C. K. Junium,
409   Onset of the aerobic nitrogen cycle during the Great Oxidation Event. *Nature* **542**, 465–467
410   (2017).

411   3.   K. O. Konhauser, S. V. Lalonde, N. J. Planavsky, E. Pecoits, T. W. Lyons, S. J. Mojzsis, O. J.
412   Rouxel, M. E. Barley, C. Rosìere, P. W. Fralick, L. R. Kump, A. Bekker, Aerobic bacterial pyrite
413   oxidation and acid rock drainage during the Great Oxidation Event. *Nature* **478**, 369–373 (2011).

414   4.   W. W. Fischer, J. Hemp, J. S. Valentine, How did life survive Earth's great oxygenation?
415   *Current Opinion in Chemical Biology* **31**, 166–178 (2016).

416   5.   D. C. Catling, C. R. Glein, K. J. Zahnle, C. P. McKay, Why O_2 Is Required by Complex Life on
417   Habitable Planets and the Concept of Planetary "Oxygenation Time." *Astrobiology* **5**, 415–438
418   (2005).

419   6.   N. Lane, How energy flow shapes cell evolution. *Current Biology* **30**, R471–R476 (2020).

420   7.   A. Bekker, H. D. Holland, P.-L. Wang, D. Rumble, H. J. Stein, J. L. Hannah, L. L. Coetzee, N.
421   J. Beukes, Dating the rise of atmospheric oxygen. *Nature* **427**, 117–120 (2004).

422   8.   G. Izon, G. Luo, B. T. Uveges, N. Beukes, K. Kitajima, S. Ono, J. W. Valley, X. Ma, R. E.
423   Summons, Bulk and grain-scale minor sulfur isotope data reveal complexities in the dynamics
424   of Earth's oxygenation. *Proceedings of the National Academy of Sciences* **119**, e2025606119
425   (2022).

426   9.   B. T. Uveges, G. Izon, S. Ono, N. J. Beukes, R. E. Summons, Reconciling discrepant minor
427   sulfur isotope records of the Great Oxidation Event. *Nat Commun* **14**, 279 (2023).

428   10.   L. A. David, E. J. Alm, Rapid evolutionary innovation during an Archaean genetic expansion.
429   *Nature* **469**, 93–96 (2011).

430   11.   J. Jabłońska, D. S. Tawfik, The evolution of oxygen-utilizing enzymes suggests early biosphere
431   oxygenation. *Nature Ecology & Evolution* **5**, 442–448 (2021).

432   12.   J. S. Boden, K. O. Konhauser, L. J. Robbins, P. Sánchez-Baracaldo, Timing the evolution of
433   antioxidant enzymes in cyanobacteria. *Nat Commun* **12**, 4742 (2021).

434   13.   J. Jabłońska, D. S. Tawfik, Innovation and tinkering in the evolution of oxidases. *Protein Science*
435   **31**, e4310 (2022).

436   14.   M. C. Weiss, F. L. Sousa, N. Mrnjavac, S. Neukirchen, M. Roettger, S. Nelson-Sathi, W. F.
437   Martin, The physiology and habitat of the last universal common ancestor. *Nature Microbiology*
438   **1**, 16116 (2016).

439   15.   J. Castresana, M. Lübben, M. Saraste, D. G. Higgins, Evolution of cytochrome oxidase, an
440   enzyme older than atmospheric oxygen. *EMBO J* **13**, 2516–2525 (1994).

441   16.   A.-L. Ducluzeau, B. Schoepp-Cothenet, R. van Lis, F. Baymann, M. J. Russell, W. Nitschke,
442   The evolution of respiratory O_2/NO reductases: an out-of-the-phylogenetic-box perspective.
443   *Journal of The Royal Society Interface* **11**, 20140196–20140196 (2014).

444   17.   C. Brochier-Armanet, E. Talla, S. Gribaldo, The Multiple Evolutionary Histories of Dioxygen
445   Reductases: Implications for the Origin and Evolution of Aerobic Respiration. *Mol Biol Evol* **26**,
446   285–297 (2009).

18.   F. L. Sousa, R. J. Alves, M. A. Ribeiro, J. B. Pereira-Leal, M. Teixeira, M. M. Pereira, The
superfamily of heme–copper oxygen reductases: Types and evolutionary considerations.
Biochimica et Biophysica Acta (BBA) - Bioenergetics **1817**, 629–637 (2012).

19.   S. Gribaldo, E. Talla, C. Brochier-Armanet, Evolution of the haem copper oxidases superfamily:
a rooting tale. *Trends in Biochemical Sciences* **34**, 375–381 (2009).

20.   R. Murali, J. Hemp, R. B. Gennis, Evolution of quinol oxidation within the heme-copper
oxidoreductase superfamily. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1863**, 148907
(2022).

21.   J. B. Glass, C. E. Elbon, L. D. Williams, Something old, something new, something borrowed,
something blue: the anaerobic microbial ancestry of aerobic respiration. *Trends in Microbiology*
31, 135–141 (2023).

22.   M. D. Collins, D. Jones, Distribution of Isoprenoid Quinone Structural Types in Bacteria and
Their Taxonomic Implications. **45**, 40 (1981).

23.   B. Nowicka, J. Kruk, Occurrence, biosynthesis and function of isoprenoid quinones. *Biochimica*
et Biophysica Acta (BBA) - Bioenergetics **1797**, 1587–1605 (2010).

24.   B. Schoepp-Cothenet, R. van Lis, A. Atteia, F. Baymann, L. Capowiez, A.-L. Ducluzeau, S.
Duval, F. ten Brink, M. J. Russell, W. Nitschke, On the universal core of bioenergetics.
Biochimica et Biophysica Acta (BBA) - Bioenergetics **1827**, 79–93 (2013).

25.   B. Schoepp-Cothenet, C. Lieutaud, F. Baymann, A. Vermeglio, T. Friedrich, D. M. Kramer, W.
Nitschke, Menaquinone as pool quinone in a purple bacterium. *Proceedings of the National*
Academy of Sciences **106**, 8549–8554 (2009).

26.   L. Bergdoll, F. ten Brink, W. Nitschke, D. Picot, F. Baymann, From low- to high-potential
bioenergetic chains: Thermodynamic constraints of Q-cycle function. *Biochimica et Biophysica*
Acta (BBA) - Bioenergetics **1857**, 1569–1579 (2016).

27.   A. Anand, K. Chen, L. Yang, A. V. Sastry, C. A. Olson, S. Poudel, Y. Seif, Y. Hefner, P. V.
Phaneuf, S. Xu, R. Szubin, A. M. Feist, B. O. Palsson, Adaptive evolution reveals a tradeoff
between growth rate and oxidative stress during naphthoquinone-based aerobic respiration.
PNAS **116**, 25287–25292 (2019).

28.   A. Oren, G. M. Garrity, Valid publication of the names of forty-two phyla of prokaryotes.
International Journal of Systematic and Evolutionary Microbiology **71**, 005056 (2021).

29.   P. López-García, L. Eme, D. Moreira, Symbiosis in eukaryotic evolution. *Journal of Theoretical*
Biology **434**, 20–33 (2017).

30.   K. W. Becker, F. J. Elling, J. M. Schröder, J. S. Lipp, T. Goldhammer, M. Zabel, M. Elvert, J.
Overmann, K.-U. Hinrichs, Isoprenoid Quinones Resolve the Stratification of Redox Processes
in a Biogeochemical Continuum from the Photic Zone to Deep Anoxic Sediments of the Black
Sea. *Applied and Environmental Microbiology* **84**, e02736-17 (2018).

31.   N. Finke, R. L. Simister, A. H. O'Neil, S. Nomosatryo, C. Henny, L. C. MacLean, D. E. Canfield,
K. Konhauser, S. V. Lalonde, D. A. Fowle, S. A. Crowe, Mesophilic microorganisms build
terrestrial mats analogous to Precambrian microbial jungles. *Nat Commun* **10**, 4323 (2019).

32. M. Delgado-Baquerizo, A. M. Oliverio, T. E. Brewer, A. Benavent-González, D. J. Eldridge, R.
D. Bardgett, F. T. Maestre, B. K. Singh, N. Fierer, A global atlas of the dominant bacteria found
in soil. *Science* **359**, 320–325 (2018).

33. N. Lang-Yona, J. M. Flores, R. Haviv, A. Alberti, J. Poulain, C. Belser, M. Trainic, D. Gat, H.-
J. Ruscheweyh, P. Wincker, S. Sunagawa, Y. Rudich, I. Koren, A. Vardi, Terrestrial and marine
influence on atmospheric bacterial diversity over the north Atlantic and Pacific Oceans. *Commun*
Earth Environ **3**, 121 (2022).

34. S. Nayfach, S. Roux, R. Seshadri, D. Udwary, N. Varghese, F. Schulz, D. Wu, D. Paez-Espino,
I.-M. Chen, M. Huntemann, K. Palaniappan, J. Ladau, S. Mukherjee, T. B. K. Reddy, T. Nielsen,
E. Kirton, J. P. Faria, J. N. Edirisinghe, C. S. Henry, S. P. Jungbluth, D. Chivian, P. Dehal, E.
M. Wood-Charlson, A. P. Arkin, S. G. Tringe, A. Visel, T. Woyke, N. J. Mouncey, N. N.
Ivanova, N. C. Kyrpides, E. A. Eloe-Fadrosh, A genomic catalog of Earth's microbiomes. *Nat*
Biotechnol **39**, 499–509 (2021).

35. S. Sunagawa, L. P. Coelho, S. Chaffron, J. R. Kultima, K. Labadie, G. Salazar, B. Djahanschiri,
G. Zeller, D. R. Mende, A. Alberti, F. M. Cornejo-Castillo, P. I. Costea, C. Cruaud, F. d'Ovidio,
S. Engelen, I. Ferrera, J. M. Gasol, L. Guidi, F. Hildebrand, F. Kokoszka, C. Lepoivre, G. Lima-
Mendez, J. Poulain, B. T. Poulos, M. Royo-Llonch, H. Sarmento, S. Vieira-Silva, C. Dimier, M.
Picheral, S. Searson, S. Kandels-Lewis, Tara Oceans coordinators, C. Bowler, C. de Vargas, G.
Gorsky, N. Grimsley, P. Hingamp, D. Iudicone, O. Jaillon, F. Not, H. Ogata, S. Pesant, S. Speich,
L. Stemmann, M. B. Sullivan, J. Weissenbach, P. Wincker, E. Karsenti, J. Raes, S. G. Acinas,
P. Bork, E. Boss, C. Bowler, M. Follows, L. Karp-Boss, U. Krzic, E. G. Reynaud, C. Sardet, M.
Sieracki, D. Velayoudon, Structure and function of the global ocean microbiome. *Science* **348**,
1261359–1261359 (2015).

36. S. S. Abby, K. Kazemzadeh, C. Vragniau, L. Pelosi, F. Pierrel, Advances in bacterial pathways
for the biosynthesis of ubiquinone. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1861**,
148259 (2020).

37. H. Daims, "The Family *Nitrospiraceae*" in *The Prokaryotes: Other Major Lineages of Bacteria*
and The Archaea, E. Rosenberg, E. F. DeLong, S. Lory, E. Stackebrandt, F. Thompson, Eds.
(Springer, Berlin, Heidelberg, 2014; https://doi.org/10.1007/978-3-642-38954-2_126), pp. 733–
749.

38. H. Yu, G. L. Chadwick, U. F. Lingappa, J. R. Leadbetter, Comparative Genomics on Cultivated
and Uncultivated Freshwater and Marine "*Candidatus* Manganitrophaceae" Species Implies
Their Worldwide Reach in Manganese Chemolithoautotrophy. *mBio* **13**, e03421-21 (2022).

39. H. Daims, S. Lücker, M. Wagner, A New Perspective on Microbes Formerly Known as Nitrite-
Oxidizing Bacteria. *Trends in Microbiology* **24**, 699–712 (2016).

40. G. M. Garrity, J. G. Holt, "Phylum BVIII. Nitrospirae phy. nov." in *Bergey's Manual of*
Systematic Bacteriology: Volume One: The Archaea and the Deeply Branching and
Phototrophic Bacteria (Springer New York, New York, NY, 2001; https://doi.org/10.1007/978-
0-387-21609-6_25), pp. 451–464.

41. H. Yu, J. R. Leadbetter, Bacterial chemolithoautotrophy via manganese oxidation. *Nature* **583**,
453–458 (2020).

42. S. Lücker, M. Wagner, F. Maixner, E. Pelletier, H. Koch, B. Vacherie, T. Rattei, J. S. Sinninghe
Damsté, E. Spieck, D. Le Paslier, H. Daims, A *Nitrospira* metagenome illuminates the
physiology and evolution of globally important nitrite-oxidizing bacteria. *Proceedings of the*
National Academy of Sciences **107**, 13479–13484 (2010).

43.   G. Levicán, M. J. Gómez, R. Chávez, O. Orellana, M. Moreno-Paz, V. Parro, Comparative
532   Genomic Analysis Reveals Novel Facts about *Leptospirillum* spp. Cytochromes. *MIP* **22**, 94–
533   104 (2012).

534   44.   Y. A. Frank, V. V. Kadnikov, A. P. Lukina, D. Banks, A. V. Beletsky, A. V. Mardanov, E. I.
535   Sen'kina, M. R. Avakyan, O. V. Karnachuk, N. V. Ravin, Characterization and Genome
536   Analysis of the First Facultatively Alkaliphilic *Thermodesulfovibrio* Isolated from the Deep
537   Terrestrial Subsurface. *Frontiers in Microbiology* **7** (2016).

538   45.   S. Zecchin, R. C. Mueller, J. Seifert, U. Stingl, K. Anantharaman, M. von Bergen, L. Cavalca,
539   M. Pester, Rice Paddy *Nitrospirae* Carry and Express Genes Related to Sulfate Respiration:
540   Proposal of the New Genus "*Candidatus* Sulfobium." *Applied and Environmental Microbiology*
541   **84** (2018).

542   46.   F. J. Elling, T. W. Evans, V. Nathan, J. D. Hemingway, J. J. Kharbush, B. Bayer, E. Spieck, F.
543   Husain, R. E. Summons, A. Pearson, Marine and terrestrial nitrifying bacteria are sources of
544   diverse bacteriohopanepolyols. *Geobiology* **20**, 399–420 (2022).

545   47.   R. Sadre, C. Pfaff, S. Buchkremer, Plastoquinone-9 biosynthesis in cyanobacteria differs from
546   that in plants and involves a novel 4-hydroxybenzoate solanesyltransferase. *Biochemical Journal*
547   **442**, 621–629 (2012).

548   48.   C. Pfaff, N. Glindemann, J. Gruber, M. Frentzen, R. Sadre, Chorismate Pyruvate-Lyase and 4-
549   Hydroxy-3-solanesylbenzoate Decarboxylase Are Required for Plastoquinone Biosynthesis in
550   the Cyanobacterium *Synechocystis* sp. PCC6803. *J. Biol. Chem.* **289**, 2675–2686 (2014).

551   49.   Y. Sakuragi, D. A. Bryant, "Genetic Manipulation of Quinone Biosynthesis in Cyanobacteria"
552   in *Photosystem I: The Light-Driven Plastocyanin:Ferredoxin Oxidoreductase*, J. H. Golbeck,
553   Ed. (Springer Netherlands, Dordrecht, 2006; https://doi.org/10.1007/978-1-4020-4256-
554   0_15)*Advances in Photosynthesis and Respiration*, pp. 205–222.

555   50.   P. T. Lee, A. Y. Hsu, H. T. Ha, C. F. Clarke, A C-methyltransferase involved in both ubiquinone
556   and menaquinone biosynthesis: isolation and identification of the *Escherichia coli ubiE* gene.
557   *Journal of Bacteriology* **179**, 1748–1754 (1997).

558   51.   Z. Cheng, S. Sattler, H. Maeda, Y. Sakuragi, D. A. Bryant, D. DellaPenna, Highly Divergent
559   Methyltransferases Catalyze a Conserved Reaction in Tocopherol and Plastoquinone Synthesis
560   in Cyanobacteria and Photosynthetic Eukaryotes. *The Plant Cell* **15**, 2343–2356 (2003).

561   52.   L. Pelosi, A.-L. Ducluzeau, L. Loiseau, F. Barras, D. Schneider, I. Junier, F. Pierrel, Evolution
562   of Ubiquinone Biosynthesis: Multiple Proteobacterial Enzymes with Various Regioselectivities
563   To Catalyze Three Contiguous Aromatic Hydroxylation Reactions. *mSystems* **1**, e00091-16
564   (2016).

565   53.   H. Nagatani, Y. Mae, M. Konishi, M. Matsuzaki, K. Kita, F. Daldal, K. Sakamoto, UbiN, a novel
566   *Rhodobacter capsulatus* decarboxylative hydroxylase involved in aerobic ubiquinone
567   biosynthesis. *FEBS Open Bio* **13**, 2081–2093 (2023).

568   54.   K. Kazemzadeh, L. Pelosi, C. Chenal, S.-C. Chobert, M. Hajj Chehade, M. Jullien, L. Flandrin,
569   W. Schmitt, Q. He, E. Bouvet, M. Jarzynka, N. Varoquaux, I. Junier, F. Pierrel, S. S. Abby,
570   Diversification of Ubiquinone Biosynthesis via Gene Duplications, Transfers, Losses, and
571   Parallel Evolution. *Molecular Biology and Evolution* **40**, msad219 (2023).

55. M. T. Huynh, C. W. Anson, A. C. Cavell, S. S. Stahl, S. Hammes-Schiffer, Quinone 1 e⁻ and 2
e⁻/2 H⁺ Reduction Potentials: Identification and Analysis of Deviations from Systematic Scaling
Relationships. *J. Am. Chem. Soc.* **138**, 15903–15910 (2016).

56. F. ten Brink, B. Schoepp-Cothenet, R. van Lis, W. Nitschke, F. Baymann, Multiple Rieske/cyt*b*
complexes in a single organism. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1827**,
1392–1406 (2013).

57. M. Degli Esposti, A Journey across Genomes Uncovers the Origin of Ubiquinone in
Cyanobacteria. *Genome Biol Evol* **9**, 3039–3053 (2017).

58. D. H. Parks, M. Chuvochina, C. Rinke, A. J. Mussig, P.-A. Chaumeil, P. Hugenholtz, GTDB:
an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank
normalized and complete genome-based taxonomy. *Nucleic Acids Research* **50**, D785–D794
(2022).

59. G. A. Coleman, A. A. Davín, T. A. Mahendrarajah, L. L. Szánthó, A. Spang, P. Hugenholtz, G.
J. Szöllősi, T. A. Williams, A rooted phylogeny resolves early bacterial evolution. *Science* **372**,
eabe0511 (2021).

60. L. Stutts, S. Latimer, Z. Batyrshina, G. Dickinson, H. Alborn, A. K. Block, G. J. Basset, The
evolution of strictly monofunctional naphthoquinol C-methyltransferases is vital in
cyanobacteria and plastids. *The Plant Cell*, koad202 (2023).

61. B. Eickmann, A. Hofmann, M. Wille, T. H. Bui, B. A. Wing, R. Schoenberg, Isotopic evidence
for oxygenated Mesoarchaean shallow oceans. *Nature Geosci* **11**, 133–138 (2018).

62. F. Ossa Ossa, A. Hofmann, J. E. Spangenberg, S. W. Poulton, E. E. Stüeken, R. Schoenberg, B.
Eickmann, M. Wille, M. Butler, A. Bekker, Limited oxygen production in the Mesoarchean
ocean. *Proc Natl Acad Sci USA* **116**, 6647–6652 (2019).

63. S. L. Olson, L. R. Kump, J. F. Kasting, Quantifying the areal extent and dissolved oxygen
concentrations of Archean oxygen oases. *Chemical Geology* **362**, 35–43 (2013).

64. A. A. Davín, B. J. Woodcroft, R. M. Soo, B. Morel, R. Murali, D. Schrempf, J. Clark, B.
Boussau, E. R. R. Moody, L. L. Szánthó, E. Richy, D. Pisani, J. Hemp, W. Fischer, P. C. J.
Donoghue, A. Spang, P. Hugenholtz, T. A. Williams, G. J. Szöllősi, An evolutionary timescale
for Bacteria calibrated using the Great Oxidation Event. bioRxiv [Preprint] (2023).
https://doi.org/10.1101/2023.08.08.552427.

65. G. P. Fournier, K. R. Moore, L. T. Rangel, J. G. Payette, L. Momper, T. Bosak, The Archean
origin of oxygenic photosynthesis and extant cyanobacterial lineages. *Proceedings of the Royal
Society B: Biological Sciences* **288**, 20210675 (2021).

66. T. Oliver, P. Sánchez-Baracaldo, A. W. Larkum, A. W. Rutherford, T. Cardona, Time-resolved
comparative molecular evolution of oxygenic photosynthesis. *Biochimica et Biophysica Acta
(BBA) - Bioenergetics* **1862**, 148400 (2021).

67. C. Magnabosco, K. R. Moore, J. M. Wolfe, G. P. Fournier, Dating phototrophic microbial
lineages with reticulate gene histories. *Geobiology* **16**, 179–189 (2018).

68. P. M. Shih, J. Hemp, L. M. Ward, N. J. Matzke, W. W. Fischer, Crown group Oxyphotobacteria
postdate the rise of oxygen. *Geobiology* **15**, 19–29 (2017).

69. L. M. Ward, D. T. Johnston, P. M. Shih, Phanerozoic radiation of ammonia oxidizing bacteria.
Scientific Reports **11**, 2070 (2021).

70. X.-Y. Zhi, J.-C. Yao, S.-K. Tang, Y. Huang, H.-W. Li, W.-J. Li, The Futalosine Pathway Played
an Important Role in Menaquinone Biosynthesis during Early Prokaryote Evolution. *Genome*
Biology and Evolution **6**, 149–160 (2014).

71. D. A. Ravcheev, I. Thiele, Genomic Analysis of the Human Gut Microbiome Suggests Novel
Enzymes Involved in Quinone Biosynthesis. *Front. Microbiol.* **7** (2016).

72. C. A. Cotrim, A. Weidner, N. Strehmel, T. B. Bisol, D. Meyer, W. Brandt, L. A. Wessjohann,
M. T. Stubbs, A Distinct Aromatic Prenyltransferase Associated with the Futalosine Pathway.
ChemistrySelect **2**, 9319–9325 (2017).

73. W. W. Fischer, J. Hemp, J. E. Johnson, Manganese and the Evolution of Photosynthesis. *Orig*
Life Evol Biosph **45**, 351–357 (2015).

74. T. Cardona, J. W. Murray, A. W. Rutherford, Origin and Evolution of Water Oxidation before
the Last Common Ancestor of the Cyanobacteria. *Molecular Biology and Evolution* **32**, 1310–
1328 (2015).

75. A.-L. Ducluzeau, R. van Lis, S. Duval, B. Schoepp-Cothenet, M. J. Russell, W. Nitschke, Was
nitric oxide the first deep electron sink? *Trends in Biochemical Sciences* **34**, 9–15 (2009).

76. R. Riding, P. Fralick, L. Liang, Identification of an Archean marine oxygen oasis. *Precambrian*
Research **251**, 232–237 (2014).

77. K. M. Kim, T. Qin, Y.-Y. Jiang, L.-L. Chen, M. Xiong, D. Caetano-Anollés, H.-Y. Zhang, G.
Caetano-Anollés, Protein Domain Structure Uncovers the Origin of Aerobic Metabolism and the
Rise of Planetary Oxygen. *Structure* **20**, 67–76 (2012).

78. W. W. Fischer, J. Hemp, J. E. Johnson, Evolution of Oxygenic Photosynthesis. *Annual Review*
of Earth and Planetary Sciences **44**, 647–683 (2016).

79. M. Havaux, Plastoquinone In and Beyond Photosynthesis. *Trends in Plant Science* **25**, 1252–
1265 (2020).

80. R. M. Soo, J. Hemp, D. H. Parks, W. W. Fischer, P. Hugenholtz, On the origins of oxygenic
photosynthesis and aerobic respiration in Cyanobacteria. *Science* **355**, 1436–1440 (2017).

81. F. Garcia-Pichel, J. Lombard, T. Soule, S. Dunaj, S. H. Wu, M. F. Wojciechowski, Timing the
Evolutionary Advent of Cyanobacteria and the Later Great Oxidation Event Using Gene
Phylogenies of a Sunscreen. *mBio* **10**, e00561-19 (2019).

82. S. Korshunov, J. A. Imlay, Detection and Quantification of Superoxide Formed within the
Periplasm of *Escherichia coli*. *Journal of Bacteriology* **188**, 6326–6334 (2006).

83. A. Nitzschke, K. Bettenbrock, All three quinone species play distinct roles in ensuring optimal
growth under aerobic and fermentative conditions in *E. coli* K12. *PLoS One* **13**, e0194699
(2018).

84. M. Khademian, J. A. Imlay, How Microbes Evolved to Tolerate Oxygen. *Trends in*
Microbiology **29**, 428–440 (2021).

85.　E. Nakamaru-Ogiso, M. Narayanan, J. A. Sakyiama, Roles of semiquinone species in proton
pumping mechanism by complex I. *J Bioenerg Biomembr* **46**, 269–277 (2014).

86.　U. Brandt, A two-state stabilization-change mechanism for proton-pumping complex I.
Biochimica et Biophysica Acta (BBA) - Bioenergetics **1807**, 1364–1369 (2011).

87.　W. F. Martin, S. Garg, V. Zimorski, Endosymbiotic theories for eukaryote origin. *Philosophical*
Transactions of the Royal Society B: Biological Sciences **370**, 20140330 (2015).

88.　A. Hiraishi, M. Iwasaki, T. Kawagishi, N. Yoshida, T. Narihiro, K. Kato, Significance of
Lipoquinones as Quantitative Biomarkers of Bacterial Populations in the Environment.
Microbes and Environments **18**, 89–93 (2003).