Sex Dimorphism in Expression of Immune Response Genes in Drosophila

MD Mursalin Khan[1*] and Rita M. Graze[1]

[1]Department of Biological Sciences, Auburn University, Alabama, USA.

[*]Corresponding author email: mursalinkhan0018@gmail.com, mzk0086@auburn.edu

Mailing Address: [1]101 Rouse Life Sciences Building, 120 W Samford Ave, Auburn, AL 36849.

Abstract

Sex dimorphism in immunity is commonly observed in a wide variety of taxa and is thought to arise from

fundamental life history differences between females and males. In *Drosophila melanogaster*, infection

with different pathogens typically results in different modes of immune sex dimorphism, with male- or

female-bias, likely due to specific disease-causing mechanisms of pathogens and host-pathogen

interactions. Studies showed that some pathways, such as IMD and Toll, can explain these sex-dimorphic

immune responses in Drosophila. However, it is unclear if sex differences in the immune response

observed in *D. melanogaster* are conserved, even in closely related species. One window into identifying

conserved and evolving sex differences in the immune response is to examine the sex-differential

expression of immunity-related genes. Here, we aim to understand whether two closely related species,

D. melanogaster and *D. simulans*, show conserved sex dimorphism in innate immunity, focusing on

associated changes in gene expression in response to infection with a gram-negative bacterium,

Providencia rettgeri. Survival, bacterial load, and bacterial load upon death (BLUD) were investigated to

assess overall sex differences. *D. melanogaster* females and males differed significantly in survival,

whereas *D. simulans* did not. Enrichment analyses revealed that both sexes and species upregulate genes

involved in similar immune-related biological processes, but downregulated groups differed. We

identified conserved sex differential gene expression of genes in the bacterial infection response

pathways IMD, Toll, Jak/STAT, their regulators, and other immune-related gene classes (e.g., BOMs), as

well as sex and species differences. In *D. melanogaster,* the effector antimicrobial peptides (AMPs)

regulated by IMD were more highly upregulated relative to *D. simulans* in both sexes. Moreover, *D.

melanogaster* females uniquely initiated high levels of gene expression that were involved in negative

feedback mechanisms that controlled the overstimulation of IMD. Genes in the Toll pathway were also

sex-differentially expressed with a higher level of upregulation in *D. melanogaster*. Remarkably,

comparing expression across species, we find that *D. simulans* likely employs both the conventional

peptidoglycan recognition-driven PRR-SPE-Spz pathway and the microbial protease recognition-based

Psh-dependent activation of Toll; in contrast, *D. melanogaster* appears to solely rely on the PRR-SPE-Spz

pathway in this context. In summary, our findings indicate that sex differences are conserved in both

species for the majority of upregulated genes, while downregulation patterns and specific gene subsets

show notable differences between sexes or species.

Keywords: Sex dimorphism, sex-biased gene expression, innate immunity, *Drosophila melanogaster*,

Drosophila simulans, Toll, IMD, BOMs, PGRPs, Bacterial infection.

42 1. Introduction

Sex dimorphism influences a diverse set of biological processes, affecting nearly all aspects of an

organism's biology (Morrow 2015; Millington and Rideout 2018; Lopes-Ramos *et al.* 2020). Sex

dimorphism of easily observable characteristics such as external coloration or anatomy has been widely

described and studied, but dimorphism also extends to less readily apparent traits, including immune

responses, infection outcomes, and disease prognosis (Carton et al. 1992; Fischer et al. 2015; Morrow

2015; McDonald et al. 2021; Wilkinson et al. 2022). Sex-biased immune responses have been observed

at different stages of life history and may also be present in local tissue-level responses such as in the

brain, intestinal layer, and hemolymph (Mackenzie et al. 2011; Deng and Jasper 2016; Millington and

Rideout 2018; Belmonte et al. 2020; Khodursky et al. 2020; Wilkinson et al. 2022). The magnitude of sex

differences may be mediated by the involvement of distinct immune compartments, such as hemocytes,

melanization, epithelial immunity, and systemic immunity, depending on the pathogen and route of

infection (Belmonte et al. 2020; Yu et al. 2022). At a mechanistic level, this may be explained by

regulatory differences in key pathways, such as the Toll and IMD pathways, or more broadly by sex

dimorphism of key tissues (Duneau et al. 2017b; Vincent and Dionne 2021). Even though immunological

differences between sexes are observed frequently and impact fundamental mechanisms underlying

immunity, these differences have received relatively little attention, including in model organisms and

human studies (Morrow 2015; Klein and Flanagan 2016; Belmonte et al. 2020; Wilkinson et al. 2022).

Sex is a central axis of variation, explaining a significant amount of variation within species and

differences between them (Bateman, 1948; Darwin, 1871; Lande, 1980). Sex differences have been

observed in the intensity of the response against pathogens, specific outcomes of infections, and

susceptibility, resistance, and tolerance to pathogens (Belmonte et al., 2020; Fischer et al., 2015; Klein &

Flanagan, 2016). The evolution of differences in the immune response between males and females is

most frequently attributed to differences in reproductive strategies and life history traits (Lande 1980;

Zuk 2009; Morrow 2015). For instance, females typically invest more in offspring production and care,

leading to the prediction that more robust immune responses will be associated with higher fitness in

females (Lande 1980; Zuk 2009; Morrow 2015). Conversely, males may invest more in traits that

enhance competitiveness, potentially at the expense of immune function (Lande 1980; Zuk 2009;

Morrow 2015). Behavioral differences between males and females may also impact the frequency,

intensity, and timing of exposure to various challenges such as viral, bacterial, fungal, and chemical,

which could contribute to the evolution of sex differences or environmentally induced sex differences in

immunity observed in natural contexts (Oertelt-Prigione 2012; Asahina 2018). In addition, other

processes may intersect with sex dimorphism in immunity, contributing to variation within populations.

For example, emerging work has detected sex differences in the aging of immune tissues, immune

senescence, and inflammation (Mondal and Rai 1999; Metcalf et al. 2020; Abancens et al. 2020;

Shepherd et al. 2021). Differences in food intake and diet are also observed between males and females

and could modulate immunity in a sex-differential fashion (Oertelt-Prigione 2012; Asahina 2018; Rai et

al. 2023).

Hormone signaling pathways, including sex hormones, have a role in the transcriptional regulation of

the immune genes (Graze et al. 2018; Gal-Oz et al. 2019; Shepherd et al. 2021). Moreover, previous

studies found that steroid hormones can directly regulate the function of the immune cells, such as the

absence of ecdysone led to the inability of the macrophage-like cells to phagocytize bacteria in

Drosophila (Regan *et al.* 2013). The peptide hormone IIS/TOR signaling disruption also reported the sex-

differential immune gene expression in Drosophila (Graze *et al.* 2018). However, hormone signaling

likely does not fully explain sex dimorphism in immune responses, as sex differences in immune

response have been observed even at sexually immature life stages (Klein and Flanagan 2016; Shepherd

et al. 2021). Sex chromosomes and dosage compensation mechanisms can also contribute to the

differential expression of immune genes as well as the function of immune cells (Scofield et al. 2008;

Libert et al. 2010). For instance, in mammals, the distinct composition of sex chromosomes can lead to

female-specific mosaicism due to cell-specific X chromosome inactivation and may influence the

expression of immune genes (Klein and Flanagan 2016; Pinheiro and Heard 2017). In humans, significant

proportions of primary B lymphocytes, monocytes, and plasmacytoid dendritic cells express TLR7 on

both X chromosomes by escaping X inactivation of XX individuals (Libert et al. 2010; Souyris et al. 2018;

Heintze 2018). Overall, multiple aspects of sex determination and development impact sex dimorphism

in immunity and often interact in complex ways.

Drosophila is a crucial model in the study of development and physiology and is growing as a model for

sex differences in body size, aging, stress responses, and disease progression (Rideout et al. 2015;

Millington and Rideout 2018; Belmonte et al. 2020). As a model organism for innate immunity,

Drosophila showcases sex differences in both the cellular and humoral responses (Reviewed in

(Belmonte *et al.* 2020)). These two central components contribute to sex-specific immune responses

depending on the type of pathogen and the context of the infection (Hoffmann 2003; Lazzaro and Little

2009; Millington and Rideout 2018; Belmonte *et al.* 2020). Thus, *D. melanogaster* exhibits sexual

dimorphism in the outcome of infectious diseases (Vincent and Dionne 2021; Rai *et al.* 2023; Khan and

Graze 2024).

The Drosophila immune system comprises various cellular and humoral components that work together

to recognize, signal proliferation of, and eliminate pathogens (Hoffmann 2003; Vlisidou and Wood 2015;

Yu *et al.* 2022). One of the key cellular components of the *Drosophila* innate immune system is a

specialized cell type called the plasmatocyte, the functional equivalent of mammalian

macrophages/monocytes, which accounts for around 95% of the circulating hemocytes (Evans *et al.*

2003; Vlisidou and Wood 2015; Banerjee *et al.* 2019). Plasmatocytes can engulf and destroy invading

microorganisms and maintain communication with the humoral response, such as response to

antimicrobial peptides (AMPs) binding (Evans *et al.* 2003; Vlisidou and Wood 2015; Banerjee *et al.* 2019).

The rest of the hemocytes (5%) primarily consist of crystal cells that produce phenol oxidase (PO) for

melanization, primarily as an immune response involved in wound healing (Evans *et al.* 2003; Dudzic *et*

al. 2015). Finally, upon pathogenic attack, fruit flies generate lamellocytes, which are larger than

plasmatocytes and specialize in the encapsulation of large pathogens such as wasp eggs (Evans *et al.*

2003; Vlisidou and Wood 2015).

In the humoral response, the most critical aspect of the *Drosophila* innate immune response is the

consistent production of small peptides called antimicrobial peptides (AMPs) and their assistance in

recruiting additional immune cells such as plasmatocytes (Lai and Gallo 2009; Vlisidou and Wood 2015;

Yu *et al.* 2022). Upon immune signal activation, AMPs are produced in the fat body and released into the

hemolymph (Tzou *et al.* 2000; Hoffmann 2003). Due to their positive charge, AMPs are attracted to

microbes with negatively charged cell membranes, where they embed in the hydrophobic region and

cause membrane destabilization, resulting in cell death (Lai and Gallo 2009; Lin *et al.* 2020). Moreover,

certain tissues, such as the trachea, midgut, oviduct, spermatheca, ganglia, and a subpopulation of

hemocytes, produce AMPs in response to local infection (Tzou *et al.* 2000; Lai and Gallo 2009). In

general, the level of AMP expression directly reflects the strength of the immune response (Lai and Gallo

2009; Lin *et al.* 2020).

The core response is initiated by pattern recognition receptors (PRRs) that recognize PAMPs, for

example, specific molecular patterns on the surface of microorganisms, and initiate a signaling cascade

that leads to the production of AMPs via the IMD and Toll pathways (Tanji and Ip 2005). The IMD and

Toll pathways are exceptionally well studied in this system (Valanne *et al.* 2011; Myllymäki *et al.* 2014).

They are homologous to the vertebrate TNF and Toll-like pathways, respectively (Tanji and Ip 2005; Yu *et*

al. 2022; Valanne *et al.* 2022). In general, the IMD pathway is activated upon the detection of

peptidoglycan produced by gram-negative bacteria, whereas the Toll pathway responds to the

peptidoglycan of most gram-positive bacteria, and fungal cell wall components such as β-(1,3)-glucan

(Tanji and Ip 2005; Valanne *et al.* 2022). Moreover, the Drosophila STING pathway activates upon

detection of bacterial cyclic dinucleotides (CDNs) to initiate an innate immune response by triggering the

NF-κB transcription factor Relish of IMD pathway to produce AMPs (Martin *et al.* 2018). Of note, a

second cascade besides PRR-based activation, the danger signal cascade, can activate the Toll pathway

(Ligoxygakis *et al.* 2002; Gottar *et al.* 2006; Chamy *et al.* 2008). The danger cascade initiates upon the

detection of the abnormal activity of bacterial or fungal proteases (danger signal) and activates the Toll

pathway by using Clip-Serine Protease Persephone (Psh) via Psh-SPE-Spz or Psh-Protease-Spz (SPE

independent) cascades (Gottar *et al.* 2006; Yamamoto-Hino and Goto 2016; Issa *et al.* 2018; Dudzic *et al.*

2019). Upon activation, each pathway launches a rapid and efficient transcriptional response to

infection via nuclear translocation of NF-κB family transcription factors, Dif and Dorsal (Tanji and Ip

2005; Valanne *et al.* 2022).

Recent studies have reported that *Drosophila* males and females differ in response to localized and

systemic bacterial, fungal, viral, and parasitic infections (Reviewed in (Klein and Flanagan 2016;

Belmonte *et al.* 2020)). In many cases, for example, Kalithea (viral), *Candida albicans* (fungal),

Tubulinosema ratisbonensis (microsporidial), and *Serratia marcescens* (bacterial), there was a bias

toward females in the effectiveness of the immune response as measured by survival (Kutch and

Fedorka 2017; Palmer *et al.* 2018; Franchet *et al.* 2019; Chowdhury *et al.* 2019). At the same time, this

pattern is not consistently observed even among similar types of pathogens; male flies infected with

Beauveria bassiana (fungal), *Providencia alcalifaciens* (bacterial), and *Enterococcus faecalis* (bacterial)

have higher survival rates relative to females (Taylor and Kimbrell 2007; Le Bourg 2011; Duneau *et al.*

2017b; Shahrestani *et al.* 2018). Collectively, these studies, along with others, have also revealed

considerable context dependency. Sex dimorphism in immunity can be influenced by various factors

such as hormonal regulation, environmental conditions, dietary habits, and metabolic processes (Leech

et al. 2019; Shepherd *et al.* 2021; Rai *et al.* 2023). Conditional impacts are especially evident when the

same pathogen shows different directions of sex dimorphism under different experimental conditions,

either in different laboratories or in the same study with different treatments (Rai *et al.* 2023). For

example, two independent studies of *Pseudomonas aeruginosa* infections demonstrate opposite-sex

bias (Vincent and Sharp 2014; Chowdhury *et al.* 2019). Moreover, recent findings suggest that sex-

determination pathway genes such as *transformer (tra)* might influence immune sex dimorphism in

Drosophila (Khan and Graze 2024).

Several recent reviews have highlighted a significant gap in our understanding of sex differences in

immunity, particularly in adult animals and in the context of evolutionary biology (Beery and Zucker

2011; Belmonte *et al.* 2020). Further, there is a relative lack of comparative studies to understand which

aspects of sex dimorphism in the immune response are conserved. Comparative studies are also crucial

for understanding the evolutionary forces influencing sex differences across the species. For instance,

comparative studies of closely related species can help to expound whether observed sex differences

are lineage-specific and/or adaptive (Hill-Burns and Clark 2010; Khodursky *et al.* 2020; Nanni *et al.*

2023).

In *D. melanogaster*, both the IMD and Toll pathways play pivotal roles in immune dimorphism (Duneau

et al. 2017b; Vincent and Dionne 2021). The Toll pathway and its regulators may be responsible for a

significant portion of immune sex dimorphism in *D. melanogaster* (Duneau *et al.* 2017b). In addition,

regulation of the IMD pathway also plays a role in sex differences (Vincent and Dionne 2021). While this

is a growing area of study, there are many remaining questions regarding the complex molecular basis

of sex dimorphism in immune responses and the degree to which the role of the Toll and IMD pathways

in sex dimorphism is conserved. The intricate crosstalk among evolutionarily conserved immune

pathways, including Toll, IMD, Jak/STAT, and JNK, introduces further complexities to the investigation of

sex dimorphism (Dudzic *et al.* 2019; Tafesh-Edwards and Eleftherianos 2020; Alejandro *et al.* 2022).

Overall, a comprehensive understanding of sexual dimorphism in immunity can provide valuable insights

into the complex interplay among evolution, diversity, immunity, and sex, which also has important

implications for the treatment and prevention of disease.

To comprehensively characterize sex dimorphism in immunity in *Drosophila*, we conducted a

comparative study, profiling outcomes of infections and gene expression in unmated-adult females and

males in two species, *D. melanogaster* and *D. simulans*. This study allows a direct comparison of the

response to infection in *D. melanogaster*, where most of the immunological studies have been

completed, and *D. simulans*, a relatively well-studied and closely related species of the same sub-group

(Figure 1A) (Bhutkar *et al.* 2007; Paris *et al.* 2013). In addition, to the best of our knowledge, this is the

first time bacterial infection in *D. simulans* has been reported for both sexes, and it is the first

comparative immune response study with *D. melanogaster.* This work identifies sex differences that are

in common, as well as divergence of sex differences in immune gene expression, laying the groundwork

for an improved understanding of the evolutionary context of sex dimorphism in the immune response.

2. Methods and Materials

2.1 Fruit fly stocks and sample collection

D. melanogaster and *D. simulans* were reared on a modified version of the Bloomington Drosophila

Stock Center (BDSC) standard cornmeal medium recipe (33 L H_2O, 237 g agar, 825 g dried deactivated

yeast, 1560 g cornmeal, 3300 g dextrose, 52.5 g Tegosept in 270 mL 95% ethanol and 60 mL propionic

acid). Flies were reared at 25°C with a 12:12 light: dark cycle. Density-controlled crosses were set up for

each species, *D. melanogaster* (Canton S, Gift Michelle Arbeitman) and *D. simulans* (Dsim w[501],

14021-0251.195, National Drosophila Species Stock Center, Cornell University); newly emerged progeny

was collected as virgins and separated by sex. Female and male individuals were isolated in standard

vials, and all experimental flies were unmated, adult, and aged 5-9 days prior to infection.

2.2 Experimental treatment

A strain of the gram-negative *Providencia rettgeri* species isolated from wild-caught *D. melanogaster* (*P.*

rettgeri strain: Dmel, Gift Brian Lazzaro) was used as an infectious agent, which is known as moderately

virulent (Juneja and Lazzaro 2009; Galac and Lazzaro 2011; Duneau *et al.* 2017b). In this study,

stationary phase *P. rettgeri* bacterial culture (OD600nm = 1.2) was used to inject bacteria into the

thoracic region (soft spot near the wings) of the fly to introduce around 1200-1500 cells directly into the

hemolymph using the needle prick method (Figure 1B; following (Khalil *et al.* 2015)). For control flies,

sterile phosphate buffer solution (1x PBS) was injected as a wound-only control (Figure 1B).

2.3 Bacterial culture preparation

Glycerol stocks of *P. rettgeri* were stored at -80 °C prior to use. Stocks were used to streak LB-Agar

plates (Teknova, California) and incubated at 37°C for 12 hours to isolate single colonies. A single colony

was then transferred to 5 mL Lysogeny broth LB-Broth (Teknova, California) and incubated at 37°C at

220 rotation per minute (rpm) for 9-10 hours. Stationary phase bacteria were centrifuged at 4000 rpm

for 3 minutes to collect bacterial pellets. Pellets were washed with sterile 1x PBS using pipette aspiration

(5 times) to remove residual LB Broth. After five washes, the pellet was resuspended with sterile 1x PBS

to achieve OD600 nm of 1.2.

2.4 Bacterial injection of flies

225   The flies were anesthetized by CO_2 during the injection. The injections were made in the thoracic region

226   beside the wings in the soft spot using the needle pricking techniques previously described by Khalil et

227   al. 2015 to introduce approximately 1200-1500 viable bacteria into each fly. This approach introduces

228   bacteria directly into the hemolymph. The fly vials were kept horizontal to allow recovery until the flies

229   were completely active.

2.5 Survival rate

For each sample type, the survival assay was performed in 5 replicates, each of which contained 30 flies.

Survival was scored at 2, 6, 12, 18, 24, and 36 hours after bacterial injection. Flies were scored as dead if

the fly was on its back and was not moving or could not turn by itself (Duneau *et al.* 2017a). Scoring

ceased when all the flies were dead in any of the experimental groups or after the 36-hour time point.

Survival analysis was conducted in R using the package "survival" (Therneau 2023). Kaplan-Meier

survival curves were generated to visualize the survival proportion. The Cox regression was performed

to determine the statistical difference among experimental groups (Therneau and Grambsch 2000).

2.6 Bacterial load time series

Bacterial load was assessed for each sample in six replicates, with each replicate consisting of one fly at

each time point. The bacterial load was determined at 4, 8, 12, 16, 20, and 24 hours after injections,

with six replicate measurements following Duneau et al. (Duneau *et al.* 2017a). A single fly per replicate

measurement was homogenized in 200 μL 1X PBS using a sterile pestle. The homogenized sample (10

μL) was transferred to a 96-well plate and diluted with 90 μL 1X PBS. From the first dilution (1/10), 10 μL

of the solution was transferred successively, and serial dilution was carried out 8 times. Subsequently,

20 µL of the dilution $1:10^2$, $1:10^4$, $1:10^6$, and $1:10^8$ were spread on LB agar plates to count CFU. Plates

were incubated at 37 °C for 12-14 hours to count the CFU before gut microbes typically form colonies

that occur at approximately 36 hours (Khalil *et al.* 2015). To test the differences between infected and

control animals for each time point and to get the differences across groups, the linear model was used

(Sparks 2017).

2.7 Bacterial load upon death (BLUD)

The bacterial load upon death was determined for each sample in four replicates, with each replicate

comprising three flies per pool. For the BLUD experiment, flies were determined to be dead flies if the

fly was on its side or back without any movement (Duneau *et al.* 2017a). A single fly was homogenized in

200 µl of 1x PBS within 30 minutes of determining death using a sterile pestle. Serial dilutions and CFU

counts were obtained, as described in the previous section. To find the differences in groups for each

time point, the linear model (CFU ~ Groups) was fit and pairwise differences tested for species and sex

comparisons.

2.8 RNA extraction and prep

Four replicate pools of 5–9 day old flies with 20 flies per pool were collected for each sex and each

group. Only live flies were used for RNA isolation. Replicate pools of wound control and bacteria-

infected flies were flash-frozen in liquid nitrogen 16 hrs post-injection (hpi). The frozen samples were

stored at -80°C prior to total RNA isolation. The 20 flies (per pooled sample) were homogenized entirely

in 700 µL of TRI-Reagent using a pellet pestle motor and RNase-Free pellet pestles in a 2 mL

microcentrifuge tube. Total RNA was extracted using the Direct-zol RNA Miniprep Plus Kit (Zymo

Research) according to the manufacturer's instructions. The TURBO DNase-free kit (Thermo Fisher

Scientific) was used to remove genomic DNA contamination following the manufacturer's instructions.

The concentration was determined using Nanodrop 2.0 (Thermo Fisher Scientific), and RNA integrity

Number (RIN) was measured by using a TapeStation system with an RNA ScreenTape Analysis kit

(Agilent Technologies).

2.9 TagSeq library prep

3' TagSeq was performed at the University of Texas Genomic Sequencing and Analysis Facility, Austin

(UT-GSAF), based on the protocols from Lohman et al. (2016) and Meyer et al. (2011) (Lohman et al.,

2016; Meyer et al., 2011). Libraries were quantified using the Quant-it PicoGreen dsDNA assay

(ThermoFisher) and pooled equally for subsequent size selection at 350-550 bp on a 2% gel using the

Blue Pippin (Sage Science). The final pools were checked for size and quality with the Bioanalyzer High

Sensitivity DNA Kit (Agilent Technologies), and their concentrations were measured using the KAPA SYBR

Fast qPCR kit (Roche). Libraries were sequenced on a NovaSeq 6000 (Illumina), 100 bp single-end reads,

with 5-10 million reads per replicate (File S1).

2.10 Preprocessing and alignment

Single-end 100 bp reads were generated using 3' Tag RNA Seq for each experimental group with 4

replicates each. The total raw reads generated by 3' Tag RNA Seq was between 5.7 and 8.4 million (M)

reads per sample, with an average of 6.99 M reads for *D. melanogaster* and 7.16 M reads for *D. simulans*

(File S1). The quality assessment of raw data was performed by FastQC (Babraham Institute, University

of Cambridge) (Anders 2010). The FastQC analysis showed that all 32 samples had high quality

throughout, and no trimming was performed except the removal of the adapter by the sequencing

facility (Trimmomatic; Bolger et al., 2014).

The reads were aligned against the respective *Drosophila* genomic references (FlyBase Releases:

dmel_r6.47 for *D. melanogaster* and dsim_r2.02 for *D. simulans*) using the read alignment tool STAR

(Gelbart *et al.* 1996; Gramates *et al.* 2022). The alignments were used to generate raw read counts for

each gene in the reference genome annotation by using the "genomeDir", "readFilesIn", and

"outSAMtype" options in STAR (Dobin *et al.* 2013). Gene expression levels were then quantified using

the "quantMode GeneCounts" option of STAR (Dobin *et al.* 2013).

The mean uniquely mapped reads percentage for *D. melanogaster* samples was 74.48%, and for *D.*

simulans samples, it was 76.81%, which is typical for 7 M reads of 3' Tag RNA Seq (File S1) (Ma *et al.*

2019). Upon mapping, raw read counts of 17,633 genes in *D. melanogaster* and 15,266 genes in *D.*

simulans were generated by STAR. Of 17,633 genes of *D. melanogaster* and 15,266 genes of *D. simulans*,

13,588 and 13,010 genes passed the detection cutoff, respectively (File S2-S5). Each experimental

sample passed the quality control analysis prior to normalization and DEG analysis.

An edgeR-limma-voom workflow in R, was used to normalize raw counts (Log2CPM) and fit a cell means

model, with defined contrasts used to determine differential expression between experimental groups,

focusing on sex and species differences in gene expression resulting from infection (Robinson *et al.*

2010; Ritchie *et al.* 2015). Differences were classified as up or downregulated relative to wound control.

The interaction of treatment by sex was tested to identify significant differences between the sexes,

within each species (Law *et al.* 2014; Ritchie *et al.* 2015). A significant treatment by sex interaction term

means that the sexes differ in the way that the gene expression for a gene changes in response to the

infection. Additional tests for interactions were included to investigate species differences. In between-

species comparisons, significant interaction terms for a gene indicate differences between species in the

changes in gene expression observed in comparisons for a single-sex (treatment by species) or in

comparisons between sexes (treatment by sex by species).

2.11 Differential gene expression analysis

Gene expression differences were first evaluated for each species individually to allow analysis of all

expressed annotated genes in a species. Genes with fewer than 2 read counts (summed across samples

within a group) were excluded from the analysis. Expression was estimated as the Log2CPM normalized

counts per gene for *D. melanogaster* (mel) or *D. simulans* (sim), females (F) or males (M), with infection

by *P. rettgeri* (pret) or PBS-only injected controls (PBS). Voom variance modeling was implemented in

limma, with a means model and comparisons between groups tested using contrasts (Law *et al.* 2014;

Ritchie *et al.* 2015). In *D. melanogaster,* the comparisons tested were mel_f_pret = mel_f_pbs and

mel_m_pret = mel_m_pbs, the effect of treatment for each sex, and the interaction of treatment by sex

mel_f_pret - mel_f_pbs = mel_m_pret - mel_m_pbs. Likewise, for the *D. simulans*, the corresponding

tests were conducted. For the enrichment, gene class heatmaps, and pathway analysis in *D. simulans,*

the annotated *D. melanogaster* orthologs were used (dmel_orthologs_in_dsim_Flybase_2017_05; File

S13). An FDR-adjusted *p*-value (*p*-adj) cutoff of 0.05 was used for all tests of differential gene expression.

For comparisons between species, unique ortholog pairs were identified and included in the analysis. If

two putative orthologs were found for one gene, then only one representative was kept for the analysis,

meaning a single representative pair for each set of ambiguous homologs, for which orthology has not

been clearly defined, was included in the analysis. As mentioned above, raw counts per gene were

normalized using edgeR-limma-voom. For the comparison between species, normalization also included

TMM density normalization of Log2CPM transformed counts. The contrast used to test the interaction

of treatment by species in females is dmel_f_pret - dmel_f_pbs = dsim_f_pret - dsim_f_pbs, and the

effect of treatment by species in males is dmel_m_pret - dmel_m_pbs = dsim_m_pret - dsim_m_pbs

(Law *et al.* 2014). Additionally, contrasts were used to test the three-way interaction of treatment by sex

by species ((dmel_f_pret - dmel_f_pbs) - (dmel_m_pret - dmel_m_pbs)) = ((dsim_f_pret - dsim_f_pbs) -

(dsim_m_pret - dsim_m_pbs)) (Law *et al.* 2014; Ritchie *et al.* 2015). An FDR-adjusted *P*-value (*p*-adj)

cutoff of 0.05 was used for all tests conducted. Unless otherwise mentioned, all the reported

log2FoldChanges (log2FC) and interactions are significant at p-adj < 0.05.

2.12 Enrichment and pathways

The "clusterProfiler" R program package was used for gene ontology (GO) based gene set enrichment analysis (Wu *et al.* 2021). GO-Biological Process terms with adjusted *p*-value (FDR) less than 0.05 were considered significantly enriched. GO-BP annotations were performed separately for each contrast. The enrichment of DEGs in the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways was also tested using the "clusterProfiler" package in the R (FDR < 0.05). Expression differences of DEGs for selected pathways were visualized using "Pathvisio" and the wiki pathways plugin. The wiki pathway WP3830 was used to generate the Toll-IMD-Jak/STAT gene list and pathways.

2.13 Functional groups

Flybase was used to identify the functional groups/class gene list using key terms for Bomanin (BOMs), Drosophila immune-induced molecules (DIMs), Turandot (Tots), Thioester-containing protein (Teps), Insulin, Antimicrobial peptides (AMPs), Jun N-terminal kinase (JNK)/TNF-alpha-Eiger Pathway and Regulator, Positive and Negative Regulators of Toll, Positive and Negative Regulators of immune deficiency (IMD), and Spätzle (Spz) Activating Pathway with Regulators (Gelbart *et al.* 1996; Gramates *et al.* 2022). The Jak/STAT and melanization gene list was created manually from earlier studies and review articles (Tang 2009; Myllymäki and Rämet 2014; Yu *et al.* 2022). Heat maps of functional groups/classes were generated by "ggplot2" in the R program.

3. Results

Sex dimorphism is commonly observed in immune responses but is variable based on the pathogen, host, and other environmental factors (Fischer *et al.* 2015; Klein and Flanagan 2016). There is clear evidence of sex differences in immunity in *Drosophila melanogaster*, both at the physiological and molecular level, with a few genes identified as sex-differentially induced or repressed in response to infection (Taylor and Kimbrell 2007; Duneau *et al.* 2017b; Shahrestani *et al.* 2018). However, the range and causes of variation in sex dimorphism in immune responses, how and why these differences have evolved, and the underlying molecular mechanisms are unknown. This study aimed to understand if sex differences in the immune response are conserved between closely related species of *Drosophila* (Figure 1A). The study characterized the sex differences between unmated males and females associated with the immune response against a gram-negative extracellular bacterium, *P. rettgeri*, in *D. melanogaster* and its close relative *D. simulans*, by combining survival assays with RNA-seq-based gene expression estimation (Figure 1B). DEGs were analyzed within species, for all annotated genes, and between species for orthologs.

3.1 Sex dimorphism in survival

Fruit flies initiate rapid and efficient innate immune responses in response to pathogens that are crucial for survival, which influence the eventual outcome of bacterial infections. As one metric of relative effectiveness of the immune response, survival assays were conducted for females and males of *D. melanogaster* and *D. simulans* upon *P. rettgeri* infection over 36 hours at 6 to 12-hour intervals in five replicates where each replicate contained 30 flies (Figure 2). The survival assay demonstrated significant sex dimorphism in *D. melanogaster* (*p*-value < 0.001), but no significant differences in survival between the sexes were identified in *D. simulans.* This demonstrates that there are species differences in sex dimorphism in response to bacterial infection. Primarily, the initial stages (0-12hpi) showed no significant difference in survival between sexes for either species (Figure 2A). Nonetheless, survival started to differ by sex around 12hpi when the infection was expected to be established. At 18hpi, there were significant differences between males and females, where females exhibited higher survival than males in *D. melanogaster*. This difference in survival rate was persistent between sexes from 18hpi to 36hpi. Overall, the Kaplan Meier survival curve and Cox-regression model for *D. melanogaster* showed significant differences between males and females for *P. rettgeri* infection (Figure 2A).

For *D. simulans*, a sister species and member of the *D. melanogaster* subgroup, comparable results were observed in the initial stages of the infection (up to 12hpi) (Figure 2B). However, survival data began to diverge at 18hpi relative to *D. melanogaster.* In *D. simulans,* females showed no survival differences compared to males. While males and females continued to have no survival differences for 18hpi to 36hpi, the *D. simulans* sexes did not differ significantly in overall survival. Unlike *D. melanogaster* flies, all the *D. simulans* infected flies had died at 36hpi for both sexes (Figure 2B).

Comparing between species, female *D. melanogaster* had significantly higher survival after the establishment of the infection relative to *D. simulans* (*p*-value < 0.001). In all data points after infection initiation, *D. melanogaster* females consistently showed significantly higher survival rates than *D. simulans* females. Males of both species had similar survival during the initial stages, but around 24- and 36hpi, the survival of *D. simulans* males is significantly lower (*p*-value < 0.001). Overall, *D. melanogaster* females had markedly higher survival than *D. melanogaster* males and both males and females of *D. simulans* (Figure 2).

3.2 Bacterial load and BLUD

Bacterial load determination is a measure of the ability of the immune response to ease bacterial growth rates. The bacterial load data showed no statistical differences among experimental groups

when comparing the samples across sexes by each interval for either species (Figure 3, File S14).

Previous studies in the females of these two species also detected no significant bacterial load

differences upon gram-negative *Serratia marcescens* infection at 12hpi and 25hpi (Hill-Burns and Clark

2010). The data showed that the loads were similar for most individuals in the early stages, at 4- and

12hpi. Overall, the load increased in all the experimental groups for *D. melanogaster* and *D. simulans* at

16- and 20hpi; however, a subset of flies in each group showed a higher load than other individual flies.

At 20hpi, both the females and males showed a binary outcome of the infection clustering into two

groups. This can be explained if some individuals are able to suppress the bacterial infection progression

and keep the bacterial count lower than average, whereas others are less able to inhibit the infection

(Duneau *et al.* 2017a).

The results of the BLUD assay showed no sexually dimorphic pattern in bacterial burden at the time of

death upon *P. rettgeri* infection in *D. melanogaster* and *D. simulans*; moreover, there was no significant

difference in the species comparison (*p*-value > 0.05) (Figure 4). Overall, the bacterial load was unable to

explain the sex differences observed in survival.

3.3 Identification of DEGs in each sex and species

Within species in *D. melanogaster*, after normalization, 13,588 genes were tested. In response to

infection, 2269 genes were downregulated in males compared to 977 in females; with 629 genes

downregulated in both sexes (Figure 5A, Table S1). A higher number of genes (1640) were uniquely

downregulated in males, compared to females (348), which might indicate higher stress levels in males

upon infection. The number of upregulated genes was more comparable between males and females.

The number of genes that were upregulated in males was 530, and in females, 599, and 264 upregulated

genes were in common (Figure 5A).

After normalization, 13,010 genes were tested in *D. simulans,* which also showed a higher number of

genes downregulated in males (1871 genes) compared to females (1683 genes), but the difference is

less than in *D. melanogaster*, where males had roughly 2-fold higher numbers of downregulated genes

relative to females (Figure 5B, Table S1). In females, 633 genes were upregulated, whereas 503 genes

increased expression under infection in males. Both sexes downregulated 850 genes and upregulated

270 genes in common in *D. simulans*. The treatment by sex interaction, was tested separately for *D.*

melanogaster and *D. simulans* (comparison within species). There were 994 genes (*D. melanogaster*)

and 1041 genes (*D. simulans*) that showed sex differences in expression in response to infection with

FDR < 0.05 (Figure 5B, Table S1).

To compare across species for each sex, the interaction for the effect of treatment by species was tested

separately for females and males (File S6-S7). There were 11,849 genes for which the species could be

compared directly. In females, 569 genes were differentially regulated upon infection in the comparison

between *D. melanogaster* and *D. simulans* at FDR < 0.05 (File S8). In males, 477 genes have differential

regulation in the comparison between these two species (FDR < 0.05) (File S8). Additionally, 159 of the

differentially regulated genes are shared between females and males; thus, females showed unique

differential regulation for 410 genes and males for 318 genes (File S8).

3.4 Comparison of gene expression profiles in female and male adults

The analysis of the DEGs in the experimental groups indicated a complex pattern of sex dimorphism for

both species (Figure 6, File S4-S5). However, all experimental groups expressed immune response genes

such as AMPs at significantly higher levels than wound control, as expected upon infection. The volcano

plots of the four experimental groups showed that *D. melanogaster* females broadly expressed key

systemic immune and stress response genes at higher levels than the rest of the three experimental

groups. In *D. melanogaster*, both males and females upregulated AMPs, PAMP recognition proteins,

regulatory proteins, and stress response genes, but females expressed them at higher levels relative to

males. The *D. simulans* females also upregulated immune genes at higher levels than males, but

generally, the increase in expression was much lower than that for *D. melanogaster* females (Figure 6).

The comparison between species showed that females of the closely related species expressed quite

different sets of genes (Figure 6). Comparing across males and females of the species, males suppress

different genes than females do in both species. Downregulated genes in females of both species

included several female reproductive genes, for example, *Yp1* and *Yp2*. This pattern is explored further

in the enrichment analysis. Moreover, as mentioned earlier, the sex differences are visible in the case of

the up, and down-regulated gene numbers for females and males in both species; however, the sex

differences are more prominent when the data was separated based on the fold changes (FC) (Figure S1,

File S4-S5).

3.5 Gene set enrichment analysis

Gene set enrichment analysis (GSEA) was conducted using the Gene Ontology Biological Process (GO-BP)

annotation. The enriched GO-BP terms showed that males and females expressed defense and stress

response gene sets, but they used distinct strategies to different degrees in response to the bacterial

infection (File S9).

In *D. melanogaster*, 50 GO-BP terms were significantly enriched in females; among them, 49 were

upregulated, and only 1 was downregulated (Figure 7, Table S2, File S9). In males, a total of 101

significantly enriched GO-BP terms were identified, with 84 corresponding to upregulation and 17 to

downregulation (Figure 7, Table S2, File S9). Interestingly, 46 out of 49 GO-BP terms for upregulated

genes in females are also significant for upregulated genes in males, with 1 out of 1 GO-BP sets of genes

commonly suppressed in both sexes.

In *D. melanogaster*, the top 20 enriched GO-BP terms for upregulated genes in females showed immune

response terms such as defense response and response to gram-negative and -positive bacterium,

response to fungus, and antimicrobial humoral response (Figure 8, File S9). *D. melanogaster* males also

upregulated gene sets related to similar immune responses, such as response to gram-positive

bacterium, antibacterial humoral response, and response to fungus. Females (5 GO-BP terms) and males

(7 GO-BP terms) upregulated genes related to negative regulation of the immune and defense responses

(Figure 8, File S9). This could correspond to the modulation of immune responses through feedback

mechanisms (Tanji and Ip 2005; Vincent and Dionne 2021). The enriched terms corresponding to

downregulated genes showed that *D. melanogaster* females downregulate only one GO-BP term, lipid

metabolic process; however, in males, 17 GO-BP were identified where almost all of the terms were

related to metabolism (14 metabolic processes, 2 catabolic processes, 1 homeostasis) (Figure 8, File S9).

In *D. simulans*, 68 GO-BPs were significantly enriched in females; among them, 47 corresponded to

upregulated genes, and 21 downregulated genes (Figure 7, Table S2, File S9). In males, 72 total

significantly enriched GO-BP terms were identified, with 57 for upregulated genes and 15 for those that

were downregulated (Figure 7, Table S2, File S9). In *D. simulans*, the enriched groups among

upregulated genes of the males and females showed 42 commonly activated gene sets, whereas no

common GO terms were identified as enriched among downregulated genes (Figure 7, File S9). In *D.*

simulans, the top 20 GO-BP terms for the males and females corresponded to activation of the immune

response, such as response against gram-negative and positive bacteria, antimicrobial humoral

response, innate immune response, etc. (Figure 9, File S9). In addition, both males (6) and females (6)

had positively regulated immune-related terms among the top 20 most enriched categories. However,

females expressed less terms related to negative regulation of the immune response (3) than males (6)

in *D. simulans*. The males of *D. simulans* showed suppression of metabolism-related genes (11 metabolic

processes, 4 catabolic processes), which was also observed in *D. melanogaster* males (Figure 8-9). On

the other hand, females of *D. simulans* downregulated predominantly reproduction-related gene sets

such as eggshell formation, developmental processes involved in reproduction, etc. Thus, although

females of both species downregulated the iconic yolk protein genes, widespread downregulation of

reproductive genes occurred only in *D. simulans* females (Figure 8-9, File S9).

In comparisons between species, for females in *D. simulans*, there were 3 upregulated and 21 uniquely

downregulated gene sets, whereas for *D. melanogaster* females, there were 5 upregulated terms and 1

downregulated term (File S9). In addition, 44 upregulated terms showed overlap between females of *D.*

melanogaster and *D. simulans*, but there is no overlap among the downregulated terms. However,

males in both species had 56 upregulated and 12 downregulated GO-BP terms in common. There were

28 uniquely upregulated terms in *D. melanogaster* males and 5 downregulated terms, whereas in *D.*

simulans only 1 upregulated term and 3 downregulated terms were unique. Overall, the enrichment

analysis showed distinct sex dimorphism with respect to up- or downregulated functional categories of

genes in *D. melanogaster* and *D. simulans*. The analysis revealed differences between males and females

not only in upregulated biological processes but also in downregulated biological processes, which might

reflect different infection levels and control strategies for the sexes (File S9).

3.6 Infection-induced differences in the expression of IMD pathway genes

In the IMD pathway, the signal receptors and cellular response showed sex dimorphism in expression in

response to infection (Figure 10-11, File S10). This could be caused by differences in the regulation of

the IMD pathway between sexes and may be associated with sex differences in the effectiveness of the

immune response. In *D. melanogaster*, peptidoglycan recognition proteins (PGRP) of the IMD pathway

showed significantly higher expression in females for PGRP-LB and PGRP-LC than in males upon infection

(p-adj < 0.05) (Figure 10, File S10). The signal transducers for the IMD pathways were not significantly

different between sexes; however, the key IMD transcription factor Relish was significantly upregulated

more in males (log2FC = 2.16) than in females (log2FC = 1.54) in *D. melanogaster* (p-adj < 0.05). The

effector genes of IMD showed broad upregulation in both sexes, which includes AMPs such as *Attacins*

(*Att*), *Cecropins* (*Cec*), and *Diptericin-B* (*DptB*). In both sexes, *AttD* was the most highly expressed AMP; it

was upregulated around 20-fold in females, as compared to 15-fold in males (p-adj < 0.05) (File S10). In

addition, *CecB*, *CecA1*, *AttB*, *AttA*, and *DptB* were upregulated at significantly higher levels in females

relative to males (p-adj < 0.05) (File S10). Moreover, females exhibited upregulation in *Dro* (log2FC =

2.89), *DptA* (log2FC = 3.66), *CecC* (log2FC = 6.22), *CecA2* (log2FC = 6.08), and *AttC* (log2FC = 3.78), and

these genes are also strongly upregulated in males, *Dro* (log2FC = 2.63), *DptA* (log2FC = 3.38), *CecC*

(log2FC = 4.34), *CecA2* (log2FC = 4.46), and *AttC* (log2FC = 3.29) (File S10).

In *D. simulans*, the IMD pathway also showed female-biased upregulation in response to bacterial

infection, but not all sex differences in specific genes are conserved between species (Figure 11, Figure

S2, File S10). The focal signal-transducing innate immune signal receptor PGRP-LC is expressed

significantly higher in females relative to males in *D. simulans* when each group is compared with its

wound control (p-adj < 0.05) (Choe *et al.* 2005). Interestingly, PGRP-LF, which inhibits the activation of

IMD pathways by capturing circulating peptidoglycan, is highly upregulated in *D. simulans* females

(log2FC = 3.08) compared to the other three experimental groups, and also showed significant sex

differences with *D. simulans* males (log2FC = 1.34) (p-adj < 0.05) (Figure 11, File S10) (Maillet *et al.*

2008). The key IMD transcription factor Relish was upregulated in females (log2FC = 2.02) and males

(log2FC = 1.84) in *D. simulans*. The effector genes of IMD showed broad upregulation in both sexes,

which includes AMPs such as *Attacins* (*Att*), *Cecropins* (*Cec*), and *Diptericin-B* (*DptB*). In both sexes, *AttD*

was the most highly expressed AMP; it was upregulated around 14.6-fold in females, as compared to

13.3-fold in males (p-adj < 0.05) (File S10). In addition, *AttA*, *AttB*, *AttC*, *CecA1*, *CecA2*, *CecB*, *CecC*, *Dro*,

DptA, and *DptB* were upregulated at significantly higher levels in females relative to males (p-adj < 0.05)

(File S10). Overall, the IMD cellular effector molecules of *D. simulans* showed higher expression in

females than males. However, the degree of upregulation of these AMPs in *D. simulans* females is lower

on average than in *D. melanogaster* females (File S10).

3.7 Regulators of IMD pathway genes

The positive and negative regulators of the IMD pathway were analyzed for each sex in both species. In

the case of the IMD positive regulators (FBgg0001197), PGRP-SD, which can significantly contribute as a

'positive regulator' of the IMD pathway, showed significant differential expression between females

(7.64-Fold) and males (5.56-Fold) of *D. melanogaster* (p-adj < 0.05) (Figure S3, File S11) (Gelbart *et al.*

1996). Expression patterns of this gene also differed between females and males in *D. simulans* (p-adj <

0.05); however, both sexes of *D. simulans* showed lower levels of upregulation than their *D.*

melanogaster counterparts. In addition, expression of the gene encoding Charon*,* which can directly

activate Rel/NF-kB, was upregulated in males (log2FC = 1.19) to a higher degree than in females (log2FC

= 0.36) in only *D. melanogaster* (p-adj < 0.05). Differences in the 'negative regulators' of IMD

(FBgg0001196) also included significantly higher expression of PGRP-SC2, a down-regulator of the IMD

pathway, and PGRP-LB in *D. melanogaster* females relative to males (p-adj < 0.05; Figure S3, File S11)

(Bischoff et al. 2006; Guo et al. 2014). While PGRP-LB is one of the prominent receptors involved in the

recognition and activation of the IMD pathway, researchers have found that it enables females to

reduce IMD activity upon a decrease in bacterial numbers (Zaidman-Rémy et al. 2006). Thus, the

significant upregulation of two negative regulators of IMD, PGRP-LB and PGRP-SC2, in *D. melanogaster*

females tempts us to speculate that in females, there is more controlled IMD expression relative to

males (Bischoff et al. 2006; Zaidman-Rémy et al. 2006; Guo et al. 2014). Additionally, the negative

regulator of IMD, PGRP-LB and PGRP-SC2, were not highly upregulated in *D. simulans* compared to *D.*

melanogaster sexes (p-adj < 0.05). However, another negative regulator, PGRP-LF, is highly upregulated

in *D. simulans* females (log2FC = 3.08) compared to males (log2FC = 1.34) (p-adj < 0.05) (Figure S3)

(Maillet et al. 2008). Overall, the IMD pathway is involved in the response to *P. rettgeri* via effector

molecules for females and males in a sex differential manner in *D. melanogaster* and *D. simulans*;

however, higher upregulation of the negative regulators such as PGRP-LB, PGRP-SC2, and PGRP-LF in

females might indicate that these flies activated IMD pathways at a level where suppression was

initiated, whereas other experimental groups did not reach that elevated level.

3.8 Infection-induced differences in the expression of Toll pathway genes

The Toll pathway was originally identified as an effector pathway activated upon gram-positive bacterial

and fungal infection, but recent studies show it also plays a crucial role in the responses to gram-

negative bacterial infections and viral infections (Valanne et al. 2022). In *D. melanogaster*, signal

receptor PGRP-SD was more elevated in females than males (p-adj < 0.05) (Figure 10, Figure S2, File

S10). In addition, the extracellular molecule *Serine Protease Immune Response Integrator* (*spirit*) showed

significant differences between females (log2FC = 3.81) and males (log2FC = 1.73) (p-adj < 0.05). The

males significantly upregulated *spz;* Spz is a core ligand of Toll receptor (Tl)*,* relative to females in *D.*

melanogaster (p-adj <0.05) (Valanne et al. 2011). The 'intracellular signal transducers' of Toll had no

significant changes in regulation between sexes, including the key NF-κB-related transcription factors

Dorsal-related immunity factor (Dif) and Dorsal (dl) (Valanne et al. 2011). In contrast, the effector

molecules showed higher upregulation of *Drosomycin* (*Drs*), *Defensin* (*Def*), and *Metchnikowin-like*

(*Mtkl*) in females relative to males in *D. melanogaster* (p-adj < 0.05). *Mtkl* was the most highly expressed

Toll-AMP for both sexes in *D. melanogaster*, which was upregulated by approximately 10-fold in females

and 7-fold in males (p-adj < 0.05). Interestingly, the gene encoding focal antifungal peptide-producing

AMP, *Drosomycin* (*Drs*), was also upregulated in both sexes, but other antifungal AMP genes such as

Daishos (*Dso1* and *Dso2*) and *Drosomycin-like* (*Drs1-6*) showed no significant increase in expression

upon bacterial infection, instead some of them including *Dso1* and *Dso2* showed significant

downregulation, where these genes were downregulated in males to a higher degree than in females (p-

adj < 0.05) (Figure 10, File S10).

D. simulans showed analogous expression patterns to *D. melanogaster* in the extracellular portion of the

Toll Pathway. PGRP-SA and PGRP-SD were significantly upregulated at higher levels in females relative to

males (p-adj < 0.05), but the degree of upregulation is lower than in *D. melanogaster* (Figure 10-11, File

S10). This also holds true for the expression of *spirit*, which exhibits significant sex differences (p-adj <

0.05), although the expression level is lower compared to *D. melanogaster*. The gene *spz* was

significantly overexpressed in males relative to females (p-adj < 0.05), as also identified in *D.*

melanogaster (Figure 10-11). In *D. simulans*, the key NF-κB-related transcription factors *Dorsal-related*

immunity factor (*Dif*) and *Dorsal* (*dl*) showed no significant differences between sexes (Valanne et al.

2011). The effector molecules showed higher upregulation of *Drosomycin* (*Drs*), *Defensin* (*Def*), and

Metchnikowin (*Mtk*) in females relative to males In *D. simulans* (p-adj < 0.05). Interestingly, *Drs* was also

upregulated in both sexes, but other antifungal AMPs such as *Drosomycin-like* (*Drsl*) *2*, *Drsl 4*, and *Drsl 5*

showed significant downregulation upon bacterial infection, where females showed higher levels of

downregulation than males did (p-adj < 0.05) (Figure 10, File S10).

In between species comparisons, *D. simulans* showed significant downregulation of *SPE*, unlike *D.*

melanogaster, with females (log2FC = -1.3) exhibiting more downregulation than males (log2FC = -0.44)

(p-adj < 0.05). Another key difference was observed in the expression of Clip-SP *Persephone (psh)*, *D.*

simulans females (7-Fold) and males (4-Fold) upregulated *psh*, where the reverse direction is observed

in females (log2FC = -0.81) and males (log2FC = -0.45) of *D. melanogaster*. Overall, these findings

suggest that *D. simulans* might be using a different route to stimulate Toll pathways other than PRR base

activation.

3.9 Regulators of Toll pathway genes

The regulation of the Toll pathway is much more complex than the regulation of IMD, as there are

diverse functions performed by the Toll pathway in addition to immunity. Drosophila possesses six

distinct Toll receptors, each contributing individually or in concert to accomplish complex

multidimensional functions; the data showed none of the Toll receptors were sex differentially

expressed in either species (Valanne et al. 2011) (File S11). However, the *spz* activation pathway, which

is critical to the regulation of the Toll pathway as Spz can be activated in multiple ways, such as PRR-

based, Psh-SPE-Spz, and Psh-Protease-Spz, shows apparent sex differences. Multiple genes in this

pathway are expressed in a sex differential manner, such as *spirit, psh*, *Hayan*, and *PGRP-SC2* (Figure S4,

File S11). The sex differential regulation of *spirit* and *psh* has previously been discussed with respect to

the overall Toll pathways (Duneau et al. 2017b). There were also female-biased sex differences for

wound-related serine protease Hayan in *D. simulans* (p-adj < 0.05), which is involved in the activation of

the melanization cascade as well as Spz based Toll pathway activation (Yamamoto-Hino and Goto 2016;

Dudzic et al. 2019). *Hayan* was also upregulated in females (log2FC = 3.22) and in males (log2FC = 2.45)

in *D. melanogaster*. The gene encoding PGRP-SC2, which acts as a negative regulator of the IMD

pathway, is expressed at higher levels upon infection in females (log2FC = 3.59) than males (log2FC =

3.15) in *D. melanogaster* (p-adj < 0.05) (File S12) (Bischoff et al. 2006). *PGRP-SC2* was also upregulated in

both females (log2FC = 1.60) and males (log2FC = 1.56) in *D. simulans*. (File S12). Aside from the *spz*

activating pathway, positive (FBgg0001057) and negative (FBgg0001053) regulators of the Toll pathway

showed no significant sex differences in either species (Figure S4, File S11).

Overall, considering the Toll pathway and its regulators, suggests there is female-biased sex differential

upregulation of Toll-AMPs for both species; however, the degree of upregulation is higher in *D.*

melanogaster than in *D. simulans* like IMD-AMPs. A comparison between species detected a remarkable

divergence in the Toll activation process, where *D. simulans* employed both PRR-dependent and

independent mechanisms, while the closely related *D. melanogaster* solely upregulated genes related to

PRR-dependent activation.

3.10 Other immune response molecules and pathways

The innate immune response of Drosophila is a complex interplay of conserved innate immune signaling

pathways and other immune molecules, along with the involvement of immune cells. Several Drosophila

immune-induced molecules (DIMs) have been identified in the fly hemolymph (Uttenweiler-Joseph et al.

1998). Recent research revealed that twelve related immune-induced molecules, known as the

Bomanins (Boms), constitute a family that plays a role in defending against certain fungi and bacteria

(Clemmons et al. 2015). Three of the twelve Bomanins (Boms) in *D. melanogaster*—*BomT1*, *BomS5*, and

BomBc3—are expressed in a sex-differential manner (p-adj < 0.05). (Figure S5, File S11). *BomT1* is

upregulated to a greater degree in females (log2FC = 2.59) relative to males (log2FC = 1.53) (p-adj <

0.05). Opposing sex differences were observed for *BomS5*, where females upregulated, and males

downregulated expression (p-adj < 0.05). *BomBc3* was downregulated in both sexes in *D. melanogaster*,

but in males, it was downregulated to a greater extent than in females (p-adj < 0.05). However, only

BomS2 and *BomS4* showed significant sex differences in *D. simulans* (p-adj < 0.05). In the case of *BomT1*,

upregulation was observed in both females (log2FC = 1.18) and males (log2FC = 1.03) in *D. simulans*.

BomBc3 expression is different from that of *D. melanogaster,* where *BomBc3* is downregulated in

females (log2FC = -1.36) and males (log2FC = -1.10) (Figure S5, File S11). The expression of the Boms and

their regulators showed marked species differences among the sexes.

Melanization is one of the extraordinary features of insect innate immunity and is regulated by Toll

pathways in terms of the gene expression of serine proteases (SPs) and serpins (Spn) (Dudzic et al.

2019). PGRP-SA and GNBP1/ β-Glucan Recognition Proteins (βGRP) can directly modulate the expression

of the melanization cascade genes (Matskevich et al. 2010). In the case of core serine proteases, *Hayan*

and *SP7/MP3* both show significant sex differential expression in *D. simulans* (p-adj < 0.05) (Figure S5,

File S11). Although *D. melanogaster* females and males upregulated Hayan and SP7 more than wound

control, no significant sex differences were observed. In both species, the level of *SP7* upregulation is

lower than that of *Hayan*. Although key SPs are upregulated, there is downregulation of the key

prophenoloxidases (*PPO*), *PPO1*, and *PPO2* (Figure S5, File S11). In addition, there is significant

upregulation of the negative regulator of melanization, *Spn28Dc*, in both species. With higher

upregulation in females (log2FC = 2.99) relative to males (log2FC = 1.15) in *D. melanogaster* (p-adj <

0.05) and a similar pattern found in the females (log2FC = 3.78) and males (log2FC = 2.07) of *D. simulans*

(*p*-adj < 0.05) (Figure S5, File S11). These data strongly suggest that the melanization cascade was not

activated; thus, Hayan and SP7 might be utilized by other immunological processes, such as Toll

activation, as has been observed in other studies (Yamamoto-Hino and Goto 2016; Dudzic et al. 2019).

In *Drosophila*, the Jak-STAT pathway is crucial for regulating antimicrobial peptide (AMP) production in

response to bacterial infection and antiviral response (Myllymäki & Rämet, 2014). The cytokine-like

ligands *upds*, signal receptor *domeless* (*Dome*), *Janus kinase* (*JAK*), *hopscotch (hop)*, and the Drosophila

STAT transcription factor, *stat92E*, were not sex-differentially expressed in either species (Figure S7, File

S10-11). Despite the lack of significant sex differences in transcript level in upstream parts of this

pathway, some of the effector molecules, such as *Turandot C* (*TotC*), *TotA*, *TotX*, and *TotM*, were

significantly upregulated compared to the wound control in males of *D. melanogaster (p*-adj < 0.001*);

this pattern is absent in females of *D. melanogaster* and both sexes of *D. simulans* (Figure 10-11). The

study found no sex differences in components of negative regulators in either species (Figure S7, File

S11). JNK is also known as the TNFalpha-Eiger pathway in Drosophila, which, like Jak-STAT, has a diverse

set of functions with a key role in immune responses, development, and morphogenesis (Gan et al.

2021). The gene expression analysis found that JNK receptors, protein kinase, and regulators showed no

significant sex differences (Tafesh-Edwards and Eleftherianos 2020)(Figure S7, File S11).

Finally, apart from PGRPs of the IMD and Toll pathways, our data found *PGRP-SB1*, an antibacterial

PGRP, was more upregulated in females (log2FC = 2.90) relative to males (log2FC = 2.11) in *D.*

melanogaster (*p*-adj < 0.05) (File S11). The upregulation is also conserved in *D. simulans*, where females

(log2FC = 1.71) upregulate *PGRP-SB1* more than males (log2FC = 0.84) (*p*-adj < 0.05) (File S11). Prior

works on *PGRP-SB1* report that it can be significantly upregulated upon IMD pathway activation by the

gram-negative bacterial infection, which is consistent with our findings (Zaidman-Rémy et al. 2011). The

analysis of *thioester-containing proteins* (*Teps*) identified that only *Tep2* showed differential expression

compared to wound control in females (log2FC = 2.38) and males (log2FC = 1.98) in *D. melanogaster*

(Figure S6, File S11). The females (log2FC = 0.44) and males (log2FC = 0.32) also upregulated *Tep2* in *D.*

simulans, but the degree of upregulation is significantly lower than in *D. melanogaster* counterparts

(Figure S6, File S11).

4. Discussion

The emergence of sex differences in immune responses is thought to originate from complex

evolutionary dynamics including trade-offs and sexual conflict (Zuk and Stoehr 2002; Scharf et al. 2013;

Schwenke et al. 2016). The interplay between the immune system and various physiological processes

can result in trade-offs between immunity and other energetically costly life-history traits, such as

reproduction (Zuk and Stoehr 2002; Schwenke et al. 2016). Immune responses can also be sexually

dimorphic, with males and females showing different responses to the same pathogens in identical

environments (Lazzaro and Little 2009; Morrow 2015). Recently, there has been a growing interest in

adult immunity and the role of sex-biased gene expression in immune response pathways in generating

sex dimorphism in the overall immune response due to its impact on the medical sector (Fischer et al.

2015; Wilkinson et al. 2022). However, many studies in immunology have focused predominantly on one

sex, often males (Belmonte et al. 2020). The importance of including both sexes in immunological

studies is increasingly acknowledged and enables researchers to achieve a comprehensive

understanding of sex dimorphism, evolution, and molecular mechanisms of the immune response.

Our investigation of sex differences in immune response against *P. rettgeri* in *D. melanogaster* and *D.*

simulans found that survival differences may not be conserved, and sexes differ in survival in *D.*

melanogaster. Interestingly, we also find that although both sexes express similar immune response

genes for conserved immune pathways and some other immune response components, there is a strong

pattern of a robust response in *D. melanogaster* females relative to males. In *D. simulans*, females also

mounted higher immune gene expression, but the expression difference between the two sexes is not

as high as in *D. melanogaster* and did not confer a significant survival advantage. Moreover, both species

upregulate similar genes to inhibit infection, but broadly, the degree of up or downregulation is lower in

D. simulans than in *D. melanogaster*. Overall, these findings highlight species-specific variations in the

effectiveness of the immune response and the physiological impact of the infection. In addition, the

downregulation of metabolism-related gene sets in males of both species, and the downregulation of

reproductive genes in *D. simulans* suggests that each sex uses different strategies to modulate other

processes in response to the infection. Lastly, the evolutionarily conserved bacterial pathways IMD and

Toll exhibited sex-specific variation, encompassing not only effector molecules but also regulators, such

as those involved in negative feedback, in the context of infection progression. This suggests that the

IMD pathway responds more in a controlled manner in *D. melanogaster* females than in the rest of the

three groups. Moreover, in *D. simulans,* infected animals used two distinct strategies at sixteen hours

post-infection, canonical PRR-SPE-Spz, and microbial proteases-based Persephone cascade, to activate

the Toll pathways.

4.1 Survival of unmated males and females differs between the sexes and species

The survival rates of unmated male and female flies were monitored over a 36-hour period following

infection. In *D. melanogaster*, females showed significantly higher survival rates than males from 18hpi

onward. In *D. simulans*, while females had slightly higher mean survival rates than males, the difference

was not statistically significant. In previous studies, females have been observed to represent higher

survival rates than males, for example, in gram-positive *S. aureus* infection in *D. melanogaster* (Duneau

et al. 2017b). However, this same study also found that *D. melanogaster* females were more susceptible

to *P. rettgeri* (gram-negative), *P. alcalifaciens* (gram-negative), and *E. faecalis* (gram-positive) infections

(Duneau et al. 2017b). In addition, two individual studies of gram-negative *Pseudomonas aeruginosa*

infections showed opposite-sex survival bias direction (Chowdhury et al., 2019; Vincent & Sharp, 2014).

Sex dimorphism in survival rates appears to depend on sex, host strains, type of infection, and external

conditions in *Drosophila* (Howick and Lazzaro 2014; Rai et al. 2023). It has also been shown that even

when the pathogen and host are identical, mating status, diet, age, and injection site, among other

differences, can alter the outcome of the infection (Lazzaro and Little 2009; Short and Lazzaro 2010;

Howick and Lazzaro 2014; Rai et al. 2023). Therefore, results from different studies may vary due to

differences between laboratories or due to specific differences in the study design.

In our study, both males and females were unmated, whereas in the Duneau et al. 2017 study, males

were mated, and then compared to mated or unmated females (Duneau et al. 2017b). Studies have

indicated that mated females exhibit increased susceptibility to infection, elevated pathogen loads, and

reduced capacity to initiate immune responses, which could explain why unmated females in our study

had better outcomes than in studies that focused on mated animals (Short and Lazzaro 2010; Schwenke

and Lazzaro 2017; Rai et al. 2023).

To the best of our knowledge, the post-infection survival of *D. simulans* has not been previously

reported; thus, a direct comparison to existing studies is not possible. However, when comparing

between species, *D. melanogaster* females had significantly higher survival rates relative to *D. simulans*

females after the infection was established. Similarly, *D. melanogaster* males showed significant

differences in survival rates compared to *D. simulans* males from 18hpi. Overall, there are both sex- and

species-based differences in survival rates, with *D. melanogaster* females showing the highest survival

rates. Previous studies have shown variation within the same species based on different conditions, as

our results also show compared to prior studies. Here, we have different species in the same

environment and with concurrent experiments where both sexes and species can be directly compared.

This means that the difference we observe between *D. melanogaster* and *D. simulans* is likely to be

genetic, although we note that evaluation of additional strains is needed to establish fixed species-level

differences. This study showed that the impact of sex on survival may have diverged between these

species and both sex and species effects may be contributing to differences observed.

4.2 Distinct patterns of up and down gene regulation are concordant with differences in survival

The up and downregulation of gene expression in the context of stress is a common phenomenon

(Lecheta et al. 2020; Girgenti et al. 2021). There are fewer upregulated genes than downregulated for

both females and males in both species. *D. melanogaster* females downregulated 977 genes compared

to 2269 in males, which is consistent with lower survival rates of males and may indicate higher stress

conditions in males relative to females. Compared to the 1871 genes downregulated in *D. simulans*

males and 1683 genes in females, *D. melanogaster* females downregulated fewer genes. A similar

pattern of up and down-regulation was observed upon Sigma Virus infection in *D. melanogaster*, where

males downregulated 336 genes compared to 88 genes in females (Carpenter et al. 2009).

The enrichment analysis indicated differences between sexes in which types of genes were

downregulated in *D. melanogaster*. Only one GO-BP category was significantly enriched among

downregulated genes in females, whereas 17 metabolism-related downregulated categories were

identified in males. This is consistent with the idea of higher stress levels in infected males relative to

females in *D. melanogaster*. Likewise, *D. simulans* males downregulated 15 GO-BP categories, among

which metabolism-related genes were predominant.

Previous studies also found a trade-off between immune and metabolic processes in males upon

bacterial infection in *D. melanogaster* (Schlamp et al. 2021). Schlamp et al. showed that the activation

of the IMD pathways affects the metabolic homeostasis of *Drosophila* males (Schlamp et al. 2021). *D.*

simulans females may also have been under increased levels of stress because of infection, given that

there was extensive downregulation of reproductive GO: BP terms, such as eggshell formation and

reproductive developmental processes. Multiple studies have reported trade-offs between immune

response and reproduction in both males and females in *Drosophila* as well as other insects (McKean

and Nunney 2001; Short et al. 2012; Schwenke et al. 2016). The previous research on *Drosophila*

infected with *P. rettgeri* revealed that females halt egg-laying during the acute phase of infection, with a

return to normalcy observed during the chronic phase (Howick and Lazzaro 2014) and upon DNA virus

Kalithea (KV) infection, females experience degeneration of ovarian tissue and reduced fecundity

(Palmer et al. 2018). Additionally, viral-induced female infertility was found upon flock house virus (FHV)

infection due to the destruction of oocytes (Thomson et al. 2012). Overall, *D. melanogaster* females

downregulated fewer genes, and had fewer functional categories that were consistently downregulated,

while at the same time upregulating immune-related genes to a greater degree, which may be related to

their higher survival rates.

4.3 Sex and species differences in the IMD pathway

IMD receptors and effector molecules were upregulated in females and males in both species. In

general, females showed higher levels of expression than males, especially the effector molecules such

as AMPs. There is a noticeable degree of higher upregulation in *D. melanogaster* flies compared to *D.*

simulans. As the IMD pathway is specialized in response to gram-negative bacterial infection, *P. rettgeri*

infection is expected to elicit a higher level of IMD-regulated AMP expression than other core immune

response pathways. There was a consistent pattern of female-biased sex differential expression in both

species for IMD-AMPs; however, it was also evident that the sexes of *D. simulans* consistently showed

lower levels of upregulation than *D. melanogaster* sexes.

The species-level expression difference is also visible in the positive and negative regulators of the IMD

pathway; for instance, PGRP-SD, primarily associated with the Toll pathway, significantly contributes to

the positive regulation of the IMD pathway by amplifying the recognition of peptidoglycans (Tanji and Ip

2005; Myllymäki et al. 2014). Our results showed that *D. melanogaster* males significantly upregulate

the gene *Charon,* which can directly activate the NF-kB/Relish at AMP promoters in males via a PARP-

1/Relish/Charon-dependent innate immune response (Ji et al. 2016). Alternatively, the knockdown of

Charon expression plays a vital role in the translocation of Relish to the nucleoplasm, and blocks that

function making flies vulnerable to bacterial attack (Ji et al. 2016). The influence of the PARP-

1/Relish/Charon-dependent response might act in conjunction with greater upregulation of NF-

kB/Relish in *D. melanogaster* males, relative to females; males may amplify the response to compensate

for lower upregulation of signal receptors (PRRs) (File S11). This indicates that sex differences are likely

to be complex, and interactions between different genes and pathways will need to be untangled to

delineate key mechanisms (Ji et al., 2016).

The precise modulation of energetically demanding immune pathways is crucial to halt the progression

of infection. Over-activation of the IMD pathways can severely affect survival (Bischoff et al. 2006). A

comprehensive analysis of the negative regulators of IMD pathways was conducted, revealing that *D.*

melanogaster and *D. simulans* exhibited sex differences in expression of the genes encoding certain

proteins, namely PGRP-LB, PGRP-SC2, PGRP-LF, and Pirk, which play roles in the negative regulation of

IMD pathways. *D. melanogaster* females and males both upregulated PGRP-LB, PGRP-SC2, PGRP-LF, and

Pirk, but differed in the degree of upregulation. PGRP-LB works as an activator of the IMD pathway, but

excessive upregulation of PGRP-LB can trigger a negative feedback loop for the IMD pathway (Bischoff et

al. 2006; Vincent and Dionne 2021). *D. melanogaster* females significantly upregulated PGRP-LB more

than males, and both sexes of *D. simulans. D. melanogaster* females also upregulated IMD-regulated

AMPs at a significantly higher level than the rest of the groups. The significant overexpression of PGRP-

LB could either activate the IMD pathway to more efficiently eliminate pathogens, or lead to the

pathway's downregulation via negative feedback in *D. melanogaster* females (Figure S2, File S11).

In addition to PGRP-LB, PGRP-SC2, another negative regulator of IMD pathways, also showed

significantly higher upregulation in *D. melanogaster* females relative to others. Another member of the

PGRP, PGRP-LF, is a membrane-associated protein, but it cannot bind to peptidoglycans directly (Persson

et al. 2007). PGRP-LF interacts with PGRP-LC and functions as a negative regulator of IMD signaling

(Maillet et al. 2008; Basbous et al. 2011). Both females (log2FC = 2.61) and males (log2FC =2.27)

upregulated PGRP-LF higher than control in *D. melanogaster*, with no significant difference between the

sexes. Interestingly, PGRP-LF is highly upregulated in *D. simulans* females compared to the other three

groups (Maillet et al. 2008). Another PGRP-LC interacting protein, Pirk, which is involved in the negative

regulation of IMD, was upregulated at significantly higher levels in females and males of both species

(Vincent and Dionne 2021).

Overall, the IMD pathway was employed in a sex-differential manner in *D. melanogaster* and *D.*

simulans. However, upon comparing these four "brakes" (PGRP-LB, PGRP-SC2, PGRP-LF, and Pirk) of IMD

across experimental groups, a noticeable trend emerges where *D. melanogaster* females exhibit higher

upregulation compared to other groups; this might indicate that *D. melanogaster* females activated the

IMD pathways at a level where negative regulation was initiated, whereas other experimental groups

were unable to reach that elevated level. The reason behind the significant upregulation of PGRP-LF and

Pirk in *D. simulans* remains unclear. This is particularly ambiguous considering the survival data, where it

is evident that *D. simulans* was not effectively handling the infection at 16hpi (Figure 2-3, Figure 10-11,

File S11).

Despite the upregulation of negative regulators, *D. melanogaster* females upregulate effector genes of

the IMD pathway to a higher degree than males or *D. simulans;* this may reflect greater control of

expression of effectors, as well as specific dynamics in female *D. melanogaster* in which upregulation of

negative regulators is a response to the high levels of effector expression at 16hpi. This would then be

expected to reduce effector expression at later time points, which was not possible to test from this

experimental data. The key transcription factor Relish/NF-κB is upregulated less than males at 16hpi for

D. melanogaster females; this might suggest there would be downregulation of the AMPs at later time

points than those measured (Zaidman-Rémy et al. 2006). Eventually, the increased expression of genes

involved in negative feedback loops regulating IMD in *D. melanogaster* females would be expected to

result in reduced signaling and potentially reduced expression of other genes in the IMD pathway (Tanji

and Ip 2005; Zaidman-Rémy et al. 2006; Vincent and Dionne 2021). These findings provide valuable

insights into the intricate mechanisms underlying sex-specific immune responses and suggest a balance

between gene activation and suppression as part of an adaptive strategy that contributes to increased

survival in females.

4.4 Sex and species differences in the Toll pathway

Toll is another conserved signaling pathway and is involved in many functionally different processes,

including innate immunity. Broadly, Toll-regulated AMPs were also upregulated in a female-biased

manner in both species. The comparison between species revealed higher upregulation in *D.*

melanogaster relative to *D. simulans,* which is also observed for IMD-regulated AMPs. However, the

level of Toll-AMP upregulation is lower than it is for IMD-regulated AMPs in both sexes for *D.*

melanogaster and *D. simulans.* This is expected because Toll is a more specialized response against

gram-positive bacterial and fungal infections (Valanne et al. 2011). In the IMD pathway, the core

transcription factors Relish showed sex differential expression for both species, but neither of the key

transcription factors (Dif and Dorsal) in the Toll pathway were expressed differentially in either species.

This suggests that the expression of Dif and Dorsal are not directly responsible for sex differences in the

expression of AMPs, although sex differences may result from post-transcriptional regulation or

differences in transcript levels prior to 16hpi (Schlamp et al., 2021). In addition, the Toll-associated

AMPs might be regulated by other noncanonical transcriptional mechanisms to elicit sex dimorphism

(Tanji and Ip 2005).

Sex differential regulation was detected for signal receptors and cellular response genes for both

species. Among these genes involved in *spz* activation, PGRP-SA, PGRP-SD, and Spirit were upregulated

in females and males of both species. This is expected to be associated with PRR-dependent Toll

activation. In addition, Clip-SP Hayan, one of the serine proteases of the melanization cascade, was

significantly overexpressed in females of both species relative to males (p-adj < 0.001) (Figure S5, File

S11). The analysis also detected a notable trend where females (7-Fold) and males (4-Fold) of *D.*

simulans drive significant upregulation of Clip-SP Psh (File S10). Both Clip-SP Hayan and Clip-SP Psh play

a vital role in regulating the Toll pathway downstream of Clip-SP Grass, ModSP, and PRRs (Yamamoto-

Hino and Goto 2016; Issa et al. 2018; Dudzic et al. 2019).

Persephone, a circulating serine protease, functions as an immune sensor for microbial proteolytic

activities upstream of the Toll pathway by detecting exogenous proteases (Chamy et al. 2008; Issa et al.

2018). Upon detection, Persephone can elicit Toll activation in two different ways, through Psh-SPE-Spz

or Psh-Protease-Spz (SPE independent) activation (Ligoxygakis et al. 2002; Gottar et al. 2006; Chamy et

al. 2008; Yamamoto-Hino and Goto 2016). Although it is not established that Hayan can detect microbial

proteases, it plays a vital role in both PRR-independent activation mechanisms (Dudzic et al. 2019).

Previous studies showed that animals with SPE null mutations can perform the pro-Spz to Spz cleavage,

suggesting that another SP can cleave Spz beyond SPE (Yamamoto-Hino and Goto 2016; Dudzic et al.

2019).

It is also possible that Hayan is upregulated to promote the melanization cascade, but the other key

members of the Hayan-dependent blackening reaction, PPO1, and PPO2, were downregulated in *D.*

simulans (Binggeli et al. 2014; Dudzic et al. 2015, 2019). In addition, the regulation of Hayan is strictly

controlled by the serpins (Spn) for melanization (Binggeli et al. 2014). Analysis of the negative regulators

of the melanization cascade reveals that Spn28Dc, a member of the Serpin family, which can negatively

regulate the melanization process, is significantly upregulated in females and males of *D. simulans* and

D. melanogaster (Ligoxygakis 2002; Tang 2009; Binggeli et al. 2014). This indicates that Hayan is more

likely to be involved in the PRR-independent Toll activation rather than melanization.

Based on the observation, we hypothesize that significant downregulation of SPE in females (log2FC = -

1.31) and males (log2FC = -0.44) in *D. simulans* coincides with the use of the alternate Toll activation

pathways, whereas females (log2FC = 0.91) and males (log2FC = 0.65) of *D. melanogaster* might

sufficiently upregulate SPE; thus *D. melanogaster* is not depending on the microbial proteases-based

Psh-SPE-Spz or Psh-Protease-Spz (SPE independent) activation at 16hpi (Issa et al. 2018; Dudzic et al.

2019). In contrast, Toll responses in *D. simulans* might be enhanced by the additional microbial protease

based Psh-SPE-Spz or Psh-Protease-Spz (SPE independent) activation allowing maintenance of

appropriate Spz levels (Yamamoto-Hino and Goto 2016; Issa et al. 2018; Dudzic et al. 2019). We note

that it is not possible to make conclusions from our single timepoint data, although our results support

the investigation of the hypothesis that SPE at 16hpi is downregulated, with *D. simulans* flies using Psh-

Protease-Spz (SPE independent) activation.

Immune responses incur significant energetic costs for organisms. Therefore, the success of an organism

depends not only on the activation of immune pathways but also on their effective suppression. Among

the suppressors of Spz activation and the Toll pathway, two of the Serpins family members named

Spn88Ea and Spn42Dd were overexpressed in both sexes of *D. melanogaster* (*p*-adj < 0.001). The role of

the Spn88Ea and Spn42Dd is to negatively regulate the production of antifungal peptides via the Toll

pathway (Reichhart et al. 2011). As mentioned earlier, there is less expression of the fungal-related

AMPs except for *Drosomycin (Drs),* which has multiple roles as antipathogenic peptides, including

function against bacterial infection (Hanson and Lemaitre 2020; Lin et al. 2020). These findings also

explain the downregulation of the fungal cell wall pattern recognition receptor GNBP3 in the infected

females (log2FC = -0.60) and males (-0.61) relative to wound control in *D. melanogaster* (Figure S2, File

S10).

5. Conclusion

Our data showed that similar genes and pathways are upregulated in males and females, but the degree

of upregulation differs between sexes and species. The comparative analysis revealed that these sex

differences in the upregulation of immune-related biological processes are conserved between the two

closely related species examined; interestingly, the sexes appear to suppress different sets of genes,

indicating potential differences in trade-offs and resulting strategies. However, there are differences

between species, both *D. melanogaster* sexes show higher upregulation of immune genes relative to *D.*

simulans. Moreover, among the four experimental groups, the *D. melanogaster* female upregulated

immune genes at the highest level relative to males and *D. simulans* of both sexes – which is concordant

with the differences observed in survival. The downregulation of biological processes in females and

males showed significant differences in both species. Although the pattern of downregulated GO terms

is similar in males between species, the categories corresponding to downregulated genes in females

are significantly different.

It has been reported in prior studies that Toll and IMD are responsible for the sex differences in *D.*

melanogaster, and our results showed there were sex differences in the regulation of both conserved

bacterial infection response pathways, IMD and Toll, which supports these previous studies (Duneau et

al. 2017b; Vincent and Dionne 2021). Intriguingly, our data strongly suggest that *D. melanogaster* and *D.*

simulans employ two mechanisms to activate Toll pathways differently during infection by *P. rettgeri*,

with *D. melanogaster* using the canonical PRR-based Toll activation, and *D. simulans,* the microbial

protease recognition-based Persephone pathway to stimulate Toll effectors along with the canonical

path. The AMP expression level of *D. melanogaster* is consistent with the hypothesis that PRR-SPE-Spz

activation alone can mount a heightened immune response in *D. melanogaster* relative to both

processes in *D. simulans*. The IMD pathway genes showed sex-differential expression and varied in the

degree of expression across species. The analysis of the negative regulators showed *D. melanogaster*

females were upregulating four key "brakes" (PGRP-LB, PGRP-SC2, PGRP-LF, and Pirk) to prevent the

overstimulation of IMD pathways at 16hpi. Overall, our finding suggests that sex differences in the

expression of immune genes are typically conserved in the two species, for most of the upregulated

genes with the key exception of negative regulators but noticeably different for downregulation; also,

there were significant sex differences in the fold-change of up- or downregulation between *D.*

melanogaster and *D. simulans*.

This study will significantly contribute to our understanding of the evolution of sex differences and point

to specific testable hypotheses that might explain observed species and sex differences in survival.

Further research involving both within and between-species comparisons is essential to ascertain

whether these differences are consistent across various contexts. The observed variation in infection

outcomes among different reports, even when using the same species and pathogens, emphasizes the

significance of considering multiple sources of variation within individual studies. Additionally,

conducting comparisons across a diverse range of time points, rather than relying on a single snapshot,

would provide a more comprehensive view of sex dimorphisms occurring at different stages of disease

progression.

Data availability

All raw and processed sequencing data generated in this study can be found on NCBI Gene Expression

Omnibus (GEO; https://www. ncbi.nlm.nih.gov/geo/) (in processing). The authors confirm that all data

supporting this article's conclusions are included in the article, figures, supplement figures, and

supplement tables.

Acknowledgments

We extend our gratitude to the Bloomington Stock Center for providing the flies. We also thank Brian

Lazzaro from the Department of Ecology and Evolutionary Biology at Cornell University for supplying the

bacterial strain. Our appreciation goes to Blake Rochester, Bayley Atkins, Isabella Blair, and Audrey

Powell for their assistance with fly work, and to the other members of the Graze lab for their valuable

discussions. RNA sequencing was conducted at the Genomic Sequencing and Analysis Facility, University

of Texas at Austin.

Funding

This work was supported by grants to RMG from the National Science Foundation (NSF), grant number

DEB-1751296.

Conflict of Interest

The author(s) declare no conflict of interest.

Figures and Tables

Figure 1: Phylogenetic relationship and experimental design. **A.** Phylogeny (represented as a schematic

diagram) of *D. melanogaster* and *D. simulans*, based on (Bhutkar et al. 2007; Paris et al. 2013; Li et al.

2022). *D. melanogaster* and *D. simulans* were used for the experiments, and *D. yakuba* is shown as an

outgroup. Mya: millions of years ago **B.** Experimental Design. The unmated adult flies were infected with

P. rettgeri (PRET) for the treatments and phosphate buffer solutions (PBS) for wound control. RNAseq

was performed from RNA isolated from pooled flies (whole body) at 16hpi in four replicates.

Figure 2: Determination of survival. **A.** Survival curve of *D. melanogaster* (Mel) upon infection with *P.*
rettgeri (PRET) and wound control (PBS). **B.** Survival curve of *D. simulans* (Sim) upon infection with *P.*
rettgeri (PRET) and wound control (PBS). The survival curve was generated with the Kaplan–Meier
estimator, and the statistical analysis was performed using Cox regression. The asterisks (***)
demonstrated the *p*-values < 0.001, and NS represented no significance. All the experimental flies of the
D. simulans were identified dead at 36hpi for females and males.

Figure 3: Bacterial load for *D. melanogaster* and *D. simulans* sexes. Determination of bacterial load for

females and males at 4-hour intervals upon *P. rettgeri* (PRET) infection in *D. melanogaster* and *D.*

simulans. The data were represented for the initial (4, 8, 12hpi) and later (16, 20, 24hpi) stages of the

infection in log10(CFU/ML) between the initial and later stages. None of the sample groups showed

significant differences at any individual time points (4, 8, 12, 16, 20, and 24hpi) for the bacterial load at

p-value < 0.05. Polynomial regression was employed to fit a polynomial curve, where bacterial loads

above the curve were marked in red and those below in blue. All *D. simulans* males died at 24hpi. There

are 6 replicates for each group, except the dmel female at 4 hrs, which has 5 Replicates (one replicate

was dropped as an outlier). The wound control data is not shown; it is available in File S14.

Figure 4: BLUD for *D. melanogaster* and *D. simulans* sexes. Determination of Bacterial Load Upon Death (BLUD) for females and males upon *P. rettgeri* (PRET) infection. The wound control (PBS) showed no death for the experimental groups. The asterisks (*) demonstrated the *p*-values < 0.05, and NS represented no significance.

1004

1005 **Figure 5:** The upset plots represent up- and down-regulated gene counts. **A.** Comparison within species

1006 for females and males in *D. melanogaster*, **B.** Comparison within species for females and males in *D.*

1007 *simulans*. The plots also show the synergistically and antagonistically expressed gene numbers for each

1008 sex in *D. melanogaster* and *D. simulans*. The figure uses an intersection-based representation (for

1009 details, File S4-S5, Table S1).

1010

Figure 6: The volcano plots represent DEGs. **A.** *D. melanogaster* females, **B.** *D. melanogaster* males, **C.** *D.*

simulans females, and **D.** *D. simulans* males represented upon *P. rettgeri* (PRET) infection compared to

wound control (PBS). For *D. melanogaster*, 13588 Normalized Genes, and for *D. simulans*, 13588

Normalized Genes were used to represent DEGs with a cutoff: log2Fold Change ≥ 1; FDR < 0.05. The *D.*

melanogaster female also expressed *AttD* (log2FC = 10.52), which is out of the figure range (for details,

File S4-S5).

1017

1018 **Figure 7:** Up- and down-regulated enriched gene ontology terms biological process (GO: BP) in males

1019 and females. Mel=*D. melanogaster*, Sim=*D. simulans* (for details, Table S2, File S9). The figure uses an

1020 intersection-based representation.

1021

1022

Figure 8: The top 20 up- and down-regulated enriched gene ontology terms (gseGO: BP) in females and

males of *D. melanogaster* upon P. rettgeri (PRET) infection (for details, File S9). Mel = *D. melanogaster*,

UpReg = Up-Regulation, DownReg = Down-Regulation.

1026

Figure 9: The top 20 up- and down-regulated enriched gene ontology terms (gseGO: BP) in females and

males of *D. simulans* upon P. rettgeri (PRET) infection (for details, File S9). Sim = *D. simulans*, UpReg =

Up-Regulation, DownReg = Down-Regulation.

1030

Figure 10: DEGs in females and males of *D. melanogaster* in response to *P. rettgeri* infection for genes in the Toll-IMD-Jak/Stat signaling pathway (wiki pathway ID: WP3830). The expression (z-score) of the pathway is represented without statistical cut-off. Red represents upregulation, green represents downregulation, and white represents no change compared to wound control (for details, File S10, Figure S2).

1036

1037 **Figure 11:** DEGs in females and males of *D. simulans* in response to *P. rettgeri* infection for genes in the

1038 Toll-IMD-Jak/Stat signaling pathway (wiki pathway ID: WP3830). The expression (z-score) of the pathway

1039 was represented without statistical cut-off. Red represents upregulation, green represents

1040 downregulation, and white represents no change compared to wound control (for details, File S10,

1041 Figure S2).

Supplement Information, Figures, and Tables

Table S1: The number of genes upregulated, downregulated, and not significant at FDR < 0.05 for *D. melanogaster* and *D. simulans*. f=Female, m=make, pbs=phosphate buffer solution, pret= *P. rettgeri*, Mel= *D. melanogaster*, Sim= *D. simulans*, DownReg= Down-Regulation, UpReg= Up-Regulation, NotSig= Not significant, FDR= False Discovery Rate (for details, File S4-S5)

Experimental Group (Contrast)	Number of Genes	Species	Sex	Regulation /Types
f_pret_VS_f_pbs_FDR0.05_down	977	Mel	Female	DownReg
f_pret_VS_f_pbs_FDR0.05_up	599	Mel	Female	UpReg
f_pret_VS_f_pbs_FDR0.05_not_sig	12012	Mel	Female	NotSig
m_pret_VS_m_pbs_FDR0.05_down	2269	Mel	Male	DownReg
m_pret_VS_m_pbs_FDR0.05_up	530	Mel	Male	UpReg
m_pret_VS_m_pbs_FDR0.05_not_sig	10789	Mel	Male	NotSig
f_pret_VS_f_pbs_FDR0.05_down	1683	Sim	Female	DownReg
f_pret_VS_f_pbs_FDR0.05_up	633	Sim	Female	UpReg
f_pret_VS_f_pbs_FDR0.05_not_sig	10694	Sim	Female	NotSig
m_pret_VS_m_pbs_FDR0.05_down	1871	Sim	Male	DownReg
m_pret_VS_m_pbs_FDR0.05_up	503	Sim	Male	UpReg
m_pret_VS_m_pbs_FDR0.05_not_sig	10636	Sim	Male	NotSig

Table S2: Up- and down-regulated enriched gene ontology terms biological process (GO: BP) in males and females. Mel=*D. melanogaster*, Sim=*D. simulans*, Com. En. w= Commonly Enriched with, UpReg=Up Regulation, DwReg=Down Regulation (for details, File S9)

Exp Groups	Total Enriched	Up Reg	Com. En. w Opposite Sex-UpReg	Com. En. w compared Species-UpReg	Dn Reg	Com. En. w Opposite Sex-DnReg	Com. En. w compared Species-DnReg
Mel Female	50	49	46	44	1	1	0
Mel Male	101	84	46	56	17	1	12
Sim Female	68	47	42	44	21	0	0
Sim Male	72	57	42	56	15	0	12

1052

Figure S1: Percent of differentially expressed gene (DEGs) in the females (F) and males (M) in *D. melanogaster* (Dmel) and *D. simulans* (Dsim) upon *P. rettgeri* (PRET) infection for different ranges of fold change (FC). **A.** Percentage of up- and downregulated DEGs (without FC cutoff), **B.** Percentage of up- and downregulated DEGs at > 2FC, **C.** Percentage of up- and downregulated DEGs at 2 to 4FC, **D.** Percentage of up and downregulated DEGs at > 4FC. The percentage of expressed genes showed differential expression with cut-off FDR < 0.05 when compared to the wound control (PBS). The expressed genes were defined after filtering for low expression, the number of expressed genes was 13588 and 13010 for *D. melanogaster* and *D. simulans*, respectively (for details, File S4-S5). The percent count of the more than 2-FC downregulated genes showed the difference between females and males in *D. melanogaster* is much more comparable, in contrast to the pattern for downregulation overall (without FC) (Figure S1). In *D. simulans,* downregulation at more than 2-FC showed the opposite direction than overall downregulation (without FC). For the greater than 2-FC cutoff, upregulation is comparable in both species for females and males. For genes with the largest differences in expression, > 4 FC, in *D. simulans*, many more genes are downregulated in females than in males, and a similar pattern is revealed in upregulation. These suggest that while overall downregulation was more prevalent in males for *D. simulans*, when the degree of suppression is dependent on sex, females showed more highly downregulated genes (at > 2-FC, 2-FC < FC < 4-FC, > 4-FC). This is also true for the *D. melanogaster* (at > 2-FC and > 4-FC), although the magnitude of the difference is not as high as in *D. simulans* (Figure S1).

Toll IMD JAK/STAT (modified WP3830)

1071

Figure S2: Sex differences in the Toll-IMD-Jak/Stat signaling pathway (wiki pathway ID: WP3830) in *D. melanogaster* and *D. simulans* upon *P. rettgeri* infection. Mel=*D. melanogaster*, Sim=*D. simulans*, F=Female, M= Male. The expression (z-score) of the pathway was represented without statistical cut-off (for details, File S10)

1076

Figure S3: Sex differences in **A.** positive and **B.** negative regulators of the IMD pathways in *D. melanogaster* and *D. simulans* upon *P. rettgeri* infection. Mel=*D. melanogaster*, Sim=*D. simulans*, F=Female, M= Male. The expression (z-score) of the pathway was represented without statistical cut-off (for details, File S11).

1081

1082

1083 **Figure S4:** Sex differences in **A.** *spz* activating pathway, **B.** positive and **C.** negative regulators of the Toll
1084 pathways in *D. melanogaster* and *D. simulans* upon *P. rettgeri* infection. Mel=*D. melanogaster*, Sim=*D.*
1085 *simulans*, F=Female, M= Male. The expression (z-score) of the pathway was represented without
1086 statistical cut-off (for details, File S11).

1087

1088 **Figure S5:** Sex differences in **A.** Bomanins (Boms) and **B.** melanization in *D. melanogaster* and *D.*
1089 *simulans* upon *P. rettgeri* infection. Mel=*D. melanogaster*, Sim=*D. simulans*, F=Female, M= Male. The
1090 expression (z-score) of the pathway is represented without statistical cut-off (for details, File S11).

1091

1092

Figure S6: Sex differences in **A.** DIMS and IMs, and **B.** Tots, and Teps in *D. melanogaster* and *D. simulans* upon *P. rettgeri* infection. Mel= *D. melanogaster*, Sim= *D. simulans*, F= Female, M= Male, DIMs= Drosophila immune-induced molecules, IMs= Immune Molecules, Tots= Turandot, Teps= Thioester-containing protein, AMPs= Antimicrobial peptides. The expression (z-score) of the pathway was represented without statistical cut-off (for details, File S11).

1098

1099   **Figure S7:** Sex differences in **A.** JAK-STAT (hop-STAT92E) and **B.** Jnk (TNFa-Eiger) pathway with regulators
1100   in *D. melanogaster* and *D. simulans* upon *P. rettgeri* infection. Mel=*D. melanogaster*, Sim=*D. simulans*,
1101   F=Female, M= Male. The expression (z-score) of the pathway was represented without statistical cut-off
1102   (for detail, File S10-S11).

References

Abancens M., V. Bustos, H. Harvey, J. McBryan, and B. J. Harvey, 2020 Sexual Dimorphism in Colon Cancer. Front Oncol 10. https://doi.org/10.3389/fonc.2020.607909

Alejandro A.-D., J.-P. Lilia, M.-B. Jesús, and R. M. Henry, 2022 The IMD and Toll canonical immune pathways of Triatoma pallidipennis are preferentially activated by Gram-negative and Gram-positive bacteria, respectively, but cross-activation also occurs. Parasit Vectors 15: 256. https://doi.org/10.1186/s13071-022-05363-y

Anders S., 2010 Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data. Soil 5: http://www.bioinformatics.babraham.ac.uk/projects/.

Asahina K., 2018 Sex differences in Drosophila behavior: qualitative and quantitative dimorphism. Curr Opin Physiol 6: 35–45. https://doi.org/10.1016/j.cophys.2018.04.004

Banerjee U., J. R. Girard, L. M. Goins, and C. M. Spratford, 2019 Drosophila as a genetic model for hematopoiesis. Genetics 211: 367–417. https://doi.org/10.1534/genetics.118.300223

Basbous N., F. Coste, P. Leone, R. Vincentelli, J. Royet, et al., 2011 The Drosophila peptidoglycan-recognition protein LF interacts with peptidoglycan-recognition protein LC to downregulate the Imd pathway. EMBO Rep 12: 327–333. https://doi.org/10.1038/embor.2011.19

Bateman A. J., 1948 Intra-sexual selection in Drosophila. Heredity (Edinb) 2: 349–368. https://doi.org/10.1038/hdy.1948.21

Beery A. K., and I. Zucker, 2011 Sex bias in neuroscience and biomedical research. Neurosci Biobehav Rev 35: 565–572. https://doi.org/10.1016/j.neubiorev.2010.07.002

Belmonte R. L., M.-K. Corbally, D. F. Duneau, and J. C. Regan, 2020 Sexual Dimorphisms in Innate Immunity and Responses to Infection in Drosophila melanogaster. Front Immunol 10. https://doi.org/10.3389/fimmu.2019.03075

Bhutkar A., S. M. Russo, T. F. Smith, and W. M. Gelbart, 2007 Genome-scale analysis of positionally relocated genes. Genome Res 17: 1880–1887. https://doi.org/10.1101/gr.7062307

Binggeli O., C. Neyen, M. Poidevin, and B. Lemaitre, 2014 Prophenoloxidase Activation Is Required for Survival to Microbial Infections in Drosophila, (D. S. Schneider, Ed.). PLoS Pathog 10: e1004067. https://doi.org/10.1371/journal.ppat.1004067

Bischoff V., C. Vignal, B. Duvic, I. G. Boneca, J. A. Hoffmann, et al., 2006 Downregulation of the Drosophila Immune Response by Peptidoglycan-Recognition Proteins SC1 and SC2, (D. Schneider, Ed.). PLoS Pathog 2: e14. https://doi.org/10.1371/journal.ppat.0020014

Bolger A. M., M. Lohse, and B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30: 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Bourg É. Le, 2011 The NF-kB like factor DIF has weaker effects on Drosophila melanogaster immune defenses than previously thought. Journal of Comparative Physiology B 181: 741–750. https://doi.org/10.1007/s00360-011-0567-1

Carpenter J., S. Hutter, J. F. Baines, J. Roller, S. S. Saminadin-Peter, et al., 2009 The Transcriptional Response of Drosophila melanogaster to Infection with the Sigma Virus (Rhabdoviridae), (S. Bereswill, Ed.). PLoS One 4: e6838. https://doi.org/10.1371/journal.pone.0006838

Carton Y., F. Frey, and A. Nappi, 1992 Genetic determinism of the cellular immune reaction in Drosophila melanogaster. Heredity (Edinb) 69: 393–399. https://doi.org/10.1038/hdy.1992.141

Chamy L. El, V. Leclerc, I. Caldelari, and J.-M. Reichhart, 2008 Sensing of "danger signals" and pathogen-associated molecular patterns defines binary signaling pathways "upstream" of Toll. Nat Immunol 9: 1165–1170. https://doi.org/10.1038/ni.1643

Choe K.-M., H. Lee, and K. V. Anderson, 2005 Drosophila peptidoglycan recognition protein LC (PGRP-LC) acts as a signal-transducing innate immune receptor. Proceedings of the National Academy of Sciences 102: 1122–1126. https://doi.org/10.1073/pnas.0404952102

Chowdhury M., C.-F. Li, Z. He, Y. Lu, X.-S. Liu, et al., 2019 Toll family members bind multiple Spätzle proteins and activate antimicrobial peptide gene expression in Drosophila. Journal of Biological Chemistry 294: 10172–10181. https://doi.org/10.1074/jbc.RA118.006804

Clemmons A. W., S. A. Lindsay, and S. A. Wasserman, 2015 An Effector Peptide Family Required for Drosophila Toll-
Mediated Immunity, (N. Silverman, Ed.). PLoS Pathog 11: e1004876.
https://doi.org/10.1371/journal.ppat.1004876
Darwin C., 1871 Descent of man:Part II Sexual Selection. New York: Prometheus Books.
Deng H., and H. Jasper, 2016 Sexual Dimorphism: How Female Cells Win the Race. Current Biology 26: R212–R215.
https://doi.org/10.1016/j.cub.2016.01.062
Dobin A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, et al., 2013 STAR: ultrafast universal RNA-seq aligner.
Bioinformatics 29: 15–21. https://doi.org/10.1093/bioinformatics/bts635
Dudzic J. P., S. Kondo, R. Ueda, C. M. Bergman, and B. Lemaitre, 2015 Drosophila innate immunity: regional and
functional specialization of prophenoloxidases. BMC Biol 13: 81. https://doi.org/10.1186/s12915-015-0193-6
Dudzic J. P., M. A. Hanson, I. Iatsenko, S. Kondo, and B. Lemaitre, 2019 More Than Black or White: Melanization
and Toll Share Regulatory Serine Proteases in Drosophila. Cell Rep 27: 1050-1061.e3.
https://doi.org/10.1016/j.celrep.2019.03.101
Duneau D., J.-B. Ferdy, J. Revah, H. Kondolf, G. A. Ortiz, et al., 2017a Stochastic variation in the initial phase of
bacterial infection predicts the probability of survival in D. melanogaster. Elife 6.
https://doi.org/10.7554/eLife.28298
Duneau D., H. Kondolf, J. H. Im, G. Ortiz, C. Chow, et al., 2017b The Toll pathway underlies host sexual dimorphism
in resistance to both Gram-negative and Gram-positive bacteria in mated Drosophila. BMC Biol 15: 124.
https://doi.org/10.1186/s12915-017-0466-3
Evans C. J., V. Hartenstein, and U. Banerjee, 2003 Thicker than blood: Conserved mechanisms in Drosophila and
vertebrate hematopoiesis. Dev Cell 5: 673–690. https://doi.org/10.1016/S1534-5807(03)00335-6
Fischer J., N. Jung, N. Robinson, and C. Lehmann, 2015 Sex differences in immune responses to infectious diseases.
Infection 43: 399–403. https://doi.org/10.1007/s15010-015-0791-9
Franchet A., S. Niehus, G. Caravello, and D. Ferrandon, 2019 Phosphatidic acid as a limiting host metabolite for the
proliferation of the microsporidium Tubulinosema ratisbonensis in Drosophila flies. Nat Microbiol 4: 645–
655. https://doi.org/10.1038/s41564-018-0344-y
Galac M. R., and B. P. Lazzaro, 2011 Comparative pathology of bacteria in the genus Providencia to a natural host,
Drosophila melanogaster. Microbes Infect 13: 673–683. https://doi.org/10.1016/j.micinf.2011.02.005
Gal-Oz S. T., B. Maier, H. Yoshida, K. Seddu, N. Elbaz, et al., 2019 ImmGen report: sexual dimorphism in the
immune system transcriptome. Nat Commun 10: 4295. https://doi.org/10.1038/s41467-019-12348-6
Gan T., L. Fan, L. Zhao, M. Misra, M. Liu, et al., 2021 JNK Signaling in Drosophila Aging and Longevity. Int J Mol Sci
22: 9649. https://doi.org/10.3390/ijms22179649
Gelbart W. M., W. P. Rindone, J. Chillemi, S. Russo, M. Crosby, et al., 1996 FlyBase: The Drosophila database.
Nucleic Acids Res 24: 53–56. https://doi.org/10.1093/nar/24.1.53
Girgenti M. J., S. Pothula, and S. S. Newton, 2021 Stress and Its Impact on the Transcriptome. Biol Psychiatry 90:
102–108. https://doi.org/10.1016/j.biopsych.2020.12.011
Gottar M., V. Gobert, A. A. Matskevich, J.-M. Reichhart, C. Wang, et al., 2006 Dual Detection of Fungal Infections in
Drosophila via Recognition of Glucans and Sensing of Virulence Factors. Cell 127: 1425–1437.
https://doi.org/10.1016/j.cell.2006.10.046
Gramates L. S., J. Agapite, H. Attrill, B. R. Calvi, M. A. Crosby, et al., 2022 FlyBase: a guided tour of highlighted
features, (V. Wood, Ed.). Genetics 220. https://doi.org/10.1093/genetics/iyac035
Graze R. M., R. Y. Tzeng, T. S. Howard, and M. N. Arbeitman, 2018 Perturbation of IIS/TOR signaling alters the
landscape of sex-differential gene expression in Drosophila. BMC Genomics 19: 893.
https://doi.org/10.1186/s12864-018-5308-3
Guo L., J. Karpac, S. L. Tran, and H. Jasper, 2014 PGRP-SC2 Promotes Gut Immune Homeostasis to Limit Commensal
Dysbiosis and Extend Lifespan. Cell 156: 109–122. https://doi.org/10.1016/j.cell.2013.12.018
Hanson M. A., and B. Lemaitre, 2020 New insights on Drosophila antimicrobial peptide function in host defense
and beyond. Curr Opin Immunol 62: 22–30. https://doi.org/10.1016/j.coi.2019.11.008
Heintze J. M., 2018 TLR7 evades silencing in human immune cells. Nat Rev Nephrol 14: 214–214.
https://doi.org/10.1038/nrneph.2018.13
Hill-Burns E. M., and A. G. Clark, 2010 Functional Regulatory Divergence of the Innate Immune System in
Interspecific Drosophila Hybrids. Mol Biol Evol 27: 2596–2605. https://doi.org/10.1093/molbev/msq146

Hoffmann J. A., 2003 The immune response of Drosophila. Nature 426: 33–38.
https://doi.org/10.1038/nature02021
Howick V. M., and B. P. Lazzaro, 2014 Genotype and diet shape resistance and tolerance across distinct phases of
bacterial infection. BMC Evol Biol 14: 56. https://doi.org/10.1186/1471-2148-14-56
Issa N., N. Guillaumot, E. Lauret, N. Matt, C. Schaeffer-Reiss, et al., 2018 The Circulating Protease Persephone Is an
Immune Sensor for Microbial Proteolytic Activities Upstream of the Drosophila Toll Pathway. Mol Cell 69:
539-550.e6. https://doi.org/10.1016/j.molcel.2018.01.029
Ji Y., C. Thomas, N. Tulin, N. Lodhi, E. Boamah, et al., 2016 Charon Mediates Immune Deficiency–Driven PARP-1–
Dependent Immune Responses in Drosophila. The Journal of Immunology 197: 2382–2389.
https://doi.org/10.4049/jimmunol.1600994
Juneja P., and B. P. Lazzaro, 2009 Providencia sneebia sp. nov. and Providencia burhodogranariea sp. nov., isolated
from wild Drosophila melanogaster. Int J Syst Evol Microbiol 59: 1108–1111.
https://doi.org/10.1099/ijs.0.000117-0
Khalil S., E. Jacobson, M. C. Chambers, and B. P. Lazzaro, 2015 Systemic Bacterial Infection and Immune Defense
Phenotypes in Drosophila Melanogaster. Journal of Visualized Experiments. https://doi.org/10.3791/52613
Khan M., and R. Graze, 2024 Regulation of sex differences in innate immunity by sex-determining gene transformer
in Drosophila melanogaster. bioRxiv. https://doi.org/doi: https://doi.org/10.1101/2024.05.28.596272
Khodursky S., N. Svetec, S. M. Durkin, and L. Zhao, 2020 The evolution of sex-biased gene expression in the
Drosophila brain. Genome Res 30: 874–884. https://doi.org/10.1101/gr.259069.119
Klein S. L., and K. L. Flanagan, 2016 Sex differences in immune responses. Nat Rev Immunol 16: 626–638.
https://doi.org/10.1038/nri.2016.90
Kutch I. C., and K. M. Fedorka, 2017 A test for Y-linked additive and epistatic effects on surviving bacterial
infections in Drosophila melanogaster. J Evol Biol 30: 1400–1408. https://doi.org/10.1111/jeb.13118
Lai Y., and R. L. Gallo, 2009 AMPed up immunity: how antimicrobial peptides have multiple roles in immune
defense. Trends Immunol 30: 131–141. https://doi.org/10.1016/j.it.2008.12.003
Lande R., 1980 Sexual Dimorphism, Sexual Selection, and Adaptation in Polygenic Characters. Evolution (N Y) 34:
292. https://doi.org/10.2307/2407393
Law C. W., Y. Chen, W. Shi, and G. K. Smyth, 2014 voom: precision weights unlock linear model analysis tools for
RNA-seq read counts. Genome Biol 15: R29. https://doi.org/10.1186/gb-2014-15-2-r29
Lazzaro B. P., and T. J. Little, 2009 Immunity in a variable world. Philosophical Transactions of the Royal Society B:
Biological Sciences 364: 15–26. https://doi.org/10.1098/rstb.2008.0141
Lecheta M. C., D. N. Awde, T. S. O'Leary, L. N. Unfried, N. A. Jacobs, et al., 2020 Integrating GWAS and
Transcriptomics to Identify the Molecular Underpinnings of Thermal Stress Responses in Drosophila
melanogaster. Front Genet 11. https://doi.org/10.3389/fgene.2020.00658
Leech T., S. E. F. Evison, S. A. O. Armitage, S. M. Sait, and A. Bretman, 2019 Interactive effects of social
environment, age and sex on immune responses in Drosophila melanogaster. J Evol Biol 32: 1082–1092.
https://doi.org/10.1111/jeb.13509
Li F., R. V. Rane, V. Luria, Z. Xiong, J. Chen, et al., 2022 Phylogenomic analyses of the genus Drosophila reveals
genomic signals of climate adaptation. Mol Ecol Resour 22: 1559–1581. https://doi.org/10.1111/1755-
0998.13561
Libert C., L. Dejager, and I. Pinheiro, 2010 The X chromosome in immune functions: when a chromosome makes
the difference. Nat Rev Immunol 10: 594–604. https://doi.org/10.1038/nri2815
Ligoxygakis P., N. Pelte, J. A. Hoffmann, and J.-M. Reichhart, 2002 Activation of Drosophila Toll During Fungal
Infection by a Blood Serine Protease. Science (1979) 297: 114–116. https://doi.org/10.1126/science.1072391
Ligoxygakis P., 2002 A serpin mutant links Toll activation to melanization in the host defence of Drosophila. EMBO J
21: 6330–6337. https://doi.org/10.1093/emboj/cdf661
Lin S. J. H., L. B. Cohen, and S. A. Wasserman, 2020 Effector specificity and function in Drosophila innate immunity:
Getting AMPed and dropping Boms, (N. Silverman, Ed.). PLoS Pathog 16: e1008480.
https://doi.org/10.1371/journal.ppat.1008480
Lohman B. K., J. N. Weber, and D. I. Bolnick, 2016 Evaluation of TagSeq, a reliable low-cost alternative for RNAseq.
Mol Ecol Resour 16: 1315–1321. https://doi.org/10.1111/1755-0998.12529

Lopes-Ramos C. M., C. Y. Chen, M. L. Kuijjer, J. N. Paulson, A. R. Sonawane, et al., 2020 Sex Differences in Gene
Expression and Regulatory Networks across 29 Human Tissues. Cell Rep 31.
https://doi.org/10.1016/j.celrep.2020.107795

Ma F., B. K. Fuqua, Y. Hasin, C. Yukhtman, C. D. Vulpe, et al., 2019 A comparison between whole transcript and 3'
RNA sequencing methods using Kapa and Lexogen library preparation methods. BMC Genomics 20: 9.
https://doi.org/10.1186/s12864-018-5393-3

Mackenzie D. K., L. F. Bussière, and M. C. Tinsley, 2011 Senescence of the cellular immune response in Drosophila
melanogaster. Exp Gerontol 46: 853–859. https://doi.org/10.1016/j.exger.2011.07.004

Maillet F., V. Bischoff, C. Vignal, J. Hoffmann, and J. Royet, 2008 The Drosophila Peptidoglycan Recognition Protein
PGRP-LF Blocks PGRP-LC and IMD/JNK Pathway Activation. Cell Host Microbe 3: 293–303.
https://doi.org/10.1016/j.chom.2008.04.002

Martin M., A. Hiroyasu, R. M. Guzman, S. A. Roberts, and A. G. Goodman, 2018 Analysis of Drosophila STING
Reveals an Evolutionarily Conserved Antimicrobial Function. Cell Rep 23: 3537-3550.e6.
https://doi.org/10.1016/j.celrep.2018.05.029

Matskevich A. A., J. Quintin, and D. Ferrandon, 2010 The Drosophila PRR GNBP3 assembles effector complexes
involved in antifungal defenses independently of its Toll -pathway activation function. Eur J Immunol 40:
1244–1254. https://doi.org/10.1002/eji.200940164

McDonald J. M. C., P. Nabili, L. Thorsen, S. Jeon, and A. W. Shingleton, 2021 Sex-specific plasticity and the
nutritional geometry of insulin-signaling gene expression in Drosophila melanogaster. Evodevo 12: 6.
https://doi.org/10.1186/s13227-021-00175-0

McKean K. A., and L. Nunney, 2001 Increased sexual activity reduces male immune function in Drosophila
melanogaster. Proceedings of the National Academy of Sciences 98: 7904–7909.
https://doi.org/10.1073/pnas.131216398

Metcalf C. J. E., O. Roth, and A. L. Graham, 2020 Why leveraging sex differences in immune trade-offs may
illuminate the evolution of senescence, (J. Lemaître, Ed.). Funct Ecol 34: 129–140.
https://doi.org/10.1111/1365-2435.13458

MEYER E., G. V. AGLYAMOVA, and M. V. MATZ, 2011 Profiling gene expression responses of coral larvae (Acropora
millepora) to elevated temperature and settlement inducers using a novel RNA-Seq procedure. Mol Ecol no-
no. https://doi.org/10.1111/j.1365-294X.2011.05205.x

Millington J. W., and E. J. Rideout, 2018 Sex differences in Drosophila development and physiology. Curr Opin
Physiol 6: 46–56. https://doi.org/10.1016/j.cophys.2018.04.002

Mondal S., and U. Rai, 1999 Sexual Dimorphism in Phagocytic Activity of Wall Lizard's Splenic Macrophages and Its
Control by Sex Steroids. Gen Comp Endocrinol 116: 291–298. https://doi.org/10.1006/gcen.1999.7370

Morrow E. H., 2015 The evolution of sex differences in disease. Biol Sex Differ 6: 5.
https://doi.org/10.1186/s13293-015-0023-0

Myllymäki H., S. Valanne, and M. Rämet, 2014 The Drosophila Imd Signaling Pathway. The Journal of Immunology
192: 3455–3462. https://doi.org/10.4049/jimmunol.1303309

Myllymäki H., and M. Rämet, 2014 JAK/STAT Pathway in Drosophila Immunity. Scand J Immunol 79: 377–385.
https://doi.org/10.1111/sji.12170

Nanni A. V, N. Martinez, R. Graze, A. Morse, J. R. B. Newman, et al., 2023 Sex-Biased Expression Is Associated With
Chromatin State in Drosophila melanogaster and Drosophila simulans, (P. Wittkopp, Ed.). Mol Biol Evol 40.
https://doi.org/10.1093/molbev/msad078

Oertelt-Prigione S., 2012 The influence of sex and gender on the immune response. Autoimmun Rev 11: A479–
A485. https://doi.org/10.1016/j.autrev.2011.11.022

Palmer W. H., N. C. Medd, P. M. Beard, and D. J. Obbard, 2018 Isolation of a natural DNA virus of Drosophila
melanogaster, and characterisation of host resistance and immune responses, (J.-L. Imler, Ed.). PLoS Pathog
14: e1007050. https://doi.org/10.1371/journal.ppat.1007050

Paris M., T. Kaplan, X. Y. Li, J. E. Villalta, S. E. Lott, et al., 2013 Extensive Divergence of Transcription Factor Binding
in Drosophila Embryos with Highly Conserved Gene Expression, (P. Wittkopp, Ed.). PLoS Genet 9: e1003748.
https://doi.org/10.1371/journal.pgen.1003748

Persson C., S. Oldenvi, and H. Steiner, 2007 Peptidoglycan recognition protein LF: A negative regulator of
Drosophila immunity. Insect Biochem Mol Biol 37: 1309–1316. https://doi.org/10.1016/j.ibmb.2007.08.003

Pinheiro I., and E. Heard, 2017 X chromosome inactivation: new players in the initiation of gene silencing.
F1000Res 6: 344. https://doi.org/10.12688/f1000research.10707.1

Rai K. E., H. Yin, A. L. C. Bengo, M. Cheek, R. Courville, et al., 2023 Immune defense in Drosophila melanogaster
depends on diet, sex, and mating status. PLoS One 18: 1–20. https://doi.org/10.1371/journal.pone.0268415

Regan J. C., A. S. Brandão, A. B. Leitão, Â. R. Mantas Dias, É. Sucena, et al., 2013 Steroid Hormone Signaling Is
Essential to Regulate Innate Immune Cells and Fight Bacterial Infection in Drosophila, (D. S. Schneider, Ed.).
PLoS Pathog 9: e1003720. https://doi.org/10.1371/journal.ppat.1003720

Reichhart J. M., D. Gubb, and V. Leclerc, 2011 The Drosophila serpins: Multiple functions in immunity and
morphogenesis, pp. 205–225 in Methods in Enzymology,.

Rideout E. J., M. S. Narsaiya, and S. S. Grewal, 2015 The Sex Determination Gene transformer Regulates Male-
Female Differences in Drosophila Body Size, (A. Kopp, Ed.). PLoS Genet 11: e1005683.
https://doi.org/10.1371/journal.pgen.1005683

Ritchie M. E., B. Phipson, D. Wu, Y. Hu, C. W. Law, et al., 2015 limma powers differential expression analyses for
RNA-sequencing and microarray studies. Nucleic Acids Res 43: e47–e47. https://doi.org/10.1093/nar/gkv007

Robinson M. D., D. J. McCarthy, and G. K. Smyth, 2010 edgeR : a Bioconductor package for differential expression
analysis of digital gene expression data. Bioinformatics 26: 139–140.
https://doi.org/10.1093/bioinformatics/btp616

Scharf I., F. Peter, and O. Y. Martin, 2013 Reproductive Trade-Offs and Direct Costs for Males in Arthropods. Evol
Biol 40: 169–184. https://doi.org/10.1007/s11692-012-9213-4

Schlamp F., S. Y. N. Delbare, A. M. Early, M. T. Wells, S. Basu, et al., 2021 Dense time-course gene expression
profiling of the Drosophila melanogaster innate immune response. BMC Genomics 22.
https://doi.org/10.1186/s12864-021-07593-3

Schwenke R. A., B. P. Lazzaro, and M. F. Wolfner, 2016 Reproduction–Immunity Trade-Offs in Insects. Annu Rev
Entomol 61: 239–256. https://doi.org/10.1146/annurev-ento-010715-023924

Schwenke R. A., and B. P. Lazzaro, 2017 Juvenile Hormone Suppresses Resistance to Infection in Mated Female
Drosophila melanogaster. Current Biology 27: 596–601. https://doi.org/10.1016/j.cub.2017.01.004

Scofield R. H., G. R. Bruner, B. Namjou, R. P. Kimberly, R. Ramsey-Goldman, et al., 2008 Klinefelter's syndrome
(47,XXY) in male systemic lupus erythematosus patients: Support for the notion of a gene-dose effect from
the X chromosome. Arthritis Rheum 58: 2511–2517. https://doi.org/10.1002/art.23701

Shahrestani P., M. Chambers, J. Vandenberg, K. Garcia, G. Malaret, et al., 2018 Sexual dimorphism in Drosophila
melanogaster survival of Beauveria bassiana infection depends on core immune signaling. Sci Rep 8: 12501.
https://doi.org/10.1038/s41598-018-30527-1

Shepherd R., A. S. Cheung, K. Pang, R. Saffery, and B. Novakovic, 2021 Sexual Dimorphism in Innate Immunity: The
Role of Sex Hormones and Epigenetics. Front Immunol 11. https://doi.org/10.3389/fimmu.2020.604000

Short S. M., and B. P. Lazzaro, 2010 Female and male genetic contributions to post-mating immune defence in
female Drosophila melanogaster. Proceedings of the Royal Society B: Biological Sciences 277: 3649–3657.
https://doi.org/10.1098/rspb.2010.0937

Short S. M., M. F. Wolfner, and B. P. Lazzaro, 2012 Female Drosophila melanogaster suffer reduced defense against
infection due to seminal fluid components. J Insect Physiol 58: 1192–1201.
https://doi.org/10.1016/j.jinsphys.2012.06.002

Souyris M., C. Cenac, P. Azar, D. Daviaud, A. Canivet, et al., 2018 TLR7 escapes X chromosome inactivation in
immune cells. Sci Immunol 3. https://doi.org/10.1126/sciimmunol.aap8855

Sparks C., 2017 Comparing multiple groups with the linear model - ANOVA.
https://rpubs.com/corey_sparks/312725.

Tafesh-Edwards G., and I. Eleftherianos, 2020 JNK signaling in Drosophila immunity and homeostasis. Immunol Lett
226: 7–11. https://doi.org/10.1016/j.imlet.2020.06.017

Tang H., 2009 Regulation and function of the melanization reaction in Drosophila. Fly (Austin) 3: 105–111.
https://doi.org/10.4161/fly.3.1.7747

Tanji T., and Y. T. Ip, 2005 Regulators of the Toll and Imd pathways in the Drosophila innate immune response.
Trends Immunol 26: 193–198. https://doi.org/10.1016/j.it.2005.02.006

Taylor K., and D. Kimbrell, 2007 Host Immune Response and Differential Survival of the Sexes in Drosophila. Fly
(Austin) 1: 197–204. https://doi.org/10.4161/fly.5082

Therneau T., and P. Grambsch, 2000 Modeling Survival Data: Extending the Cox Model, in Springer, New York.

Therneau T., 2023 A Package for Survival Analysis in R

Thomson T. C., A. Schneemann, and J. Johnson, 2012 Oocyte destruction is activated during viral infection. genesis
50: 453–465. https://doi.org/10.1002/dvg.22004

Tzou P., S. Ohresser, D. Ferrandon, M. Capovilla, J.-M. Reichhart, et al., 2000 Tissue-Specific Inducible Expression of
Antimicrobial Peptide Genes in Drosophila Surface Epithelia. Immunity 13: 737–748.
https://doi.org/10.1016/S1074-7613(00)00072-8

Uttenweiler-Joseph S., M. Moniatte, M. Lagueux, A. Van Dorsselaer, J. A. Hoffmann, et al., 1998 Differential display
of peptides induced during the immune response of Drosophila : A matrix-assisted laser desorption
ionization time-of-flight mass spectrometry study. Proceedings of the National Academy of Sciences 95:
11342–11347. https://doi.org/10.1073/pnas.95.19.11342

Valanne S., J.-H. Wang, and M. Rämet, 2011 The Drosophila Toll Signaling Pathway. The Journal of Immunology
186: 649–656. https://doi.org/10.4049/jimmunol.1002302

Valanne S., L. Vesala, M. K. Maasdorp, T. S. Salminen, and M. Rämet, 2022 The Drosophila Toll Pathway in Innate
Immunity: from the Core Pathway toward Effector Functions. The Journal of Immunology 209: 1817–1825.
https://doi.org/10.4049/jimmunol.2200476

Vincent C. M., and N. P. Sharp, 2014 Sexual antagonism for resistance and tolerance to infection in Drosophila
melanogaster. Proceedings of the Royal Society B: Biological Sciences 281: 20140987.
https://doi.org/10.1098/rspb.2014.0987

Vincent C. M., and M. S. Dionne, 2021 Disparate regulation of IMD signaling drives sex differences in infection
pathology in Drosophila melanogaster. Proc Natl Acad Sci U S A 118.
https://doi.org/10.1073/pnas.2026554118

Vlisidou I., and W. Wood, 2015 Drosophila blood cells and their role in immune responses. FEBS J 282: 1368–1382.
https://doi.org/10.1111/febs.13235

Wilkinson N. M., H.-C. Chen, M. G. Lechner, and M. A. Su, 2022 Sex Differences in Immunity. Annu Rev Immunol
40: 75–94. https://doi.org/10.1146/annurev-immunol-101320-125133

Wu T., E. Hu, S. Xu, M. Chen, P. Guo, et al., 2021 clusterProfiler 4.0: A universal enrichment tool for interpreting
omics data. The Innovation 2: 100141. https://doi.org/10.1016/j.xinn.2021.100141

Yamamoto-Hino M., and S. Goto, 2016 Spätzle-Processing Enzyme-independent Activation of the Toll Pathway in
Drosophila Innate Immunity. Cell Struct Funct 41: 55–60. https://doi.org/10.1247/csf.16002

Yu S., F. Luo, Y. Xu, Y. Zhang, and L. H. Jin, 2022 Drosophila Innate Immunity Involves Multiple Signaling Pathways
and Coordinated Communication Between Different Tissues. Front Immunol 13.
https://doi.org/10.3389/fimmu.2022.905370

Zaidman-Rémy A., M. Hervé, M. Poidevin, S. Pili-Floury, M.-S. Kim, et al., 2006 The Drosophila Amidase PGRP-LB
Modulates the Immune Response to Bacterial Infection. Immunity 24: 463–473.
https://doi.org/10.1016/j.immuni.2006.02.012

Zaidman-Rémy A., M. Poidevin, M. Hervé, D. P. Welchman, J. C. Paredes, et al., 2011 Drosophila Immunity: Analysis
of PGRP-SB1 Expression, Enzymatic Activity and Function, (S. Bereswill, Ed.). PLoS One 6: e17231.
https://doi.org/10.1371/journal.pone.0017231

Zuk M., and A. M. Stoehr, 2002 Immune Defense and Host Life History. Am Nat 160: S9–S22.
https://doi.org/10.1086/342131

Zuk M., 2009 The Sicker Sex, (G. F. Rall, Ed.). PLoS Pathog 5: e1000267.
https://doi.org/10.1371/journal.ppat.1000267

1403