pH dependent direct sulfhydrylation pathway is required for pathogenesis of *Mycobacterium tuberculosis*

Vaibhav Kumar Nain[1,2], Vishawjeet Barik[1,2#], Manitosh Pandey[1#], Mohit Pareek[3], Taruna Sharma[1,2],

Rahul Pal[1], Shaifali Tyagi[1], Manish Bajpai[4], Prabhanjan Dwivedi[5], Bhishma Narayan Panda[5],

Yashwant Kumar[4], Shailendra Asthana[3] and Amit Kumar Pandey[1,5*]

[1]Mycobacterial Pathogenesis Laboratory, Centre for Tuberculosis Research, BRIC-Translational

Health Science and Technology Institute, Haryana, India

[2]Special Centre for Molecular Medicine, Jawaharlal Nehru University, New Delhi, India

[3]Computational Biophysics and CADD Group, Computational and Mathematical Biology Centre,

BRIC-Translational Health Science and Technology Institute, Haryana, India

[4]Biomarker Discovery Laboratory, Non-Communicable Diseases Centre, BRIC-Translational Health

Science and Technology Institute, Haryana, India

[5]Experimental Animal Facility, BRIC-Translational Health Science and Technology Institute,

Haryana, India

[#] Contributed equally to this work

* Correspondence: Amit Kumar Pandey, amitpandey@thsti.res.in

Abstract

Methionine is essential for the survival of *Mycobacterium tuberculosis* (*M. tuberculosis*) inside the

host. Inhibiting the transsulfuration pathway fails to exhibit methionine auxotrophy suggesting the

presence of an alternate methionine biosynthesis pathway in *M. tuberculosis*. Orthologues of MetZ$_{TB}$

in other bacterial species are known to regulate a redundant single step methionine biosynthesis

pathway known as *direct sulfhydrylation.* Genetic disruption of the *metZ*-mediated direct sulfhydrylation pathway impaired the ability of *M. tuberculosis* to survive inside macrophages. Further, we demonstrated that direct sulfhydrylation pathway is indispensable for the survival at low pH. Using *in-silico* molecular docking and MD simulations studies we showed that the active site of MetZ binds substrate O-succinyl homoserine (OSH) more tightly at acidic rather than neutral pH. Intriguingly, despite higher intracellular ATP levels, a relative decrease in the frequency of generation of Bedaquiline (BDQ)-induced persisters in *ΔmetZ* suggests a role of direct sulfhydrylation pathway in modulating BDQ sensitivity in *M. tuberculosis*. Finally, using an animal model, we demonstrated that absence of *metZ* impeded the ability of *M. tuberculosis* to grow inside the host. Our findings suggest that administering inhibitors of *metZ* as an adjunct can be explored as a strategy to potentiate BDQ as a novel anti-TB regimen.

Keywords: tuberculosis, methionine biosynthesis, acidic pH, drug tolerance, transsulfuration, direct sulfhydrylation.

Introduction

Tuberculosis is one of the ancient diseases that remains a global threat to this day posing a great socio-economic threat to global public healthcare. As of 2023, *Mycobacterium tuberculosis* (*M. tuberculosis*) caused 1.8 million deaths making it the highest killing disease [1]. A gradual increase in the frequency of emergence of drug resistance cases has further complicated the therapeutic landscape against tuberculosis. A majority of drugs approved for treating tuberculosis are focused on targeting essential genes critical for survival of *M. tuberculosis* such as transcription, lipid biosynthesis, ATP synthesis, protein synthesis, etc. Targeting essential amino acid biosynthesis pathways (such as tryptophan,

arginine, proline, histidine, aspartate, etc.) as a therapeutic strategy against tuberculosis has

demonstrated immense potential and might lead to molecules with the potential to become a future

drug [2].

Methionine is required for maintenance of various biochemical processes to sustain life and regulation

of methionine biosynthesis becomes more consequential as, *N-formyl*-L-methionine (*f*Met), is known

to initiate translation in prokaryotes [3] and protein degradation in eukaryotes [4]. Further, it is required

for biosynthesis of other key metabolites – cysteine, biotin, SAM (S-adenosylmethionine), etc.

involved in several physiological and regulatory mechanisms. Historically, targeting the methionine

biosynthesis pathway as a strategy to identify new drugs against tuberculosis has been extensively

explored in various studies [5-9]. The transsulfuration pathway consisting of a sequence of reactions,

enzymes and intermediates involved in methionine biosynthesis are largely conserved across various

microbial species and has been extensively reviewed [10]. However, other than *metA/X, metH* genes

which have been reported to be essential for the growth of *M. tuberculosis in-vitro* as well as *in-vivo*

[5, 7, 11], some of the genes belonging to this pathway are dispensable for growth of *M. tuberculosis*

and hence, are either redundant or are conditionally essential [12-14]. There are reports which suggest

that there exists another biochemical reaction wherein O-succinyl homoserine (OSH) is directly

converted into homocysteine [13, 15]. Interestingly, biological H_2S, instead of cysteine is used as S-

donor in this reaction driven by an enzyme O-succinyl homoserine sulfhydrylase (OSHS) that bypasses

the two-step conversion of OSH to homocysteine, via cystathionine as intermediate step. This direct

conversion of OSH into homocysteine is called direct sulfhydrylation. Still uncharacterized in *M.*

tuberculosis, direct sulfhydrylation pathway is functional in several microbial species such as

Pseudomonas aeruginosa [16], Leptospira [17] and *Corynebacterium glutamicum [10, 18]. M.*

tuberculosis harbors a putative OSHS encoded by *metZ* (*Rv0391*) which is yet to be studied for its role

in direct sulfhydrylation pathway.

In this study, we have used genetics and computational biology-based tools to address the necessity of

M. tuberculosis to harbor redundant methionine biosynthesis pathways. We find that *metZ* null mutant

in *M. tuberculosis* is dispensable for growth at neutral pH, however, it is required for survival of the

bacterium at acidic pH, *ex-vivo* and *in-vivo*. It was also observed using computational approach that

MetZ protein, although stable at both acidic and neutral pH, the active site of the protein held the

substrate more stably at acidic pH than at neutral pH. Finally, using CRISPRi we prove that *metZ* in

M. tuberculosis is independent of transsulfuration pathway and involved in methionine biosynthesis as

corroborated by methionine levels in the mutant strain lacking both transsulfuration as well as direct

sulfhydrylation. In sum, we report that *metZ* is required for methionine biosynthesis via direct

sulfhydrylation pathway at acidic pH when transsulfuration fails to keep up with the cellular

requirements.

Results

MetZ is dispensable for the *in-vitro* growth of *M. tuberculosis* and possibly involved in direct

sulfhydrylation pathway for methionine biosynthesis

While *metZ* has been reported to be essential for growth in cholesterol [19], we hypothesize that the

Rv0391 (metZ) gene encodes a putative O-succinylhomoserine sulfhydrylase (OSHS) that is

responsible for direct sulfhydrylation (Fig. 1A). We performed a bioinformatics-based analysis using

the protein sequence of $MetZ_{TB}$ and found that the putative sequence belonged to the cluster of

orthologues group ENOG502GJ5S which is composed of 1097 protein sequences orthologous to

$MetZ_{TB}$ across 644 bacterial species. This cluster of proteins comprises various enzymes involved in

cellular amino acid metabolism namely, cysteine, methionine, homocysteine, serine and aspartate

family of amino acids. Interestingly, all these proteins have common PLP (pyridoxal-5-phosphate) –

binding domain which is essentially a modified lysine residue that covalently binds the co-factor, hence

also categorized as PLP-binding superfamily of proteins. (Fig. 1B and S1A). Further, to elucidate the

role of *metZ* in *M. tuberculosis*, a deletion mutant in the genetic background of H37Rv was generated

by genetic recombineering as described previously [20]. Briefly, a hygromycin resistance cassette was

inserted into the ORF encoding *metZ* (*Rv0391*) with the help of allelic exchange substrate (AES)

consisting of 1kb regions of homology upstream and downstream to the target locus. The recombinants

were screened for hygromycin resistance and the cassette was subsequently excised out thereby

generating unmarked mutant (Fig. 1C). The loss of the genetic sequence between 470034-471188 bases

was confirmed by gene-specific PCR (Fig. S2A), DNA sequencing of the locus (Fig. S2B) and at the

transcript level by RT-qPCR (Fig 1D). In order to check the essentiality of the *metZ* in growth in

cholesterol, we compared the growth of H37Rv, *ΔmetZ* and the complemented strains in different

media containing glycerol (0.1%), cholesterol (150μM), propionate (10mM) as the sole carbon source.

As a control, we used the *M. tuberculosis* strain lacking the *prpR* gene (*ΔprpR*) which is known to be

essential for growth in cholesterol [21]. While, as expected, *ΔprpR* failed to grow on cholesterol and

propionate, the growth of all the three strains were comparable irrespective to the carbon source (Figs.

1E, 1F and S3B). These findings indicate *metZ* does not contribute to the central carbon metabolism

and is dispensable for *in-vitro* growth in *M. tuberculosis* on cholesterol and/ or propionate.

***metZ* deletion renders *M. tuberculosis* sensitive to physiological stressors and lead to reduced *ex-**

***vivo* fitness**

In order to replicate and grow inside the host, intracellular pathogens have evolved mechanisms to

neutralize the host-induced physiological stressors which include redox, cell membrane and pH stress.

These *in-vitro* induced stress conditions mimic *in-vivo* growth conditions, wherein bacterial cells were

subjected to different stress inducing reagents and their survival index was determined w.r.t. *Rv*-WT.

ΔmetZ, when compared to WT and complement strains, demonstrated a reduced ability (~40-60%) to

survive under all the stress conditions tested (Fig. 2A). Further, we determined intrabacterial redox

state using genetically encoded redox biosensor, Mrx1-roGFP2 and observed a significant rise in intracellular ROS in $\Delta metZ$ compared to WT strain (Fig. 2B) suggesting an inability of the strain to maintain redox homeostasis possibly due to impaired mycothiol recycling. Subsequently, increased levels of lipid hydroperoxides in $\Delta metZ$ as opposed to WT and complement strains corroborate with our findings that the absence of *metZ* in *M. tuberculosis* results in excess accumulation of intracellular ROS (Fig. 2C). We then checked the role of *metZ* in the intracellular survival of the *M. tuberculosis* inside the macrophages. Briefly, resting or IFN-γ induced THP-1 derived macrophages were infected with H37Rv, $\Delta metZ$ and the complemented strain and survival respective survival were monitored by CFU plating of the macrophage lysates at day 5 post infection. We observed that $\Delta metZ$ displayed a reduced bacterial burden (~60%) as compared to WT and complement strains inside resting macrophages, which was further significantly reduced (~48%) inside activated macrophages (Fig. 2D). Since restricting the phagolysosomal fusion is a common strategy employed by the pathogen to increase its chances of survival inside the cell, we checked the role of *metZ* in preventing the fusion of phagolysosomes. Briefly, IFN-γ induced THP1 macrophages infected with GFP expressing wild type, $\Delta metZ$ and complemented strains were subsequently stained with Lysotracker. The study revealed that relative to the wild type and complemented strains, $\Delta metZ$ was more likely to end up in the acidic compartment of the cell (Fig. 2E). Overall, the data suggests that absence of *metZ* weakens the ability of *M. tuberculosis* to resist host-induced physiological stress and phagosome-lysosome fusion both critical for survival of the pathogen inside the host cell.

Direct sulfhydrylation pathway is essential for *M. tuberculosis* to survive under low pH

The above data suggest an enhanced sensitivity of $\Delta metZ$ on exposure to acidic pH. To confirm this, we cultured different bacterial strains (WT, $\Delta metZ$ and complement) at various pH and observed the changes in growth kinetics. While the growth kinetics of all three strains were similar at neutral pH

(7.2), a decrease in pH led to an observable defect (pH 5.5) to complete growth arrest (pH 5.0) of the

ΔmetZ strain relative to the wild type and complemented strain (Fig. 3A and S4 A-C). Further, to

determine the effect of acidic pH on expression of genes involved in transsulfuration (*metB* and *metC*)

and direct sulfhydrylation (*metZ*), we cultured *M. tuberculosis* at pH 7.2 and pH 5.5. Relative

expression determined by RT-qPCR revealed that expression of *metZ* increased by ~ 12-fold upon

induction by acidic pH 5.5 when compared to its expression at pH 7.2 (Fig. 3B). To verify if the low

CFU counts at pH 5.5 (Fig. S4C) is due to bacterial killing or a consequence of metabolic slowdown,

we determined metabolic activity of the mycobacterial strains at pH 5.5 using Resazurin dye [22]. We

find that relative to the wild type and the complemented strain, *ΔmetZ* failed to change the color of the

dye from blue to red which corresponded to a low fluorescence readout indicating that the absence of

metZ significantly alleviated the metabolic activity of *ΔmetZ* grown in pH 5.5 (Figure 3C).

Furthermore, estimated ATP levels in all three strains when cultured at acidic pH 5.5 revealed that

ΔmetZ surprisingly had higher ATP levels relative to WT and complemented strains (Fig. 3D). It is

known that PYZ is active only at acidic pH and efficiently kills slow or non-replicating cells. We

hypothesized that the differences in the cytoplasmic pH will directly correlate with PYZ sensitivity.

As expected, we observed a ~19-fold and 4-fold increase in the sensitivity of *ΔmetZ* to PYZ treatment

in comparison to WT and complemented strain respectively (Fig. 3E). To confirm, if the observed

phenotype is a direct consequence of defect in the biosynthesis of methionine, we measured the

intracellular levels of methionine in H37Rv and *ΔmetZ* grown in media at both the pH. We found that

methionine levels were significantly lowered in WT strain when grown in acidic pH 5.5 compared to

bacteria grown at pH 7.2. Similarly, mutants had lower methionine levels at acidic pH in comparison

to mutant cells grown at pH 7.2 (Fig. 3F). Interestingly, exogenous supplementation of methionine

partially rescued the growth defect phenotype of *ΔmetZ* observed at pH5.5 (Fig. 3G). Interestingly, it

was also seen that while growing at acidic pH, *M. tuberculosis* can produce H_2S at levels significantly

higher than that growing at neutral pH (Fig. S5B). H_2S is a co-substrate and S-donor in the reaction

catalyzed by OSHS enzyme (MetZ). In sum, a defect in the methionine biosynthesis pathway at low

pH renders *ΔmetZ* metabolically less active thereby preventing the growth of the mutant at low pH.

Molecular docking and MD simulations revealed enhanced stability of MetZ at acidic pH

To enable us to perform *in-silico* molecular docking with MetZ protein (Fig. 4A), we determined the

binding site of O-succinyl homoserine (OSH) by focused molecular docking [23, 24] that provided the

final pose at the cofactor bound site. A pose with docking energy of -4.07 Kcal/mol and cluster size of

50 (out of 200) was found suitable at pH 7.0. While, at pH 5.5 the most likely pose showcased a docking

energy of -4.19 Kcal/mol with a cluster size of 80 (200). For both these poses at diverse pH, a static

thermodynamic quantification was carried out using MM-GBSA which was found to be -49.20

Kcal/mol and -56.93 Kcal/mol, respectively. Molecular Dynamics Simulation at a constant pH was

performed for these poses. Following the 300ns MD simulations, the protein exhibited stable RMSD

values at 2.98 Å and 4.34 Å at pH 7.0 and 5.5, respectively. However, significant changes were

observed in the stability of the substrate as well as the occupied binding site. On observing the substrate

in consensus with its binding site, the region was found to have higher stability and an average RMSD

of 1.87 Å was observed at pH 5.5 in comparison to an average of 2.71 Å at pH 7.0 (as shown in Fig.

4B and D). The underlying reason for this higher stability at pH 5.5 is due to the establishment of stable

durable bonds in the form of hydrogen bonds with residues V196, F197 and T319 and the O-atoms at

the terminals of the substrate. Furthermore, another factor which enhances the strength is the formation

of a salt-bridge with residue H220 and substrate. However, a similar configuration of H-bonds and salt-

bridge are not observed at pH 7.0, possibly due to conformational dynamics [25] (Fig. 4E). These

characteristics are a result of the change in pKa values as well as conformations of the binding site

forming amino acid residues. Protonation states of amino acids are highly susceptible to change in

environmental pH and thus, preparing the protein at pH 7.0 and 5.5 generates different acid dissociation

constants, which ultimately affects the bond forming tendencies of residues. The residue E165 attains

a pKa of 8.78 at pH 7.0 and donates a proton, forming a bond with the substrate. Similarly, D194 attains

a pKa of 9.11 at pH 5.5 which affects its binding to the substrate. Also, H220 is present in a protonated

state at a lower pH and thus involved in the formation of a Salt-Bridge with the substrate. While at pH

7.0, H220 remains in a deprotonated state and thus does not contribute to the stability of the binding

pocket at neutral pH. This showcases the importance of the acidic pH which contributes to relatively

greater stability of the substrate. On analysis of the binding site, we found that H220 and T319 interact

with the substrate at the acidic pH but they showcase conformational changes at pH 7.0 which hinders

their interaction. The imidazole ring sidechain of H220 rotates in anti-clockwise direction and thus

does not get involved in bond formation. Similarly, T319 forms H-bonds at both pH but the nature of

the bonds varies from transient at pH 7.0 to largely consistent at pH 5.5. Overall, conformational

changes in the binding site residues can be accredited for better stability at pH 5.5 The stability of the

protein and the substrate at acidic pH hints towards a better binding activity of the substrate. This

characteristic feature can be examined by performing the MM-GBSA of a few selected frames from

both the trajectories to compare their activity. The average binding/biological activity of the protein is

found to be -37.35 Kcal/mol and -59.40 Kcal/mol at pH 7.0 and 5.5, respectively (Fig. 4C). This

increase in the binding affinity is due to the effect of acidic pH on the protonation state as well as the

induced conformational changes. The docking scores at both pH 7.0 & 5.5 suggests that the residence

time of the substrate at the binding site must be short and the sulfur transfer reaction might be transient

in nature. Overall, focused docking and MD simulations revealed that although protein is stable at both

acidic and neutral pH, the active site of the protein holds substrate (OSH) more stably at acidic pH

rather at neutral pH.

***metZ* mediated direct sulfhydrylation pathway is critical for methionine biosynthesis at low pH**

Transsulfuration pathway is well-known and characterized for methionine biosynthesis in several

organisms including *M. tuberculosis*. However, to confirm whether *metZ*-dependent methionine

biosynthesis in *M. tuberculosis* is independent of transsulfuration pathway, we knocked down

expression of *metB* using CRISPRi in the background of Rv-WT (*metB$_{KD}$*) and *ΔmetZ (ΔZ::metB$_{KD}$)*

strains (Fig. 5A). Anhydrotetracycline (ATc) induced knockdown of *metB* gene in both the strains were

confirmed by quantifying the *metB* transcript levels by RT-qPCR (Fig. S5A). Interestingly, upon

induction with ATc, we find no difference in the growth of the H37Rv and *metB$_{KD}$*, but *ΔZ::metB$_{KD}$*,

which upon induction by ATc behaved as a double knockout strain, was found to be defective (3.5-

fold at day 9) in growth even under normal pH condition (Fig. 5B). The observed growth defect could

be due to an obstructed methionine biosynthesis pathway which was confirmed by relative methionine

levels wherein we found a ~1.5-fold decrease in the intracellular methionine in *ΔZ::metB$_{KD}$* compared

to the *metB$_{KD}$* (Fig. 5C). Further, we opined that the double mutant (*ΔZ::metB$_{KD}$*) should be attenuated

in animal model owing to its reduced methionine levels. To test this, we infected mice with

mycobacterial strains (NT, *metB$_{KD}$* and *ΔZ::metB$_{KD}$*) and fed with doxycycline to induce CRISPRi

throughout the study and bacterial burden was determined by CFU plating (Fig 5D). Compared to NT

and *ΔmetZ,* double mutant *ΔZ::metB$_{KD}$* showed reduced bacterial burden at initial stages of infection in

both lungs and spleen of mice model, which however, was rescued later by week 6 (Fig. 5E). We

believe the presence of active methionine transporter like MetM (Rv3253) [26] could possibly be

involved in rescuing the growth defect phenotype of *ΔZ::metB$_{KD}$* in animal model study. Overall, the

data suggest that direct sulfhydrylation pathway of *M. tuberculosis* which is driven by MetZ enzyme

is independent of transsulfuration pathway (driven by MetB and MetC) as shutting down of both the

arms lead to attenuation in growth both *in-vitro* as well as *in-vivo* and reduced methionine levels.

Direct sulfhydrylation pathway of *M. tuberculosis* contributes towards antibiotic persistence in

response to Bedaquiline

Next, we wanted to study the role of direct sulfhydrylation pathway in regulating the drug susceptibility

profile of *M. tuberculosis*. We first compared the Minimum Inhibitory Concentration (MIC) of various

anti-TB drugs (RIF, INH, EMB and BDQ) and found that the IC50 values for each of these drugs were

similar for all the three strains (H37Rv, *ΔmetZ* and the complemented strain) (Fig. 6A). Further, a time-

kill assay was performed to determine the role of *metZ* in modulating the formation of drug-induced

persisters in *M. tuberculosis*. Difference in the persister formation within the strains was estimated by

comparing the biphasic kill curve generated by plotting the data obtained through time-kill assay.

Briefly, each of these cultures were treated with different antibiotics (10X MIC conc.) and the bacterial

survival was estimated by CFU plating of the cultures at different time points. We did not find any

difference in the frequency of generation of persisters between different strains treated with either RIF,

INH and EMB. Surprisingly, in comparison to the WT and the complemented strain, a significant

decrease in the persister formation was observed in *ΔmetZ* treated with BDQ (Fig 6B). Since BDQ acts

by restricting the ability of the pathogen to synthesize ATP, we hypothesized that the observed decrease

in the persister population could be a result of defect in generation of ATP in *ΔmetZ*. On the contrary,

our ATP depletion kinetic analysis using 2X MIC showed that BDQ treated *ΔmetZ* had higher ATP

levels relative to the WT and the complemented strains (Fig. 6C) suggesting that BDQ treatment led

to slower ATP depletion in *ΔmetZ*.

Direct sulfhydrylation pathway supports the growth of *M. tuberculosis* inside the host

To elucidate the role of *metZ* in mycobacterial pathogenesis, a guinea pig model of tuberculosis

infection was used. Briefly, guinea pigs were infected with the respective mycobacterial strains (Rv-

WT, *ΔmetZ* and Complement) via aerosol route. Bacterial burden in the lungs and the spleen was

determined by CFU plating at different timepoints post-infection. Gross and histopathological analysis

were performed to study the damage due to infection at the tissue level (Fig. 7A). We found that, at

week 4 post infection the bacterial burden in the lungs of the infected guinea pigs was ~3.7-fold and 2-

fold lower in *ΔmetZ* when compared to WT and Complement strains respectively. Approximately a

difference of ~8.9-fold was observed between bacterial burden in lungs infected with WT and *ΔmetZ*

at week 8 post infection. It may also be noted that no significant increment in the CFU of mutant week

4 through week 8 (Fig. 7B). Similarly, there were ~6-fold lesser bacilli in the spleen of *ΔmetZ* infected

animals opposed to WT and complement strains' infected spleen at week 4 post infection. Interestingly,

there was a marked decline in the recovery of colonies of *ΔmetZ* from guinea pig spleens at week 8

post infection. This was almost near to the limit of detection (LOD) by CFU and almost 263-fold lesser

bacilli than what was recovered from week 4 time point spleens (Fig. 7C). In coherence with CFU

burden, (Fig. 7B), we found lesser granulomatous lesions in *ΔmetZ* infected lungs as compared to WT

and complement strains infected lungs from week 8 lungs (Fig 7D). To visualize and quantify the

damage to lung tissues, H&E staining of the formalin preserved lung sections were done. Microscopic

examination of lung samples collected from animals infected with wild type strain revealed severe

granulomatous inflammation throughout the lung parenchyma. Numerous well-defined necrotizing

granulomas consisting of epithelioid histiocytes, perimetric cuffing of lymphocytes and plasma cells

aggregates and a caesious necrotic mass at the center were observed. However, lung samples from

animals infected with complement strain and *ΔmetZ* strain showed only non-necrotizing granulomas

along with multiple foci of inflammatory cells aggregates and occasional alveolar hemorrhagic patches

(Fig. 7E). Comparative granuloma scoring revealed the highest score for WT infected samples

followed by *ΔmetZ* and Complement strain infected samples (Figure 7F). Overall, the data suggests

that absence of *metZ* gene in *M. tuberculosis* impedes the ability of the pathogen to replicate and

survive inside the host indicating *metZ* (or direct sulfhydrylation) for methionine biosynthesis is

required for survival inside host at later stages of infection.

metZ is involved in methionine biosynthesis in _M. tuberculosis_ at acidic pH

We hereby summarize that when *M. tuberculosis* infects macrophage, it passes through a series of

compartments starting from phagosome where pH is essentially ~6.5 and the bacteria is in replicating

state, with enough nutrients (including methionine). Further, as host immunity kicks in and IFN-γ

induces accumulation of ROS, RNS and subsequently, phagosome maturation leading to drop in pH to

5.5 and beyond, conditions are non-replicating for *M. tuberculosis* and host begins to quench nutrients

from bacteria as means to impose restriction on the trapped bacteria. At this stage bacteria undergoes

several transcriptional and metabolic changes to sustain the host induced insults, one of these includes

induction of *metZ* expression that help bacteria survive these stress conditions and helps maintain

methionine levels by utilizing host produced H_2S which acts as co-substrate for methionine

biosynthesis in acidic micro-environment and helps bacteria maintain recycling of ATP, mycothiol and

its metabolic activity going (Figure 8).

Discussion

M. tuberculosis, adapts to various conditions within macrophages, resulting in the differential

expression of proteins that support homeostasis. Under such circumstances, the pathogen needs to

evolve and device a mechanism to efficiently regulate the flux of cysteine towards either biosynthesis

of methionine and/or mycothiol. Cysteine, in addition to other metabolites, is a key amino acid essential

for biosynthesis of both methionine and mycothiol in mycobacteria. *M. tuberculosis* enhances the

biosynthesis of cysteine to most possibly to compensate for the increase in the flux of cysteine towards

generating the thiol-active metabolites (mycothiol, ergothionine, NAD+, etc.) essential for maintaining

the redox balance in conditions with heightened levels of ROS, RNS [27] and upon exposure to acidic

pH [28]. Similar findings in *E. coli* have been reported that growing demand of cytosolic redox

glutathione under higher oxidative stress boosts the synthesis of cysteine [29]. We, in the current study,

have used a combination of genetics and computational biology-based tools to probe the role of *metZ*

mediated direct sulfhydrylation pathway in *M. tuberculosis*. We found that MetZ dependent direct

sulfhydrylation is induced by several folds at acidic pH. Disrupting this pathway in *M. tuberculosis*

results in a decrease in methionine at low pH. Finally, we demonstrated that MetZ dependent direct

sulfhydrylation is required for growth of *M. tuberculosis* inside the host at later stages of infection.

In contrast to an earlier report [19], *metZ* was found to be dispensable for growth of *M. tuberculosis* on

cholesterol. The conditional essentiality phenotype observed for *ΔmetZ* under low pH further suggests

that the observed essentiality of *metZ* for growth on cholesterol was a result of relatively low pH growth

conditions used in that study. Cysteine being the common precursor for the generation of both the

methionine and the mycothiol, the observed imbalance in the redox homeostasis in *ΔmetZ* could be

attributed to the inefficient recycling of mycothiol due to a major shift of the cysteine pool towards

methionine biosynthesis. *M. tuberculosis metZ* (*Rv0391*) gene encodes a probable OSHS catalyzing

direct conversion of OSH to homocysteine using soluble H_2S instead of cysteine as S-donor. Given the

crucial role of methionine in bacterial physiology, there has been a gap in our understanding of the

rationale of harboring multiple methionine biosynthesis pathways and their role in maintaining

methionine homeostasis critical for the survival of *M. tuberculosis* in a nutrient deprived *in-vivo* growth

environment. Interestingly, other than maintaining the redox homeostasis, our data underscores the role

of methionine biosynthesis, through direct sulfhydrylation pathway, in regulating the cytosolic pH and

metabolism. The inability of the *ΔmetZ* to regulate the intracellular pH homeostasis impedes the ability

of the pathogen to utilize the cellular ATP rendering the pathogen metabolically less active. Similar

findings have been reported in *E. coli* previously where antibiotic induced oxidative stress led to

increased levels of intracellular ATP which is a consequence of non-utilization rather than over-

production [30]. This explains increased levels of ATP in *ΔmetZ* at acidic pH which is a consequence

of metabolic slowdown at acidic pH. The enhanced susceptibility of *ΔmetZ* to pyrazinamide (PYZ)

further confirms the presence of a relatively acidic cytosol and accentuates the role of *metZ* in

maintaining the pH homeostasis in *M. tuberculosis*. In an *ex-vivo* setting, IFN-γ induction leads to the

acidification of cellular compartments where the internalized bacteria is trapped and maintained at low

pH. As a consequence of this, *ΔmetZ* replication inside resting as well as activated macrophages was

reduced which was because of the comparatively higher co-localization of *ΔmetZ* mutant to acidic

compartments of the macrophages as revealed by confocal microscopy. Similar to our findings, other

studies have also observed that enhanced susceptibility to either acidic pH or oxidative stress results in

increased killing due to higher co-localization with acidic compartments of the macrophages [31-33].

Our data provides mechanistic understanding of the methionine starvation associated susceptibility in

M. tuberculosis previously reported by several groups [5, 11]. The presence of both transsulfuration

and direct sulfhydrylation pathway for the biosynthesis of methionine in *M tuberculosis* is indeed

intriguing. Although both the pathways demonstrate redundancy at normal pH, our qPCR data along

with previous studies categorically indicate that enzyme responsible for carrying out direct

sulfhydrylation is over-expressed in *M tuberculosis* upon induction with acidic pH [32, 34, 35]. These

findings are corroborating well with our pH dependent molecular dynamics simulation studies which

suggested that the binding affinity between substrate and MetZ active site is more strong at low pH.

The thorough pH-dependent MD simulations provide structural insights in terms of distinct behavior

as how pH and protonation states of key residues alter the interaction networks particularly hydrogen

bonding and electrostatic interactions within the binding site residues of the protein, leading to

significant changes in the conformation. The change in hydrogen bonding network identified as a

determinant which majorly impacts the change in binding affinity of the substrate. A comparative

analysis between two pH was evaluated in terms of MetZ structural stability using metrics such as root-

mean-square deviation (RMSD) and thermodynamic changes that further confirm the pH-dependent

behavior, indicating that *metZ*-mediated direct sulfhydrylation becomes the major methionine

producing pathway of *M. tuberculosis* under low pH.

We further confirm that *M. tuberculosis* strain that lacks both transsulfuration and direct

sulfhydrylation pathways ($\Delta metZ/metB_{KD}$) fail to generate methionine and demonstrate methionine

auxotrophy under *in-vitro* growth conditions implying that both the pathways are independent of each

other. Surprisingly, in comparison to *in-vitro* growth condition, the growth defect observed in

$\Delta metZ/metB_{KD}$ under *in-vivo* growth condition, although significant, was less pronounced. This could

be attributed to the fact that the pathogen is known to harbor potential methionine transporter encoded

by *Rv3253c* (*metM*) [26] which helps the pathogen salvage methionine from the host. It has also been

shown previously that *M. tuberculosis* has an ability to produce H_2S – a gaseous signaling molecule

upon induction with acidic pH (Fig. S5B) and [14] probably as a mechanism to counteract increasing

ROS and dropping metabolic potential of the cells. Further, it has also been shown that H_2S at sub-

inhibitory levels in *M. tuberculosis* have a positive effect on mycobacterial electron transport chain

(ETC) and this effect is reversible [36, 37]. Additionally, host produced H_2S in response to *M.*

tuberculosis infection and have a positive effect on bacterial replication such that *M. tuberculosis*

replication was hampered when host's ability to generate H_2S was abrogated either by genetically

knocking out enzymes responsible for H_2S generation – CSE and CBS (among others) or by using

pharmacological inhibitors of the same [36, 38, 39]. Co-incidentally, through literature it has also been

extensively established that sulfhydrylases in general, use H_2S as S-donor to either synthesize cysteine

or, in this case, MetZ produces homocysteine, hence, methionine.

Development of antibiotic persistence in bacteria is a major culprit responsible for the failure of anti-

TB regimen. Frequency of generation of persister population can be quantified by performing time-kill

assay by treating different culture at high dosage of antibiotic (5-10X MIC). A biphasic kill kinetics is

indicative of drug persistence [40]. Our time-kill experiments suggest that the absence of direct

sulfhydrylation pathway in *ΔmetZ* prevents the BDQ specific generation of antibiotic persistence in *M.*

tuberculosis. Intriguingly, in spite of higher cytosolic ATP levels relative to the wild type, *ΔmetZ*

demonstrated a decrease in the frequency of BDQ specific persisters formation. This is similar to previous study where authors have reported *ΔpckA* strain maintained higher ATP levels even upon treatment with BDQ grown in presence of acetate which promoted persister formation in *ΔpckA*. However, as opposed to our findings, they demonstrated that *ΔpckA* had increased drug tolerance and more ATP levels in acetate medium [41]. In sum, apart from maintaining methionine levels at acidic pH, direct sulfhydrylation pathway also modulate BDQ specific generation of antibiotic persistence during infection. Future studies on understanding this complex phenotype will help us design strategy to prevent the development of BDQ resistance in tuberculosis. It was also found that *metZ* is required for maintenance of *M. tuberculosis* infection in guinea pig model of tuberculosis infection, as indicated by a lowered CFU burden at week 8 post-infection and severe attenuation in the spleen as well. Spleen is known to have a high iron content, which itself is responsible for the generation of ROS via Fenton chemistry. Given the susceptibility of the *metZ* mutant to oxidative stress, we find a very low bacterial count in the spleen at week 8 post-infection. This suggests that *metZ* supports methionine biosynthesis and hence bacterial survival in a nutrient deprived and low pH niche inside the host.

The presented data therefore, suggest that *M. tuberculosis* is short of methionine at acidic pH when trapped inside phagolysosome. At this stage, expression of *metZ* is induced several folds as opposed to *metB* and *metC*. Absence of direct sulfhydrylation pathway curtails the growth of Mtb at acidic pH by significantly reducing the metabolic activity together with decrease in the ability to utilize the cellular ATP. Further, such conditions arise around 3–4-week post infection in animal model which is marked by onset of IFN-γ mediated bacterial control where antimicrobial mechanisms are at peak posing challenge to bacteria in the form of ROS, RNS, acidic pH and increasing H_2S biogenesis. Concordantly, *metZ* null mutant was susceptible to all these stresses *in-vitro*, verifying our findings. Our results, hence, indicate that *M. tuberculosis* adapts to the decreasing pH in its surroundings to maintain methionine biosynthesis through direct sulfhydrylation using exogenous and perhaps endogenous H2S

as S-donor. This makes MetZ an interesting drug target that can be aimed in combination with anti-TB

drugs to improve their efficacy. A pharmacological inhibitor of MetZ should be able to inhibit

methionine biosynthesis in the bacteria trapped inside a non-replicative condition (i.e. phagolysosome)

characterized by exceedingly low pH ~4.5-5.5. Another strategy could be using small molecules that

could drop intrabacterial pH of *M. tuberculosis* [42] in combination with pharmacological inhibitors

of MetZ. This would ensure bacterial growth arrest due to reduced Intrabacterial pH and block

methionine biosynthesis at the same time by inhibiting MetZ.

408 Materials and Methods

409 Bacterial strains

Knockout strain of *metZ* gene *(rv0391)* was generated in the genetic background of wild type

M. tuberculosis H37Rv by genetic recombineering. Briefly, a gene-specific allelic exchange

substrate (AES) harboring 1kb sequences of target gene upstream and downstream homology

flanking Hygromycin resistance gene cassette was constructed in the suicidal vector pJM1. AES

was transformed into H37Rv (wild-type) strain overexpressing Phage Che9c RecET proteins to

facilitate recombination. Recombinants with the Hyg^r gene were selected over Hyg plates and

screened using gene-specific PCR. The marked mutant thus prepared was unmarked to excise out

Hyg resistance cassette using Cre-recombinase expressed from plasmid pCre-SacB-Zeo. Unmarked

mutant was confirmed by PCR and RT-qPCR to confirm loss of gene as well as transcripts of metZ

gene. Unmarked mutant was then transformed with plasmid expressing a functional copy of *metZ*

gene under P$_{myc1}$*tetO* promoter replicating either episomally or integrated to generate genetic

complementation strain (denoted as *ΔmetZ Comp*). Bioluminescent mycobacterial strains were

generated by transforming respective mycobacterial strains (**List of Strains, Appendix-I**) with

integrative plasmid expressing *luxABCDE* operon (Addgene # 26161) [43] with slight

modifications [44]. Strains were checked for bioluminescence and a luminescence vs CFU plot was

prepared to determine linear relation between luminescence and CFU based read-outs of bacterial

viability for lux-based mycobacterial strains (Fig S3A).

Media and cultural conditions

The bacterial strains were maintained as frozen glycerol stocks at -80°C. Whenever required, the

stocks were thawed and diluted 1:500 in 7H9 (Difco, BBL) medium supplemented with OADS and

0.01% Tween 80. Bacterial enumeration was done on 7H11 (Difco BBL) agar plates supplemented

with OADS. Kanamycin (Sigma), 25µg/ml, Hygromycin (Sigma), 50µg/ml and Zeocin

(Invitrogen), 25µg/ml were used whenever required. For media with no carbon source, 7H9 base

medium was prepared without glycerol and OADS (or ADS) enrichment and later supplemented

with desired C-sources and 0.05% tyloxapol. 100mM stock of Cholesterol (Sigma) was prepared

by heating cholesterol in Tyloxapol (Sigma): Ethanol (Merck) mix (1:4) at 80°C for 30 mins [45].

When required, cholesterol was diluted to desired concentration (200µM) in 7H9 base medium

wherever specified. Likewise, glycerol (Sigma) was used at final concentration 0.1% v/v and

sodium propionate (Sigma) was used at concentration of 10µM (0.1%) for growth kinetics studies.

For pH dependent growth kinetics experiments, 7H9 base medium was prepared, supplemented

with 0.2% glycerol (v/v) and ADS acidified to desired pH using HCl. Phosphate buffered saline

(PBS) was used at 1X concentration supplemented with tyloxapol (0.05%) whenever mentioned.

Bacterial enumeration by CFU was done on 7H11 agar plates supplemented with OADS, wherever

mentioned. Colonies were counted after 21 days of incubation. Anhydrotetracycline, ATc (Caymen)

was used at 200 ng/mL concentration, unless mentioned otherwise.

CRISPRi knockdown

For transcriptional knockdown of *metB (Rv1079)* using CRISPR interference (CRISPRi), target specific sgRNA was cloned into *BsmBI* (NEB) digested backbone plJR965 (Addgene # 115163) to generate knockdown construct. Knockdown construct was transformed into *Rv*-WT and *ΔmetZ* strains. To achieve knockdown of the target genes, cells were grown to mid-log phase and diluted back to OD 0.01 in fresh culture medium with or without ATc. Growth was monitored by measuring OD of the cultures. Further, expression of target genes was quantified by qPCR to determine the levels of knockdown.

RT-qPCR

Total RNA was isolated from the respective bacterial strains under specific conditions using Trizol (TaKaRa) according to the manufacturer's protocol. Briefly, bacterial cells were washed with 1X PBS and resuspended in Trizol. Cells were lysed by bead-beating using 0.22μM sized zirconia beads via 5 cycles with intermittent cooling step of 1 minute. Further, RNA was extracted and precipitated using chloroform-isopropanol treatments. Total RNA thus prepared was treated with Turbo DNaseI (Ambion) to digest away the DNA contamination. 3000 μg equivalent DNase treated RNA was used to prepare cDNA using SuperScript IV Reverse Transcriptase (Invitrogen). Finally, 1μg equivalent cDNA was used to set up qPCR reactions with SYBR (TaKaRa). *sigA* was used as internal control and $\Delta\Delta C_t$ method [46] was used determine relative gene expressions w.r.t. specified control conditions. For qPCR at acidic pH, *icl1* was used as positive control.

Growth kinetics

Auto-bioluminescent bacterial strains were grown till OD 0.5 (mid-log phase) and washed once with 1X PBST to remove the spent media. The cells were resuspended in 1X PBST and diluted back

to an OD of 0.01 in 7H9 base medium supplemented with different carbon sources – glycerol

(0.1%), cholesterol (200µM) or propionate (10mM) for C-source dependent growth kinetics. For

pH dependent growth kinetics, bacterial cells were diluted back to an initial OD of 0.05 in 7H9

medium (supplemented with ADS and 0.2% glycerol) acidified to different pH – 7.2, 6.5, 5.5 and

5.0. Whenever required, L-methionine (50µg/ml, Sigma) was added to the cultures. The growth was

monitored by measuring OD and/ or auto-bioluminescence using a Multimode Microplate reader

(SpectraMax Pro ID3, Molecular Devices). At times, bacterial growth was also determined by CFU

at indicated time points by plating aliquots of cultures and plating appropriate dilutions on 7H11

agar plates.

Physiological stress survival assay

Bacterial strains were grown till OD 0.5 (mid-log phase) and washed once with 1x PBST. Aliquots

of 2ml cell suspensions equivalent to OD 0.2 were treated with 0.5mM H_2O_2 (oxidative stress),

200µM Deta-NO (nitrosative stress), 0.01% SDS (membrane stress), PBST (nutrient starvation

stress), Phosphate citrate buffer, pH 5.5 (acidic stress) *in-vitro*. The survival of the respective strains

was determined by CFU plating at indicated time points and calculating survival index (i.e., CFU

of strain/ CFU of Rv-WT).

Drug susceptibility (MIC determination)

Auto-bioluminescent bacterial strains were grown till OD 0.5 (mid-log phase) and washed with 1x

PBST. Anti-TB drugs (RIF, INH, BDQ, EMB, CFZ and PYZ) were serially diluted on a 10-points,

2-fold format and added to the 96-well plates (SPL). Cells were added at final density of OD 0.01

to the wells containing drugs and incubated at 37°C for 3 days as described previously [47]. Finally,

luminescence was recorded as an indicator of bacterial viability which was expressed as dose-

response curves and IC_{50} were calculated using non-linear regression analysis in GraphPad Prism.

Kill kinetics

Bacterial strains were grown till OD 0.5 (mid-log phase) and washed with 1x PBST. Aliquots of

cell suspensions (2ml) at bacterial density of OD 0.2 were treated with 10X MIC concentrations of

anti-TB drugs (Rif, Inh, Emb and Bdq) for mentioned time. Bacterial survival was determined by

CFU enumeration at indicated time points.

Methionine estimation

Log-phase cultures of mycobacterial strains were washed with 1X PBST and transferred to 7H9

medium maintained at pH 7.2 or 5.5 at an OD 0.4. Cultures were allowed to grow for 6 days after

which they were washed with 1X PBS once followed by washing with chilled methanol. Cells

equivalent to OD 1 ($\sim 3 \times 10^8$ cells/mL) were lysed in 1 ml extraction buffer (Acetonitrile:Methanol:

water:: 2:2:1) by bead beating with intermittent cooling of 1 min per cycle. Lysates were clarified

by spinning at 13,000 rpm, 10 mins at 4°C. The clarified supernatants were filtered through 0.22-μ

filters and dried by speed vac to evaporate the extraction buffer. Dried pellets were resuspended in

methanol:water (1:1) + 0.1% formic acid and vigorously mixed. Finally, 10μL of the reconstituted

samples were run on Hypersil Gold C8 column (100 x 2.1mm) with 0.1% formic acid in water as

mobile phase A and 0.1% formic acid in acetonitrile as mobile phase B on SCIEX 6500+ QTRAP

and peak and retention time was matched with standard (Methionine) across all the samples. Peak

intensities were normalized to CFU of the respective samples.

Total ATP estimation

Cultures of Rv-WT, *ΔmetZ* and Complement were grown at acidic pH 5.5 till OD 0.5 (mid-log

phase) and washed with 1X PBST. Cells were resuspended in 1000μL 1X PBST (with Protease

Inhibitor Cocktail) and lysed by 5 cycles: 45 seconds of bead beating with intermittent cooling on

ice. Lysates were passed through 0.5-μ sized filters to remove any bacterial cells or debris. 50μL of

clarified lysates were transferred to a 96-well plate (white, flat-bottom, SPL) and mixed (1:1) with

BacTiter-Glo (Promega) reaction buffer followed by incubation at 37°C for 30 minutes. 1X PBS

was used as blank to cancel out the background signals from the samples. Luminescence was

recorded using a Multimode Microplate reader (SpectraMax Pro ID3, Molecular Devices).

Normalization was done using total protein content of the respective samples and ATP levels were

expressed as relative units w.r.t. Rv-WT strain. For ATP estimation of BDQ treated cultures, 0.4

OD bacterial suspensions (2ml) of the respective strains were treated with 2X MIC concentration

of BDQ for the indicated time-points and samples were prepared as described above.

Metabolic activity

Auto-bioluminescent mycobacterial strains were grown till OD 0.5 (mid-log phase) and washed

with 1X PBST to remove the spent media. Cells were resuspended in 1X PBST and re-diluted to an

OD of 0.4 in media acidified to required pH (pH 5.5). 100μL of these suspensions were added to a

96-well plate (white, flat-bottom, SPL) containing 7H9 (ADS + 0.1% glycerol, pH 5.5) and serially

diluted (2-fold). The periphery wells were filled with sterile medium to prevent desiccation. The

plate was incubated at 37°C for 5 days after which 20μL of PrestoBlue (Thermo) was added to each

well followed by an additional incubation of 2 days to allow development of color. After 2 days of

incubation, luminescence read-out was taken and contents of the wells were transferred to a clear

bottom black plate followed by recording fluorescence of AlamarBlue at 560(ex)/ 590(em). Ratio

of Fluorescence/Luminescence was calculated to determine metabolic activity per cell.

Intracellular redox state

Rv-WT, *ΔmetZ* and Complement strains were transformed with plasmid encoding redox biosensor, Mrx1-roGFP2 as described previously [48]. Cells were grown to OD 0.5 and washed with 1X PSBT and resuspended in PBST in MCTs. Rv-WT cells in respective tubes were treated with either 100μM CHP or 100μM DTT as control groups for oxidized and reduced conditions respectively. Further, cells were transferred to 96-well plates (clear bottom, black wells) and excitation at 430 and 490 nm while emission at 510 nm was recorded. Ratio of fluorescence at 390 nm/490 nm were compared to determine intracellular redox state of the cells.

Lipid hydroperoxidation assay

Total lipids were extracted from the respective bacterial strains as described previously [49]. Briefly, bacterial cells were grown to an OD 0.8-1.0 and pelleted down. Pellets were suspended in methanol:chloroform (2:1), 6 ml overnight at 37°C to extract total cellular lipids. Next day, cells were spun down and supernatants were collected. Supernatants having total cellular lipids were further precipitated by saturating the supernatants with 0.8% KCl. The lower phase was collected and dried in glass vials to evaporate the solvents and concentrate the lipids. Dried lipids were resuspended in 1 ml chloroform (Sigma) and used for FOX2 assay for determining lipid hydroperoxidation in the extracted lipids as described elsewhere [50]. Briefly, 200μL of the concentrated lipids were mixed with 1000μL FOX2 reagent in MCTs in dark for 1hr. Samples were transferred to a 96-well plate and read out at 560 nm.

Intracellular sulfide estimation

Total sulfide estimation was performed as described previously [51] [37]. Briefly, Rv-WT, $\Delta metZ$

and Complement strains were grown to OD 0.5 (mid-log phase) in regular 7H9 medium

supplemented with OADS and 0.01% Tween 80. Cells equivalent to OD 0.5 were inoculated into

7H9 medium supplemented with ADS and 0.01% tyloxapol at pH 7.2 and pH 5.5. After 4 days of

incubation, the cultures were pellet down and washed with 1X PBST to remove spent media. Pellets

were resuspended in the same media with respective pH 7.2 and 5.5 at an OD 0.4. 100µL of the cell

suspensions were added to 96-well plate and mixed with 100µL 2X Bismuth chloride buffer.

Following an incubation of 4 hours at 37°C, colorimetry was done at 405nm. The values were

normalized to culture OD to determine relative sulfide levels in the respective strains at two different

pH.

Macrophage infection

THP-1 monocytic cells were grown in RPMI medium (Lonza) supplemented with 10% fetal bovine

serum (FBS, Invitrogen). Further, cells were seeded at a density of $5x10^5$ cells/ well in a 12-well

plate (Corning) and matured for 24 hours with PMA (25ng/mL, Sigma) followed by a rest of 24

hours. Whenever required, cells were activated with IFN-γ (10ng/ml, Peprotech) for 16 hrs before

infection. Single cell suspensions of the bacterial strains (Rv-WT, $\Delta metZ$ and complement) were

prepared by soft-spin method as described [44] and used for infecting resting or activated

macrophages at an MOI of 1. Bacteria were allowed to infect the cells for 4 hours after which,

extracellular bacteria were washed off by three washes with pre-warmed PBS (Gibco). At indicated

time points, macrophages were lysed with 1X PBS + 0.01% Triton X-100 to release intracellular

bacterial cells. Lysates were serially diluted and plated for bacterial enumeration.

Confocal Microscopy

THP-1 macrophages were seeded on glass coverslips in a 24-well plate (5×10^5 cells/well) and were

activated by 10ng/ml of human IFN-γ (Peprotech, USA) for 16 hours before infection. Activated

THP-1 cells were infected with mycobacterial strains (Rv-WT, *ΔmetZ* and Complement) harboring

Live-Dead reporter plasmid [52] constitutively expressing mEmerld at an MOI of 10. After 4 hours

of infection, the cells were washed thrice with 1X PBS to remove the extracellular bacteria and

replenished with fresh media. Subsequently, after 24 hours post infection, cells were incubated with

50nM LysoTracker Red DND-99 (Invitogen life Technologies, CA, USA) in complete RPMI media

for 30 minutes at 37°C with 5% CO2 and later fixed with 4% paraformaldehyde in 1X PBS.

Coverslips were mounted with ProLong Diamond antifade with DAPI (Molecular Probes by Life

Technologies, CA, USA). Finally, slides were imaged by Olympus Confocal Laser Scanning

Microscope (CLSM). Later, GFP-expressing mycobacteria that colocalized with LysoTracker Red

was determined by analyzing the ~ 100 phagosomes and co-localization index was determined.

Animal Infection

In-vivo experiment in this study was performed with approval from Institutional Animal Ethics

Committee (IAEC Approval No: IAEC/THSTI/165) as per guidelines laid by Committee for the

Purpose of Control of Experiments on Animals (CPCEA), Government of India. Briefly, bacterial

suspensions of respective strains at a density of 10^8 cells/mL were prepared in normal saline and

guinea pigs (Duncan Hartley) were infected via aerosol route using Inhalation Exposure System

(Glas-Col). Animals were euthanized at indicated time points, organs were harvested and bacterial

burdens were determined by CFU plating. Histopathological damage to the lung tissue was

determined by H&E staining of lungs fixed in 5% paraformaldehyde followed by granuloma scoring

of the infected lung tissue.

CRISPRi mediated *in-vivo* fitness

The experiment was performed with approval from Institutional Animal Ethics Committee (IAEC

Approval No: IAEC/THSTI/196) as per guidelines laid by Committee for the Purpose of Control of

Experiments on Animals (CPCEA), Government of India. Briefly, bacterial suspensions of respective

strains (NT and $\Delta Z::metB_{KD}$) at a density of $3\mathrm{x}10^8$ cells/mL were prepared in normal saline and

BALB/c mice were infected via aerosol route using Inhalation Exposure System (Glas-Col). To

induce CRISPRi mediated gene silencing of *metB* in $\Delta Z::metB_{KD}$ strain, doxycycline (1 mg/ml) was

administered to mice via oral route at indicated time points [53]. NT (non-targeting CRISPRi strain)

was maintained with doxycycline as control which is isogenic to Rv-WT. Animals were euthanized

at indicated time points, organs (lungs and spleens) were harvested and bacterial burden were

determined by CFU plating.

Bioinformatics analysis

MetZ protein sequence was retrieved from Mycobrowser

(https://mycobrowser.epfl.ch/genes/Rv0391) and the cluster of orthologous groups analysis (COG)

was performed using Eggnog v5 [54]. Further, a multiple sequence alignment of the protein sequence

from members of *Mycobacterium* genus (*M. tuberculosis, M. bovis, M. canettii, M. marinum, M.*

leprae and *M. smegmatis*) along with gram negative (*Pseudomonas aeruginosa*) and gram-positive

bacterium (*Corynebacterium glutamicum* and *Rhodococcus*) was performed using Clustal W [55] to

identify common domains and motifs amongst the protein sequences.

Molecular docking and molecular dynamics simulations

For system preparation, the crystal structure of putative O-succinylhomoserine sulfhydrylase of *M.*

tuberculosis was retrieved from RCSB PDB [56] (PDB-ID: 3NDN [57]). The protein has 4 chains and

exists in a homo-tetramer form. The receptor has a covalently bound pyridoxal-5-phosphate attached

to a modified lysine (K219). The substrate, O-succinyl homoserine, was retrieved from PubChem

(PubChemCID 439406). The substrate undergoes a sulfur transfer reaction in the presence of

pyridoxal-5-phosphate forming L-homocysteine which takes part in the methylation cycle. The

receptor protein was optimized and then minimized using the OPLS3 force field of Protein Preparation

Wizard module [58] of Schrodinger suite. Next, Binding Site Identification was done using Autodock

4 [59] wherein blind docking of the substrate in the presence and absence of the cofactor to identify

the possible binding site was performed. Autogrid4 was used to build a 126x126x126 points grid map

with a spacing of 0.586 angstrom and positioned at the center of mass of the receptor protein. The

Lamarckian genetic algorithm was employed to generate 200 runs. On docking the substrate in the

presence of the cofactor, no suitable results were produced. However, docking in the absence of the

co-factor resulted in comparatively higher binding affinity and a larger cluster size. Focused docking

on the identified site was also carried out by generating a grid map of 40x40x40 units with a spacing

of 0.375 A and 17X20X22 as grid center coordinates. MM-GBSA of few conformers based on their

binding affinity and cluster size was done using Prime module [60] of maestro. Finally, the lowest

energy conformations of the substrate were subsequently used for constant pH MD. Further, Constant

pH Molecular Dynamics Simulation (MDS) was performed with the Desmond [61] module from

Schrodinger suite. Both systems were protonated at pH 5.5 and 7.0, respectively, and minimized with

OPLS- all Atom force fields and solvated with the predefined TIP3P water solvent model. The

complexes were placed in the orthorhombic periodic boundary conditions to specify the shape and size

of the repeating unit buffered at 12 A distances. 4 Na+ and 9 Na+ ions were added to both the systems

respectively, to neutralize the systems. The NPT ensemble was employed for the simulation with a

timestep of 0.002 ps. The simulation was carried out for a total of 300 ns for each system and the

coordinates were saved at intervals of 20 ps to generate 15000 frames. Also, the protein was simulated

without the substrate at the two pH values. The post-processing of MD simulation trajectories was

carried out using VMD [62]. The root-mean-square-deviation and fluctuation were measured for the

protein, substrate and cofactor. Lastly, for thermodynamics analysis of the system we performed

Molecular Mechanics Generalized Born Surface Area (MM-GBSA) analysis to evaluate the MD

trajectories using Schrodinger software. To effectively encompass the thermodynamic properties,

every 500th frame was selected for MM-GBSA analysis.

Statistical Analysis

The data values are represented as mean ± SEM of the two-three independent experiments. Statistical

comparisons have been made with Rv-WT in most cases unless mentioned otherwise. GraphPad

Prism (ver 9.0) has been used for statistical analysis of all data with suitable statistical tests mentioned

with figure legend.

Conflict of Interest

The authors declare no conflict of interests

Acknowledgments

Authors acknowledge THSTI for infrastructural support, Department of Biotechnology, Government

of India for funding, Experimental Animal Facility for supporting animal experimentations and BSL-

3 facility for conducting all *M. tuberculosis* related experiments. Technical support of Mr. Surjit Yadav

is duly acknowledged. We acknowledge the Multi-Omics facility for LC-MS based experiments.

Funding

This study was supported by the grant (No. EM/Dev/SG/212/7864/2023) from the Indian Council of

Medical Research (ICMR), Government of India and intramural funding by BRIC-THSTI to A.K.P.

and are duly acknowledged. VKN is supported by a research fellowship from ICMR (3/1/3/JRF-

2018/HRD-043/64467).

Supplementary material

1. Table S1: List of plasmids and constructs used in the study

2. Table S2: List of bacterial used in the study

3. Table S3: List of primers used in the study

693   **List of relevant references:**
694

1.   WHO, *World TB report 2022.* 2023.

2.   Yelamanchi, S.D. and A. Surolia, *Targeting amino acid metabolism of Mycobacterium*
tuberculosis for developing inhibitors to curtail its survival. IUBMB Life, 2021. **73**(4): p. 643-
658.

3.   Adams, J.M. and M.R. Capecchi, *N-formylmethionyl-sRNA as the initiator of protein synthesis.*
Proc Natl Acad Sci U S A, 1966. **55**(1): p. 147-55.

4.   Kim, J.M., et al., *Formyl-methionine as an N-degron of a eukaryotic N-end rule pathway.*
Science, 2018. **362**(6418).

5.   Berney, M., et al., *Essential roles of methionine and S-adenosylmethionine in the autarkic*
lifestyle of Mycobacterium tuberculosis. Proc Natl Acad Sci U S A, 2015. **112**(32): p. 10008-
13.

6.   Chaton, C.T., et al., *Structural analysis of mycobacterial homoserine transacetylases central*
to methionine biosynthesis reveals druggable active site. Sci Rep, 2019. **9**(1): p. 20267.

7.   Chaudhary, D., et al., *Identification of small molecules targeting homoserine acetyl transferase*
from Mycobacterium tuberculosis and Staphylococcus aureus. Sci Rep, 2022. **12**(1): p. 13801.

8.   Berger, B.J. and M.H. Knodel, *Characterisation of methionine adenosyltransferase from*
Mycobacterium smegmatis and M. tuberculosis. BMC Microbiol, 2003. **3**: p. 12.

9.   Venos, E.S., et al., *Branched-chain amino acid aminotransferase and methionine formation in*
Mycobacterium tuberculosis. BMC Microbiol, 2004. **4**: p. 39.

10.   Lee, H.S. and B.J. Hwang, *Methionine biosynthesis and its regulation in Corynebacterium*
glutamicum: parallel pathways of transsulfuration and direct sulfhydrylation. Appl Microbiol
Biotechnol, 2003. **62**(5-6): p. 459-67.

11.   Hasenoehrl, E.J., et al., *Derailing the aspartate pathway of Mycobacterium tuberculosis to*
eradicate persistent infection. Nat Commun, 2019. **10**(1): p. 4215.

12.   Smith, D.A., et al., *Characterization of auxotrophic mutants of Mycobacterium tuberculosis*
and their potential as vaccine candidates. Infect Immun, 2001. **69**(2): p. 1142-50.

13.   Wheeler, P.R., et al., *Functional demonstration of reverse transsulfuration in the*
Mycobacterium tuberculosis complex reveals that methionine is the preferred sulfur source for
pathogenic Mycobacteria. J Biol Chem, 2005. **280**(9): p. 8069-78.

14.   Mishra, R., et al., *Targeting redox heterogeneity to counteract drug tolerance in replicating*
Mycobacterium tuberculosis. Sci Transl Med, 2019. **11**(518).

15.   Hatzios, S.K. and C.R. Bertozzi, *The regulation of sulfur metabolism in Mycobacterium*
tuberculosis. PLoS Pathog, 2011. **7**(7): p. e1002036.

16.   Foglino, M., et al., *A direct sulfhydrylation pathway is used for methionine biosynthesis in*
Pseudomonas aeruginosa. Microbiology (Reading), 1995. **141 (Pt 2)**: p. 431-9.

17.   Belfaiza, J., et al., *Direct sulfhydrylation for methionine biosynthesis in Leptospira meyeri.* J
Bacteriol, 1998. **180**(2): p. 250-5.

18.   Hwang, B.J., et al., *Corynebacterium glutamicum utilizes both transsulfuration and direct*
sulfhydrylation pathways for methionine biosynthesis. J Bacteriol, 2002. **184**(5): p. 1277-86.

19.   Griffin, J.E., et al., *High-resolution phenotypic profiling defines genes essential for*
mycobacterial growth and cholesterol catabolism. PLoS Pathog, 2011. **7**(9): p. e1002251.

20. Murphy, K.C., K. Papavinasasundaram, and C.M. Sassetti, *Mycobacterial recombineering.*
Methods Mol Biol, 2015. **1285**: p. 177-99.

21. Griffin, J.E., et al., *Cholesterol catabolism by Mycobacterium tuberculosis requires*
transcriptional and metabolic adaptations. Chem Biol, 2012. **19**(2): p. 218-27.

22. Chang, J.C., et al., *igr Genes and Mycobacterium tuberculosis cholesterol metabolism.* J
Bacteriol, 2009. **191**(16): p. 5232-9.

23. Srivastava, M., et al., *Molecular dynamics simulation reveals the possible druggable hot-spots*
of USP7. Oncotarget, 2018. **9**(76): p. 34289-34305.

24. Mittal, L., et al., *Targeting cryptic-orthosteric site of PD-L1 for inhibitor identification using*
structure-guided approach. Arch Biochem Biophys, 2021. **713**: p. 109059.

25. Kumari, A., et al., *Deciphering the Structural Determinants Critical in Attaining the FXR*
Partial Agonism. J Phys Chem B, 2023. **127**(2): p. 465-485.

26. Howe, M.D., et al., *Methionine Antagonizes para-Aminosalicylic Acid Activity via Affecting*
Folate Precursor Biosynthesis in Mycobacterium tuberculosis. Front Cell Infect Microbiol,
2018. **8**: p. 399.

27. Khan, M.Z., et al., *Redox homeostasis in Mycobacterium tuberculosis is modulated by a novel*
actinomycete-specific transcription factor. EMBO J, 2021. **40**(14): p. e106111.

28. Gouzy, A., et al., *Growth of Mycobacterium tuberculosis at acidic pH depends on lipid*
assimilation and is accompanied by reduced GAPDH activity. Proc Natl Acad Sci U S A, 2021.
118(32).

29. Loewen, P.C., *Levels of glutathione in Escherichia coli.* Can J Biochem, 1979. **57**(2): p. 107-
11.

30. Akhova, A.V. and A.G. Tkachenko, *ATP/ADP alteration as a sign of the oxidative stress*
development in Escherichia coli cells under antibiotic treatment. FEMS Microbiol Lett, 2014.
353(1): p. 69-76.

31. Singh, S., et al., *Unraveling the role of the transcriptional regulator VirS in low pH-induced*
responses of Mycobacterium tuberculosis and identification of VirS inhibitors. J Biol Chem,
2019. **294**(26): p. 10055-10075.

32.   Mehta, M., R.S. Rajmani, and A. Singh, *Mycobacterium tuberculosis WhiB3 Responds to*
Vacuolar pH-induced Changes in Mycothiol Redox Potential to Modulate Phagosomal
Maturation and Virulence. J Biol Chem, 2016. **291**(6): p. 2888-903.

33.   Banerjee, S.K., et al., *The sensor kinase MtrB of Mycobacterium tuberculosis regulates hypoxic*
survival and establishment of infection. J Biol Chem, 2019. **294**(52): p. 19862-19876.

34.   Choudhary, E., et al., *Deciphering the Proteomic Landscape of Mycobacterium tuberculosis in*
Response to Acid and Oxidative Stresses. ACS Omega, 2022. **7**(30): p. 26749-26766.

35.   Rohde, K.H., et al., *Linking the transcriptional profiles and the physiological states of*
Mycobacterium tuberculosis during an extended intracellular infection. PLoS Pathog, 2012.
8(6): p. e1002769.

36.   Saini, V., et al., *Hydrogen sulfide stimulates Mycobacterium tuberculosis respiration, growth*
and pathogenesis. Nat Commun, 2020. **11**(1): p. 557.

37.   Kunota, T.T.R., et al., *Mycobacterium tuberculosis H2S Functions as a Sink to Modulate*
Central Metabolism, Bioenergetics, and Drug Susceptibility. Antioxidants (Basel), 2021. **10**(8).

38.   Rahman, M.A., et al., *The Role of Host-Generated H2S in Microbial Pathogenesis: New*
Perspectives on Tuberculosis. Front Cell Infect Microbiol, 2020. **10**: p. 586923.

39.   Rahman, M.A., et al., *Hydrogen sulfide dysregulates the immune response by suppressing*
central carbon metabolism to promote tuberculosis. Proc Natl Acad Sci U S A, 2020. **117**(12):
p. 6663-6674.

40.   Brauner, A., et al., *Distinguishing between resistance, tolerance and persistence to antibiotic*
treatment. Nat Rev Microbiol, 2016. **14**(5): p. 320-30.

41.   Quinonez, C.G., et al., *The Role of Fatty Acid Metabolism in Drug Tolerance of Mycobacterium*
tuberculosis. mBio, 2022. **13**(1): p. e0355921.

42.   Wu, J., et al., *Derivatives of Natural Product Agrimophol as Disruptors of Intrabacterial pH*
Homeostasis in Mycobacterium tuberculosis. ACS Infect Dis, 2019. **5**(7): p. 1087-1104.

43.   Andreu, N., et al., *Optimisation of bioluminescent reporters for use with mycobacteria.* PLoS
One, 2010. **5**(5): p. e10777.

44.   Babunovic, G.H., et al., *CRISPR Interference Reveals That All-Trans-Retinoic Acid Promotes*
792   *Macrophage Control of Mycobacterium tuberculosis by Limiting Bacterial Access to*
793   *Cholesterol and Propionyl Coenzyme A.* mBio, 2022: p. e0368321.

794   45.   Pandey, A.K. and C.M. Sassetti, *Mycobacterial persistence requires the utilization of host*
795   *cholesterol.* Proc Natl Acad Sci U S A, 2008. **105**(11): p. 4376-80.

796   46.   Schmittgen, T.D. and K.J. Livak, *Analyzing real-time PCR data by the comparative C(T)*
797   *method.* Nat Protoc, 2008. **3**(6): p. 1101-8.

798   47.   Vocat, A., et al., *Bioluminescence for assessing drug potency against nonreplicating*
799   *Mycobacterium tuberculosis.* Antimicrob Agents Chemother, 2015. **59**(7): p. 4012-9.

800   48.   Bhaskar, A., et al., *Reengineering redox sensitive GFP to measure mycothiol redox potential*
801   *of Mycobacterium tuberculosis during infection.* PLoS Pathog, 2014. **10**(1): p. e1003902.

802   49.   Folch, J., M. Lees, and G.H. Sloane Stanley, *A simple method for the isolation and purification*
803   *of total lipides from animal tissues.* J Biol Chem, 1957. **226**(1): p. 497-509.

804   50.   Nambi, S., et al., *The Oxidative Stress Network of Mycobacterium tuberculosis Reveals*
805   *Coordination between Radical Detoxification Systems.* Cell Host Microbe, 2015. **17**(6): p. 829-
806   37.

807   51.   Mishra, R., et al., *Heterogeneous Host-Pathogen Encounters Coordinate Antibiotic Resilience*
808   *in Mycobacterium tuberculosis.* Trends Microbiol, 2021. **29**(7): p. 606-620.

809   52.   Cai, Y., et al., *Host immunity increases Mycobacterium tuberculosis reliance on cytochrome*
810   *bd oxidase.* PLoS Pathog, 2021. **17**(7): p. e1008911.

811   53.   Tripathi, A., et al., *Mycobacterium tuberculosis requires SufT for Fe-S cluster maturation,*
812   *metabolism, and survival in vivo.* PLoS Pathog, 2022. **18**(4): p. e1010475.

813   54.   Huerta-Cepas, J., et al., *eggNOG 5.0: a hierarchical, functionally and phylogenetically*
814   *annotated orthology resource based on 5090 organisms and 2502 viruses.* Nucleic Acids Res,
815   2019. **47**(D1): p. D309-D314.

816   55.   Thompson, J.D., D.G. Higgins, and T.J. Gibson, *CLUSTAL W: improving the sensitivity of*
817   *progressive multiple sequence alignment through sequence weighting, position-specific gap*
818   *penalties and weight matrix choice.* Nucleic Acids Res, 1994. **22**(22): p. 4673-80.

56.   Berman, H.M., et al., *The Protein Data Bank.* Nucleic Acids Res, 2000. **28**(1): p. 235-42.

57.   Baugh, L., et al., *Increasing the structural coverage of tuberculosis drug targets.* Tuberculosis
(Edinb), 2015. **95**(2): p. 142-8.

58.   Sastry, G.M., et al., *Protein and ligand preparation: parameters, protocols, and influence on
virtual screening enrichments.* J Comput Aided Mol Des, 2013. **27**(3): p. 221-34.

59.   Morris, G.M., et al., *AutoDock4 and AutoDockTools4: Automated docking with selective
receptor flexibility.* J Comput Chem, 2009. **30**(16): p. 2785-91.

60.   Jacobson, M.P., et al., *A hierarchical approach to all-atom protein loop prediction.* Proteins,
2004. **55**(2): p. 351-67.

61.   Bowers, K.J.C., E.; Xu, H.; Dror, R. O.; Eastwood, M. P.; Gregersen, B. A.; Klepeis, J. L.;
Kolossvary, I.; Moraes, M. A.; Sacerdoti, F. D.; Salmon, J. K.; Shan, Y.; Shaw, D. E., , *Scalable
algorithms for molecular dynamics simulations on commodity clusters.* Proceedings of the
ACM/IEEE Conference on Supercomputing (SC06), 2006.

62.   Humphrey, W., A. Dalke, and K. Schulten, *VMD: visual molecular dynamics.* J Mol Graph,
1996. **14**(1): p. 33-8, 27-8.

Figures and Legends.

Figure 1. *metZ* is dispensable for growth in *M. tuberculosis.* (A). Methionine biosynthesis pathways in

bacteria. There exist two alternative pathways in many bacterial species – transsulfuration where cysteine

is used as sulfur-donor (S-donor) and direct sulfhydrylation where in H_2S is used as S-donor. *Rv0391*

encoded MetZ is such putative O-succinylhomoserine sulfhydrylase (OSHS) involved in direct

sulfhydrylation in *M. tuberculosis*. (B). Cluster of orthologues analysis for *M. tuberculosis* MetZ (Rv0391)

protein sequence with member species of ENOG502GJ5S as indicated by EggNog database. (C). Schematic

for knockout generation of *ΔmetZ* in *M. tuberculosis.* (D). Confirmation of *metZ* knockout using RT-qPCR

for the loss of *metZ* transcript. Growth kinetics of auto bioluminescent *ΔmetZ* with (E) glycerol (0.1%) and

(F) cholesterol (150μM) as C-source. The data summary represents mean ± SEM for technical replicates

and is representation of two independent experiments. Statistical comparisons were made w.r.t. Rv-WT for

(D) using Unpaired *t*-test. ****, *P*<0.0001. Comparisons not marked with statistical significance were not

significant with Rv-WT vs *ΔmetZ* for (E) and (F).

Figure 2: ***metZ* deletion renders *M. tuberculosis* sensitive to physiological stressors and lead to reduced *ex-vivo* fitness.** (A). *In-vitro* stress survival assay. Mycobacterial cultures were treated with reagents mimicking host-induced stresses: H_2O_2 (Oxidative stress), DetaNO (Nitrosative stress), SDS (Surfactant stress), low pH (5.0), PSBT (nutrient starvation) *in-vitro* and survival was determined by CFU plating. *ΔmetZ* was comparatively more susceptible (~40-60%) to the stressors. (B). Intra bacterial redox state determination of *ΔmetZ* as compared to Rv-WT using genetically encoded redox biosensor Mrx1-roGFP2. (C). Lipid hydroperoxidation levels in *ΔmetZ* compared to Rv-WT and complement strains using FOX2 colorimetric assay. (D). Intra-macrophage survival assay using THP-1 monocytes derived macrophages either resting or activated using IFN-γ. Survival index was determined by CFU enumeration. (E). Intracellular co-localization of GFP labeled mycobacterial strains with LysoTracker stained acidic compartments inside IFN-γ activated macrophages. Colocalization coefficient in case of *ΔmetZ* was recorded ~1.2 fold higher than Rv-WT and complement strains. The points represent replicates per group and data represented is where *ΔmetZ* is compared to Rv-WT for statistical significance. Statistics were performed using two-way ANOVA (multiple comparisons) for (A), (B), (D) and Ordinary one-way ANOVA (Dunnett's multiple comparisons test) for (C) and (E). *, P<0.05; **, P<0.01; ***P<0.001; ****P<0.0001; ns, non-significant.

Figure 3: Direct sulfhydrylation pathway is essential for *M. tuberculosis* to survive under low pH.
(A) *In-vitro* growth kinetics of *ΔmetZ* at various pH – 7.2, 5.5 and 5.0. (B). Expression analysis of genes involved in transsulfuration (*metB* and *metC*) and direct sulfhydrylation (*metZ*) using RT-qPCR. Relative expression of *metZ* was ~12-fold higher at acidic pH as compared to neutral pH. *Icl1* was used as a positive control for acidic conditions. (C). Metabolic activity of *ΔmetZ* at acidic pH using redox dye AlamarBlue Assay. (D). Intra bacterial ATP levels estimation of *ΔmetZ* at acidic pH 5.5. (E). Pyrazinamide sensitivity assay of *ΔmetZ* at pH 5.5. IC50 of PYZ for the respective strains at acidic pH 5.5 were determined and represented as dose-response curve. (F). Methionine estimation in Rv-WT and *ΔmetZ* at pH 7.2 and pH 5.5 was done using LC-MS. (G). Growth defect in *ΔmetZ* at pH 5.5 was rescued using exogenous supplementation of methionine. Dose-response curve for PYZ was made by Non-linear regression analysis using GraphPad Prism. Data is represented as the mean ± SEM of the replicates indicated by points in the respective graphs. Comparisons for statistical significance have been made with Rv-WT (except B) using two-way ANOVA (multiple comparisons) for (A) and (G); ordinary one-way ANOVA (Dunnett's Multiple Comparisons Test) for (B), (C) and (D) and ordinary one-way ANOVA (Sidak's Multiple Comparisons Test) for (F). *, P<0.01; **, P<0.002; ***P<0.005; ****P<0.0001; ns, non-significant.

Figure 4. Molecular docking and MD simulations revealed enhanced stability of MetZ at acidic pH.

(A). Structural representation of O-succinylhomoserine sulfhydrylase, rendered in new cartoon and colored in gray (single chain) with pyridoxal-5-phosphate in licorice and rendered atom-wise. (B). Dynamic characteristics of systems elucidated through constant pH MD simulations. The system's evolution is measured via RMSD through the course of 300ns of the protein, substrate and the binding site of the substrate. Color coding is followed as a system at pH 7 (blue) and at pH 5.5 (red). (C). The Binding Free Energy (in Kcal/mol) comparison of the complexes at pH 7 (blue) and pH 5.5 (red). (D). The dynamic changes represent the positional shifts of the substrate (orange) and co-factor (cyan) throughout the simulation, with a color gradient (time-dependent manner) from red (initial), white (intermediate) to blue (final) at pH 7 (a) at pH 5.5 (b). (E). Interaction map of the substrate at pH 7 and pH 5.5. Residues lining the pocket (within 4.0 Å) are shown in licorice and colored atom-wise (C: cyan, N: blue and O: red). The residues forming hydrogen bonds (dotted line in blue) and Salt-bridges (dotted line in yellow) are shown. The Cα atoms of residues involved in hydrophobic interaction are shown in beads.

Figure 5: *metZ* mediated direct sulfhydrylation pathway is critical for methionine biosynthesis at low pH. (A). Diagrammatic representation of experimental setup to elucidate independent existence of direct sulfhydrylation pathway. CRISPRi based knockdown of *metB* (*Rv1079*) strains were generated in the genetic backgrounds of Rv-WT and *ΔmetZ* strains. These strains upon induction with ATc, were depleted of *metB* transcripts and characterized for *in-vitro* growth kinetics, methionine levels and *in-vivo* fitness. (B). *In-vitro* growth kinetics of CRISPRi knockdown strains. (C). Relative methionine levels in CRISPRi knockdown strains of *metB* and double mutant lacking both *metZ* and *metB* by LC-MS. (D). Graphical representation of *in-vivo* fitness experiment. Mice were infected with CRISPRi knockdown of *metB* in Rv-WT and *ΔmetZ* background and orally fed with doxycycline (1 mg/ml) to induce knockdown of *metB* gene in the respective strains. Bacterial burden was determined by CFU plating in lungs and spleen homogenates. (E).and (F). Bacterial burden of mycobacterial strains lacking either *metZ*, *metB* or both in mice lungs and spleen lysates respectively. Dots represent replicates per group. Statistical significance has been made using ordinary one-way ANOVA (Dunnett's Multiple Comparisons test) for (B) where ****$P<0.0001$; Unpaired t-test for (C) where ***$P=0.0001$; Mann-Whitney Test for Spleen bacterial burden in (E) where *$P=0.015$, **$P=0.007$ and two-way ANOVA (multiple comparisons) for Lungs bacterial burden in (E) where *$P=0.02$, **$P=0.004$, ***$P=0.0003$ and ****$P<0.0001$.

Figure 6: Direct sulfhydrylation pathway of *M. tuberculosis* contributes towards antibiotic persistence in response to Bedaquiline. (A). Minimum Inhibitory Concentration (MIC) determination of the *ΔmetZ* in comparison to Rv-WT and Complement strains against anti-TB drugs: Rifampicin (RIF), Isoniazid (INH), Ethambutol (EMB) and Bedaquiline (BDQ) using microtiter plate assay. IC50 was determined for the drugs with mentioned strains. (B). Kill kinetics of anti-TB drugs – RIF, INH, EMB and BDQ at 10X MIC concentrations. Survival was determined in the form of viable bacterial count expressed as CFU. (C). Kinetics of ATP depletion post BDQ treatment. *ΔmetZ* along with Rv-WT and Complement strains were treated with 2X concentration of BDQ and ATP was estimated at day 8 wherein *ΔmetZ* maintained high ATP levels. Data is represented as mean ± SEM and dots represent replicates per group. Comparison for statistical significance has been made using two-way ANOVA (Dunnett's Multiple Comparisons test) for (B) where **P<0.001 and ns, non-significance; ordinary one-way ANOVA (Dunnett's Multiple Comparisons test) where ****P<0.0001 and ns, non-significance.

Figure 7: *metZ* is required for survival of *M. tuberculosis in-vivo*. (A). Diagrammatic representation of *in-vivo* fitness experimental set-up. Guinea pigs were infected with WT, mutant and complement strains and disease progression was monitored through CFU plating at indicated time points. (B). Bacterial burden in guinea pig lungs as determined by CFU plating of the lung homogenates of the infected animals at indicated time points. Mutant strain failed to replicate in guinea pig lungs post week 4 of initial infection. (C). CFU burden in spleens of the infected animals revealed that the mutant failed to survive inside the spleen post week 4 of infection. (D). Gross lung pathology indication lesions on the lungs at week 8 time point post aerosol infection. (E). H&E staining (F). Total granuloma scores of three strains in lung sections revealed significantly lesser granulomas in *ΔmetZ* infected lungs at week 8 post infection. Dots indicate the number of animals (or samples) per group and data is represented as mean ± SEM. Comparison for determining statistical significance has been made using Mann-Whitney Test for (B) and (C) where *P<0.05, **P=0.0079; ordinary one-way ANOVA (Dunnett's Multiple Comparisons test) where **P=0.004, ***P=0.0004.

Figure 8. *metZ* driven direct sulfhydrylation is critical for methionine biosynthesis at low pH. When *M. tuberculosis* infects macrophage, it passes through a series of compartments starting from phagosome where pH is essentially ~6.5 and the bacteria is in replicating state, with enough nutrients (including methionine). Further, as host immunity kicks in and IFN-γ induces accumulation of ROS, RNS and subsequently, phagosome maturation leading to drop in pH to 5.5 and beyond, conditions are non-replicating for *M. tuberculosis* and host begins to quench nutrients from bacteria as means to impose restriction on the trapped bacteria. At this stage bacteria undergo several transcriptional and metabolic changes to sustain the host induced insults, one of these includes induction of *metZ* expression that helps bacteria survive these stress conditions and helps maintain methionine levels by utilizing endogenous as well as host generated H_2S which acts as co-substrate and S-donor for methionine biosynthesis in acidic micro-environment. This would help bacteria to maintain sufficient levels of methionine to keep adaptation going while also helps recycling of ATP, mycothiol and its metabolic activity keep going.

Supplementary Figures

pH dependent direct sulfhydrylation pathway is required for mycobacterial pathogenesis

Vaibhav Kumar Nain[1,2], Vishawjeet Barik[1,2#], Manitosh Pandey[1#], Mohit Pareek[3], Taruna Sharma[1,2], Rahul

Pal[1], Shaifali Tyagi[1], Manish Bajpai[4], Prabhanjan Dwivedi[5], Bhishma Narayan Panda[5], Yashwant Kumar[4],

Shailendra Asthana[3] and Amit Kumar Pandey[1,5*]

[1]Mycobacterial Pathogenesis Laboratory, Centre for Tuberculosis Research, BRIC-Translational Health

Science and Technology Institute, Haryana, India

[2]Special Centre for Molecular Medicine, Jawaharlal Nehru University, New Delhi, India

[3]Computational Biophysics and CADD Group, Computational and Mathematical Biology Centre, BRIC-

Translational Health Science and Technology Institute, Haryana, India

[4]Biomarker Discovery Laboratory, Non-Communicable Diseases Centre, BRIC-Translational Health

Science and Technology Institute, Haryana, India

[5]Experimental Animal Facility, BRIC-Translational Health Science and Technology Institute, Haryana, India

[#] Contributed equally to this work

* Correspondence: Amit Kumar Pandey, amitpandey@thsti.res.in

Figure S1: Rv0391 (MetZ) sequence is conserved across *Mycobacterium* genus (A). Protein sequence alignment of MetZ across other members of MTBC. Multiple sequence alignment (MSA) was done using bacterial species with gram negative (*Pseudomonas aeruginosa*) and gram-positive bacterium (*Corynebacterium glutamicum* and *Rhodococcus*) along with other mycobacterial species (*M. tuberculosis, M. bovis, M. canettii, M. marinum, M. leprae* and *M. smegmatis*). MSA revealed similarity amongst analyzed sequences wrt. Several motifs give rise to similar secondary structure (indicated by coils on top of the alignment) and conserved lysine (K219), marked by blue arrow head at which the cofactor pyridoxal-5'-phosphate is covalently attached. Presence of a characterized OSHS protein in *C. glutamicum*, in particular with similar motifs to Rv0391 indicates the probable role of MetZ (Rv0391) as an OSHS.

Figure S2: *Rv0391 (metZ)* genetic mutant confirmation. (A). Mutant confirmation by gene-specific PCR. Genomic DNA was isolated from the unmarked mutant culture and WT *M. tuberculosis* using which 1.4kb sized fragment encoding *metZ* gene was amplified. The unmarked mutant carried deletion of a major fragment of the said ORF thus, showing an amplicon of 153bp, confirming loss of *metZ* encoding sequence. (B). Genetic locus encoding *metZ* was sequenced in the unmarked mutant wherein, loss of the functional ORF confirmed the generation of mutant when compared to WT sequence. Post unmarking, a 34bp loxP sequence remains in the bacterial chromosome within the target sequence which serves as indicator of gene deletion and unmarking of the mutant, thereby confirming loss of Hyg[r] cassette.

Figure S3: Generation of auto-bioluminescent Rv, *ΔmetZ* and complement stains and growth kinetics. (A). Linear regression between Luminescence and CFU in *M. tuberculosis*. Log-phase culture of *M. tuberculosis*-lux strain was grown and ~10 OD equivalent cells were serially diluted (1:10) through 0.0001. For each dilution, bacterial enumeration was done by CFU plating and luminescence was recorded. Linear relationship between luminescence and CFU was determined using linear regression analysis categorically indicating that luminescence can be used as a marker for bacterial viability and an alternative to CFU. (B) *In-vitro* growth kinetics of *ΔmetZ* in propionate (10mM) as sole C-source. Propionate is a short, odd chain fatty acid and an end product of cholesterol and fatty acid catabolism in *M. tuberculosis*.

Figure S4: pH dependent growth kinetics and sulfide estimation. (A). *In-vitro* growth kinetics of *ΔmetZ* at pH 6.5. (B). *In-vitro* growth kinetics of *ΔmetZ* at pH 5.5. (C). Viable bacterial count of *ΔmetZ* in comparison to Rv-WT and Complement strains at pH 5.5 (D). Endogenous H_2S estimation in bacterial cultures of respective strains at pH 7.2 and pH 5.5 using Bismuth chloride-based colorimetric assay. Data is represented as the mean ± SEM of the replicates indicated by points in the respective graphs. Comparisons for statistical significance have been made with Rv-WT using two-way ANOVA (Dunnett's multiple comparisons test) for (B) and (C); ordinary one-way ANOVA (Sidak's Multiple Comparisons Test) for (D). **, P=0.005; ***P=0.0003; ****P<0.0001; ns, non-significant.

Figure S5: Confirmation of *metB_{KD}* using RT-qPCR. (A). CRISPRi strains were induced with ATc (200 ng/ml) and total RNA was isolated and reverse transcribed to cDNA. Further, fold expression w.r.t. ATc untreated cultures in both *Rv::metB_{KD}* and *ΔmetZ::metB_{KD}* strains were determined using the ΔΔCt method. *metB* levels were significantly reduced by several folds. Data

87 is represented as mean ± SEM and replicates are indicated by dots. Comparison for statistical
88 significance has been made using Unpaired t-test where ****P<0.0001.

B MIC of anti-TB drugs

Drugs	Rv-WT	*ΔmetZ*	*ΔmetZ* Comp
	MIC_{50} (µg/ml)		
RIF	0.008	0.009	0.014
INH	0.075	0.060	0.061
BDQ	0.041	0.032	0.051
EMB	1.588	1.466	1.480
CFZ	0.089	0.091	0.051
PYZ	0.669	0.051	0.262

Figure S6: Drug susceptibility testing (DST) of anti-TB drugs on *Rv-WT*, *ΔmetZ* and
Complement strain. (A). Dose-response curve of Clofazimine (CFZ) against the mycobacterial
strains and its IC50. (B). Table summarizing IC50 values of anti-TB drugs with the respective
mycobacterial strains. (C). Bedaquiline dependent ATP depletion kinetics in the indicated
mycobacterial strains. Mycobacterial strains were treated with 2X BDQ for 8 days and ATP was
estimated using the BacTiter-Glo kit at the indicated time-points. Data is represented as mean±SEM.
Comparisons for statistical significance were made with *ΔmetZ* group using 2-way ANOVA
(Dunnett's Multiple Comparisons test) where ****P, <0.0001 and ns, non-significant.

Figure S7: Intracellular survival assay. (A). Resting or IFN-g activated THP-1 macrophages were infected with Rv-WT, *ΔmetZ* and Complement strains and bacterial survival was determined by plating macrophages lysates at the indicated time-points. CFU counts show that *ΔmetZ* has compromised fitness inside resting macrophages. *ΔmetZ* CFU count is further reduced inside activated macrophages. Dots represent the replicates per group and data is shown as mean ± SEM. Comparisons have been made with Rv-WT for statistical significance using 2-way ANOVA (Dunnett's Multiple Comparisons test) where ****P <0.0001. The Survival Index calculated from the CFU values is shown in Figure 2D.

Figure S8: Panel showing intracellular co-localization of the Rv-WT, *ΔmetZ* and Complement strains. mEmerald expressing mycobacterial strains were used to infect IFN-γ activated THP-1 macrophages. Two days after infection, the cells were fixed and stained with DAPI (for nucleus) and LysoTracker Red. *ΔmetZ* cells were co-localized with acidic regions of the cells (stained in red) as also indicated by higher colocalization coefficients as shown in Figure 2E. White bars indicate the scale corresponding to 20μM length.

List of bacterial strains used in the study

S. No.	Strain	Marker	Feature	Reference
1.	*E. coli* XL1Blue			This lab
2.	*M. tuberculosis* H37Rv		Wild type *Mtb*.	Kind gift from Christopher M. Sassetti
3.	*ΔprpR::Hyg^r*	*Hyg^r*	*prpR* deletion mutant with Hyg cassette inserted in chromosomal copy of *prpR* gene (*Rv1129c*)	Kind gift from Christopher M. Sassetti
4.	*ΔprpR::Hyg^r::lux*	*Hyg^r, Kan^r*	*prpR* marked deletion mutant with luciferase reporter integrated at L5 attP site	This work
5.	*ΔmetZ*		*Rv0391*, *metZ* deletion mutant, umarked	This work
6.	*ΔmetZ Comp*	*Kan^r*	*ΔmetZ* mutant expressing pST-K:*metZ* plasmid	This work
7.	Rv::*lux*	*Zeo^r,*	Tweety int site, Luciferase reporter	This work
8.	*ΔmetZ::lux*	*Zeo^r,*	Tweety int site, Luciferase reporter	This work
9.	*ΔmetZ Comp::lux*	*Kan^r, Zeo^r,*	Tweety int site, Luciferase reporter	This work
10.	Rv: pMV261 mEmerald, TagRFP	*Hyg^r,*	Mycobacterial Live Dead Reporter expressing Rv-WT	This work
11.	*ΔmetZ*: pMV261 mEmerald, TagRFP	*Hyg^r,*	Mycobacterial Live Dead Reporter expressing *ΔmetZ* mutant	This work
12.	*ΔmetZ Comp*: pMV261 mEmerald, TagRFP	*Kan^r, Hyg^r*	Mycobacterial Live Dead Reporter expressing *ΔmetZ Comp* strain	This work
13.	*Rv: Mrx1-roGFP2*	*Hyg^r,*	Redox biosensor expressing Rv-WT	This work
14.	*ΔmetZ: Mrx1-roGFP21*	*Hyg^r,*	Redox biosensor expressing *ΔmetZ* mutant	This work
15.	*Rv::metB_{KD}*	*Kan^r*	WT-Mtb with *metB* gene knocked down using CRISPRi	This work
16.	*ΔmetZ::metB_{KD}*	*Kan^r*	*ΔmetZ* with *metB* gene knocked down using CRISPRi	This work
17.	*Rv::NT_{KD}*	*Kan^r*	WT-Mtb with control sgRNA (Non-targetting) CRISPRi control	This work

List of plasmids used in the study

S. No.	Plasmid	Feature	Reference
1.	pJM1	Hyg/Cm, suicidal vector for generating knockout construct for gene deletion in *M. tuberculosis*	Kind gift from Christopher M. Sassetti
2.	pST-K	Kan, episomal, mycobacterial expression vector	Kind gift from Vinay K. Nandicoori, NII, India
3.	pST-Ki	Kan, attB integrative, mycobacterial expression vector	Kind gift from Vinay K. Nandicoori, NII, India
4.	pST-K:*metZ*	Kan, pST-K backbone with metZ encoding ORF	This study
5.	pST-Ki:*metZ*	Kan, pST-Ki backbone with metZ encoding ORF	This study
6.	pMV261: mEmerald, TagRFP	Hyg, Mycobacterial Live-Dead reporter with ATc inducible RFP	Kind gift from Christopher M. Sassetti
7.	pJW276	Zeo, Integrative, Lux expressing plasmid	Kind gift from Sarah M. Fortune
8.	pMV306:LuxAB+G13	Kan, Integrative, Lux expressing plasmid	Addgene # 26161
9.	pMV762: Mrx1-roGFP2	Hyg, episomal, genetically encoded redox biosensor	Kind gift from Amit Singh, IISc, India
10.	plJR965	Kan, attB integrative, target sgRNA and *dCas9$_{Sth}$* expressing plasmid for transcriptional knockdown of target gene(s) in *M. tuberculosis*	Kind gift from Sarah M. Fortune
11.	plJR965::*metB*_sgRNA	CRISPRi knockdown construct for targeting *metB* (*Rv1079*) gene in *M. tuberculosis*	This study

List of primers used in the study

S. No.	Primer Name	Sequence 5'-3'	Feature
1.	*metZ*-F1	GCTTAATTAAGTCGAGATCGAGTTTTTGGTCACC	Cloning of 1000bp upstream region of *metZ* gene for knockout construct
2.	*metZ*-R1	ATGCGGCCGCGCGGACCGAAGACTCGTC	
3.	*metZ*-F2	ATCTCGAGACCGACGACCTGATTGCCGAT	Cloning of 1000bp downstream region of *metZ* gene for knockout construct
4.	*metZ*-R2	ATCTCGAGCGGAATGAAGACCATCGACGAC	
5.	*metZ*-F	AACATATGATGACCGACGAGTCTTCGGTC	Cloning of *metZ* encoding ORF in *NdeI/ EcoRI* sites
6.	*metZ*-R	AAGAATTCTTAGCTCAACGCCCGATCGATATC	
7.	*metB*-CR-F	GGGAACCGCGCCGGCGCCGTTCTG	Cloning of *metB* targeting sgRNA
8.	*metB*-CR-R	AAACCAGAACGGCGCCGGCGCGGT	
9.	*metB*-RT-F	GGCGGCACATTCCGGTTGATA	qPCR primers of *metB*
10.	*metB*-RT-R	GTGAGGCAAAGGTATTGTCCACCA	
11.	*metC*-RT-F	GACCATCTCCAACCCGCAG	qPCR primers of *metC*
12.	*metC*-RT-R	GTCGGGGGTGGTGAAGC	
13.	*metZ*-RT-F	GACGACCTCTCGCAATGGGA	qPCR primers of *metZ*
14.	*metZ*-RT-R	GTACACCACCACGTCGACC	
15.	*sigA*-RT-F	GAAGACCACGAAGACCTCGAAG	qPCR primers of *sigA*
16.	*sigA*-RT-R	GTTTGAGGTAGGCGCGAACC	
17.	*Icl1*-RT-F		qPCR primers of *icl1*
18.	*Icl1*-RT-R		