pH-dependent direct sulfhydrylation pathway is required for the pathogenesis of *Mycobacterium*

tuberculosis

Vaibhav Kumar Nain[1,2], Vishawjeet Barik[1,2#], Manitosh Pandey[1#], Mohit Pareek[3], Taruna Sharma[1,2],

Rahul Pal[1], Shaifali Tyagi[1], Manish Bajpai[4], Prabhanjan Dwivedi[5], Bhishma Narayan Panda[5],

Yashwant Kumar[4], Shailendra Asthana[3] and Amit Kumar Pandey[1,5*]

[1]Mycobacterial Pathogenesis Laboratory, Centre for Tuberculosis Research, BRIC-Translational

Health Science and Technology Institute, Haryana, India

[2]Special Centre for Molecular Medicine, Jawaharlal Nehru University, New Delhi, India

[3]Computational Biophysics and CADD Group, Computational and Mathematical Biology Centre,

BRIC-Translational Health Science and Technology Institute, Haryana, India

[4]Biomarker Discovery Laboratory, Non-Communicable Diseases Centre, BRIC-Translational Health

Science and Technology Institute, Haryana, India

[5]Experimental Animal Facility, BRIC-Translational Health Science and Technology Institute,

Haryana, India

[#] Contributed equally to this work

* Correspondence: Amit Kumar Pandey, amitpandey@thsti.res.in

Abstract

Methionine is essential for the survival of *Mycobacterium tuberculosis* (*M. tuberculosis*) inside the

host. However, the transsulfuration pathway, a major contributor of methionine, is dispensable for the

growth of *M. tuberculosis* suggesting redundancy in the methionine biosynthesis pathway. Orthologues

of MetZ$_{TB}$ in other bacterial species are known to operate a redundant single-step methionine

biosynthesis pathway called *direct sulfhydrylation.* In this study, we demonstrate that genetic disruption of the *metZ*-mediated direct sulfhydrylation pathway in *M. tuberculosis* hinders growth at low pH, an effect mitigated by methionine supplementation. Computational analyses, including *in-silico* molecular docking and molecular dynamics (MD) simulations, reveal enhanced binding of the MetZ substrate, O-succinyl homoserine (OSH), to the active site of MetZ at acidic pH. Intriguingly, despite increased intracellular ATP levels, a relative decrease in the frequency of Bedaquiline (BDQ)-induced persisters is observed in *metZ*-deficient strain, suggesting a role of direct sulfhydrylation pathway in modulating BDQ sensitivity. Finally, we demonstrated that the absence of *metZ* impedes the ability of *M. tuberculosis* to grow inside the host.

Keywords: tuberculosis, methionine biosynthesis, acidic pH, drug tolerance, transsulfuration, direct sulfhydrylation.

Introduction

Tuberculosis, an ancient disease, continues to pose a serious global health challenge, negatively impacting economies and the public health systems worldwide. In 2023, *Mycobacterium tuberculosis* (*M. tuberculosis*) caused 1.8 million deaths, making it the leading cause of death from a single infectious agent [1]. A steady increase in the frequency of drug-resistant tuberculosis cases has significantly complicated the therapeutic landscape. A majority of the approved drugs target pathways that are essential for the survival of *M. tuberculosis*. Targeting biosynthesis of essential amino acids as a therapeutic strategy has demonstrated immense potential and has yielded several molecules with the potential to become a future drug [2].

Methionine plays a pivotal role in cellular function, serving as the initiator amino acid, *N-formyl*-L-methionine (*f*Met), in protein synthesis in prokaryotes   as well as protein degradation in eukaryotes [3]. Methionine also acts as a precursor for biosynthesis of essential metabolites that includes cysteine, biotin and SAM (S-adenosyl methionine). Given its critical importance, inhibiting some of the essential

proteins involved in the methionine biosynthesis pathway has been extensively investigated as a potential strategy to identify new drugs against tuberculosis [4-8]. The transsulfuration pathway, comprising a series of enzymatic reactions and intermediates involved in methionine biosynthesis, is largely conserved across various microbial species and has been extensively reviewed [9]. Although biosynthesis of methionine is known to be essential for the growth of *M. tuberculosis* [4, 6, 10], some genes in this pathway are dispensable, indicating potential redundancy or conditional essentiality [11-13]. Several reports suggest the existence of an alternate biochemical reaction in which the metabolite O-succinyl homoserine (OSH) is directly converted into homocysteine [12, 14]. Interestingly, instead of cysteine, host-generated hydrogen sulfide (H_2S) acts as the S-donor in this reaction catalyzed by the enzyme O-succinyl homoserine sulfhydrylase (OSHS). The direct sulfhydrylation step allows several microbial species to bypass the two-step conversion of OSH to homocysteine via cystathionine, a crucial intermediate in the canonical methionine biosynthesis pathway [9, 15-17]. The presence of a putative OSHS encoded by *metZ* (*Rv0391*) in *M. tuberculosis* suggests the existence of a yet uncharacterized active direct sulfhydrylation pathway in *M. tuberculosis* [18].

In this study, employing genetic and computational approaches, we investigate the necessity of redundant methionine biosynthesis pathways in *M. tuberculosis*. We demonstrate that *metZ* gene is required for methionine biosynthesis under acidic conditions. Using computational approach, we further demonstrate that the binding of the substrate to the active site of the enzyme MetZ is more stable at acidic pH. Using CRISPR-mediated gene silencing, we provide evidence that *metZ*-dependent direct sulfhydrylation pathway in *M tuberculosis* functions independently of the transsulfuration pathway. The failure of *M. tuberculosis* strain, lacking both the transsulfuration and direct sulfhydrylation pathways, to grow in methionine-lacking medium further validates the redundancy of these pathways and suggests that the *metZ*-mediated direct sulfhydrylation pathway becomes conditionally essential for the pathogen to survive under acidic and *in-vivo* growth conditions. In

conclusion, we report that *metZ*-mediated direct sulfhydrylation pathway is the primary driver of the

methionine biosynthesis under acidic conditions and is therefore essential for the survival of *M.*

tuberculosis within the host.

Results

MetZ is dispensable for the *in-vitro* growth of *M. tuberculosis* and possibly involved in direct

sulfhydrylation pathway for methionine biosynthesis

While *metZ* has been reported to be essential for the growth of *M. tuberculosis* on cholesterol [19], we

hypothesized that the *Rv0391 (metZ)* gene encodes a putative O-succinylhomoserine sulfhydrylase

(OSHS) that is responsible for direct sulfhydrylation (Fig. 1A). We performed a bioinformatics-based

analysis using the protein sequence of MetZ$_{TB}$ and found that the putative sequence belonged to the

cluster of orthologues group ENOG502GJ5S which is composed of 1097 protein sequences

orthologous to MetZ$_{TB}$ across 644 bacterial species. This cluster of proteins comprises various enzymes

involved in cellular amino acid metabolism namely, cysteine, methionine, homocysteine, serine and

aspartate. Co-incidently, all these proteins have a common PLP (pyridoxal-5-phosphate) – binding

domain which is essentially a modified lysine residue that covalently binds the co-factor, hence also

categorized as PLP-binding superfamily of proteins (Fig. 1B and S1A). Further, to elucidate the role

of *metZ* in *M. tuberculosis*, a deletion mutant in the genetic background of H37Rv (*Rv-WT*) was

generated by genetic recombineering as described previously [20]. Briefly, a hygromycin resistance

cassette was inserted into the ORF encoding *metZ* (*Rv0391*) with the help of allelic exchange substrate

(AES) consisting of 1kb regions of homology upstream and downstream to the target locus. The

recombinants were screened for hygromycin resistance and the cassette was subsequently excised out

thereby generating an unmarked mutant (Fig. 1C). The loss of the genetic sequence between 470034-

471188 bases was confirmed by gene-specific PCR (Fig. S2A), DNA sequencing of the locus (Fig.

S2B) and at the transcript level by RT-qPCR (Fig 1D). To check the essentiality of the *metZ* gene in

growth on cholesterol, we compared the growth of *Rv-WT*, *ΔmetZ* and the complement luminescence reporter strains in different media containing glycerol (0.1%), cholesterol (200μM), propionate (10mM) as the sole carbon source. As a control, we used the *M. tuberculosis* strain lacking the *prpR* gene (*ΔprpR)* which is known to be essential for growth on cholesterol [21]. While, as expected, *ΔprpR* failed to grow on cholesterol and propionate*,* the growth of all the three strains were comparable irrespective to the carbon source (Figs. 1E, 1F and S3B). These findings indicate *metZ* does not contribute to the central carbon metabolism and is dispensable for *in-vitro* growth of *M. tuberculosis* on cholesterol and/ or propionate.

metZ deletion renders *M. tuberculosis* sensitive to physiological stressors and leads to reduced *ex-vivo* fitness

To replicate and grow inside the host, intracellular pathogens have evolved mechanisms to neutralize the host-induced physiological stressors imposed by the host environment, including oxidative stress, insult to the cell membrane, acidic pH and nutrient limitation. To study the role of *metZ* in mitigating these host-induced stress, the *M. tuberculosis* strains were exposed to a range of stress-inducing reagents. The effect of these stresses on each strain was subsequently assessed by calculating the survival index relative to the *Rv-WT* strain. In comparison to the *Rv-WT* and complement strains, *ΔmetZ,* demonstrated a reduced ability (~40-60%) to survive under all the stress conditions tested (Fig. 2A). Further, we determined intra-bacterial redox state using respective *M. tuberculosis* strains expressing genetically encoded redox biosensor (Mrx1-roGFP2). To establish the dynamic range of the biosensor readout, *Rv-WT* and the *ΔmetZ* strains expressing Mrx-roGFP2 were treated with H_2O_2 and Dithiothreitol (DTT) to achieve maximum oxidation and reduction states, respectively. This defined the upper and lower limits of the biosensor response (dotted lines in Fig. 2B). We demonstrate that in comparison to *Rv-WT*, the *ΔmetZ* strain exhibited significantly higher intracellular ROS levels

(Fig. 2B) suggesting an inability of the mutant to maintain redox homeostasis possibly due to impaired

mycothiol recycling. Notably, unlike *Rv-WT*, we did not observe any additional increase in

fluorescence signal in H_2O_2 treated *ΔmetZ* strain, further supporting the presence of elevated ROS

levels that may have saturated the biosensor (Fig. 2B). Subsequently, increased levels of lipid

hydroperoxides were detected in the *ΔmetZ,* strain compared to the *Rv-WT* and complement strains,

further corroborating our previous finding that the absence of *metZ* gene in *M. tuberculosis* results in

excessive accumulation of intracellular reactive oxygen species (Fig. 2C). Further, to assess the role of

metZ in the intracellular survival of the *M. tuberculosis* inside the macrophages, resting or IFN-γ

induced THP1-derived macrophages were infected with *Rv-WT*, *ΔmetZ* and the complement strains

and survival indices (w.r.t. *Rv-WT*) were determined by enumerating the bacterial load by CFU (Colony

Forming Unit) plating of the infected cell lysates at day 5 post-infection. We observed a significant

reduction in bacterial burden in the *ΔmetZ* strain compared to both *Rv-WT* and complement strains in

both resting (~60%) and activated (~48%) macrophages (Fig. 2D and S4A). Since restricting the

phagosome-lysosome fusion is a common strategy employed by the pathogen to enhance its

intracellular survival, we further investigated the role of *metZ* in regulating this process. Briefly, IFN-

γ induced THP1 macrophages infected with mEmerald (GFP) expressing *Rv-WT*, *ΔmetZ* and

complement strains were subsequently stained with Lysotracker. The study revealed that in comparison

to the *Rv-WT* and complement strains, the *ΔmetZ* strain exhibited a higher propensity to localize within

the acidic compartments of the cell (Fig. 2E and S5A). Overall, these data suggest that absence of *metZ*

significantly weakens the ability of *M. tuberculosis* to resist host-induced physiological stresses and

evade phago-lysosome fusion, both of which are crucial for the survival of the pathogen within the host

cell.

Direct sulfhydrylation pathway is essential for *M. tuberculosis* to survive under low pH

The observed data suggest that the *ΔmetZ* strain might exhibit increased sensitivity on exposure to an

acidic environment. To investigate this, we cultured different bacterial strains (*Rv-WT*, *ΔmetZ* and

complement) at varying pH values and monitored their growth kinetics. While the growth kinetics were

comparable at neutral pH (7.2) for all the strains, a decrease in pH resulted in a discernible growth

defect in the *ΔmetZ* strain, ranging from observable impairment at pH 5.5 to complete growth arrest at

pH 5.0, relative to the *Rv-WT* and complement strain (Fig. 3A and S6A and B). To further understand

the impact of acidic pH on the expression of genes involved in transsulfuration (*metB* and *metC*) and

direct sulfhydrylation (*metZ*), we quantified the relative expression of these genes in *M. tuberculosis*

cultured at pH 7.2 and pH 5.5. RT-qPCR analysis revealed an ~12-fold increase in the expression of

metZ gene at pH 5.5 compared to pH 7.2, suggesting transcriptional induction of the *metZ* gene under

acidic conditions (Fig. 3B). To differentiate between bacterial killing and metabolic slowdown as the

cause of reduced CFU counts in the *ΔmetZ* strain at pH 5.5 (Fig. S6C), we assessed metabolic activity

using Resazurin dye [22]. We found that, in contrast to the *Rv-WT* and complement strains, the *ΔmetZ*

strain exhibited significantly reduced metabolic activity at pH 5.5, as indicated by lack of color change

and low fluorescence, suggesting a significant impairment in metabolic function (Fig. 3C).

Surprisingly, despite this metabolic impairment, ATP levels in the *ΔmetZ* strain were found to be higher

than those in the *Rv-WT* and complement strains when cultured at acidic pH 5.5 (Fig. 3D).

Given that the first-line anti-TB drug - Pyrazinamide (PZA) is active only at acidic pH and efficiently

kills slow or non-replicating cells, we hypothesized that the differences in the cytoplasmic pH will

correlate with PZA sensitivity. As expected, we observed a ~19-fold and 4-fold increase in the

sensitivity of the *ΔmetZ* strain to PZA treatment compared to *Rv-WT* and complement strain,

respectively (Fig. 3E). To confirm, if the observed phenotype is a direct consequence of a defect in the

methionine biosynthesis, we quantified the intracellular methionine levels. We found significantly

lower levels of methionine in both *Rv-WT* and *ΔmetZ* strains grown in acidic pH 5.5 compared to those

grown at pH 7.2 (Fig. 3F). Exogenous supplementation of methionine rescued the growth defect

phenotype of *ΔmetZ* observed at pH 5.5 (Fig. 3G). Importantly, the addition of methionine did not alter

the pH of the culture media (Fig. S6D and E). Furthermore, we observed a reduction in the intracellular

ROS levels in *ΔmetZ* upon exogenous supply of methionine at pH 7.2 while this effect was less

pronounced at pH 5.5 (Fig 3H). Concurrently, methionine supplementation not only restored ATP

levels but also increased the IC50 of PZA in the *ΔmetZ* at pH 5.5 (Fig 3I and S6F). *M. tuberculosis*

produced significantly higher levels of hydrogen sulfide (H_2S) at acidic pH compared to the neutral pH

(Fig S6G). H_2S serves as a co-substrate and S-donor in the reaction catalyzed by OSHS enzyme (MetZ),

an enzyme belonging to the PLP-dependent superfamily of proteins which require PLP as co-factor

(Fig 1B and S1). To assess the role of *metZ* in PLP homeostasis, we examined the growth kinetics of

different mycobacterial strains in minimal medium supplemented with or without PLP at both pH 7.2

and pH 5.5. We found that knocking out *metZ* had no apparent consequences on the growth kinetics of

the *M. tuberculosis* with or without PLP at either pH conditions (Fig. S7A and B). However, we did

observe a low pH-based growth defect in *ΔmetZ* even in minimal medium (Fig. S7B). In summary, a

defect in the methionine biosynthesis pathway at low pH renders the *ΔmetZ* strain metabolically less

active, thereby inhibiting its growth at acidic pH.

Molecular docking and Molecular dynamics (MD) simulations revealed enhanced stability of MetZ at acidic pH

To investigate the molecular mechanisms underlying MetZ substrate binding and their kinetics at

physiologically relevant pH (7.0 and 5.4), we employed computational approaches, including

molecular docking and molecular dynamics simulations studies. To perform *in-silico* molecular

docking with MetZ protein (Fig. 4A), we determined the binding site of O-succinyl homoserine (OSH)

by focused molecular docking [23, 24] that provided the final pose at the cofactor bound site. A pose

with docking energy of -4.07 Kcal/mol and cluster size of 50 (out of 200) was found suitable at pH 7.0.

While, at pH 5.5 the most likely pose showcased a docking energy of -4.19 Kcal/mol with a cluster

size of 80 (200). For both these poses at diverse pH, a static thermodynamic quantification was carried

out using MM-GBSA which was found to be -49.20 Kcal/mol and -56.93 Kcal/mol, respectively.

Molecular Dynamics Simulation at a constant pH was performed for these poses. Following the 300ns

MD simulations, the protein exhibited stable RMSD values at 2.98 Å and 4.34 Å at pH 7.0 and 5.5,

respectively. However, significant changes were observed in the stability of the substrate as well as the

occupied binding site. On observing the substrate in consensus with its binding site, the region was

found to have higher stability and an average RMSD of 1.87 Å was observed at pH 5.5 in comparison

to an average of 2.71 Å at pH 7.0 (as shown in Fig. 4B and D). The underlying reason for this higher

stability at pH 5.5 is due to the establishment of stable durable bonds in the form of hydrogen bonds

with residues V196, F197 and T319 and the O-atoms at the terminals of the substrate. Furthermore,

another factor which enhances the strength is the formation of a salt-bridge with residue H220 and

substrate. However, a similar configuration of H-bonds and salt-bridge are not observed at pH 7.0,

possibly due to conformational dynamics [25] (Fig. 4E). These characteristics are a result of the change

in pKa values as well as conformations of the binding site forming amino acid residues. Protonation

states of amino acids are highly susceptible to change in environmental pH and thus, preparing the

protein at pH 7.0 and 5.5 generates different acid dissociation constants, which ultimately affects the

bond forming tendencies of residues. The residue E165 attains a pKa of 8.78 at pH 7.0 and donates a

proton, forming a bond with the substrate. Similarly, D194 attains a pKa of 9.11 at pH 5.5 which affects

its binding to the substrate. Also, H220 is present in a protonated state at a lower pH and thus involved

in the formation of a Salt-Bridge with the substrate. While at pH 7.0, H220 remains in a deprotonated

state and thus does not contribute to the stability of the binding pocket at neutral pH. This showcases

the importance of the acidic pH which contributes to relatively greater stability of the substrate. On

analysis of the binding site, we found that H220 and T319 interact with the substrate at the acidic pH

but they showcase conformational changes at pH 7.0 which hinders their interaction. The imidazole

ring sidechain of H220 rotates in anti-clockwise direction and thus does not get involved in bond

formation. Similarly, T319 forms H-bonds at both pH but the nature of the bonds varies from transient

at pH 7.0 to largely consistent at pH 5.5. Overall, conformational changes in the binding site residues

can be accredited for better stability at pH 5.5 The stability of the protein and the substrate at acidic

pH hints towards a better binding activity of the substrate. This characteristic feature can be examined

by performing the MM-GBSA of a few selected frames from both the trajectories to compare their

activity. The average binding/biological activity of the protein is found to be -37.35 Kcal/mol and -

59.40 Kcal/mol at pH 7.0 and 5.5, respectively (Fig. 4C). This increase in the binding affinity is due to

the effect of acidic pH on the protonation state as well as the induced conformational changes. The

docking scores at both pH 7.0 & 5.5 suggest that the residence time of the substrate at the binding site

must be short and the sulfur transfer reaction might be transient. Overall, focused docking and MD

simulations revealed that although protein is stable at both acidic and neutral pH, the active site of the

protein holds substrate (OSH) more stably at acidic pH rather than at neutral pH which sheds light on

its physiological role in methionine biosynthesis at acidic pH instead of neutral pH in concordance

with our *in-vitro* studies.

metZ-mediated direct sulfhydrylation pathway is critical for methionine biosynthesis at acidic

pH

To investigate the independence of the *metZ*-dependent methionine biosynthesis pathway from the

transsulfuration pathway in *M. tuberculosis*, we employed CRISPRi to knockdown *metB* (*Rv1079*)

expression in both *Rv-WT* (*Rv::metB_{KD}*) and *ΔmetZ (ΔZ::metB_{KD})* strains (Fig. 5A).

Anhydrotetracycline (ATc)-induced knockdown of *metB* expression in both strains was confirmed by

RT-qPCR analysis (Fig. S8A). While ATc-induced knockdown of *metB* had no significant effect on

the growth of the *Rv-WT* strain, it resulted in a growth defect in the *ΔZ::metB*$_{KD}$ strain, effectively

mimicking a double knockout phenotype (Fig. 5B and S8B). This growth defect is likely attributed to

a compromised methionine biosynthesis pathway, as evidenced by a ~1.5-fold decrease in the

intracellular methionine levels in the *ΔZ::metB*$_{KD}$ strain compared to the *Rv::metB*$_{KD}$ strain (Fig. 5C).

Based on these findings we hypothesized that the *ΔZ::metB*$_{KD}$ double mutant would exhibit attenuated

growth in an animal model due to reduced methionine availability. To test this hypothesis, mice were

infected with *Rv::NT* (Non-targeting control sgRNA) and *ΔZ::metB*$_{KD}$ strains, and doxycycline was

administered to the specific groups to induce CRISPRi-mediated knockdown of the respective gene.

Bacterial load was subsequently estimated by CFU plating (Fig 5D). We found that compared to NT

and *ΔmetZ*, the *ΔZ::metB*$_{KD}$ double mutant exhibited reduced bacterial burden during the initial stages

of infection in both lungs and spleen which was partially rescued by week 6 post infection (Fig. 5E).

We hypothesized that the presence of an active methionine transporter, such as MetM (Rv3253c) [26]

might contribute to the partial rescue of the growth defect observed in the animal infected with

ΔZ::metB$_{KD}$ double mutant strain of *M tuberculosis*. Collectively, the above data demonstrate the

functional independence of the MetZ-mediated direct sulfhydrylation pathway in *M. tuberculosis*. The

observed growth defects and reduced methionine levels in the *ΔZ::metB*$_{KD}$ strain both *in-vitro* and *in-*

vivo models highlight the critical role of these redundant pathways in supporting bacterial growth and

survival inside the host.

Direct sulfhydrylation pathway of *M. tuberculosis* contributes towards antibiotic persistence in

response to Bedaquiline

Next, we wanted to study the role of direct sulfhydrylation pathway in regulating the drug susceptibility

profile of *M. tuberculosis*. We first compared the Minimum Inhibitory Concentration (MIC) of various

anti-TB drugs. We found that the IC50 values for each of these drugs were similar for all three strains

(*Rv-WT*, *ΔmetZ* and the complement strain) (Fig. 6A and 9A-D, F). Further, a time-kill assay was

performed to determine the role of *metZ* in modulating the formation of drug-induced persisters in *M.*

tuberculosis. The frequency of persisters formation was assessed by comparing the biphasic kill curves

generated from the time-kill assays using various antibiotics (RIF, INH, EMB and BDQ) at 10X MIC

concentrations. No significant differences in the persisters formation were observed among the strains

when treated with RIF, INH or EMB (Fig. 6B). Interestingly, compared to the *Rv-WT* and the

complement strains, a significant decrease in the persisters formation was observed in the *ΔmetZ* upon

treatment with BDQ at both neutral (Fig. 6B) and acidic pH (Fig. S9E). While a biphasic kill curve

was observed for the *ΔmetZ* treated with BDQ at pH 7.2, the kill curve at pH 5.5 was predominantly

linear, suggesting a lower frequency of persisters formation at acidic pH. Given BDQ inhibits ATP

synthesis, we hypothesized that the observed decrease in the persisters formation in the *ΔmetZ* might

be due to impaired generation of ATP. However, ATP depletion kinetics analysis using 2X MIC BDQ

revealed higher ATP levels in the *ΔmetZ* compared to the *Rv-WT* and the complement strains (Fig. 6C),

suggesting that BDQ-mediated ATP depletion was slower in the *ΔmetZ* strain. In sum, we found that

the *metZ*-mediated direct sulfhydrylation pathway modulates the susceptibility of *M. tuberculosis* to

BDQ and that the reduction in the BDQ-dependent persisters generation observed in the *M.*

tuberculosis strain lacking *metZ* is independent of the ATP levels present inside the bacteria.

Direct sulfhydrylation pathway supports the growth of *M. tuberculosis* inside the host

To elucidate the role of *metZ* in mycobacterial pathogenesis, a guinea pig model of tuberculosis

infection was employed. Briefly, guinea pigs were infected with the respective mycobacterial strains

(*Rv-WT*, *ΔmetZ* and complement) via the aerosol route. Bacterial burden in the lungs and the spleen

was determined by CFU plating at different time points post-infection. Gross and histopathological

analyses were performed to study the damage due to infection at the tissue level (Fig. 7A). We found

that at week 4 post-infection the bacterial burden in the lungs of guinea pigs infected with *ΔmetZ* was

significantly lower than that observed in animal infected with the *Rv-WT* and complement strains,

exhibiting a ~3.7-fold and 2-fold reduction, respectively. By 8 weeks post-infection, a more

pronounced difference was observed, with approximately an 8.9-fold reduction in the bacterial burden

in the lungs of *ΔmetZ*-infected animals compared to those infected with the *Rv-WT* strain. Notably,

only a modest increase in CFU counts was observed in the *ΔmetZ*-infected animals between weeks 4

and 8 (Fig. 7B). Similarly, a 6-fold reduction in bacterial burden was observed in the spleens of animals

infected with the *ΔmetZ* compared to those infected with the WT and complement strains at week 4

post-infection. By week 8, bacterial burden in the spleens of the *ΔmetZ*-infected animals was

significantly reduced, approaching the limit of detection by CFU and exhibiting a nearly 263-fold

decrease compared to week 4 post-infection (Fig. 7C). Consistent with the reduced bacterial burden

(Fig. 7B), gross pathology revealed fewer granulomatous lesions in the lungs of animal infected with

the *ΔmetZ* compared to those infected with the *Rv-WT* and complement strains at 8 weeks post-infection

(Fig. 7D). To visualize and quantify the damage to lung tissues, Hematoxylin & Eosin (H&E) staining

of the formalin preserved lung sections was done. Microscopic examination of lung samples collected

from animals infected with *Rv-WT* strain revealed severe granulomatous inflammation throughout the

lung parenchyma. Numerous well-defined necrotizing granulomas consisting of epithelioid histiocytes,

perimetric cuffing of lymphocytes and plasma cell aggregates and a caseous necrotic mass at the centre

were observed. However, lung samples from animals infected with complement strain and *ΔmetZ* strain

showed only non-necrotizing granulomas along with multiple foci of inflammatory cell aggregates and

occasional alveolar hemorrhagic patches (Fig. 7E). Granuloma scoring further confirmed these

observations, with the highest scores observed in lungs sections from animals infected with the *Rv-WT*

strain, followed by those infected with the complement and *ΔmetZ* strains (Fig. 7F). Overall, these data

suggests that absence of *metZ* gene significantly impairs the ability of *M. tuberculosis* to replicate and

survive within the host, suggesting that the *metZ*-dependent direct sulfhydrylation pathway is crucial

for the survival of the pathogen inside the host.

metZ is involved in methionine biosynthesis in *M. tuberculosis* at acidic pH

We hereby summarize that upon infecting macrophages, *M. tuberculosis* traverses a series of

intracellular compartments, initially residing in the phagosomes where a relatively neutral pH (~6.5)

and abundant nutrients, including methionine that support bacterial growth. However, as host immunity

mounts, IFN-γ-mediated induction of reactive oxygen species (ROS) and reactive nitrogen

intermediates (RNI) triggers phagosomal maturation leading to a severe drop in the pH (5.5 or lower).

These acidic conditions, coupled with nutrient restrictions imposed by the host, create a hostile

environment that inhibits the growth of the pathogen. In response to these stressors, *M. tuberculosis*

undergoes substantial transcriptional and metabolic reprogramming. One critical adaptation involves

the induction of *metZ* expression, which enables the pathogen to survive these challenging conditions.

The MetZ-mediated direct sulfhydrylation pathway utilizes host-derived H_2S as a co-substrate and S-

donor for methionine biosynthesis within this acidic microenvironment, facilitating the maintenance

of ATP levels, mycothiol recycling and overall metabolic activity (Fig. 8).

Discussion

M. tuberculosis, adapts to various hostile conditions within host macrophages resulting in the

differential expression of proteins crucial for its survival. One such key adaptation involves increase in

the biosynthesis of cysteine which compensates for the increase in the flux of cysteine towards

generating the thiol-active metabolites (mycothiol, ergothioneine, NAD+, etc.). These metabolites are

essential for maintaining the redox balance upon exposure to heightened levels of ROS, RNI [27] and

acidic pH [28]. The same has been reported in *E. coli*, where increased oxidative stress leads to higher

demand for the cytosolic glutathione which stimulates cysteine biosynthesis [29]. Under such

demanding circumstances, the pathogen needs to evolve and devise a mechanism to efficiently regulate

the flux of cysteine towards either biosynthesis of methionine and/or mycothiol which would maintain

the total thiol pool thereby maintaining redox homeostasis. Likewise, given the crucial role of

methionine in bacterial physiology, there has been a gap in our understanding of the rationale of

harbouring multiple methionine biosynthesis pathways and their role in maintaining methionine

homeostasis critical for the survival of *M. tuberculosis* in a nutrient-deprived *in-vivo* growth

environment. We, in the current study, have used a combination of genetics and computational biology-

based tools to probe the role of the *metZ*-mediated direct sulfhydrylation pathway in *M. tuberculosis*.

We found that MetZ-dependent direct sulfhydrylation is significantly induced at acidic pH. Disrupting

this pathway in *M. tuberculosis* results in a decrease, but not complete elimination, of cellular

methionine at low pH. Finally, we demonstrated that MetZ-dependent direct sulfhydrylation is required

for growth of *M. tuberculosis* inside the host.

In contrast to an earlier report [19], *metZ* was found to be dispensable for the growth of *M. tuberculosis*

on cholesterol. The conditional essentiality phenotype observed for *ΔmetZ* under low pH further

suggests that the earlier observation of the *metZ* essentiality on cholesterol likely resulted from the low

pH growth conditions used in that study. High demand for cysteine, a common precursor for the

generation of both methionine and mycothiol, during stress resulted in observed imbalance in the redox

homeostasis in *ΔmetZ*. Exogenous supplementation of methionine alleviates the stress by reducing the

need to divert the intracellular cysteine pool towards methionine production, thereby enhancing the

mycothiol generation critical for mitigating the ROS levels at acidic pH. Beyond redox homeostasis,

our data underscores the role of methionine biosynthesis, through the direct sulfhydrylation pathway,

in regulating the cytosolic pH and metabolism. The inability of the *ΔmetZ* to maintain the intracellular

pH homeostasis impedes the ability of the pathogen to utilize the cellular ATP, leading to a metabolic

slowdown. The elevated levels of ATP observed in *ΔmetZ* at acidic pH, is likely, a consequence of

reduced ATP utilization rather than overproduction as reported in an earlier study using *E. coli* [30].

The enhanced susceptibility of *ΔmetZ* to Pyrazinamide (PZA), an anti-TB prodrug that gets activated

upon exposure to acidic pH, further confirms the presence of a relatively acidic cytosol and accentuates

the role of *metZ* in maintaining the pH homeostasis in *M. tuberculosis*. In an *ex-vivo* setting, IFN-γ-

induced activation leads to acidification of the pathogen containing phagosomes in macrophages.

Consequently, *ΔmetZ* replication inside both resting and the activated macrophages were reduced,

likely due to increased co-localization of *ΔmetZ* to acidic compartments as revealed by confocal

microscopy. This finding aligns with previous studies suggesting that enhanced susceptibility to

oxidative stress results in increased bacterial killing in the acidified macrophage compartments [31-

33].

Our data provides a mechanistic understanding of the methionine starvation-associated susceptibility

in *M. tuberculosis* in line with reports published by other groups [4, 10]. The presence of both

transsulfuration and direct sulfhydrylation pathways for the biosynthesis of methionine in *M*

tuberculosis is indeed intriguing. Although both the pathways demonstrate redundancy at normal pH,

our qPCR data, along with previous studies, categorically indicate that enzyme responsible for carrying

out direct sulfhydrylation is over-expressed in *M tuberculosis* upon induction with acidic pH [32, 34,

35]. These findings are corroborating well with our pH dependent molecular dynamics (MD)

simulation studies which suggested that the binding affinity between the substrate and MetZ active site

is stronger at low pH. The MD simulation data provide structural insights into how pH and the

protonation states of key residues alter the hydrogen bonding and electrostatic interaction network

leading to significant changes in the conformation of the MetZ protein. The change in the hydrogen

bonding network, identified as a determinant, majorly impacts the change in the binding affinity of the

substrate. Furthermore, a comparative analysis of MetZ structural stability at different pH values, using

metrics such as root-mean-square deviation (RMSD) and thermodynamic changes, confirms the pH-

dependent behaviour of MetZ. The observed changes in the hydrogen bonding network significantly

impact the substrate binding affinity of the MetZ, indicating that, under low pH conditions, *metZ*-

mediated direct sulfhydrylation majorly contributes towards the biosynthesis of methionine in *M.*

tuberculosis. We further confirm that *M. tuberculosis* strain that lacks both transsulfuration and direct

sulfhydrylation pathways (*ΔmetZ/metB$_{KD}$*) fails to generate methionine, demonstrating methionine

auxotrophy under *in-vitro* conditions. This finding confirms the redundancy that exists between these

two pathways. Surprisingly, the growth defect observed in the *ΔZ::metB$_{KD}$* under *in-vivo* conditions,

while significant, was less pronounced than that observed *in-vitro*. This may be attributed to the

presence of a potential methionine transporter, encoded by Rv3253c (MetM), which allows the

pathogen to salvage methionine from the host environment [26]. Similar to our finding (Fig. S6G), it

has been reported previously that *M. tuberculosis* can produce H_2S, a gaseous signalling molecule,

upon induction with acidic pH [13] probably as a mechanism to counteract increasing ROS and

dropping metabolic potential of the cells. Further, it has also been shown that H_2S at sub-inhibitory

levels in *M. tuberculosis* have a positive effect on the mycobacterial electron transport chain (ETC)

and this effect is reversible [36, 37]. Additionally, host-generated H_2S has been shown to enhance *M.*

tuberculosis replication inside the host. This effect is evident in studies demonstrating impaired *M.*

tuberculosis replication when the hosts' ability to generate H_2S is compromised, either through genetic

knockout of the genes encoding the enzyme cystathionine-γ-lyase (CSE) and cystathionine-β-synthase

(CBS) or by using pharmacological inhibitors of these enzymes [36, 38, 39]. Co-incidentally, it is well

established in the literature that sulfhydrylases, including MetZ, utilize H_2S as a S-donor for various

reactions, such as cysteine biosynthesis or, in this case of MetZ, for the production of homocysteine.

The development of antibiotic persistence in bacteria is a significant contributor to the failure of anti-

TB treatment. The frequency of generation of the persisters population can be quantified through a

time-kill assay using high antibiotic concentration (5-10X MIC) [40]. Our time-kill experiments

suggest that the absence of a direct sulfhydrylation pathway in *ΔmetZ* prevents the BDQ-specific

generation of antibiotic persistence in *M. tuberculosis*. Intriguingly, despite exhibiting higher cytosolic

ATP levels compared to the wild-type strains, *ΔmetZ* demonstrated a decrease in the BDQ-mediated

persisters formation. This observation contrast with a previous study on *ΔpckA* mutant, wherein

elevated cytosolic ATP levels increased drug tolerance when grown in acetate medium [41]. In

summary, our findings suggest that beyond its role in maintaining methionine levels at acidic pH, the

direct sulfhydrylation pathway also modulates the generation of BDQ-specific generation of antibiotic

persistence during infection. Further investigation into this complex phenotype will be crucial for

developing new strategies to prevent the emergence of BDQ resistance in tuberculosis. Furthermore,

we found that *metZ* is essential for the maintenance of *M. tuberculosis* infection in guinea pig model,

as evident by a reduced bacterial burden and severe attenuation in the spleen. Spleen is known to be

iron rich, which can contribute to the generation of reactive oxygen species via Fenton chemistry.

Given the observed susceptibility of the *ΔmetZ* to oxidative stress, the low bacterial count in the spleen

at week 8 post-infection likely reflects the inability of the mutant to survive in this nutrient-deprived,

low pH and oxidative stress-inducing environment within the host. Our results indicate that *M.*

tuberculosis adapts to the decreasing pH in its surroundings by maintaining methionine biosynthesis

through direct sulfhydrylation utilizing exogenous and endogenous H_2S as S-donor.

This study provides a foundation for designing novel anti-tuberculosis treatment strategies. Using

pharmacological inhibitors of MetZ as an adjunct to the existing anti-tuberculosis regimens could offer

a promising approach. This synergistic strategy can potentially enhance the efficacy of the existing

treatment regimen against tuberculosis.

Materials and Methods

Bacterial strains

The knockout strain of *metZ* gene *(Rv0391)* was generated in the genetic background of wild type

M. tuberculosis H37Rv by genetic recombineering. Briefly, a gene-specific allelic exchange substrate

(AES) harboring 1kb sequences of target gene upstream and downstream homology flanking

Hygromycin resistance gene cassette was constructed in the suicidal vector pJM1. AES was

transformed into H37Rv (wild-type, *Rv*-WT) strain overexpressing Phage Che9c RecET proteins to

facilitate recombination. Recombinants with the Hygromycin resistance (Hyg^r) gene were selected

over Hygromycin (Hyg) plates and screened using gene-specific PCR. The marked mutant thus

prepared was unmarked to excise out Hygromycin resistance cassette using Cre-recombinase

expressed from plasmid pCre-SacB-Zeo. The unmarked mutant was confirmed by PCR and RT-

qPCR to confirm the loss of gene as well as transcripts of *metZ* gene. The unmarked mutant was then

transformed with a plasmid expressing a functional copy of the *metZ* gene under P$_{myc1}tetO$ promoter

replicating either episomally or integrated to generate genetic complementation strain (denoted as

ΔmetZ Comp). Bioluminescent mycobacterial strains were generated by transforming respective

mycobacterial strains (**Table S3, List of strains used in the study**) with integrative plasmid

expressing *luxABCDE* operon (Addgene# 26161) [42] with slight modifications [43]. Strains were

checked for bioluminescence and a luminescence vs CFU plot was prepared with *Rv-WT*

luminescence reporter strain to determine the relation between luminescence and CFU-based read-

outs of bacterial viability (Fig. S3A). Henceforth, where ever appropriate, luminescence was used as

proxy for the bacterial viability and growth.

Media and cultural conditions

The bacterial strains were maintained as frozen glycerol stocks at -80°C. Whenever required, the stocks were thawed and diluted 1:500 in 7H9 (Difco, BBL) medium supplemented with Oleic acid, Albumin, Dextrose, Salt (NaCl) (OADS) and 0.01% Tween 80. Bacterial enumeration was done on 7H11 (Difco BBL) agar plates supplemented with OADS. Kanamycin (Sigma), 25µg/ml, Hygromycin (Sigma), 50µg/ml and Zeocin (Invitrogen), 25µg/ml were used whenever required. For media with no carbon source, 7H9 base medium was prepared without glycerol and OADS (or Albumin, Dextrose, Salt: ADS) enrichment and later supplemented with desired Carbon (C)-sources and 0.05% tyloxapol. 100mM stock of Cholesterol (Sigma) was prepared by heating cholesterol in Tyloxapol (Sigma): Ethanol (Merck) mix (1:4) at 80°C for 30 mins [44]. When required, cholesterol was diluted to desired concentration (200µM) in 7H9 base medium wherever specified. Likewise, glycerol (Sigma) was used at a final concentration 0.1% v/v and sodium propionate (Sigma) was used at a concentration of 10mM (0.1%) for growth kinetics studies. For pH-dependent growth kinetics experiments, 7H9 base medium was prepared, supplemented with 0.1% glycerol (v/v) and ADS acidified to desired pH using HCl. For Pyridoxal-5'-Phosphate (PLP) supplementation-based growth kinetics, minimal medium (0.5g/litre asparagine, 1g/litre KH_2PO_4, 2.5g/litre Na_2HPO_4, 50mg/litre ferric ammonium citrate, 0.5g/litre $MgSO_4.7H_2O$, 0.5mg/litre $CaCl_2$, and 0.1mg/litre $ZnSO_4$) containing 0.1% glycerol (vol/vol) was used with or without 50µg/ml PLP (Sigma) supplementation. Phosphate buffered saline (PBS) was used at 1X concentration supplemented with tyloxapol (0.05%), PBST whenever mentioned. Bacterial enumeration by CFU was done on 7H11 agar plates supplemented with OADS, wherever mentioned. Colonies were counted after 21 days of incubation. Anhydrotetracycline, ATc (Caymen) was used at 200-300ng/mL concentration unless mentioned otherwise. L-methionine (Sigma) was supplemented at 100µg/ml concentration wherever required.

CRISPRi knockdown

For transcriptional knockdown of *metB (Rv1079)* using CRISPR interference (CRISPRi), target-specific sgRNA was cloned into *BsmBI* (NEB) digested backbone plJR965 (Addgene # 115163) to generate knockdown construct. The knockdown construct was transformed into *Rv-WT* and *ΔmetZ* strains through electroporation. To achieve knockdown of the target genes, cells were grown to mid-log phase and diluted back to OD 0.01 in fresh culture medium with or without ATc. Growth was monitored by measuring OD of the cultures. Further, expression of target genes was quantified by qPCR to determine the levels of knockdown.

RT-qPCR

Total RNA was isolated from the respective bacterial strains under specific conditions using Trizol (TaKaRa) according to the manufacturer's protocol. Briefly, bacterial cells were washed with 1X PBS and resuspended in Trizol. Cells were lysed by bead-beating using 0.22μM sized zirconia beads via 5 cycles with intermittent cooling step of 1 minute. Further, RNA was extracted and precipitated using chloroform-isopropanol treatments. Total RNA thus prepared was treated with Turbo DNaseI (Ambion) to digest away the DNA contamination. 3000 μg equivalent DNase treated RNA was used to prepare cDNA using SuperScript IV Reverse Transcriptase (Invitrogen). Finally, 1μg equivalent cDNA was used to set up qPCR reactions with SYBR (TaKaRa). *sigA* was used as internal control and $\Delta\Delta C_t$ method [45] was used determine relative gene expressions w.r.t. specified control conditions. For qPCR at acidic pH, *icl1* was used as positive control.

Growth kinetics

Auto-bioluminesc ent bacterial strains were grown till OD 0.5 (mid-log phase) and washed once

with 1X PBST to remove the spent media. The cells were resuspended in 1X PBST and diluted back

to an OD of 0.01 in 7H9 base medium supplemented with different carbon sources – glycerol

(0.1%), cholesterol (200µM) or propionate (10mM) for C-source dependent growth kinetics. For

pH-dependent growth kinetics, bacterial cells were diluted back to an initial OD of 0.05 in 7H9

medium (supplemented with ADS and 0.2% glycerol) acidified to different pH – 7.2, 6.5, 5.5 and

5.0. Whenever required, L-methionine (100µg/ml, Sigma) was added to the cultures and 50µg/ml

L-methionine was replenished every 4 days until the conclusion of the experiment. For CRISPRi

based growth kinetics, Anhydrotetracycline chloride (ATc) was added to the cultures (200ng/ml)

and replenished every 4 days to maintain the suppression of the target gene until the conclusion of

experiment. The growth was monitored by measuring OD and/ or auto-bioluminescence using a

Multimode Microplate reader (SpectraMax Pro ID3, Molecular Devices). At times, bacterial growth

was also determined by CFU at indicated time points by plating aliquots of cultures and plating

appropriate dilutions on 7H11 agar plates.

Physiological stress survival assay

Bacterial strains were grown till OD 0.5 (mid-log phase) and washed once with 1x PBST. Aliquots

of 2ml cell suspensions equivalent to OD 0.2 were treated with 0.5mM H_2O_2 (oxidative stress),

200µM Deta-NO (nitrosative stress), 0.01% SDS (membrane stress), PBST (nutrient starvation

stress), Phosphate citrate buffer, pH 5.5 (acidic stress) *in-vitro*. The survival of the respective strains

was determined by CFU plating at indicated time points and calculating survival index (i.e., CFU

of strain/ CFU of *Rv-WT*).

Drug susceptibility (MIC determination)

Auto-bioluminescent bacterial strains were grown till OD 0.5 (mid-log phase) and washed with 1x

PBST. Anti-TB drugs: Rifampicin (RIF), Isoniazid (INH), Bedaquiline (BDQ), Ethambutol (EMB),

Clofazimine (CFZ) and Pyrazinamide (PZA) were serially diluted on a 10-points, 2-fold format and

added to the 96-well plates (Corning). Cells were added at final density of OD 0.01 to the wells

containing drugs and incubated at 37°C for 3 days as described previously [46]. Finally,

luminescence was recorded as an indicator of bacterial viability which was expressed as dose-

response curves and IC_{50} were calculated using non-linear regression analysis in GraphPad Prism.

For MIC determination at pH 5.5, the drug dilutions and the bacterial suspensions (final density of

OD 0.05) were prepared in 7H9 medium supplemented ADS (pH adjusted to 5.5). For MIC

determination of PZA at pH 5.5 upon supplementation with methionine, an additional group of

ΔmetZ + Methionine was maintained. Dose-response curves were prepared similarly as described

above.

Kill kinetics

Bacterial strains were grown till OD 0.5 (mid-log phase) and washed with 1x PBST. Aliquots of

cell suspensions (2ml) at a bacterial density of OD 0.2 were treated with 10X MIC concentrations

of anti-TB drugs- Rifampicin (RIF), Isoniazid (INH), Ethambutol (EMB) and Bedaquilline (BDQ)

for mentioned time intervals. Bacterial survival was determined by CFU enumeration at indicated

time points. For pH-dependent kill kinetics of BDQ, 7H9 (+ADS) medium acidified to pH 5.5 was

employed and 10X MIC BDQ treatment was given as described above followed by CFU plating at

indicated time points.

Methionine estimation

Log-phase cultures of mycobacterial strains were washed with 1X PBST and transferred to 7H9

medium maintained at pH 7.2 or 5.5 at an OD 0.4. Cultures were allowed to grow for 6 days after

which they were washed with 1X PBS once followed by washing with chilled methanol. Cells

equivalent to OD 1 ($\sim$3x10^8 cells/mL) were lysed in 1 ml extraction buffer (Acetonitrile:Methanol:

water:: 2:2:1) by bead beating with intermittent cooling of 1 min per cycle. Lysates were clarified

by spinning at 13,000 rpm, 10 mins at 4°C. The clarified supernatants were filtered through 0.22-μ

filters and dried by speed vac to evaporate the extraction buffer. Dried pellets were resuspended in

methanol:water (1:1) + 0.1% formic acid and vigorously mixed. Finally, 10μL of the reconstituted

samples were run on Hypersil Gold C8 column (100 x 2.1mm) with 0.1% formic acid in water as

mobile phase A and 0.1% formic acid in acetonitrile as mobile phase B on SCIEX 6500+ QTRAP

and peak and retention time was matched with standard (Methionine) across all the samples. Peak

intensities were normalized to CFU of the respective samples.

Total ATP estimation

Cultures of *Rv-WT*, *ΔmetZ* and complement were grown at acidic pH 5.5 till OD 0.5 (mid-log phase)

and washed with 1X PBST. Cells were resuspended in 1000μL 1X PBST (with Protease Inhibitor

Cocktail) and lysed by 5 cycles: 45 seconds of bead beating with intermittent cooling on ice. Lysates

were passed through 0.5-μ sized filters to remove any bacterial cells or debris. 50μL of clarified

lysates were transferred to a 96-well plate (white, flat-bottom, SPL) and mixed (1:1) with BacTiter-

Glo (Promega) reaction buffer followed by incubation at 37°C for 30 minutes. 1X PBS was used as

blank to cancel out the background signals from the samples. Luminescence was recorded using a

Multimode Microplate reader (SpectraMax Pro ID3, Molecular Devices). Normalization was done

using total protein content of the respective samples and ATP levels were expressed as relative units

w.r.t. *Rv-WT* strain. For ATP estimation of BDQ treated cultures, 0.4 OD bacterial suspensions

(2ml) of the respective strains were treated with 2X MIC concentration of BDQ for the indicated

time-points and samples were prepared as described above.

For determination of total ATP levels upon supplementation of methionine, the bacterial strains

were grown to OD 0.5, washed and transferred to 7H9 (+ADS) pH 5.5 with an additional group of

ΔmetZ +Methionine. The cultures were allowed to grow for 5 days, washed with PBST and lysates

were prepared by bead-beating the cells. ATP estimation was performed as described above.

Metabolic activity

Auto-bioluminescent mycobacterial strains were grown till OD 0.5 (mid-log phase) and washed

with 1X PBST to remove the spent media. Cells were resuspended in 1X PBST and re-diluted to an

OD of 0.4 in media acidified to the required pH (pH 5.5). 100μL of these suspensions were added

to a 96-well plate (white, flat-bottom, SPL) containing 7H9 (ADS + 0.1% glycerol, pH 5.5) and

serially diluted (2-fold). The periphery wells were filled with sterile medium to prevent desiccation.

The plate was incubated at 37°C for 5 days after which 20μL of PrestoBlue (Thermo) was added to

each well followed by an additional incubation of 2 days to allow the development of color. After

2 days of incubation, luminescence read-out was taken and contents of the wells were transferred

to a clear bottom black plate followed by recording fluorescence of AlamarBlue at 560(ex)/

590(em). Ratio of Fluorescence/Luminescence was calculated to determine metabolic activity per

cell.

Intracellular redox state

Rv-WT, *ΔmetZ* and complement strains were transformed with plasmid encoding redox biosensor,

Mrx1-roGFP2 as described previously [47]. Cells were grown to OD 0.5 and washed with 1X PSBT

and resuspended in PBST in MCTs. *Rv-WT* cells in respective tubes were treated with either 100μM

H_2O_2 or 100μM DTT as control groups for oxidized and reduced conditions respectively. Further,

cells were transferred to 96-well plate (clear bottom, black wells) and excitation at 390 and 490 nm

while emission at 510 nm was recorded. Ratio of fluorescence at 390 nm/490 nm were compared to

determine intracellular redox state of the cells.

For determination of intracellular ROS levels upon supplementation of methionine, the bacterial

strains were grown to OD 0.5, washed and transferred to 7H9 (+ADS) pH 7.2 and pH 5.5 with an

additional group of *ΔmetZ* +Methionine. The cultures were allowed to grow for 5 days, washed with

PBST and resuspended in a final volume of 1ml PBST at density OD 0.5. The samples were

transferred to the 96-well plate (clear bottom, black wells) and excitation at 390 and 490 nm while

emission at 510 nm was recorded. Ratio of fluorescence at 390 nm/490 nm were compared to

determine intracellular redox state of the cells.

Lipid hydroperoxidation assay

Total lipids were extracted from the respective bacterial strains as described previously [48]. Briefly,

bacterial cells were grown to an OD 0.8-1.0 and pelleted down. Pellets were suspended in

methanol:chloroform (2:1), 6 ml overnight at 37°C to extract total cellular lipids. Next day, cells

were spun down and supernatants were collected. Supernatants having total cellular lipids were

further precipitated by saturating the supernatants with 0.8% KCl. The lower phase was collected

and dried in glass vials to evaporate the solvents and concentrate the lipids. Dried lipids were

resuspended in 1 ml chloroform (Sigma) and used for FOX2 assay for determining lipid

hydroperoxidation in the extracted lipids as described elsewhere [49]. Briefly, 200μL of the

concentrated lipids were mixed with 1000μL FOX2 reagent in MCTs in dark for 1hr. Samples were

transferred to a 96-well plate and read out at 560 nm.

Intracellular sulfide estimation

Total sulfide estimation was performed as described previously [50] [37]. Briefly, *Rv-WT*, *ΔmetZ* and complement strains were grown to OD 0.5 (mid-log phase) in regular 7H9 medium supplemented with OADS and 0.01% Tween 80. Cells equivalent to OD 0.5 were inoculated into 7H9 medium supplemented with ADS and 0.01% tyloxapol at pH 7.2 and pH 5.5. After 4 days of incubation, the cultures were pellet down and washed with 1X PBST to remove spent media. Pellets were resuspended in the same media with respective pH 7.2 and 5.5 at an OD 0.4. 100μL of the cell suspensions were added to 96-well plate and mixed with 100μL 2X Bismuth chloride buffer. Following an incubation of 4 hours at 37°C, colorimetry was done at 405nm. The values were normalized to culture OD to determine relative sulfide levels in the respective strains at two different pH.

Extracellular pH (pH$_{Ex}$) measurement

Extracellular pH (pH$_{Ex}$) of the bacterial cultures was measured using pH indicator dye – Chlorophenol red (CPR) as described previously [28]. Briefly, 7H9 medium supplemented with ADS and 0.01% Tween 80 was used to prepare pH standards – pH 7.0, 6.5, 6.0, 5.5, 5.0, 4.5 and 4.0. To 1ml of these standards, 0.004% v/v CPR was added and colorimetric read-outs at 430nm and 590nm were recorded. Ratio of readouts at 430/590nm were used to prepare a pH calibration curve. For pH of cultures, 250μL of bacterial cultures at indicated time points were taken and spun down at 12,000 rpm for 10 minutes to pellet down bacterial cells. To the supernatants, 0.004% v/v CPR was added and colorimetric assay was performed as described above. Ratio of 430/590nm of cultures was used to extrapolate pH$_{Ex}$ using pH calibration curve. Fresh pH standards were prepared for each time-point.

Macrophage infection

THP-1 monocytic cells were grown in RPMI medium (Lonza) supplemented with 10% fetal bovine

serum (FBS, Invitrogen). Further, cells were seeded at a density of 5×10^5 cells/ well in a 12-well

plate (Corning) and matured for 24 hours with PMA (25ng/mL, Sigma) followed by a rest of 24

hours. Whenever required, cells were activated with IFN-γ (10ng/ml, Peprotech) for 16 hrs before

infection. Single cell suspensions of the bacterial strains (*Rv-WT*, *ΔmetZ* and complement) were

prepared by soft-spin method as described [43] and used for infecting resting or activated

macrophages at an MOI of 1. Bacteria were allowed to infect the cells for 4 hours after which,

extracellular bacteria were washed off by three washes with pre-warmed PBS (Gibco). At indicated

time points, macrophages were lysed with 1X PBS + 0.01% Triton X-100 to release intracellular

bacterial cells. Lysates were serially diluted and plated for bacterial enumeration.

Confocal Microscopy

THP-1 macrophages were seeded on glass coverslips in a 24-well plate (5×10^5 cells/well) and were

activated by 10ng/ml of human IFN-γ (Peprotech, USA) for 16 hours before infection. Activated

THP-1 cells were infected with mycobacterial strains (*Rv-WT*, *ΔmetZ* and complement) harboring

Live-Dead reporter plasmid [51] constitutively expressing mEmerld at an MOI of 10. After 4 hours

of infection, the cells were washed thrice with 1X PBS to remove the extracellular bacteria and

replenished with fresh media. Subsequently, after 24 hours post infection, cells were incubated with

50nM LysoTracker Red DND-99 (Invitogen life Technologies, CA, USA) in complete RPMI media

for 30 minutes at 37°C with 5% CO2 and later fixed with 4% paraformaldehyde in 1X PBS.

Coverslips were mounted with ProLong Diamond antifade with DAPI (Molecular Probes by Life

Technologies, CA, USA). Finally, slides were imaged by Olympus Confocal Laser Scanning

Microscope (CLSM). Later, GFP-expressing mycobacteria that colocalized with LysoTracker Red

was determined by analyzing the ~ 100 phagosomes and co-localization index was determined.

Animal Infection

In-vivo experiment in this study was performed with approval from the Institutional Animal Ethics

Committee (IAEC Approval No: IAEC/THSTI/165) as per guidelines laid by the Committee for the

Purpose of Control of Experiments on Animals (CPCEA), Government of India. Briefly, bacterial

suspensions of respective strains at a density of 10^8 cells/mL were prepared in normal saline and

guinea pigs (Duncan Hartley) were infected via aerosol route using Inhalation Exposure System

(Glas-Col). To determine baseline infection in all groups, a few representative animals (n=3 per

group) were sacrificed, followed by an evaluation of disease progression at indicated time points

with animals (n=5) per group; organs were harvested and bacterial burdens were determined by

CFU plating. Histopathological damage to the lung tissue was determined by H&E staining of lungs

fixed in 5% paraformaldehyde followed by granuloma scoring of the infected lung tissue.

CRISPRi-mediated *in-vivo* fitness

The experiment was performed with approval from the Institutional Animal Ethics Committee (IAEC

Approval No: IAEC/THSTI/196) as per guidelines laid by the Committee for the Purpose of Control

of Experiments on Animals (CPCEA), Government of India. Briefly, bacterial suspensions of

respective strains (***Rv***::NT and *ΔZ::metB$_{KD}$*) at a density of $3x10^8$ cells/mL were prepared in normal

saline and BALB/c mice were infected via aerosol route using Inhalation Exposure System (Glas-

Col). To induce CRISPRi-mediated gene silencing of *metB* in *ΔZ::metB$_{KD}$* strain, doxycycline

(1mg/ml) was administered to mice via oral route at indicated time points [52]. NT (non-targeting

CRISPRi strain) was maintained with doxycycline as control which is isogenic to *Rv-WT*. Animals

were euthanized at indicated time points, organs (lungs and spleens) were harvested and bacterial

burden were determined by CFU plating.

Bioinformatics analysis

MetZ   protein   sequence   was   retrieved   from   Mycobrowser

(https://mycobrowser.epfl.ch/genes/Rv0391) and the cluster of orthologous groups analysis (COG)

was performed using Eggnog v5 [53]. Further, a multiple sequence alignment of the protein sequence

from members of *Mycobacterium* genus (*M. tuberculosis, M. bovis, M. canettii, M. marinum, M.*

leprae and *M. smegmatis*) along with gram-negative (*Pseudomonas aeruginosa*) and gram-positive

bacterium (*Corynebacterium glutamicum* and *Rhodococcus*) was performed using Clustal W [54] to

identify common domains and motifs amongst the protein sequences.

Molecular docking and molecular dynamics (MD) simulations

For system preparation, the crystal structure of putative O-succinylhomoserine sulfhydrylase of *M.*

tuberculosis was retrieved from RCSB PDB [55] (PDB-ID: 3NDN [56]). The protein has 4 chains and

exists in a homo-tetramer form. The receptor has a covalently bound pyridoxal-5-phosphate attached

to a modified lysine (K219). The substrate, O-succinyl homoserine, was retrieved from PubChem

(PubChemCID 439406). The substrate undergoes a sulfur transfer reaction in the presence of

pyridoxal-5-phosphate forming L-homocysteine which takes part in the methylation cycle. The

receptor protein was optimized and then minimized using the OPLS3 force field of Protein Preparation

Wizard module [57] of the Schrodinger suite. Next, Binding Site Identification was done using

AutoDock 4 [58] wherein blind docking of the substrate in the presence and absence of the cofactor to

identify the possible binding site was performed. Autogrid4 was used to build a 126x126x126 points

grid map with a spacing of 0.586 angstrom and positioned at the center of mass of the receptor protein.

The Lamarckian genetic algorithm was employed to generate 200 runs. On docking the substrate in the

presence of the cofactor, no suitable results were produced. However, docking in the absence of the

co-factor resulted in comparatively higher binding affinity and a larger cluster size. Focused docking

on the identified site was also carried out by generating a grid map of 40x40x40 units with a spacing

of 0.375 A and 17X20X22 as grid center coordinates. MM-GBSA of few conformers based on their

binding affinity and cluster size was done using Prime module [59] of the maestro. Finally, the lowest

energy conformations of the substrate were subsequently used for constant pH MD. Further, Constant

pH Molecular Dynamics Simulation (MDS) was performed with the Desmond [60] module from

Schrodinger suite. Both systems were protonated at pH 5.5 and 7.0, respectively, and minimized with

OPLS- all Atom force fields and solvated with the predefined TIP3P water solvent model. The

complexes were placed in the orthorhombic periodic boundary conditions to specify the shape and size

of the repeating unit buffered at 12 A distances. 4 Na+ and 9 Na+ ions were added to both the systems

respectively, to neutralize the systems. The NPT ensemble was employed for the simulation with a

timestep of 0.002 ps. The simulation was carried out for a total of 300 ns for each system and the

coordinates were saved at intervals of 20 ps to generate 15000 frames. Also, the protein was simulated

without the substrate at the two pH values. The post-processing of MD simulation trajectories was

carried out using VMD [61]. The root-mean-square-deviation and fluctuation were measured for the

protein, substrate and cofactor. Lastly, for thermodynamics analysis of the system we performed

Molecular Mechanics Generalized Born Surface Area (MM-GBSA) analysis to evaluate the MD

trajectories using Schrodinger software. To effectively encompass the thermodynamic properties,

every 500th frame was selected for MM-GBSA analysis.

Statistics and Reproducibility

The data values are represented as mean ± SEM of the replicates from two-three independent

experiments. Wherever applicable, the number of replicates per group have been indicated by the

dots. Statistical comparisons have been made with *Rv-WT* in most cases unless mentioned otherwise.

GraphPad Prism (ver 9.0) has been used for statistical analysis of all data with suitable statistical tests

mentioned with figure legend.

Conflict of Interests

The authors declare no conflict of interests

Acknowledgements

The authors acknowledge THSTI for infrastructural support, Department of Biotechnology,

Government of India for funding, the Experimental Animal Facility (and staff) for supporting animal

experimentations and BSL-3 facility (and staff) for conducting all *M. tuberculosis* related experiments.

Technical support of Mr. Surjit Yadav is duly acknowledged. We acknowledge the Multi-Omics

facility for LC-MS based experiments. We also duly acknowledge the receipt of plasmids and

constructs used in the study (as listed in Table S1).

Funding

This study was supported by the grant (No. EM/Dev/SG/212/7864/2023) from the Indian Council of

Medical Research (ICMR), Government of India and intramural funding by BRIC-THSTI to A.K.P.

and are duly acknowledged. VKN was supported by the research fellowship from ICMR (3/1/3/JRF-

2018/HRD-043/64467). VB is supported by the research fellowship from CSIR :

09/1049(11494)/2021-EMR-I. TS was supported by the research fellowship from CSIR :

09/1049(0036)/2019-EMR-I.

Supplementary material

1. Table S1: List of plasmids and constructs used in the study

2. Table S2: List of bacterial strains used in the study

3. Table S3: List of primers used in the study

4. Supplementary data

List of relevant references:

1.   WHO, *World TB report 2022.* 2023.

2.   Yelamanchi, S.D. and A. Surolia, *Targeting amino acid metabolism of Mycobacterium tuberculosis for developing inhibitors to curtail its survival.* IUBMB Life, 2021. **73**(4): p. 643-658.

3.   Kim, J.M., et al., *Formyl-methionine as an N-degron of a eukaryotic N-end rule pathway.* Science, 2018. **362**(6418).

4.   Berney, M., et al., *Essential roles of methionine and S-adenosylmethionine in the autarkic lifestyle of Mycobacterium tuberculosis.* Proc Natl Acad Sci U S A, 2015. **112**(32): p. 10008-13.

5.   Chaton, C.T., et al., *Structural analysis of mycobacterial homoserine transacetylases central to methionine biosynthesis reveals druggable active site.* Sci Rep, 2019. **9**(1): p. 20267.

6.   Chaudhary, D., et al., *Identification of small molecules targeting homoserine acetyl transferase from Mycobacterium tuberculosis and Staphylococcus aureus.* Sci Rep, 2022. **12**(1): p. 13801.

7.   Berger, B.J. and M.H. Knodel, *Characterisation of methionine adenosyltransferase from Mycobacterium smegmatis and M. tuberculosis.* BMC Microbiol, 2003. **3**: p. 12.

8.   Venos, E.S., et al., *Branched-chain amino acid aminotransferase and methionine formation in Mycobacterium tuberculosis.* BMC Microbiol, 2004. **4**: p. 39.

9.   Lee, H.S. and B.J. Hwang, *Methionine biosynthesis and its regulation in Corynebacterium*
glutamicum: parallel pathways of transsulfuration and direct sulfhydrylation. Appl Microbiol
Biotechnol, 2003. **62**(5-6): p. 459-67.

10.   Hasenoehrl, E.J., et al., *Derailing the aspartate pathway of Mycobacterium tuberculosis to*
eradicate persistent infection. Nat Commun, 2019. **10**(1): p. 4215.

11.   Smith, D.A., et al., *Characterization of auxotrophic mutants of Mycobacterium tuberculosis*
and their potential as vaccine candidates. Infect Immun, 2001. **69**(2): p. 1142-50.

12.   Wheeler, P.R., et al., *Functional demonstration of reverse transsulfuration in the*
Mycobacterium tuberculosis complex reveals that methionine is the preferred sulfur source for
pathogenic Mycobacteria. J Biol Chem, 2005. **280**(9): p. 8069-78.

13.   Mishra, R., et al., *Targeting redox heterogeneity to counteract drug tolerance in replicating*
Mycobacterium tuberculosis. Sci Transl Med, 2019. **11**(518).

14.   Hatzios, S.K. and C.R. Bertozzi, *The regulation of sulfur metabolism in Mycobacterium*
tuberculosis. PLoS Pathog, 2011. **7**(7): p. e1002036.

15.   Foglino, M., et al., *A direct sulfhydrylation pathway is used for methionine biosynthesis in*
Pseudomonas aeruginosa. Microbiology (Reading), 1995. **141 (Pt 2)**: p. 431-9.

16.   Belfaiza, J., et al., *Direct sulfhydrylation for methionine biosynthesis in Leptospira meyeri.* J
Bacteriol, 1998. **180**(2): p. 250-5.

17.   Hwang, B.J., et al., *Corynebacterium glutamicum utilizes both transsulfuration and direct*
sulfhydrylation pathways for methionine biosynthesis. J Bacteriol, 2002. **184**(5): p. 1277-86.

18.   Cole, S.T., et al., *Deciphering the biology of Mycobacterium tuberculosis from the complete*
genome sequence. Nature, 1998. **393**(6685): p. 537-44.

19.   Griffin, J.E., et al., *High-resolution phenotypic profiling defines genes essential for*
mycobacterial growth and cholesterol catabolism. PLoS Pathog, 2011. **7**(9): p. e1002251.

20.   Murphy, K.C., K. Papavinasasundaram, and C.M. Sassetti, *Mycobacterial recombineering.*
Methods Mol Biol, 2015. **1285**: p. 177-99.

21.   Griffin, J.E., et al., *Cholesterol catabolism by Mycobacterium tuberculosis requires*
transcriptional and metabolic adaptations. Chem Biol, 2012. **19**(2): p. 218-27.

22.   Chang, J.C., et al., *igr Genes and Mycobacterium tuberculosis cholesterol metabolism.* J
Bacteriol, 2009. **191**(16): p. 5232-9.

23.   Srivastava, M., et al., *Molecular dynamics simulation reveals the possible druggable hot-spots*
of USP7. Oncotarget, 2018. **9**(76): p. 34289-34305.

24.   Mittal, L., et al., *Targeting cryptic-orthosteric site of PD-L1 for inhibitor identification using*
structure-guided approach. Arch Biochem Biophys, 2021. **713**: p. 109059.

25.   Kumari, A., et al., *Deciphering the Structural Determinants Critical in Attaining the FXR*
Partial Agonism. J Phys Chem B, 2023. **127**(2): p. 465-485.

26.   Howe, M.D., et al., *Methionine Antagonizes para-Aminosalicylic Acid Activity via Affecting*
Folate Precursor Biosynthesis in Mycobacterium tuberculosis. Front Cell Infect Microbiol,
2018. **8**: p. 399.

27.   Khan, M.Z., et al., *Redox homeostasis in Mycobacterium tuberculosis is modulated by a novel*
actinomycete-specific transcription factor. EMBO J, 2021. **40**(14): p. e106111.

28.   Gouzy, A., et al., *Growth of Mycobacterium tuberculosis at acidic pH depends on lipid*
assimilation and is accompanied by reduced GAPDH activity. Proc Natl Acad Sci U S A, 2021.
118(32).

29.   Loewen, P.C., *Levels of glutathione in Escherichia coli.* Can J Biochem, 1979. **57**(2): p. 107-
11.

30.   Akhova, A.V. and A.G. Tkachenko, *ATP/ADP alteration as a sign of the oxidative stress*
development in Escherichia coli cells under antibiotic treatment. FEMS Microbiol Lett, 2014.
353(1): p. 69-76.

31.   Singh, S., et al., *Unraveling the role of the transcriptional regulator VirS in low pH-induced*
responses of Mycobacterium tuberculosis and identification of VirS inhibitors. J Biol Chem,
2019. **294**(26): p. 10055-10075.

32.   Mehta, M., R.S. Rajmani, and A. Singh, *Mycobacterium tuberculosis WhiB3 Responds to*
Vacuolar pH-induced Changes in Mycothiol Redox Potential to Modulate Phagosomal
Maturation and Virulence. J Biol Chem, 2016. **291**(6): p. 2888-903.

33.   Banerjee, S.K., et al., *The sensor kinase MtrB of Mycobacterium tuberculosis regulates hypoxic*
survival and establishment of infection. J Biol Chem, 2019. **294**(52): p. 19862-19876.

34.   Choudhary, E., et al., *Deciphering the Proteomic Landscape of Mycobacterium tuberculosis in*
837   *Response to Acid and Oxidative Stresses.* ACS Omega, 2022. **7**(30): p. 26749-26766.

838   35.   Rohde, K.H., et al., *Linking the transcriptional profiles and the physiological states of*
839   *Mycobacterium tuberculosis during an extended intracellular infection.* PLoS Pathog, 2012.
840   **8**(6): p. e1002769.

841   36.   Saini, V., et al., *Hydrogen sulfide stimulates Mycobacterium tuberculosis respiration, growth*
842   *and pathogenesis.* Nat Commun, 2020. **11**(1): p. 557.

843   37.   Kunota, T.T.R., et al., *Mycobacterium tuberculosis H2S Functions as a Sink to Modulate*
844   *Central Metabolism, Bioenergetics, and Drug Susceptibility.* Antioxidants (Basel), 2021. **10**(8).

845   38.   Rahman, M.A., et al., *The Role of Host-Generated H2S in Microbial Pathogenesis: New*
846   *Perspectives on Tuberculosis.* Front Cell Infect Microbiol, 2020. **10**: p. 586923.

847   39.   Rahman, M.A., et al., *Hydrogen sulfide dysregulates the immune response by suppressing*
848   *central carbon metabolism to promote tuberculosis.* Proc Natl Acad Sci U S A, 2020. **117**(12):
849   p. 6663-6674.

850   40.   Brauner, A., et al., *Distinguishing between resistance, tolerance and persistence to antibiotic*
851   *treatment.* Nat Rev Microbiol, 2016. **14**(5): p. 320-30.

852   41.   Quinonez, C.G., et al., *The Role of Fatty Acid Metabolism in Drug Tolerance of Mycobacterium*
853   *tuberculosis.* mBio, 2022. **13**(1): p. e0355921.

854   42.   Andreu, N., et al., *Optimisation of bioluminescent reporters for use with mycobacteria.* PLoS
855   One, 2010. **5**(5): p. e10777.

856   43.   Babunovic, G.H., et al., *CRISPR Interference Reveals That All-Trans-Retinoic Acid Promotes*
857   *Macrophage Control of Mycobacterium tuberculosis by Limiting Bacterial Access to*
858   *Cholesterol and Propionyl Coenzyme A.* mBio, 2022: p. e0368321.

859   44.   Pandey, A.K. and C.M. Sassetti, *Mycobacterial persistence requires the utilization of host*
860   *cholesterol.* Proc Natl Acad Sci U S A, 2008. **105**(11): p. 4376-80.

861   45.   Schmittgen, T.D. and K.J. Livak, *Analyzing real-time PCR data by the comparative C(T)*
862   *method.* Nat Protoc, 2008. **3**(6): p. 1101-8.

863   46.   Vocat, A., et al., *Bioluminescence for assessing drug potency against nonreplicating*
864   *Mycobacterium tuberculosis.* Antimicrob Agents Chemother, 2015. **59**(7): p. 4012-9.

47. Bhaskar, A., et al., *Reengineering redox sensitive GFP to measure mycothiol redox potential*
of Mycobacterium tuberculosis during infection. PLoS Pathog, 2014. **10**(1): p. e1003902.

48. Folch, J., M. Lees, and G.H. Sloane Stanley, *A simple method for the isolation and purification*
of total lipides from animal tissues. J Biol Chem, 1957. **226**(1): p. 497-509.

49. Nambi, S., et al., *The Oxidative Stress Network of Mycobacterium tuberculosis Reveals*
Coordination between Radical Detoxification Systems. Cell Host Microbe, 2015. **17**(6): p. 829-
37.

50. Mishra, R., et al., *Heterogeneous Host-Pathogen Encounters Coordinate Antibiotic Resilience*
in Mycobacterium tuberculosis. Trends Microbiol, 2021. **29**(7): p. 606-620.

51. Cai, Y., et al., *Host immunity increases Mycobacterium tuberculosis reliance on cytochrome*
bd oxidase. PLoS Pathog, 2021. **17**(7): p. e1008911.

52. Tripathi, A., et al., *Mycobacterium tuberculosis requires SufT for Fe-S cluster maturation,*
metabolism, and survival in vivo. PLoS Pathog, 2022. **18**(4): p. e1010475.

53. Huerta-Cepas, J., et al., *eggNOG 5.0: a hierarchical, functionally and phylogenetically*
annotated orthology resource based on 5090 organisms and 2502 viruses. Nucleic Acids Res,
2019. **47**(D1): p. D309-D314.

54. Thompson, J.D., D.G. Higgins, and T.J. Gibson, *CLUSTAL W: improving the sensitivity of*
progressive multiple sequence alignment through sequence weighting, position-specific gap
penalties and weight matrix choice. Nucleic Acids Res, 1994. **22**(22): p. 4673-80.

55. Berman, H.M., et al., *The Protein Data Bank.* Nucleic Acids Res, 2000. **28**(1): p. 235-42.

56. Baugh, L., et al., *Increasing the structural coverage of tuberculosis drug targets.* Tuberculosis
(Edinb), 2015. **95**(2): p. 142-8.

57. Sastry, G.M., et al., *Protein and ligand preparation: parameters, protocols, and influence on*
virtual screening enrichments. J Comput Aided Mol Des, 2013. **27**(3): p. 221-34.

58. Morris, G.M., et al., *AutoDock4 and AutoDockTools4: Automated docking with selective*
receptor flexibility. J Comput Chem, 2009. **30**(16): p. 2785-91.

59. Jacobson, M.P., et al., *A hierarchical approach to all-atom protein loop prediction.* Proteins,
2004. **55**(2): p. 351-67.

60.   Bowers, K.J.C., E.; Xu, H.; Dror, R. O.; Eastwood, M. P.; Gregersen, B. A.; Klepeis, J. L.;

Kolossvary, I.; Moraes, M. A.; Sacerdoti, F. D.; Salmon, J. K.; Shan, Y.; Shaw, D. E., , *Scalable*

algorithms for molecular dynamics simulations on commodity clusters. Proceedings of the

ACM/IEEE Conference on Supercomputing (SC06), 2006.

61.   Humphrey, W., A. Dalke, and K. Schulten, *VMD: visual molecular dynamics.* J Mol Graph,

1996. **14**(1): p. 33-8, 27-8.

Figure 1. *metZ* is dispensable for growth in *M. tuberculosis.* (A). Methionine biosynthesis pathways

in bacteria. Two alternate pathways exist in many bacterial species – transsulfuration where cysteine is

used as an S-donor and direct sulfhydrylation wherein H_2S is used as an S-donor. *Rv0391* encoded MetZ

is such putative O-succinyl homoserine sulfhydrylase (OSHS) involved in direct sulfhydrylation in *M.*

tuberculosis. (B). Cluster of Orthologues analysis for *M. tuberculosis* MetZ (Rv0391) protein sequence

with member species of ENOG502GJ5S as indicated by EggNog database. (C). Schematic for knockout

generation of *ΔmetZ* in *M. tuberculosis.* (D). Confirmation of *metZ* knockout using RT-qPCR for the

loss of *metZ* transcript. Growth kinetics of auto-bioluminescent *ΔmetZ* with (E) glycerol (0.1%) and (F)

cholesterol (200µM) as C-source. The data summary represents means ± SEM for technical replicates

from two independent experiments. Statistical comparisons were made w.r.t. *Rv-WT* for (D) using

Unpaired *t*-test. ****$P<0.0001$. Comparisons not marked with statistical significance were not

significant with *Rv-WT* vs *ΔmetZ* for (E) and (F).

Figure 2: *metZ* deletion renders *M. tuberculosis* sensitive to physiological stressors and reduced *ex-vivo* fitness. (A). *In-vitro* stress survival assay. Mycobacterial cultures were treated with reagents mimicking host-induced stresses: H_2O_2 (Oxidative stress), DetaNO (Nitrosative stress), SDS (Surfactant stress), low pH (5.0), PSBT (nutrient starvation) *in-vitro* and survival was determined by CFU plating. (B). Intrabacterial redox state determination of *ΔmetZ* compared to *Rv-WT* using genetically encoded redox biosensor Mrx1-roGFP2. (C). Lipid hydroperoxidation levels in *ΔmetZ* compared to *Rv-WT* and complement strains using FOX2 colorimetry assay. (D). Intra-macrophage survival assay using THP-1 monocyte-derived macrophages resting or activated using IFN-γ. The survival index was determined by CFU enumeration. (E). Intracellular co-localization of GFP labelled mycobacterial strains with LysoTracker stained acidic compartments inside IFN-γ activated macrophages. Data is represented as mean ± SEM of the replicates. Comparisons for statistical significance have been made with *Rv-WT* (Control) using two-way ANOVA (Dunnett's multiple comparisons test) for (A) where **P = 0.0084, ****P<0.0001; two-way ANOVA (Sidak's multiple comparisons test) for (B) where **P = 0.0034; ordinary one-way ANOVA (Dunnett's multiple comparisons test) for (C) where ****P<0.0001; two-way ANOVA (Dunnett's multiple comparisons test) for (D) where ***P = 0.0002, ****P<0.0001; comparison of *ΔmetZ* (Resting vs. Activated macrophages) using unpaired t-test for (D) where **P = 0.0034 and ordinary one-way ANOVA for (E) where, *P = 0.0166, ****P<0.0001.

Figure 3: Direct sulfhydrylation pathway is essential for *M. tuberculosis* to survive at low pH. (A) *In-vitro* growth kinetics of *ΔmetZ* at various pH – 7.2, 5.5 and 5.0. (B). Expression analysis of genes involved in transsulfuration (*metB* and *metC*) and direct sulfhydrylation (*metZ*) using RT-qPCR. (C). Metabolic activity of *ΔmetZ* at acidic pH using redox dye AlamarBlue Assay. (D). Intrabacterial ATP levels estimation of *ΔmetZ* at acidic pH 5.5. (E). Pyrazinamide (PZA) sensitivity assay of *ΔmetZ* at pH 5.5. IC50 of PZA for the respective strains at acidic pH 5.5 was determined and represented as a dose-response curve. (F). Methionine estimation at pH 7.2 and pH 5.5 was performed using LC-MS. (G). Growth defect in *ΔmetZ* at pH 5.5 was rescued using exogenous methionine (Met) supplementation. (H). The intrabacterial redox state of *ΔmetZ* was determined using the Mrx1-roGFP2 redox biosensor upon supplementation with methionine (Met) at pH 7.2 and 5.5. (I). Bacterial ATP levels at pH 5.5, upon supplementation of methionine (Met). The dose-response curve for PZA was made by Non-linear regression analysis using GraphPad Prism. Data is represented as mean ± SEM of the replicates.

Comparisons for statistical significance have been made using two-way ANOVA (Dunnett's multiple comparisons test) for (A) where, ****P<0.0001, ns = non-significant; ordinary one-way ANOVA (Dunnett's multiple comparisons test) for (B) where, ***P = 0.004, ****P<0.0001, ns = non-significant; ordinary one-way ANOVA (Dunnett's multiple comparisons test) for (C) where *P = 0.011, ns = non-significant; ordinary one-way ANOVA (Dunnett's multiple comparisons test) for (D) where ****P<0.0001; ordinary one-way ANOVA (Dunnett's multiple comparisons test) for (E) where *P=0.0155, ns = non-significant; two-way ANOVA (Dunnett's multiple comparisons test) for (F) where **P = 0.0030, ns = non-significant; two-way ANOVA (Dunnett's multiple comparisons test) for (G) w.r.t. $\Delta metZ$ where *P=0.0157, # = ****P<0.0001; ordinary one-way ANOVA (Dunnett's multiple comparisons test) for (H) w.r.t. $\Delta metZ$ where **P<0.006, ***P<0.0004, ****P<0.0001, ns = non-significant; ordinary one-way ANOVA (Dunnett's multiple comparisons test) for (I) w.r.t. $\Delta metZ$ where ****P<0.0001.

Figure 4. Molecular docking and Molecular dynamics (MD) simulations revealed enhanced stability of MetZ at acidic pH. (A). Structural representation of O-succinyl homoserine sulfhydrylase, rendered in new cartoon and colored in grey (single chain) with pyridoxal-5-phosphate in licorice and rendered atom-wise. (B). Dynamic characteristics of systems elucidated through constant pH MD simulations. The system evolution was measured via RMSD through the course of 300ns of the protein, substrate and the binding site of the substrate. Color coding is followed as the system at pH 7 (blue) and

at pH 5.5 (red). (C). The Binding Free Energy (in Kcal/mol) comparison of the complexes at pH 7 (blue)

and pH 5.5 (red). (D). The dynamic changes represent the positional shifts of the substrate (orange) and

co-factor (cyan) throughout the simulation, with a color gradient (time-dependent manner) from red

(initial), and white (intermediate) to blue (final) at pH 7 at pH 5.5. (E). Interaction map of the substrate

at pH 7 and pH 5.5. Residues lining the pocket (within 4.0 Å) are shown in licorice and colored atom-

wise (C: cyan, N: blue and O: red). The residues forming hydrogen bonds (dotted line in blue) and Salt-

bridges (dotted line in yellow) are shown. The Cα atom of residues involved in hydrophobic interaction

are shown in beads.

Figure 5: *metZ mediated* **direct sulfhydrylation pathway is critical for methionine biosynthesis at**

low pH. (A). Diagrammatic representation of experimental setup to elucidate independent existence of

direct sulfhydrylation pathway. CRISPRi-based knockdown of *metB* (*Rv1079*) strains was achieved in

the genetic backgrounds of *Rv-WT* and *ΔmetZ* strains. Upon induction with ATc, these strains were

depleted of *metB* transcripts and characterised for *in-vitro* growth kinetics, methionine levels and *in-vivo*

fitness. (B). *In-vitro* growth kinetics of CRISPRi knockdown strains. (C). Relative methionine levels in

CRISPRi knockdown strains either lacking *metB* ($Rv::metB_{KD}$) or both *metZ* and *metB* ($ΔZ::metB_{KD}$) by

LC-MS. (D). Graphical representation of *in-vivo* fitness experiment. Mice were infected with CRISPRi

knockdown of *metB* in Rv-WT and *ΔmetZ* background and orally fed with doxycycline (1mg/ml) to

induce knockdown of *metB* gene in the respective strains. Bacterial burden was determined by CFU

plating in lungs and spleen homogenates. (E). and (F). The bacterial burden of mycobacterial strains lacking either *metZ, metB,* or both in mice lungs and spleen lysates respectively. Data is represented as mean ± SEM of replicates. Comparisons for statistical significance have been made using ordinary one-way ANOVA (Dunnett's multiple comparisons test) for (B) where ***P<0.0003, ****P<0.0001; Unpaired t-test for (C) where ***P=0.0001; Mann-Whitney Test for Spleen bacterial burden in (E) where *P=0.015, **P=0.007 and two-way ANOVA (Dunnett's multiple comparisons) for Lungs bacterial burden in (E) where **P=0.0064, ***P=0.0003 and ****P<0.0001.

Figure 6: *metZ* is required for persister formation in the presence of BDQ. (A). Minimum Inhibitory Concentration (MIC) determination of the *ΔmetZ* in comparison to Rv-WT and Complement strains against anti-TB drugs: Rifampicin (RIF), Isoniazid (INH), Ethambutol (EMB) and Bedaquilline (BDQ) using microtiter plate assay. IC50 was determined for the drugs with the mentioned strains. (B). Kill kinetics of anti-TB drugs – RIF, INH, EMB, and BDQ at 10X MIC concentrations. Survival was determined in the form of viable bacterial count expressed as CFU. (C). Kinetics of ATP depletion post BDQ treatment. *ΔmetZ* along with Rv-WT and Complement strains were treated with 2X concentration of BDQ and ATP was estimated at indicated time points. *ΔmetZ* maintained high ATP levels. Data is represented as mean ± SEM of replicates. Comparison for statistical significance has been made using two-way ANOVA (Dunnett's Multiple Comparisons test) for (B) where *P<0.004 and ns = non-significant; two-way ANOVA (Dunnett's multiple comparisons test) for (B) where *P = 0.029, **P<0.005, ****P<0.0001 and ns, non-significant; using two-way ANOVA (Dunnett's multiple comparisons test) for (C) where ****P < 0.0001.

Figure 7: *metZ* **is required for survival of** *M. tuberculosis in-vivo***.** (A). Diagrammatic representation
of *in-vivo* fitness experimental set-up. Guinea pigs were infected with WT, mutant, and complement
strains, and disease progression was monitored through CFU plating at indicated time points. (B).
Bacterial burden in guinea pigs' lungs as determined by CFU plating of the lung homogenates of the
infected animals at indicated time points. The mutant strain failed to replicate in guinea pig lungs post-
week 4 of initial infection. (C). CFU burden in spleens of the infected animals revealed that the mutant
could not survive inside the spleen post week 4 of infection. (D). Gross lung pathology indicates
granulomatous lesions on the infected lungs at week 8-time point post initial infection. (E). H&E staining
(F). Total granuloma scores of three strains in lung sections revealed significantly lesser granulomas in
ΔmetZ infected lungs at week 8 post-infection. Data is represented as mean ± SEM of 5 replicates
(animals per group). Comparison for determining statistical significance has been made using the Mann-
Whitney Test for (B) and (C) where *P<0.05, **P=0.0079; ordinary one-way ANOVA (Dunnett's
Multiple Comparisons test) for (F) where **P=0.004, ***P=0.0004.

Figure 8. *MetZ-driven* direct sulfhydrylation is critical for methionine biosynthesis at low pH.
When *M. tuberculosis* infects macrophage, it passes through a series of compartments starting from the phagosome where pH is essentially ~6.5 and the bacteria are replicating, with enough nutrients (including methionine). Further, as host immunity kicks in IFN-γ induced production of ROS and RNS followed by phagosome maturation leads to a drop in pH to 5.5 and beyond. Such conditions are unfavourable for *M. tuberculosis* and the host begins to quench nutrients as a means to restrict the trapped bacteria. At this stage bacteria undergo several transcriptional and metabolic changes to sustain the host-induced insults, one of these, as per our findings, includes the induction of *metZ* expression. This helps bacteria survive these stress conditions and maintain methionine levels by utilizing endogenous as well as host-generated H_2S which acts as co-substrate and S-donor for methionine biosynthesis in acidic micro-environment. This enables bacteria to maintain sufficient levels of methionine to keep adaptation going while also helping the recycling of ATP, mycothiol, at pace with its metabolic activity.