Coordinated regulation of chemotaxis and resistance to copper by CsoR in *Pseudomonas putida*

Meina He, Yongxin Tao, Kexin Mu, Haoqi Feng, Ying Fan, Tong Liu, Qiaoyun Huang,

Yujie Xiao*, Wenli Chen*

National Key Laboratory of Agricultural Microbiology, Huazhong Agricultural

University, Wuhan 430070, China.

*To whom correspondence should be addressed. Tel: +86 27 87280670; Fax: +86 27

87280670; Email: wlchen@mail.hzau.edu.cn; yjxiao@mail.hzau.edu.cn

Abstract

Copper is an essential enzyme cofactor in bacteria, but excess copper is highly toxic. Bacteria can cope with copper stress by increasing copper resistance and initiating chemorepellent response. However, it remains unclear how bacteria coordinate chemotaxis and resistance to copper. By screening proteins that interacted with the chemotaxis kinase CheA, we identified a copper-binding repressor CsoR that interacted with CheA in *Pseudomonas putida*. CsoR interacted with the HPT (P1), Dimer (P3), and HATPase_c (P4) domains of CheA and inhibited CheA autophosphorylation, resulting in decreased chemotaxis. The copper-binding of CsoR weakened its interaction with CheA, which relieved the inhibition of chemotaxis by CsoR. In addition, CsoR bound to the promoter of copper-resistance genes to inhibit gene expression, and copper-binding released CsoR from the promoter, leading to increased gene expression and copper resistance. *P. putida* cells exhibited a chemorepellent response to copper in a CheA-dependent manner, and CsoR inhibited the chemorepellent response to copper. Besides, the CheA-CsoR interaction also existed in proteins from several other bacterial species. Our results revealed a mechanism by which bacteria coordinately regulated chemotaxis and resistance to copper by CsoR.

Keywords: chemotaxis; kinase CheA; CsoR; copper resistance; protein-protein interaction; *Pseudomonas putida*

Introduction

Chemotaxis is a widespread ability of motile bacteria to direct their movement

towards higher concentrations of beneficial chemicals or lower concentrations of toxic

chemicals[1,2]. Chemotaxis plays a vital role in bacterial exploration and adaptation to

complex environments[3,4]. The chemotaxis signaling pathway is extensively studied in

the enteric bacteria *Escherichia coli* and *Salmonella enterica* serovar Typhimurium[2,5].

In *E. coli*, the chemotaxis system consists of five methyl-accepting chemotaxis

proteins (MCPs) and six core components (the kinase CheA, the response regulator

CheY, the coupling protein CheW, the methylesterase CheB, the phosphatase CheZ,

and the methyltransferase CheR). In response to decreased attractant or increased

repellent, methylated MCPs and the coupling protein CheW activate CheA

autophosphorylation[6,7]. The phosphorylated CheA (CheA-P) is a phosphodonor for

the response regulator CheY and the methylesterase CheB. After CheY accepts the

phosphate group from CheA, it binds FliM and FliN of the motor-switch complex,

resulting in a switch in the flagellar rotational direction and bacteria tumbling[8].

Meanwhile, on a slower timescale, phosphorylated CheB demethylates the active

MCPs, thus reducing their ability to activate CheA, leading to decreased levels of

phosphorylated CheY and less tumbling[9]. The phosphatase CheZ contributes to signal

termination by removing the phosphoryl group from phosphorylated CheY, and the

methyltransferase CheR contributes to signal adaptation by catalyzing the methylation

of MCPs[2,5].

Among the bacterial chemotaxis system components, the kinase CheA is a

five-domain enzyme central to the chemotaxis signaling pathway[10,11]. The five

domains (P1-P5) of CheA each have distinct functions. The P1 domain (HPT domain)

contains the phosphoryl-accepting histidine that becomes phosphorylated, the P2

domain (CheY-binding domain) docks the response regulator proteins CheY and

CheB, the P3 domain (Dimer domain) dimerizes the CheA protein, the P4 domain

(HATPase_c domain) binds ATP and catalyzes phosphoryl transfer to the histidine

residue on P1, and the P5 domain (CheW-binding domain) couples CheA to other

chemotaxis components by binding both CheW and the chemoreceptors[10,11]. Except

for interacting with the components in chemotaxis system, CheA is reported to

interact with proteins from other systems. For example, in the plant pathogen

Xanthomonas oryzae pv. *oryzicola*, CheA interacts with and phosphorylates the

response regulator VemR to regulate bacterial virulence, motility, and EPS

production[12]. In *Vibrio parahaemolyticus*, a polarly localized protein ParP interacts

with CheA and prevents its dissociation from chemotaxis signaling arrays, facilitating

proper chemotaxis and accurate inheritance of these macromolecular chemotactic

machines[13]. In *Comamonas testosteroni*, CheA interacts with and phosphorylates the

response regulator FlmD, resulting in decreased biofilm formation[14]. In *Azospirillum*

brasilense with two chemotaxis signaling systems, the kinase CheA from each

chemotaxis signaling system physically interacts with the CheY response regulator of

another system[15]. In *Pseudomonas aeruginosa*, CheA interacts with the

phosphodiesterase DipA to regulate its subcellular localization and activity, leading to

individual cell heterogeneity and motility behavior diversity in bacterial populations[16].

These studies suggest that the CheA-mediated crosstalk between chemotaxis and

other systems coordinates complex behaviors in diverse bacteria.

In most living organisms, copper is an essential cofactor for enzymes involved in

fundamental processes such as respiration and photosynthesis[17,18]. However, copper

also has toxic effects on cells, and bacteria have several strategies to increase their

resistance to copper[19-21]. The direct bacterial response associated with copper

resistance is highly conserved and generally involves (a) sensing of the increased

copper concentration by sensors, (b) activation of bespoke transcriptional networks, (c)

overproduction of copper efflux pumps that secrete copper out of the cells, and (d)

recruitment of copper-binding and copper-oxidizing proteins that prevent copper from

interacting with cellular components[22-26]. The genome of *Pseudomonas putida* was

predicated to encode a dozen of proteins involved in copper resistance[27], including

copper sensor proteins (CopS-I, CopS-II), P-type ATPases for copper efflux (CueA,

CopA-I, CopA-II, CopB-I, and CopB-II), heavy-metal efflux complex components

(CusA, CusB, CusC, and CusF). Expression of these proteins were regulated by

copper-responsive positive regulatory proteins (CopR-I, CopR-II, CueR)[28,29]. CueR

regulated the expression of genes implicated in cytoplasmic copper homeostasis,

whereas CopR controlled the expression of genes involved in maintaining periplasmic

metal level[30]. Meanwhile, some proteins/systems appeared to be duplicated, and some

were proved to be functionally redundant[28,30]. Besides, extracellular polymeric

substances (EPS) in *P. putida* biofilms showed high affinity for most heavy metals,

including copper, thereby providing a protective barrier under copper stress[31-33].

In addition to the above copper resistance strategies, bacteria can avoid copper

stress through chemotaxis. For example, in *Caulobacter crescentus*, the reactive

oxygen species derived from cytoplasmic copper ions mediate the bacterial

chemotaxis to copper, and a potential cytoplasmic MCP McpR regulates bacterial

chemotaxis in response to cellular copper content, enabling bacteria to escape from

copper-rich environment[34,35]. However, unlike the widely reported mechanisms of

copper resistance in diverse bacterial species, the mechanism(s) of bacterial

chemotaxis to copper is poorly studied. Besides, it remains unclear how bacteria

coordinate chemotaxis and resistance to copper.

Since the kinase CheA plays a central role in chemotaxis signaling, identifying

CheA-interacting proteins would extend the knowledge of chemotaxis regulation. In

this study, by screening proteins that interacted with CheA in *Pseudomonas putida*,

we obtained 16 novel CheA-interacting proteins. Among the 16 proteins, CsoR, a

copper-binding transcription regulator, inhibited the autophosphorylation of CheA,

leading to decreased chemotaxis. Meanwhile, CsoR functioned as a DNA-binding

repressor to inhibit the expression of copper-resistance genes. Copper-binding of

CsoR relieved its inhibition of gene expression and chemotaxis.

Results

Identification of new CheA-interacting proteins in *P. putida*

We performed a pull-down assay to identify protein(s) interacting with CheA. Purified

$6 \times$ His-tagged CheA bound onto a Ni-NTA agarose column was used as "bait"

protein to pull out potential CheA-interacting "prey" protein(s) from whole cell lysate

of *P. putida* KT2440, and a blank Ni-NTA agarose column was used as a negative

control. All "prey" proteins from the CheA-binding column and control column were

collected, resolved by SDS–PAGE (Fig. 1a), and analyzed by mass spectrometry

(MS). In MS analysis, 43 proteins showed a significantly higher amount in the

CheA-binding column than in the control column (Log_2(fold change) > 2) (Fig. 1b,

Table S1). As expected, the "bait" protein CheA showed the highest amount, with a

Log_2(fold change) value close to 8. Meanwhile, the response regulator CheY and the

phosphatase CheZ (two proteins known to be associated with CheA) also showed high

Log_2(fold change) values. In addition to these three proteins (CheA, CheY, and CheZ),

the remaining 40 proteins were considered potential new CheA-interacting proteins.

To verify the above result, we performed bacterial two-hybrid (BTH) assay to test

the interactions between CheA and the 40 proteins. The results from the BTH assay

revealed that 19 proteins showed apparent interaction with CheA, including PP_2969

(CsoR), PP_1612 (Eno), PP_4111 (FusB), PP_5046 (GlnA), PP_1074 (GlpR),

PP_4728 (GrpE), PP_0853 (IspG), PP_1877 (MsrC), PP_2378 (NfuA), PP_1023

(Pgl), PP_5006 (PhaD), PP_0691 (ProB), PP_0148, PP_1644, PP_2683, PP_3177,

PP_3227, PP_3501, and PP_4460 (Fig. 1c). Meanwhile, the rest 21 proteins displayed

no significant interaction with CheA in BTH assay (Fig. S1a and S1b), suggesting that

they were false positive results in the pull-down assay. We further tested the

interactions between CheA and the 19 proteins in *P. putida* using bimolecular

fluorescence complementation (BiFC) assay. The results displayed that except three

proteins (PP_2683, PP_3227, and PP_4460) showed no apparent interaction with

CheA, the other 16 proteins displayed obvious interaction with CheA in the BiFC

assay (Fig. 1d, Fig. S2). Collectively, using pull-down, BTH, and BiFC assays, we

identified 16 new CheA-interacting proteins in *P. putida*, including CsoR, Eno, FusB,

GlnA, GlpR, GrpE, IspG, MsrC, NfuA, Pgl, PhaD, ProB, PP_0148, PP_1644,

PP_3177, and PP_3501.

Effect of CheA-interacting proteins on bacterial chemotaxis

Since CheA played a crucial role in bacterial chemotaxis, we wondered whether the

16 CheA-interacting proteins were involved in regulating chemotaxis. Thus, we

overexpressed each of the 16 proteins in wild-type *P. putida* and tested the chemotaxis

ability. The wild-type *P. putida* harboring empty vector (WT+pVec) was used as the

control. Chemotaxis ability was assessed using semisolid nutrient agar plates on

which bacteria formed large colonies (swimming zone) by generating and following

attractant gradients leading outward from the colony origin[36]. Growth of the 16 strains

in liquid medium showed a similar trend as that of the control strain (Fig. S3),

suggesting that the 16 proteins had no noticeable effect on bacterial growth. As shown

in Fig. 2a, five strains (WT+p*csoR*, WT+p*ispG*, WT+p*nfuA*, WT+p*phaD*, and

WT+p*PP_1644*) displayed smaller colony than the control strain (WT+pVec),

indicating a weaker chemotaxis ability in these five strains. The other 11 strains

showed a similar chemotaxis ability as WT+pVec. These results suggested that five

CheA-interacting proteins (CsoR, IspG, NfuA, PhaD, and PP_1644) inhibited

chemotaxis in *P. putida*.

CsoR and PhaD inhibit CheA autophosphorylation

CheA has autophosphorylation activity, and it can phosphorylate its cognate response

regulator[11]. To test whether phosphate transfer existed between CheA and the 16

proteins, we performed a phosphate transfer assay using purified proteins and

[^{32}P]ATP[γP]. We successfully purified 14 out of the 16 proteins but failed to purify

two proteins (GlpR and PP_3177) after several attempts. Thus, the two proteins were

not included in the phosphate transfer assay. The results showed that CheA exhibited a

strong autophosphorylation signal after incubation with [^{32}P]ATP[γP] for 45 min (Fig.

S4a). Then, each of the 14 proteins or CheY (positive control) was added to

phosphorylated CheA to investigate the phosphate transfer. The addition of CheY to

phosphorylated CheA led to a labeling of CheY and a reduction in the

phospholabeling of CheA (Fig. S4a), indicating a phosphate transfer happened

between CheA and CheY. However, no labeling of the 14 target proteins was observed,

and there was no apparent change in the phospholabeling of CheA after adding each

of the 14 proteins (Fig. S4a), suggesting no phosphate transfer happened between

CheA and the 14 tested proteins.

Then, we wondered whether the 14 proteins influenced CheA autophosphorylation.

To answer this question, we mixed CheA with each of the 14 proteins before adding

the substrate [^{32}P]ATP[γP]. The mixture containing CheA and bovine serum albumin

(BSA) was used as a negative control. As shown in Fig. 2b, CsoR/PhaD significantly

decreased the phospholabeling of CheA. In contrast, the other 12 proteins and BSA

had no apparent influence on the phospholabeling of CheA. We further tested the

impact of CsoR/PhaD on CheA autophosphorylation with a more detailed assay, in

which CheA was mixed with an increased amount of CsoR/PhaD. The results showed

that CheA phospholabeling decreased as CsoR/PhaD increased. In contrast, the

increase of BSA had no obvious influence on CheA phospholabeling (Fig. 2c and 2d).

These results indicated that CsoR and PhaD inhibited CheA autophosphorylation. It

was also possible that CsoR and PhaD degraded the substrate$[^{32}P]ATP[\gamma P]$ in the

reaction mixture, resulting in decreased CheA autophosphorylation. To test this

possibility, we examined whether CsoR and PhaD had ATPase activity. The known

ATPase FleQ was used as a positive control, and BSA was used as a negative control.

The results showed that adding FleQ into the reaction mixture caused a decrease in

ATP level (Fig. S4b), indicating the existence of ATPase activity. Meanwhile, the

addition of CsoR/PhaD exhibited no apparent influence on ATP level as the addition

of BSA (Fig. S4b), suggesting that CsoR and PhaD had no ATPase activity.

In the chemotaxis signaling pathway, CheA transfers the phosphate group to the

response regulator CheY to modulate flagellar rotation[2,5]. Since CsoR and PhaD

interacted with CheA, we wondered whether CsoR/PhaD influenced the phosphate

transfer between CheA and CheY. Thus, we incubated CsoR/PhaD with the

phosphorylated CheA before adding CheY to the reaction mixture. The results

revealed that CheA phospholabeling in the CsoR/PhaD-adding group reduced to a

similar level as that in the group without CsoR/PhaD (Fig. S4c), implying that

CsoR/PhaD had no apparent influence on the phosphate transfer between CheA and

CheY.

The domains of CheA involved in interacting with CsoR and PhaD

To further explore how CsoR and PhaD affect CheA autophosphorylation, we determined the domain(s) of CheA involved in interacting with CsoR and PhaD. Similar to the *E. coli* CheA, the *P. putida* CheA consists of five domains with distinct functions (Fig. 3a). We constructed five truncated CheA variants, with each missing one domain (termed CheA$_{\Delta HPT}$/CheA$_{\Delta YB}$/CheA$_{\Delta Dim}$/CheA$_{\Delta HATPase}$/CheA$_{\Delta WB}$) (Fig. 3a). Then we performed BTH assay to test the interaction between CsoR/PhaD and each of these truncated CheAs. The results revealed that three truncated CheAs (CheA$_{\Delta HPT}$, CheA$_{\Delta Dim}$, and CheA$_{\Delta HATPase}$) showed no interaction with CsoR and PhaD. In comparison, the other two truncated CheAs (CheA$_{\Delta YB}$ and CheA$_{\Delta WB}$) interacted with CsoR and PhaD with a similar intensity as the wild-type CheA did (Fig. 3b and 3c), indicating that the HPT (P1), Dimer (P3), and HATPase_c (P4) domains were essential for interacting with CsoR and PhaD, while the YB (P2) and WB (P5) domains were not required to interact with CsoR and PhaD. To further test this result, we cloned each of the five domains into a BTH vector and tested the interaction between CsoR/PhaD and each of the five domains. The result showed that the Dimer domain (Dim$_{CheA}$) interacted with CsoR and PhaD like the whole-length CheA did (Fig. 3b and 3c). The HPT domain (HPT$_{CheA}$) and the HATPase_c domain (HATPase$_{CheA}$) showed a weaker interaction with CsoR and PhaD. Meanwhile, the YB domain (YB$_{CheA}$) and the WB domain (WB$_{CheA}$) displayed no interaction with CsoR and PhaD (Fig. 3b and 3c). Together, these results revealed that the Dimer domain (P3) of CheA played a significant role, the HPT (P1) and HATPase_c (P4) domains played a minor role, while the CheY-binding (P2) and CheW-binding (P5) domains played no role in the interaction between CheA and CsoR/PhaD. Since the three domains involved in interacting with CsoR/PhaD were also essential for CheA

autophosphorylation　activity[10,11],　CsoR/PhaD　might　inhibit　CheA

autophosphorylation by inhibiting the function of the three domains.

Of the two proteins that inhibit CheA autophosphorylation, CsoR is annotated as a

metal-binding transcriptional repressor, and PhaD is annotated as a TetR family

transcriptional regulator[37]. Although the two proteins were quite different in size

(CsoR 10.8 kDa, PhaD 23.1 kDa), they interacted with the same domains of CheA to

inhibit its autophosphorylation, we wondered whether the two proteins shared

similarity in sequence and structure. BLAST result revealed no sequence similarity

between CsoR and PhaD (data not shown). We predicted structures of CsoR and PhaD

using AlphaFold (Fig. S5a and S5b), and compared their structures using Pymol.

However, the result revealed no significant structural similarity between the two

proteins (Fig. S5c).

CsoR is a metal-binding transcriptional repressor for metal-resistance genes

Previous research revealed that the expression of CsoR (also named MreA) was

induced by cadmium, nickel, zinc, and cobalt[38]. The homolog of CsoR in

Mycobacterium tuberculosis binds to copper and regulates the expression of

copper-resistance genes[39,40]. However, the metal-binding ability and the function of

the *P. putida* CsoR were not experimentally characterized. Thus, we tested the

metal-binding ability of the *P. putida* CsoR using MicroScale Thermophoresis (MST).

Six metal ions were involved in the assay, including copper (Cu^{2+}), zinc (Zn^{2+}), nickel

(Ni^{2+}), cobalt (Co^{2+}), cadmium (Cd^{2+}), and magnesium (Mg^{2+}). The results revealed

that CsoR bound to three out of the six tested metal ions (Cu^{2+}, Zn^{2+}, Ni^{2+}), and the

binding to Cu^{2+} was the strongest with a calculated binding constant (K_d) of 5.5 ± 1.98

μM. In contrast, the binding to Ni^{2+}/Zn^{2+} was weak (with K_d value of 125 ± 38 μM

and 253 ± 42 μM, respectively) (Fig. 4a). Meanwhile, CsoR showed no apparent

binding to $Co^{2+}/Cd^{2+}/Mg^{2+}$ under the same condition (Fig. 4a). We further constructed

a *csoR* deletion mutant ($\Delta csoR$) and investigated the effect of *csoR* deletion on the

expression of copper resistance genes using quantitative real-time PCR (qRT-PCR).

Three key copper resistance genes (*copA-I*, *copA-II*, and *copB-II*) from two operons

were chosen as targets in qRT-PCR. The results showed that *csoR* deletion

($\Delta csoR$+pVec) led to a weak but significant increase (about 1.3-fold) in the

expression of the three genes, and complementation ($\Delta csoR$+p*csoR*) decreased the

expression of the three genes (Fig. 4b). The addition of $CuCl_2$ (final concentration 10

μM) induced the expression of the three genes (about 5-fold) (Fig. 4b). However, no

noticeable difference in gene expression level was observed between WT+pVec and

$\Delta csoR$+pVec, as well as between $\Delta csoR$+pVec and $\Delta csoR$+p*csoR* (Fig. 4b), implying

that CsoR was not required for the copper-induced genes expression. Besides, the

deletion of *cheA* ($\Delta cheA$+pVec) displayed no obvious effect on the expression of the

three genes in either the presence or absence of copper (Fig. 4b). Since CsoR also

bound Zn^{2+} and Ni^{2+}, we investigated the influence of CsoR on the expression of

several other metal resistance genes, including three nickel-resistance genes (*nikA*,

nikB, and *nikD*), one zinc-resistance genes (*znuC*), two cadmium-resistance genes

(*cadA-I* and *cadA-III*), two cobalt-resistance gene (*cbiD* and *cbtA*), and three multiple

metal-resistance genes (*czcC-I*, *czcB-II*, and *PP_0026*). The results reveled that CsoR

repressed the expression of *nikB*, *cadA-I*, *cadA-III*, *cbtA*, *czcC-I*, *czcB-II*, and

PP_0026, and the inhibition degree was close to the inhibition degree of copper

resistance genes (Fig. S6a). Together, these results demonstrated that CsoR functioned

as a metal-binding repressor for metal resistance genes in *P. putida*.

Previous studies reported that CsoR bound the promoter of target genes to inhibit

gene expression[39,40]. We further tested the interaction between CsoR and the promoter

of *copA-I* using electrophoretic mobility shift assay (EMSA). The fragment of *copA-I*

promoter exhibited a stepwise increase in the shifted DNA amount (CsoR-DNA

complex), with the CsoR protein amount increasing from 5 to 20 μM (Fig. 4c).

Adding Cu^{2+} to the reaction mixture decreased the CsoR-DNA complex (Fig. 4c),

indicating that Cu^{2+} inhibited the interaction between CsoR and promoter DNA.

Previous study in *M. tuberculosis* showed that CsoR bound a single-monomer mole

equivalent of Cu^+ to form a trigonally coordinated complex[39], but our results indicated

that CsoR bound to Cu^{2+}. To further test whether Cu^+ bind to CsoR and affect its

DNA-binding ability, we added dithiothreitol (DTT) to the EMSA reaction mixture.

DTT can reduce Cu^{2+} to Cu^+ in solution[41]. As shown in the following Fig. 4d, the

addition of DTT (0.1 and 1 mM) decreased CsoR-DNA interaction in the presence of

0.2 mM Cu^{2+}, and the addition of DTT alone had no apparent influence on

CsoR-DNA interaction, indicating that DTT enhanced the inhibition of Cu^{2+} on

CsoR-DNA interaction. These results suggested that the Cu^+ converted from Cu^{2+} by

DTT had stronger inhibitory effect than Cu^{2+} on CsoR-DNA interaction, indicating

that CsoR bound more strongly to Cu^+ than to Cu^{2+}. Besides, Ni^{2+}, Zn^{2+}, and Cd^{2+} also

exhibited an inhibitory effect on the interaction between CsoR and promoter DNA,

but to a much lower extent compared with Cu^{2+} (Fig. 4e). Meanwhile, Co^{2+} and Mg^{2+}

displayed no obvious effect on the CsoR-DNA interaction (Fig. 4e). These results

showed that CsoR was a promoter-binding transcriptional repressor, and binding to

metal (especially copper) decreased the interaction between CsoR and promoter

DNA.

To test the effect of CsoR and CheA on bacterial copper resistance, we tested the

growth of Δ*csoR* and Δ*cheA* under different copper concentrations using both solid

agar plate and liquid medium. However, in both cases, there was no significant

difference between the growth trend of Δ*csoR*/Δ*cheA* and the WT strain at different

copper concentrations (Fig. S6b and S6c). This might attribute to the fact that CsoR

was a repressor, and the expression of copper resistance genes in WT was similar to

that in Δ*csoR* under copper stress (Fig. 4b).

Copper inhibits the interaction between CheA and CsoR

Since CsoR interacted with CheA and bound copper, we wondered whether copper

affected the interaction between CheA and CsoR. Thus, we investigated the

interaction between CheA and CsoR under different $CuCl_2$ concentrations using MST.

As revealed in Fig. 5a, CheA showed strong interaction with CsoR in the absence of

Cu^{2+} (K_d= 0.17 ± 0.1 μM), and the addition of Cu^{2+} (20 and 200 μM) led to increased

binding constant (0.59 ± 0.2 μM and 2.15 ± 0.97 μM), indicating that copper

decreased the interaction between CheA and CsoR. A similar trend was observed in

the pull-down assay, in which the amount of CsoR bound by CheA gradually

decreased with the concentration of Cu^{2+} increased from 2 to 20 μM (Fig. 5b).

A previous study on *M. tuberculosis* CsoR revealed three residues that played a

vital role in copper-binding[39]. The alignment assay showed that the three residues

were conserved among CsoR homologs from several bacterial species, including the *P.*

putida CsoR (Fig. S7a). To test the role of the three residues in copper-binding, we

individually replaced each of the three residues (Cys40, His65, and Cys69) with

alanine and tested the copper-binding ability of these mutated CsoRs ($CsoR_{C40A}$,

$CsoR_{H65A}$, and $CsoR_{C69A}$). MST results showed that the $CsoR_{C69A}$ displayed

significantly decreased copper-binding ability compared to wild-type CsoR. In

contrast, $CsoR_{C40A}$ and $CsoR_{H65A}$ showed a slight decrease in copper-binding ability

compared with that of the wild-type CsoR (Fig. S7b), suggesting that the Cys69

residue of CsoR played a critical positive role in copper-binding. Besides, the results

from EMSA also supported this conclusion, in which the mutation of C69A, but not

C40A/H65A, noticeably inhibited the effect of Cu^{2+} on CsoR-DNA interaction (Fig.

S7c and S7d). BTH assay revealed that the three point-mutated CsoRs interacted with

CheA with a similar intensity to the wild-type CsoR (Fig. S7e). Results from the

pull-down assay showed that the addition of Cu^{2+} significantly decreased the amount

of $CsoR_{C40A}$ and $CsoR_{H65A}$ bound by CheA, but had less effect on the amount of

$CsoR_{C69A}$ bound by CheA under the same condition (Fig. 5b). Together, these results

revealed that copper bound CsoR and inhibited its interaction with CheA.

Copper relives the inhibition of CsoR on bacterial chemotaxis

Since CsoR interacted with CheA and inhibited its autophosphorylation, and copper

hindered the interaction between CheA and CsoR, we speculated that copper might

relieve the inhibition of CsoR on CheA autophosphorylation. To test this hypothesis,

we added CsoR and copper to the reaction mixture containing CheA and then

analyzed the autophosphorylation of CheA using $[^{32}P]ATP[\gamma P]$. CsoR alone

significantly inhibited CheA autophosphorylation, while adding CsoR and copper

showed weaker inhibition on CheA autophosphorylation (Fig. 5c). Besides, $CsoR_{C69A}$

showed similar inhibitory effects on CheA autophosphorylation in the presence and

absence of Cu^{2+} (Fig. 5c), indicating that binding with CsoR was the prerequisite for

Cu^{2+} to relieve the inhibition of CsoR on CheA autophosphorylation.

We further tested the effect of CsoR on chemotaxis in the presence of copper using

the semisolid plate. As shown in Fig. 5d, in the absence of Cu^{2+}, the overexpression of

point-mutated CsoR ($CsoR_{C40A}$/$CsoR_{H65A}$/$CsoR_{C69A}$) led to a similar decrease (about

40%) in chemotaxis as the overexpression of wild-type CsoR. In comparison, in the

presence of 200 μM Cu^{2+}, the inhibitory effect of CsoR/$CsoR_{C40A}$/$CsoR_{H65A}$

overexpression on chemotaxis was weaker (about 20%). However, the inhibitory

effect of $CsoR_{C69A}$ overexpression on chemotaxis was not affected by Cu^{2+} (Fig. 5d).

These results demonstrated that binding copper relieved the inhibition of CsoR on

bacterial chemotaxis.

CsoR inhibits chemorepellent response to copper

Excess copper is toxic to cells, and bacteria avoid high copper concentrations through

chemotaxis. Since CsoR interacted with CheA and inhibited chemotaxis, we

wondered about the role of CsoR in bacterial chemotaxis to copper. Thus, we tested

the chemotaxis response of *P. putida* to copper gradient and the role of CsoR and

CheA in this response using semisolid plates. The copper gradient was achieved by

placing an agar plug containing 200 mM $CuCl_2$ in the center of a semisolid plate, and

bacterial cells were spotted two centimeters away from the plug to test their

chemotaxis to Cu^{2+}. An agar plug without $CuCl_2$ was also placed in the center of a

semisolid plate and used as a negative control. As shown in Fig. 6a, the swimming

zone of all tested strains was a circle shape in the control plate without $CuCl_2$.

However, all swimming zones showed an oval shape in the plate with $CuCl_2$ gradient

(Fig. 6a), in which the bacterial movement distance near the plug (D1) was short, and

the movement distance from the plug (D2) was long, indicating that the strains

showed chemorepellent response to $CuCl_2$. The RI value (Response index value,

RI=D1/(D1+D2)) was further calculated to characterize the strength of the

chemorepellent response to $CuCl_2$. The results showed that WT+p*csoR* displayed a

higher RI value (0.428 ± 0.015) than WT+pVec (0.373 ± 0.021) (Fig. 6b). Meanwhile,

Δ*csoR*+pVec showed a lower RI value (0.324 ± 0.013) than WT+pVec, and

complementation increased the RI value to wild-type level (Fig. 6b). These results

suggested that CsoR inhibited the chemorepellent response to copper. Besides, both

cheA deletion mutant (Δ*cheA*+pVec) and *cheA csoR* double deletion mutant

($\Delta csoR\Delta cheA$+pVec) displayed no chemotaxis ability in either the presence or

absence of copper gradient (Fig. 6a and 6b), indicating that CsoR inhibited

chemotaxis ability in a CheA-dependent manner.

Using time-lapse microscopy experiments and cell-tracking analysis, we further

examined the bacterial chemorepellent response to Cu^{2+}. In the control group without

chemokine (Cu^{2+} gradient), cells of all tested strains swam randomly in all directions

(Fig. 6c). The center of mass (defined as the average of all single cell endpoints, and it

reflects the movements of target strain and the strength of chemotaxis response) of all

tested strains showed no apparent difference (Fig. 6d). In the group with chemokine

(Cu^{2+} gradient), cells of WT+pVec, WT+p$csoR$, $\Delta csoR$+pVec, and $\Delta csoR$+p$csoR$

migrated towards the lower concentration of Cu^{2+} (Fig. 6c). In comparison, cells of

$\Delta cheA$+pVec and $\Delta csoR\Delta cheA$+pVec still swam randomly in all directions (Fig. 6c),

indicating that *P. putida* cells showed a chemorepellent response to Cu^{2+} in a

CheA-dependent manner. The center of mass value of WT+p$csoR$ (13.85 ± 1.92 μm)

was smaller than that of WT+pVec (21.77 ± 3.60 μm) (Fig. 6d). Meanwhile, the center

of mass value of $\Delta csoR$+p$csoR$ (14.16 ± 1.41 μm) was smaller than that of

$\Delta csoR$+pVec (21.39 ± 2.02 μm) (Fig. 6d), suggesting that CsoR inhibited bacterial

chemorepellent response to Cu^{2+}. The velocities (cell migration speeds) of cells from

WT+p$csoR$, $\Delta csoR$+pVec, and $\Delta csoR$+p$csoR$ were similar to that from WT+pVec both

in the presence and absence of Cu^{2+} gradient, implying that CsoR had no evident

influence on bacterial migration speed. In contrast, the velocities of cells from

$\Delta cheA$+pVec and $\Delta csoR\Delta cheA$+pVec were lower than that from WT+pVec (Fig. 6e),

indicating that CheA played a positive role in bacterial migration speed. Together,

these results demonstrated that CsoR inhibited the chemorepellent response to copper

in a CheA-dependent manner.

The interaction between CheA and CsoR exists in several bacterial species

The role of CsoR in regulating copper resistance has been reported in several bacterial

species, including *Acidithiobacillus caldus*[42], *Bacillus subtilis*[43], *Bradyrhizobium*

diazoefficiens[44], *Corynebacterium glutamicum*[45], *Listeria monocytogenes*[46], *M.*

tuberculosis[40], *Staphylococcus aureus*[47], *Streptomyces lividans*[48], and *Thermus*

thermophilus[49]. BLAST results showed that four of the nine above species (*A. caldus*,

B. diazoefficiens, *B. subtilis*, and *L. monocytogenes*) had both *cheA* and *csoR* on their

genomes. Besides, in addition to *P. putida*, *cheA* and *csoR* coexist in other

Pseudomonas species, including *Pseudomonas fluorescens*, *Pseudomonas syringae*,

and *Pseudomonas stutzeri*. We wondered whether the CheA-CsoR interaction also

occurred between proteins from these strains. Thus, we tested the interaction between

CheA and CsoR of the same strain via BTH assay. The result showed that the

CheA-CsoR interaction existed between proteins from *A. caldus*, *B. subtilis*, *P.*

syringae, and *P. stutzeri* (Fig. 7). However, CheA and CsoR from *B. diazoefficiens*, *L.*

monocytogenes, and *P. fluorescens* showed no apparent interaction (Fig. 7). Besides,

the intensity of CheA-CsoR interaction was more vigorous between the two proteins

from *B. subtilis*, but weaker between that from *A. caldus*, *P. syringae*, and *P. stutzeri*

(Fig. 7). These results suggested that except in *P. putida*, the interaction between

CheA and CsoR also existed in other bacterial species.

Discussion

Integrating components from different systems provides a straightforward mechanism

for coordinating signaling from various systems. This study identified an interaction

between the chemotaxis kinase CheA and the copper-responsive transcriptional

repressor CsoR in *P. putida*. Further analysis revealed that CsoR inhibited bacterial

chemotaxis via interacting with CheA and hindering its autophosphorylation.

Meanwhile, CsoR regulated copper resistance by modulating the expression of

copper-resistance genes. Together with previous reports[50-52], we proposed a potential

model to describe the function of CsoR in regulating copper resistance and bacterial

chemotaxis. As shown in Fig. 8, under low copper levels, CsoR molecules exist

mainly in none copper-binding status (free CsoR) in the cell, and the free CsoR forms

tetramer and binds to promoters of copper-resistance genes (such as *copA-I*), leading

to repressed gene transcription and copper resistance ability. Meanwhile, the free

CsoR interacts with CheA and inhibits its autophosphorylation, decreasing bacterial

chemotaxis ability. Under high copper levels, more copper-binding CsoR molecules

exist, and copper-binding changes the conformation of the CsoR tetramer and releases

CsoR from promoters, leading to increased gene transcription and copper resistance.

Besides, the copper-binding of CsoR decreases the interaction between CsoR and

CheA, which relieves the inhibition of CsoR on CheA autophosphorylation, resulting

in increased chemotaxis ability.

In classical chemotaxis signaling, MCP on the cell membrane senses external

signaling molecules and regulates bacterial chemotaxis by mediating CheA

autophosphorylation activity[5,7]. MCPs can directly bind diverse external signaling

molecules, such as amino acids, dipeptides, sugars, tricarboxylic acid cycle

intermediates, aromatic molecules, and inorganic phosphate[53]. However, there are

only a few reports on the relationship between MCP and metal ions[54-56], and no

evidence supports that MCP senses metal ions by direct binding. It is possible that

bacteria sense and trigger chemotaxis to metal ions differently from that they sense

and trigger chemotaxis to external signaling molecules like amino acids. Our results

provide a mechanism by which bacteria sense copper and regulate chemotaxis via the

copper-responsive repressor CsoR. Through the interaction between CsoR and CheA,

bacteria coordinately regulated chemotaxis and resistance to copper stress, which

would favor the bacteria to better adapt to complex environments. Besides, the

interaction between CsoR and CheA was not limited to the proteins from *P. putida*,

and it was also found in proteins from several other bacterial species (Fig. 7),

implying that the regulation of chemotaxis and resistance to copper via the interaction

between CsoR and CheA might be a widespread regulatory mechanism.

Although the *P. putida* CsoR functioned as a copper-responsive regulator to

modulate the expression of copper-resistance genes, its effect on gene expression was

much weaker than its homologous protein in other bacterial species. In *M.*

tuberculosis, *B. subtilis*, *C. glutamicum*, *L. monocytogenes*, and *S. aureus*, deletion of

csoR resulted in an about 10-fold increase in the expression of target genes in the

absence of copper[40,43,45,46,47]. In contrast, deletion of *csoR* in *P. putida* led to a slight

but reproducible increase (about 1.3-fold) in gene expression in the absence of copper

(Fig. 4b). This difference might be attributed to the existence of several key regulators

that activated the expression of copper-resistance genes in response to copper in *P.*

putida, such as CueR and CopR. CueR positively regulated the expression of *cueA*,

encoding a copper-transporting P1-type ATPase that played a crucial role in copper

resistance[28]. CopR was essential for expressing several genes implicated in

cytoplasmic copper homeostasis, such as *copA-II*, *copB-II*, and *cusA*[30,57]. The

existence of these positive regulators makes the function of CosR a secondary or even

dispensable insurance in the expression of copper-resistance genes. Consistent with

this, there is no CosR homolog in *P. aeruginosa*, and copper homeostasis in *P.*

aeruginosa is mainly controlled by CueR and CopR[29,30].

Through pull-down, BTH, and BiFC assays, we obtained 16 new CheA-interacting

proteins involved in different physiological processes (Table S1). Among the 16

proteins, five proteins (CsoR, IspG, NfuA, PhaD, and PP_1644) inhibited bacterial

chemotaxis on semisolid plates (Fig. 2a). Our study here focused on the physiological

role of CsoR-CheA interaction. Still, the function of other interactions remained

unclear. PhaD is a TetR family transcriptional regulator that behaves as a carbon

source-dependent activator of the *pha* cluster related to polyhydroxyalkanoates (PHAs)

biosynthesis[58,59]. Bacterial PHAs are isotactic polymers synthesized under

unfavorable growth conditions in the presence of excess carbon sources. PHAs are

critical in central metabolism, acting as dynamic carbon reservoirs and reducing

equivalents[60]. The interaction between PhaD and CheA leads one to speculate that

there might be some connection between PHA synthesis and bacterial chemotaxis.

Another CheA-interacting protein, PP_1644, also attracts our interest. PP_1644 is

annotated as a NAD(P)H dehydrogenase involved in cyclic electron transport and

respiration processes. Exploring the physiological role of the interaction between

CheA and these proteins in the future helps to reveal the association between the

chemotaxis process and other physiological metabolisms.

Materials and Methods

Bacterial strains and growth conditions

All strains and plasmids used in this study are listed in Supplementary Table S2.

Unless specifically mentioned, *E. coli* strains were grown in lysogeny broth (LB)

medium at 37°C. *P. putida* KT2440 and its derivative strains were cultured at 28°C in

LB medium or chemically defined M9 minimal medium supplemented with 40 mM

glucose as carbon source. Antibiotics were used, when required for plasmid

maintenance or transformants screening, at the following concentrations: kanamycin

(50 mg/L), carbenicillin (50 mg/L), chloramphenicol (25 mg/L), and gentamycin (20

mg/L for *E. coli* or 40 mg/L for *P. putida*).

Plasmid and strain construction

The routine cloning of DNA fragments into plasmid was performed by following a T5 exonuclease-dependent method[61]. Briefly, a 14 base pairs (bp) homologous end was added to the 5' of primer during synthesizing. Amplified DNA fragments and linearized plasmid containing the same homologous end were incubated in a reaction buffer containing T5 exonuclease (0.04 U) and left at 30□ for 40 min before transforming into *E. coli* competent cells. Primers used for plasmid construction are listed in Supplementary Table S3. All cloning steps involving PCR were verified by commercial sequencing (Tsingke, Wuhan, China).

Gene deletion mutant was constructed by homologous recombination using the suicide plasmid pBBR401. For example, to construct a markerless *P. putida csoR* deletion mutant, ∼ 800 bp from the chromosomal regions flanking *csoR* (upstream region and downstream region) were PCR-amplified. The PCR products were cloned into pBBR401 to create pBBR401-*csoR*UP-DOWN. Then, the final plasmid was transferred to *P. putida* by electroporation. The integration strain was selected on plates containing gentamicin. After subculturing the integration strain in LB medium without antibiotics six times (12 h each time), single colonies were obtained by plate streaking. Then, colonies losing gentamicin resistance were kept for further verification. The *csoR* delete mutant was confirmed by PCR and sequencing.

To generate a *csoR/cheA* overexpression plasmid, a DNA fragment containing the complete *csoR/cheA* was PCR amplified. The product was cloned into expressional vector pBBR403 to yield pBBR403-*csoR/cheA*. The expression of *csoR/cheA* on pBBR403-*csoR/cheA* was controlled by an inducible *tac* promoter. To construct a vector for target protein expression and purification, we amplified and cloned the

target gene into pET-28a with $6 \times$ His tag or pHS-Strep with Strep II tag. Overlapping

PCR was used to create point mutations of CsoR. To construct a C40A point mutation

in CsoR, we amplified two fragments with two primer pairs (CsoR$_{C40A}$ s1/CsoR$_{C40A}$

a1 and CsoR$_{C40A}$ s2/CsoR$_{C40A}$ a2). The CsoR$_{C40A}$ a1 and the CsoR$_{C40A}$ s2 shared

reverse complementary sequences containing the point mutation in which the original

TGC codon of arginine was replaced by GCC of alanine. The two fragments were

mixed in a 1:1 ratio to perform overlapping extension. The final PCR product was

cloned into pET-28a and pBBR403. The mutation in *csoR* was confirmed by

sequencing.

Expression and purification of His/Strep II-tagged proteins

For the expression of His/Strep II-tagged protein, overnight culture of *E. coli* BL21

carrying the construct of target proteins was 1:100 diluted into LB medium and

incubated for 4 h at 37°C. Then, 0.4 mM IPTG (isopropyl-D-thiogalactopyranoside)

was added to induce protein expression. After 4 h incubation at 16°C, cells were

harvested and resuspended in lysing buffer (10 mM Tris-Cl [pH 7.8], 300 mM KCl,

and 10% (w/v) glycerol). The harvested cells were lysed using a pressure cell

breaking apparatus, and cell debris was removed by centrifugation at 15000 rpm for

20 min. The supernatants were then filtered through a 0.22-μm-pore-size filter and

loaded onto a Ni-NTA Resin column (for His-tagged protein) or Strep-Tactin Resin

column (for Strep II-tagged protein). Target proteins were eluted using an imidazole

gradient (50/100/150/250mM imidazole for His-tagged protein) or 5 mM biotin (for

Strep II-tagged protein) and then dialyzed overnight against lysing buffer to remove

imidazole. The concentrations of obtained proteins were determined using BCA assay.

Protein-protein pull-down assay

Protein-protein pull-down assay was used to identify CheA-interacting protein and

test the effect of copper on CheA-CsoR interaction. Briefly, $6 \times$ His/Strep II-tagged

CheA was induced and loaded to a Ni-NTA/Strep-Tactin column as described above.

Then, overnight cultured wild-type KT2440/BL21 strain expressing CsoR was

harvested, lysed, and filtered before adding to the same column. For the pull-down

assay to identify CheA-interacting protein, the same volume of wild-type extract was

added to a blank Ni-NTA column as a negative control. For the pull-down assay to

test the effect of copper on CheA-CsoR interaction, various amounts of $CuCl_2$ were

mixed with the cell extract containing CsoR before being added to the Strep-Tactin

column. Then, the columns were sealed and incubated at 4°C with 40 rpm shaking.

After 2 h incubation, the supernatant was removed, and the columns were washed

with lysing buffer containing 20 mM imidazole (for Ni-NTA column) or lysing buffer

(for Strep-Tactin column) to wash the unspecific binding protein away. Then, elution

buffer containing 250 mM imidazole (for the Ni-NTA column) or 5 mM biotin (for

the Strep-Tactin column) was added to wash down all proteins on the column. The

eluted proteins were collected and resolved by 12.5% SDS–PAGE followed by

Coomassie blue staining and mass spectrometry analyses.

Mass spectrometry (MS)-based protein sequencing

After Coomassie blue staining of eluted proteins obtained from pull-down assay, the

whole lane of the experimental or control sample was excised from gels and prepared

for MS analysis. Protein from the excised gels was extracted with a Micro Protein

PAGE Recovery Kit (Sangon Biotech, China) following the operating instructions.

Then, trypsin digestion of extracted protein was performed with 1 g trypsin (Promega,

USA) and incubated at 37°C overnight. The digestion was terminated by adding

trifluoroacetic acid (TFA). Then, desalting was subsequently performed using Zip-tip

(Merck Millipore, Ireland). Peptides were eluted from the Zip-tip with 50 µL of

matrix solution (5 mg/mL α-cyano-4-hydroxycinnamic acid, 50% acetonitrile, 0.1%

TFA). The supernatant was collected and concentrated to a final volume of 10 μL in a

centrifugal concentrator.

The samples were analyzed using the MALDI-TOF/TOF mass spectrometer

(Applied Biosystems, USA). Mass spectra were recorded in the positive-ion mode,

averaging 2500 laser shots per spectrum. Mass spectra (excluding trypsin autolytic

peptides and other known background ions) were searched against the *P. putida*

proteome from the UniProt database to identify the proteins. The search was

performed using trypsin digestion, allowing two missed cleavages, specifying

carbamidomethyl-Cys as a fixed modification, and setting a peptide mass tolerance of

± 1.6 Da. The global false discovery rate (FDR) cutoff for peptide and protein

identification was set to 0.01. An intensity-based absolute quantification (iBAQ)

algorithm was used to rank the relative abundance of different proteins as previously

described[62]. iBAQ percentage of specific proteins in the experimental sample

(iBAQ_T (%)) and control sample (iBAQ_CK (%)) were used to represent relative

protein concertation. Log_2 (iBAQ_T/iBAQ _CK) fold change of ≥ 2 or ≤-2 and a *P*

value of ≤ 0.05 was considered significantly different.

Bacterial two-hybrid (BTH) assay

For bacterial two-hybrid analysis of protein-protein interactions of *P. putida* proteins

expressed in *E. coli*, each ORF was cloned in-frame with the T18 and T25 fragments

of adenylate cyclase ORF in vectors pUT18C and pKT25. Primers used to amplify

each ORF are listed in Table S3. The resulting vectors were co-transformed into *E.*

coli BTH101 and plated onto LB agar plate supplemented with 50 mg/L carbenicillin,

50 mg/L kanamycin, 40 mg/L 5-bromo-4-chloro-3-indolyl-b-D-galactopyranoside

(X-gal), and 0.5 mM IPTG, and incubated at 28°C for 48 h. Three co-transformants

for each assay were cultured to stationary phase in LB broth at 28°C, then spotted

onto an LB agar plate supplemented as above, and incubated for 60 h at 28°C. Plates

were then photoed on a Tanon 2500 scanner. After the photographs were taken, the

colonies on the plates were scraped off, and the LacZ activities of obtained cells were

measured using o-nitrophenyl-β-galactopyranoside (ONPG) as substrate, as described

before[63]. The experiments were repeated three times with three technical repeats per

sample, and the data are presented as Miller units.

Bimolecular fluorescence complementation (BiFC) assay

BiFC was used to analyze protein-protein interactions as previously described[64].

Briefly, to determine the interaction between two interested proteins (such as CheA

and CsoR), CheA was fused to KN151 (the N-terminal of mLumin), and CsoR was

fused to LC151 (the C-terminal of mLumin), yielding a recombinant plasmid

pBBR403-CheA-KN151-CsoR-LC151. Then, the recombinant plasmid was

transformed into the wild-type *P. putida* strain. Transformants were picked and

cultured in LB medium containing 40 mg/L gentamycin and 0.5 mM IPTG for 24 h at

28□. Then, images of the dark-filed and bright field of the transformant cells were

obtained using FV1000 CLSM (Olympus, Japan) equipped with a 100×/1.4 oil

immersion objective lens. Besides, the transformant cells were washed twice with 0.9%

NaCl, and then fluorescence intensities and OD_{600} were measured using a Spark

microplate reader (Tecan, Switzerland). Fluorescence intensity and OD_{600} were

detected using the black and transparent microplate, respectively. The excitation and

the emission wavelength to detect fluorescence were 587 nm and 620 nm, respectively,

and the experiment was repeated twice with triplicates.

Microscale thermophoresis (MST) assay

MST was performed to analyze the interaction between two proteins or proteins and

metal ions. Briefly, to test the interaction between CsoR and copper. A green

fluorescent protein (GFP) encoding gene was fused to the end of *csoR* in

pET-28a-*csoR* to achieve fusion expression. The fusion protein CsoR-GFP was

induced and purified as described above in protein induction and purification. The

obtained CsoR-GFP was dialyzed with MST buffer (50 mM Tris-HCl [pH 7.8], 150

mM NaCl, 0.05% Tween 20). The MST assay was performed on a Monolith

Instrument NT.115 device using standard treated capillaries (NanoTemper

Technologies, Germany). The concentration of CsoR-GFP was constant at 250 nM,

and the $CuCl_2$ concentration was varied from 0.031 to 1000 μM with a 2-fold gradient.

The experiment was recorded using the Nano-BLUE fluorescent detector.

Measurements were performed in MST buffer. The MO. Affinity Analysis software

(version 2.3) was used to calculate the dissociation constant (K_d) from triplicate reads

of measurements.

***In vitro* phosphorylation assays**

For the autophosphorylation reaction, purified CheA (3 μM) was incubated in

phosphorylation buffer containing 50 mM Tris–HCl [pH 7.5], 15 mM $MgCl_2$, and 50

mM NaCl. The reaction was initiated by adding 0.03 μCi of $[^{32}P]ATP[\gamma P]$

(PerkinElmer, USA) to the mixture. SDS-PAGE loading buffer containing SDS and

EDTA was added to the mixture to terminate the reaction at the indicated time. To test

the effect of target proteins on CheA autophosphorylation, target protein (10 μM) was

mixed with CheA (3 μM) for 10 min before adding $[^{32}P]ATP[\gamma P]$. To test the

transphosphorylation reaction, CheA was autophosphorylated before CheY/target

protein (10 μM) was added to the mixture, and the reaction mixture was incubated at

30°C for different time intervals before being terminated with SDS-PAGE loading

buffer. Samples were heated at 95°C for 5 min and then resolved by 12.5%

SDS–PAGE. After drying of the gels, products were visualized by autoradiography.

RNA extraction and real-time RT-PCR (qRT-PCR) assay

P. putida cells were cultured in M9 minimal medium supplemented with 40 mM

glucose as carbon source for 24 h. Then, cells were harvested and washed thrice with

sterilized phosphate buffer saline (PBS) before being divided into two equal parts.

One part was resolved with fresh M9 medium and another with M9 medium

containing 10 μM $CuCl_2$. After 30 min incubation, cells were harvested for RNA

extraction using a total RNA extraction reagent (Vazyme R401-01, China) as

recommended by the manufacturer. 1 μg extracted RNA was digested with DNase I

and reverse transcribed to cDNA using a reverse transcription kit (Takara RR047A,

Japan), and cDNA was used as the template for qRT-PCR analysis. The qRT-PCR

assay was performed using Power SYBRTM Green PCR mix (Applied Biosystems

4367659, USA) and analyzed using a QuantStudio™ 3 Real-Time PCR System

(Applied Biosystems, USA). The *rpoD* gene was selected as an internal control. The

primers used in qRT-PCR analysis are listed in Supplementary Table S3. All

experiments were performed thrice with three technical repeats per sample.

Electrophoretic mobility shift assay (EMSA)

EMSA was used to test the interaction between CsoR and *copA-I* promoter DNA.

Equal amounts of DNA (60 ng) were added to binding reactions with various

quantities of CsoR in binding buffer (10 mM Tris-Cl [pH 7.8], 50 mM KCl, 20 mM

$MgCl_2$, 5% glycerol, 20 μL total reaction volume). CsoR was incubated with promoter

DNA for 20 min at room temperature. Reaction mixtures containing

Cu^{2+}/DTT/Cu^{2+}+DTT were performed as described above, except

Cu^{2+}/DTT/Cu^{2+}+DTT was incubated with CsoR for 10 min before adding DNA. All

reaction solutions were loaded onto 5% acrylamide gel and electrophoresed at 150 V

for 40 min in $0.5 \times$ TBE buffer (45 mM Tris-Cl [pH 7.8], 45 mM borate, 1 mM

EDTA). Gels were stained with ethidium bromide before being digitized using a

scanner (Tanon 2500, China).

Bacterial chemotaxis assay

The chemotaxis ability of *P. putida* strains was assessed by using semisolid plate and

μ-slide Chemotaxis plate (Ibidi 80326, Germany). For the method with semisolid

plates, an agar plug (1% agar) containing 200 mM $CuCl_2$ was placed in the center of a

LB semisolid plate (0.25% agar) and left at room temperature for 12 h to achieve a

$CuCl_2$ gradient on the semisolid plate. An agar plug without $CuCl_2$ was also placed in

the center of a semisolid plate and used to test bacterial chemotaxis without copper.

For the assay to investigate the effect of copper on bacterial chemotaxis, 200 μM

$CuCl_2$ (final concentration) was mixed with semisolid LB before making a semisolid

plate. Overnight growth *P. putida* cells were washed and resuspended with fresh M9

medium and adjusted to the same optical density ($OD_{600} = 0.5$). Then, 2 μL of

resuspended cultures was spotted 2 centimeters away from the agar plug/plate center,

and the plate was incubated for 16 h (for the control plate) or 18 h (for the

copper-containing plate) at 28 °C before digital photographs were taken. The distance

from the point of inoculation to the edge of the colony growth closest to the agar plug

(D1) and the distance from the point of inoculation to the edge of the colony growth

farthest from the agar plug (D2) were measured. The response index (RI) value was

calculated to characterize the bacterial response to $CuCl_2$ as previously described[36].

The RI was calculated using the following equation: RI = D1/(D1 + D2). RI values

greater than 0.52 and less than 0.48 correspond to attractant and repellent responses,

and intermediate values represent nonresponses.

For the method with μ-slide Chemotaxis plate (a commercially available

microfluidic device with a channel connecting two reservoirs), the reservoirs were

filled with the two bacterial solutions (with or without $CuCl_2$) following a modified

version of the manufacturer's "Fast Method" protocol. Briefly, overnight cultures

were inoculated with M9 medium and grown at 28 °C, 180 rpm, until they reached an

optical density (OD_{600}) between 0.3 and 0.35. 1 ml of bacterial culture was washed

twice with fresh M9 medium by centrifugation (3 min at 3000 rpm), and finally

diluted to a target OD_{600} of 0.015 with fresh M9 supplemented with 0.01% Tween 20,

with or without 2 mM $CuCl_2$ for injection into the chemotaxis device. First, the entire

device was overfilled with buffer free of $CuCl_2$ or bacteria through the filling ports,

and then the central channel's ports were closed with plugs. About 65 μl was removed

from one reservoir, replaced by 65 μl of $CuCl_2$-free bacterial solution, and then this

reservoir's ports were closed. Finally, all liquid was removed from the other reservoir

and replaced with a bacterial solution containing $CuCl_2$. For control measurements,

neither bacterial solution contained $CuCl_2$. Phase contrast microscopy recordings were

obtained at room temperature on a Nikon Ti-E inverted microscope using an sCMOS

camera (PCO Edge 4.2) and a 40 x objective lens. Recordings were obtained starting

from 20 min after filling the device. Three 15 s-long recordings of cells in the

observation area were obtained at 15 fps. Then, single-cell tracking analysis was

performed using ImageJ's Manual Tracking and Chemotaxis Tool plugins.

Statistical analysis

Statistical analyses were performed using Graph-Pad Prism (version 9.0.0). Student's

t-test or an analysis of variance (ANOVA) was used to analyze the significance of

differences in LacZ activity, fluorescence intensity, swimming zone diameter, gene

expression, response index, center of mass, and velocity. A P value less than 0.05 was

considered statistically significant.

Acknowledgment

This work was supported by the National Key Research and Development Program of

China [2020YFC1806803], the Hubei Provincial Natural Science Foundation of

China [2024AFB691], the National Natural Science Foundation of China [31900054],

and the Fundamental Research Funds for the Central Universities

[2662022SKQD002].

Competing interests

The authors declare no competing interests.

Data availability

Relevant data supporting the critical findings of this study are available within the

article and the Supplementary Information file. The absorbance, microscopic

photographs, gels, plates, and EMSA data generated in this study have been provided

in the Source data file.

References

1. Keegstra, J.M., Carrara, F. & Stocker, R. The ecological roles of bacterial

chemotaxis. *Nat. Rev. Microbiol.* **20**, 491–504 (2022).

2. Wadhams, G.H. & Armitage, J.P. Making sense of it all: bacterial chemotaxis. *Nat.*

Rev. Mol. Cell. Biol. **5**, 1024–1037 (2004).

3. Zboralski, A. & Filion, M. Genetic factors involved in rhizosphere colonization by

phytobeneficial *Pseudomonas* spp. *Comput. Struct. Biotechnol. J.* **18**, 3539–3554

(2020).

4. Boin, M.A., Austin, M.J. & Häse, C.C. Chemotaxis in *Vibrio cholerae*. *FEMS.*

Microbiol. Lett. **239**, 1–8 (2004).

5. Porter, S.L., Wadhams, G.H. & Armitage, J.P. Signal processing in complex
chemotaxis pathways. *Nat. Rev. Microbiol.* **9**, 153–165 (2011).

6. Parkinson, J.S., Hazelbauer, G.L. & Falke, J.J. Signaling and sensory adaptation in
Escherichia coli chemoreceptors: 2015 update. *Trends. Microbiol.* **23**, 257–266
(2015).

7. Ortega, A., Zhulin, I.B. & Krell, T. Sensory repertoire of bacterial chemoreceptors.
Microbiol. Mol. Biol. Rev. **81**, e00033-17 (2017).

8. Welch, M., Oosawa, K., Aizawa, S. & Eisenbach, M. Phosphorylation-dependent
binding of a signal molecule to the flagellar switch of bacteria. *Proc. Natl. Acad.*
Sci. USA. **90**, 8787–8791 (1993).

9. Stewart, R.C. Activating and inhibitory mutations in the regulatory domain of
CheB, the methylesterase in bacterial chemotaxis. *J. Biol. Chem.* **268**, 1921–1930
(1993).

10. Bilwes, A.M., Alex, L.A., Crane, B.R. & Simon, M.I. Structure of CheA, a
signal-transducing histidine kinase. *Cell.* **96**, 131–141 (1999).

11. Muok, A.R., Briegel, A. & Crane, B.R. Regulation of the chemotaxis histidine
kinase CheA: A structural perspective. *Biochim. Biophys. Acta. Biomembr.* **1862**,
183030 (2020).

12. Cai, L., Ma, W., Zou, L., Xu, X., Xu, Z., Deng, C., Qian, W., Chen, X. & Chen, G.
Xanthomonas oryzae pv. *oryzicola* response regulator VemR is co-opted by the
sensor kinase CheA for phosphorylation of multiple pathogenicity-related targets.
Front. Microbiol. **13**, 928551 (2022).

13. Ringgaard, S., Zepeda-Rivera, M., Wu, X., Schirner, K., Davis, B.M. & Waldor,
M.K. ParP prevents dissociation of CheA from chemotactic signaling arrays and
tethers them to a polar anchor. *Proc. Natl. Acad. Sci. USA.* **111**, E255–264 (2014).

14. Huang, Z., Wang, Y.H., Zhu, H.Z., Andrianova, E.P., Jiang, C.Y., Li, D., Ma, L.,

Feng, J., Liu, Z.P., Xiang, H., Zhulin, I.B. & Liu, S.J. Cross talk between

chemosensory pathways that modulate chemotaxis and biofilm formation. *mBio.*

10, e02876-18 (2019).

15. O'Neal, L., Gullett, J.M., Aksenova, A., Hubler, A., Briegel, A., Ortega, D., Kjær,

A., Jensen, G. & Alexandre, G. Distinct chemotaxis protein paralogs assemble

into chemoreceptor signaling arrays to coordinate signaling output. *mBio.* **10**,

e01757-19 (2019).

16. Kulasekara, B.R., Kamischke, C., Kulasekara, H.D., Christen, M., Wiggins, P.A.

& Miller, S.I. c-di-GMP heterogeneity is generated by the chemotaxis machinery

to regulate flagellar motility. *eLife.* **2**, e01402 (2013).

17. Kim, B.E., Nevitt, T. & Thiele, D.J. Mechanisms for copper acquisition,

distribution and regulation. *Nat. Chem. Biol.* **4**, 176–185 (2008).

18. Tsang, T., Davis, C.I. & Brady, D.C. Copper biology. *Curr. Biol.* **31**, R421–R427

(2021).

19. Giachino, A. & Waldron, K.J. Copper tolerance in bacteria requires the activation

of multiple accessory pathways. *Mol. Microbiol.* **114**, 377–390 (2020).

20. Andrei, A., Öztürk, Y., Khalfaoui-Hassani, B., Rauch, J., Marckmann, D., Trasnea,

P.I., Daldal, F. & Koch, H.G. Cu homeostasis in bacteria: the ins and outs.

Membranes (Basel). **10**, 242 (2020).

21. Hyre, A., Casanova-Hampton, K. & Subashchandrabose, S. Copper homeostatic

mechanisms and their role in the virulence of *Escherichia coli* and *Salmonella*

enterica. EcoSal. Plus. **9**, eESP00142020 (2021).

22. Novoa-Aponte, L., Ramírez, D. & Argüello, J.M. The interplay of the

metallosensor CueR with two distinct CopZ chaperones defines copper

homeostasis in *Pseudomonas aeruginosa*. *J. Biol. Chem.* **294**, 4934–4945 (2019).

23. Öztürk, Y., Andrei, A., Blaby-Haas, C.E., Daum, N., Daldal, F. & Koch, H.G. Metabolic sensing of extracytoplasmic copper availability via translational control by a nascent exported protein. *mBio.* **14**, e0304022 (2023).

24. Roy, G., Antoine, R., Schwartz, A., Slupek, S., Rivera-Millot, A., Boudvillain, M. & Jacob-Dubuisson, F. Posttranscriptional regulation by copper with a new upstream open reading frame. *mBio.* **13**, e0091222 (2022).

25. Zuily, L., Lahrach, N., Fassler, R., Genest, O., Faller, P., Sénèque, O., Denis, Y., Castanié-Cornet, M.P., Genevaux, P., Jakob, U., Reichmann, D., Giudici-Orticoni, M.T. & Ilbert, M. Copper induces protein aggregation, a toxic process compensated by molecular chaperones. *mBio.* **13**, e0325121 (2022).

26. Dennison, C., David, S. & Lee, J. Bacterial copper storage proteins. *J. Biol. Chem.* **293**, 4616–4627 (2018).

27. Cánovas, D., Cases, I. & de Lorenzo, V. Heavy metal tolerance and metal homeostasis in *Pseudomonas putida* as revealed by complete genome analysis. *Environ. Microbiol.* **5**, 1242-1256 (2003).

28. Adaikkalam, V. & Swarup, S. Molecular characterization of an operon, *cueAR*, encoding a putative P1-type ATPase and a MerR-type regulatory protein involved in copper homeostasis in *Pseudomonas putida*. *Microbiology.* **148**, 2857-2867 (2002).

29. Hofmann, L., Hirsch, M. & Ruthstein, S. Advances in understanding of the copper homeostasis in *Pseudomonas aeruginosa*. *Int. J. Mol. Sci.* **22**, 2050 (2021).

30. Quintana, J., Novoa-Aponte, L. & Argüello, J.M. Copper homeostasis networks in the bacterium *Pseudomonas aeruginosa*. *J. Biol. Chem.* **292**, 15691-15704 (2017).

31. Fang, L., Wei, X., Cai, P., Huang, Q., Chen, H., Liang, W. & Rong, X. Role of

extracellular polymeric substances in Cu(II) adsorption on *Bacillus subtilis* and

Pseudomonas putida. Bioresour. Technol. **102**, 1137-1141 (2011).

32. Lin, H., Wang, C., Zhao, H., Chen, G. & Chen, X. A subcellular level study of

copper speciation reveals the synergistic mechanism of microbial cells and EPS

involved in copper binding in bacterial biofilms. *Environ. Pollut*. **263**, 114485.

(2020)

33. Lin, H., Wang, C., Zhao, H., Chen, G. & Chen, X. Interaction between copper and

extracellular nucleic acids in the EPS of unsaturated *Pseudomonas putida* CZ1

biofilm. *Environ. Sci. Pollut. Res. Int*. **25**, 24172-24180 (2018).

34. Lawarée, E., Gillet, S., Louis, G., Tilquin, F., Le, Blastier.S., Cambier, P. &

Matroule, J.Y. *Caulobacter crescentus* intrinsic dimorphism provides a prompt

bimodal response to copper stress. *Nat. Microbiol.* **1**, 16098 (2016).

35. Louis, G., Cherry, P., Michaux, C., Rahuel-Clermont, S., Dieu, M., Tilquin, F.,

Maertens, L., Van, Houdt.R., Renard, P., Perpete, E. & Matroule, J.Y. A

cytoplasmic chemoreceptor and reactive oxygen species mediate bacterial

chemotaxis to copper. *J. Biol. Chem.* **299**, 105207 (2023).

36. Pham, H.T. & Parkinson, J.S. Phenol sensing by *Escherichia coli* chemoreceptors:

a nonclassical mechanism. *J. Bacteriol.* **193**, 6597-6604 (2011).

37. Winsor, G.L., Griffiths, E.J., Lo, R., Dhillon, B.K., Shay, J.A. & Brinkman, F.S.

Enhanced annotations and features for comparing thousands of *Pseudomonas*

genomes in the Pseudomonas genome database. *Nucleic. Acids. Res.* **44(D1)**,

D646-D653 (2016).

38. Haritha, A., Sagar, K.P., Tiwari, A., Kiranmayi, P., Rodrigue, A., Mohan, P.M. &

Singh, S.S. MrdH, a novel metal resistance determinant of *Pseudomonas putida* KT2440, is flanked by metal-inducible mobile genetic elements. *J. Bacteriol.* **191**, 5976-5987 (2009).

39. Liu, T., Ramesh, A., Ma, Z., Ward, S.K., Zhang, L., George, G.N., Talaat, A.M., Sacchettini, J.C. & Giedroc, D.P. CsoR is a novel *Mycobacterium tuberculosis* copper-sensing transcriptional regulator. *Nat. Chem. Biol.* **3**, 60-68 (2007).

40. Marcus, S.A., Sidiropoulos, S.W., Steinberg, H. & Talaat, A.M. CsoR is essential for maintaining copper homeostasis in *Mycobacterium tuberculosis*. *PLoS. One.* **11**, e0151816 (2016).

41. Kr zel, A., Lesniak, W., Jezowska-Bojczuk, M., Mlynarz, P., Brasuñ, J., Kozlowski, H. & Bal, W. Coordination of heavy metals by dithiothreitol, a commonly used thiol group protectant. *J. Inorg. Biochem.* **84**, 77-88 (2001).

42. Hou, S., Tong, Y., Yang, H. & Feng, S. Molecular insights into the copper-sensitive operon repressor in *Acidithiobacillus caldus*. *Appl. Environ. Microbiol.* **87**, e0066021 (2021).

43. Smaldone, G.T. & Helmann, J.D. CsoR regulates the copper efflux operon *copZA* in *Bacillus subtilis*. *Microbiology.* **153**, 4123-4128 (2007).

44. Pacheco, P.J., Cabrera, J.J., Jiménez-Leiva, A., Torres, M.J., Gates, A.J., Bedmar, E.J., Richardson, D.J., Mesa, S., Tortosa, G. & Delgado, M.J. The copper-responsive regulator CsoR is indirectly involved in *Bradyrhizobium diazoefficiens* denitrification. *FEMS. Microbiol. Lett.* **370**, fnad084 (2023).

45. Teramoto, H., Yukawa, H. & Inui, M. Copper homeostasis-related genes in three separate transcriptional units regulated by CsoR in *Corynebacterium glutamicum*. *Appl. Microbiol. Biotechnol.* **99**, 3505-3517 (2015).

46. Corbett, D., Schuler, S., Glenn, S., Andrew, P.W., Cavet, J.S. & Roberts, I.S. The

combined   actions   of   the   copper-responsive   repressor   CsoR   and

copper-metallochaperone CopZ modulate CopA-mediated copper efflux in the

intracellular pathogen *Listeria monocytogenes*. *Mol. Microbiol.* **81**, 457-472

(2011).

47. Baker, J., Sengupta, M., Jayaswal, R.K. & Morrissey, J.A. The *Staphylococcus*

aureus CsoR regulates both chromosomal and plasmid-encoded copper resistance

mechanisms. *Environ. Microbiol.* **13**, 2495-2507 (2011).

48. Dwarakanath, S., Chaplin, A.K., Hough, M.A., Rigali, S., Vijgenboom, E. &

Worrall, J.A.R. Response to copper stress in *Streptomyces lividans* extends

beyond genes under direct control of a copper-sensitive operon repressor protein

(CsoR). *J. Biol. Chem.* **287**, 17833-17847 (2012).

49. Sakamoto, K., Agari, Y., Agari, K., Kuramitsu, S. & Shinkai, A. Structural and

functional characterization of the transcriptional repressor CsoR from *Thermus*

thermophilus HB8. *Microbiology.* **156**, 1993-2005 (2010).

50. Chang, F.M., Coyne, H.J., Cubillas, C., Vinuesa, P., Fang, X., Ma, Z., Ma, D.,

Helmann, J.D., García-de, los.Santos.A., Wang, Y.X., Dann, III.CE. & Giedroc,

D.P. Cu(I)-mediated allosteric switching in a copper-sensing operon repressor

(CsoR). *J. Biol. Chem.* **289**, 19204-19217 (2014).

51. Jacobs, A.D., Chang, F.M., Morrison, L., Dilger, J.M., Wysocki, V.H., Clemmer,

D.E. & Giedroc, D.P. Resolution of stepwise copperativities of copper binding by

the homotetrameric copper-sensitive operon repressor (CsoR): Impact on

structure and stability. *Angew. Chem. Int. Ed. Engl.* **54**, 12795-12799 (2015).

52. Tan, B.G., Vijgenboom, E. & Worrall, J.A. Conformational and thermodynamic

hallmarks of DNA operator site specificity in the copper sensitive operon

repressor from *Streptomyces lividans*. *Nucleic. Acids. Res*. **42**, 1326-1340 (2014).

53. Bi, S. & Lai, L. Bacterial chemoreceptors and chemoeffectors. *Cell. Mol. Life. Sci.*

72, 691-708 (2015).

54. Chandrashekhar, K., Srivastava, V., Hwang, S., Jeon, B., Ryu, S. & Rajashekara,

G. Transducer-like protein in *Campylobacter jejuni* with a role in mediating

chemotaxis to iron and phosphate. *Front. Microbiol.* **9**, 2674 (2018).

55. Li, X., Sun, M., Zhang, L., Finlay, R.D., Liu, R. & Lian, B. Widespread bacterial

responses and their mechanism of bacterial metallogenic detoxification under

high concentrations of heavy metals. *Ecotoxicol. Environ. Saf.* **246**, 114193

(2022).

56. Martín-Mora, D., Reyes-Darias, J.A., Ortega, Á., Corral-Lugo, A., Matilla, M.A.

& Krell, T. McpQ is a specific citrate chemoreceptor that responds preferentially

to citrate/metal ion complexes. *Environ. Microbiol.* **18**, 3284-3295 (2016).

57. Quaranta, D., McEvoy, M.M. & Rensing, C. Site-directed mutagenesis identifies a

molecular switch involved in copper sensing by the histidine kinase CinS in

Pseudomonas putida KT2440. *J. Bacteriol.* **191**, 5304-5311 (2009).

58. de, Eugenio.L.I., Galán, B., Escapa, I.F., Maestro, B., Sanz, J.M., García, J.L. &

Prieto, M.A. The PhaD regulator controls the simultaneous expression of the *pha*

genes involved in polyhydroxyalkanoate metabolism and turnover in

Pseudomonas putida KT2442. *Environ. Microbiol.* **12**, 1591-1603 (2010).

59. Tarazona, N.A., Hernández-Arriaga, A.M., Kniewel, R. & Prieto, M.A. Phasin

interactome reveals the interplay of PhaF with the polyhydroxyalkanoate

transcriptional regulatory protein PhaD in *Pseudomonas putida*. *Environ.*

Microbiol. **22**, 3922-3936 (2020).

60. Gregory, D.A., Taylor, C.S., Fricker, A.T.R., Asare, E, Tetali, S.S.V., Haycock,

J.W. & Roy, I. Polyhydroxyalkanoates and their advances for biomedical

applications. *Trends. Mol. Med.* **28**, 331-342 (2022).

61. Xia, Y., Li, K., Li, J., Wang, T., Gu, L. & Xun, L. T5 exonuclease-dependent
assembly offers a low-cost method for efficient cloning and site-directed
mutagenesis. *Nucleic. Acids. Res.* **47**, e15 (2019).

62. Schwanhäusser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen,
W. & Selbach, M. Global quantification of mammalian gene expression control.
Nature. **473**, 337-342 (2011).

63. Schaefer, J., Jovanovic, G., Kotta-Loizou, I. & Buck, M. Single-step method for
β-galactosidase assays in *Escherichia coli* using a 96-well microplate reader. *Anal.*
Biochem. **503**, 56-57 (2016).

64. Chu, J., Zhang, Z., Zheng, Y., Yang, J., Qin, L., Lu, J., Huang, Z.L., Zeng, S. &
Luo, Q. A novel far-red bimolecular fluorescence complementation system that
allows for efficient visualization of protein interactions under physiological
conditions. *Biosens. Bioelectron.* **25**, 234-239 (2009).

Figure legends

Fig. 1 Screening and verifying proteins interacting with CheA. (a) Protein samples

obtained in pull-down assay and detected by SDS/PAGE. The "bait" protein CheA on

the gel was indicated. Lanes 1, 2, and 3 are samples from the control column, and

lanes 4, 5, and 6 are samples from the CheA-binding column. M represents a protein

marker. (b) The volcano plot shows the *P*-value and fold-change of all proteins

identified in MS analyses. Red spots represent proteins that showed two or higher

folds in the experimental group compared with the control group ($P < 0.05$). Blue

spots represent proteins with a higher amount in the control group ($P < 0.05$). Grey

spot proteins showed no apparent difference between the two groups ($P > 0.05$). (c)

Detect the interaction between CheA and indicated proteins via BTH. Blue indicates

protein-protein interaction in the colony after 60 h of incubation, while white

indicates no protein-protein interaction. A colony containing T25a-zip and T18C-zip

plasmids was used as a positive control (CK+), and a colony containing empty T25a

and T18C plasmids was used as a negative control (CK-). The LacZ activities of

colonies were shown above the colonies. (d) The red fluorescence intensities in BiFC

assay. The results in panels c and d are the average of three independent assays. Error

bars represent standard deviations. The asterisks above the column denote significant

differences (**$P < 0.01$) between indicated strains and CK- strain. "ns." represents

none statistically significant between indicated strain and CK- strain.

Fig. 2 CsoR and PhaD inhibit CheA autophosphorylation. (a) Chemotaxis of

indicated strains on semisolid plates. Photos of colonies on the top were taken after 16

h incubation at 28°C. Diameters of colonies (swimming zone) shown at down were

calculated from three replicates. The asterisks above the column denote significant

differences (*$P < 0.05$, **$P < 0.01$) between indicated strains and control strain

(WT+pVec) analyzed by Student's t-test. (b) Effect of the 14 proteins on the CheA

autophosphorylation. The name/ID and the concentration of tested proteins added in

each lane are indicated above the gel. CK represents CheA alone in the reaction

mixture. BSA is used as a negative control. The ladder represents a protein marker. (c

and d) CsoR (c) and PhaD (d) affect CheA autophosphorylation. The concentration of

tested proteins added in each lane is indicated above the gel. The time represents the

time of the CheA autophosphorylation reaction. The SDS-PAGE gels in panels b, c,

and d were detected by Coomassie Blue Staining (Above) and autoradiograph

(Below). The experiments for panels b, c, and d were repeated three times, and a

representative photo was shown. The relative autoradiograph intensity of the CheA

band was calculated using Image J software and shown below each lane.

Fig. 3 CheA domains involved in interacting with CsoR and PhaD. (a) Schematic

diagram of CheA and truncated CheA proteins. The predicted domains are based on

the Pfam database and the amino acid positions where the predicted domains start and

end are shown. (b) The interaction between CheA domains and CsoR/PhaD was tested

using BTH. Blue indicates protein-protein interaction in the colony after 60 h of

incubation, while white indicates no protein-protein interaction. A colony containing

T25a-zip and T18C-zip plasmids was used as a positive control (CK+), and a colony

containing empty T25a and T18C plasmids was used as a negative control (CK-). (c)

Confirmation of BTH interactions in panel B by LacZ activity assay. The results are

the average of three independent assays. Error bars represent standard deviations. The

asterisks above the column denote significant differences (**$P < 0.01$) between

indicated strains and CK- strain analyzed by Student's t-test. "ns." represents none

statistically significant between indicated strain and CK- strain.

Fig. 4 CsoR is a metal-binding repressor. (a) MST analysis of the interaction

between CsoR-GFP and metal ions. CsoR-GFP (250 nM) was incubated with

increasing concentrations of metal ions. (b) Analysis of relative transcription level of

target genes in wild-type (WT+pVec), *csoR* mutant (Δ*csoR*+pVec), complemented

strain (Δ*csoR*+p*csoR*), and *cheA* mutant (Δ*cheA*+pVec) in the presence and absence of

$CuCl_2$ (10 μM) by qRT-PCR. The results are the average of three independent assays.

Error bars represent standard deviations. The asterisks represent statistically

significant differences between the two compared strains (*$P < 0.05$, **$P < 0.01$). "ns."

represents none statistically significant between two compared strains. (c) Analysis

for interactions between CsoR and *copA-I* promoter DNA using EMSA. (d) The effect

of DTT/$CuCl_2$+DTT on the interaction between CsoR and *copA-I* promoter DNA. (e)

The effect of indicated metal ions on the interaction between CsoR and *copA-I*

promoter DNA. The concentrations of CsoR, metal ions, and DTT in panels c, d, and

e added in each lane are shown above the gel. Free DNA and CsoR-DNA complex are

indicated.

1010

Fig. 5 Copper inhibits the interaction between CheA and CsoR. (a) MST analysis

of the interaction between CheA-GFP and CsoR in the presence of Cu^{2+}. CheA-GFP

(160 nM) was incubated with increasing concentrations of CsoR. (b) SDS-PAGE

detected protein samples obtained in a pull-down assay. The "bait" protein

Strep-CheA and the "prey" protein His-CsoR on the gel were indicated. The gel

showed the influence of Cu^{2+} on the amount of "prey" protein His-CsoR. The

concentration of Cu^{2+} added in each pull-down assay was displayed above the gel.

The SDS-PAGE gel was detected by Coomassie Blue Staining. (c) CheA

autophosphorylation in the presence of CsoR and Cu^{2+}. The tested proteins and Cu^{2+}

concentrations added in each lane are indicated above the gel. The SDS-PAGE gel

was detected by Coomassie Blue Staining (Above) and autoradiograph (Below). The

relative intensity of the CsoR band in panel b and the autoradiograph intensity of the

CheA band in panel c were calculated using Image J software and shown below each

lane. (d) Chemotaxis of indicated strains on semisolid plates supplied with or without

$CuCl_2$. Photos of colonies on the top were taken after 16 h (for the control plate) or 18

h (for the copper-adding plate) incubation at 28°C. The diameters of colonies were

measured and normalized to the diameters of WT+pVec, shown below. The asterisks

above the column denote significant differences (**$P < 0.01$) between two indicated

strains analyzed by Student's t-test. "ns." represents none statistically significant

between two compared strains.

1031

Fig. 6 Role of CsoR in bacterial chemotaxis to copper. (a) Chemotaxis of indicated

strains in the absence and presence of copper gradient. The chemotaxis rings and

chemotaxis distance (D1/D2) were indicated by arrows. The red arrows pointed at the

agar plug with or without copper in the center of the plate. The green dots represented

the sites where the bacteria were initially inoculated on the semisolid plate. The assay

was performed with three repeats, and a representative photo was shown. (b) RI value

(D1/(D1+D2)) of indicated strains shown in panel a. (c) Aggregated trajectories of

individual tested strain cells in the absence and presence of copper gradient. The

tracking data presented is a composite of two experiments performed in duplicate (n =

100 cells). The overall directionality of migration is depicted in the rose diagram in

the upper right corner of each single-track summary. (d) Center of mass of tested

strains in the presence of copper gradients. It represents the average of all single-cell

endpoints. The results of panels b and d are the average of three independent assays.

Error bars represent standard deviations. The asterisks represent statistically

significant differences between the two indicated strains (*$P < 0.05$, **$P < 0.01$). "ns."

represents no statistically significant between the two indicated strains. (e) Velocity

analysis of indicated strains in the presence or absence of copper gradient (n = 100

cells). The lowercase letters above each bar in panel e indicate significant differences

($P < 0.05$).

1051

Fig. 7 The interaction between CheA and CsoR from indicated bacterial species.

The interaction between CsoR and CheA was tested by using BTH. The LacZ

activities of colonies were shown above the colonies. The results are the average of

three independent assays. Error bars represent standard deviations. The asterisks

above the column denote significant differences (**$P < 0.01$) between indicated

strains and CK- strain analyzed by Student's t-test. "ns." represents none statistically

significant between indicated strain and CK- strain.

1059

Fig. 8 A proposed model for describing how CsoR coordinates chemotaxis and

resistance to copper in *P. putida*. Under low Cu^+/Cu^{2+} levels, more none

Cu^+/Cu^{2+}-binding CsoR molecules (free CsoR) exist in the cell, and the free CsoR

forms tetramer and binds to promoters of copper-resistance genes (such as *copA-I* and

copA-II), leading to repressed gene transcription and low copper resistance.

Meanwhile, free CsoR interacts with CheA and inhibits its autophosphorylation

activity, decreasing chemotaxis ability. In contrast, more Cu^+/Cu^{2+}-binding CsoR

molecules exist under high Cu^+/Cu^{2+} levels, and the Cu^+/Cu^{2+}-binding changes the

conformation of the CsoR tetramer and releases CsoR from target promoters, leading

to increased gene transcription and copper resistance. Besides, Cu^+/Cu^{2+}-binding of

CsoR decreases the interaction between CsoR and CheA, which relieves the inhibition

of CsoR on CheA autophosphorylation, resulting in increased chemotaxis ability.

1072

Supplementary Information

Fig. S1 Detect the interaction between CheA and the 40 proteins using BTH. CheA

was cloned into T25a (a) and T18C (b). Blue indicates protein-protein interaction, and

white indicates no interaction in the colony after 60 h of incubation. A colony

containing T25a-zip and T18C-zip plasmids was used as a positive control (CK+), and

a colony containing empty T25a and T18C plasmids was used as a negative control

(CK-).

1080

1081 **Fig. S2** Detect the interaction between CheA and indicated proteins using BiFC.

1082 Representative images for each pair of proteins were shown. Jun-KN151+Fos-LC151

1083 and KN151+LC151 were used as CK+ and CK-, respectively. RFP channel images

1084 (fluorescent field), bright field images, and overlay images of the same field are

1085 shown. Scale bar = 2 μm.

1086

1087 **Fig. S3** Growth curve of the 16 overexpression strains and wild-type strain in liquid

1088 LB broth (100 mL in a 250 mL triangular glass flask, at 28°C with 180 rpm shaking).

1089 The optical density at 600 nm (OD_{600}) was used to characterize the growth of bacterial

1090 cells in the medium.

1091

Fig. S4 Role of target proteins in the CheA-mediated transphosphorylation. (a)

Transphosphorylation between CheA and the 14 proteins. Target proteins were added

to the phosphorylated CheA and incubated for 10 s before adding termination buffer.

CheY was added as a positive control. (b) ATPase activity of indicated proteins. The

amount of remaining ATP after incubation with 10 μM indicated proteins for 30 min.

BSA was used as a negative control. The asterisks represent statistically significant

differences between FleQ and BSA (**P < 0.01). "ns." represents no statistically

significant between the indicated protein and BSA. (c) Effect of CsoR and PhaD on

the transphosphorylation between CheA and CheY. The PT time in seconds (2 s and

10 s) represents the time of transphosphorylation. The bands of CheA, CheY, CsoR,

and PhaD on the gel were indicated with arrows. The relative autoradiograph intensity

of the CheA band was calculated using Image J software, shown below each lane.

1104

Fig. S5 Predicted structure of the *P. putida* CsoR (a) and PhaD (b) obtained using an

online AlphaFold server (https://www.alphafold.ebi.ac.uk). Position of the first and

the last amino acid residue of CsoR and PhaD are indicated on the structure. (c) The

structure alignment of CsoR and PhaD was performed with PyMOL software. The

protein name and its structure were shown with the same color. The differences

between protein structures were quantified using the Root Mean Square Deviation

(RMSD) index. An RMSD value lower than 2 was considered a high similarity

between two compared proteins.

1113

Fig. S6 Function of CsoR in the expression of metal resistant genes and the

bacterial growth under copper stress. (a) Analysis of relative transcription level of

metal resistant genes in wild-type (WT+pVec), *csoR* mutant (Δ*csoR*+pVec), and

complemented strain (Δ*csoR*+p*csoR*) by qRT-PCR. The results are the average of

three independent assays. Error bars represent standard deviations. The asterisks

represent statistically significant differences between the two compared strains (*$P <$

0.05, **$P < 0.01$). (b and c) Effect of *csoR*/*cheA* deletion on bacterial growth under

copper stress. Growth of WT, Δ*csoR*, and Δ*cheA* on M9 minimal medium agar plate

(b) and in liquid 1/4 LB medium (c) containing different $CuCl_2$ concentrations. A

same amount of bacterial culture (OD600=0.5) was spotted onto the plate and

incubated at 28°C for 24 h. The concentrations of $CuCl_2$ used in the assay were

indicated above.

1126

Fig. S7 Role of the three conserved residues in the Cu^{2+}-binding ability of CsoR.

(a) *P. putida* CsoR and CsoR homologs of indicated bacterial species are aligned

using an online ClustalW server (https://www.genome.jp/tools-bin/clustalw). The

number above the sequence represents the amino acid order of *S. saprophyticus* CsoR.

The identical residues are shown as white on red letters. Similar residues are shown in

red in blue boxes. Asterisks indicated the three conserved residues involved in

Cu^{2+}-binding. The sequence identity of CsoR homologs to the *P. putida* CsoR was

calculated. (b) MST analysis of the interaction between wild-type and mutated CsoRs

and Cu^{2+}. Wild-type/mutated CsoR (250 nM) was incubated with increasing

concentrations of Cu^{2+}. (c and d) Effect of point mutation on the binding of CsoR to

copA-I promoter in the presence of Cu^{2+} in the EMSA assay. Free DNA and

CsoR-DNA complex are indicated. (e) Detect the interaction between CheA and

point-mutated CsoR using BTH. The LacZ activities of colonies are shown below.

The result is the average of three independent assays. The data represent mean values

with standard deviations. "ns." represents none statistically significant between

indicated strain and CK- strain analyzed by Student's t-test.

1143

1144   **Table S1** Target proteins identified in a pull-down assay. iBAQ_T (%) and iBAQ_CK

1145   (%) represent the percentage of certain proteins in experimental and control samples,

1146   respectively. The *P* values are for three technical replicates. Protein names and

1147   descriptions are based on the annotated genome as indicated at

1148   http://www.pseudomonas.com.

1149

1150   **Table S2** Strains and plasmids used in this work.

1151

1152   **Table S3** Primers used in this work.

a

CheA — HPT (8–100) | CheY-binding (111–320) | Dimer (361–420) | HATPase_c (468–607) | CheW-binding (614–741) — 747

CheA$_{\Delta HTP}$ — CheY-binding | Dimer | HATPase_c | CheW-binding — 660

CheA$_{\Delta YB}$ — HPT | Dimer | HATPase_c | CheW-binding — 590

CheA$_{\Delta Dim}$ — HPT | CheY-binding | HATPase_c | CheW-binding — 694

CheA$_{\Delta HATPase}$ — HPT | CheY-binding | Dimer | CheW-binding — 613

CheA$_{\Delta WB}$ — HPT | CheY-binding | Dimer | HATPase_c — 627

HPT$_{CheA}$ 1 — HPT — 113   **YB$_{CheA}$** 1 — CheY-binding — 230

Dim$_{CheA}$ — Dimer — 80   **HATPase$_{CheA}$** 1 — HATPase_c — 165

WB$_{CheA}$ 1 — CheW-binding — 147

b

c

Legend:
- T18C-CheA
- T18C-CheA$_{\Delta HPT}$
- T18C-CheA$_{\Delta YB}$
- T18C-CheA$_{\Delta Dim}$
- T18C-CheA$_{\Delta HATPase}$
- T18C-CheA$_{\Delta WB}$
- T18C-HPT$_{CheA}$
- T18C-YB$_{CheA}$
- T18C-Dim$_{CheA}$
- T18C-HATPase$_{CheA}$
- T18C-WB$_{CheA}$

Y-axis: Miller Units

X-axis: T25a-*csoR*, T25a-*phaD*

a

b

c

d